PIECEWISE LINEAR HOMEOMORPHISMS OF THE REAL LINE

NICK GILL AND IAN SHORT

ABSTRACT. Brin and Squier described how to classify the elements of the group of piecewise linear homeomorphisms of the real line according to conjugacy. We supply a short account of the key step in their classification. The results in this document are unoriginal. The proofs are a little sketchy in places.

Let $\operatorname{PLF}^+([0,1])$ denote the group of orientation preserving piecewise linear homeomorphisms of [0,1] that are locally affine at all but a finite number of points in [0,1]. Let \mathcal{F} denote the subset of $\operatorname{PLF}^+([0,1])$, which consists of homeomorphisms f such that f(x) > x for each $x \in (0,1)$. Membership of \mathcal{F} is preserved by conjugation in $\operatorname{PLF}^+([0,1])$. Identifying the conjugacy classes that make up \mathcal{F} is a key step in understand conjugacy in $\operatorname{PLF}^+([0,1])$ and similar groups of piecewise linear homeomorphisms.

A node of a map f from $\text{PLF}^+([0,1])$ is a point x in (0,1) at which f is not locally affine. Let x_f denote the smallest node of f in (0,1). We also define $f'_-(x)$ and $f'_+(x)$ to be the left and right derivatives of f at x. Define $f^*(x) = f'_+(x)/f'_-(x)$ so that $f^*(x) \neq 1$ if and only if x is a node of f. By applying the chain-rule to the left and right derivatives of f, we deduce that there is a chain-rule for the operation $f \mapsto f^*$ which resembles the usual chain-rule for differentiation.

We associate to f two objects α_f and β_f that will be seen to be the only quantities necessary for a conjugacy classification in $\text{PLF}^+([0,1])$ of elements of \mathcal{F} . Define $\alpha_f = f'_-(x_f)$; this is the slope of the line segment in the graph of f that protrudes from 0. We can apply the chain-rule near 0 to see that α is invariant under conjugation. To define β_f , first consider the the function $\phi_f : [x_f, f(x_f)) \to \mathbb{R}^+$, given by

$$x \mapsto \prod_{n=0}^{\infty} f^*(f^n(x)). \tag{1}$$

Almost all terms $f^*(f^n(x))$ equal 1. Hence, by the chain-rule, if N is chosen to be sufficiently large then the image of x in (1) is equal to $(f^N)^*(x)$. The value of the product remains unchanged if we extend to $n = -\infty$ because $f^*(x) = 1$ for all $x \in (0, x_f)$. Let \mathbb{S} denote the topological space [0, 1] with 0 and 1 identified. Let $\psi_f : \mathbb{S} \to \mathbb{R}^+$ be given by the equation

$$\psi_f(s) = \phi_f(\alpha_f^s x_f).$$

We consider the set \mathcal{K} of functions from \mathbb{S} to \mathbb{R}^+ that take the value 1 at all but a finite number of points. We define an equivalence relation on \mathcal{K} such that, for functions u and v in \mathcal{K} , u is equivalent to v if and only if there is a translation $t(x) = x + p \pmod{1}$, $p \in \mathbb{R}$, such that u = vt. Finally, we define β_f to be the equivalence class of ψ_f in \mathcal{K} . That β_f is invariant under conjugation in $\mathrm{PLF}^+([0, 1])$ follows from the next two lemmas.

Date: October 7, 2018.

In both lemmas we use the notation ϕ_f^a to denote the map from [a, f(a)) to \mathbb{R}^+ given by (1). For $a \in (0, x_f)$, we define $\psi_f^a : \mathbb{S} \to \mathbb{R}^+$ by $\psi_f^a(s) = \phi_f^a(\alpha_f^s a)$.

Lemma 1. If $a \in (0, x_f)$ then $\psi_f^a = \psi_f$.

Proof. Let $m = \alpha_f$. For a positive integer n, let $b = f^{-n}(x_f)$. For $s \in \mathbb{S}$,

$$\psi_f^b(s) = \phi_f^b(m^s b) = \phi_f(f^n(m^s b)) = \phi_f(m^s f^n(b)) = \psi_f(s).$$

If $a \in (b, f(b))$ we obtain

 $\psi_f^a(s) = \psi_f(s+t),$

 $t = \log_m(a/b)$, for $s \in \mathbb{S}$.

Lemma 2. If f and g are two members of \mathcal{F} that are conjugate in $PLF^+([0,1])$ then $\beta_f = \beta_g$.

Proof. Suppose that $h \in \text{PLF}^+([0,1])$ and $hfh^{-1} = g$. We work in a small enough neighbourhood U of 0 that f, g, and h are linear in $U \cup f(U) \cup g(U)$. Let $\lambda > 0$ be such that $h(x) = \lambda x$ in U. Choose $a \in U$ and let b = h(a). For $x \in [b, g(b))$ and suitably large integers N we have,

$$(g^{N})^{*}(x) = (hf^{N}h^{-1})^{*}(x) = h^{*}(f^{N}h^{-1}(x))(f^{N})^{*}(h^{-1}(x)(h^{-1})^{*}(x)) = (f^{N})^{*}(x/\lambda)$$

Let $m = \alpha_f$. Using Lemma 1 we obtain,

$$\psi_g(s) = \phi_g^b(m^s b) = \phi_f^a(m^s b/\lambda) = \phi_f^a(m^s a) = \psi_f(s).$$

The remainder of this document is a proof of the following theorem.

Theorem 3. Two maps $f, g \in \mathcal{F}$ are conjugate in $PLF^+([0, 1])$ if and only if $\alpha_f = \alpha_g$ and $\beta_f = \beta_g$.

It remains to show that, for $f, g \in \mathcal{F}$, if $\alpha_f = \alpha_g$ and $\beta_f = \beta_g$ then f and g are conjugate in $\mathrm{PLF}^+([0,1])$. To prove this statement, we repeatedly conjugate by piecewise linear homeomorphisms with a single node. The result of this repeated conjugation will be a *corner function*, that is, a member f of \mathcal{F} for which all nodes of f occur in $[x_f, f(x_f))$. Given $p \in (0,1)$ and $0 < \lambda < 1$, there is a unique $h \in \mathrm{PLF}^+([0,1])$ with $h^*(p) = \lambda$ such that h has only the single node p in (0,1). If z_1, \ldots, z_m are the nodes of f in increasing order, then the nodes of hfh^{-1} are contained within the set $\{h(z_1), \ldots, h(z_m), h(p), hf^{-1}(p)\}$. Choose $p = z_m$ and $\lambda = f^*(z_m)$ then

$$(hfh^{-1})^*(h(p)) = h^*(f(p))f^*(p)(h^{-1})^*(h(p)) = f^*(p)(h^{-1})^*(h(p)) = 1.$$

Thus hfh^{-1} has nodes $h(z_1), \ldots, h(z_{m-1}), hf^{-1}(z_m)$, and $(hfh^{-1})^*$ takes values $f^*(z_1), \ldots, f^*(z_{m-1}), f^*(z_m)$ at these nodes. (Possibly $hf^{-1}(z_m)$ coincides with one of the $h(z_i)$.) We describe conjugation by h as the *elementary conjugation* of f. If f is already a corner function and we apply the elementary conjugation, then the nodes of f are permuted. (The last node of f becomes the first node of hfh^{-1} and all other nodes are shifted one place to the right. The values of the nodes remain unchanged.)

Lemma 4. Each member of \mathcal{F} is conjugate to a corner function.

Proof. For a map $f \in \mathcal{F}$ that is not a corner function, let x_f and y_f be the smallest and largest nodes of f. Let $n \in \mathbb{N}$ be such that $y_f \in [f^n(x_f), f^{n+1}(x_f))$. Elementary conjugation of f yields a function $g = hfh^{-1}$, and the node y_f of f is replaced by a node $hf^{-1}(y_f)$ of g that lies in the interval $[g^{n-1}(x_g), g^n(x_g))$. Therefore repeatedly applying elementary conjugations yields a corner function. \Box

Lemma 5. A corner function is uniquely specified by α and ψ .

Proof. Certainly a corner function f is uniquely specified by α_f and ϕ_f , because $(x_f, \alpha_f x_f)$ is the first turning point in the graph of f, and all subsequent turning points are determined by ϕ_f . If f and g are corner functions with $\alpha_f = \alpha_g$, $\psi_f = \psi_g$, but $x_f < x_g$, then g must coincide with the function $x \mapsto \lambda f(\lambda^{-1}x)$, where $\lambda = x_g/x_f$. This function does not fix 1, which is a contradiction. Therefore $x_f = x_g$. Hence $\phi_f = \phi_g$, hence f = g. \Box

To complete the proof of Theorem 3, suppose that f and g are maps in \mathcal{F} such that $\alpha_f = \alpha_g$ and $\beta_f = \beta_g$. By Lemma 4 we may assume that f and g are corner functions. We apply elementary conjugations to f, cycling the nodes of f, until $\psi_f = \psi_g$. From Lemma 5 we deduce that f = g. This completes the proof of Theorem 3.

References

 Brin, M. G. and Squier, C. C., Presentations, conjugacy, roots, and centralizers in groups of piecewise linear homeomorphisms of the real line, *Comm. Algebra* (10) 29 (2001), 4557–4596.

E-mail address: Ian.Short@nuim.ie