

A collection of simple proofs of Sharkovsky's theorem

Bau-Sen Du
 Institute of Mathematics
 Academia Sinica
 Taipei 11529, Taiwan
 dubs@math.sinica.edu.tw

Abstract

This is a revision of the previous paper "More simple proofs of Sharkovsky's theorem". New proofs are included, details of the proofs are provided, and references are expanded.

1 Introduction

Throughout this note, I is a compact interval, and $f : I \rightarrow I$ is a continuous map. For each integer $n \geq 1$, let f^n be defined by: $f^1 = f$ and $f^n = f \circ f^{n-1}$ when $n \geq 2$. For x_0 in I , we call x_0 a periodic point of f with least period m (or a period- m point of f) if $f^m(x_0) = x_0$ and $f^i(x_0) \neq x_0$ when $0 < i < m$. If $f(x_0) = x_0$, then we call x_0 a fixed point of f .

For discrete dynamical systems defined by iterated interval maps, one of the most remarkable results is Sharkovsky's theorem which states as follows:

Theorem (Sharkovsky [23, 26]). Let the Sharkovsky's ordering of the natural numbers be defined as follows:

$$3 \quad 5 \quad 7 \quad 9 \quad \dots \quad 2 \quad 3 \quad 2 \quad 5 \quad 2 \quad 7 \quad 2 \quad 9 \quad 2^2 \quad 3 \quad 2^2 \quad 25 \quad 2^2 \quad 7 \quad 2^2 \quad 9$$

$$\dots \quad 3^2 \quad 2 \cdot 2^2 \quad 2 \quad 1:$$

Then the following three statements hold:

- (1) Assume that $f : I \rightarrow I$ is a continuous map. If f has a period- m point and if $m \prec n$, then f also has a period- n point.

- (2) For each positive integer n there exists a continuous map $f : I \rightarrow I$ that has a period- n point but has no period- m point for any m with $m \mid n$.
- (3) There exists a continuous map $f : I \rightarrow I$ that has a period- 2^i point for $i = 0, 1, 2, \dots$ but has no periodic point of any other period.

There have been a number of different proofs [1–5, 7–14, 16–31] of it in the past 30 years providing various viewpoints of this beautiful theorem. Especially, some of these proofs are excellent for the discrete dynamical systems course. So, why do we need another proof? Sharkovsky's theorem is well-known for its simplicity in assumptions and yet abundance in conclusions. Furthermore, what makes it more appealing is that its proof uses only the intermediate value theorem (in an indirect way through, say, directed graph arguments). The idea was clear except that the details were a little messy [23]. Even the seemingly simplest case that if f has a period- n point with $n \geq 3$ then f has a period-2 point cannot be explained in a few words [1, 3, 11, 29]. Our aim is to find a proof of this theorem which uses the intermediate value theorem in a more intuitive and direct way (see section 3) so that this beautiful result can be someday introduced in the calculus curriculum.

It is well-known [13, 14, 29] that (1) is equivalent to the following three statements:

- (a) if f has a period- m point with $m \geq 3$, then f has a period-2 point;
- (b) if f has a period- m point with $m \geq 3$ and odd, then f has a period- $(m + 2)$ point; and
- (c) if f has a period- m point with $m \geq 3$ and odd, then f has a period-6 point and a period- $(2m)$ point.

These three statements explain clearly why Sharkovsky's ordering is defined as it is. Note that in (c) we don't require the existence of periodic points of all even periods. Only the existence of period-6 and period- $(2m)$ points suffices. In this note, we present several strategies on how to prove (a), (b) and (c). In section 2, we state some preliminary results. In section 3, the proof is based on the existence of a point v such that $f^2(v) < v < f(v)$ and $f(v)$ is in the given period- m orbit. In sections 4 and 5, we do some surgery to the map f and then use Lemma 6 to prove (c). In section 6, we examine how the iterates of a given period- m point jump around a fixed point of f . In section 7, we investigate how the points in a given period- m orbit which lie on one side of a fixed point of f correspond under the action of f . In section 8, we combine the strategies in sections 6 and 7 to find the structure of some periodic orbits. In section 9, we concentrate our attention on the specific point m in P in any given period- m orbit P . In section 10, we based our proof on Lemma 6. The proof in each section is independent of the other. Some proofs of (a), (b), or (c) in these sections can be combined to give various complete proofs of (a), (b) and (c). For completeness, we also include some modifications of the proofs of Sharkovsky's theorem in [13, 14].

2 Preliminary results

To make this paper self-contained, we include the following results and some of their proofs.

Lemma 1. If $f^m(x_0) = x_0$, then the least period of x_0 under f divides m .

Lemma 2. If J is a closed subinterval of I such that $f(I) \subset J$, then f has a fixed point in J .

Lemma 3. Let k, m, n , and s be positive integers. Then the following statements hold:

- (1) If y is a periodic point of f with least period m , then it is a periodic point of f^n with least period $m = (m, n)$, where (m, n) is the greatest common divisor of m and n .
- (2) If y is a periodic point of f^n with least period k , then it is a periodic point of f with least period $kn = s$, where s divides n and is relatively prime to k .

Proof. (1) Let t denote the least period of x_0 under f^n . Then m divides nt since $x_0 = (f^n)^t(x_0) = f^{nt}(x_0)$. Consequently, $\frac{m}{(m, n)}$ divides $\frac{n}{(m, n)}t$. Since $\frac{m}{(m, n)}$ and $\frac{n}{(m, n)}$ are coprime, $\frac{m}{(m, n)}$ divides t . On the other hand, $(f^n)^{(m/(m, n))}(x_0) = (f^m)^{(n/(m, n))}(x_0) = x_0$. So, t divides $\frac{m}{(m, n)}$. This shows that $t = \frac{m}{(m, n)}$.

(2) Since $x_0 = (f^n)^k(x_0) = f^{kn}(x_0)$, the least period of x_0 under f is $\frac{kn}{s}$ for some positive integer s . By (1), $(\frac{kn}{s}) = ((\frac{kn}{s}), n) = k$. So, $\frac{n}{s} = ((\frac{n}{s})k, n) = ((\frac{n}{s})k, (\frac{n}{s})s) = (\frac{n}{s})(k, s)$. This shows that s divides n and $(s, k) = 1$.

Lemma 4. Let J and L be closed subintervals of I with $f(J) \subset L$. Then there exists a closed subinterval K of J such that $f(K) = L$.

Proof. Let $L = [a; b]$. Then since $f(J) \subset L \subset f(J)$, there are two points p and q in J such that $f(p) = a$ and $f(q) = b$. If $p < q$, let $c = \max\{p, x : f(x) = a\}$ and let $d = \min\{c, x : f(x) = b\}$. If $p > q$, let $c = \max\{q, x : f(x) = b\}$ and let $d = \min\{c, x : f(x) = a\}$. In any case, let $K = [c; d]$. It is clear that $f(K) = L$.

If there are closed subintervals $J_0, J_1, \dots, J_{n-1}, J_n$ of I with $J_n = J_0$ such that $f(J_i) \subset J_{i+1}$ for $i = 0; 1; \dots; n-1$, then we say that $\bigcup_{i=0}^{n-1} J_i$ is a cycle of length n . We require the following result.

Lemma 5. If $J_0, J_1, J_2, \dots, J_{n-1}, J_n = J_0$ is a cycle of length n , then there exists a periodic point y of f such that $f^i(y)$ belongs to J_i for $i = 0; 1; \dots; n-1$ and $f^n(y) = y$.

Proof. Let $Q_n = J_0$. Since $f(J_{n-1}) \subset J_0 = Q_n$, there is, by Lemma 4, a closed subinterval Q_{n-1} of J_{n-1} such that $f(Q_{n-1}) = Q_n = J_0$. Continuing this process one by one, we obtain, for each $0 \leq i \leq n-1$, a closed subinterval Q_i of J_i such that $f(Q_i) = Q_{i+1}$. Consequently, $f^i(Q_0) = Q_i$ for all $0 \leq i \leq n$. In particular, $f^n(Q_0) = Q_n = J_0 = Q_0$. By Lemma 2,

there is a point y in $Q_0 \cap J_0$ such that $f^n(y) = y$. Since $y \notin Q_0$, we also obtain that $f^i(y) \notin f^i(Q_0) = Q_i \cap J_i$ for all $0 < i < n - 1$.

We call f turbulent if there exist two closed subintervals I_0 and I_1 of I with at most one point in common such that $f(I_0) \setminus f(I_1) \subset I_0 \cap I_1$ and call f strictly turbulent if such I_0 and I_1 are disjoint.

Lemma 6. If there exist a point c , a fixed point z of f , and an integer $k \geq 2$ such that $f(c) < c < z \leq f^k(c)$ ($f(c) < c < z < f^k(c)$ respectively), then there exist a point d , a fixed point \hat{z} of f such that $f(d) < d < \hat{z} \leq f^2(d)$ ($f(d) < d < \hat{z} < f^2(d)$ respectively) and f has periodic points of all periods and is turbulent (strictly turbulent respectively).

Proof. If $f(c) < c < z \leq f^2(c)$, by taking $I_0 = [f(c); c]$ and $I_1 = [c; z]$, we obtain that f is turbulent. If $f(c) < c < z < f^2(c)$, then there is a point r such that $f(c) < f(r) < c < r < z < f^2(r) \leq f^2(c)$. By taking $I_0 = [f(r); c]$ and $I_1 = [r; z]$, we see that f is strictly turbulent. In either case, the cycle $I_0(I_1)^{i-1}I_0$ of length $i - 1$ gives a period- i point of f . Now assume that the lemma is true respectively for $k = n - 2$ and let $k = n + 1$. If $c < f^2(c) < z$ and $f^2(c) < f^3(c)$, let z be a fixed point of f in $(c; f^2(c))$. If $z < f^2(c)$, let $z = z$. In either case, we have $f(c) < c < z < f^2(c)$ and we are done as before. If $f^2(c) < f(c)$, let $u = f(c)$. If $f(c) < f^2(c) < c < z$, let u be a point in $(c; z)$ such that $f(u) = f^2(c)$. If $f(c) < c < f^2(c) < z$ and $f^3(c) < f^2(c)$, let u be a point in $(f^2(c); z)$ such that $f(u) = f^2(c)$. Then, in either case, we have $f(u) < u < z \leq f^{n+1}(u) = f^n(u)$ (or $f(u) < u < z < f^{n+1}(u) = f^n(u)$ respectively) and the lemma follows from induction hypothesis. This completes the proof.

3 A non-directed graph proof of (a), (b) and (c)

The proof we present here is slightly different from the one in [14].

Let P be a period- m orbit of f with $m \geq 3$. Let v be a point such that $m \in P$, $f^2(v) < v < f(v) = b \in P$. Let z be a fixed point of f in $[v; b]$. Since $f^2(m \in P) > m \in P$ and $f^2(v) < v$, the point $y = \max\{m \in P : x \leq v : f^2(x) = x\}$ exists. Furthermore, $f(x) > z$ on $[y; v]$ and $f^2(x) < x$ on $(y; v]$. Therefore, y is a period-2 point of f . (a) is proved.

For the proofs of (b) and (c), assume that $m \geq 3$ is odd and note that $f(x) > z > x > f^2(x)$ on $(y; v]$. Since $f^{m+2}(y) = f(y) > y$ and $f^{m+2}(v) = f^2(v) = f(b) < v$, the point $p_{m+2} = \min\{x \leq v : f^{m+2}(x) = x\}$ exists. Let k denote the least period of p_{m+2} with respect to f . Then k divides $m + 2$ and so k is odd. Furthermore, $k > 1$ because f has no fixed points in $(y; v)$. If $k < m + 2$, let $x_k = p_{m+2}$, then x_k is a solution of the equation $f^k(x) = x$ in $(y; v)$. Since $f^{k+2}(y) = f(y) > y$ and $f^{k+2}(x_k) = f^2(f^k(x_k)) = f^2(x_k) < x_k$, the equation $f^{k+2}(x) = x$ has a solution x_{k+2} in $(y; x_k)$. Inductively, for each $n \geq 1$, the equation $f^{k+2n}(x) = x$ has a solution x_{k+2n} such that $y < x_{k+2} < x_{k+4} < \dots < x_{k+2n} < x_k < v$. Consequently, the equation $f^{m+2}(x) = x$ has a solution x_{m+2} such that $y < x_{m+2} < x_k = p_{m+2}$. This

contradicts the minimality of p_{m+2} . So, p_{m+2} is a period- $(m+2)$ point of f . This proves (b).

We now prove (c). Let $z_0 = \min\{x : v < x < z, f^2(x) = xg\}$. Then $f^2(x) < x$ and $f(x) > z$ when $y < x < z_0$. If $f^2(x) < z_0$ whenever $m \in I$, $x < z_0$, then $f^2(\min\{I; z_0\}) \in \{m \in I; z_0\}$, which contradicts the fact that $(f^2)^{(m+1)/2}(v) = b > z_0$. So, the point $d = \max\{x : m \in I, x < y; f^2(x) = z_0g\}$ exists and $f(x) > z > z_0 > f^2(x)$ for all x in $(d; y)$. Therefore $f(x) > z > z_0 > f^2(x)$ whenever $d < x < z_0$. Let $s = \min\{f^2(x) : d < x < z_0g\}$. If $s = d$, then $f^2(\{d; z_0\}) \subset \{d; z_0\}$, which again contradicts the fact that $(f^2)^{(m+1)/2}(v) = b > z_0$. Thus $s < d$. Let $u = \min\{d < x < z_0 : f^2(x) = dg\}$. Since $f^2(d) = z_0 > d$ and $f^2(u) = d < u$, the point $c_2 = \min\{d < x < u : f^2(x) = xg\}$ (y) exists and $d < f^2(x) < z_0$ on $(d; c_2]$. Let w be a point in $(d; c_2)$ such that $f^2(w) = u$. Then since $f^4(d) = z_0 > d$ and $f^4(w) = d < w$, the point $c_4 = \min\{d < x < w : f^4(x) = xg\}$ ($< c_2$) exists and $d < f^4(x) < z_0$ on $(d; c_4]$. Inductively, for each $n \geq 1$, let $c_{2n+2} = \min\{d < x < c_{2n} : f^{2n+2}(x) = xg\}$. Then $d < c_{2n+2} < c_{2n} < \dots < c_2 < y$ and $d < (f^2)^k(x) < z_0$ on $(d; c_{2k}]$ for all $1 \leq k \leq n$. Since $f(x) > z_0$ on $(d; z_0)$, we have $f^i(c_{2n}) < z_0 < f^j(c_{2n})$ for all even $2 \leq i \leq 2n$ and all odd $1 \leq j \leq 2n-1$. So, each c_{2n} is a period- $(2n)$ point of f . Therefore, f has periodic points of all even periods. This establishes (c).

Remarks. (1) In the proof of (a), there are many ways to choose such point v . One is: Let $a = \max\{x \in P : f(x) > xg\}$ and $b = \min\{x \in P : a < xg\}$. Then $f(b) = a < b = f(a)$. We can choose v to be a point in $[a; b]$ such that $f(v) = b$. The other is: Let b be the point in P such that $f(b) = \min P$ and let v be a point in $[\min P; b]$ such that $f(v) = b$ (see section 9).

(2) The arguments in the proofs of (a) and (b) can be used to give a simpler proof of the main result of Block in [6] on the stability of periodic orbits in Sharkovsky's theorem. Indeed, assume that f has a period- 2^n point. Let $F = f^{2^n}$. Then F has a period-4 orbit P . Arguing as in the proof of (a), there exists a point v such that $\min P < F^2(v) < v < F(v)$ and $F^2(\min P) > \min P$. So, there is an open neighborhood N of f in $C^0(I; I)$ such that, for each g in N , the map $G = g^{2^n}$ satisfies $G^2(v) < v < G(v)$ and $G^2(\min P) > \min P$. Thus, the point $y = \max\{\min P < x < v : G^2(x) = xg\}$ is a period-2 point of G . By Lemma 3, y is a period- 2^{n-1} point of g . On the other hand, assume that f has a period- 2^i point with $m \geq 3$ and odd and $i \geq 0$. Let $F = f^{2^i}$. Then F has period- m points. Arguing as in the proof of (b), there exist a period-2 point y of F and a point v with $y < v$ such that $F^{m+2}(y) > y$ and $F^{m+2}(v) < v$ and F has no fixed points in $[y; v]$. So, there is an open neighborhood N of f in $C^0(I; I)$ such that, for each g in N , the map $G = g^{2^i}$ satisfies that $G^{m+2}(y) > y$, $G^{m+2}(v) < v$, and G has no fixed points in $[y; v]$. By Sharkovsky's theorem, g has period- $(2^i(m+2))$ points.

(3) In proving (c), we implicitly use the fact that $f^2(v) < v < z < f(v) = b = f^{m+1}(v) = (f^2)^{(m+1)/2}(v)$. Surprisingly, these inequalities imply, by Lemma 6, the existence of periodic points of all periods for f^2 . However, the existence of periodic points of all periods for f^2 does not automatically guarantee the existence of periodic points of all even periods for f . It only guarantees the existence of periodic points of f with least period $2k$ for each even $k \geq 2$ and least period k or $2k$ for each odd $k \geq 1$. We need a little more work to ensure the existence of periodic points of all even periods for f as we have done in the previous

section and in [14]. In the next two sections, we present two different strategies by doing some suitable "surgery" to the map f to achieve our goal. These two proofs of (c) can be combined with the proofs of (a) and (b) in the previous section to give more non-directed graph proofs of (a), (b) and (c).

In the following two sections, we shall let $m \geq 3$ be odd and let a and b be any two adjacent points in P (so that $(a;b) \setminus P = \emptyset$) such that $f(b) < a < b < f(a)$. We also let v be a point in $(a;b)$ such that $f(v) = b$.

4 The first 'surgical' proof of (c)

Let $z_1 < z_2$ be the smallest and largest fixed points of f in $(v;b)$ respectively. Let g be the continuous map on I defined by $g(x) = \max\{f(x); z_2\}$ if $x < z_1$; $g(x) = \min\{f(x); z_1\}$ if $x > z_2$; and $g(x) = x + z_1 + z_2$ if $z_1 < x < z_2$. Then $g([m \text{ in } I; z_1]) \subset [z_2; \max I]$ and $g([z_2; \max I]) \subset [m \text{ in } I; z_1]$ and $g^2(x) = x$ on $[z_1; z_2]$. So, g has no periodic points of any odd periods ≥ 3 . On the other hand, it follows from the definition of g that if the iterates $t; g(t); g^2(t); \dots; g^k(t)$ or the iterates $t; f(t); f^2(t); \dots; f^k(t)$ of some point t are jumping alternatively between $[m \text{ in } I; z_1)$ and $(z_2; \max I]$, then $g^i(t) = f^i(t)$ for all $1 \leq i \leq k$.

Since $m \geq 3$ is odd, for some $1 \leq i \leq m-1$, both $f^i(b)$ and $f^{i+1}(b)$ lie on the same side of $[z_1; z_2]$. Let k be the smallest among these i 's. Then the iterates $b; f(b); f^2(b); \dots; f^k(b)$ are jumping alternatively between $[m \text{ in } I; z_1)$ and $(z_2; \max I]$ and since $f(v) = b$, so are the iterates $v; f(v); f^2(v); \dots; f^k(v)$. Consequently, $g^i(v) = f^i(v)$ for all $0 \leq i \leq k+1$. If k is odd, then $f^k(b) < z_1$ and $f^{k+1}(b) > z_2$ and so $g^{k+1}(b) = z_2$ and $g^2(v) = f^2(v) = f(b) < v < z_1 = g^{k+3}(v)$. If k is even, then $f^k(b) > z_2$ and $f^{k+1}(b) < z_1$ and so $g^{k+1}(b) = z_1$ and $g^2(v) = f^2(v) = f(b) < v < z_1 = g^{k+2}(v)$. In either case, we have $g^2(v) < v < z_1 = (g^2)^n(v)$ for some $n \geq 2$. By Lemma 6, g^2 has periodic points of all periods. So, g^2 has period- j points for all odd $j \geq 3$. By Lemma 3, g has either period- j points or period- $(2j)$ points for any odd $j \geq 3$. Since g has no periodic points of any odd periods ≥ 3 , g has period- $(2j)$ points for all odd $j \geq 3$ which are also periodic points of f with the same periods. This proves (c).

5 The second 'surgical' proof of (c)

We first consider the special case when P is a period- m orbit of f such that $(m \text{ in } P; \max P)$ contains no period- m orbits of f . By (b), $[m \text{ in } P; \max P]$ contains a period- $(m+2)$ orbit Q of f . Let g be the continuous map from I into itself defined by $g(x) = \min Q$ if $f(x) \in m \text{ in } Q$; $g(x) = \max Q$ if $f(x) \in \max Q$; and $g(x) = f(x)$ elsewhere. Then g has no period- m points and by (b) g has no period- j points for any odd $3 \leq j \leq m-2$. Since g has the period- $(m+2)$ orbit Q , there exist a fixed point z of g and a point v such that $g(v) \in 2Q$ and $g^2(v) < v < z < g(v) = (g^2)^{(m+3)/2}(v)$. By Lemma 6, g^2 has periodic points of all periods.

Since g has no period- j points for any odd $3 \leq j \leq m-2$, g has period- $(2j)$ points for all odd $3 \leq j \leq m$ which are also periodic points of f with the same periods. Thus, (c) is proved in this special case. The rest of the proof is to reduce the general case to this special one.

For the general case, we may assume that $I = [m \text{ in } P; \text{max } P]$ and v is the largest point in $[m \text{ in } P; b]$ such that $f(v) = b$. By Lemma 6, we may also assume that f has no fixed points in $[m \text{ in } P; v] \cup [b; \text{max } P]$. Let $z_1 \leq z_2$ be the smallest and largest fixed points of f in $[v; b]$ respectively. Without loss of generality, we may assume that $f(x) = x$ for all $z_1 \leq x \leq z_2$. If f has a period-2 point \hat{y} in $[v; z_1]$, then $z_2 < f(\hat{y}) < b$. In this case, let $t = \hat{y}$ and $u = f(\hat{y})$ and let h be the continuous map from I into itself defined by $h(x) = f(x)$ for x not in $[t; u]$ and $h(x) = x + t + u$ for x in $[t; u]$. Otherwise, let $t = v$ and $u = z_2$ and let $h(x) = f(x)$ for x in I . Then it is clear that if S is a period- k orbit of h with $k \geq 3$, then it is also a period- k orbit of f . For each period- k orbit S of h with $k \geq 3$, since $[m \text{ in } S; \text{max } S]$ must contain a fixed point and a period-2 point of h , we have $[m \text{ in } S; \text{max } S] \subset [t; u]$. Let $w = \min_{m \text{ in } C} C$: C is a period- m orbit of hg . Then by Lemma 1 w is a periodic point of h whose least period divides m . It is obvious that w cannot be a fixed point of h . If the least period of w is $< m$, let W denote the orbit of w under h . Then by (b), $(m \text{ in } W; \text{max } W)$ contains a period- m orbit of h which contradicts the minimality of w . Therefore, W is a period- m orbit of h such that $(m \text{ in } W; \text{max } W)$ contains no period- m orbits of h . It follows from what we have just proved above that h has period- $(2j)$ points for all odd $3 \leq j \leq m$ which are also periodic points of f with the same periods. (c) is proved.

In the following, we present three different directed graph proofs of (a), (b) and (c).

6 The first directed graph proof of (a), (b) and (c)

The proof we present here is different from the one in [13].

Let $P = \{x_i : 1 \leq i \leq m\}$, with $x_1 < x_2 < \dots < x_m$ be a period- m orbit of f with $m \geq 3$. Let $x_s = \max\{x \in P : f(x) > x\}$. Then $f(x_s) = x_{s+1}$ and $f(x_{s+1}) = x_s$. So, f has a fixed point z in $[x_s; x_{s+1}]$. If for all integers i such that $1 \leq i \leq m-1$ the points $f^i(x_s)$ and $f^{i+1}(x_s)$ lie on opposite sides of z , then $f([x_1; x_s] \setminus P) \subset [x_{s+1}; x_m] \setminus P$ and $f([x_{s+1}; x_m] \setminus P) \subset [x_1; x_s] \setminus P$. Since f is one-to-one on P , this implies that $f([x_1; x_s] \setminus P) = [x_{s+1}; x_m] \setminus P$ and $f([x_{s+1}; x_m] \setminus P) = [x_1; x_s] \setminus P$. In particular, $f([x_1; x_s]) \subset [x_{s+1}; x_m]$ and $f([x_{s+1}; x_m]) \subset [x_1; x_s]$. By applying Lemma 5 to the cycle $[x_1; x_s][x_{s+1}; x_m][x_1; x_s]$, we obtain a period-2 point of f .

Now assume that for some $1 \leq r \leq m-1$ the points $f^r(x_s)$ and $f^{r+1}(x_s)$ lie on the same side of z (this includes the case when m is odd). We may also assume that r is the smallest such integer (and so the iterates $x_s; f(x_s); f^2(x_s); \dots; f^r(x_s)$ are jumping around z alternatively). Suppose they both lie on the right side of z (if they both lie on the left side of z , then we consider the iterates of the point x_{s+1} instead of the point x_s and the proof is similar). Then $r \leq m-2$ and since $f(x_{s+1}) = x_s < x_{s+1}$, we have $f^r(x_s) > x_{s+1}$. Let k be

the smallest positive integer such that $f^k(x_s) = f^r(x_s)$. Then $1 \leq k \leq r$. If $k = 1$, then for each $n \geq 2$, by appealing to Lemma 5 for the cycle

$$[x_{s+1}; f^r(x_s)]([x_s; x_{s+1}])^{n-1} [x_{s+1}; f^r(x_s)]$$

of length n (here $([x_s; x_{s+1}])^{n-1}$ represents $n-1$ copies of $[x_s; x_{s+1}]$), we see that f has periodic points of all periods ≥ 2 . So, suppose $k \geq 2$. Since the iterates $x_s; f(x_s); f^2(x_s); \dots; f^{k-1}(x_s)$ are jumping around z alternatively, we actually have $k \geq 3$. So, $f^{k-1}(x_s) < z < f^{k-2}(x_s) < f^r(x_s) = f^k(x_s)$. Choose points $w; v; u$ with $f^{k-1}(x_s) < v < z < u < f^{k-2}(x_s) < w < f^r(x_s)$ such that $f(w) = z, f(v) = w$ and $f(u) = v$. For each even integer $n \geq 2$, we apply Lemma 5 to the cycle

$$[u; w][v; z]([z; u][v; z])^{\frac{n-2}{2}} [u; w]$$

of length n to establish the existence of periodic points of all even periods, including period-2, for f . This proves (a) and (c).

Finally, let $f^j(x_s)$ be the unique point in $ff^i(x_s) : 1 \leq i \leq k-2$ and i is odd g which is closest to the point $f^r(x_s)$. Then $1 \leq j \leq k-2$ and $x_{s+1} = f^{k-2}(x_s) = f^j(x_s) < f^r(x_s)$. Let $L = [f^j(x_s); f^r(x_s)]$ and let $[a; b]$ denote the closed interval with a and b as endpoints. For each integer $n \geq m+1$, we apply Lemma 5 to the cycle

$$L([x_s; x_{s+1}])^{n-j-3} [x_s; z][f(x_s); z][f^2(x_s); z] \dots [f^j(x_s); z][z; f^{k-2}(x_s)][f^{k-1}(x_s); z]L$$

of length n to obtain a period- n point of f (note that when $m+4$ is even, this means that f has periodic points of odd period $m+1$). This proves (b) and hence completes the proof of (a), (b) and (c).

7 The second directed graph proof of (a), (b) and (c)

The proof we present here is a variant of that in [13] except the proof of (a).

Let $P = \{x_i : 1 \leq i \leq m\}$, with $x_1 < x_2 < \dots < x_m$, be a period- m orbit of f with $m \geq 3$. Let $x_s = \max\{x \in P : f(x) > x\}$. Then $f(x_s) = x_{s+1}$ and $f(x_{s+1}) = x_s$. So, f has a fixed point z in $[x_s; x_{s+1}]$. If for all integers i such that $1 \leq i \leq m-1$ and $i \notin s$ the points $f(x_i)$ and $f(x_{i+1})$ lie on the same side of z , then $f([x_1; x_s] \setminus P) \subseteq [x_{s+1}; x_m] \setminus P$ and $f([x_{s+1}; x_m] \setminus P) \subseteq [x_1; x_s] \setminus P$. Arguing as in section 6, we obtain a period-2 point of f .

On the other hand, if for some integer t such that $1 \leq t \leq m-1$ and $t \notin s$ the points $f(x_t)$ and $f(x_{t+1})$ lie on opposite sides of z (this includes the case when m is odd), then $f([x_t; x_{t+1}]) \subseteq [x_s; x_{s+1}]$. For simplicity, we assume that $x_t < x_s$. If $x_{s+1} = x_t$, the proof is similar.

Let q be the smallest positive integer such that $f^q(x_s) = x_t$. Then $2 \leq q \leq m-1$. For $i = 0; 1; \dots; q-1$, let $t_i = [z; f^i(x_s)]$. For each $n \geq m+1$, we appeal to Lemma 5 for

the cycle $J_0 J_1 \dots J_{q-1} J [x_t; x_{t+1}] ([x_s; x_{s+1}])^{n-q-1} J_0$ of length n to confirm the existence of a period- n point of f . This proves (b).

On the other hand, since $x_t < x_s$ and $f(x_s) = x_{s+1}$, we may also assume that t is the largest integer in $[1; s-1]$ such that $f(x_t) = x_s$. So, $f(x_i) = x_{s+1}$ for all $t+1 \leq i \leq s$. If there is an integer k in $[s+1; m-1]$ such that $f(x_{k+1}) = x_{s+1}$, let r be the smallest such integer k , otherwise, let $r = m$. Then $f(x_i) = x_s$ for all $s+1 \leq i \leq r$. Let $f(x \cdot) = m \text{ inf } f(x_i) : s+1 \leq i \leq r$, where $s+1 \leq \cdot \leq r$. Then $f(x \cdot) = x_s$. We have two cases to consider:

Case 1. $x_{t+1} = f(x \cdot) = x_s$. In this case, if $x_{s+1} = f(x) = x_r$ for all x in $[f(x \cdot); x_s] \setminus P$, then $f(x \cdot) = f^n(x_s) = x_r$ for all $n \geq 1$, contradicting the fact that $f^i(x_s) = x_t$ for some $1 \leq i \leq m-1$. So, $s+1 \leq r \leq m-1$ and for some x_j in $[f(x \cdot); x_s] \setminus P$, $f(x_j) = x_{r+1}$. Let w, v and u be points with $f(x \cdot) = x_j < v < z < u < x \cdot$, $x_r < w < f(x_j)$ such that $f(w) = z, f(v) = w$ and $f(u) = v$.

Case 2. $f(x \cdot) = x_t$. In this case, let k be the smallest integer in $[s+1; \cdot]$ such that $f(x_k) = x_t$. Then $f(x_i) = x_{t+1}$ for all $s+1 \leq i \leq k-1$. If $f(x_i) = x_{k-1}$ for all $t+1 \leq i \leq s$, then $x_{t+1} = f^n(x_s) = x_{k-1}$ for all $n \geq 1$. This is again a contradiction. So, for some $t+1 \leq j \leq s$, $f(x_j) = x_k$. Let w, v and u be points with $x_t < w < x_{t+1} < x_j < u < z < v < x_k$ such that $f(w) = z, f(v) = w$ and $f(u) = v$.

In either case, for each even integer $n \geq 2$, we apply Lemma 5 to the cycle

$$[u : w][v : z]([z : u][v : z])^{\frac{n-2}{2}} [u : w]$$

of length n to establish the existence of periodic points of all even periods, including period-2, for f . This proves (a) and (c) and hence completes the proofs of (a), (b) and (c).

8 The third directed graph proof of (a), (b) and (c)

The proof we present here follows the line of the usual proof [1] which involves periodic orbits of some special types called Stefan orbits. However, our argument is different and is interesting in its own right. Let P be a period- m orbit of f with $m \geq 3$ and odd. If there is a point $p \in P$ such that either

$$f^{m-2}(p) < \dots < f(p) < p = f^m(p) < f^2(p) < f^4(p) < \dots < f^{\frac{m-1}{2}}(p)$$

or

$$f^{m-1}(p) < \dots < f^2(p) < p = f^m(p) < f(p) < f^3(p) < \dots < f^{\frac{m-1}{2}}(p);$$

then we say that P is a Stefan orbit of f with least period m .

Let $P = \{x_i : 1 \leq i \leq m\}$, with $x_1 < x_2 < \dots < x_m$ be a period- m orbit of f with $m \geq 3$. Let $x_s = \max\{x \in P : f(x) > x\}$. Then $f(x_s) = x_{s+1}$ and $f(x_{s+1}) = x_s$ and so f has

a fixed point z in $[x_s; x_{s+1}]$. If for all $1 \leq i \leq m-1$, the points $f^i(x_s)$ and $f^{i+1}(x_s)$ lie on opposite sides of z , then arguing as in section 6, we obtain a period-2 point of f . Now assume that there is a smallest integer r in $[1; m-1]$ such that the points $f^r(x_s)$ and $f^{r+1}(x_s)$ lie on the same side of z . If r is even, let J_0 be the smallest closed interval which contains $f^i(x_s)$ for all even i in $[0; r]$ and let J_1 be the smallest closed interval which contains x_s and $f^j(x_s)$ for all odd j in $[1; r-1]$. If r is odd, let J_0 be the smallest closed interval which contains $f^i(x_s)$ for all even i in $[0; r-1]$ and let J_1 be the smallest closed interval which contains $f^j(x_s)$ for all odd j in $[1; r]$. By applying Lemma 5 to the cycle $J_0J_1J_0$, we obtain a period-2 point of f . This proves (a).

In the following, we show that if $m \geq 3$ is odd and f has no periodic points of odd period with $1 < p < m$, then f has periodic points of all periods $\leq m$ and all even periods and P is a Stefan orbit (see also [15]). If $m = 3$, the proof is easy. So, suppose $m > 3$. Since m is odd, there is a smallest integer r in $[1; m-1]$ such that the points $f^r(x_s)$ and $f^{r+1}(x_s)$ lie on the same side of z and for some integer t such that $1 \leq t \leq m-1$ and $t \notin s$ the points $f^t(x_s)$ and $f^{t+1}(x_s)$ lie on opposite sides of z (and so $f^t([x_t; x_{t+1}]) \cap [x_s; x_{s+1}] = \emptyset$). For simplicity, we assume that $x_t < x_s$. If $x_{s+1} < x_t$, then we consider the iterates of the point x_{s+1} instead of the point x_s and the proof is similar. Let q be the smallest positive integer such that $f^q(x_s) < x_t$. Then $2 \leq q \leq m-1$. Let $J_i = [f^i(x_s); z]$ for all $0 \leq i \leq q-1$. For each $n \geq 1$, we apply Lemma 5 to the cycle $J_0J_1 \dots J_{q-1}J[x_t; x_{t+1}](J_0)^n$ of length $n+q$ to obtain a period- $(n+q)$ point with $n+q \leq m$. If $2 \leq q \leq m-3$, then by letting $n = m-2-q$, we obtain a period- $(m-2)$ point with $m-2$ odd and $1 < m-2 < m$. This is a contradiction. So, either $q = m-1$ or $q = m-2$.

If $1 \leq r \leq m-3$, then since f maps each of x_s and x_{s+1} to the other side of z , $f^r(x_s) \notin [x_s; x_{s+1}]$. Let $J_i = [f^i(x_s); z]$ for all $0 \leq i \leq r-1$ and $J_r = [f^r(x_s); x_s]$ if $f^r(x_s) < x_s$ and $J_r = [x_{s+1}; f^r(x_s)]$ if $x_{s+1} < f^r(x_s)$. Then we apply Lemma 5 to the cycle $J_0J_1 \dots J_r(J_0)^{m-2-r}$ of odd length $m-2$ to obtain a periodic point of odd period with $1 < p < m$. This is a contradiction. (Note that this argument can also be used to show that if $m \geq 4$ is even and f has no periodic points of odd period with $1 < p < m$, then the iterates $x_s; f(x_s); f^2(x_s); \dots; f^{m-1}(x_s)$ are jumping alternately around a fixed point of f). Therefore, $f^{m-1}(x_s) = x_t < f^i(x_s) < z < f^j(x_s)$ for all even i in $[0; m-3]$ and all odd j in $[1; m-2]$. Let $U = [f^{m-1}(x_s); f^{m-3}(x_s)]$, $V = [f^{m-3}(x_s); z]$ and $W = [z; f^{m-2}(x_s)]$. By applying Lemma 5 to the cycles $U(WV)^iWU, i \geq 0$, we obtain that f has periodic points of all even periods.

For $0 \leq i \leq m-1$, let $J_i = [z; f^i(x_s)]$. If, for some $1 \leq j < k \leq m-1$ with $k-j$ even, we have $[z; f^k(x_s)] \cap [z; f^j(x_s)] = \emptyset$, then by applying Lemma 5 to the cycle $J_0J_1 \dots J_{j-1}J_kJ_{k+1} \dots J_{m-2}[x_t; x_{t+1}]J_0$ of odd length $m-k+j < m$, we obtain a periodic point of f of odd period strictly between 1 and m . This contradicts the assumption. So, we must have

$$f^{m-1}(x_s) < f^2(x_s) < f^2(x_s) < x_s < f(x_s) < f^3(x_s) < f^4(x_s) < f^2(x_s):$$

That is, the orbit of x_s is a Stefan orbit.

Now suppose f has a periodic point of odd period $n \geq 3$. Then there is an odd integer $1 < m \leq n$ such that f has a period- m point and no periodic points of odd periods with

$1 < j < m$. It follows from what we have just proved that f has periodic points of all even periods and for each $j \leq m + 1$, in particular, for each $j \leq n + 1$, f has a period- j point. This completes the proofs of (a), (b) and (c).

9 The fourth directed graph proof of (a), (b) and (c)

These directed-graph proofs of (a), (b) and (c) are different from the one given in [13] and depend heavily on the choice of the particular point m in P in the given period- m orbit P .

Let P be a period- m orbit of f with $m \geq 3$ and let b be a point in P such that $f(b) = m$ in P . If $f(x) < b$ for all m in P $x < b$, then $(m$ in P) $f(m$ in $P) < b$. So, $(m$ in P) $f^2(m$ in $P) = f(f(m$ in $P)) < b$. By induction, $(m$ in P) $f^k(m$ in $P) < b$ for each $k \geq 1$. This contradicts the fact that $b = f^m(b) = f^{m-1}(f(b)) = f^{m-1}(m$ in $P) < b$. Therefore, there exists a point v in $(m$ in $P; b)$ such that $f(v) = b$. Let z be a fixed point of f in $(v; b)$.

If for some point x in $[m$ in $P; v]$, $f(x) \leq z$, then we consider the cycle $[x; v][z; b][x; v]$. If for some x in $[m$ in $P; v]$, $f(x) > z$ and $v < f^2(x)$, then we consider the cycle $[x; v][b : f(x)][x; v]$. If $f^2(x) \leq v < z < f(x)$ for all x in $[m$ in $P; v]$, let $q = \max([m$ in $P; v] \setminus P) (v)$. Then $f^2([m$ in $P; q] \setminus P) \subseteq [m$ in $P; q] \setminus P$. Since f^2 is one-to-one on P , this implies that $f^2([m$ in $P; q] \setminus P) = [m$ in $P; q] \setminus P$ and so $f^2([m$ in $P; q]) \subseteq [m$ in $P; q]$. In this case, we consider the cycle $[m$ in $P; q][f([m$ in $P; q])][m$ in $P; q]$. In either of the above three cases, by Lemma 5, we obtain a period-2 point y of f such that $y < v < z < f(y)$. This proves (a).

Here is the second proof of (a). If there is a fixed point z of f in $[m$ in $P; v]$, then by applying Lemma 5 to the cycle $[v; b][z; v]^{n-1}[v; b]$ of length $n-1$, we obtain periodic points of all periods for f and we are done. So, in the following, we assume that f has no fixed points in $[m$ in $P; v]$. If $f^2(x) \leq v$ for all x in $[m$ in $P; v]$, then in particular, $f^2([m$ in $P; v] \setminus P) \subseteq [m$ in $P; v] \setminus P$. So, if $q = \max([m$ in $P; v] \setminus P) (v)$, then $f^2([m$ in $P; q]) \subseteq [m$ in $P; q]$. On the other hand, if $f^2(x) > v$ for some x in $[m$ in $P; v]$, then $f^2([m$ in $P; v]) \subseteq [m$ in $P; v]$. In either case, by Lemma 2, there is a point y in $[m$ in $P; v]$ such that $f^2(y) = y$. Since f has no fixed points in $[m$ in $P; v]$ and $f(v) = b > v$, y is a period-2 point of f such that $y < v < f(y)$. In this case, we may choose z to be any fixed point of f in $[v; f(y)]$. (a) is proved.

In the following, let $m \geq 3$ be odd. We shall give two different proofs of (c) which are related to each other in some way. However, each is interesting in its own right. Let $z_0 = \min\{x \in b : f^2(x) = x\}$. Then f has neither fixed points nor period-2 points in $[v; z_0)$. Since $f(v) > v$ and $f^2(v) < v$, we have $f(x) > x$ and $f^2(x) < x < z_0$ on $[v; z_0)$. If $f^2(x) < z_0$ for all m in P $x \leq v$, then $f^2(x) < z_0$ for all m in P $x < z_0$. In particular, $(m$ in P) $f^2(m$ in $P) < z_0$ and so $(m$ in P) $f^2(f^2(m$ in $P)) < z_0$. By induction, $(m$ in P) $(f^2)^k(m$ in $P) < z_0$ for each $k \geq 1$. Since $m \geq 3$ is odd, this contradicts the fact that $(f^2)^{(m-1)/2}(m$ in $P) = b > z_0$. Hence $\max\{f^2(x) : m$ in P $x \leq v\} = z_0$. Let $I_0 = [m$ in $P; v]$ and $I_1 = [v; z_0]$. Then $f^2(I_0) \setminus f^2(I_1) \subseteq I_0 \setminus I_1$. For each $n \geq 1$, we apply Lemma 5 to the cycle $I_1(I_0)^n I_1$ (with respect to f^2) of length $n+1$ to obtain a point w in $[v; z_0]$

such that $(f^2)^{n+1}(w) = w$ and $(f^2)^i(w) \notin I_0$ for all $1 \leq i \leq n$. Thus $f^{2i}(w) < v < w < f(w)$ for all $1 \leq i \leq n$. Consequently, w is a period- $(2n+2)$ point of f . Therefore, f has periodic points of all even periods ≥ 4 . This proves (c).

We now give the second proof of (c). For this proof, we choose v , in the beginning of this section, to be the largest point in $[y; z]$ such that $f(v) = b$ and then let $z_0 = \min\{x \in [v; z] : f^2(x) = x\}$. Then $v < z_0 < z$, $f(z_0) < b$ and $f^2(x) < x < z_0$ on $[v; z_0]$. We choose a sequence x_1, x_2, x_3, \dots of points so that $v < x_1 < x_2 < x_3 < \dots < z_0 < f(z_0) < x_4 < x_3 < x_2 < x_1 < b$ and $f(x_k) = x_{k+1}$ for $k = 1, 2, 3, \dots$ with $x_1 = v$. If $f^2(x) < z_0$ whenever $m \in P \cap [x, v]$, then $f^2(x) < z_0$ for all $m \in P \cap [x, z_0]$. In particular, $(m \in P) \implies f^{2k}(m \in P) < z_0$ for all $k \geq 1$, contradicting the fact that $z_0 < b = (f^2)^{(m \in P)/2}(m \in P)$. So, $\max\{f^2(x) : m \in P \cap [x, v] \cap [z_0, b]\} = z_0$. For each even integer $n \geq 2$, we appeal to Lemma 5 for the cycle

$$[m \in P; v]f([m \in P; v])[x_{n+2}; x_n][x_{n+3}; x_{n+1}] \dots [x_2][x_1; b][m \in P; v]$$

of length $n+2$ to confirm the existence of periodic points of all even periods ≥ 4 for f . This also proves (c).

Finally we prove (b). Let $m \geq 3$ be odd. Let j be the largest even integer in $[0; m-1]$ such that $f^j(m \in P) < y$. Then $0 \leq j < m-1$ since $f^{m-1}(m \in P) = b > y$. For $0 \leq i < m-1$, let $J_i = [f^i(m \in P); f^i(y)]$. If there is an odd integer k with $j < k < m-1$ such that $f^k(m \in P) < y$, let k denote the smallest such k . Then $k \geq j+2$ and $y < f^i(m \in P)$ for all $j < i < k$. We apply Lemma 5 to the cycle

$$J_j J_{j+1} J_{j+2} \dots J_{k-1} [v; f(y)] ([v; b])^{n+j-k+1} J_j$$

of length n for each $n \geq m+1$. If $y < f^i(m \in P)$ for all odd integers $j < k < m-1$, we apply Lemma 5 to the cycle

$$J_j J_{j+1} J_{j+2} \dots J_m [v; b]^{n-m+j+1} J_j$$

of length n for each $n \geq m+1$. In either case, for each $n \geq m+1$, we obtain a point p_n in $[m \in P; y]$ such that $f^n(p_n) = p_n$ and $y < f^i(p_n)$ for all $1 \leq i \leq n-1$. Therefore, in either case, p_n is a period- n point of f for each $n \geq m+1$. This establishes (b).

10 The fifth directed graph proof of (a), (b) and (c)

The following proofs of (a) and (c) depend on Lemma 6. Let P be a period- m orbit of f with $m \geq 3$ and let a and b be points in P such that $f(a) = \max P$ and $f(b) = \min P$. If $b < a$, then by Lemma 6, f has periodic points of all periods and we are done. So, suppose $a < b$.

Let z be a fixed point of f in $(a; b)$. Let v be the unique point in P such that $f(v) = b$. If $b < v$, then by applying Lemma 5 to the cycle $[b; v]([a; b])^i [b; v]$, $i \geq 1$, we obtain that f has

periodic points of all periods. 2. If $z < v < b$ or, $m \in P$ $v < z$ and there is a fixed point of f in $[m \in P; v]$, then Lemma 6 implies that f has periodic points of all periods. So, assume that $m \in P < v < z$ and f has no fixed points in $[m \in P; v]$. Then since f^2 is one-to-one on P , we have $f^2([m \in P; v] \setminus P) = [m \in P; v] \setminus P$. In particular, $f^2([m \in P; v]) = [m \in P; v]$. By Lemma 2, there is a point y in $[m \in P; v]$ such that $f^2(y) = y$. Since f has no fixed points in $[m \in P; v]$, y is a period-2 point of f in $[m \in P; v]$ and $f(y) > v$. This proves (a). In the sequel, we assume that y is a period-2 point of f in $[m \in P; v]$ such that $y < v < f(y)$.

We now give a proof of (b) which is different from those given in the previous sections and can be seen as a directed graph version of the proof of (b) in section 3 with the choice of the particular point $m \in P$. Let $m \geq 3$ be odd. Since $f^{m+2}(y) = f(y) > v$ and $f^{m+2}(v) = m \in P < y$, we have $[f^{m+2}(y) : f^{m+2}(v)] \subset [y; v]$. Hence, by applying Lemma 5 to the cycle

$$[y; v][f(y) : f(v)][f^2(y) : f^2(v)][f^3(y) : f^3(v)] \dots [f^m(y) : f^{m+1}(v)][y; v]$$

of length $m + 2$, we obtain a point r in $[y; v]$ such that $f^{m+2}(r) = r$, $f(r) \notin [f(y) : f(v)]$ and $f^2(r) \notin [f^2(y) : f^2(v)] = [m \in P; y]$. If r has least period $m + 2$, we are done. Otherwise, let $n \geq 3$ be the least period of r under f . Then n is odd and $m + 2 \leq n < (m + 2) = 2$. Since $[f^n(y) : f^n(r)] = [r; f(y)] \subset [v; f(y)]$ and $f([v; b]) \subset [v; b]$, we apply Lemma 5 to the cycle

$$[y; r][f(y) : f(r)][f^2(y) : f^2(r)] \dots [f^n(y) : f^{n+1}(r)][v; f(y)][v; b]^{m+1-n} [y; r]$$

of length $m + 2$ to obtain a point s in $[y; r]$ such that $f^{m+2}(s) = s$, $f^2(s) \notin [f^2(y) : f^2(r)]$, $f^n(s) \notin [v; f(y)]$ and $f^j(s) \notin [v; b]$ for all $n + 1 \leq j \leq m + 1$. Since $f^2(r) < y$, $f^2(s) \notin [f^2(y) : f^2(r)] = [f^2(r); y]$. Thus, $f^2(s) < y < s < v < f^j(s)$ for all $n \leq j \leq m + 1$. Since there are at least $(m + 1) - n + 1 \geq (m + 2) = 2$ consecutive iterates of s under f lying to the right of y and $f^2(s) < y$, the least period of s under f is $m + 2$. This establishes (b).

Finally we prove (c). Let $m \geq 3$ be odd. If $m = 3$, then since $f^2(m \in P) = f^2(f(b)) = b$, by applying Lemma 5 to the cycle $[y; v][b : f(y)][m \in P; y][f(m \in P); f(y)][v; z][z; b][y; v]$ of length 6, we obtain a period-6 point of f . If $m \geq 5$ and odd, let $J_i = [f^i(m \in P) : f^i(y)]$; $i = 0; 1; 2; \dots; m - 1$. Then since $[m \in P) : f^m(y)] \subset [y; f(y)]$, we apply Lemma 5 to the cycle

$$[y; v][b : f(y)]J_0J_1J_2 \dots J_{m-1}([y; f(y)])^{m-2} [y; v]$$

of length $2m$ to obtain a point t such that $f^2(t) < y < t = f^{2m}(t) < v < f(t)$ and $f^i(t) \notin [y; f(y)]$ for all $m + 2 \leq i \leq 2m - 1$. Consequently, these $m + 1$ points $t; f(t); f^2(t); f^i(t); m + 2 \leq i \leq 2m - 1$ in the orbit of t are all distinct. So, t is a period- $(2m)$ point of f . Furthermore, since $f^2(v) = m \in P < v < z < b = (f^2)^{(m+1)/2}(v)$, f^2 has, by Lemma 6, a period-3 point. So, f has either a period-6 point or a period-3 point. If f has a period-3 point, then as shown above, f has a period-6 point. This, together with the above, shows that if f has period- m points with $m \geq 3$ and odd then f has period- $(2m)$ and period-6 points. (c) is proved.

For the sake of completeness, we include a proof of Sharkovsky's theorem which is slightly different from those in [13], [14].

11 A proof of Sharkovsky's theorem

If f has period- m points with $m \geq 3$ and odd, then by (b) f has period- $(m+2)$ points and by (c) f has period- $(2m-3)$ points. If f has period- $(2m-3)$ points with $m \geq 3$ and odd, then by Lemma 3, f^2 has period- m points. By (b), f^2 has period- $(m+2)$ points, which implies by Lemma 3 that f has either period- $(m+2)$ points or period- $(2(m+2))$ points. If f has period- $(m+2)$ points, then according to (c) f has period- $(2(m+2))$ points. In either case, f has period- $(2(m+2))$ points. On the other hand, since f^2 has period- m points, by (c) f^2 has period- $(2m-3)$ points, hence by Lemma 3, f has period- $(2m-3)$ points. Now if f has period- $(2^k m)$ points with $m \geq 3$ and odd and if $k \geq 2$, then by Lemma 3, $f^{2^{k-1}}$ has period- $(2m)$ points. It follows from what we have just proved that $f^{2^{k-1}}$ has period- $(2(m+2))$ points and period- (2^{2k-3}) points. So, by Lemma 3, f has period- $(2(m+2))$ points and period- $(2^{k+1}m-3)$ points. Consequently, if f has period- $(2^k m)$ points with $m \geq 3$ and odd and if $i \geq 0$, then by Lemma 3, f^{2^i} has period- m points. For each $i \geq 0$, by Lemma 3, $f^{2^i} = (f^{2^{i-1}})^2$ has period- m points. By (c), f^{2^i} has period-6 points. So, $f^{2^{i+1}}$ has period-3 points and hence has period-2 points. This implies that f has period- 2^{i+2} points for all $i \geq 0$. Finally, if f has period- 2^k points for some $k \geq 2$, then $f^{2^{k-2}}$ has period-4 points and, by (a), has period-2 points. So, f has period- 2^{k-1} points. This proves (1).

For the proofs of (2) and (3), it suffices to assume that $I = [0;1]$. Let $T(x) = 1 - 2|x - 1/2|$ be the tent map on I . Then for each positive integer n the equation $T^n(x) = x$ has exactly 2^n distinct solutions in I . It follows that T has finitely many period- n orbits. Among these period- n orbits, let P_n be one with the smallest diameter $\max P_n = m$ in P_n (see also [1, pp 32-34]). For any x in I , let $T_n(x) = m$ in P_n if $T(x) = m$ in P_n , $T_n(x) = \max P_n$ if $T(x) = \max P_n$, and $T_n(x) = T(x)$ elsewhere. It is then easy to see that T_n has exactly one period- n orbit (i.e., P_n) but has no period- m orbit for any m with $m \leq n$ in the Sharkovsky ordering. Now let Q_3 be the unique period-3 orbit of T of smallest diameter. Then $\{m \in Q_3; \max Q_3\}$ contains finitely many period-6 orbits of T . If Q_6 is one of smallest diameter, then $\{m \in Q_6; \max Q_6\}$ contains finitely many period-12 orbits of T . We choose one, say Q_{12} , of smallest diameter and continue the process inductively. Let $q_0 = \sup\{m \in Q_{2^i-3} : i \geq 0\}$ and $q_1 = \inf\{\max Q_{2^i-3} : i \geq 0\}$. Let $T_1(x) = q_0$ if $T(x) = q_0$, $T_1(x) = q_1$ if $T(x) = q_1$, and $T_1(x) = T(x)$ elsewhere. Then it is easy to see that T_1 has a period- 2^i point for $i = 0;1;2;\dots$ but has no periodic points of any other periods. This completes the proof of Sharkovsky's theorem.

References

- [1] L. Alsedà, J. Llibre and M. Misiurewicz, Combinatorial dynamics and entropy in dimension one, Second edition, Advanced Series in Nonlinear Dynamics, vol. 5, World Scientific, River Edge, NJ, 2000.
- [2] A. Amedo, P. Ferrero and C. Tresser, Sharkovski's order for the appearance of superstable cycles of one-parameter families of simple real maps: an elementary proof,

- Comm. Pure Appl. Math. 37 (1984), 13–17.
- [3] R. Barton and K. Burns, A simple special case of Sharkovskii's theorem, *Amer. Math. Monthly* 107 (2000), 932–933.
- [4] C. Bernhardt, A proof of Sharkovsky's theorem, *J. Diff. Equ. Appl.* 9 (2003), 373–379.
- [5] N. Bhatia and W. Egerland, New proof and extension of Sarkovskii's theorem, *Far East J. Math. Sci. Special volume, Part I*, 1996, 53–67.
- [6] L. Block, Stability of periodic orbits in the theorem of Sarkovskii, *Proc. Amer. Math. Soc.* 81 (1981), 333–336.
- [7] L. Block and W. A. Coppel, Dynamics in one dimension, *Lecture Notes in Mathematics*, Vol. 1513, Springer-Verlag, Berlin, 1992.
- [8] L. Block, J. Guckenheimer, M. Misiurewicz and L.-S. Young, Periodic points and topological entropy of one-dimensional maps, in *Global Theory of Dynamical Systems*, *Lecture Notes in Math.*, no. 819, Springer-Verlag, Berlin, 1980, pp. 18–34.
- [9] U. Burkart, Interval mapping graphs and periodic points of continuous functions, *J. Combin. Theory Ser. B* 32 (1982), 57–68.
- [10] K. Burns and B. Hasselblatt, Sharkovsky's theorem, *Preprint* (2007).
- [11] W. A. Coppel, The solution of equations by iteration, *Proc. Cambridge Philos. Soc.* 51 (1955), 41–43.
- [12] B.-S. Du, The minimal number of periodic orbits of periods guaranteed in Sharkovskii's theorem, *Bull. Austral. Math. Soc.* 31 (1985), 89–103. *Corrigendum*: 32 (1985), 159.
- [13] B.-S. Du, A simple proof of Sharkovsky's theorem, *Amer. Math. Monthly* 111 (2004), 595–599.
- [14] B.-S. Du, A simple proof of Sharkovsky's theorem revisited, *Amer. Math. Monthly* 114 (2007), 152–155.
- [15] B.-S. Du, On the invariance of Li-Yorke chaos of interval maps, *J. Diff. Equ. Appl.* 11 (2005), 823–828.
- [16] S. Elyadi, On a converse of Sharkovsky's theorem, *Amer. Math. Monthly* 103 (1996), 386–392.
- [17] B. Gaweł, On the theorems of Sarkovskii and Stefan on cycles, *Proc. Amer. Math. Soc.* 107 (1989), 125–132.
- [18] J. Guckenheimer, On the bifurcation of maps of the interval, *Invent. Math.* 39 (1977), 165–178.
- [19] C.-W. Ho and C. Morris, A graph-theoretic proof of Sharkovsky's theorem on the periodic points of continuous functions, *Pacific J. Math.* 96 (1981), 361–370.

- [20] X.C.Huang, From intermediate value theorem to chaos, *Math. Mag.* 65 (1992), 91-103.
- [21] H.Kaplan, A cartoon-assisted proof of Sarkovskii's theorem, *Amer. J. Phys.* 55 (1987), 1023-1032.
- [22] J.P.Keener, The Sarkovskii sequence and stable periodic orbits of maps of the interval, *SIAM J. Numer. Anal.* 23 (1986), 976-985.
- [23] M.Misiurewicz, Remarks on Sharkovsky's theorem, *Amer. Math. Monthly* 104 (1997), 846-847.
- [24] M.Osikawa and Y.Oono, Chaos in C^0 endomorphism of interval, *Publ. RIMS Kyoto Univ.* 17 (1981), 165-177.
- [25] S.Patinkin, Stirring our way to Sharkovsky's theorem, *Bull. Austral. Math. Soc.* 56 (1997), 453-458.
- [26] A.N.Sharkovsky, Coexistence of cycles of a continuous map of a line into itself, *Ukrain. Mat. Zh.* 16 (1964) 61-71 (Russian); English translation, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* 5 (1995), 1263-1273.
- [27] A.N.Sharkovsky, S.F.Kolyada, A.G.Sivak and V.V.Fedorenko, Dynamics of one-dimensional maps, Translated from the 1989 Russian original by A.Sivak, P.Malyshv and D.Malyshv and revised by the authors, *Mathematics and its Applications*, vol. 407, Kluwer Academic Publishers Group, Dordrecht, 1997.
- [28] P.Stefan, A theorem of Sarkovskii on the existence of periodic orbits of continuous endomorphisms of the real line, *Comm. Math. Phys.* 54 (1977), 237-248.
- [29] P.D.Stran Jr., Periodic points of continuous functions, *Math. Mag.* 51 (1978), 99-105.
- [30] J.C.Xiong, A simple proof of Sharkovskii's theorem on the existence of periodic points of continuous self-maps of an interval, *J. China Univ. Sci. Tech.* 12 (1982), 17-20 (Chinese).
- [31] J.Z.Zhang and L.Yang, Some theorems on the Sarkovskii order, *Adv. in Math. (Beijing)* 16 (1987), 33-48 (Chinese).