

A PROOF OF THE SMOOTHNESS OF THE FINITE TIME HORIZON AMERICAN PUT OPTION FOR JUMP DIFFUSIONS

ERHAN BAYRAKTAR

ABSTRACT. We give a new proof of the fact that the value function of the finite time horizon American put option for a jump diffusion, when the jumps are from a compound Poisson process, is the classical solution of a free boundary equation. We also show that the value function is C^1 across the optimal stopping boundary. Our proof, which only uses the classical theory of parabolic partial differential equations of [7, 8], is an alternative to the proof that uses the theory of viscosity solutions (see [14]). This new proof relies on constructing a monotonous sequence of functions, each of which is a value function of an optimal stopping problem for a geometric Brownian motion, converging to the value function of the American put option for the jump diffusion uniformly and exponentially fast. This sequence is constructed by iterating a functional operator that maps a certain class of convex functions to classical solutions of corresponding free boundary equations. On the other hand, since the approximating sequence converges to the value function exponentially fast, it naturally leads to a good numerical scheme.

1. INTRODUCTION

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space hosting a Wiener process $W = \{W_t; t \geq 0\}$ and a Poisson random measure N on $\mathbb{R}_+ \times \mathbb{R}_+$, with mean measure $\lambda \nu(dx)dt$ (in which ν is a probability measure on \mathbb{R}_+), independent of the Wiener process. We will consider a Markov process $S = \{S_t; t \geq 0\}$ of the form

$$dS_t = \mu S_t dt + \sigma S_t dW_t + S_{t-} \int_{\mathbb{R}_+} (z - 1) N(dt, dz). \quad (1.1)$$

In this model, if the stock price jumps at time t , then it moves from S_{t-} to $S_t = Z S_{t-}$, in which Z is a positive random variable whose distribution is given by ν . Note that when $Z < 1$ the stock price jumps down and when $Z > 1$ the stock price jumps up. In the Merton jump diffusion model $Z = \exp(Y)$, in which Y is a Gaussian random variable. We will take $\mu = r + \lambda - \lambda \xi$, in which $\xi = \int_{\mathbb{R}_+} xv(dx) < \infty$, so that $(e^{-rt} S_t)_{t \geq 0}$ is a martingale, i.e., \mathbb{P} is a risk neutral measure. The constant $r \geq 0$ is the interest rate, and the constant $\sigma > 0$ is the volatility. We assume the risk neutral pricing measure \mathbb{P} , and hence the parameters of the problem, are fixed as a result of a calibration to historical data. The value function of the American put option pricing problem is

$$V(x, T) := \sup_{\tau \in \tilde{\mathcal{S}}_{0, T}} \mathbb{E}^x \{e^{-r\tau} (K - S_\tau)^+\}, \quad (1.2)$$

in which $\tilde{\mathcal{S}}_{0, T}$ is the set of stopping times (of the filtration generated by W and N) that take values in $[0, T]$, and \mathbb{E}^x is the expectation under the probability measure \mathbb{P} , given that $S_0 = x$.

2000 *Mathematics Subject Classification.* 60G40, 62L15, 60J75.

Key words and phrases. Optimal stopping, Markov Processes, Jump Diffusions, American Options, Integro-Differential Equations, Parabolic Free Boundary Equations.

This research is supported in part by the National Science Foundation.

I would like to thank the two referees and the anonymous AE for their insightful comments. I also would like to express my gratitude to Farid Aitsahlia, Tom Bielecki, Masahiko Egami, Sebastian Jaimungal, Sergei Levendorskii, Mihai Sirbu, Mete Soner, Hao Xing, Virginia R. Young, Thaleia Zariphopoulou and Gordan Zitkovic for their feedback.

We will show that V is the classical solution of a free boundary equation and that it satisfies the *smooth fit principle*, i.e., V is continuously differentiable with respect to its first variable at the optimal stopping boundary. We argue these facts by showing that V is the fixed point of an operator, which we will denote by J , that maps a given function to the value function of an optimal stopping problem for a geometric Brownian motion. This operator acts as a regularizer: As soon as the given function f has some certain regularity properties, we show that Jf is the unique classical solution of a corresponding free boundary equation and that it satisfies the smooth fit principle. The proof of the main result concludes once we show that V has these certain regularity properties. In this last step we make use of a sequence (which is constructed by iterating J starting with the pay-off function of the put option) that converges to V uniformly and exponentially fast. Incidentally, this sequence yields a numerical procedure, whose accuracy versus speed characteristics can be controlled. Each element of this sequence is an optimal stopping problem for geometric Brownian motion and can be readily calculated using classical finite difference methods (see e.g. [18] for the implementation of these methods). An alternative proof of the regularity of V was given in [14]. This proof used a combination of the results in [8] and the theory of viscosity solutions. In particular the proof of Proposition 3.1 in [14] is carried out (details are not provided but hinted) using arguments similar to those used in the proof of Proposition 5.3 in [15]. The latter proof uses the uniqueness results of [9] for viscosity solutions.

The infinite horizon American put option for jump diffusions were analyzed in [3] using the iterative scheme we describe here. The main technical difficulty in the current paper stems from the fact that each element in the approximating sequence solves a parabolic rather than an elliptic problem. In fact, in the infinite horizon case one can obtain a closed form representation for the value function, which is not possible in the finite horizon case. We make use of the results of [8], and Chapter 2 of [10] (also see Chapter 7 of [13]) to study the properties of the approximating sequence. For example, we show that the approximating sequence is bounded with respect to the Hölder semi-norm (see page 61 in [7] for a definition), which is used to argue that the limit of the approximating sequence (which is a fixed point of J) solves a corresponding free boundary equation.

Somewhat similar approximation techniques to the one we employ were used to solve optimal stopping problems for *diffusions*: see e.g. [2] for perpetual optimal stopping problems with non-smooth pay-off functions; and [6], [5] for finite time horizon American put option pricing problems for geometric Brownian motion. On the other hand, [1] and [11] consider the smooth fit principle for the infinite horizon American put option pricing problems for one-dimensional exponential Lévy processes using the fluctuation theory. Also see [4] for the analysis of the smooth fit principle for a multi-dimensional infinite horizon optimal stopping problem.

The next two sections prepare the proof of our main result, Theorems 3.1, in a sequence of lemmata and corollaries. In the next section, we introduce the functional operator J , that maps a given function to the value function of an optimal stopping problem for a geometric Brownian motion. We then analyze the properties of J . For example, J preserves convexity with respect to the first variable; the increase in the Hölder semi-norm after the application of J can be controlled; J maps certain class of functions to the classical solutions of free boundary equations. In Section 3, we construct a sequence of functions that converge to the smallest fixed point of the operator J . We show that the sequence is bounded in the Hölder norm, and satisfies certain regularity properties using results of Section 2. We eventually arrive at the fact that the smallest fixed point of J is equal to V . As a result the regularity properties of V follow.

2. A FUNCTIONAL OPERATOR AND ITS PROPERTIES

Let us define an operator J through its action on a test function $f : \mathbb{R}_+ \times \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}_+$: The operator J takes the function f to the value function of the following optimal stopping problem

$$Jf(x, T) = \sup_{\tau \in \mathcal{S}_{0, T}} \mathbb{E}^x \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot Pf(S_t^0, T-t) dt + e^{-(r+\lambda)\tau} (K - S_\tau^0)^+ \right\}, \quad (2.1)$$

in which

$$Pf(x, T-t) = \int_{\mathbb{R}_+} f(xz, T-t) \nu(dz), \quad x \geq 0. \quad (2.2)$$

We will extend $T \rightarrow Jf(x, T)$ onto $[0, \infty]$ by letting

$$Jf(x, \infty) = \lim_{T \rightarrow \infty} Jf(x, T). \quad (2.3)$$

Here, $S^0 = \{S_t^0; t \geq 0\}$ is the solution of

$$dS_t^0 = \mu S_t^0 dt + \sigma S_t^0 dW_t, \quad S_0^0 = x, \quad (2.4)$$

whose infinitesimal generator is given by

$$\mathcal{A} := \frac{1}{2} \sigma^2 x^2 \frac{d^2}{dx^2} + \mu x \frac{d}{dx}. \quad (2.5)$$

In (2.1), $\mathcal{S}_{[0, T]}$ denotes the set of stopping times of S^0 which take values in $[0, T]$. Note that

$$S_t^0 = xH_t, \quad (2.6)$$

where

$$H_t = \exp \left\{ \left(\mu - \frac{1}{2} \sigma^2 \right) t + \sigma W_t \right\}. \quad (2.7)$$

The next remark characterizes the optimal stopping times of (2.1) using the Snell envelope theory.

Remark 2.1. *Let us denote*

$$Y_t := \int_0^t e^{-(r+\lambda)s} \lambda \cdot Pf(S_s^0, T-s) ds + e^{-(r+\lambda)t} (K - S_t^0)^+. \quad (2.8)$$

Using the strong Markov property of S^0 , we can determine the Snell envelope of Y as

$$\xi_t := \sup_{\tau \in \mathcal{S}_{t, T}} \mathbb{E} \{ Y_\tau | \mathcal{F}_t \} = e^{-(\lambda+r)t} Jf(S_t^0, T-t) + \int_0^t e^{-(r+\lambda)s} \lambda Pf(S_s^0, T-s) ds, \quad t \in [0, T]. \quad (2.9)$$

Theorem D.12 in [10] implies that the stopping time

$$\tau_x := \inf \{ t \in [0, T] : \xi_t = Y_t \} \wedge T = \inf \{ t \in [0, T] : Jf(S_t^0, T-t) = (K - S_t^0)^+ \}, \quad (2.10)$$

satisfies

$$Jf(x, T) = \mathbb{E}^x \left\{ \int_0^{\tau_x} e^{-(r+\lambda)t} \lambda \cdot Pf(S_t^0, T-t) dt + e^{-(r+\lambda)\tau_x} (K - S_{\tau_x}^0)^+ \right\}. \quad (2.11)$$

Moreover, the stopped process $(e^{-(r+\lambda)(t \wedge \tau_x)} Jf(S_{t \wedge \tau_x}^0, T-t \wedge \tau_x) + \int_0^{t \wedge \tau_x} e^{-(r+\lambda)s} \lambda \cdot Pf(S_s^0, T-s) ds)_{t \geq 0}$ is a martingale. The second infimum in (2.10) is less than T because $Jf(S_T^0, 0) = (K - S_T^0)^+$.

When f is bounded, it follows from the bounded convergence theorem that (using the results of [3] and arguments similar to the ones used in Corollary 7.3 in Chapter 2 of [10])

$$Jf(x, \infty) = \sup_{\tau \in \mathcal{S}_{0, \infty}} \mathbb{E}^x \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot Pf(S_t^0, \infty) dt + e^{-(r+\lambda)\tau} (K - S_\tau^0)^+ \right\}. \quad (2.12)$$

The next three lemmas on the properties of J immediately follow from the definition in (2.1). The first lemma states that J preserves monotonicity.

Lemma 2.1. *Let $T \rightarrow f(x, T)$ be non-decreasing, and $x \rightarrow f(x, T)$ be non-increasing. Then $T \rightarrow Jf(x, T)$ is non-decreasing and $x \rightarrow Jf(x, T)$ is non-increasing.*

The operator J preserves boundedness and order.

Lemma 2.2. *Let $f : \mathbb{R}_+ \times \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}_+$ be a bounded function. Then Jf is also bounded. In fact,*

$$0 \leq \|Jf\|_\infty \leq K + \frac{\lambda}{r + \lambda} \|f\|_\infty. \quad (2.13)$$

Lemma 2.3. *For any $f_1, f_2 : \mathbb{R}_+ \times \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}_+$ that satisfy $f_1(x, T) \leq f_2(x, T)$, we have that $Jf_1(x, T) \leq Jf_2(x, T)$ for all $(x, T) \in \mathbb{R}_+ \times \bar{\mathbb{R}}_+$.*

As we shall see next, the operator J preserves convexity (with respect to the first variable).

Lemma 2.4. *If $f : \mathbb{R}_+ \times \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}_+$ is a convex function in its first variable, then so is $Jf : \mathbb{R}_+ \times \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}_+$.*

Proof. Note that Jf can be written as

$$Jf(x, T) = \sup_{\tau \in \mathcal{S}_{0, T}} \mathbb{E} \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot Pf(xH_t, T-t) dt + e^{-(r+\lambda)\tau} (K - xH_\tau)^+ \right\}. \quad (2.14)$$

Since $f(\cdot, T-t)$ is convex, so is $Pf(\cdot, T-t)$. As a result the integral with respect to time in (2.14) is also convex in x . On the other hand, note that $(K - xH_\tau)^+$ is also a convex function of x . Taking the expectation does not change the convexity with respect to x . Since the upper envelope (supremum) of convex functions is convex, the result follows. \square

Remark 2.2. *Since $x = 0$ is an absorbing boundary for the process S^0 , for any $f : \mathbb{R}_+ \times \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}_+$,*

$$\begin{aligned} Jf(0, T) &= \sup_{t \in \{0, T\}} \left\{ \int_0^t e^{-(r+\lambda)s} \lambda f(0, T-s) ds + e^{-(\lambda+r)t} K \right\} \\ &= \max \left\{ K, \int_0^T e^{-(r+\lambda)s} \lambda f(0, T-s) ds + e^{-(\lambda+r)T} K \right\}, \quad T \geq 0. \end{aligned} \quad (2.15)$$

If we further assume $f \leq K$, then $Jf(0, T) = K$, $T \geq 0$.

Lemma 2.5. *Let us assume that $f : \mathbb{R}_+ \times \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}_+$ is convex in its first variable and $\|f\|_\infty \leq K$. Then $x \rightarrow Jf(x, t)$ satisfies*

$$|Jf(x, T) - Jf(y, T)| \leq |x - y|, \quad (x, y) \in \mathbb{R}_+ \times \bar{\mathbb{R}}_+, \quad (2.16)$$

and all $T \geq 0$.

Proof. First note that a positive convex function that is bounded above has to be non-increasing. Therefore f is non-increasing. As a result of Lemma 2.1, $x \rightarrow Jf(x, t)$ is non-increasing. This function is convex (by Lemma 2.4), and it satisfies

$$Jf(x, T) \geq (K - x)^+ \quad Jf(0, T) = K. \quad (2.17)$$

Consequently, the left and right derivatives of Jf satisfy

$$-1 \leq D_-^x Jf(x, T) \leq D_+^x Jf(x, T) \leq 0, \quad x > 0, T \geq 0. \quad (2.18)$$

Now, the result follows since the derivatives are bounded by 1 (also see Theorem 24.7 (on page 237) in [17]). \square

Remark 2.3. Let $T_0 \in (0, \infty)$ and denote

$$F(x, T) = \sup_{\tau \in \mathcal{S}_{0, T}} \mathbb{E} \left\{ e^{-(r+\lambda)\tau} (K - xH_\tau)^+ \right\}, \quad x \in \mathbb{R}_+, T \in [0, T_0]. \quad (2.19)$$

Then for $S \leq T \leq T_0$

$$F(x, T) - F(x, S) \leq C \cdot |T - S|^{1/2}, \quad (2.20)$$

for all $x \in \mathbb{R}_+$ and for some C that depends only on T_0 . See e.g. equation (2.4) in [14].

The next lemma, which is very crucial for our proof of the smoothness of the American option price for jump diffusions, shows that the increase in the Hölder semi-norm that the operator J causes can be controlled.

Lemma 2.6. Let us assume that for some $L \in (0, \infty)$

$$|f(x, T) - f(x, S)| \leq L|T - S|^{1/2}, \quad (T, S) \in [S_0, T_0] \times [S_0, T_0], \quad (2.21)$$

for all $x \in \mathbb{R}_+$, for $0 \leq S_0 < T_0 < \infty$. Then

$$|Jf(x, T) - Jf(x, S)| \leq (aL + C) |T - S|^{1/2}, \quad (T, S) \in [S_0, T_0] \times [S_0, T_0], \quad (2.22)$$

for some $a \in (0, 1)$ whenever

$$|T - S| < \left(\frac{r}{r + \lambda} \frac{L}{\lambda K} \right)^2. \quad (2.23)$$

Here, $C \in (0, \infty)$ is as in Remark 2.3.

Proof. Without loss of generality we will assume that $T > S$. Then we can write

$$\begin{aligned} Jf(x, T) - Jf(x, S) &\leq \sup_{\tau \in \mathcal{S}_{0, T}} \left[\mathbb{E} \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda Pf(xH_t, T-t) dt + e^{-(r+\lambda)\tau} (K - xH_\tau)^+ \right\} \right. \\ &\quad \left. - \mathbb{E} \left\{ \int_0^{\tau \wedge S} e^{-(r+\lambda)t} \lambda Pf(xH_t, S-t) dt + e^{-(r+\lambda)(\tau \wedge S)} (K - xH_{\tau \wedge S})^+ \right\} \right] \\ &= \sup_{\tau \in \mathcal{S}_{0, T}} \left[\mathbb{E} \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda (Pf(xH_t, T-t) - Pf(xH_t, S-t)) dt \right. \right. \\ &\quad \left. \left. + \mathbb{1}_{\{S < \tau\}} \left[\int_S^\tau e^{-(r+\lambda)t} \lambda Pf(xH_t, S-t) dt + \left(e^{-(r+\lambda)\tau} (K - xH_\tau)^+ - e^{-(r+\lambda)S} (K - xH_S)^+ \right) \right] \right\} \right] \quad (2.24) \\ &\leq \frac{\lambda}{r + \lambda} L (T - S)^{1/2} + \frac{\lambda}{r + \lambda} K \left(e^{-(r+\lambda)S} - e^{-(r+\lambda)T} \right) \\ &\quad + \sup_{\tau \in \mathcal{S}_{S, T}} \mathbb{E} \left\{ \left(e^{-(r+\lambda)\tau} (K - xH_\tau)^+ \right) - \mathbb{E} \left\{ e^{-(r+\lambda)S} (K - xH_S)^+ \right\} \right\} \\ &\leq \frac{\lambda}{r + \lambda} L (T - S)^{1/2} + \lambda K (T - S) + e^{-(r+\lambda)S} (F(H_S, T - S) - F(H_S, 0)), \\ &\leq \left(\frac{\lambda}{r + \lambda} L + C \right) (T - S)^{1/2} + \lambda K (T - S) \end{aligned}$$

in which F is given by (2.19). To derive the second inequality in (2.24), we use the fact that

$$|Pf(xH_t, T-t) - Pf(xH_t, S-t)| \leq \int_{\mathbb{R}_+} \nu(dz) |f(xzH_t, T-t) - f(xzH_t, S-t)| \leq L |T - S|^{1/2}, \quad (2.25)$$

which follows from the assumption in (2.21), and that

$$\mathbb{E} \left\{ 1_{\{S < \tau\}} \int_0^{\tau \wedge S} e^{-(r+\lambda)t} \lambda Pf(xH_t, S-t) dt \right\} \leq \lambda K \mathbb{E} \left\{ \int_S^T e^{-(r+\lambda)t} dt \right\} \leq \frac{\lambda K}{\lambda + K} \left(e^{-(r+\lambda)S} - e^{-(r+\lambda)T} \right). \quad (2.26)$$

To derive the third inequality in (2.24), we use

$$e^{-(r+\lambda)S} - e^{-(r+\lambda)T} \leq e^{-(r+\lambda)S} (r + \lambda)(T - S) \leq (r + \lambda)(T - S). \quad (2.27)$$

The last inequality in (2.24) follows from (2.20). Equation (2.22) follows from (2.24) whenever T and S satisfy (2.23). \square

Let us define the continuation region and its sections by

$$\mathcal{C}^{Jf} := \{(T, x) \in (0, \infty)^2 : Jf(x, T) > (K - x)^+\}, \text{ and } \mathcal{C}_T^{Jf} := \{x \in (0, \infty) : Jf(T, x) > (K - x)^+\}, \quad (2.28)$$

$T > 0$, respectively.

Lemma 2.7. *Suppose that $f : \mathbb{R}_+ \times \bar{\mathbb{R}}_+ \rightarrow \bar{\mathbb{R}}_+$ is such that $x \rightarrow f(x, T)$ is a positive convex function, $T \rightarrow f(x, T)$ is non-decreasing, and $\|f\|_\infty \leq K$. Then for every $T > 0$ there exists $c^{Jf}(T) \in (0, K)$ such that $\mathcal{C}_T^{Jf} = (c^{Jf}(T), \infty)$. Moreover, $T \rightarrow c^{Jf}(T)$ is non-increasing.*

Proof. Let us first show that if $x \geq K$, then $x \in \mathcal{C}_T^{Jf}$ for all $T \geq 0$. Let $\tau_\varepsilon := \inf\{0 \leq t \leq T : S_t^0 \leq K - \varepsilon\}$. Since $\mathbb{P}\{0 < \tau_\varepsilon < T\} > 0$ for $x \geq K$, for all $T > 0$, we have that

$$\mathbb{E}^x \left\{ \int_0^{\tau_\varepsilon} e^{-(r+\lambda)t} \lambda Pf(S_t^0, T-t) dt + e^{-(r+\lambda)\tau_\varepsilon} (K - S_{\tau_\varepsilon}^0)^+ \right\} > 0, \quad (2.29)$$

which implies that $x \in \mathcal{C}_T^{Jf}$. On the other hand, it is clear that

$$(K - x)^+ \leq Jf(x, T) \leq Jf(x, \infty), \quad (x, T) \in \mathbb{R}_+ \times \bar{\mathbb{R}}_+. \quad (2.30)$$

Thanks to in Lemma 2.6 of [3], there exist $l^f \in (0, K)$ such that

$$Jf(x, \infty) = (K - x)^+, \quad x \in [0, l^f]; \quad Jf(x, \infty) > (K - x)^+, \quad x \in (l^f, \infty). \quad (2.31)$$

Since $x \rightarrow Jf(x, \infty)$ and $x \rightarrow Jf(x, T)$, $T \geq 0$, are convex functions (from Lemma 2.2 in [3] and Lemma 2.4 respectively), (2.29), (2.30) and (2.31) imply that there exists a point $c^{Jf}(T) \in (l^f, K)$ such that

$$Jf(x) = (K - x)^+, \quad x \in [0, c^{Jf}(T)]; \quad Jf(x, T) > (K - x)^+, \quad x \in (c^{Jf}(T), \infty), \quad (2.32)$$

for $T > 0$. This proves the first statement of the Lemma. The fact that $T \rightarrow c(T)$ is non-increasing follows from the fact that $T \rightarrow Jf(x, T)$ is non-decreasing. \square

In the following lemma we will argue that if f has certain regularity properties, then Jf is the classical solution of a parabolic free boundary equation.

Lemma 2.8. *Let us assume that $f : \mathbb{R}_+ \times \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}_+$ is convex in its first variable, $\|f\|_\infty \leq K$ and $T \rightarrow f(x, T)$ is non-increasing. Moreover, we will assume that f satisfies*

$$|f(x, T) - f(x, S)| \leq A |T - S|^{1/2} \quad \text{whenever } |T - S| < B, \quad (2.33)$$

for all $x \in \mathbb{R}_+$, where A, B are strictly positive constants that do not depend on x . Then the function $Jf : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is the unique bounded solution (in the classical sense) of

$$Au(x, T) - (r + \lambda) \cdot u(x, T) + \lambda \cdot (Pf)(x, T) - \frac{\partial}{\partial T} u(x, T) = 0 \quad x > c^{Jf}(T), \quad (2.34)$$

$$u(x, T) = (K - x) \quad x \leq c^{Jf}(T), \quad (2.35)$$

in which \mathcal{A} is as in (2.5) and c^{Jf} is as in Lemma 2.7. Moreover,

$$\mathcal{A}Jf(x, T) - (r + \lambda) \cdot Jf(x, T) + \lambda \cdot (Pf)(x, T) - \frac{\partial}{\partial T}Jf(x, T) \leq 0, \quad x < c^{Jf}(T). \quad (2.36)$$

Proof. The proof is motivated by Theorem 2.7.7 of [10]. Equation (2.35) is clearly satisfied by Jf . In what follows, we will first show that Jf satisfies (2.34). Let us take a point in $(t, T) \in \mathcal{C}^{Jf}$ and consider a bounded rectangle $R = (t_1, t_2) \times (x_1, x_2)$ containing this point. We will let

$$t_2 - t_1 < B \wedge \left(\frac{rA}{(r + \lambda)\lambda K} \right)^2. \quad (2.37)$$

Let $\partial_0 R$ be the parabolic boundary of R and consider the parabolic partial differential equation

$$\begin{aligned} \mathcal{A}u(x, T) - (r + \lambda) \cdot u(x, T) + \lambda \cdot (Pf)(x, T) - \frac{\partial}{\partial T}u(x, T) &= 0 \quad \text{in } R, \\ u(x, T) &= Jf(x, T) \quad \text{on } \partial_0 R. \end{aligned} \quad (2.38)$$

As a result of Lemmas 2.5 and 2.6, Jf satisfies the uniform Lipschitz and Hölder continuity conditions, which implies that Jf is continuous. On the other hand, for any $(T, x) \in R$

$$\begin{aligned} |Pf(x, T) - Pf(y, S)| &\leq |Pf(x, T) - Pf(x, S)| + |Pf(x, S) - Pf(y, S)| \\ &\leq \int_{\mathbb{R}_+} \nu(dz) (|f(xz, T) - f(xz, S)| + |f(xz, S) - f(yz, S)|) \\ &\leq A |T - S|^{1/2} + \xi |x - y|, \end{aligned} \quad (2.39)$$

Now, Theorem 5.2 in [8] implies that (2.38) has a unique classical solution. We will show that this unique solution coincides with Jf using optional sampling theorem. Let us introduce the stopping time

$$\tau := \inf\{\theta \in [0, t_0 - t_1] : (t_0 - \theta, x_0 H_\theta) \in \partial_0 R\} \wedge (t_0 - t_1), \quad (2.40)$$

which is the first time S^0 hits the parabolic boundary when S^0 starts from (x_0, t_0) . Let us also define the process $N_\theta := e^{-(r+\lambda)\theta} u(x_0 H_\theta, t_0 - \theta) + \int_0^\theta e^{-(r+\lambda)t} \lambda \cdot Pf(S_t^0, t_0 - t) dt$, $\theta \in [0, t_0 - t_1]$. From the classical Itô's formula it follows that the stopped process $N_{\theta \wedge \tau}$ is a bounded martingale. As a result

$$u(x_0, t_0) = N_0 = \mathbb{E}^x \{N_\tau\} = \mathbb{E} \left\{ e^{-(r+\lambda)\tau} Jf(x H_\tau, t_0 - \tau) + \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot Pf(S_t^0, t_0 - t) dt \right\}. \quad (2.41)$$

Clearly $\tau \leq \tau_x$. Since the stopped process $(e^{-(r+\lambda)(t \wedge \tau_x)} Jf(S_{t \wedge \tau_x}^0, t_0 - t \wedge \tau_x) + \int_0^{t \wedge \tau_x} e^{-(r+\lambda)t} \lambda \cdot Pf(S_s^0, t_0 - s) ds)_{t \geq 0}$ is a bounded martingale, another application of the optional sampling theorem yields

$$\mathbb{E} \left\{ e^{-(r+\lambda)\tau} Jf(x_0 H_\tau, t_0 - \tau) + \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot Pf(S_t^0, t_0 - t) dt \right\} = Jf(x_0, t_0). \quad (2.42)$$

Combining (2.41) and (2.42), we see that (2.34) is satisfied in the classical sense since the choice of $(x_0, t_0) \in \mathcal{C}^{Jf}$ is arbitrary.

We still need to show uniqueness among bounded functions. Fix $x > c^{Jf}(T)$. Let u be a bounded function satisfying (2.34) and (2.35). Let us define $M_t := e^{-(r+\lambda)t} u(x H_t, T - t) + \int_0^t e^{-(r+\lambda)s} \lambda \cdot Pf(S_s^0, T - s) ds$. Using the classical Itô formula it can be seen that $M_{t \wedge \tau_x}$ is a bounded martingale. Since τ_x is optimal (see (2.11)), by the optional sampling theorem, we have

$$\begin{aligned} u(x, T) &= M_0 = \mathbb{E}^x \{M_{\tau_x}\} = \mathbb{E} \left\{ e^{-(r+\lambda)\tau_x} u(x H_{\tau_x}, T - \tau_x) + \int_0^{\tau_x} e^{-(r+\lambda)t} \lambda \cdot Pf(S_s^0, T - s) ds \right\} \\ &= \mathbb{E} \left\{ e^{-(r+\lambda)\tau_x} (K - x H_{\tau_x})^+ + \int_0^{\tau_x} e^{-(r+\lambda)t} \lambda \cdot Pf(S_s^0, T - s) ds \right\} = Jf(x, T). \end{aligned} \quad (2.43)$$

Next, we will prove (2.36). To this end, let $x < c^{Jf}(t)$. Let U a closed interval centered at x such that $U \subset (0, c^{Jf}(T))$. Let $\tau_U = \{t \geq 0 : xH_t \notin U\}$. Since $(e^{-(r+\lambda)t} Jf(S_t^0, T-t) + \int_0^t e^{-(r+\lambda)s} \lambda \cdot Pf(S_s^0, T-s) ds)_{t \geq 0}$ is a supermartingale we can write

$$\mathbb{E} \left[e^{-(r+\lambda)(\tau_U \wedge t)} Jf(xH_{\tau_U \wedge t}, T - \tau_U \wedge t) + \int_0^{\tau_U \wedge t} e^{-(r+\lambda)u} \lambda Pf(xH_u, T-u) du \right] \leq Jf(x, T), \quad (2.44)$$

for all $t \geq 0$. Since $Jf(x, t) = K - x$ when $(T, x) \in \mathbb{R}_+^2 - C^{Jf}$, we can apply Itô's formula to obtain that

$$\lim_{t \rightarrow 0} \mathbb{E} \left[\frac{1}{t} \int_0^{\tau_U \wedge t} e^{-(r+\lambda)u} \left(\left(\mathcal{A} - (r+\lambda) \cdot -\frac{\partial}{\partial T} \right) Jf(xH_u, t-u) + \lambda Pf(xH_u, T-u) \right) du \right] \leq 0. \quad (2.45)$$

Now, (2.36) follows thanks to dominated convergence theorem, which allows us to exchange the limit and the expectation. We can apply the dominated convergence theorem thanks to the fact that U is a compact domain. \square

Lemma 2.9. *For a given $T > 0$, let $x \rightarrow f(x, T)$ be a convex and non-increasing function. Then the convex function $x \rightarrow Jf(x, T)$ is of class C^1 at $x = c(T)$, i.e.,*

$$\left. \frac{\partial}{\partial x} Jf(x, T) \right|_{x=c(T)} = -1. \quad (2.46)$$

Proof. The proof is similar to the proof of Lemma 7.8 on page 74 of [10], but we will provide it here for the sake of completeness. If we let $x = c(T)$, then

$$\begin{aligned} Jf(x + \varepsilon, T) &= \mathbb{E} \left\{ \int_0^{\tau_{x+\varepsilon}} e^{-(r+\lambda)t} \lambda \cdot Pf((x + \varepsilon)H_t, T-t) dt + e^{-(r+\lambda)\tau_{x+\varepsilon}} (K - (x + \varepsilon)H_{\tau_{x+\varepsilon}})^+ \right\} \\ &= \mathbb{E} \left\{ \int_0^{\tau_{x+\varepsilon}} e^{-(r+\lambda)t} \lambda \cdot Pf(xH_t, T-t) dt + e^{-(r+\lambda)\tau_{x+\varepsilon}} (K - xH_{\tau_{x+\varepsilon}})^+ \right\} \\ &\quad + \mathbb{E} \left\{ \int_0^{\tau_{x+\varepsilon}} e^{-(r+\lambda)t} \lambda \cdot [Pf((x + \varepsilon)H_t, T-t) - Pf(xH_t, T-t)] dt \right\} \\ &\quad + \mathbb{E} \left\{ e^{-(r+\lambda)\tau_{x+\varepsilon}} [(K - (x + \varepsilon)H_{\tau_{x+\varepsilon}})^+ - (K - xH_{\tau_{x+\varepsilon}})^+] \right\} \\ &\leq Jf(x, T) + \mathbb{E} \left\{ 1_{\{\tau_{x+\varepsilon} < T\}} e^{-(r+\lambda)\tau_{x+\varepsilon}} [(K - (x + \varepsilon)H_{\tau_{x+\varepsilon}}) - (K - xH_{\tau_{x+\varepsilon}})] \right\} \\ &\quad + \mathbb{E} \left\{ 1_{\{\tau_{x+\varepsilon} = T\}} e^{-(r+\lambda)\tau_{x+\varepsilon}} [(K - (x + \varepsilon)H_{\tau_{x+\varepsilon}})^+ - (K - xH_{\tau_{x+\varepsilon}})^+] \right\} \\ &\leq Jf(x, T) - \varepsilon \mathbb{E}^x \left\{ 1_{\{\tau_{x+\varepsilon} < T\}} e^{-(r+\lambda)\tau_{x+\varepsilon}} H_{\tau_{x+\varepsilon}} \right\} \\ &= Jf(x, T) - \varepsilon \mathbb{E}^x \left\{ e^{-(r+\lambda)\tau_{x+\varepsilon}} H_{\tau_{x+\varepsilon}} \right\} + \varepsilon \mathbb{E}^x \left\{ 1_{\{\tau_{x+\varepsilon} = T\}} e^{-(r+\lambda)T} H_T \right\}. \end{aligned} \quad (2.47)$$

The first inequality follows since $\tau_{x+\varepsilon}$ is not optimal when S^0 starts at x and $x \rightarrow Pf(x, T)$ is a decreasing function for any $T \geq 0$. From (2.47) it follows that

$$D_+^x Jf(x, T) \leq -1, \quad (2.48)$$

since $e^{-(r+\lambda)t} H_t$ is a uniformly integrable martingale and $\tau_{x+\varepsilon} \downarrow 0$. Convexity of $Jf(t, x)$ (Lemma 2.4) implies that

$$-1 = D_-^x Jf(x-, t) \leq D_+^x Jf(x+, t) \leq -1, \quad (2.49)$$

which yields the desired result. \square

3. A SEQUENCE OF FUNCTIONS APPROXIMATING V

Let us define a sequence of functions by the following iteration:

$$v_0(x, T) = (K - x)^+, \quad v_{n+1}(x, T) = Jv_n(x, T), \quad n \geq 0, \quad \text{for all } (x, T) \in \mathbb{R}_+ \times \mathbb{R}_+. \quad (3.1)$$

We extend these functions onto $\mathbb{R}_+ \times \bar{\mathbb{R}}_+$ by letting

$$v_n(x, \infty) = \lim_{T \rightarrow \infty} v_n(x, T). \quad (3.2)$$

This sequence of functions is a bounded sequence as the next lemma shows.

Corollary 3.1. *For all $n \geq 0$,*

$$(K - x)^+ \leq v_n(x, T) \leq \left(1 + \frac{\lambda}{r}\right) K, \quad (x, T) \in \mathbb{R}_+ \times \bar{\mathbb{R}}_+. \quad (3.3)$$

Proof. The first inequality follows since it may not be optimal to stop immediately. Let us prove the second inequality using an induction argument: Observe that $v_0(x, T) = (K - x)^+$, $(x, T) \in \mathbb{R}_+ \times \bar{\mathbb{R}}_+$, satisfies (3.3). Let us assume that (3.3) holds for n and show that it holds for $n + 1$. Using (2.13), we get that

$$\|v_{n+1}\|_\infty = \|Jv_n\|_\infty \leq K + \frac{\lambda}{r + \lambda} \left(1 + \frac{\lambda}{r}\right) K = \left(1 + \frac{\lambda}{r}\right) K. \quad (3.4)$$

□

As a corollary of Lemmas 2.3 and 2.4 we can state the following corollary, whose proof can be carried out by induction.

Corollary 3.2. *The sequence $(v_n(x, T))_{n \geq 0}$ is increasing for all $(x, T) \in \mathbb{R}_+ \times \bar{\mathbb{R}}_+$. For each n , the function $x \rightarrow v_n(x, T)$, $x \geq 0$, is convex for all $T \in \bar{\mathbb{R}}_+$.*

Remark 3.1. *Let us define,*

$$v_\infty(x, T) := \sup_{n \geq 0} v_n(x, T), \quad (x, T) \in \mathbb{R}_+ \times \bar{\mathbb{R}}_+. \quad (3.5)$$

This function is well defined as a result of (3.3) and Corollary 3.2. In fact, it is convex, because it is the upper envelope of convex functions, and it is bounded by the right-hand-side of (3.3).

Corollary 3.3. *For each $n \geq 0$ and $t \in \mathbb{R}_+$, $x \rightarrow v_n(x, T)$, is a decreasing function on $[0, \infty)$. Moreover, $T \rightarrow v_n(x, T)$ is non-decreasing. The same statements hold for $x \rightarrow v_\infty(x, T)$, and $T \rightarrow v_\infty(x, T)$, respectively.*

Proof. The behaviour with respect to the first variable is a result of Corollary 3.2 and Remark 3.1 since any positive convex function that is bounded from above is decreasing. For each n , the fact that $T \rightarrow v_n(x, T)$ is non-decreasing is a corollary of Lemma 2.1. On the other hand, for any $T \geq S \geq 0$, we have that $v_\infty(x, T) = \sup_n v_n(x, T) \geq \sup_n v_n(x, S) = v_\infty(x, S)$. □

Next, we will sharpen the upper bound in Corollary 3.1. This improvement has some implications for the continuity of $x \rightarrow v_n(x, T)$, $n \geq 1$, and $x \rightarrow v_\infty(x, T)$ at $x = 0$.

Remark 3.2. *The upper bound in (3.1) can be sharpened using Corollary 3.3 and Remark 2.2. Indeed, we have*

$$(K - x)^+ \leq v_n(x, T) < K, \quad \text{for each } n, \quad \text{and} \quad (K - x)^+ \leq v_\infty(x, T) < K, \quad (x, T) \in (0, \infty)^2. \quad (3.6)$$

It follows from this observation that for every $T \in \bar{\mathbb{R}}_+$, $x \rightarrow v_n(x, T)$, for every n , and $x \rightarrow v_\infty(x, T)$, are continuous at $x = 0$ since $v_n(0, T) = v_\infty(0, T) = K$ and these functions are convex. (Note that convexity already guarantees continuity for $x > 0$.)

Lemma 3.1. *The function v_∞ is the smallest fixed point of the operator J .*

Proof.

$$\begin{aligned}
v_\infty(x, T-t) &= \sup_{n \geq 1} v_n(x, T-t) \\
&= \sup_{n \geq 1} \sup_{\tau \in \mathcal{S}_{0,T}} \mathbb{E}^x \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot P v_n(S_t^0, T-t) dt + e^{-(r+\lambda)\tau} (K - S_\tau^0)^+ \right\} \\
&= \sup_{\tau \in \mathcal{S}_{0,T}} \sup_{n \geq 1} \mathbb{E}^x \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot P v_n(S_t^0, T-t) dt + e^{-(r+\lambda)\tau} (K - S_\tau^0)^+ \right\} \\
&= \sup_{\tau \in \mathcal{S}_{0,T}} \mathbb{E}^x \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot P(\sup_{n \geq 1} v_n)(S_t^0, T-t) dt + e^{-(r+\lambda)\tau} (K - S_\tau^0)^+ \right\} \\
&= J v_\infty(x, T-t),
\end{aligned} \tag{3.7}$$

in which the fourth equality follows by applying the monotone convergence theorem three times. Let $w : \mathbb{R}_+ \times \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}_+$ be another fixed point of the operator J . We will argue by induction that $w \geq v_\infty$. For $(x, t) \in \mathbb{R}_+ \times \bar{\mathbb{R}}_+$, $w(x, T-t) = Jw(x, T-t)$, which implies that $w(x, T-t) = Jw(x, T-t) \geq (K-x)^+ = v_0(\cdot)$. If we assume that $w(x, T-t) \geq v_n(x, T-t)$, then $w(x, T-t) = Jw(x, T-t) \geq Jv_n(x, T-t) = v_{n+1}(x, T-t)$. Consequently $w(x, T-t) \geq v_n(x, T-t)$ for all $n \geq 0$. As a result $w(x, T-t) \geq \sup_{n \geq 0} v_n(x, T-t) = v_\infty(x, T-t)$. \square

Lemma 3.2. *The sequence $\{v_n(\cdot, \cdot)\}_{n \geq 0}$ converges uniformly to v_∞ . In fact, the rate of convergence is exponential:*

$$v_n(x, T) \leq v_\infty(x, T) \leq v_n(x, T) + \left(\frac{\lambda}{\lambda+r} \right)^n K, \quad (x, T) \in \mathbb{R}_+ \times \bar{\mathbb{R}}_+. \tag{3.8}$$

Proof. The first inequality follows from the definition of v_∞ . The second inequality can be proved by induction. The inequality holds when we set $n = 0$ by Remark 3.2. Assume that the inequality holds for $n > 0$. Then

$$\begin{aligned}
v_\infty(x, T) &= \sup_{\tau \in \mathcal{S}_{0,T}} \mathbb{E}^x \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot P v_\infty(S_t^0, T-t) dt + e^{-(r+\lambda)\tau} (K - S_\tau^0)^+ \right\} \\
&\leq \sup_{\tau \in \mathcal{S}_{0,T}} \mathbb{E}^x \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot P v_n(S_t^0, T-t) dt + e^{-(r+\lambda)\tau} (K - S_\tau^0)^+ \right\} + \int_0^\infty dt e^{-(\lambda+r)t} \lambda \left(\frac{\lambda}{\lambda+r} \right)^n K \\
&= v_{n+1}(x, T) + \left(\frac{\lambda}{\lambda+r} \right)^{n+1} K.
\end{aligned} \tag{3.9}$$

\square

Remark 3.3. *Note that, for a fixed $T_0 > 0$,*

$$v_n(x, T) \leq v_\infty(x, T) \leq v_n(x, T) + \left(1 - e^{-(r+\lambda)T_0} \right)^n \left(\frac{\lambda}{\lambda+r} \right)^n K, \quad x \in \mathbb{R}_+, T \in (0, T_0). \tag{3.10}$$

This can be derived using an induction argument similar to the one used in the proof of Lemma 3.2. We simply replace (3.9) by

$$\begin{aligned} v_\infty(x, T) &\leq \sup_{\tau \in S_{0, T}} \mathbb{E}^x \left\{ \int_0^\tau e^{-(r+\lambda)t} \lambda \cdot P v_n(S_t^0, T-t) dt + e^{-(r+\lambda)\tau} (K - S_\tau^0)^+ \right\} \\ &+ \int_0^{T_0} dt e^{-(\lambda+r)t} \left(1 - e^{-(r+\lambda)T_0}\right)^n \lambda \left(\frac{\lambda}{\lambda+r}\right)^n K = v_{n+1}(x, T) + K \left(1 - e^{-(r+\lambda)T_0}\right)^{n+1} \left(\frac{\lambda}{\lambda+r}\right)^{n+1} \end{aligned} \quad (3.11)$$

Observe that one can replace K in (3.10) by $\|v_\infty - v_0\|_\infty$. Note that the convergence rate in (3.10) is fast. This will lead to a numerical scheme, whose error versus accuracy characteristics can be controlled, for pricing American options.

Remark 3.4. Let $T_0 \in (0, \infty)$. It can be shown using similar arguments to the ones used in the proof of Lemma 2.6 that

$$|v_1(x, T) - v_1(x, S)| \leq \lambda K |T - S| + C |T - S|^{1/2}, \quad T, S \in (0, T_0], \quad (3.12)$$

for all $x \in \mathbb{R}_+$, in which $C \in (0, \infty)$ is as in Remark 2.3. In fact

$$|v_1(x, T) - v_1(x, S)| \leq L |T - S|^{1/2}, \quad (3.13)$$

for all $x \in \mathbb{R}_+$ and for some L that depends only on T_0 .

The next lemma shows that the functions v_n , $n \geq 0$, and v_∞ are locally Hölder continuous with respect to the time variable.

Lemma 3.3. Let $T_0 \in (0, \infty)$ and $L \in (0, \infty)$ be as in Remark 3.4 and $C \in (0, \infty)$ be as in Remark 2.3. Then for $T, S \in (0, T_0)$, we have that

$$|v_n(x, T) - v_n(x, S)| \leq \left(L + \frac{C}{1-a}\right) |T - S|^{1/2} \quad \text{whenever } |T - S| \leq \left(\frac{r}{r+\lambda} \frac{L}{\lambda K}\right)^2, \quad (3.14)$$

for all $x \in \mathbb{R}_+$ and for all $n \geq 1$. Here, $a \in (0, 1)$ is as in Lemma 2.6. Moreover,

$$|v_\infty(x, T) - v_\infty(x, S)| \leq \left(L + \frac{C}{1-a}\right) |T - S|^{1/2} \quad \text{whenever } |T - S| \leq \left(\frac{r}{r+\lambda} \frac{L}{\lambda K}\right)^2, \quad (3.15)$$

for all $x \in \mathbb{R}_+$.

Proof. The proof of (3.14) will be carried out using an induction argument. Observe from Remark 3.4 that (3.14) holds for $n = 1$. Let us assume that (3.14) holds for n and show that it holds for $n + 1$. Using Lemma 2.6, we have that

$$|v_{n+1}(x, T) - v_{n+1}(x, S)| \leq \left(a \left(L + \frac{C}{1-a}\right) + C\right) |T - S|^{1/2}, \quad (3.16)$$

for $|T - S| \leq \left(\frac{r}{r+\lambda} \frac{L+C/(1-a)}{\lambda K}\right)^2$. It is clear that the right-hand-side of (3.16) is less than that of (3.14), and

$$\frac{r}{r+\lambda} \frac{L+C/(1-a)}{\lambda K} \geq \frac{r}{r+\lambda} \frac{L}{\lambda K}, \quad (3.17)$$

from which the first statement of the lemma follows. Now let us prove (3.15). To this end observe that

$$\begin{aligned} |v_\infty(x, T) - v_\infty(x, S)| &\leq |v_\infty(x, T) - v_n(x, T)| + |v_n(x, T) - v_n(x, S)| + |v_\infty(x, S) - v_n(x, S)| \\ &\leq 2 \left(\frac{\lambda}{\lambda+r}\right)^n K + \left(L + \frac{C}{1-a}\right) |T - S|^{1/2}, \end{aligned} \quad (3.18)$$

for any $n > 1$, which follows from (3.14) and Lemma 3.2. The result follows since n on the right-hand-side of (3.18) is arbitrary. \square

Lemma 3.4. For $n \geq 0$, $|v_n(x, T) - v_n(y, T)| \leq |x - y|$, and $|v_\infty(x, T) - v_\infty(y, T)| \leq |x - y|$, $(x, y) \in \mathbb{R}_+ \times \mathbb{R}_+$, for all $T \geq 0$.

Proof. It follows from Remark 3.2 that $\|v_n\|_\infty \leq K$, for all $n \geq 0$, and $\|v_\infty\|_\infty \leq K$. Moreover, for each $n \geq 0$, $v_n(\cdot, T)$ is convex (for all $T \in \mathbb{R}_+$) as a result of Corollary 3.2. On the other hand, it was pointed in Remark 3.1 that $v_\infty(\cdot, T)$ is convex for all $T \in \mathbb{R}_+$. Since

$$v_{n+1}(x, T) = Jv_n(x, T) \quad \text{and} \quad v_\infty(x, T) = Jv_\infty(x, T), \quad (3.19)$$

the statement of the lemma follows from Lemma 2.5. \square

Lemma 3.5. For all $T \geq 0$ and $n \geq 0$, $\mathcal{C}_T^{v_{n+1}} = (c^{v_{n+1}}(T), \infty)$ for some $c^{v_{n+1}}(T) \in (0, K)$ and $\mathcal{C}_T^{v_\infty} = (c^{v_\infty}(T), \infty)$ for some $c^{v_\infty} \in (0, K)$. The function v_{n+1} is the unique bounded solution (in the classical sense) of

$$\begin{aligned} \mathcal{A}v_{n+1}(x, T) - (r + \lambda) \cdot v_{n+1}(x, T) + \lambda \cdot (Pv_n)(x, T) - \frac{\partial}{\partial T}v_{n+1}(x, T) &= 0, \quad x > c^{v_{n+1}}(T), \\ v_{n+1}(x, T) &= (K - x), \quad x \leq c^{v_{n+1}}(T), \end{aligned} \quad (3.20)$$

and it satisfies

$$\left. \frac{\partial}{\partial x}v_{n+1}(x, T) \right|_{x=c^{v_{n+1}}(T)} = -1, \quad T > 0. \quad (3.21)$$

Moreover, v_∞ is the unique bounded solution (in the classical sense) of

$$\begin{aligned} \mathcal{A}v_\infty(x, T) - (r + \lambda) \cdot v_\infty(x, T) + \lambda \cdot (Pv_\infty)(x, T) - \frac{\partial}{\partial T}v_\infty(x, T) &= 0 \quad x > c^{v_\infty}(T), \\ v_\infty(x, T) &= (K - x) \quad x \leq c^{v_\infty}(T), \end{aligned} \quad (3.22)$$

and it satisfies

$$\left. \frac{\partial}{\partial x}v_\infty(x, T) \right|_{x=c^{v_\infty}(T)} = -1, \quad T > 0. \quad (3.23)$$

On the other hand,

$$\mathcal{A}v_\infty(x, T) - (r + \lambda) \cdot v_\infty(x, T) + \lambda \cdot (Pv_\infty)(x, T) - \frac{\partial}{\partial T}v_\infty(x, T) \leq 0 \quad x < c^{v_\infty}(T). \quad (3.24)$$

Proof. The fact that $\mathcal{C}^{v_{n+1}} = (c^{v_{n+1}}, \infty)$ and $\mathcal{C}^{v_\infty} = (c^{v_\infty}, \infty)$ for some $c^{v_{n+1}} \in (0, K)$ and $c^{v_\infty} \in (0, K)$ follows from Lemma 2.7 since the assumptions in that lemma hold thanks to Corollaries 3.2, 3.3; Remarks 3.1 and 3.2; and Lemma 3.1.

The partial differential equations (3.20), (3.22); and the inequality in (3.24) are satisfied as a corollary of Lemma 2.8; Corollaries 3.2 and 3.3, Remarks 3.1, 3.2; Lemmas 3.1, 3.3.

Observe that since v_n is convex (Corollary 3.2) and non-increasing (Corollary 3.3) with respect to its first variable, v_{n+1} ($= Jv_n$) satisfies the smooth fit condition in (3.21) as a result of Lemma 2.9. The smooth fit condition in (3.23) holds for v_∞ as a result of Lemma 2.9 since v_∞ ($= Jv_\infty$) (Lemma 3.1) and $x \rightarrow v_\infty(x, T)$ is non-increasing and convex. \square

The next lemma will be used to verify the fact that $V = v_\infty$. The classical Itô's rule can not be applied to the process $t \rightarrow v_\infty(S_t, T - t)$ since the function v_∞ may fail to be $C^{2,1}$ at $T \rightarrow c^{v_\infty}(T)$. As a result, the semi-martingale decomposition of the process $t \rightarrow v_\infty(S_t, T - t)$ may contain an extra term due to the local time of the process S at the free boundary.

Lemma 3.6. *Let $X = \{X_t; t \geq 0\}$ be a semi-martingale and $b : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a continuous function of bounded variation. Let $F : \mathbb{R} \times \mathbb{R}_+ \rightarrow \mathbb{R}$ be a continuous function that is $C^{2,1}$ on \bar{C} and \bar{D} (it may not be necessarily $C^{1,2}$ across the boundary curve b), in which*

$$C \triangleq \{(x, t) \in \mathbb{R} \times \mathbb{R}_+ : x < b(t)\}, \quad D \triangleq \{(x, t) \in \mathbb{R} \times \mathbb{R}_+ : x > b(t)\}.$$

That is, there exist two functions $F^1, F^2 : \mathbb{R} \times \mathbb{R}_+ \rightarrow \mathbb{R}$, that $C^{2,1}$ on $\mathbb{R} \times \mathbb{R}_+$, and $F(x, t) = F^1(x, t)$ when $(x, t) \in C$ and $F(x, t) = F^2(x, t)$ when $(x, t) \in D$. Moreover, $F^1(b(t), t) = F^2(b(t), t)$. Then the following generalization of Itô's formula holds:

$$\begin{aligned} F(X_t, t) &= F(X_0, 0) + \int_0^t \frac{1}{2} [F_t(X_{s-+}, s) + F_t(X_{s--}, s)] ds \\ &\quad + \frac{1}{2} \int_0^t [F_x(X_{s-+}, s) + F_x(X_{s--}, s)] dX_s + \frac{1}{2} \int_0^t 1_{\{X_{s-} \neq b(s)\}} F_{xx}(X_{s-}, s) d\langle X, X \rangle_s^c \\ &\quad + \sum_{0 < s \leq t} \left\{ F(X_s, s) - F(X_{s-}, s) - \frac{1}{2} \Delta X_s [F_x(X_{s-}, s) + F_x(X_{s-+}, s)] \right\} \\ &\quad + \frac{1}{2} \int_0^t [F_x(X_{s-+}, s) - F_x(X_{s--}, s)] 1_{\{X_{s-} = b(s)\}} dL_t^b, \end{aligned} \tag{3.25}$$

where L_t^b is the local time of the semi-martingale $X_t - b(t)$ at zero (see the definition on page 216 in [16]).

Lemma 3.6 was stated in Theorem 2.1 of [12] for continuous semimartingales. The generalization for the case when the underlying process is not necessarily continuous is intuitively clear and just technical, but we will prove it in the Appendix for the sake of completeness. We are now ready to state the main results.

Theorem 3.1. *The value function V is the unique bounded solution (in the classical sense) of the integro-partial differential equation in (3.22). Given $(x, T) \in \mathbb{R}_+ \times \mathbb{R}_+$ belongs to the optimal continuation region if $x > c^{v_\infty}(T)$. Moreover, it satisfies the smooth fit condition at the optimal stopping boundary, i.e.,*

$$\left. \frac{\partial}{\partial x} V(x, T) \right|_{x=c^{v_\infty}(T)} = -1, \quad T > 0.$$

Proof. The proof is a corollary of the optional sampling theorem and the generalized Itô's formula given above. Let $T \in (0, \infty)$ and define

$$\widetilde{M}_t = e^{-rt} v_\infty(S_t, T - t), \quad \text{and} \quad \widetilde{\tau}_x := T \wedge \inf\{t \in [0, T] : S_t \leq c^{v_\infty}(T - t)\}. \tag{3.26}$$

It follows from (3.22) and the classical Itô's lemma that $\{\widetilde{M}_{t \wedge \widetilde{\tau}_x}\}_{0 \leq t \leq T}$ is a bounded \mathbb{P} -martingale. Using the optional sampling theorem, one obtains

$$v_\infty(x, T) = \widetilde{M}_0 = \mathbb{E}^x \left\{ \widetilde{M}_{\widetilde{\tau}_x} \right\} = \mathbb{E}^x \left\{ e^{-r\widetilde{\tau}_x} v_\infty(S_{\widetilde{\tau}_x}, T - \widetilde{\tau}_x) \right\} = \mathbb{E}^x \left\{ e^{-r\widetilde{\tau}_x} (K - S_{\widetilde{\tau}_x})^+ \right\} \leq V(x, T). \tag{3.27}$$

In the rest of the proof we will show that $v_\infty(x, T) \geq V(x, T)$. Since v_∞ satisfies the smooth fit principle across the free boundary, when we apply the generalized Itô's formula to $v_\infty(S_t, T - t)$, the local time term drops. Thanks to (3.22) and (3.24), $v_\infty(S_t, T - t)$ is a positive \mathbb{P} -supermartingale. Again, using the optional sampling theorem, for any $\tau \in \widetilde{\mathcal{S}}_{0, T}$

$$v_\infty(x, T) = \widetilde{M}_0 \geq \mathbb{E}^x \left\{ \widetilde{M}_\tau \right\} = \mathbb{E}^x \left\{ e^{-r\tau} v_\infty(S_\tau, T - \tau) \right\} \geq \mathbb{E}^x \left\{ e^{-r\tau} (K - S_\tau)^+ \right\}. \tag{3.28}$$

As a result $v_\infty(x, T) \geq V(x, T)$. □

Remark 3.5. *We have that*

$$\mathcal{C}_T^{v_\infty} = \{x \in (0, \infty) : v_\infty > (K - x)^+\} = (c^{v_\infty}(T), \infty). \quad (3.29)$$

On the other hand, $v_\infty = K - x$ for $x \leq c^{v_\infty}$. Since $V = v_\infty$, by Theorem 3.1, it follows that

$$\mathcal{C}_T^V = \{x \in (0, \infty) : V > (K - x)^+\} = (c^{v_\infty}(T), \infty). \quad (3.30)$$

A. APPENDIX

Proof of Lemma 3.6. As in [12] we will define $Z_t^1 = X_t \wedge b(t)$, $Z_t^2 = X_t \vee b(t)$, and observe that

$$F(X_t, t) = F^1(Z_t^1, t) + F^2(Z_t^2, t) - F(b(t), t). \quad (A.1)$$

On the other hand, applying the Meyer-Itô formula (see Theorem 70 in [16]) to the semi-martingale $X_t - b(t)$, we obtain

$$\begin{aligned} |X_t - b(t)| &= |X_0 - b(0)| + \int_0^t \text{sign}(X_{s-} - b(s)) d(X_s - b(s)) \\ &\quad + 2 \sum_{0 < s \leq t} [1_{\{X_{s-} > b(s)\}} (X_s - b(s))^- + 1_{\{X_{s-} \leq b(s)\}} (X_s - b(s))^+] + L_t^b. \end{aligned} \quad (A.2)$$

Since $Z_t^1 = \frac{1}{2}(X_t + b(t) - |X_t - b(t)|)$ and $Z_t^2 = \frac{1}{2}(X_t + b(t) + |X_t - b(t)|)$, using (A.2), we get

$$\begin{aligned} dZ_t^1 &= \frac{1}{2} \left\{ (1 - \text{sign}(X_{t-} - b(t))) dX_t + (1 + \text{sign}(X_{t-} - b(t))) db(t) - dL_t^b \right\} \\ &\quad - [1_{\{X_{t-} > b(t)\}} (X_t - b(t))^- + 1_{\{X_{t-} \leq b(t)\}} (X_t - b(t))^+], \end{aligned} \quad (A.3)$$

$$\begin{aligned} dZ_t^2 &= \frac{1}{2} \left\{ (1 + \text{sign}(X_{t-} - b(t))) dX_t + (1 - \text{sign}(X_{t-} - b(t))) db(t) - dL_t^b \right\} \\ &\quad + [1_{\{X_{t-} > b(t)\}} (X_t - b(t))^- + 1_{\{X_{t-} \leq b(t)\}} (X_t - b(t))^+]. \end{aligned} \quad (A.4)$$

It follows from the dynamics of Z^i , $i \in \{1, 2\}$ that

$$d\langle Z^i, Z^i \rangle_t^c = \left(1_{\{X_{t-} < b(t)\}} + \frac{1}{4} 1_{\{X_{t-} = b(t)\}} \right) d\langle X, X \rangle_t^c = 1_{\{X_{t-} < b(t)\}} d\langle X, X \rangle_t^c, \quad (A.5)$$

where the second equality follows from the occupation density formula, see e.g. Corollary 1 on page 219 of [16]. Applying the classical Itô's formula to $F^1(Z_t^1, t)$ and $F^2(Z_t^2, t)$ and using the dynamics of Z^1 and

Z^2 , we get

$$\begin{aligned}
F^1(Z_t^1, t) &= F^1(Z_0^1, 0) + \int_0^t F_t^1(Z_{s-}^1, s) ds + \int_0^t F_x^1(Z_{s-}^1, s) dZ_s^1 + \frac{1}{2} \int_0^t F_{xx}^1(s, Z_{s-}^1) d\langle Z^1, Z^1 \rangle_s^c \\
&+ \sum_{0 \leq s \leq t} [F^1(Z_s^1, s) - F^1(Z_{s-}^1, s) - \Delta Z_s^1 F_x^1(Z_{s-}^1, s)] \\
&= F^1(Z_0^1, 0) + \int_0^t F_t^1(Z_{s-}^1, s) ds + \frac{1}{2} \int_0^t (1 - \text{sign}(X_{s-} - b(s))) F_x^1(Z_{s-}^1, s) dX_s \\
&+ \frac{1}{2} \int_0^t (1 + \text{sign}(X_{s-} - b(s))) F_x^1(Z_{s-}^1, s) db(s) \\
&- \sum_{0 < s \leq t} [1_{\{X_{s-} > b(s)\}} (X_s - b(s))^- + 1_{\{X_s \leq b(s)\}} (X_s - b(s))^+] F_x^1(Z_{s-}^1, s) \\
&- \frac{1}{2} \int_0^t F_x^1(Z_{s-}^1, s) dL_t^b + \frac{1}{2} \int_0^t 1_{\{X_{s-} < b(s)\}} F_{xx}^1(Z_{s-}^1, s) d\langle X^c, X^c \rangle_s \\
&\sum_{0 < s \leq t} [F^1(Z_s^1, s) - F(Z_{s-}^1, s) - \Delta Z_s^1 F_x^1(Z_{s-}^1, s)], \quad \text{and}
\end{aligned} \tag{A.6}$$

$$\begin{aligned}
F^2(Z_t^2, t) &= F^2(Z_0^2, 0) + \int_0^t F_t^2(Z_{s-}^2, s) ds + \int_0^t F_x^2(Z_{s-}^2, s) dZ_s^2 + \frac{1}{2} \int_0^t F_{xx}^2(s, Z_{s-}^2) d\langle Z^2, Z^2 \rangle_s^c \\
&+ \sum_{0 \leq s \leq t} [F^2(Z_s^2, s) - F^2(Z_{s-}^2, s) - \Delta Z_s^2 F_x^2(Z_{s-}^2, s)] \\
&= F^2(Z_0^2, 0) + \int_0^t F_t^2(Z_{s-}^2, s) ds + \frac{1}{2} \int_0^t (1 + \text{sign}(X_{s-} - b(s))) F_x^2(Z_{s-}^2, s) dX_s \\
&+ \frac{1}{2} \int_0^t (1 - \text{sign}(X_{s-} - b(s))) F_x^2(Z_{s-}^2, s) db(s) \\
&+ \sum_{0 < s \leq t} [1_{\{X_{s-} > b(s)\}} (X_s - b(s))^- + 1_{\{X_s \leq b(s)\}} (X_s - b(s))^+] F_x^2(Z_{s-}^2, s) \\
&- \frac{1}{2} \int_0^t F_x^2(Z_{s-}^2, s) dL_t^b + \frac{1}{2} \int_0^t 1_{\{X_{s-} < b(s)\}} F_{xx}^2(Z_{s-}^2, s) d\langle X^c, X^c \rangle_s \\
&+ \sum_{0 < s \leq t} [F^2(Z_s^2, s) - F(Z_{s-}^2, s) - \Delta Z_s^2 F_x^2(Z_{s-}^2, s)].
\end{aligned} \tag{A.7}$$

By splitting each term to their respective values on the sets $\{X_{s-} < b(s)\}$, $\{X_{s-} = b(s)\}$ and $\{X_{s-} > b(s)\}$, it can be seen that the following four equations are satisfied:

$$F^1(Z_0^1, 0) + F^2(Z_0^2, 0) = F(X_0, 0) + F(b(0), 0), \tag{A.8}$$

$$\begin{aligned}
&\int_0^t F_t^1(Z_{s-}, s) ds + \int_0^t F_t^2(Z_{s-}^2, s) ds = \frac{1}{2} \int_0^t F_t(X_{s-}, s) + F_t(X_{s-}, s) ds \\
&+ \int_0^t \left[F_t(b(s)+, s) 1_{\{X_{s-} < b(s)\}} + \frac{1}{2} (F_t(b(s)-, s) + F_t(b(s)+, s)) 1_{\{X_{s-} = b(s)\}} + F_t(b(s)-, s) 1_{\{X_{s-} > b(s)\}} \right] ds,
\end{aligned} \tag{A.9}$$

$$\begin{aligned} & \frac{1}{2} \int_0^t (1 - \text{sign}(X_{s-} - b(s))) F_x^1(Z_{s-}^1, s) dX_s + \frac{1}{2} \int_0^t (1 + \text{sign}(X_{s-} - b(s))) F_x^2(Z_{s-}^2, s) dX_s \\ &= \frac{1}{2} \int_0^t [F_x(X_{s-}, s) + F_x(X_{s-}, s)] dX_s, \end{aligned} \quad (\text{A.10})$$

$$\begin{aligned} & \frac{1}{2} \int_0^t (1 + \text{sign}(X_{s-} - b(s))) F_x^1(Z_{s-}^1, s) db(s) + \frac{1}{2} \int_0^t (1 - \text{sign}(X_{s-} - b(s))) F_x^2(Z_{s-}^2, s) db(s) = \\ & \int_0^t \left[F_x(b(s)+, s) 1_{\{X_{s-} < b(s)\}} + \frac{1}{2} [F_x(b(s)+, s) + F_x(b(s)-, s)] 1_{\{X_{s-} = b(s)\}} + F_x(b(s)-, s) 1_{\{X_{s-} > b(s)\}} \right] db(s). \end{aligned} \quad (\text{A.11})$$

On the other hand, (3.15) of [12] still holds:

$$\begin{aligned} F(b(t), t) &= F(b(0), 0) + \int_0^t \left[F_t(b(s)+, s) 1_{\{X_{s-} < b(s)\}} + \frac{1}{2} [F_t(b(s)-, s) + F_t(b(s)+, s) 1_{\{X_{s-} = b(s)\}}] \right. \\ & \quad \left. + F_t(b(s)-, s) 1_{\{X_{s-} > b(s)\}} \right] ds \\ & \int_0^t \left[F_x(b(s)+, s) 1_{\{X_{s-} < b(s)\}} + \frac{1}{2} [F_x(b(s)-, s) + F_x(b(s)+, s) 1_{\{X_{s-} = b(s)\}}] \right. \\ & \quad \left. + F_x(b(s)-, s) 1_{\{X_{s-} > b(s)\}} \right] db(s), \end{aligned} \quad (\text{A.12})$$

whose proof is carried out by using the uniqueness of finite measures on p-systems.

Let us analyze the jump terms in (A.6) and (A.7). We will denote

$$\begin{aligned} A &:= - [1_{\{X_{s-} > b(s)\}} (X_s - b(s))^- + 1_{\{X_s \leq b(s)\}} (X_s - b(s))^+] F_x^1(Z_{s-}^1, s) \\ & \quad + [F^1(Z_s^1, s) - F(Z_{s-}^1, s) - \Delta Z_s^1 F_x^1(Z_{s-}^1, s)], \end{aligned} \quad (\text{A.13})$$

$$\begin{aligned} B &:= [1_{\{X_{s-} > b(s)\}} (X_s - b(s))^- + 1_{\{X_s \leq b(s)\}} (X_s - b(s))^+] F_x^2(Z_{s-}^2, s) \\ & \quad + \sum_{0 < s \leq t} [F^2(Z_s^2, s) - F(Z_{s-}^2, s) - \Delta Z_s^2 F_x^2(Z_{s-}^2, s)]. \end{aligned} \quad (\text{A.14})$$

Depending on the whereabouts of X_{s-} and X_s with respect to the boundary curve b , A and B take four different values:

(1) $X_{s-} > b(s)$ and $X_t \geq b(t)$. In this case

$$A = 0, \quad B = F^2(X_s, s) - F^2(X_{s-}, s) - \Delta X_s F_x^2(X_{s-}, s), \quad (\text{A.15})$$

and

$$A + B = F(X_s, s) - F(X_{s-}, s) - \Delta X_s F_x(X_{s-}, s). \quad (\text{A.16})$$

(2) $X_{s-} > b(s)$ and $X_s < b(s)$. In this case

$$\begin{aligned} A &= -(b(s) - X_s) F_x^1(b(s), s) + F^1(X_s, s) - F^1(b(s), s) - (X_s - b(s)) F_x^1(b(s), s) \\ &= F^1(X_s, s) - F^1(b(s), s), \end{aligned} \quad (\text{A.17})$$

$$\begin{aligned} B &= (b(s) - X_s) F_x^2(b(s), s) + F^2(b(s), s) - F^2(X_{s-}, s) - (b(s) - X_{s-}) F_x^2(X_{s-}, s) \\ &= F^2(b(s), s) - F^2(X_{s-}, s) - \Delta X_s F_x^2(X_{s-}, s), \quad \text{and} \end{aligned} \quad (\text{A.18})$$

$$A + B = F(X_s, s) - F(X_{s-}, s) - \Delta X_s F_x(X_s, s). \quad (\text{A.19})$$

(3) $X_{s-} \leq b(s)$ and $X_s \geq b(s)$. We have that

$$\begin{aligned} A &= -(X_s - b(s))F_x^1(X_{s-}, s) + F^1(b(s), s) - F^1(X_{s-}, s) - (b(s) - X_{s-})F_x^1(X_{s-}, s) \\ &= F^1(b(s), s) - F^1(X_{s-}, s) - \Delta X_s F^1(X_{s-}, s), \end{aligned} \quad (\text{A.20})$$

$$\begin{aligned} B &= (X_s - b(s))F_x^2(b(s), s) + F^2(X_s, s) - F^2(b(s), s) - (X_s - b(s))F_x^2(b(s), s) \\ &= F^2(X_s, s) - F^2(b(s), s). \end{aligned} \quad (\text{A.21})$$

As a result

$$A + B = F(X_s, s) - F(X_{s-}, s) - \Delta X_s F_x(X_{s-}, s). \quad (\text{A.22})$$

(4) $X_{s-} \leq b(s)$ and $X_s < b(s)$. Clearly,

$$A = F^1(X_s, s) - F^1(X_{s-}, s) - \Delta X_s F_x^1(X_{s-}, s) \quad \text{and} \quad B = 0. \quad (\text{A.23})$$

As a result

$$A + B = F(X_s, s) - F(X_{s-}, s) - \Delta X_s F_x(X_{s-}, s). \quad (\text{A.24})$$

Now combining (A.1), (A.5), (A.6), (A.7), (A.8), (A.9), (A.10), (A.11), (A.12), (A.16), (A.19), (A.22), (A.24), we obtain

$$\begin{aligned} F(X_t, t) &= F(X_0, 0) + \frac{1}{2} \int_0^t [F_t(X_{s-}, +, s) + F_t(X_{s-}, -, s)] ds \\ &\quad + \frac{1}{2} \int_0^t [F_x(X_{s-}, +, s) + F_x(X_{s-}, -, s)] dX_s + \frac{1}{2} \int_0^t 1_{\{X_{s-} \leq b(s)\}} F_{xx}(s, X_{s-}) d\langle X, X \rangle_{s-}^c \\ &\quad + \sum_{0 < s \leq t} \left[F(X_s, s) - F(X_{s-}, s) - \Delta X_s F_x(X_{s-}, -, s) 1_{\{X_{s-} \leq b(s)\}} - \Delta X_s F_x(s, X_{s-}, +) 1_{\{X_{s-} > b(s)\}} \right] \\ &\quad + \frac{1}{2} \int_0^t [F_x^2(Z_{s-}^2, s) - F^1(Z_{s-}^1, s)] dL_t^b. \end{aligned} \quad (\text{A.25})$$

The last term on the right-hand-side of (A.25) can be written as

$$\frac{1}{2} \int_0^t [F_x^2(Z_{s-}^2, s) - F^1(Z_{s-}^1, s)] dL_t^b = \frac{1}{2} \int_0^t [F_x(X_{s-}, +, s) - F_x(X_{s-}, -, s)] 1_{\{X_{s-} = b(s)\}} dL_t^b, \quad (\text{A.26})$$

using Theorem 69 of [16]. On the other hand, the jump term in (A.25) can be written as

$$\begin{aligned} &\sum_{0 < s \leq t} \left[F(X_s, s) - F(X_{s-}, s) - \Delta X_s F_x(X_{s-}, -, s) 1_{\{X_{s-} \leq b(s)\}} - \Delta X_s F_x(s, X_{s-}, +) 1_{\{X_{s-} > b(s)\}} \right] \\ &= \sum_{0 < s \leq t} \left[F(X_s, s) - F(X_{s-}, s) - \frac{1}{2} \Delta X_s [F_x(X_{s-}, -, s) + F_x(X_{s-}, +, s)] \right]. \end{aligned} \quad (\text{A.27})$$

This completes the proof. \square

REFERENCES

- [1] Larbi Alili and Andreas E. Kyprianou. Some remarks on first passage of Lévy processes, the American put and pasting principles. *Ann. Appl. Probab.*, 15(3):2062–2080, 2005.
- [2] Luis H. R. Alvarez. Solving optimal stopping problems of linear diffusions by applying convolution approximations. *Math. Methods Oper. Res.*, 53(1):89–99, 2001.
- [3] Erhan Bayraktar. On the perpetual American put options for level dependent volatility models with jumps. Technical report, University of Michigan, 2008. Available at <http://arxiv.org/pdf/math/0703538>.

- [4] Erhan Bayraktar, Savas Dayanik, and Ioannis Karatzas. Adaptive Poisson disorder problem. *Ann. Appl. Probab.*, 16(3):1190–1261, 2006.
- [5] Bruno Bouchard, Nicole El Karoui, and Nizar Touzi. Maturity randomization for stochastic control problems. *Ann. Appl. Probab.*, 15(4):2575–2605, 2005.
- [6] Peter Carr. Randomization and the American put. *Review of Financial Studies*, 11 (3):597–626, 1998.
- [7] Avner Friedman. *Partial differential equations of parabolic type*. Prentice-Hall Inc., Englewood Cliffs, N.J., 1964.
- [8] Avner Friedman. *Stochastic differential equations and applications*. Dover Publications Inc., Mineola, NY, 2006. Two volumes bound as one, Reprint of the 1975 and 1976 original published in two volumes.
- [9] Hitoshi Ishii. On uniqueness and existence of viscosity solutions of fully nonlinear second-order elliptic PDEs. *Comm. Pure Appl. Math.*, 42(1):15–45, 1989.
- [10] Ioannis Karatzas and Steven E. Shreve. *Methods of mathematical finance*, volume 39 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, 1998.
- [11] Ernesto Mordecki and Paavo Salminen. Optimal stopping of Hunt and Lévy processes. *Stochastics*, 79(3-4):233–251, 2007.
- [12] Goran Peskir. A change-of-variable formula with local time on curves. *J. Theoret. Probab.*, 18(3):499–535, 2005.
- [13] Goran Peskir and Albert Shiryaev. *Optimal stopping and free-boundary problems*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2006.
- [14] Huyên Pham. Optimal stopping, free boundary, and American option in a jump-diffusion model. *Appl. Math. Optim.*, 35(2):145–164, 1997.
- [15] Huyên Pham. Optimal stopping of controlled jump diffusion processes: a viscosity solution approach. *J. Math. Systems Estim. Control*, 8(1):27 pp. (electronic), 1998.
- [16] Philip E. Protter. *Stochastic integration and differential equations*, volume 21 of *Stochastic Modelling and Applied Probability*. Springer-Verlag, Berlin, 2005. Second edition. Version 2.1, Corrected third printing.
- [17] R. Tyrrell Rockafellar. *Convex analysis*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1997. Reprint of the 1970 original, Princeton Paperbacks.
- [18] Paul Wilmott, Sam Howison, and Jeff Dewynne. *The mathematics of financial derivatives*. Cambridge University Press, Cambridge, 1995. A student introduction.

(E. Bayraktar) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109
E-mail address: erhan@umich.edu