

A Proof of the Smoothness of the Finite Time Horizon American Put Option for Jump Diffusions

Y

Erhan Bayraktar^{zx}

Abstract

We give a new proof of the fact that the value function of the finite time horizon American put option for a jump diffusion, when the jumps are from a compound Poisson process, is the classical solution of a quasi-variational inequality and it is C^1 across the optimal stopping boundary. Our proof only uses the classical theory of parabolic partial differential equations of Friedman (1964, 2006) and does not use the theory of viscosity solutions, since our proof relies on constructing a sequence of functions, each of which is a value function of an optimal stopping time for a diffusion. The sequence is constructed by iterating a functional operator that maps a certain class of convex functions to smooth functions satisfying variational inequalities (or to value functions of optimal stopping problems involving only a diffusion). The approximating sequence converges to the value function exponentially fast, therefore it constitutes a good approximation scheme, since the optimal stopping problems for diffusions can be readily solved. Our technique also lets one see why the jump-diffusion control problems may be smoother than the control problems with piece-wise deterministic Markov processes: In the former case the sequence of functions that converge to the value function is a sequence of value functions of control problems for diffusions, and in the latter case the converging sequence is a sequence of the value functions of deterministic optimal control problems. The first of these sequences is known to be smoother than the second one.

1 Introduction

Let $(\Omega; \mathcal{F}; P)$ be a complete probability space hosting a Wiener process $W = (W_t; t \geq 0)$ and a Poisson random measure on $\mathbb{R}_+ \times \mathbb{R}_+$ with mean measure $(dx)dt$ (in which dx is a probability measure on \mathbb{R}_+) independent of the Wiener process. We will consider a Markov process $S = (S_t; t \geq 0)$ of the form

$$dS_t = -S_t dt + S_t dW_t + S_t \int_{\mathbb{R}_+} (z - 1) N(dt; dz) \quad (1.1)$$

AMS 2000 Subject Classification. 60G40, 62L15, 60J75.

^yKey Words. Optimal stopping, Markov Processes, Jump Diffusions, American Options, Integro-Differential Equations, Parabolic Variational Inequalities

^zE. Bayraktar is with the Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA, email: erhan@umich.edu

^xThis research is partially supported by NSF Research Grant, DMS-0604491.

We will take $\sigma = r + \frac{1}{2}\sigma^2$, in which

$$\int_{\mathbb{R}_+} x v(dx) < 1; \quad (1.2)$$

(a standing assumption). Here, S is the price of a security/stock and the dynamics in (1.1) are stated under a risk neutral measure. The constant $r \geq 0$ is the interest rate, and the constant $\sigma > 0$ is the volatility.

The value function of the American put option pricing problem is

$$V(x; T) := \sup_{\tau \in \mathcal{S}_{0,T}} E^x [e^{-r\tau} (K - S_\tau)^+]; \quad (1.3)$$

in which $\mathcal{S}_{0,T}$ is the set of stopping times (of the filtration generated by W and N) taking values in $[0; T]$, and E^x is the expectation under the probability measure P , given that $S_0 = x$.

We will show that V is the classical solution of a quasi-variational inequality and that it satisfies smooth fit principle across the optimal stopping boundary, i.e., $V(\cdot; \cdot)$ is continuously differentiable with respect to its first variable at the optimal stopping boundary. We argue this by showing that $V(\cdot; \cdot)$ is the fixed point of an operator, which we will denote by J , that maps functions to the value functions of optimal stopping problems for dividends. As soon as a given function, $f(\cdot; \cdot)$ has some regularity properties, we show that $Jf(\cdot; \cdot)$ is the unique (classical) solution of a quasi-variational inequality and satisfies the smooth fit principle (it is C^1 across the optimal stopping boundary). We show that V has these regularity properties by making use a sequence (iterating J starting at the pay-off function of the put option) that converges to V uniformly and exponentially fast. This incidentally gives a numerical procedure, whose accuracy versus speed characteristics can be controlled, since the optimal stopping problems for dividends can be solved readily using the several numerical procedures that are available (e.g. Monte-Carlo methods of Longstaff and Schwartz (2001), Glasermann and Yu (2004) or Finite Difference Methods in Williams et al. (1995) or Kusner and Dupuis (2003)).

Pham (1997) provided a proof that the value function V is the classical solution of a quasi-variational inequality using a combination of Friedman (2006) and the theory of viscosity solutions. See Proposition 3.1 in Pham (1997) which carries out its proof (details are not provided but hinted) using similar line of arguments to the proof of Proposition 5.3 in Pham (1998), which uses the uniqueness results for viscosity solutions, see e.g. Ishii (1989). Our proof does not make use the theory of viscosity solutions and relies on constructing a sequence of functions, each of which is a value function of an optimal stopping time for a dividend. This technique provides an intuitive explanation as to why the jump-dividend optimal stopping problem may be smoother than the control problems with piece-wise deterministic Markov processes. The failure of the smooth fit principle was observed for example in optimal stopping problem for multi-dimensional piece-wise deterministic processes in Bayraktar et al. (2006). In the case of optimal stopping problems for jump dividends the sequence of functions that converge to the value function is a sequence of value function of control problems for dividends, and in the case of optimal stopping problems for piece-wise deterministic processes the converging sequence is a sequence of the value functions of deterministic optimal control problems. The first of these two sequences is known to be smoother. Also, as we mentioned before, our proof procedure gives an accurate numerical procedure as an added benefit.

The finite horizon American put option for jump dividends were analyzed in Bayraktar (2007) by similar means. The main difficulty in this paper stems from the fact that we are

dealing with two dimensional parabolic variational inequalities. For example, we make use of Friedman (2006), Karatzas and Shreve (1998) (Chapter 2) (also see Peskir and Shiryaev (2006) (Chapter 7)) to study the properties of the operator J , and we show that the approximating sequence of functions are bounded with respect to the Hölder semi-norm (see Friedman (1964), page 61) to argue that the limit of the approximating sequence (which is a fixed point of J) solves a partial differential equation in a bounded parabolic domain, etc. Alili and Kyprianou (2005) and Mordecki and Salminen (2006) considered the smooth fit principle (the fact that the value function is C^1 across the optimal stopping boundary) for one-dimensional exponential Levy processes using fluctuation theory. See Bayraktar (2007) for a detailed discussion.

We represent the value function of the American put option for a jump diffusion process as a limit of a sequence of optimal stopping problems for a diffusion (by taking the horizon of the problem to be the times of jumps of the compound Poisson process). Somewhat similar, approximation techniques were used to solve optimal stopping problems for diffusions (not jump diffusions), see e.g. Alvarez (2004) for perpetual optimal stopping problems with non-smooth pay-off functions, and Carr (1998) and Bouchard et al. (2005) for finite time horizon American put option pricing problems.

The next two sections prepare the proof of our main result Theorem 3.1 in a sequence of Lemmata and Corollaries. Our contribution is not the result itself but our proof of it. In the next section, we introduce the functional operator J , that maps a function to the value function of an optimal stopping time for a diffusion, and analyze its properties: for e.g. it preserves convexity with respect to the first variable, the increase in the Hölder semi-norm of the function can be controlled, it maps certain class of functions to the classical solutions of quasi-variational inequalities. In Section 3, we construct a sequence of functions that converge to the value function uniformly and exponentially to the smallest fixed point of the operator J that is greater than the pay-off of the put option. We show that the sequence has bounded in the Hölder norm, convex with respect to the first variable, etc using results of Section 2. We eventually arrive at the fact that the fixed point of the operator we consider is the value function itself and is the classical solution of the quasi-variational inequalities and satisfies smooth fit principle.

2 A Functional Operator and Its Properties

Let us define an operator J through its action on a test function $f : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$, as the value function of the following optimal stopping problem

$$Jf(x; T) = \sup_{\tau \in \mathcal{S}_{0,T}} E^x \int_0^{\tau} e^{-(x+\tau)t} P f(S_t^0; T-t) dt + e^{-(x+\tau)\tau} (K - S_\tau^0)^+; \quad (2.1)$$

in which

$$P f(x; T-t) = \int_{\mathbb{R}_+} f(xz; T-t) (dz); \quad x \geq 0; \quad (2.2)$$

We will extend $T \mapsto Jf(x; T)$ onto $[0; 1]$ by letting

$$Jf(x; 1) = \lim_{T \rightarrow 1} Jf(x; T); \quad (2.3)$$

Here, $S_t^0 = f(S_t^0; t) = 0$ is the solution of

$$dS_t^0 = S_t^0 dt + S_t^0 dW_t; \quad S_0^0 = x; \quad (2.4)$$

whose infinitesimal is given by

$$A = \frac{1}{2} x^2 \frac{d^2}{dx^2} + x \frac{d}{dx}; \quad (2.5)$$

and $S_{[0;T]}$ is the set of stopping times of S^0 which take values in $[0;T]$. We will denote

$$S_t^0 = xH_t; \quad (2.6)$$

where

$$H_t = \exp \left(\frac{1}{2} t + W_t \right); \quad (2.7)$$

The next remark characterizes the optimal stopping times of (2.1) using the Snell envelope theory.

Remark 2.1 Let us denote

$$Y_t = \int_0^{Z_t} e^{(x^+)^s} P f(S_t^0; T-s) ds + e^{(x^+)^t} (K - S_t^0)^+; \quad (2.8)$$

Using the strong Markov property of S_0 we can determine the Snell envelope of Y as

$$t \mapsto \sup_{S_t; T} E f Y | \mathcal{F}_t g = e^{(x^+)^t} J f(S_t^0; T-t) + \int_0^{Z_t} e^{(x^+)^s} P f(S_s^0; T-s) ds; \quad t \in [0; T]; \quad (2.9)$$

in which $S_{t;T}$ is the set of stopping times taking value in $[t;T]$. Then using Theorem D.12 in Karatzas and Shreve (1998), the stopping time

$$x \mapsto \inf_{t \in [0;T]} J f(S_t^0; T-t) = (K - S_T^0)^+; \quad (2.10)$$

satisfies

$$J f(x; T) = E^x \int_0^Z e^{(x^+)^t} P f(S_t^0; T-t) dt + e^{(x^+)^x} (K - S_x^0)^+; \quad (2.11)$$

The second infimum in (2.10) is less than T because $J f(S_T^0; 0) = (K - S_T^0)^+$.

When f is bounded it follows from the bounded convergence theorem that (using the results of Bayraktar (2007) and arguments similar to the ones used in Corollary 7.3 in Chapter 2 of Karatzas and Shreve (1998))

$$J f(x; 1) = \sup_{S_0; 1} E^x \int_0^Z e^{(x^+)^t} P f(S_t^0; 1-t) dt + e^{(x^+)^x} (K - S_x^0)^+; \quad (2.12)$$

which is the value function of the infinite horizon version of (2.1).

The next lemma on the monotonicity properties of $J f(\cdot; \cdot)$ immediately follows from 2.4).

Lemma 2.1 Let $T \mapsto f(x; T)$ be non-decreasing, and $x \mapsto f(x; T)$ be non-increasing. Then the functions $T \mapsto J f(x; T)$ is non-decreasing and $x \mapsto J f(x; T)$ is non-increasing.

As we shall see next, the operator J in (2.1) preserves convexity (with respect to the first variable).

Lemma 2.2 Let J be as in (2.1). Then if $f : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a convex function in its first variable, then so is $Jf : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$.

Proof: Note that if $f(\cdot; T-t)$ is convex, so is $Pf(\cdot; T-t)$. Because $\int_0^t e^{-(r+x)s} ds = xH_t$ is linear in x , it follows that $x \mapsto Pf(xH_t; T-t)$ is a convex function of x for all $t \in \mathbb{R}_+$. Therefore, the integral in (2.1) is also convex in x . Also note that $(K - xH)^+$ is also convex in x . Since the upper envelope (supremum) of convex functions is convexity $Jf : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ follows.

Remark 2.2 Since $x = 0$ is an absorbing boundary for the process S^0 , for any $f : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$

$$\begin{aligned} Jf(0; T) &= \sup_{t \in [0, T]} \int_0^t e^{-(r+x)s} f(0; T-s) ds + e^{-(r+x)T} K \\ &= \max_{0 \leq t \leq T} \int_0^t e^{-(r+x)s} f(0; T-s) ds + e^{-(r+x)T} K, \quad T \geq 0. \end{aligned} \quad (2.13)$$

If we further assume $f(\cdot; \cdot) \leq K$, then $Jf(0; T) = K$, $T \geq 0$. Moreover if f is convex with respect to its first variable, then

$$D_+^x Jf(\cdot; T) \leq 1; \quad \text{for all } T \in [0; 1]. \quad (2.14)$$

Moreover, $D_+^x Jf(x; T) \leq 1$, $x > 0$ and $T \in [0; 1]$. Here D_+^x and D_-^x denote the right-derivative and left-derivative operators with respect to the first variable.

The next two lemmas are very crucial for our proof of the smoothness of the American option price for jump diffusions, shows that the increase in the Hölder seminorm that the operator J causes can be controlled.

Lemma 2.3 Let us assume that $f : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is convex in its first variable and $\|f\|_1 < K$. Then $x \mapsto Jf(x; t)$ satisfies

$$|Jf(x; T) - Jf(y; T)| \leq |x - y|, \quad (x, y) \in \mathbb{R}_+ \times \mathbb{R}_+; \quad (2.15)$$

and all $T \geq 0$.

Proof: Observe that $x \mapsto Jf(x; t)$ is non-increasing (by Lemma 2.1), convex (Lemma 2.2), and satisfies

$$Jf(x; T) \leq (K - x)^+ \quad Jf(0; T) = K; \quad (2.16)$$

This implies that the left and right derivatives satisfy

$$-1 \leq D_-^x Jf(x; T) \leq D_+^x Jf(x; T) \leq 0; \quad x > 0; T \geq 0. \quad (2.17)$$

Now, (2.15) follows from Theorem 24.7 (on page 237) in Rockafellar (1997) (especially the proof of the second statement of the theorem).

Remark 2.3 Let $T_0 \in (0; 1)$ and denote the value function of the American put option (for the diffusion S^0) by

$$F(x; T) = \sup_{2S_0, T}^n E e^{-(r+)^t} (K - xH_t)^+; \quad x \in \mathbb{R}_+; T \in [0; T_0]; \quad (2.18)$$

Then for $S = T = T_0$

$$F(x; T) = F(x; S) + C \mathbb{P}(S_j^{1=2}); \quad (2.19)$$

for all $x \in \mathbb{R}_+$ and for some C that depends only on T_0 . See e.g. equation (2.4) in Pham (1997).

Lemma 2.4 Let us assume that for some $L \in (0; 1)$

$$|f(x; T) - f(x; S)| \leq L \mathbb{P}(S_j^{1=2}); \quad (T; S) \in \mathbb{S}_0; T_0]^2; \quad (2.20)$$

for all $x \in \mathbb{R}_+$, for $0 \leq S_0 < T_0 < 1$. Then

$$|Jf(x; T) - Jf(x; S)| \leq (aL + C) \mathbb{P}(S_j^{1=2}); \quad (T; S) \in \mathbb{S}_0; T_0]^2; \quad (2.21)$$

for some $a \in (0; 1)$, whenever,

$$\mathbb{P}(S_j) < \frac{r}{r+} \frac{L}{K}^2; \quad (2.22)$$

Here, $C \in (0; 1)$ is as in Remark 2.3.

Proof: Without loss of generality we will assume that $T > S$ (and use Lemma 2.1).

$$\begin{aligned} Jf(x; T) - Jf(x; S) &= \sup_{2S_0, T}^n E \int_0^T e^{-(r+)^t} (Pf(xH_t; T) - Pf(xH_t; S)) dt + e^{-(r+)^T} (K - xH_T)^+ \\ &\quad - \sup_{2S_0, S}^n E \int_0^S e^{-(r+)^t} (Pf(xH_t; T) - Pf(xH_t; S)) dt + e^{-(r+)^S} (K - xH_S)^+ \\ &= \sup_{2S_0, T}^n E \int_0^T e^{-(r+)^t} (Pf(xH_t; T) - Pf(xH_t; S)) dt \\ &\quad + 1_{FS < g_S} \int_S^T e^{-(r+)^t} (Pf(xH_t; S) - Pf(xH_t; T)) dt + e^{-(r+)^T} (K - xH_T)^+ - e^{-(r+)^S} (K - xH_S)^+ \\ &\quad + \frac{L}{r+} (T - S)^{1=2} + \frac{K}{r+} e^{-(r+)^S} e^{-(r+)^T} \\ &\quad + \sup_{2S_S, T}^n E e^{-(r+)^t} (K - xH_t)^+ - \sup_{2S_S, T}^n E e^{-(r+)^t} (K - xH_t)^+ \\ &\quad + \frac{L}{r+} (T - S)^{1=2} + K (T - S) + e^{-(r+)^S} (F(H_S; T - S) - F(H_S; 0)); \\ &\quad + \frac{L + C}{r+} (T - S)^{1=2} + K (T - S) \end{aligned} \quad (2.23)$$

in which $F(\cdot; \cdot)$ is given by (2.18). To derive the second inequality, we used the fact that

$$\int_{\mathbb{R}_+} |Pf(xH_t; T) - Pf(xH_t; S)| (dz) \mathbb{P}(xH_t; T) - \mathbb{P}(xH_t; S) \leq L \mathbb{P}(S_j^{1=2}); \quad (2.24)$$

which follows from the assumption in (2.20); and that

$$E \int_0^{\tau_{fS} \wedge g} e^{-(r+)^t} P f(x; H_t; S_t) dt = E \int_0^{\tau_T} e^{-(r+)^t} dt K - \frac{K}{r+ + K} e^{-(r+)^S} - e^{-(r+)^T} \quad (2.25)$$

and also that

$$e^{-(r+)^S} - e^{-(r+)^T} = e^{-(r+)^S} (r+)^S (T - S) - (r+)^T (T - S): \quad (2.26)$$

The second equality in (2.23), on the other hand, follows from the strong Markov property of S^0 and the form of the optimal stopping times in (2.10) (i.e. the fact that the optimal stopping times are hitting times is used here).

Now, observe that since T and S satisfy (2.22) we obtain (2.21) using (2.23) and (2.19).

Recalling Remark 2.1, let us define the continuation region and its sections by

$$C_T^{Jf} := \{x \in (0; 1) : Jf(x; T) > (K - x)^+ g\}; \text{ and } C_T^{Jf} := \{x \in (0; 1) : Jf(T; x) > (K - x)^+ g\}; \quad (2.27)$$

$T > 0$, respectively.

Lemma 2.5 Suppose that $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is such that $x \mapsto f(x; T)$ is a positive convex function, $T \mapsto f(x; T)$ is non-decreasing, and $\|f\|_1 \leq K$. Moreover f satisfies, $D_+^x f(x; 1) \leq 1$, for all $x \in \mathbb{R}_+$. For every $T > 0$, there exists $c^f(T) \in (0; K)$ such that $C_T^{Jf} = (c^f(T); 1)$.

Moreover, $T \mapsto c^f(T)$ is non-increasing.

Proof: Let us first show that if $x \leq K$, then $x \in C_T^{Jf}$, for all $T \geq 0$. Let

$$\mu := \inf_{0 \leq t \leq T} S_t^0 \leq K - \mu g: \quad (2.28)$$

Since

$$Pf(0) < \mu < Tg > 0 \quad \text{for } x \leq K; \quad (2.29)$$

for all $T > 0$, we have that

$$E^x \int_0^{\tau_{\mu}} e^{-(r+)^t} P f(S_t^0; T - t) dt + e^{-(r+)^{\tau_{\mu}}} (K - S_{\tau_{\mu}}^0)^+ > 0; \quad (2.30)$$

which implies that $x \in C_T^{Jf}$.

On the other hand, it is clear that

$$(K - x)^+ \leq Jf(x; T) - Jf(x; 1); \quad (x; T) \in \mathbb{R}_+ \times \mathbb{R}_+: \quad (2.31)$$

Recall (2.12), which shows that $Jf(\cdot; 1)$ is the value function of an infinite horizon problem. It was shown in Bayraktar (2007) (see Corollary 2.4) under the condition on the right derivative of f that there exist $\bar{1}^f \in (0; K)$ such that

$$Jf(x; 1) = (K - x)^+; \quad x \in [0; \bar{1}^f]; \quad Jf(x; 1) > (K - x)^+; \quad x \in (\bar{1}^f; 1): \quad (2.32)$$

Since, $x \mapsto Jf(x; 1)$ and $x \mapsto Jf(x; T)$, $T \geq 0$, are convex functions (from Lemma 2.2 in Bayraktar (2007) and Lemma 2.2 respectively), (2.30), (2.31) and (2.32) imply that there exists a point $c^f(T) \in (1^f; K)$ such that

$$Jf(x) = (K - x)^+; x \in [0; c^f(T)]; \quad Jf(x; T) > (K - x)^+; x \in (c^f(T); 1); \quad (2.33)$$

for $T > 0$. The first statement of the Lemma immediately follows from (2.33).

The fact that $T \mapsto c(T)$ is non-increasing follows from the fact that $T \mapsto Jf(x; T)$ is non-decreasing.

We will argue in the following lemma that, if $f(\cdot; \cdot)$ has certain regularity properties, then $Jf(\cdot; \cdot)$ is the classical solution of a parabolic variational inequality.

Lemma 2.6 Let us assume that $f: R_+ \times R_+ \rightarrow R_+$ is convex in its first variable, $\|f\|_1 \leq K$ and $D_+^x f(\cdot; 1) \leq 1$, $T \geq 0$. Moreover, $T \mapsto f(x; T)$ is non-increasing, Hölder continuous uniformly in T on compact subsets of R_+ . Then the function $Jf: R_+ \times R_+ \rightarrow R_+$ (the restriction of Jf onto R_+^2) is the unique solution of

$$Au(x; T) - (r + \delta)u(x; T) + (Pf)(x; T) - \frac{\partial}{\partial T}u(x; T) = 0 \quad x > c^f(T); \quad (2.34)$$

$$u(x; T) = (K - x) - x - c^f(T); \quad (2.35)$$

in which A is as in (2.5) and c^f is as in Lemma 2.5.

Proof: Let us take a point in $(t; T) \in C^{Jf}$ (recall the notation in (2.27) and Lemma 2.5) and a rectangle $R = (t_1; t_2) \times (x_1; x_2)$, the parabolic boundary of R and consider parabolic partial differential equation (PDE)

$$Au(x; T) - (r + \delta)u(x; T) + (Pf)(x; T) - \frac{\partial}{\partial T}u(x; T) = 0 \quad \text{in } R; \quad (2.36)$$

$$u(x; T) = Jf(x; T) \quad \text{on } \partial R_0;$$

Since by Lemmas 2.3 and 2.4, $f(\cdot; \cdot)$ satisfies the uniform Lipschitz and Hölder continuity conditions, with respect to the first and second variables respectively, the function $Jf(\cdot; \cdot)$ is continuous. On the other hand

$$\begin{aligned} |Pf(x; T) - Pf(y; S)| &\leq |Pf(x; T) - Pf(x; S)| + |Pf(x; S) - Pf(y; S)| \\ &\leq \int_{R_+} (dz) (|f(xz; T) - f(xz; S)| + |f(xz; S) - f(yz; S)|) \\ &\leq L |T - S|^{\frac{1}{2}} + |x - y| \end{aligned} \quad (2.37)$$

in which L is a constant that depends on t_1 and t_2 and \cdot is as in (1.2). Then parabolic partial differential equation in (2.36) has a unique solution as a corollary of Theorem 5.2 in Friedman (2006). The rest of the proof follows as a result of optional sampling theorem using the same line of arguments as in the proof of Theorem 7.7 on page 73 of Karatzas and Shreve (1998).

Lemma 2.7 For a given $T > 0$, let $x \mapsto f(x; T)$ be a convex and non-increasing function. Then the convex function $x \mapsto Jf(x; T)$ is of class C^1 at $x = c(T)$, i.e.,

$$\frac{\partial}{\partial x} Jf(x; T) \Big|_{x=c(T)} = 1; \quad (2.38)$$

Proof: The proof is similar to the proof of Lemma 7.8 on page 74 of Karatzas and Shreve (1998), but we will provide it here for the sake of completeness. Let $x = c(T)$, then

$$\begin{aligned} Jf(x+; T) &= E \int_0^{x+} e^{-(x+)^+ t} P f((x+)^+ H_t; T - t) dt + e^{-(x+)^+ x+} (K - (x+)^+ H_{x+})^+ \\ &= E \int_0^{x+} e^{-(x+)^+ t} P f(x H_t; T - t) dt + (K - x H_{x+})^+ \\ &\quad + E \int_0^{x+} e^{-(x+)^+ t} [P f((x+)^+ H_t; T - t) - P f(x H_t; T - t)] dt \\ &\quad + E e^{-(x+)^+ x+} (K - (x+)^+ H_{x+})^+ - (K - x H_{x+})^+ \\ &\quad + E \int_{x+}^{\infty} 1_{f_{x+} < T} e^{-(x+)^+ x+} (K - (x+)^+ H_{x+}) - (K - x H_{x+}) \\ &\quad + E \int_{x+}^{\infty} 1_{f_{x+} = T} e^{-(x+)^+ x+} (K - (x+)^+ H_{x+})^+ - (K - x H_{x+})^+ \\ &= Jf(x; T) - E^x \int_0^{x+} 1_{f_{x+} < T} e^{-(x+)^+ x+} H_{x+} + E^x \int_{x+}^{\infty} 1_{f_{x+} = T} e^{-(x+)^+ T} H_T : \end{aligned} \quad (2.39)$$

The second inequality follows since $x+$ is not optimal when one starts at x and that $x \mapsto P f(x; T)$ is a decreasing function for any $T \geq 0$. From (2.39) it follows that

$$D_+^x Jf(x+; T) = 1; \quad (2.40)$$

since $e^{-(x+)^+ t} H_t$ is a uniformly integrable martingale and $x+ \neq 0$. Convexity of $Jf(t; x)$ (Lemma 2.2) implies that

$$1 = D^x Jf(x-; t) \leq D_+^x Jf(x+; t) = 1; \quad (2.41)$$

which yields the desired result.

The operator J in (2.1) preserves boundedness and order.

Lemma 2.8 Let $f: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a bounded function. Then Jf is also bounded. In fact,

$$0 \leq Jf \leq K + \frac{1}{r+} \|f\|_1; \quad (2.42)$$

Moreover, if $T \mapsto f(x; T)$ is non-decreasing, then so is $T \mapsto Jf(x; T)$.

Proof: The proof follows directly from the definition in (2.1).

Lemma 2.9 The operator J in (2.1) preserves order, i.e. whenever for any $f_1, f_2: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfy $f_1(x; T) \leq f_2(x; T)$, then $Jf_1(x; T) \leq Jf_2(x; T)$ for all $(x; T) \in \mathbb{R}_+ \times \mathbb{R}_+$.

Proof: The fact that J preserves order is evident from (2.1).

3 A Sequence of Value Functions of Optimal Stopping Problems for S^0 Approximating V

Let us define a sequence of functions by

$$v_0(x; T) = (K - x)^+; \quad v_{n+1}(x; T) = Jv_n(x; T); \quad n \geq 0; \quad \text{for all } (x; T) \in \mathbb{R}_+ \times \mathbb{R}_+. \quad (3.1)$$

We extend these functions onto $\mathbb{R}_+ \times \mathbb{R}_+$ by letting

$$v_n(x; 1) = \lim_{T \downarrow 1} v_n(x; T): \quad (3.2)$$

This sequence of functions is a bounded sequence as the next lemma shows.

Corollary 3.1 Let $(v_n)_{n \geq 0}$ be as in (3.1). For all $n \geq 0$,

$$(K - x)^+ \leq v_n(x; T) \leq 1 + \frac{K}{r}; \quad (x; T) \in \mathbb{R}_+ \times \mathbb{R}_+: \quad (3.3)$$

Proof: The first inequality follows it may not be optimal to stop immediately. Let us prove the second inequality using an induction argument: Observe that $v_0(x; T) = (K - x)^+$, $(x; T) \in \mathbb{R}_+ \times \mathbb{R}_+$, satisfies (3.3). Assume (3.3) holds for $n = n$ and let us show that it holds for $n = n + 1$. Then using (2.42)

$$\|v_{n+1}\|_1 = \|Jv_n\|_1 \leq K + \frac{1}{r+} \left(1 + \frac{K}{r}\right) = 1 + \frac{K}{r}: \quad (3.4)$$

As a corollary of Lemmas 2.2 and 2.9 we can state the following corollary, whose proof can be carried out by induction.

Corollary 3.2 Recall the sequence of functions, $(v_n(\cdot; n))_{n \geq 0}$, defined in (3.1). $(v_n(x; T))_{n \geq 0}$ is increasing for all $(x; T) \in \mathbb{R}_+ \times \mathbb{R}_+$. The function $x \mapsto v_n(x; T)$, $x \geq 0$, is convex for all $T \in \mathbb{R}_+$.

Remark 3.1 Let us define,

$$v_1(x; T) := \sup_{n \geq 0} v_n(x; T); \quad (x; T) \in \mathbb{R}_+ \times \mathbb{R}_+: \quad (3.5)$$

This function is well defined as a result of (3.3) and Corollary 3.2. In fact, it is convex because it is the upper envelope of convex functions and it is bounded by the right-hand-side of (3.3).

Corollary 3.3 For each $n \geq 0$ and $t \in \mathbb{R}_+$, $x \mapsto v_n(x; T)$, is a decreasing function on $[0; 1)$. Moreover, $T \mapsto v_n(x; T)$ is non-decreasing. The same holds for $x \mapsto v_1(x; T)$, $x \geq 0$, for all $T \in \mathbb{R}_+$.

Proof: The behaviour with respect to the first variable is a corollary of Corollary 3.2 and Remark 3.1 since any convex function that is bounded from above is decreasing. The behavior with respect to the second variable is a corollary of the second statement in Lemma 2.8.

Now, we can sharpen the upper bound in Corollary 3.1, and this has some implications about the continuity of $x \mapsto (v_n(x; T))_{n \geq 1}$ and $x \mapsto v_1(x; T)$ at $x = 0$.

Remark 3.2 The upper bound in (3.1) can be sharpened using Corollary 3.3 and Remark 2.2. Indeed, we have

$$(K - x)^+ - v_n(x; T) < K; \text{ for each } n; \text{ and } (K - x)^+ - v_1(x; T) < K; \quad (x; T) \in (0; 1)^2: \quad (3.6)$$

It follows from this observation and Corollary 3.3 that for every $T \in \mathbb{R}_+$, $x \mapsto v_n(x; T)$, for every n , and $x \mapsto v_1(x; T)$, are continuous at $x = 0$. Since they are convex, these functions are continuous on $[0; 1]$.

Lemma 3.1 The function $v_1(\cdot; \cdot)$ is the smallest fixed point of the operator J .

Proof:

$$\begin{aligned} v_1(x; T - t) &= \sup_{n \geq 1} v_n(x; T - t) \\ &= \sup_{n \geq 1} \sup_{2S_{0,T}} E^x \int_0^{Z^{(x+)}t} e^{-(x+)t} P_{\mathbb{Y}}(S_t^0; T - t) dt + e^{-(x+)} (K - S^0)^+ \\ &= \sup_{2S_{0,T}} \sup_{n \geq 1} E^x \int_0^{Z^{(x+)}t} e^{-(x+)t} P_{\mathbb{Y}}(S_t^0; T - t) dt + e^{-(x+)} (K - S^0)^+ \quad (3.7) \\ &= \sup_{2S_{0,T}} E^x \int_0^{Z^{(x+)}t} e^{-(x+)t} P_{\sup_{n \geq 1} v_n}(S_t^0; T - t) dt + e^{-(x+)} (K - S^0)^+ \\ &= Jv_1(x; T - t); \end{aligned}$$

in which last line follows by applying the monotone convergence theorem twice. Let $w : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be another fixed point of the operator J . For $(x; t) \in \mathbb{R}_+ \times \mathbb{R}_+$, $w(x; T - t) = Jw(x; T - t)$, which implies that $w(x; T - t) = Jw(x; T - t) - (K - x)^+ = v_0(\cdot)$. Now assume that $w(x; T - t) \leq v_n(x; T - t)$. Then $w(x; T - t) = Jw(x; T - t) \leq Jv_n(x; T - t) = v_{n+1}(x; T - t)$. Therefore, $w(x; T - t) \leq v_n(x; T - t)$ for all $n \geq 0$. Therefore, $w(x; T - t) \leq \sup_{n \geq 0} v_n(x; T - t) = v_1(x; T - t)$.

Lemma 3.2 The sequence $fv_n(\cdot; \cdot)g_0$ converges uniformly to $v_1(\cdot; \cdot)$. In fact, the rate of convergence is exponential:

$$v_n(x; T) - v_1(x; T) \leq v_n(x; T) + \frac{e^{-n}}{1 + r} K; \quad (x; T) \in \mathbb{R}_+ \times \mathbb{R}_+: \quad (3.8)$$

Proof: The first inequality follows from the definition of $v_1(\cdot; \cdot)$. The second inequality can be proved by induction. The inequality holds when we set $n = 0$ by Remark 3.2. Assume that the inequality holds for $n = n > 0$. Then

$$\begin{aligned} v_1(x; T) &= \sup_{2S_{0,T}} E^x \int_0^{Z^{(x+)}t} e^{-(x+)t} P_{\mathbb{Y}}(S_t^0; T - t) dt + e^{-(x+)} (K - S^0)^+ \\ &\leq \sup_{2S_{0,T}} E^x \int_0^{Z^{(x+)}t} e^{-(x+)t} P_{\mathbb{Y}}(S_t^0; T - t) dt + e^{-(x+)} (K - S^0)^+ \quad (3.9) \\ &\quad + \int_0^{Z^{(x+)}t} e^{-(x+)t} \frac{e^{-n}}{1 + r} K dt = v_{n+1}(x; T) + \frac{e^{-(n+1)}}{1 + r} K: \end{aligned}$$

Remark 3.3 Note that, for a fixed $T_0 > 0$,

$$v_n(x; T) - v_1(x; T) = v_n(x; T) + 1 - e^{(r+)^{T_0} n} \frac{1}{1+r} K; \quad x \in \mathbb{R}_+; T \in (0; T_0); \quad (3.10)$$

This can be derived using an induction argument as in the proof of Lemma 3.2, when one replaces (3.9) by

$$\begin{aligned} v_1(x; T) &= \sup_{S_0; T} E^x \int_0^T e^{(r+)^t} P_M(S_t^0; T-t) dt + e^{(r+)^T} (K - S^0)^+ \\ &+ \int_0^{T_0} dt e^{(r+)^t} 1 - e^{(r+)^{T_0} n} \frac{1}{1+r} K = v_{n+1}(x; T) + K - 1 - e^{(r+)^{T_0} n+1} \frac{1}{1+r} K^{n+1} \end{aligned} \quad (3.11)$$

Observe that one can replace K in (3.10) by $kv_1 - v_0 k_1$. Note that the convergence rate in (3.10) is fast. This will lead to a numerical scheme for the price of the American option whose error versus accuracy characteristics can be controlled.

Remark 3.4 Let $T_0 \in (0; 1)$. Then it can be shown using similar arguments that we used in the proof of Lemma 2.4 that for $(T; S) \in (0; T_0]^2$,

$$|v_1(x; T) - v_1(x; S)| \leq K |T - S| + C |T - S|^{1/2} \quad (3.12)$$

for all $x \in \mathbb{R}_+$, in which $C \in (0; 1)$ is as in Remark 2.3. In fact

$$|v_1(x; T) - v_1(x; S)| \leq L |T - S|^{1/2}; \quad (3.13)$$

for all $x \in \mathbb{R}_+$ and for some L that depends only on T_0 .

Lemma 3.3 Let $T_0 \in (0; 1)$ and $L \in (0; 1)$ be as in Remark 3.4 and $C \in (0; 1)$ be as in Remark 2.3. Then for $T; S \in (0; T_0)$ we have that

$$|v_n(x; T) - v_n(x; S)| \leq L + \frac{C}{1-a} |T - S|^{1/2} \quad \text{whenever } |T - S| \leq \frac{r}{r+} \frac{L}{K}^2; \quad (3.14)$$

for all $x \in \mathbb{R}_+$ and for all $n \geq 1$. Here, $a \in (0; 1)$ is as in Lemma 2.4. Moreover,

$$|v_1(x; T) - v_1(x; S)| \leq L + \frac{C}{1-a} |T - S|^{1/2} \quad \text{whenever } |T - S| \leq \frac{r}{r+} \frac{L}{K}^2; \quad (3.15)$$

for all $x \in \mathbb{R}_+$.

Proof: The proof of (3.14) will be carried out using an induction argument. Observe from Remark 3.4 that (3.14) holds for $n = 1$. Let us assume that (3.14) holds for n and show that it holds for $n + 1$. Using Lemma 2.4 we have that

$$|v_{n+1}(x; T) - v_{n+1}(x; S)| \leq a L + \frac{C}{1-a} + C |T - S|^{1/2}; \quad |T - S| \leq \frac{r}{r+} \frac{L + C(1-a)}{K}^2; \quad (3.16)$$

It is clear that the right-hand-side of (3.16) is less than that of (3.14), and

$$\frac{r}{r+} \frac{L + C(1-a)}{K} \leq \frac{r}{r+} \frac{L}{K}; \quad (3.17)$$

from which (3.14) follows.

Let us prove (3.15).

$$\begin{aligned} & |j_1(x;T) - v_1(x;S)| \leq |j_1(x;T) - v_n(x;T)| + |j_n(x;T) - v_n(x;S)| + |j_1(x;S) - v_n(x;S)| \\ & \leq \frac{2}{r+} \left(\frac{1}{K} + \frac{L}{1-a} \right) T \leq S^{\frac{1}{2}}; \end{aligned} \quad (3.18)$$

for any $n > 1$, which follows from (3.14) and Lemma 3.2. The result follows after taking the limit of right-hand-side of (3.18).

Lemma 3.4 Let $(v_n(\cdot; n))_0$ and $(v_1(\cdot;))$ be as in (3.4) and (3.5), respectively. Then for $n \geq 0$

$$|j_n(x;T) - v_n(y;T)| \leq |x - y| \quad \text{and} \quad |j_1(x;T) - v_1(y;T)| \leq |x - y| \quad (x,y) \in \mathbb{R}_+ \times \mathbb{R}_+; \quad (3.19)$$

and for all $T \geq 0$.

Proof: It follows from Remark 3.2 that $kv_n k_1 \leq K$, for all $n \geq 0$, and $kv_1 k_1 \leq K$. Moreover, for each $n \geq 0$, $v_n(\cdot; T)$ is convex (for all $T \in \mathbb{R}_+$) as a result of Corollary 3.2. On the other hand, it was pointed in Remark 3.1 that $v_1(\cdot; T)$ is convex for all $T \in \mathbb{R}$. Since

$$v_{n+1}(x;T) = Jv_n(x;T) \quad \text{and} \quad v_1(x;T) = Jv_1(x;T); \quad (3.20)$$

(3.19) follows from Lemma 2.2. Note that the second equation in (3.20) is due to Lemma 3.1.

Lemma 3.5 The sequence of functions $(v_n(\cdot; n))_0$ (defined in (3.1)) and its limit $v_1(\cdot;)$ satisfy

$$D_+^x v_n(\cdot;) \leq 1; \quad \text{for all } n \text{ and } Jv_1(\cdot;) \leq 1; \quad (3.21)$$

Proof: The proof follows from the fact that $v_n(0;T) = v_1(0;T) = K$ (see Remark 3.2), and $v_1(x;T) - v_n(x;T) \leq (K - x)^+$ (see Remarks 3.1 and 3.2) for all $(x;T) \in \mathbb{R}_+ \times \mathbb{R}_+$ and $n \geq 0$ and that the functions $v_n(\cdot;)$, $n \geq 0$, and $v_1(\cdot;)$ are convex with respect to their first variable (see Remark 3.2).

Lemma 3.6 Let us denote by $(v_n(\cdot; n))_0$ the sequence of functions defined in (3.1) and let $v_1(\cdot;)$ denote its limit. Recall (2.7). Then for all $T \geq 0$ and $n \geq 0$, $C_T^{v_{n+1}} = (c^{v_{n+1}}(T); 1)$ for some $c^{v_{n+1}}(T) \in (0; K)$ and $C_T^{v_1} = (c^{v_1}(T); 1)$ for some $c^{v_1} \in (0; K)$. Then for $n \geq 0$, the restriction of $v_{n+1}(\cdot;)$ onto ${}^2\mathbb{R}$ is the unique solution of

$$\begin{aligned} & A v_{n+1}(x;T) - (r+) v_{n+1}(x;T) + (P - H)(x;T) - \frac{\partial}{\partial T} v_{n+1}(x;T) = 0; \quad x > c^{v_{n+1}}(T); \\ & v_{n+1}(x;T) = (K - x); \quad x \leq c^{v_{n+1}}(T); \end{aligned} \quad (3.22)$$

in which A is as in (2.5). Also for $n \geq 1$, $v_n(\cdot; \cdot)$ satisfies

$$\frac{\partial}{\partial x} v_{n+1}(x; T) \Big|_{x=c^{v_{n+1}}(T)} = 1; \quad T > 0; \quad (3.23)$$

Moreover, the restriction of $v_1(\cdot; \cdot)$ onto \mathbb{R}_+ is the unique solution of

$$\begin{aligned} & A v_1(x; T) - (r + \frac{1}{2}\sigma^2) v_1(x; T) + \frac{1}{2}\sigma^2 (v_1''(x; T)) - \frac{\partial}{\partial T} v_1(x; T) = 0 \quad x > c^{v_1}(T); \\ & v_1(x; T) = (K - x)^+ \quad x \leq c^{v_1}(T); \end{aligned} \quad (3.24)$$

and satisfies

$$\frac{\partial}{\partial x} v_1(x; T) \Big|_{x=c^{v_1}(T)} = 1; \quad T > 0; \quad (3.25)$$

Proof: The fact that $C^{v_{n+1}} = (c^{v_{n+1}}; 1)$ and $C^{v_1} = (c^{v_1}; 1)$ for some $c^{v_{n+1}} \in (0; K)$ and $c^{v_1} \in (0; K)$ follows from Lemma 2.5 using the fact that the assumptions in that lemma hold as a result of Corollaries 3.2, 3.3; Remarks 3.1 and 3.2; and Lemmas 3.1 and 3.5.

The partial differential equations (3.22) and (3.24) are satisfied as a corollary of Lemma 2.6; Corollaries 3.2 and 3.3, Remarks 3.1, 3.2; Lemmas 3.1, 3.3, and 3.5.

Observe that since $v_n(\cdot; \cdot)$ is convex (Corollary 3.2) and non-increasing (Corollary 3.3) with respect to its first variable then $v_{n+1}(\cdot; \cdot) = \mathbb{P} v_n(\cdot; \cdot)$ satisfies the smooth fit condition in (3.23) as a result of Lemma 2.7. The smooth fit condition in (3.25) holds for $v_1(\cdot; \cdot)$, since $v(\cdot; \cdot) = \mathbb{P} v_1(\cdot; \cdot)$ (Lemma 3.1), and since $x \mapsto v_1(x; T)$ is convex and non-increasing, as a result of Lemma 2.7.

Theorem 3.1 Let $V(\cdot; \cdot)$ as in (1.3), which is the value function of the American put option for the jump diffusion S , whose dynamics are given in (1.1). Then $V(\cdot; \cdot)$ is the unique solution (in the classical sense of) the integro-partial differential equation in (3.24) and $(x; T) \in \mathbb{R}_+ \times \mathbb{R}_+$ belongs to the optimal continuation region if $x > c^{v_1}$. Moreover, it satisfies the smooth fit condition at the optimal stopping boundary, i.e.

$$\frac{\partial}{\partial x} V(x; T) \Big|_{x=c^{v_1}(T)} = 1; \quad T > 0; \quad (3.26)$$

Proof: Let $T \in (0; 1)$ and define

$$M_t = e^{-rt} v_1(S_t; T - t); \quad (3.27)$$

Let us define

$$x := T \wedge \inf_{t \in [0; T]} S_t - c^{v_1}(T - t); \quad (3.28)$$

It follows from (3.24) and Itô's lemma that $\mathbb{E} M_{t \wedge x} g_0 - t - T$ is a bounded \mathbb{P} -martingale. Using the optional sampling theorem one obtains,

$$v_1(x; T) = M_0 = \mathbb{E}^x \mathbb{E} M_{t \wedge x} g = \mathbb{E}^x e^{-rx} v_1(S_x; T - x) = \mathbb{E}^x e^{-rx} (K - S_x)^+ = V(x; T); \quad (3.29)$$

On the other hand, it is easy to show that for any stopping time $\tau \leq S$, $fM_{\tau} - g_0 - \tau$ is a positive P -supermartingale. Again, using the optional sampling theorem

$$v_1(x; T) = M_0 - E^x fM_{\tau} - g = E^x e^{-r} v_1(S; T) - E^x e^{-r} (K - S)^+; \quad (3.30)$$

therefore $v_1(x; T) = V(x; T)$. Now, the statement of the theorem immediately follows.

References

- Aili, L. and Kyprianou, A. E. (2005). Some remarks on first passage of Levy processes, the American put and the pasting principles, *Annals of Applied Probability* 15: 2062{2080.
- Alvarez, L. H. R. (2004). Solving optimal stopping problems of linear diffusions by applying convolution approximations, *Mathematical Methods of Operations Research* 53 (1): 89{99.
- Bayraktar, E. (2007). Remarks on the American put option for jump diffusions, submitted, <http://arxiv.org/abs/math/0703538>.
- Bayraktar, E., Dayanik, S. and Karatzas, I. (2006). Adaptive poisson disorder problem, *Annals of Applied Probability* 16 (3): 1190{1261.
- Bouchard, B., Karoui, N. E. and Touzi, N. (2005). Maturity randomisation for stochastic control problems, *Annals of Applied Probability* 15 (4): 2575{2605.
- Carr, P. (1998). Randomization and the American put, *Review of Financial Studies* 11 (3): 597{626.
- Friedman, A. (1964). *Partial Differential Equations of Parabolic Type*, Prentice Hall, Englewood Cliffs, N.J.
- Friedman, A. (2006). *Stochastic Differential Equations and Applications*, Dover Publications, Mineola, NY.
- Glasserman, P. and Yu, B. (2004). Simulation for american options: regression now or regression later?, *Monte Carlo and quasi-Monte Carlo methods 2002*, Springer, Berlin: 213{226.
- Ishii, H. (1989). On uniqueness and existence of viscosity solutions of fully non-linear second-order elliptic PDE's, *Communications in Pure and Applied Mathematics* 42: 14{45.
- Karatzas, I. and Shreve, S. E. (1998). *Methods of Mathematical Finance*, Springer, New York.
- Kusner, H. and Dupuis, P. G. (2003). *Numerical Methods for Stochastic Control Problems in Continuous Time*, Springer Verlag, New York.
- Longstaffe, F. A. and Schwartz, E. (2001). Valuing american options by simulation: a simple least squares approach, *Review of Financial Studies* 14: 113{148.
- Mordecki, E. and Salminen, P. (2006). Optimal stopping of hunt and Levy processes, preprint, <http://arxiv.org/abs/math.PR/0605054>.
- Peskir, G. and Shiryaev, A. (2006). *Optimal Stopping and Free Boundary Problems*, Birkhauser, Basel, Switzerland.

- Pham, H. (1997). Optimal stopping, free boundary, and american option in a jump-diffusion model, *Applied Mathematics and Optimization* 35:145{164.
- Pham, H. (1998). Optimal stopping of controlled jump diffusion processes: A viscosity solution approach, *Journal of Mathematical Systems, Estimation and Control* 8 (1):1{27.
- Rockafellar, R. T. (1997). *Convex Analysis*, Princeton University Press, Princeton Landmarks in Mathematics, Princeton, NJ.
- Wilmott, P., Howison, S. and Dewynne, J. (1995). *The Mathematics of Financial Derivatives*, Cambridge University Press, New York, NY.