Vector-valued L_p -convergence of orthogonal series and Lagrange interpolation.

H. König (Kiel)^{*} N. J. Nielsen (Odense)[†]

Abstract

We give necessary and sufficient conditions for interpolation inequalities of the type considered by Marcinkiewicz and Zygmund to be true in the case of Banach space-valued polynomials and Jacobi weights and nodes. We also study the vector-valued expansion problem of L_p -functions in terms of Jacobi polynomials and consider the question of unconditional convergence. The notion of type p with respect to orthonormal systems leads to some characterizations of Hilbert spaces. It is also shown that various vector-valued Jacobi means are equivalent.

1 Introduction and results

Let X be a Banach space, $1 \leq p < \infty$ and $L_p(\mathbb{R}; \mathbb{X})$ denote the space of (classes of) p-th power integrable functions with norm $|| f || := (\int_{\mathbb{R}} || f(t) ||^p dt)^{1/p}$. A Banach space is a *UMD-space* provided that the Hilbert transform on \mathbb{R} ,

$$Hf(t) := \text{p.v.} \int_{\mathbb{R}} \frac{f(s)}{t-s} ds, \ f \in L_p(\mathbb{R}; \mathbb{X}),$$
(1)

^{*}Supported in part by the DFG, Ko 962/3-1.

[†]Supported in part by the DFG, Ki 51/98-1 and the Danish Natural Science Council, grant 11-8622.

defines a bounded operator $H: L_p(\mathbb{R}; \mathbb{X}) \longrightarrow \mathbb{L}_1(\mathbb{R}; \mathbb{X})$ for some 1 .It is well-known that this holds for*some*<math>1 if and only if it holds for*all* $<math>1 , see e.g. Schwarz [28]. All <math>L_q(\mu)$ -spaces with $1 < q < \infty$ or all reflexive Orlicz spaces are UMD-spaces, cf. Fernandez and Garcia [7].

Let $I = (-1, 1), \alpha, \beta > -1$ and $w_{\alpha\beta}(t) := (1-t)^{\alpha}(1+t)^{\beta}$ for $t \in I$. Let

$$L_p(I, w_{\alpha\beta}; X) := \{ f : I \longrightarrow X | \| f \|_p := \| f \|_{p;\alpha,\beta} := (\int_I \| f(t) \|^p w_{\alpha\beta}(t) dt)^{1/p} < \infty \}.$$

The scalar product in $L_2(I, w_{\alpha\beta}) := L_2(I, w_{\alpha\beta}; \mathbb{R})$ will be denoted by $\langle \cdot, \cdot \rangle$ or $\langle \cdot, \cdot \rangle_{\alpha\beta}$. For $\alpha = \beta$ we just write w_{α} and $\langle \cdot, \cdot \rangle_{\alpha}$. By $\Pi_n(X)$ we denote the space of polynomials of degree $\leq n$ with coefficients in X. Let $\Pi_n := \Pi_n(\mathbb{R})$. The $L_2(I, w_{\alpha\beta})$ -normalized Jacobi polynomials with respect to $(I, w_{\alpha\beta})$ will be denoted by $p_n^{(\alpha,\beta)}$, $n \in \mathbb{N}_{\Bbbk}$. Hence $p_n^{(\alpha,\beta)} \in \Pi_n$ and

$$\langle p_n^{(\alpha,\beta)}, p_m^{(\alpha,\beta)} \rangle_{\alpha,\beta} = \int_I p_n^{(\alpha,\beta)}(t) p_m^{(\alpha,\beta)}(t) w_{\alpha\beta}(t) dt = \delta_{nm}.$$
 (2)

This normalization is more convenient for us than the standard one of Szegö [30]. For $\alpha = \beta = -\frac{1}{2} \left(\frac{1}{2}\right)$ one gets the Tchebychev polynomials of the first (second) kind, for $\alpha = \beta = 0$ the Legendre polynomials. Let $t_1 > \cdots > t_{n+1}$ denote the zeros of $p_{n+1}^{(\alpha,\beta)}$, all of which are in I, and $\lambda_1, \cdots, \lambda_{n+1} > 0$ the Gaussian quadrature weights. Thus for any real polynomial q of degree $\leq 2n + 1$, one has

$$\int_{I} q(t) w_{\alpha\beta}(t) dt = \sum_{j=1}^{n+1} \lambda_j q(t_j).$$
(3)

Clearly, λ_j and t_j depend on n, j, α and β but not on q. One has for $\alpha, \beta > -1$

$$\lambda_j = (2n + \alpha + \beta + 3)((1 - t_j^2)p_{n+1}^{(\alpha,\beta)'}(t_j)^2)^{-1} \sim \left\{ \begin{array}{cc} j^{2\alpha+1}/n^{2\alpha+2} & j \le \frac{n}{2} \\ (n+2-j)^{2\beta+1}/n^{2\beta+2} & j > \frac{n}{2} \end{array} \right\}$$
(4)

$$1 - t_j^2 \sim (j/n)^2, and \quad p_{n+1}^{(\alpha,\beta)'}(t_j) \sim n^{5/2+\alpha}/j^{3/2+\alpha} \quad \text{for all } j \le \frac{n}{2}.$$
(5)
$$p_n^{(\alpha,\beta)}(-x) = (-1)^n p_n^{(\alpha,\beta)}(x).$$

See Szegö [30, 3.4, 4.1, 4.3, 8.9, 15.3], taking into account the different normalization there. Here $\lambda_j \sim f_j$ means that there are constants $c_1, c_2 > 0$ independent of j and n such that $c_1 f_j \leq \lambda_j \leq c_2 f_j$ for all n and j concerned. For $\alpha = \beta = -\frac{1}{2}, \lambda_j = \pi/(n+1)$.

Marcinkiewicz and Zygmund [31, ch. X] proved interpolation inequalities for trigonometric polynomials of degree $\leq n$ which for even trigonometric polynomials g, after a transformation $g(x) = q(\text{cost}), x = \text{cost}, q \in \Pi_n$, can be restated as

$$\frac{1}{3} \left(\sum_{j=1}^{n+1} |q(t_j)|^p / (n+1)\right)^{\frac{1}{p}} \leq \left(\frac{1}{2} \int_I |q(t)|^p (1-t^2)^{-\frac{1}{2}} dt\right)^{\frac{1}{p}} \leq c_p \left(\sum_{j=1}^{n+1} |q(t_j)|^p / (n+1)\right)^{1/p}.$$

Here (t_j) are the zeros of the Jacobi polynomial $p_{n+1}^{\alpha,\beta}$ in the Tchebychev case $\alpha = \beta = -1/2$, and c_p depends on 1 only. The left inequality $holds for <math>p = 1, \infty$ as well whereas the right one fails, in general. For p = 2, (3) gives more precise information since $\lambda_j = \pi/(n+1)$. The Marcinkiewicz-Zygmund inequalities extend to the Jacobi case of general $\alpha, \beta > -1$ and to the vector-valued setting in the following sense:

Theorem 1 Let X be a Banach space, $1 \le p \le \infty$, $\alpha, \beta > -1$, (t_j) the zeros of $p_{n+1}^{\alpha,\beta}$ and (λ_j) the corresponding quadrature weights.

a) There is c > 0 such that for all $1 \le p \le \infty$, $n \in \mathbb{N}$ and $q \in \Pi_{2n}(X)$

$$c^{-1} (\sum_{j=1}^{n+1} \lambda_j \parallel q(t_j) \parallel^p)^{1/p} \le (\int_{-1}^1 \parallel q(t) \parallel^p w_{\alpha\beta}(t) dt)^{1/p}.$$
(6)

b) Let

$$\mu(\alpha,\beta): = \max(1,4(\alpha+1)/(2\alpha+5),4(\beta+1)/(2\beta+5))$$

$$m(\alpha,\beta): = \max(1,4(\alpha+1)/(2\alpha+3),4(\beta+1)/(2\beta+3))$$

and $M(\alpha, \beta) := m(\alpha, \beta)'$, i.e. $m(\alpha, \beta)^{-1} + M(\alpha, \beta)^{-1} = 1$. Then the following are equivalent.

(1) There is $c_p > 0$ such that for all $n \in \mathbb{N}$ and $q \in \Pi_n(X)$

$$\left(\int_{-1}^{1} \| q(t) \|^{p} w_{\alpha\beta}(t) dt\right)^{1/p} \le c_{p} \left(\sum_{j=1}^{n+1} \lambda_{j} \| q(t_{j}) \|^{p}\right)^{1/p}.$$
 (7)

(2) X is a UMD-space and p satisfies $\mu(\alpha, \beta) .$

Part (a) is proved just as the scalar result which goes back to Askey [2], Nevai [19] and Zygmund [32]. The converse inequality (7) was shown in the scalar case (for $\alpha = \beta$) by Askey [2] under the more restrictive assumption $m(\alpha, \beta) using (a) and duality; the duality method, however,$ $fails if <math>\mu(\alpha, \beta) . The question whether (7) in the vector-valued$ case requires X to be a UMD-space was raised by Pietsch in the case of $trigonometric polynomials (corresponding to <math>\alpha = \beta = -1/2$) and solved by him in this case by a different method [22].

In terms of Banach spaces, Theorem 1 states that the spaces $\Pi_n(X)_p$ as subspaces of $L_p(I, w_{\alpha\beta}; X)$ are uniformly isomorphic to $l_p^{n+1}(X)$ -spaces, by evaluating the polynomials q at the zeros (t_j) , provided that (b), (2) holds; i.e. the Banach-Mazur distances $d(\Pi_n(X)_p, l_p^{n+1}(X))$ are uniformly bounded.

For $f \in L_p(I, w_{\alpha\beta}; X)$, let $Q_n f := \sum_{j=0}^n \langle f, p_j^{\alpha,\beta} \rangle > p_j^{(\alpha,\beta)} \in \Pi_n(X)_p$ denote the orthogonal projection of f onto the space of polynomials of degree $\leq n$. The following vector-valued expansion theorem for Jacobi polynomials generalizes the classical scalar result of Pollard [26] and Muckenhaupt [18].

Theorem 2 Let X be a Banach space, $1 \le p \le \infty$, $\alpha, \beta > -1$ and $m(\alpha, \beta)$ and $M(\alpha, \beta)$ as before. Then the following are equivalent:

- (1) For all $f \in L_p(I, w_{\alpha\beta}; X)$ $Q_n f$ converges to f in the L_p -norm.
- (2) X is a UMD-space and $m(\alpha, \beta) .$

The necessity of the UMD-condition on X will be proved using Theorem 1; the interval for p is "symmetric" with respect to p = 2 and smaller than the one exhibited in Theorem 1, (b). Analogues of Theorems 1 and 2 in the case of the Hermite polynomials are proved in [12]. Using the results of Gilbert [8], we also prove that various vector-valued Jacobi means are equivalent:

Proposition 3 Let $\alpha, \beta > -1$, $1 , <math>\gamma \in \mathbb{R}$ with $|\frac{\gamma}{2} + \frac{1}{p} - \frac{1}{2}| < \frac{1}{4}$. Let X be a UMD-space. Then there is $M = M(\alpha, \beta, \gamma, p) \ge 1$ such that for all $n \in \mathbb{N}$ and all $x_1, \ldots, x_n \in X$

$$(\int_{-1}^{1} \| \sum_{j=0}^{n} p_{j}^{(\alpha,\alpha)}(t) x_{j} \|^{p} (1-t^{2})^{(\alpha+\gamma)p/2} dt)^{1/p}$$

$$\stackrel{M}{\sim} (\int_{-1}^{1} \| \sum_{j=0}^{n} p_{j}^{(\beta,\beta)}(t) x_{j} \|^{p} (1-t^{2})^{(\beta+\gamma)p/2} dt)^{1/p}.$$
(8)

Here $\stackrel{M}{\sim}$ means that the quotient of the two expressions is between 1/Mand M. Instead of (α, α) and (β, β) , one could consider (α_1, α_2) and (β_1, β_2) as Jacobi-indices, provided the weight functions are changed accordingly. The convergence of the Jacobi series in Theorem 2 is not unconditional unless p = 2 and X is a Hilbert space, as will follow from the following general result. Recall that a series $\sum_{n \in \mathbb{N}} y_n$ in a Banach space Y converges unconditionally if $\sum_{n \in \mathbb{N}} \varepsilon_n y_n$ converges in Y for all choices of signs $\varepsilon_n = \pm 1$.

Proposition 4 Let (Ω, μ) be a measure space and (p_n) be a complete orthonormal system in $L_2(\Omega, \mu)$, assumed to be infinite dimensional. Let X be a Banach space and $1 \le p < \infty$. Assume that for all $f \in L_p(\Omega, \mu; X)$, the series $\sum_n < f, p_n > p_n$ converges unconditionally in $L_p(\Omega, \mu; X)$. Then:

- (i) If $|| p_j ||_2 \sim || p_j ||_{\max(p,p')}$ and (Ω, μ) is a finite measure space, one has p = 2.
- (ii) If $\sup_{j} |p_{j}| \in L_{2}(\Omega, \mu)$, X is isomorphic to a Hilbert space.

Statement (ii) was also shown by Defant and Junge [6]. Both conditions (i) and (ii) are satisfied in the Jacobi case provided that the condition $m(\alpha, \beta) holds (necessary for convergence). Without an as$ $sumption like <math>\sup_j |p_j| \in L_2(\Omega, \mu)$, X is not isomorphic to a Hilbert space in general, as the Haar system shows. However, one has:

Proposition 5 Let $1 and <math>(p_n)_{n \in \mathbb{N}}$ be an unconditional basis of $L_p(0,1)$. Let X be a Banach space such that for any $f \in L_p(0,1;X)$, the series $\sum_{n \in \mathbb{N}} \langle f, p_n \rangle p_n$ converges unconditionally in $L_p(0,1;X)$. Then X is a UMD-space.

The proof shows that the Haar basis is unconditional in $L_p(0, 1; X)$ which by Maurey [16], Burkholder [5] and Bourgain [4] is equivalent to X being a UMD-space. It was shown by Aldous [1] that X is a UMD-space if $L_p(X)$ has an unconditional basis.

Let (Ω, μ) be a measure space and $(p_n)_{n \in \mathbb{N}}$ be a complete orthonormal system in $L_2(\Omega, \mu)$. We say that a Banach space X has $(\mathbf{p_n}) - \mathbf{type} \ 2$ provided there is c > 0 such that for all $m \in \mathbb{N}$ and all $x_1, \ldots, x_m \in X$

$$(\int_{\Omega} \| \sum_{j=1}^{m} p_j(t) x_j \|^2 d\mu(t))^{1/2} \le c (\sum_{j=1}^{m} \| x_j \|^2)^{1/2}.$$

X has $(\mathbf{p_n}) - \mathbf{cotype}\ 2$ if the reverse inequality holds. In [24], Pisier showed for the Haar system (h_n) , that (h_n) -type 2 of X is equivalent to X being 2-smooth, e.g. has an equivalent uniformly convex norm with modulus of convexity of power type 2. In Pisier and Xu [25] the related notion of H-type $p (\leq 2)$ is considered for all orthonomal systems (p_n) . Kwapień [13] studied this notion for the trigonometric system (e_n) in $L_2(0, 2\pi)$ showing that (e_n) -type 2 (also called Fourier-type 2, $e_n(t) = exp(int)$) of X implies that X is isomorphic to a Hilbert space. This result generalizes to the case of Jacobi polynomials.

Proposition 6 Let X be a Banach space which is Jacobi $(p_n^{(\alpha,\beta)})$ -type 2 for some $\alpha, \beta > -1$. Then X is isomorphic to a Hilbert space.

The proof uses the interpolation inequalities of Theorem 1. In general, (p_n) -type 2 implies type 2 in the usual sense [17], i.e. with respect to the Rademacher system (r_n) , $r_n(t) = \operatorname{sgn sin} 2^n \pi t$.

Proposition 7 Let X be a Banach space which is of (p_n) -type 2 for some complete orthonormal system (p_n) in $L_2(0,1)$. Then X is of Haar type 2, hence 2-smooth and of type 2.

There is a partial converse to this result.

Proposition 8 Let X be a Banach space and $(p_n)_{n\in\mathbb{N}} \subset L_2(0,1)$ be a complete orthonormal system such that for any $f \in L_2(0,1;X)$ the series $\sum_{n\in\mathbb{N}} < f, p_n > p_n$ converges unconditionally in $L_2(0,1;X)$. Then, if X has type 2, it also has (p_n) -type 2.

It follows from Proposition 5 that the unconditionality assumption in Proposition 8 implies that the space X in question has UMD. On the other hand, if X has UMD the unconditionality assumption in Proposition 8 is satisfied for the Haar system, and thus by Pisier's result mentioned above, type 2 and UMD of X implies that X is 2-smooth. This in turn implies type 2 but does not imply the UMD-property, since by Bourgain [4] there exists a Banach lattice satisfying an upper-p and lower-q estimate and failing the UMD-property; choosing 2 there, such a lattice is 2-smooth,cf. [15].

2 The interpolation inequalities

For the proof of Theorem 1, we need a well-known fact about continuity in L_p , cf. Pollard [26] or Benedek, Murphy and Panzone [3]. In the scalar case, it is a special case of the theory of weighted singular integral operators with weights in the Muckenhaupt class A_p , cf. Garcia-Cuerva and de Francia [9, chap. IV].

Lemma 1 Let X be a Banach space, $1 \le p \le \infty$, $b \in \mathbb{R}$ and $k : \mathbb{R}^{\nvDash} \longrightarrow \mathbb{R}$ be defined by $k(u, v) := ||u/v|^b - 1|/|u - v|$. Then the integral operator T_k given by $T_k f(u) := \int_{\mathbb{R}} k(u, v) f(v) dv$ defines a bounded operator $T_k : L_p(\mathbb{R}; \mathbb{X}) \longrightarrow$ $\mathbb{L}_1(\mathbb{R}; \mathbb{X})$ provided that -1/p < b < 1 - 1/p (actually if and only if).

Proof: We sketch the simple proof. Let $r(u, v) := |u/v|^{1/pp'}$. It suffices to show that

$$\sup_{u} \int_{\mathbb{R}} k(u,v) r(u,v)^{p'} dv \le M, \ \sup_{v} \int_{\mathbb{R}} k(u,v) r(u,v)^{-p} du \le M.$$
(9)

An application of Hölder's inequality then shows that T_k is continuous as a map $T_k : L_p(\mathbb{R}; \mathbb{X}) \longrightarrow \mathbb{L}_{\mathbb{I}}(\mathbb{R}; \mathbb{X})$ with norm $\leq M$. To check the first inequality in (9), substitute v/u = t to find

$$\sup_{u \neq 0} \int_{\mathbb{R}} k(u, v) \ r(u, v)^{p'} dv = \int_{\mathbb{R}} |t^{-b} - 1| \ |t|^{-1/p} / |t - 1| dt.$$

This is finite since integrability at 0 is assured by b < 1 - 1/p, and integrability at $\pm \infty$ by b > -1/p. Note that for $t \longrightarrow 1$, there is no singularity,

the integrand tends to |b|. The second condition in (9) is checked similarly. \Box

By Szegö [30, 7.32, 4.3], for any pair of indices $\alpha, \beta > -1$, there is $c = c_{\alpha,\beta}$ such that for all $n \in \mathbb{N}$ and $t \in [-1, 1]$, the L_2 -normalized Jacobi polynomials $p_n^{(\alpha,\beta)}$ satisfy the estimate

$$|p_n^{(\alpha,\beta)}(t)| \le c (1-t+n^{-2})^{-(\alpha/2+1/4)} (1+t+n^{-2})^{-(\beta/2+1/4)}$$
(10)

Proof of theorem 1: We start with

(b), $(2) \Rightarrow (1)$.

Assume that X is a UMD space and that p satisfies $\mu(\alpha, \beta) .$ $Let <math>q \in \prod_n(X)$ and put $y_j := q(t_j)/p'_{n+1}(t_j)$. The Lagrange functions $\ell_j \in \prod_n$,

$$\ell_j(t) := p_{n+1}^{(\alpha,\beta)}(t) / (p_{n+1}^{(\alpha,\beta)'}(t_j)(t-t_j))$$

satisfy $\ell_j(t_i) = \delta_{ji}$ for $i, j = 1, \dots, n+1$ and thus q coincides with its interpolating polynomial $q = \sum_{j=1}^{n+1} q(t_j)\ell_j$. We have to estimate

$$L := \left(\int_{-1}^{1} \| q(t) \|^{p} w_{\alpha\beta}(t) dt\right)^{1/p} = \left(\int_{-1}^{1} \| \sum_{j=1}^{n+1} y_{j} \frac{p_{n+1}^{(\alpha,\beta)}(t)}{t - t_{j}} \|^{p} w_{\alpha\beta}(t) dt\right)^{1/p}$$

from above. Let $I_j = (t_j, t_{j-1})$, $|I_j| = (t_{j-1} - t_j)$ and χ_j be the characteristic function of I_j , for $j = 1, \dots, n+1$, with $t_o := 1$. The proof relies on the fact that $1/(t - t_j)$ is sufficiently close to the Hilbert transform of $-\chi_j/|I_j|$ at t which is

$$H(-\frac{\chi_j}{|I_j|})(t) = \frac{1}{|I_j|} \log |\frac{t - t_{j-1}}{t - t_j}| = \frac{1}{|I_j|} \log |1 - \frac{|I_j|}{t - t_j}|.$$

Let $J_n = [a_n, b_n]$ where

$$a_n = \begin{cases} -1 & \text{if } \beta > -1/2 \\ -1 + dn^{-2} & \text{if } \beta \le -1/2, \end{cases}$$

$$b_n = \begin{cases} 1 & \text{if } \alpha > -1/2 \\ 1 - dn^{-2} & \text{if } \alpha \le -1/2; \end{cases}$$

and d is chosen such that $\min(1 - t_1, 1 + t_{n+1}) \ge 2dn^{-2}$. By [30] this is possible.

It follows from (10) that for $n \in \mathbb{N}$ and $t \in J_n$

$$|p_n^{(\alpha,\beta)}(t)| \le c(1-t)^{-(\alpha/2+1/4)}(1+t)^{-(\beta/2+1/4)}.$$
(11)

In the following, constants c_1, c_2, \cdots may depend on α, β and p, but not on n, j and t. We claim that for $n \in \mathbb{N}$ and $t \in J_n$

$$|p_{n+1}^{(\alpha,\beta)}(t)||\frac{1}{t-t_j} + H(\frac{\chi_j}{|I_j|})(t)| \le f_j(t),$$
(12)

where

$$f_j(t) := c_1 \min(\frac{1}{|I_j|}, \frac{|I_j|}{(t-t_j)^2})(1-t)^{-(\alpha/2+1/4)}(1+t)^{-(\beta/2+1/4)}$$

If t is such that $|t-t_j| > 2|I_j|$, (12) follows from (11) and $|x-\log(1+x)| \le x^2$ for $|x| \le 1/2$, i.e. $|\frac{1}{t-t_j} + H(\frac{\chi_j}{|I_j|})(t)| \le \frac{|I_j|}{(t-t_j)^2}$. For $|t-t_j| \le 2|I_j|$, one uses that $p_{n+1}^{(\alpha,\beta)}$ has a zero in t_j . By the mean-value theorem there is a θ between t and t_j such that

$$\frac{p_{n+1}^{(\alpha,\beta)}(t)}{t-t_j}| = |\frac{p_{n+1}^{(\alpha,\beta)}(t) - p_{n+1}^{(\alpha,\beta)}(t_j)}{t-t_j}| = |p_{n+1}^{(\alpha,\beta)'}(\theta)| \le f_j(t),$$

using that by Szegö [30, 8.9] and (5), e.g. for $j \leq n/2$,

$$|p_{n+1}^{(\alpha,\beta)'}(\theta)| \le c_3 n^{\alpha+5/2} / j^{\alpha+3/2} \sim f_j(t_j) \sim f_j(t).$$

We note that (only) for j = 1, the logarithmic singularity of $H(x_j/|I_j|)$ at t = 1 is not compensated by a zero of $p_{n+1}^{(\alpha,\beta)}$ $(t_0 = 1)$, but (12) is true in this case and $\alpha > -1/2$ too, since by (10) for $(1-t) \le n^{-2}$

$$|p_n(t)H(\frac{\chi_j}{|I_j|})(t)| \leq c_4 n^{5/2+\alpha} |\log(n^2(1-t))|$$

$$\leq c_5 n^2 (1-t)^{-(\alpha/2+1/4)}$$

using $|\log v| \le c_3 v^{-\varepsilon}$ for $\varepsilon = \frac{\alpha}{2} + \frac{1}{4} > 0$ and $0 < v \le 1$. Hence (12) holds. Applying this we find

$$L_n := \left(\int_{J_n} \|\sum_{j=1}^{n+1} y_j \frac{p_{n+1}^{(\alpha,\beta)}(t)}{t-t_j} \|^p w_{\alpha\beta}(t) dt\right)^{1/p} \le M_1 + M_2.$$

where

$$M_{1} := c_{6} \left(\int_{J_{n}} \| H(\sum_{j=1}^{n+1} y_{j} \chi_{j} / |I_{j}|)(t) \|^{p} (1-t)^{\gamma p} (1+t)^{\delta p} dt \right)^{1/p},$$

$$M_{2} := \left(\int_{J_{n}} (\sum_{j=1}^{n+1} \| y_{j} \| f_{j}(t))^{p} w_{\alpha \beta}(t) dt \right)^{1/p},$$

and $\gamma := \alpha(1/p - 1/2) - 1/4$, $\delta := \beta(1/p - 1/2) - 1/4$. The restrictions on p are equivalent to $-1/p < \gamma$, $\delta < 1 - 1/p$ and $1 . In particular, <math>|\gamma| < 1$, $|\delta| < 1$. We estimate the "main" term M_1 and the "error" term M_2 separately.

We claim that the kernel $K(t,s) := 1/(t-s)((1-t)/(1-s))^{\gamma}((1+t)/(1+s))^{\delta}$, $t,s \in [-1,1]$ defines a bounded integral operator $T_k : L_p(-1,1;X) \longrightarrow$ $L_p(-1,1;X)$. Indeed, by lemma 1, the kernel $|((1-t)/(1-s))^{\gamma} - 1|/|t-s|$ defines a bounded operator $L_p(\mathbb{R};\mathbb{X}) \longrightarrow \mathbb{L}_{\mathbb{I}}(\mathbb{R};\mathbb{X})$, replacing t and s by (1-t) and (1-s). Since X is a UMD-space, so does 1/(t-s) and hence also $1/(t-s)((1-t)/(1-s))^{\gamma}$. Since $(\frac{1+t}{1+s})^{\delta}$ is bounded from above and below by positive constants for $s, t \in [0, 1]$, the kernel K defines a bounded operator $T_k: L_p(0,1;X) \longrightarrow L_p(0,1;X)$. The same holds on the interval [-1,0]. The kernel is less singular for t and s of different sign: if $s \in [-1, 0], t \in [0, 1], t \in [0, 1]$ the substitution $s \to -s$ yields a kernel of the type $(1/(t+s))w_1(t)w_2(s)$ on $[0,1]^2$, where w_1 and w_2 are integrable over [0,1] and bounded near 0. By Hardy, Littlewood and Polya [10], the kernel 1/(t+s) defines an operator $L_p(0,\infty) \longrightarrow L_p(0,\infty)$ of norm $\pi/\sin(\pi/p)$, for any 1 .Since 1/(t+s) is positive, this also holds for X-valued functions and hence $T_k: L_p(-1,0;X) \longrightarrow L_p(0,1;X)$ is bounded as well. The case $s \in [0,1]$, $t \in [-1,0]$ is treated similarly. Together these facts prove the claim. Hence there is a c_7 such that for all $f \in L_p(-1, 1; w_{\gamma\delta}; X)$

$$\left(\int_{-1}^{1} \|\int_{-1}^{1} \frac{f(s)}{t-s} ds \|^{p} (1-t)^{\gamma p} (1+t)^{\delta p} dt\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} (1+s)^{\delta p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} (1-s)^{\gamma p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{p} ds\right)^{1/p} \le c_{7} \left(\int_{-1}^{1} \|f(s)\|^{$$

and thus

$$M_{1} \leq c_{6}c_{7}\left(\int_{-1}^{1} \|\sum_{j=1}^{n+1} y_{j}\chi_{j}(s)/|I_{j}| \|^{p} w_{\gamma p,\delta p}(s)ds\right)^{1/p}$$

$$\leq c_{8}\left(\sum_{j=1}^{n+1} \|y_{j}\|^{p}/|I_{j}|^{p-1}w_{\gamma p,\delta p}(t_{j})\right)^{1/p} \qquad (13)$$

$$\leq c_{9}\left(\sum_{j=1}^{n+1} \lambda_{j} \|q(t_{j})\|^{p}\right)^{1/p},$$

using that by (4) and (5)

$$\lambda_j \sim |p_{n+1}^{(\alpha,\beta)'}(t_j)|^{-p} |I_j|^{1-p} (1-t_j)^{\gamma p} (1+t_j)^{\delta p}.$$

The error term M_2 can be discretized in view of the monotonicity properties of the f_j 's. The integration with respect to t for $|t - t_j| \leq 2|I_j|$ leads to another term M_{21} of the form (13), and $M_2 \leq M_{21} + M_{22}$ with

$$M_{22} = c_{10} \left(\sum_{i=1}^{n+1} \left(\sum_{j=1, j \neq i}^{n+1} \| y_j \| |I_j| / (t_i - t_j)^2)^p |I_i| w_{\gamma p, \delta p}(t_i) \right)^{1/p} \right)$$

$$= c_{10} \left[\sum_{i=1}^{n+1} \left(\sum_{j=1, j \neq i}^{n+1} a_{ij} \lambda_j^{1/p} \| q(t_j) \| \right)^p \right]^{1/p},$$
(14)

where $a_{ij} = (|I_i|/\lambda_j)^{1/p} w_{\gamma,\delta}(t_i) |I_j|/(|p'_{n+1}(t_j)|(t_i - t_j)^2)$ for $i \neq j$ and γ, δ as before. We claim that $A_n = (a_{ij})_{i,j=1}^{n+1}$ defines a map $A_n : \ell_p^{n+1} \longrightarrow \ell_p^{n+1}$ with norm bounded by a *C* independent of $n \in \mathbb{N}$. Then (14) is bounded by $c_{10}C(\sum_{j=1}^{n+1} \lambda_j || q(t_j) ||^p)^{1/p}$ as required. Calculation using (4) and (5) shows that for $i, j \leq n/2$

$$a_{ij} \sim (\frac{i}{j})^{\eta} \frac{j^2}{(i^2 - j^2)^2}, \qquad \eta := (\alpha + 1/2)(2/p - 1).$$

The restriction on α gives that $-1/2 \leq \eta \leq 2$. This easily implies $|a_{ij}| \leq c_{11}/(i-j)^{3/2}$; for $\eta \geq 0$ or $(\eta < 0 \text{ and } i > j/2)$ one even has the

bound $c_{11}/(i-j)^2$. In any case

$$\sup_{i \le n/2} \sum_{j \le n/2, i \ne j} |a_{ij}| \le C, \qquad \qquad \sup_{j \le n/2} \sum_{i \le n/2, i \ne j} |a_{ij}| \le C,$$

and hence $(a_{ij})_{i,j \le n/2}$ is uniformly bounded on $\ell_1^{[n/2]}$ and $\ell_{\infty}^{[n/2]}$ and thus by interpolation on $\ell_p^{[n/2]}$. The three other cases of pairs (i, j), e.g. $i > n/2 \ge j$, are treated similarly, using the assumption on β as well.

This proves $(2) \Rightarrow (1)$ except for the case of $\alpha \leq -1/2$ or $\beta \leq -1/2$ when (11) and (12) do not hold for $t \notin J_n$. Assume e.g. $\alpha \leq -1/2$. In this case, we estimate the remaining term

$$M_3 := \left(\int_{b_n}^1 \|\sum_{j=1}^{n+1} y_j \frac{p_{n+1}^{(\alpha,\beta)}(t)}{t-t_j} \|^p w_{\alpha\beta}(t) dt\right)^{1/p}$$

by the triangle and the Hölder inequality, using (4), (5), (10) and the fact that for $t \ge b_n |t - t_j|^{-1} \le d(n/j)^2$. We find

$$M_{3} \leq c_{12} n^{-2(1+\alpha)/p} \left(\sum_{j=1}^{n+1} j^{\alpha-1/2} \| x_{j} \|\right)$$

$$\leq c_{13} \left(\sum_{j=1}^{n+1} j^{-(\alpha(2/p-1)+1/p+1/2)p'}\right)^{1/p'} \left(\sum_{j=1}^{n+1} \lambda_{j} \| x_{j} \|^{p}\right)^{1/p}$$

$$\leq c_{14} \left(\sum_{j=1}^{n+1} \lambda_{j} \| x_{j} \|^{p}\right)^{1/p}.$$

where we have used that $\alpha(\frac{2}{p}-1) + \frac{1}{p} - \frac{1}{2} > \frac{1}{p'}$. (b) (1) \Rightarrow (2). For the converse, assume the interpolation inequality (7) to be true.

We claim that (7) implies that the Hilbert matrix $A = ((i-j+1/2)^{-1})_{i,j\in\mathbb{N}}$ defines a bounded operator $A : \ell_p(X) \to \ell_p(X)$. A well-known approximation and scaling argument shows that this is equivalent to the boundedness of the Hilbert transform H in $L_p(\mathbb{R}; \mathbb{X})$, i.e. X is a UMD-space and necessarily 1 . In this sense <math>A is a discrete version of H. For $n \in \mathbb{N}$ we need the zeros $(t_j^{n+1})_{j=1}^{n+1}$ of $p_{n+1}^{(\alpha,\beta)}$ and $(t_i^n)_{i=1}^n$ of $p_n^{(\alpha,\beta)}$, ordered decreasingly as before and the corresponding quadrature weights λ_j^{n+1} and λ_i^n . Let $J_n := \{j \in \mathbb{N} \mid \ltimes / \not \geq \exists \leq \nvDash \ltimes / \not \geq\}$. For any sequence $(x_j)_{j \in J_n} \subseteq X$, consider the X-valued polynomial

$$q := \sum_{j \in J_n} \lambda_j^{-1/p} x_j \ell_j^{n+1} \in \Pi_n(X) \quad , \quad \ell_j^{n+1}(t_i^{n+1}) = \delta_{ij}$$

Applying (7) to q and inequality (6) with (n + 1) replaced by n we find

$$\left(\sum_{i\in J_n}\lambda_i^n \parallel q(t_i^n)\parallel^p\right)^{1/p} \le c_1\left(\int_{-1}^1 \parallel q(t)\parallel^p w_{\alpha\beta}(t)dt\right)^{1/p} \le c_2\left(\sum_{j\in J_n}\lambda_j^{n+1}\parallel q(t_j^n)\parallel^p\right)^{1/p}$$

i.e.

$$\left(\sum_{i\in J_n} \|\sum_{j\in J_n} b_{ij} x_j \|^p\right)^{1/p} \le c_2 \left(\sum_{j\in J_n} \|x_j\|^p\right)^{1/p} \tag{15}$$

with

$$b_{ij} := (\lambda_i^n / \lambda_j^{n+1})^{1/p} \ell_j^{n+1}(t_i^n) = (\lambda_i^n / \lambda_j^{n+1})^{1/p} \frac{p_{n+1}^{(\alpha,\beta)}(t_i^n)}{p_{n+1}^{(\alpha,\beta)'}(t_j^{n+1})(t_j^n - t_j^{n+1})}$$

Let $k_n := |J_n| \sim n/2$. By (15), $B_n := (b_{ij})_{i,j\in J_n}$ satisfies $|| B_n : \ell_p^{k_n}(X) \longrightarrow \ell_p^{k_n}(X) || \le c_2, c_2$ being independent of $n \in \mathbb{N}$. B_n is close to the block $A_n := ((i - j + 1/2)^{-1})_{i,j\in J_n}$ of the Hilbert matrix A. To show this we first evaluate b_{ij} . By Szegö [30, (4,5.7)]

$$(1-t^2)p_n^{(\alpha,\beta)'}(t) = (\eta_n't + \eta_n'')(p_n^{(\alpha,\beta)}(t) - \eta_n p_{n+1}^{(\alpha,\beta)}(t),$$

where $\eta_n, \eta_n', \eta_n'' \in \mathbb{R}$ depend on n and (α, β) , with $\eta_n/n \longrightarrow 1$ for $n \longrightarrow \infty$. Hence, using (5), for $i \leq 3n/4$

$$p_{n+1}^{(\alpha,\beta)}(t_i^n) = \eta_n^{-1} (1 - (t_i^n)^2) p_n^{(\alpha,\beta)'}(t_i^n) \sim (-1)^i (n/i)^{\alpha - 1/2}.$$
 (16)

Thus $b_{ij} = \gamma_i^n / (\delta_j^n n(t_i^n - t_j^{n+1}))$ where $0 < c_3 \leq |\gamma_i^n|, |\delta_j^n| \leq c_4 < \infty$ for $i, j \in J_n$ and the matrices $C_n := (c_{ij})_{i,j\in J_n}, c_{ij} := n^{-1}(t_i^n - t_j^{n+1})^{-1}$, are uniformly bounded on $\ell_p^{k_n}(X)$ as well. By Szegö [30, (8.9.8],

$$t_i^n = \cos \theta_i^n, \qquad \qquad \theta_i^n = \frac{(i+k+\varepsilon_{ni})\pi}{n+\alpha+1/2}, \qquad (17)$$

where k depends on (α, β) only and $\sup_{i \in J_n} |\varepsilon_{ni}| \longrightarrow 0$ for $n \longrightarrow \infty$. For $k_n \times k_n$ matrices D_n and E_n , we write $D_n \approx E_n$ provided that the matrices $D_n - E_n$ are uniformly bounded as maps on $\ell_p^{k_n}(X)$, for any $1 \le p \le \infty$. It suffices to show that $C_n \approx A_n$, then $\sup_{n \in \mathbb{N}} || A_n : \ell_p^{k_n}(X) \longrightarrow \ell_p^{k_n}(X) || \le c_5$, i.e. X is a UMD-space. We have

$$c_{ij} = \frac{1}{2n} \frac{1}{\sin(\theta_i^n - \theta_j^{n+1})/2} \frac{1}{\sin(\theta_i^n + \theta_j^{n+1})/2}.$$

If

$$d_{ij} := \frac{1}{\pi(i-j+1/2)\sin(\theta_i^n + \theta_j^{n+1})/2}, \qquad D_n := (d_{ij})_{i,j \in J_n},$$

then $C_n \approx D_n : 1/\sin[(\theta_i^n + \theta_j^{n+1})/2]$ is uniformly bounded for $i, j \in J_n$, and hence the estimates $|1/\sin x - 1/x| \leq |x|$ for $|x| \leq \pi/4$ and $|1/x - 1/(x+\varepsilon)| \leq 2\varepsilon/x^2$ for x = i - j + 1/2 and $|\varepsilon| \leq 1/4$ yield

$$\begin{aligned} |c_{ij} - d_{ij}| &\leq c_6(|\frac{1}{2n\sin(\theta_i^n - \theta_j^{n+1})/2} - \frac{1}{n(\theta_j^n - \theta_j^{n+1})}| + |\frac{1}{n(\theta_i^n - \theta_j^{n+1})} - \frac{1}{\pi(i - j + 1/2)}|) \\ &\leq c_7(\frac{|i - j + 1/2|}{n^2} + \frac{1}{|i - j + 1/2|^2}). \end{aligned}$$

Hence $\sup_i \sum_j |c_{ij} - d_{ij}| \le c_8$, $\sup_j \sum_i |c_{ij} - d_{ij}| \le c_8$ uniformly in $n \in \mathbb{N}$, i.e. $C_n \approx D_n$ (first for $p = 1, \infty$, then by interpolation for general p). Next D_n is transformed into $E_n = (e_{ij})$ with $D_n \approx E_n$,

$$e_{ij} := \frac{1}{\pi(i-j+1/2)\sin\frac{i+j}{2n}\pi}; \qquad i,j \in J_n.$$

By the Lipschitz continuity of $1/\sin x$ in $\pi/4 \le x \le 3\pi/4$,

$$|d_{ij} - e_{ij}| \le c_9/(n|i - j + 1/2|),$$

which again is uniformly row- and column- summable, i.e. $D_n \approx E_n$. Finally, let

$$f_{ij} := \frac{1}{\pi(i - j + 1/2) \sin \frac{i}{n}\pi}, \qquad F_n := (f_{ij})_{i,j \in J_n}.$$

For $y \in [\pi/8, 3\pi/8]$, $g(x) = 1/\sin(x+y)$ is Lipschitz-continuous in $x \in [\pi/8, 3\pi/8]$ with constant ≤ 2 . Hence for $i, j \in J_n$

$$|e_{ij} - f_{ij}| \le \frac{|i-j|}{n|i-j+1/2|} \le \frac{2}{n}, \qquad E_n \approx F_n.$$

Since $(\sin(\frac{i\pi}{n})^{-1})_{i \in J_n}$ is bounded away from zero, this implies that $A_n = ((i - j + 1/2)^{-1})_{i,j \in J_n}$ defines uniformly bounded maps on $\ell_p^{k_n}(X)$. Hence X is a UMD-space and 1 .

We now prove that (7) implies $\mu(\alpha, \beta) . This is a purely$ $scalar argument, <math>X = \mathbb{R}$. By symmetry, we may assume that $\alpha \geq \beta$; the case of $\alpha \leq -1/2$ $(1 is known already. So let <math>-1/2 < \alpha$. Take $q = p_n^{(\alpha,\beta)} \in \prod_n$ in (7) to show that necessarily $p < M(\alpha,\beta) = 4\frac{\alpha+1}{2\alpha+1}$: Using the second formula of Szegö [30, 4.5.7], one shows similarly as in (16) that $|p_n^{(\alpha,\beta)}(t_i^{n+1})| \sim (n/i)^{\alpha-1/2}$ for $i \leq n/2$ and $\sim (n/(n+2-i))^{\beta-1/2}$ for i > n/2. Thus by (4) and Newman-Rudin [20], cf. also (10),

$$\left(\int_{-1}^{1} |p_{n}^{(\alpha,\beta)}(t)|^{p} w_{\alpha\beta}(t) dt\right)^{1/p} \sim \left\{\begin{array}{cc} 1 & p < M(\alpha,\beta) \\ (\log n)^{1/p} & p = M(\alpha,\beta) \\ n^{p(\alpha+1/2)-2(\alpha+1)} & p > M(\alpha,\beta) \end{array}\right\}, \quad (18)$$

$$\left\{\begin{array}{ll} \sum_{j+1}^{n+1} \lambda_j |p_n^{(\alpha,\beta)}(t_j)|^p)^{1/p} \sim \\ \left\{\begin{array}{ll} 1 & (p < \infty \text{ and } \alpha \le 1/2) \text{ or } (p < 4(\alpha+1)/(2\alpha-1) \text{ and } \alpha > 1/2) \\ (\log n)^{1/p} & p = 4(\alpha+1)/(2\alpha-1) \text{ and } \alpha > 1/2 \\ n^{(\alpha-1/2)-2/p(\alpha+1)} & p > 4(\alpha+1)/(2\alpha-1) \text{ and } \alpha > 1/2 \end{array}\right\}.$$
(19)

Hence for $p \ge M(\alpha, \beta)$, the order of growth (in *n*) in (18) is faster than in (19) and (7) cannot hold. To prove that necessarily $p > \mu(\alpha, \beta) = 4(\alpha + 1)/(2\alpha + 5)$ for $\alpha > 1/2$ ($\mu(\alpha, \beta) = 1$ for $\alpha \le 1/2$), we take

$$q = \ell_1 \in \Pi_n, \qquad \ell_1(t) = p_{n+1}^{(\alpha,\beta)}(t) / ((t-t_1)p_{n+1}^{(\alpha,\beta)'}(t_1)).$$

Clearly, the right side of (7) is $\sim \lambda_1^{1/p} \sim n^{-2/p(\alpha+1)}$ by (4) whereas the asymptotic formulas for $p_n^{(\alpha,\beta)}$ of Szegö [30, 8.21] and (5) yield

$$\left(\int_{-1}^{1} |\ell_{1}(t)|^{p} w_{\alpha\beta}(t) dt\right)^{1/p} \sim \left(\int_{0}^{1-n^{-2}} (1-t)^{\alpha-p/4(2\alpha+5)} dt\right)^{1/p} n^{-(\alpha+5/2)} \\ \sim \left\{ \begin{array}{l} n^{-(\alpha+5/2)} & p < \mu(\alpha,\beta) \\ (\log n)^{1/p} & p = \mu(\alpha,\beta) \\ n^{-2/p(\alpha,\beta)} & p > \mu(\alpha,\beta) \end{array} \right\}.$$
(20)

Hence (20) grows faster in *n* than $\lambda_1^{1/p} \sim n^{-2/p(\alpha+1)}$ if $p \leq \mu(\alpha, \beta)$, i.e. $p > \mu(\alpha, \beta)$ is necessary for (7) to hold. This proves (b) of Theorem 1.

(a). The left interpolation inequality (6) in Theorem 1 is proved as in the scalar case. Nevai's proof in [19] using the mean value theorem, Hölder's inequality and some weighted form of Bernstein's inequality in the p-norm generalizes directly to the vector-valued setting. Just as the scalar result of Khalilova [11] and Potapov [27], the vector-valued form of the Bernstein L_p inequality (lemma 2 in [19]) is proved by interpolating at Tchebychev nodes, using an averaging technique, the triangle inequality in L_p and the Bernstein inequality for the sup-norm. In the vector valued case the latter follows from the scalar version, applying linear functionals and using the Hahn-Banach theorem. We do not give the details, since the proofs of [19], [11] and [27] directly generalize. \Box

Remarks.

(1). If the validity of (7) of Theorem 1 (b),

$$(\int_{I} \| q(t) \|^{p} w_{\alpha\beta}(t) dt)^{1/2} \le c_{p} (\sum_{j=1}^{n+1} \lambda_{j}^{n+1} \| q(t_{j}^{n+1}) \|^{p})^{1/p},$$

is required only for all polynomials $q \in \Pi_k(X)$ with $k \leq n/2$, this holds for all $1 \leq p \leq \infty$ and all Banach spaces, at least if $\alpha, \beta \geq -1/2$. This follows from the boundedness of the generalized de la Vallée-Poussain means in $L_p(X)$ along similar lines as in Zygmund [32], Stein [29] and Askey [2]. Thus the restriction on p and X in Theorem 1 comes from requiring the number of nodes to equal the dimension of Π_n , namely (n + 1). In this way, however, one isomorphically identifies $\Pi_n(X) \subset L_p(X)$ with the space $\ell_p^{n+1}(X)$.

(2). The proof of the necessity of the UMD-condition for inequality (7) of Theorem 1 (b) will work for more general orthogonal polynomials provided

that sufficiently precise information on a fairly large part of the zeros of these is known, like in (17).

The restriction $p < M(\alpha, \beta)$ means geometrically (for $\alpha > -1/2$) that the value $|p_n^{(\alpha,\beta)}(t_1^{n+1})|^p$ is much smaller than the dominating mean value of $|p_n^{(\alpha,\beta)}(t)|^p$ over $(t_1^{n+1}, 1)$ with respect to $w_{\alpha\beta}(t)dt$, if $p \ge M(\alpha, \beta)$.

An immediate corollary to Theorem 1 is the following result on the convergence of interpolating polynomials which in the scalar case is due to Askey [2] and Nevai [19].

Proposition 9 Let X be a UMD-space, $\alpha, \beta > -1$ and $p < M(\alpha, \beta)$. Let $f: (-1, 1) \longrightarrow X$ be continuous. Then the interpolating polynomials of f at the zeros $(t_j)_1^{n+1}$ of $p_{n+1}^{(\alpha,\beta)}$, $I_n f := \sum_{j=1}^{n+1} f(t_j) \ell_j \in \Pi_n(X)$, converge to f in the p-norm,

$$\| f - I_n f \|_{p;\alpha,\beta} = \left(\int_{-1}^1 \| f(t) - I_n f(t) \|^p w_{\alpha\beta}(t) dt \right)^{1/p} \longrightarrow 0.$$

Proof: Approximate f by polynomials $q_n \in \Pi_n(X)$ in the sup-norm, $\| f - q_n \|_{\infty} \longrightarrow 0$. We may assume that $\mu(\alpha, \beta) , since the p-norms get weaker for smaller <math>p$. Using (b) of Theorem 1 and

$$\sum_{j=1}^{n+1} \lambda_j = \int_{-1}^1 w_{\alpha\beta}(t) dt =: M < \infty,$$

we find

$$\| f - I_n f \|_{p;\alpha,\beta} \leq \| f - q_n \|_{p;\alpha,\beta} + \| q_n - I_n f \|_{p;\alpha,\beta}$$

$$\leq M^{1/p} \| f - q_n \|_{\infty} + c_p (\sum_{j=1}^{n+1} \lambda_j \| q_n(t_j) - f(t_j) \|^p)^{1/p}$$

$$\leq (1 + c_p) M^{1/p} \| f - q_n \|_{\infty} \longrightarrow 0.$$

3 Convergence of vector-valued Jacobi series

Proof of Theorem 2: Recall that $Q_n f := \sum_{j=0}^n \langle f, p_j^{(\alpha,\beta)} \rangle p_j^{(\alpha,\beta)}$ for $f \in L_p(I, w_{\alpha,\beta}; X)$. Thus Q_n is the integral operator induced by the ker-

nel $k_n(x,y) = \sum_{j=0}^n p_j^{(\alpha,\beta)}(x) p_j^{(\alpha,\beta)}(y)$ with respect to the measure $d\mu(t) = w_{\alpha\beta}(t) dt$.

 $(2) \Rightarrow (1)$. We sketch the straightforward generalization of the scalar proof of Pollard [26] and Muckenhaupt [18] to the UMD-case. Since $Q_n f \rightarrow f$ on the dense set of X-valued polynomials f, (1) of Theorem 2 is equivalent to

$$\sup_{n \in \mathbb{N}} \| Q_n : L_p(I, w_{\alpha,\beta}; X) \longrightarrow L_p(I, w_{\alpha\beta}, X) \| = c_p < \infty.$$
 (21)

Using the Christoffel-Darboux formula for k_n and the classical analysis of Pollard [26], (21) will follow from the uniform boundedness of the integral operators T_{n1} , T_{n2} , T_{n3} induced by the following kernels as maps in $L_p(I, w_{\alpha\beta}; X)$:

$$k_{n1}(x,y) := p_{n+1}^{(\alpha,\beta)}(x)q_n^{(\alpha,\beta)}(y)/(x-y) , \quad q_n^{(\alpha,\beta)}(y) := (1-y^2)p_n^{(\alpha+1,\beta+1)}(y)$$
$$k_{n2}(x,y) := k_{n1}(y,x) , \quad k_{n3}(x,y) := p_n^{(\alpha,\beta)}(x)p_n^{(\alpha,\beta)}(y).$$

The proof of the uniform boundedness of T_{n1} and T_{n2} is similar to the proof of Theorem 1, (b), (2) \Rightarrow (1). On the intervals J_n defined there (for $\alpha, \beta \geq -1/2, J_n = I$), T_{n1} and T_{n2} are uniformly bounded in p-norm provided that the weighted Hilbert transform kernels $(x - y)^{-1}(w_{\alpha\beta}(x)w_{\alpha\beta}(y))^{1/2}((1 - x^2)/(1 - y^2))^{\pm 1/4}$ (+ for T_{n2} , - for T_{n1}) define bounded operators on $L_p(I; X)$, as follows from (10) the same way as in (b), (2) \Rightarrow (1). In view of the UMD-assumption on X, this will follow from the boundedness of the kernel operator defined by

$$\frac{1}{|x-y|} |(w_{\alpha\beta}(x)w_{\alpha\beta}(y))^{1/2}((1-x^2)/(1-y^2))^{\pm 1/4} - 1|$$

on $L_p(I;X)$. Using again lemma 1, the latter fact is a consequence of

$$-\frac{1}{p} < \alpha(\frac{1}{p} - \frac{1}{2}) \pm \frac{1}{4}, \beta(\frac{1}{p} - \frac{1}{2}) \pm \frac{1}{4} < 1 - \frac{1}{p},$$

i.e. $m(\alpha, \beta) . If e.g. <math>\alpha < -1/2$, the part of $|| T_{ni}f ||_p$, $i \in \{1, 2\}$, on the interval $(1 - n^{-2}, 1)$ outside J_n has to be estimated separately. However, $p_n^{(\alpha,\beta)}$ and $q_n^{(\alpha,\beta)}$ are uniformly bounded in $n \in \mathbb{N}$ there, and a direct application of the continuity of the (unweighted) Hilbert transform suffices. The uniform boundedness of T_{n3} follows from $\sup_{n \in \mathbb{N}} || p_n^{(\alpha,\beta)} ||_p || p_n^{(\alpha,\beta)} ||_{p'} < \infty$ if $m(\alpha, \beta) .$ $(1) \Rightarrow (2)$. Assume that $Q_n f \longrightarrow f$ for all $f \in L_p(I, w_{\alpha\beta}; X)$. By the Banach-Steinhaus theorem, this is equivalent to (21). Using (21), we prove (7) of Theorem 1, which then implies that X is an UMD-space and, in view of the self-duality of (21), that $m(\alpha, \beta) . To show (7), we dualize (6) which holds for all X and p. Let <math>q \in \Pi_n(X)$. Then there is a $g \in L_{p'}(I, w_{\alpha\beta}; X^*)$ which $||g||_{p';\alpha,\beta} = 1$ and

$$J := \left(\int_{-1}^{1} \| q(t) \|^{p} w_{\alpha\beta}(t) dt\right)^{1/p} = \int_{-1}^{1} \langle q(t), g(t) \rangle_{(X,X^{*})} w_{\alpha\beta}(t) dt$$
$$= \int_{-1}^{1} \langle q(t), Q_{n}g(t) \rangle_{(X,X^{*})} w_{\alpha\beta}(t) dt.$$

Since $\langle q, Q_n g \rangle \in \Pi_{2n}$, Gaussian quadrature, Hölder's inequality and (6) as well as (the dual form of) (21) yield

$$J = \sum_{j=1}^{n+1} \lambda_j < q(t_j), Q_n g(t_j) >$$

$$\leq (\sum_{j=1}^{n+1} \lambda_j \| Q_n g(t_j) \|_{X^*}^{p'})^{1/p'} (\sum_{j=1}^{n+1} \lambda_j \| q(t_j) \|_X^p)^{1/p}$$

$$\leq c \| Q_n g \|_{p';\alpha,\beta} (\sum_{j=1}^{n+1} \lambda_j \| q(t_j) \|^p)^{1/p}$$

$$\leq c c_p (\sum_{j=1}^{n+1} \lambda_j \| q(t_j) \|^p)^{1/p}$$

which is (7).

We turn to the equivalence of vector-valued Jacobi means.

Proof of proposition 3: For $\alpha > -1$ and $w_{\alpha} = w_{\alpha,\alpha}$, the map $\psi : L_2(I, w_{\alpha}; X) \longrightarrow L_2(0, \pi; X)$ defined by

$$\psi(g)(s) = (\sin s)^{\alpha + 1/2} g(\cos s) \qquad g \in L_2(I, w_\alpha; X), \qquad s \in [0, \pi],$$

is an isometry. Let $q_n^{(\alpha)} := \psi(p_n^{(\alpha,\alpha)})$ and

$$\mathcal{K}_n^{(\alpha,\beta)}(t,s) := \sum_{j=0}^n q_j^{(\alpha)}(t)q_j^{(\beta)}(s); \qquad \alpha,\beta > -1.$$

These kernels induce uniformly bounded operators on $L_p(0, \pi; X)$ for any $1 , e.g. there is <math>c_p$ such that for all $n \in \mathbb{N}$ and $h \in L_p(0, \pi; X)$

$$\left(\int_{0}^{\pi} \| \int_{0}^{\pi} \mathcal{K}_{n}^{(\alpha,\beta)}(t,s)h(s)ds \|^{p} dt\right)^{1/p} \leq c_{p}\left(\int_{0}^{\pi} \| h(s) \|^{p} ds\right)^{1/p}.$$
 (22)

This follows from the proofs of Theorem 1 and 3 of Gilbert [8]: The scalar proof given there directly generalizes to the X-valued UMD-case since only the L_p -uniform boundedness of the Dirichlet and conjugate Dirichlet kernel operators is used, which holds X-valued for UMD-spaces. In effect, $\mathcal{K}_n^{(\alpha,\beta)}$ is shown in [8] to behave very similar to the Dirichlet kernel. In particular

$$|\mathcal{K}_n^{(\alpha,\beta)}(t,s)| \le d_1/|t-s| \tag{23}$$

where d_1 is independent of $n \in \mathbb{N}$ and $t, s \in [0, \pi]$. We claim that also

$$\left(\int_{0}^{\pi} \|\int_{0}^{\pi} \mathcal{K}_{n}^{(\alpha,\beta)}(t,s)(\sin t/\sin s)^{\gamma+1/p-1/2}h(s)ds \|^{p} dt\right)^{1/p} \leq c_{p}'\left(\int_{0}^{\pi} \|h(s)\|^{p} ds\right)^{1/p}$$
(24)

provided that $|\gamma/2 + 1/p - 1/2| < 1/4$. By (22), this will follow from the uniform boundedness of the difference kernel operators

$$\mathcal{L}_n^{(\alpha,\beta)}(t,s) := \mathcal{K}_n^{(\alpha,\beta)}(t,s)((\sin t / \sin s)^{\gamma+1/p-1/2} - 1)$$

in $L_p(0, \pi; X)$. Using (23) and elementary estimates we obtain the existence of a d_2 such that for $n \in \mathbb{N}$ and $t, s \in [0, \pi/2]$

$$|\mathcal{L}_{n}^{(\alpha,\beta)}(t,s)| \leq \frac{d_{1}}{|t-s|} |(\frac{\sin t}{\sin s})^{\gamma+1/p-1/2} - 1| \leq \frac{d_{2}}{|t-s|} |(\frac{t}{s})^{\gamma+1/p-1/2} - 1|.$$

Hence by lemma 1, the $\mathcal{L}_n^{(\alpha,\beta)}$ -kernels define uniformly bounded integral operators in $L_p(0, \pi/2; X)$ since $-1/p < \gamma + 1/p - 1/2 < 1 - 1/p$. On $(\pi/2, \pi)$, the estimate is similar; for $t \in [\pi/2, \pi], s \in [0, \pi/2]$, there are only point singularities and the transformation $t \to \pi - t$ reduces the $\mathcal{L}_n^{(\alpha,\beta)}$ -boundedness to the one of the positive kernel 1/(t+s) in $L_p(0, \pi/2; X)$. Hence (24) holds.

For functions $f \in L_p(I, w_{(\beta+\gamma)p/2}; X)$ on the interval I = (-1, 1) and the kernel

$$k_n^{(\alpha,\beta)}(x,y) := \sum_{j=0}^n p_j^{(\alpha,\alpha)}(x) p_j^{(\beta,\beta)}(y),$$

(24) is equivalent to

$$(\int_{-1}^{1} \| \int_{-1}^{1} k_{n}^{(\alpha,\beta)}(x,y) f(y) w_{\beta}(y) dy \|^{p} w_{(\alpha+\gamma)p/2}(x) dx)^{1/p} \\ \leq c_{p}' \int_{-1}^{1} \| f(y) \|^{p} w_{(\beta+\gamma)p/2}(y) dy)^{1/p}$$
(25)

as the transformation $h(s) = (\sin s)^{\beta+\gamma+1/p} f(\cos s)$ shows. Applying (25) to $f(y) = \sum_{j=0}^{n} p_j^{(\beta,\beta)}(y) x_j$, where $x_j \in X$, yields a one-sided estimate of (8); the converse direction follows from the symmetry of the statement in α and β . The argument also shows that the convergence of the series $\sum_j p_j^{(\alpha,\alpha)} \otimes x_j$ in $L_p(I, w_{(\alpha+\gamma)p/2}; X)$ is equivalent to the convergence of the series $\sum_j p_j^{(\beta,\beta)} \otimes x_j$ in $L_p(I, w_{(\beta+\gamma)p/2}; X)$, provided that $|\gamma/2 + 1/p - 1/2| < 1/4$. \Box

The choice of p = 2 and $\gamma = 0$ shows that the means

$$\left(\int_{-1}^{1} \|\sum_{j=0}^{n} p_{j}^{(\alpha,\alpha)}(t) x_{j} \|^{2} w_{\alpha}(t) dt\right)^{1/2}$$

are essentially independent of α , the choice of $\gamma = 0$ for 4/3 showsa similar statement for the means

$$\left(\int_{-1}^{1} \|\sum_{j=0}^{n} p_{j}^{(\alpha,\alpha)}(t) x_{j} \|^{p} w_{\alpha p/2}(t) dt\right)^{1/p}.$$

4 Unconditional convergence

We now show that under the conditions of Proposition 4, vector-valued convergence of orthonormal series is unconditional only in the case of Hilbert spaces.

Proof of Proposition 4:

(i). Let (Ω, μ) be a finite measure space and (p_n) be a complete orthonormal system in $L_2(\Omega, \mu)$ such that $\sum_n \langle f, p_n \rangle p_n$ converges unconditionally for all $f \in L_p(\Omega, \mu)$. By duality, the same holds in $L_{p'}(\Omega, \mu)$. Thus we may assume that $p \geq 2$. Using the unconditionality and the Khintchine inequality, we find for any finite sequence $(a_n) \subset \mathbb{K}$

$$\sum_{n} |a_{n}|^{2} = \|\sum_{n} a_{n} p_{n}\|_{2}^{2} \leq c_{1} \|\sum_{n} a_{n} p_{n}\|_{p}^{2}$$

$$\leq c_{2} \left(\int_{0}^{1} \int_{\Omega} |\sum_{n} a_{n} r_{n}(t) p_{n}(w)|^{p} d\mu(w) dt \right)^{2/p}$$

$$\leq c_{3} \left(\int_{\Omega} (\sum_{n} |a_{n}|^{2} |p_{n}(w)|^{2})^{p/2} d\mu(w) \right)^{2/p}$$

$$\leq c_{3} \sum_{n} (\int_{\Omega} |a_{n}|^{p} |p_{n}(w)|^{p} d\mu(w))^{2/p}$$

$$= c_{3} \sum_{n} |a_{n}|^{2} ||p_{n}||_{p}^{2} \leq c_{3} \sup_{n} ||p_{n}||_{p}^{2} \sum_{n} |a_{n}|^{2}$$

$$\leq c_{4} \sum_{n} |a_{n}|^{2},$$

where we have used the triangle inequality in $L_{p/2}$ and the assumption on $|| p_n ||_p$.

Hence $\|\sum_{n} a_n p_n\|_p \sim \|(a_n)\|_{l_2}$, which implies p = 2 since (p_n) was assumed to be a *complete* orthonormal system and $L_p(\Omega, \mu) \sim \ell_2$ only for p = 2. Thus (i) holds, even for $X = \mathbb{K}$.

(ii). We give a modification of the argument of Defant and Junge [6]. Let $x_1, \dots x_m \in X$. By the unconditionality assumption on the (p_j) , the hypothesis that $\sup_j |p_j| \in L_2(\Omega, \mu)$, and the contraction principle, cf. Maurey and Pisier [17] we get,

$$\begin{aligned} (\int_{\Omega} \|\sum_{j=1}^{m} p_j(w) x_j \|^2 \, d\mu(w))^{1/2} &\leq c_1 (\int_{\Omega} \int_{0}^{1} \|\sum_{j=1}^{m} r_j(t) p_j(w) x_j \|^2 \, dt d\mu(w))^{1/2} \\ &\leq c_2 (\int_{\Omega} (\sup_j |p_j(w)|^2) (\int_{0}^{1} \|\sum_{j=1}^{m} r_j(t) x_j \|^2 \, dt) d\mu(w))^{1/2} \\ &\leq c_3 (\int_{0}^{1} \|\sum_{j=1}^{m} r_j(t) x_j \|^2 \, dt)^{1/2}. \end{aligned}$$

Let (γ_j) be a sequence of independent standard N(0, 1) Gaussian variables on a probability space (Γ, ν) . By Pisier [23] with $c_4 = \sqrt{\pi/2} c_3$

$$\left(\int_{\Omega} \|\sum_{j=1}^{m} p_j(w) x_j \|^2 d\mu(w)\right)^{1/2} \le c_4 \left(\int_{\Gamma} \|\sum_{j=1}^{m} \gamma_j(s) x_j \|^2 d\nu(s)\right)^{1/2}$$
(26)

Since $L_2(\Omega, \mu)$ is infinite dimensional, for any $n \in \mathbb{N}$ there is a unitary map

 $U: L_2(\Omega, \mu) \longrightarrow L_2(\Omega, \mu)$ such that with $f_j := Up_j$ the functions f_1, \cdots, f_n are mutually disjointly supported. Let $f_k = \sum_j u_{jk} p_j \in L_2(\Omega, \mu)$, (u_{jk}) unitary. Applying (26) for arbitrary $y_1, \cdots, y_n \in X$ with $x_j := \sum_{k=1}^n u_{jk} y_k$ we find using the unitary invariance of the right side of (26)

$$\begin{split} (\sum_{k=1}^{n} \| y_{k} \|^{2})^{1/2} &= (\int_{\Omega} \| \sum_{k=1}^{n} f_{k}(w) y_{k} \|^{2} d\mu(w))^{1/2} \\ &= (\int_{\Omega} \| \sum_{j} p_{j}(w) x_{j} \|^{2} d\mu(w))^{1/2} \\ &\leq c_{4} (\int_{\Gamma} \| \sum_{j} \gamma_{j}(s) x_{j} \|^{2} d\nu(s))^{1/2} \\ &= c_{4} (\int_{\Gamma} \| \sum_{k=1}^{n} \gamma_{k}(s) y_{k} \|^{2} d\nu(s))^{1/2}, \end{split}$$

i.e. X has cotype 2. Similarly, the converse inequality to (26) will imply that X has type 2 and thus by Kwapień [13] that X is isomorphic to a Hilbert space. By Maurey and Pisier [17], the Gaussian and the Rademacher means are equivalent since X has cotype 2. Using this and Kahane's inequality [15], we get for any $x_1, \dots x_m \in X$

$$(\int_{\Gamma} \| \sum_{j=1}^{m} \gamma_j(s) x_j \|^2 d\nu(s))^{1/2} \le c_5 (\int_{0}^{1} \| \sum_{j=1}^{m} r_j(t) x_j \|^2 dt)^{1/2}$$

$$\le c_6 \int_{0}^{1} \| \sum_{j=1}^{m} r_j(t) x_j \| dt.$$

Since $|| p_j ||_2 = 1$, the contraction principle, the Hölder inequality and the unconditionality assumption yield similarly as in Defant and Junge [6], cf. also Pisier [23], that this is

$$= c_6 \int_0^1 \| \sum_{j=1}^m (\int_\Omega |p_j(w)|^2 d\mu(w)) r_j(t) x_j \| dt$$

$$\leq c_{7} \int_{0}^{1} \int_{\Omega} (\sup_{j} |p_{j}(w)|) \| \sum_{j=1}^{m} p_{j}(w) r_{j}(t) x_{j} \| d\mu(w) dt$$

$$\leq c_{7} \| \sup_{j} |p_{j}| \|_{L_{2}(\Omega,\mu)} (\int_{\Omega} \int_{0}^{1} \| \sum_{j=1}^{m} r_{j}(t) p_{j}(w) x_{j} \|^{2} dt d\mu(w))^{1/2}$$

$$\leq c_{8} (\int_{\Omega} \| \sum_{j=1}^{m} p_{j}(w) x_{j} \|^{2} d\mu(w))^{1/2},$$

i.e. the converse to (26) holds. Hence X is isomorphic to a Hilbert space. \Box As a corollary we find

Proposition 10 Let $\alpha, \beta > -1$, $1 \le p \le \infty$ and X be a Banach space. Assume that for all $f \in L_p(I, w_{\alpha\beta}; X)$, the Jacobi series $\sum_{n=0}^{\infty} \langle f, p_n^{(\alpha,\beta)} \rangle p_n^{(\alpha,\beta)}$ converges unconditionally in $L_p(I, w_{\alpha\beta}; X)$. Then p = 2 and X is isomorphic to a Hilbert space: the expansions converge unconditionally precisely in the Hilbert space situation.

Proof: By Theorem 2, necessary for convergence is $m(\alpha, \beta) .$ For these values of <math>p, the inequality (10) yields e.g. if p > 2

$$\| p_j^{(\alpha,\beta)} \|_{p;\alpha,\beta} \le c_1 \quad \int_{-1}^{1} (1-t)^{\alpha(1-p/2)-p/4} (1+t)^{\beta(1-p/2)-p/4} dt$$
$$= c_2 = c_2 \| p_j^{(\alpha,\beta)} \|_{2;\alpha,\beta} < \infty$$

and

$$\|\sup_{j} |p_{j}^{(\alpha,\beta)}| \|_{2;\alpha,\beta} \le c_{3} (\int_{-1}^{1} (1-t^{2})^{-1/2} dt)^{1/2} = c_{4}.$$

Thus p = 2 and X is a Hilbert space by Proposition 4.

The unconditionality of the Haar system in $L_p(0, 1; X)$, if 1and X is an UMD-space, shows that Proposition 4 does not hold without $conditions being imposed on the system <math>(p_j)$ as done in (i), (ii) there.

For the proof of Propositions 5 and 7, we need the following result due to Lindenstrauss and Pełczyński [14, proof of Theorem 4.2] and Olevskii [21].

Theorem 11 Let $1 \le p \le \infty$ and $(p_n)_{n \in \mathbb{N}}$ be a basis of $L_p(0, 1)$. Let $(h_j)_{j \in \mathbb{N}}$ denote the Haar system on [0, 1], normalized by $||h_j||_p = 1$. For any $0 < \delta < \infty$ 1 there is a block basis sequence $(z_j)_{j\in\mathbb{N}}$ of $(p_n)_{n\in\mathbb{N}}$ such that for every $N\in\mathbb{N}$ there is a measure preserving automorphism $\varphi_N: [0,1] \to [0,1]$ with

$$\sum_{j=1}^{N} \parallel z_j^* \parallel_{p'} \parallel h_j o \varphi_N - z_j \parallel_p \leq \delta,$$

where $(z_j^*) \subset L_{p'}(0,1)$ is biorthogonal to $(z_j) \subset L_p(0,1)$.

Thus for some increasing sequence $(m_j)_{j \in \mathbb{N}_{\mathcal{F}}}$ of integers and scalars $(a_n)_{n \in \mathbb{N}}$

$$z_j = \sum_{n=m_{j-1}+1}^{m_j} a_n p_n$$

is in the above sense close to the Haar system.

Proof of Proposition 5. We will show that the Haar system is unconditional in $L_p(0, 1; X)$. Then, by Maurey [16] X has to be a UMD-space, using also the results of [4] and [5].

Let $N \in \mathbb{N}$, $0 < \delta < 1$ and $x_1, \dots, x_N \in X$. Let φ_N be as in the theorem and put $g_j := h_j o \varphi_N$. Since $(p_n)_{n \in \mathbb{N}}$ is unconditional in $L_p(0, 1; X)$ by assumption, so is the block basic sequence $(z_j)_{j \in \mathbb{N}}$. Hence for any sequence of signs $(\varepsilon_j), \varepsilon_j \in \{+1, -1\}$,

$$\begin{split} (\int_{0}^{1} \| \sum_{j=1}^{N} \varepsilon_{j} h_{j}(t) x_{j} \|^{p} dt)^{1/p} &= (\int_{0}^{1} \| \sum_{j=1}^{N} \varepsilon_{j} g_{j}(t) x_{j} \|^{p} dt)^{1/p} \\ &\leq (\int_{0}^{1} \| \sum_{j=1}^{N} \varepsilon_{j} z_{j}(t) x_{j} \|^{p} dt)^{1/p} + (\int_{0}^{1} \| \sum_{j=1}^{N} \varepsilon_{j}(g_{j}(t) - z_{j}(t)) x_{j} \|^{p} dt \\ &\leq K (\int_{0}^{1} \| \sum_{j=1}^{N} z_{j}(t) x_{j} \|^{p} dt)^{1/p} + \sum_{j=1}^{N} \| g_{j} - z_{j} \|_{p} \| x_{j} \| \\ &\leq (K + \delta) (\int_{0}^{1} \| \sum_{j=1}^{N} z_{j}(t) x_{j} \|^{p} dt)^{1/p}, \end{split}$$

using that $x_j = \langle z_j^*, \sum_{k=1}^N z_k x_k \rangle$ and hence

$$||x_j|| \le ||z_j^*||_p (\int_0^1 ||\sum_{k=1}^N z_k(t)x_k||^p dt)^{1/p}.$$

The constant K is independent of N, (x_j) and (ε_j) . The chain of inequalities can be reversed with all $\varepsilon_j = +1$ to find

$$\begin{aligned} (\int_0^1 \| \sum_{j=1}^N \varepsilon_j h_j(t) x_j \|^p \, dt)^{1/p} &\leq (K+\delta) (\int_0^1 \| \sum_{j=1}^N z_j(t) x_j \|^p \, dt)^{1/p} \\ &\leq (K+\delta) (1+\delta) (\int_0^1 \| \sum_{j=1}^N h_j(t) x_j \|^p \, dt)^{1/p}, \end{aligned}$$

i.e. the Haar system is unconditional.

A similar procedure is used in the

Proof of Proposition 7: Let $N \in \mathbb{N}$, $x_1, \dots, x_N \in X$ and $0 < \delta < 1$. With the same notation as in the previous proof,

$$\left(\int_{0}^{1} \|\sum_{j=1}^{N} h_{j}(t)x_{j}\|^{2} dt\right)^{1/2} \leq (1+\delta)\left(\int_{0}^{1} \|\sum_{j=1}^{N} z_{j}(t)x_{j}\|^{2} dt\right)^{1/2}.$$

with $z_j = \sum_{n=m_{j-1}+1}^{m_j} a_n p_n$ and $(m_j)_{j \in \mathbb{N}_{\not\vdash}} \subset \mathbb{N}$ increasing. Note that

$$\left(\sum_{n=m_{j-1}+1}^{m_j} |a_n|^2\right)^{1/2} = \parallel z_j \parallel_2 \le (1+\delta) \parallel h_j \parallel_2 = (1+\delta).$$

Thus, using that X has (p_n) -type 2, there is K independent of $x_1, \dots x_N \in X$ such that

$$\begin{aligned} (\int_{0}^{1} \| \sum_{j=1}^{N} h_{j}(t) x_{j} \|^{2} dt)^{1/2} &\leq (1+\delta) (\int_{0}^{1} \| \sum_{j=1}^{N} (\sum_{n=m_{j-1}+1}^{m_{j}} a_{n} p_{n}(t)) x_{j} \|^{2} dt)^{1/2} \\ &\leq K(1+\delta) (\sum_{j=1}^{N} \sum_{n=m_{j-1}+1}^{m_{j}} |a_{n}|^{2} \| x_{j} \|^{2})^{1/2} \\ &\leq K(1+\delta)^{2} (\sum_{j=1}^{N} \| x_{j} \|^{2})^{1/2}. \end{aligned}$$

This shows that X has "Haar-type 2" which directly implies type 2 since the Rademacher functions form a block basis of the Haar functions,

$$r_k = \sum_{j=n_{k-1}+1}^{n_k} t_j h_j , \sum_{j=n_{k-1}+1}^{n_k} |t_j|^2 = 1.$$

where $(n_k)_{k\in\mathbb{N}_{\neq}}\subset\mathbb{N}$ is a suitable increasing sequence. Hence for any sequence $y_1,\cdots,y_\ell\in X$

$$(\int_{0}^{1} \| \sum_{k=1}^{\ell} r_{k}(t) y_{k} \|^{2} dt)^{1/2} = (\int_{0}^{1} \| \sum_{k=1}^{\ell} (\sum_{j=n_{k-1}+1}^{n_{k}} t_{j} h_{j}(t)) y_{k} \|^{2} dt)^{1/2}$$

$$\leq K(1+\delta)^{2} (\sum_{k=1}^{\ell} (\sum_{j=n_{k-1}+1}^{n_{k}} |t_{j}|^{2}) \| y_{k} \|)^{1/2}$$

$$= K(1+\delta)^{2} (\sum_{k=1}^{\ell} \| y_{k} \|^{2})^{1/2}.$$

We note that if conversely X has type 2, one has as estimate of the Rademacher against the Haar mean, i.e. there is a constant C such that for all $\ell \in \mathbb{N}$ and all $(y_k)_{k=1}^{\ell} \subseteq X$:

$$\int_0^1 \|\sum_{k=1}^{\ell} r_k(t) y_k\|^2 dt \le C \int_0^1 \|\sum_{k=1}^{\ell} h_k(t) y_k\|^2 dt.$$

Indeed, this statement is equivalent to the existence of a constant C_1 , so that for all $\ell \in \mathbb{N}$ and all $f \in L_2(0, 1; X)$

$$\left(\int_{0}^{1} \|\sum_{k=1}^{\ell} r_{k}(t) < f, h_{k} > \|^{2}\right)^{1/2} \le C_{1}(\|f\|_{2})^{1/2}$$

Let $f \in L_2(0,1;X)$ be of the form $f := \sum_{j=1}^n x_j \otimes f_j$, where $(f_j)_{j=1}^n \subseteq L_2(0,1)$ is a finite sequence of normalized, mutually disjointly supported functions and $(x_j)_{j=1}^n \subseteq X$. Since the set of such functions is a dense subspace of $L_2(0,1;X)$ it suffices to prove the inequality for those. Let (γ_j) be a sequence of standard independent N(0, 1) Gaussian variables on a probability space (Γ, ν) and let M be the type 2 constant of X. Since for every $x^* \in X^*$

$$\begin{split} \sum_{k=1}^{\ell} |x^*(\langle f, h_k \rangle)|^2 &= \sum_{k=1}^{\ell} |\langle x^*f, h_k \rangle|^2 \\ &\leq \int_0^1 |x^*(f(t))|^2 dt \\ &= \sum_{j=1}^n |x^*(x_j)|^2, \end{split}$$

it follows from the unitary invariance of the γ_j 's (see e.g. [17]) that

$$\int_{\Gamma} \|\sum_{k=1}^{\ell} \gamma_k(s) < f, h_k > \|^2 d\nu(s) \le \int_{\Gamma} \|\sum_{j=1}^{n} \gamma_j(s) x_j \|^2 d\nu(s).$$

Combining this with the fact that since X is of type 2 the Rademacher and the Gauss means are K-equivalent for a suitable K, we obtain

$$\begin{split} (\int_{0}^{1} \| \sum_{k=1}^{\ell} r_{k}(t) < f, h_{k} > \|^{2} dt)^{1/2} &\leq K(\int_{\Gamma} \| \sum_{k=1}^{\ell} \gamma_{k}(s) < f, h_{k} > \|^{2} d\nu(s))^{1/2} \\ &\leq K(\int_{\Gamma} \| \sum_{j=1}^{n} \gamma_{j}(s) x_{j} \|^{2} d\nu(s))^{1/2} \\ &\leq KM(\sum_{j=1}^{n} \| x_{j} \|^{2})^{1/2} = \| f \|_{2}, \end{split}$$

which proves the claim.

The partial converse of Proposition 7 follows easily:

Proof of Proposition 8: Let $N \in \mathbb{N}$, $x_1, \dots, x_N \in X$. By the unconditionality assumption on the (p_n) -system in $L_2(0, 1; X)$ and the type property of X there are constants c_1, c_2 independent of N and $x_1, \dots, x_N \in X$ such that

$$\left(\int_{0}^{1} \|\sum_{j=1}^{N} p_{j}(t)x_{j}\|^{2} dt\right)^{1/2} \leq c_{1}\left(\int_{0}^{1} \int_{0}^{1} \|\sum_{j=1}^{N} r_{j}(s)p_{j}(t)x_{j}\|^{2} ds dt\right)^{1/2}$$

$$\leq c_2 \left(\int_0^1 \sum_{j=1}^N |p_j(t)|^2 \| x_j \|^2 dt \right)^{1/2}$$
$$= c_2 \left(\sum_{j=1}^N \| x_j \|^2 \right)^{1/2}.$$

Hence X has (p_n) -type 2.

We still have to show that in certain cases X is isomorphic to a Hilbert space provided that it only has (p_n) -type 2. This will be another application of the interpolation inequalities of Theorem 1.

Proof of Proposition 6: For $n \in \mathbb{N}$, let $(t_j)_{j=1}^{n+1}$ denote the Gaussian quadrature weights. The $(n + 1) \times (n + 1)$ matrix $A_n = (a_{ij})$ defined by $a_{jk} := \sqrt{\lambda_j} p_k^{(\alpha,\beta)}(t_j), j = 1, \dots, n+1, k = 0, \dots, n$ is orthogonal since by Gaussian quadrature for $k, \ell \in 0, \dots, n$

$$\delta_{k\ell} = \int_{-1}^{1} p_k^{(\alpha,\beta)}(t) p_\ell^{(\alpha,\beta)}(t) w_{\alpha\beta}(t) dt$$
$$= \sum_{j=1}^{n+1} \lambda_j p_k^{(\alpha,\beta)}(t_j) p_\ell^{(\alpha,\beta)}(t_j)$$
$$= \sum_{j=1}^{n+1} a_{jk} a_{j\ell}.$$

Since the measure space $(I, w_{\alpha\beta})$ is equivalent to (0, 1), we know from Proposition 7 that X has type 2. We will now show that X also has cotype 2 and hence by Kwapień [13] is isomorphic to a Hilbert space. We use Theorem 1 (a) to discretize the notion of $(p_n^{(\alpha,\beta)})$ -type 2 and reverse the inequality using the orthogonality of the matrix A_n appearing in this way: By Theorem 1 and the $(p_n^{(\alpha,\beta)})$ -type 2 property there are c_1, c_2 such that for any $n \in \mathbb{N}$ and $x_0, \dots, x_n \in X$

$$(\sum_{j=1}^{n+1} \| \sum_{k=0}^{n} a_{jk} x_k \|^2)^{1/2} = (\sum_{j=1}^{n+1} \lambda_j \| \sum_{k=0}^{n} p_k^{(\alpha,\beta)}(t_j) x_k \|^2)^{1/2}$$

$$\leq c_1 (\int_{-1}^{1} \| \sum_{k=0}^{n} p_k^{(\alpha,\beta)}(t) x_k \|^2 w_{\alpha\beta}(t) dt)^{1/2}$$

$$\leq (\sum_{k=0}^{n} || x_k ||^2)^{1/2}.$$
(27)

Since $A_n^{-1} = A_n^{t}$, we can invert (27) easily: starting with arbitrary $y_1, \dots, y_{n+1} \in X$ and applying (27) to $x_k := \sum_{\ell=1}^{n+1} a_{\ell k} y_{\ell}, k \in 0, \dots, n$, we find

$$\left(\sum_{j=1}^{n+1} \parallel y_j \parallel^2\right)^{1/2} \le c_2 \left(\sum_{k=0}^n \parallel \sum_{\ell=1}^{n+1} a_{\ell k} y_\ell \parallel^2\right)^{1/2}.$$

If the A_n were symmetric, this and Theorem 1 (a) would yield the cotype 2 property. However, $A_n \neq A_n^{t}$, in general. To prove the cotype 2 property, we replace y_j by $r_j(s)y_j$ and apply the contraction principle to find

$$\begin{aligned} (\sum_{j=1}^{n+1} \| y_j \|^2)^{1/2} &\leq c_2 (\sum_{k=0}^n \int_0^1 \| \sum_{\ell=1}^{n+1} a_{\ell k} r_\ell(s) y_\ell \|^2 \, ds)^{1/2} \\ &\leq c_3 (\sum_{k=0}^n \sup_{\ell \leq n+1} |a_{\ell k}|^2)^{1/2} (\int_0^1 \| \sum_{\ell=1}^{n+1} r_\ell(s) y_\ell \|^2 \, ds)^{1/2}. \end{aligned}$$

Hence X will have cotype 2 provided that $\sum_{k=0}^{n} \sup_{\ell \leq n+1} |a_{\ell k}|^2$ is uniformly bounded in $n \in \mathbb{N}$. This is correct since by (4) and (10) e.g. if $\ell \leq n/2$

$$\sqrt{\lambda_{\ell}} \sim \ell^{\alpha+1/2} / n^{\alpha+1} , |p_k^{(\alpha,\beta)}(t_{\ell})| \stackrel{<}{\sim} (n/\ell)^{\alpha+1/2}$$
$$|a_{\ell k}| = \sqrt{\lambda_{\ell}} |p_k^{(\alpha,\beta)}(t)| \stackrel{<}{\sim} n^{-1/2}$$

The case of $\ell > n/2$ is similar. By the result of Kwapień, used earlier, X is isomorphic to a Hilbert space.

References

- [1] D. J. Aldous; Unconditional bases and martingales in $L_p(F)$, Math. Proc. Camb. Phil. Soc. 85 (1979),117 123.
- [2] R. Askey; Mean convergence of orthogonal series and Lagrange interpolation, Acta Math. Sci. Hungar. 23 (1972), 71 - 85.

- [3] A. I. Benedek, E. R. Murphy, R. Panzone; *Cuestions del analisis de Fourier*, Notas de Algebra y Analysis 5 (1974), Univ. Bahia Blanca, Argentina.
- [4] J. Bourgain; Some remarks on Banach spaces in which martingale difference sequences are unconditional, Arkiv för Matematik 21 (1983), 163
 - 168.
- [5] D. L. Burkholder; A geometrical condition that implies the existence of certain singular integrals of Banach-space-valued functions, Proc. Conf. Harm. Anal., Univ. of Chicago, 1981, 271 - 286.
- [6] M. Defant, M. Junge; Unconditional orthonormal systems, preprint, Kiel, 1991.
- [7] D. L. Fernandez, J. B. Garcia; Interpolation of Orlicz-valued function spaces and U.M.D. property, Stud. Math. 99 (1991), 23 - 39.
- [8] J. E. Gilbert; Maximal theorems for some orthogonal series I, Transact. AMS 145 (1969), 495 - 515.
- [9] J. Garcia-Cuerva, J. L. de Francia; Weighted norm inequalities and related topics, North Holland, 1985.
- [10] G. H. Hardy, J. E. Littlewood, G. Polya; *Inequalities*, Cambridge University Press, 1973.
- [11] B. Khalilova; On some estimates for polynomials, Izv. Akad. Nauk Azerb. SSR 2 (1974), 46 - 55 (Russian)
- [12] H. König; Vector-valued Lagrange interpolation and mean convergence of Hermite series, preprint, Kiel, 1992.
- [13] S. Kwapień; Isomorphic characterizations of inner product spaces by orthogonal series with vector-valued coefficients, Stud. Math. 44 (1972), 583 - 595.
- [14] J. Lindenstrauss, A. Pełczyński; Contributions to the theory of the classical Banach spaces, Journ. Funct. Anal. 8 (1971), 225 - 249.
- [15] J. Lindenstrauss, L. Tzafriri; *Classical Banach spaces II*, Springer, 1979.

- [16] B. Maurey; Système de Haar, Séminaire Maurey-Schwarz 1974/75, exp. I, II.
- [17] B. Maurey, G. Pisier; Séries de variables aléatoires vectorielles indépendantes et propriétés géométriques des espaces de Banach, Stud. Math. 58 (1976), 45 - 90.
- [18] B. Muckenhaupt; Mean convergence of Jacobi series, Proc. AMS 23 (1969), 306 - 310.
- [19] G. P. Nevai; Mean convergence of Lagrange interpolation I, Journ. Appr. Th. 18 (1976), 363 - 376.
- [20] J. Newman, W. Rudin; Mean convergence of orthogonal series, Proc. AMS 3 (1952), 219 - 222.
- [21] A. M. Olevskii; Fourier series and Lebesgue functions, Uspehi Mat. Nauk 22 (1967), 237 - 239 (Russian).
- [22] A. Pietsch; Personal communication
- [23] G. Pisier; *Type des espaces normés*, Compt. Rendues A 276 (1973), 1973
 1976.
- [24] G. Pisier; Martingales with values in uniformly convex spaces, Israel Journ. Math. 20 (1975), 326 - 350.
- [25] G. Pisier, Q. Xu; The strong p-th variation of martingales and orthogonal series, Probab. Th. Rel. Fields 77 (1988), 497 - 524.
- [26] H. Pollard; Mean convergence for orthogonal series: II, Transact AMS 63 (1948), 355 367 and III, Duke Journ. Math. 16 (1949), 189 191.
- [27] M. K. Potapov; Some inequalities for polynomials and their derivatives, Vestnik Mosk. Univ. Mat. Mek. 2 (1960), 10 - 20 (Russian).
- [28] J. Schwartz; A remark om inequalities of Calderon-Zygmund type for vector-valued functions, Comm. Pures Appl. Math. 14 (1961), 785 799.
- [29] E. M. Stein; Interpolation in polynomial classes and Markoff's inequality, Duke J. Math. 24 (1957), 467 - 476.

- [30] G. Szegö; Orthogonal polynomials, AMS, Providence, 1959.
- [31] A. Zygmund; Trigonometric series, Cambridge University Press, 1968.
- [32] A. Zygmund; A property of the zeros of the Legendre polynomials, Transact. AMS 54 (1943), 39 - 56.

Hermann König Mathematisches Seminar Universität Kiel Ludewig Meyn Str. 4 23 Kiel Germany Niels Jørgen Nielsen Matematisk Institut Odense Universitet Campusvej 55 5230 Odense M Denmark