## A Dvoretsky Theorem for Polynomials by Seán Dineen\* (University College Dublin)

## Abstract

We lift upper and lower estimates from linear functionals to n-homogeneous polynomials and using this result show that  $l_{\infty}$  is finitely represented in the space of n-homogeneous polynomials,  $n \geq 2$ , for any infinite dimensional Banach space. Refinements are also given.

The classical Dvoretsky spherical sections theorem [5,13] states that  $l_2$  is finitely represented in any infinite dimensional Banach space. Using this, the Riesz Representation theorem (for finite dimensional  $l_p$  spaces) and the Hahn-Banach theorem we show that  $l_{\infty}$ is finitely represented in  $\mathcal{P}(^n E)$ , for any infinite dimensional Banach space and any  $n \geq 2$ . This shows that  $\mathcal{P}(^n E)$  does not have any non-trivial superproperties and explains why spaces such as Tsirelson's space play such a positive role in the recent theory of polynomials on Banach spaces ([1,2,6,7,8,9,10]). We refer to [3,11,12] for properties of Banach spaces and to [4] for properties of polynomials.

**Theorem 1.** Suppose E is a Banach space,  $1 , <math>\{\phi_j\}_{j=1}^k$  is a finite sequence of vectors in E' and A and B are positive constants such that

$$A^p \sum_{j=1}^k |\alpha_j|^p \le \left\| \sum_{j=1}^k \alpha_j \phi_j \right\|^p \le B^p \sum_{j=1}^k |\alpha_j|^p \tag{1}$$

for any sequence of scalars  $(\alpha_j)_{j=1}^k$ . Then for any integer  $n, n \ge q$  where  $\frac{1}{p} + \frac{1}{q} = 1$ , and any sequence of scalars  $(\alpha_j)_{j=1}^k$  we have

$$A^{n} \sup_{1 \le j \le k} |\alpha_{j}| \le \left\| \sum_{j=1}^{k} \alpha_{j} \phi_{j}^{n} \right\| \le B^{n} \sup_{1 \le j \le k} |\alpha_{j}|$$

$$\tag{2}$$

<sup>\*</sup> The author thanks Universidade Federal do Rio de Janeiro for support during the period this research was initiated.

**Proof.** For any  $x \in E$ ,  $||x|| \le 1$ , we have

$$\sup_{\sum_{j=1}^{k} |\alpha_j|^p \le 1} \left| \sum_{j=1}^{k} \alpha_j \phi_j(x) \right|^p \le B^p.$$

Since  $(l_p)' = l_q$  this implies

$$\sup_{\|x\| \le 1} \sum_{j=1}^{k} \left| \phi_j(x) \right|^q \le B^q.$$

If  $n \ge q$  then

$$\sup_{\|x\|\leq 1} \left| \sum_{j=1}^{k} \alpha_{j} \phi_{j}^{n}(x) \right| \leq \sup_{j} |\alpha_{j}| \cdot B^{n} \sup_{\|x\|\leq 1} \sum_{j=1}^{k} \left| \frac{\phi_{j}(x)}{B} \right|^{n} \leq B^{n} \sup_{j} |\alpha_{j}|.$$

$$(3)$$

On the other hand

$$A^{p} \sum_{j=1}^{k} |\alpha_{j}|^{p} = A^{p} \sup_{\sum_{j=1}^{k} |\beta_{j}|^{q} \le 1} \left| \sum_{j=1}^{k} \alpha_{j} \beta_{j} \right|^{p} \le \sup_{\|x\| \le 1} \left| \sum_{j=1}^{k} \alpha_{j} \phi_{j}(x) \right|^{p}$$

Since the set  $\left\{ \left( \phi_j(x) \right)_{j=1}^k; \|x\| \le 1 \right\}$  is a convex balanced set, the Hahn-Banach theorem implies that

$$A \cdot B_{l_q^k} = A \cdot \left\{ (\beta_j)_{j=1}^k; \sum_{j=1}^k |\beta_j|^q \le 1 \right\} \subset \overline{\left\{ (\phi_j(x))_{j=1}^k; \|x\| \le 1 \right\}}$$

Hence, for any fixed integer  $l, 1 \leq l \leq k$ , there exists  $(x_n)_n$  in  $E, ||x_n|| \leq 1$ , such that

$$\phi_l(x_n) \to A \text{ and, for } j \neq l, \quad \phi_j(x_n) \to 0 \text{ as } n \to \infty.$$

This implies that

$$\sup_{\|x\| \le 1} \left| \sum_{j=1}^{k} \alpha_j \phi_j^n(x) \right| \ge A^n \cdot \sup_{1 \le j \le k} |\alpha_j| \tag{4}$$

and the inequalities (3) and (4) prove the proposition.

Note that in the proof of theorem 1 we have actually shown that the left (resp. right) hand side of the inequality (1) implies the left (resp. right) hand side of the inequality (2). Conditions (1) and (2) can be rephrased in terms of the Banach-Mazur distance d to give the following result.

**Corollary 2.** Let *E* denote a *k*-dimensional Banach space and suppose  $d(E, l_p^k) \leq C$ where  $1 \leq p < \infty$ . Then, for  $n \geq p$ ,  $\mathcal{P}(^nE)$  contains a *k*-dimensional subspace *F* such that

$$d(F, l_{\infty}^k) \le C^n.$$

**Corollary 3.** If E is an infinite dimensional Banach space and  $n \ge 2$  then  $l_{\infty}$  is finitely represented in  $\mathcal{P}(^{n}E)$ .

**Proof.** By the classical Dvoretsky theorem we can choose for any positive integer k and any  $\epsilon > 0$  vectors  $\{\phi_j\}_{j=1}^k$  in E' such that

$$\sum_{j=1}^{k} |\alpha_j|^2 \le \left\| \sum_{j=1}^{k} \alpha_j \phi_j \right\|^2 \le (1+\epsilon)^2 \sum_{j=1}^{k} |\alpha_j|^2$$

for any sequence of scalars  $(\alpha_j)_{j=1}^k$ . Hence, by theorem 1, we have, for  $n \ge 2$  and any  $(\alpha_j)_{j=1}^k$ ,

$$\sup_{1 \le j \le k} |\alpha_j| \le \left\| \sum_{j=1}^k \alpha_j \phi_j^n \right\| \le (1+\epsilon)^n \sup_{1 \le j \le k} |\alpha_j|$$

This proves the corollary.

In fact it is easily seen that the above shows that  $l_{\infty}$  is finitely represented in the space of polynomials of finite type.

**Corollary 4.** If  $l_p$ ,  $1 \le p < \infty$ , is a quotient of E then  $l_\infty$  is a subspace of  $\mathcal{P}(^nE)$ ,  $n \ge p$ , and  $l_1$  is a complemented subspace of the completed symmetric tensor product endowed with the projective topology,  $\bigotimes_{\substack{n \le \pi}} E$ .

**Proof.** Let  $\frac{1}{p} + \frac{1}{q} = 1$ . We can choose constants A and B independent of k and vectors  $(\phi_k)_k$  in E' such that

$$A^{q} \cdot \sum_{j=1}^{k} |\alpha_{j}|^{q} \le \left\| \sum_{j=1}^{k} \alpha_{j} \phi_{j} \right\|^{q} \le B^{q} \cdot \sum_{j=1}^{k} |\alpha_{j}|^{q}$$

for any sequence of scalars  $(\alpha_j)_j$ . Theorem 1 implies that, for  $n \ge p$ ,

$$A^n \sup_{1 \le j \le k} |\alpha_j| \le \left\| \sum_{j=1}^k \alpha_j \phi_j^n \right\| \le B^n \sup_{1 \le j \le k} |\alpha_j|$$

for any sequence of scalars  $(\alpha_j)_j$  and any k. Hence  $\{\phi_j^n\}_{j=1}^{\infty}$  is equivalent to the unit vector basis of  $c_0$ . Since  $\mathcal{P}(^nE)$  is a dual space this implies  $l_{\infty} \hookrightarrow \mathcal{P}(^nE)$ . Since  $\left( \bigotimes_{n,s,\pi} E \right)' \cong$   $\mathcal{P}(^{n}E)$  it follows by [3, p. 48] that  $l_{1}$  is a complemented subspace of  $\bigotimes_{n,s,\pi} E$ . This completes the proof.

This result for p = 2 is given in [8, proposition 13] and also implies the well known fact that  $\mathcal{P}({}^{n}l_{p})$  is not reflexive if  $n \geq p$ .

We now extend the result given in corollary 3 and at the same time obtain a refinement of [6, theorem 1(ii)]. The elements of  $\bigotimes_{n,s,\epsilon} E'$  are *n*-homogeneous polynomials on Ewhich are uniformly weakly continuous on bounded subsets of E. Hence they have unique extensions to E''. We use the notation  $\tilde{P}$  to denote this extension.

**Lemma 5.** A bounded sequence  $(P_j)_j$  in  $\bigotimes_{n,s,\epsilon} E'$  is a weakly null sequence if and only if  $\tilde{P}_j(x'') \to 0$  as  $j \to \infty$  for any  $x'' \in E''$ .

**Proof.** If  $\phi \in \left( \bigotimes_{n,s,\epsilon} E' \right)'$  then there exists a regular Borel measure  $\mu$  on  $\left( \overline{B}_{E''}, \sigma(E'', E') \right)$  such that

$$\phi(P) = \int_{\overline{B}_{E''}} \tilde{P}(x'') \, d\mu(x'')$$

for all  $P \in \widehat{\bigotimes}_{n,s,\epsilon} E'$ .

If  $(P_j)_j$  is bounded then  $(\tilde{P}_j)_j$  is uniformly bounded on  $\overline{B}_{E''}$ . If  $\tilde{P}_j(x'') \to 0$  as  $j \to \infty$  for each x'' in  $\overline{B}_{E''}$  then the Lebesgue dominated convergence theorem implies that  $\phi(P_j) \to 0$  as  $j \to \infty$ . Hence  $(P_j)_j$  is weakly null. The converse is obvious.

**Proposition 6.** If *E* is a Banach space and *E'* contains a weakly null sequence of unit vectors, which satisfies an upper *q* estimate,  $q < \infty$ , then for  $n \ge p$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , we have  $l_{\infty} \hookrightarrow \mathcal{P}(^{n}E)$ .

**Proof.** If suffices to show  $c_0 \hookrightarrow \bigotimes_{n,s,\epsilon} E'$ . Let  $(\phi_j)_j$  denote a weakly null sequence of unit vectors in E' which satisfies an upper q-estimate. Then  $(\phi_j^n)_j$  is a sequence of unit vectors in  $\bigotimes_{n,s,\pi} E'$ . Since  $(\phi_j)_j$  is weakly null lemma 5 implies that  $(\phi_j^n)_j$  is a weakly null sequence

in  $\bigotimes_{n,s,\epsilon} E'$ . By the Bessaga-Pelczynski selection principle [3, p. 42 and 11, p. 5] the sequence

 $(\phi_j^n)_j$  contains a subsequence which forms a basic sequence. Since upper q estimates are inherited by subsequences we may suppose that  $(\phi_j^n)_j$  is a basic sequence.

Hence there exists B > 0 such that

$$\left\|\sum_{j=1}^{k} \alpha_{j} \phi_{j}\right\|^{q} \leq B^{q} \sum_{j=1}^{k} |\alpha_{j}|^{q}$$

for any integer k and any sequence of scalars  $(\alpha_j)_j$ .

By theorem 1 we have

$$\left\|\sum_{j=1}^{k} \alpha_{j} \phi_{j}^{n}\right\| \leq B^{n} \sup_{1 \leq j \leq n} |\alpha_{j}|$$

for any sequence of scalars  $(\alpha_j)_j$ .

Since  $(\phi_j^n)_j$  is a basic sequence the closed subspace of  $\bigotimes_{n,s,\epsilon} E'$  spanned by  $(\phi_j^n)_j$  is isomorphic to  $c_0$ . This completes the proof.

Proposition 5 applies in particular to reflexive Banach lattices which satisfy a lower *p*-estimate  $\frac{1}{p} + \frac{1}{q} = 1$  ([12]).

## **Bibliography**

- R. Alencar, R. M. Aron, S. Dineen, A reflexive space of holomorphic functions in infinitely many variables, Proc. A. M. S., 90, 3, 1984, 407–411.
- [2] R. M. Aron, S. Dineen, *Q*-reflexive Banach spaces, preprint.
- [3] J. Diestel, Sequences and Series in Banach Spaces, Springer, Graduate Text in Mathematics, 1984.
- [4] S. Dineen, Complex Analysis in locally convex spaces, North Holland Math Studies, 57, 1981.
- [5] A. Dvoretzty, Some results on convex bodies and Banach Spaces, Proc. International Symposium on Linear spaces, Jerusalem, 1961, 123–160.
- [6] J. Farmer, Polynomial Reflexivity in Banach Spaces, preprint.
- [7] J. Farmer, W. B. Johnson, Polynomial Schur and Polynomial Dunford-Pettis properties, Contemporary Mathematics, 144, AMS, 1993, 95–105.
- [8] M. Gonzalez, J. Gutierrez, Unconditionally converging Polynomials on Banach spaces, preprint.
- [9] R. Gonzalo, J. A. Jarmillo, Compact Polynomials between Banach spaces, preprint.
- [10] J. Jaramillo, A. Prieto, Weak polynomial convergence on a Banach space, preprint.
- [11] J. Lindenstrauss, L. Tzafriri, Classical Banach spaces I, "Sequence Spaces", Springer Verlag, 1977.
- [12] L. Lindenstrauss, L. Tsafriri, Classical Banach Spaces II, Springer, Graduate Text in Mathematics, 1979.
- [13] G. Pisier, The volume of convex bodies and Banach space geometry, Cambridge Tracts in Mathematics, 94, 1989.

Department of Mathematics, University College Dublin, Belfield, Dublin 4, Ireland. e-mail <sdineen@irlearn.bitnet>