## EXAMPLES OF DOMAINS WITH NON-COMPACT AUTOMORPHISM GROUPS

SIQI FU, A. V. ISAEV, AND S. G. KRANTZ

ABSTRACT. We give an example of a bounded, pseudoconvex, circular domain in  $\mathbb{C}^n$  for any  $n \geq 3$  with smooth real-analytic boundary and non-compact automorphism group, which is not biholomorphically equivalent to any Reinhardt domain. We also give an analogous example in  $\mathbb{C}^2$ , where the domain is bounded, non-pseudoconvex, is not equivalent to any Reinhardt domain, and the boundary is smooth real-analytic at all points except one.

Let D be a bounded or, more generally, a hyperbolic domain in  $\mathbb{C}^n$ . Denote by  $\operatorname{Aut}(D)$  the group of biholomorphic self-mappings of D. The group  $\operatorname{Aut}(D)$ , with the topology given by uniform convergence on compact subsets of D, is in fact a Lie group [Kob].

A domain D is called Reinhardt if the standard action of the n-dimensional torus  $\mathbb{T}^n$  on  $\mathbb{C}^n$ ,

$$z_j \mapsto e^{i\phi_j} z_j, \qquad \phi_j \in \mathbb{R}, \quad j = 1, \dots, n,$$

leaves *D* invariant. For certain classes of domains with non-compact automorphism groups, Reinhardt domains serve as standard models up to biholomorphic equivalence (see e.g. [R], [W], [BP], [GK1], [Kod]).

It is an intriguing question whether any domain in  $\mathbb{C}^n$  with non-compact automorphism group and satisfying some natural geometric conditions is biholomorphically equivalent to a Reinhardt domain. The history of the study of domains with non-compact automorphism groups shows that there were expectations that the answer to this question would be positive (see [Kra]). In this note we give examples that show that the answer is in fact negative.

While the domain that we shall consider in Theorem 1 below has already been noted in the literature [BP], it has never been proved that this domain is not biholomorphically equivalent to a Reinhardt domain. Note that this domain is circular, i.e. it is invariant under the special rotations

$$z_j \mapsto e^{i\phi} z_j, \qquad \phi \in \mathbb{R}, \quad j = 1, \dots, n.$$

Our first result is the following

Mathematics Subject Classification: 32A07, 32H05, 32M05

Keywords and Phrases: Automorphism groups, Reinhardt domains, circular domains.

**Theorem 1.** There exists a bounded, pseudoconvex, circular domain  $\Omega \subset \mathbb{C}^3$  with smooth real-analytic boundary and non-compact automorphism group, which is not biholomorphically equivalent to any Reinhardt domain.

*Proof.* Consider the domain

$$\Omega = \{|z_1|^2 + |z_2|^4 + |z_3|^4 + (\overline{z_2}z_3 + \overline{z_3}z_2)^2 < 1\}.$$

The domain  $\Omega$  is invariant under the action of the two-dimensional torus  $\mathbb{T}^2$ 

$$z_1 \mapsto e^{i\phi_1} z_1, \qquad \phi_1 \in \mathbb{R},$$
  
 $z_j \mapsto e^{i\phi_2} z_j, \qquad \phi_2 \in \mathbb{R}, \quad j = 2, 3,$ 

and therefore is circular. It is also a pseudoconvex, bounded domain with smooth real-analytic boundary. The automorphism group  $\operatorname{Aut}(\Omega)$  is non-compact since it contains the following subgroup

$$z_{1} \mapsto \frac{z_{1} - a}{1 - \overline{a}z_{1}},$$

$$z_{2} \mapsto \frac{(1 - |a|^{2})^{\frac{1}{4}}z_{2}}{(1 - \overline{a}z_{1})^{\frac{1}{2}}},$$

$$z_{3} \mapsto \frac{(1 - |a|^{2})^{\frac{1}{4}}z_{3}}{(1 - \overline{a}z_{1})^{\frac{1}{2}}},$$

$$(1)$$

for a complex parameter a with |a| < 1.

The above argument also shows that any linear automorphism of  $\Omega$  can be written as

$$z_1 \mapsto e^{i\phi_1} z_1,$$
  

$$z_2 \mapsto az_2 + bz_3,$$
  

$$z_3 \mapsto cz_3 + dz_3,$$

where  $\phi_1 \in \mathbb{R}$ ,  $a, b, c, d \in \mathbb{C}$ , and the transformation in the variables  $(z_2, z_3)$  is an automorphism of the section  $\Omega \cap \{z_1 = 0\}$ . Further, since the only points of  $\partial \Omega$  where rank  $\mathcal{L}_{\partial \Omega} = 1$  are those of the form  $(z_1, w, \pm w)$  with  $w \neq 0$  and since automorphisms of  $\Omega$  preserve such points, it follows that any linear automorphism of  $\Omega$  is in fact given by

$$z_1 \mapsto e^{i\phi_1} z_1,$$

$$z_2 \mapsto e^{i\phi_2} z_{\sigma(2)},$$

where  $\phi_1, \phi_2 \in \mathbb{R}$ , and  $\sigma$  is a permutation of the set  $\{2, 3\}$ .

The preceding description of  $\operatorname{Aut}(\Omega)$  implies that  $\dim \operatorname{Aut}(\Omega) = 4$ . That is to say, each of the four connected components of  $\operatorname{Aut}(\Omega)$  is parametrized by the point a from the unit disc and by the rotation parameters  $\phi_1, \phi_2$ .

Suppose now that  $\Omega$  is biholomorphically equivalent to a Reinhardt domain  $D \subset \mathbb{C}^3$ . Since  $\Omega$  is bounded, it follows that D is hyperbolic. It follows from [Kru] that any hyperbolic Reinhardt domain  $G \subset \mathbb{C}^n$  can be biholomorphically mapped onto its normilized form  $\tilde{G}$  for which the identity component  $\operatorname{Aut}_0(\tilde{G})$  of  $\operatorname{Aut}(\tilde{G})$  is described as follows. There exist integers  $0 \le s \le t \le p \le n$  and  $n_i \ge 1, i = 1, \ldots, p$ , with  $\sum_{i=1}^p n_i = n$ , and real numbers  $\alpha_i^k$ ,  $i = 1, \ldots, s$ ,  $k = t+1, \ldots, p$ , and  $\beta_j^k$ ,  $j = s+1, \ldots, t$ ,  $k = t+1, \ldots, p$  such that if we set  $z^i = (z_{n_1+\cdots+n_{i-1}+1}, \ldots, z_{n_1+\cdots+n_i})$ ,  $i = 1, \ldots, p$ , then  $\operatorname{Aut}_0(\tilde{G})$  is given by the mappings

$$z^{i} \mapsto \frac{A^{i}z^{i} + b^{i}}{c^{i}z^{i} + d^{i}}, \quad i = 1, \dots, s,$$

$$z^{j} \mapsto B^{j}z^{j} + e^{j}, \quad j = s + 1, \dots, t,$$

$$z^{k} \mapsto C^{k} \frac{\prod_{j=s+1}^{t} \exp\left(-\beta_{j}^{k} \left(2\overline{e^{j}}^{T} B^{j}z^{j} + |e^{j}|^{2}\right)\right) z^{k}}{\prod_{i=1}^{s} (c^{i}z^{i} + d^{i})^{2\alpha_{i}^{k}}}, \quad k = t + 1, \dots, p,$$

$$(2)$$

where

$$\begin{pmatrix} A^i & b^i \\ c^i & d^i \end{pmatrix} \in SU(n_i, 1), \quad i = 1, \dots, s,$$

$$B^j \in U(n_j), \quad e^j \in \mathbb{C}^{n_j}, \quad j = s + 1, \dots, t,$$

$$C^k \in U(n_k), \quad k = t + 1, \dots, p.$$

The normalized form  $\tilde{G}$  is written as

$$G = \left\{ \left| z^{1} \right| < 1, \dots, \left| z^{s} \right| < 1, \right.$$

$$\left( \frac{z^{t+1}}{\prod_{i=1}^{s} \left( 1 - \left| z^{i} \right|^{2} \right)^{\alpha_{i}^{t+1}} \prod_{j=s+1}^{t} \exp\left( -\beta_{j}^{t+1} \left| z^{j} \right|^{2} \right)}, \dots, \right.$$

$$\left. \frac{z^{p}}{\prod_{i=1}^{s} \left( 1 - \left| z^{i} \right|^{2} \right)^{\alpha_{i}^{p}} \prod_{j=s+1}^{t} \exp\left( -\beta_{j}^{p} \left| z^{j} \right|^{2} \right)} \right) \in \tilde{G}_{1} \right\},$$
(3)

where  $\tilde{G}_1 := \tilde{G} \cap \{z^i = 0, i = 1, ..., t\}$  is a hyperbolic Reinhardt domain in  $\mathbb{C}^{n_{t+1}} \times \cdots \times \mathbb{C}^{n_p}$ .

It is now easy to see that, for any hyperbolic Reinhardt domain  $D \subset \mathbb{C}^3$  written in a normilized form  $\tilde{D}$ ,  $\operatorname{Aut}_0(\tilde{D})$  given by formulas (2) cannot have dimension equal to 4.

This completes the proof.

**Remark.** The theorem can be easily extended to  $\mathbb{C}^n$  for any  $n \geq 3$  (just replace

There is considerable evidence that, in complex dimension two, an example such as that constructed in Theorem 1 does not exist. Certainly the example provided above depends on the decoupling, in the domain  $\Omega$ , of the variables  $z_2, z_3$  from the variable  $z_1$ . Such decoupling is not possible when the dimension is only two.

The work of Bedford and Pinchuk (see [BP] and references therein) suggests that the only smoothly bounded domains in  $\mathbb{C}^2$  with non-compact automorphism groups are (up to biholomorphic equivalence) the complex ellipsoids

$$\Omega_{\alpha} = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1|^2 + |z_2|^{2\alpha} < 1\},\$$

where  $\alpha$  is a positive integer. It is also a plausible conjecture that any bounded domain in  $\mathbb{C}^2$  with non-compact automorphism group and a boundary of finite smoothness  $C^k$  for  $k \geq 1$ , is biholomorphically equivalent to some  $\Omega_{\alpha}$ , where  $\alpha \geq 1$  and is not necessarily an integer. Of course all the domains  $\Omega_{\alpha}$  are pseudoconvex and Reinhard.

However, as the following theorem shows, if we allow the boundary to be non-smooth at just one point, then the domain may be non-pseudoconvex and be non-equivalent to any Reinhardt domain.

**Theorem 2.** There exists a bounded, non-pseudoconvex domain  $\Omega \subset \mathbb{C}^2$  with non-compact automorphism group such that  $\partial\Omega$  is smooth real-analytic everywhere except one point (this exceptional point is an orbit accumulation point for the automorphism group action), and such that  $\Omega$  is not biholomorphically equivalent to any Reinhardt domain.

For the proof of Theorem 2, we first need the following lemma.

**Lemma A.** If  $\Omega \subset \mathbb{C}^2$  is a bounded, non-pseudoconvex, simply-connected domain such that the identity component  $Aut_0(\Omega)$  of the automorphism group  $Aut(\Omega)$  is non-compact, then  $\Omega$  is not biholomorphically equivalent to any Reinhardt domain.

Proof of Lemma A. Suppose that  $\Omega$  is biholomorphically equivalent to a Reinhardt domain D. Since  $\Omega$  is bounded, it follows that D is hyperbolic. Also, since  $\operatorname{Aut}_0(\Omega)$  is non-compact, then so is  $\operatorname{Aut}_0(D)$ . We are now going to show that any such domain D is either pseudoconvex, or not simply-connected, or cannot be biholomorphically equivalent to a bounded domain. This result clearly implies the lemma.

We can now assume that the domain D is written in its normalized form  $\tilde{D}$  as in (3), and  $\operatorname{Aut}_0(\tilde{D})$  is given by formulas (2). Then, since  $\operatorname{Aut}_0(\tilde{D})$  is non-compact, it must be that t>0. Next, if p=t, then  $\tilde{D}$  is either non-hyperbolic (for s< t), or (for s=t) is the unit ball or the unit polydisc and therefore is pseudoconvex. Thus we can assume that  $t=1, p=2, n_1=n_2=1$ .

Let  $\tilde{D}_1 \subset \mathbb{C}$  be the hyperbolic Reinhardt domain analogous to  $\tilde{G}_1$  that was defined above (see (3)). Clearly, there are the following possibilities for  $\tilde{D}_1$ :

- (i)  $\tilde{D}_1 = \{0 < |z_2| < R\}, \ 0 < R < \infty;$
- (ii)  $\tilde{D}_1 = \{r < |z_2| < R\}, \ 0 < r < R \le \infty;$
- (iii)  $\tilde{D}_1 = \{|z_2| < R\}, \ 0 < R < \infty.$

For the cases (i), (ii), D is always not simply-connected, and therefore we will concentrate on the case (iii). If s = 0, then  $\tilde{D}$  is not hyperbolic since it contains the complex line  $\{x_1, \dots, 0\}$ . Thus we can assume that  $x_1 = 1$ . Next charge that for

 $\alpha_1^2 \geq 0$ , the domain  $\tilde{D}$  is always pseudoconvex. Thus we may take  $\alpha_1^2 < 0$ . Then the domain  $\tilde{D}$  has the form

$$\tilde{D} = \left\{ |z_1| < 1, \, |z_2| < \frac{R}{(1 - |z_1|^2)^{\gamma}} \right\}, \qquad \gamma > 0.$$

We will now show that the above domain  $\tilde{D}$  cannot be biholomorphically equivalent to a bounded domain. More precisely, we will show that any bounded holomorphic function on  $\tilde{D}$  is independent of  $z_2$ .

Let  $f(z_1, z_2)$  be holomorphic on  $\tilde{D}$  and |f| < M for some M > 0. For every  $\rho$  such that  $|\rho| \le \frac{R}{2}$ , the disc  $\Delta_{\rho} = \{|z_1| < 1, z_2 = \rho\}$  is contained in  $\tilde{D}$ . We will show that  $\partial f/\partial z_2 \equiv 0$  on every such  $\Delta_{\rho}$ , which implies that  $\partial f/\partial z_2 \equiv 0$  everywhere in  $\tilde{D}$ .

Fix a point  $(\mu, \rho) \in \Delta_{\rho}$  and restrict f to the disc  $\Delta'_{\mu} = \{z_1 = \mu, |z_2| < R_{\mu}\}$ , where  $R_{\mu} = R/2(1 - |\mu|^2)^{\gamma}$ . Clearly,  $(\mu, \rho) \in \Delta'_{\mu}$  and  $\overline{\Delta'_{\mu}} \subset \tilde{D}$ . By the Cauchy Integral Formula

$$f(\mu, z_2) = \frac{1}{2\pi i} \int_{\partial \Delta'_{\mu}} \frac{f(\mu, \zeta)}{\zeta - z_2} d\zeta,$$

for  $|z_2| < R_{\mu}$ , and therefore

$$\frac{\partial f}{\partial z_2}(\mu, \rho) = \frac{1}{2\pi i} \int_{\partial \Delta'} \frac{f(\mu, \zeta)}{(\zeta - \rho)^2} d\zeta.$$

Hence

$$\left| \frac{\partial f}{\partial z_2}(\mu, \rho) \right| \le \frac{MR_{\mu}}{(R_{\mu} - |\rho|)^2}.$$

Letting  $|\mu| \to 1$  and taking into account that  $R_{\mu} \to \infty$ , we see that  $|\partial f/\partial z_2(\mu, \rho)| \to 0$  as  $|\mu| \to 1$ . Therefore,  $\partial f/\partial z_2 \equiv 0$  on  $\Delta_{\rho}$ .

The lemma is proved.

*Proof of Theorem 2.* We will now present a domain that satisfies the conditions of the lemma. Set

$$\Omega = \left\{ |z_1|^2 + |z_2|^4 + 8|z_1 - 1|^2 \left( \frac{z_2^2}{z_1 - 1} - \frac{3}{2} \frac{|z_2|^2}{|z_1 - 1|} + \frac{\overline{z_2}^2}{\overline{z_1} - 1} \right)^2 < 1 \right\}.$$

The domain  $\Omega$  is plainly bounded since the third term on the left is non-negative. Next, the identity component  $\operatorname{Aut}_0(\Omega)$  of its automorphism group is non-compact since it contains the subgroup

$$z_1 \mapsto \frac{z_1 - a}{1 - az_1},$$

$$z_2 \mapsto \frac{(1 - a^2)^{\frac{1}{4}} z_2}{(1 - az_1)^{\frac{1}{2}}},$$

Further,  $\Omega$  is simply-connected, since the family of mappings  $F_{\tau}(z_1, z_2) = (z_1, \tau z_2)$ ,  $0 \le \tau \le 1$ , retracts  $\Omega$  inside itself, as  $\tau \to 0$ , to the unit disc  $\{|z_1| < 1, z_2 = 0\}$  (which is simply-connected).

To show that  $\Omega$  is not pseudoconvex, consider its unbounded realization. Namely, under the mapping

$$z_1 \mapsto \frac{z_1 + 1}{z_1 - 1},$$

$$z_2 \mapsto \frac{\sqrt{2}z_2}{\sqrt{z_1 - 1}},$$

$$(4)$$

the domain  $\Omega$  is transformed into the domain

$$\Omega' = \left\{ \operatorname{Re} z_1 + \frac{1}{4} |z_2|^4 + 2\left(z_2^2 - \frac{3}{2} |z_2|^2 + \overline{z_2}^2\right)^2 < 0 \right\}.$$

It is easy to see that at the boundary point  $(-\frac{3}{4},1) \in \partial\Omega'$  the Levi form of  $\partial\Omega'$  is equal to  $-|z_2|^2$ , and thus is negative-definite. Therefore,  $\Omega$  is non-pseudoconvex.

Hence, by Lemma A,  $\Omega$  is not biholomorphically equivalent to any Reinhardt domain.

Next, if  $\phi$  denotes the defining function of  $\Omega$ , the following holds at every boundary point of  $\Omega$  except (1,0):

$$\frac{\partial \phi}{\partial z_1} = \frac{1}{z_1 - 1} \left( -\frac{z_2}{2} \frac{\partial \phi}{\partial z_2} + 1 - \overline{z_1} \right),\,$$

and therefore grad  $\phi$  does not vanish at every such point. Hence,  $\partial\Omega$  is smooth real-analytic everywhere except at (1,0).

The theorem is proved.

## Remarks.

- 1. The hypothesis of simple connectivity in Lemma A is automatically satisfied if, for example, the boundary of the domain is locally variety-free and smooth near some orbit accumulation point for the automorphism group of the domain (see e.g. [GK2]). For a smoothly bounded domain it would follow from a conjecture of Greene/Krantz [GK3].
- 2. Tedious calculations show that the boundary of the domain  $\Omega$  in Theorem 2 is quite pathological near the exceptional point (1,0). It is not Lipschitz-smooth of any positive degree. It would be interesting to know whether there is an example with Lipschitz-1 boundary at the bad point.

In fact, many more examples similar to that in Theorem 2 can be constructed in the following way. Let

$$\Omega' = \{ (z_1, z_2) \in \mathbb{C}^2 : \operatorname{Re} z_1 + P(z_2) < 0 \}, \tag{5}$$

where  $P = |z_2|^{2m} + Q(z_2)$  is a homogeneous non-plurisubharmonic polynomial, m is a positive integer, and  $Q(z_2)$  is positive away from the origin. Then, by a mapping analogous to (4),  $\Omega'$  can be transformed into a bounded domain  $\Omega$ . The domain  $\Omega$  is given by connected, non-paper by approximately non-paper by  $\Omega$ .

real-analytic everywhere except at the point (1,0). For all such examples,  $\partial\Omega$  is not Lipschitz-smooth of any positive degree at (1,0).

It is also worth noting that, in the example contained in Theorem 2, the point (-1,0) is also an orbit accumulation point, but  $\partial\Omega$  is smooth real-analytic at this point.

3. It is conceivable that the domain  $\Omega$  as in Theorem 2 has an alternative, smoothly bounded realization, but it looks plausible that if in formula (5) we allow  $P(z_2)$  to be an arbitrary homogeneous polynomial positive away from the origin with no harmonic term, then domain (5) does not have a bounded realization with  $C^1$ -smooth boundary, unless  $P(z_2) = c|z_2|^{2m}$ , where c > 0 and m is a positive integer.

**Acknowledgements.** This work was completed while the second author was an Alexander von Humboldt Fellow at the Research at MSRI by the third author supported in part by NSF Grant DMS-9022140. University of Wuppertal.

## References

- [BL] Bell, S., Ligocka, E., A simplification and extension of Fefferman's theorem on biholomorphic mappings, Invent. Math. 57 (1980), 283–289.
- [BP] Bedford, E., Pinchuk, S., Domains in  $\mathbb{C}^{n+1}$  with non-compact automorphism groups, J. Geom. Anal. 1 (1991), 165–191.
- [C] Cartan, H., Les transformations analytiques des domaines cerclés les uns dans les autres, Compt. Rendus de l'Académie des Sciences de Paris 190 (1930), 718–720.
- [GK1] Greene, R. E., Krantz, S. G., Characterization of certain weakly pseudoconvex domains with non-compact automorphism groups, Lecture Notes in Mathematics 1268, Springer-Verlag, 1987, 121–157.
- [GK2] Greene, R. E, Krantz, S. G., Invariants of Bergman geometry and the automorphism groups of domains in  $\mathbb{C}^n$ , Proceedings of a Conference on Complex Analysis and Geometry held in Cetraro, 1989, Mediterranean Press, 1992, 107–136.
- [GK3] Greene, R. E., Krantz, S. G., Techniques for studying automorphisms of weakly pseudoconvex domains, in J. E. Fornaess (Ed.) Several Complex Variables: Proceedings of the Mittag-Leffler Institute, 1987-1988, Math. Notes, Vol. 38, Princeton University Press, 1993, 389-410.
- [Kob] Kobayashi, S., Intrinsic distances, measures and geometric function theory, Bull. Amer. Math. Soc. 82 (1976), 357–416.
- [Kod] Kodama, A., A characterization of certain domains with good boundary points in the sense of Greene-Krantz, Kodai. Math. J. 12 (1989), 257–269.
- [Kra] Krantz, S. G., Convexity in complex analysis, Several Complex Variables and Complex Geometry, Part 1 (Santa Cruz, CA, 1989), Proc. Symp. Pure Math. **52**, Part 1 (1991), 119–137.
- [Kru] Kruzhilin, N. G., Holomorphic automorphisms of hyperbolic Reinhardt domains (translated from Russian), Math. USSR-Izv. 32 (1989), 15–38.
- [R] Rosay, J. P., Sur une caractérisation de la boule parmi les domaines de  $\mathbb{C}^n$  par son groupe d'automorphismes, Ann. Inst. Fourier (Grenoble) **29** (1979), 91–97.
- [W] Wong, B., Characterization of the unit ball in  $\mathbb{C}^n$  by its automorphism group, Invent.

Siqi Fu

DEPARTMENT OF MATHEMATICS

University of California, Irvine, CA 92717

USA

E-MAIL ADDRESS: SFU@MATH.UCI.EDU

A. V. Isaev

CENTRE FOR MATHEMATICS AND ITS APPLICATIONS

THE AUSTRALIAN NATIONAL UNIVERSITY

CANBERRA, ACT 0200

AUSTRALIA

E-MAIL ADDRESS: ALEXANDER.ISAEV@ANU.EDU.AU

AND

BERGISCHE UNIVERSITÄT

GESAMTHOCHSCHULE WUPPERTAL

Матнематік (FB 07)

Gaussstrasse 20

42097 Wuppertal

GERMANY

E-MAIL ADDRESS: ALEXANDER.ISAEV@MATH.UNI-WUPPERTAL.DE

## S. G. Krantz

DEPARTMENT OF MATHEMATICS

Washington University, St.Louis, MO 63130

TISΔ

E-mail address: sk@math.wustl.edu

AND

MSRI

1000 CENTENNIAL DRIVE

Berkeley, California 94720

USA

E-MAIL ADDRESS: KRANTZ@MSRI.ORG