VANISHING SUMS OF *m* TH ROOTS OF UNITY IN FINITE FIELDS

T. Y. LAM AND K. H. LEUNG

ABSTRACT. In an earlier work, the authors have determined all possible weights n for which there exists a vanishing sum $\zeta_1 + \cdots + \zeta_n = 0$ of m th roots of unity ζ_i in characteristic 0. In this paper, the same problem is studied in finite fields of characteristic p. For given m and p, results are obtained on integers n_0 such that all integers $n \ge n_0$ are in the "weight set" $W_p(m)$. The main result (1.3) in this paper guarantees, under suitable conditions, the existence of solutions of $x_1^d + \cdots + x_n^d = 0$ with all coordinates not equal to zero over a finite field.

1. INTRODUCTION

By a vanishing sum of m th roots of unity, we mean an equation $\alpha_1 + \cdots + \alpha_n = 0$ where $\alpha_i^m = 1$ for each i. The integer n is said to be the *weight* of this vanishing sum. In [LL], considering m th roots of unity in \mathbb{C} , we defined W(m) to be the set of integers $n \ge 0$ for which there exists a vanishing sum $\alpha_1 + \cdots + \alpha_n = 0$ as above. The principal result in [LL] gives a complete determination of the weight set W(m)(in characteristic 0), as follows.

Theorem 1.1. For any natural number m with prime factorization $p_1^{a_1} \cdots p_r^{a_r}$, the weight set W(m) is exactly given by $\mathbb{N} p_1 + \cdots + \mathbb{N} p_r$. (Here and in the following, $\mathbb{N} := \{0, 1, 2, \cdots\}$).

In this paper, we study vanishing sums of m th roots of unity in characteristic p. In analogy to the characteristic 0 case, we define $W_p(m)$ to be the set of weights n for which there exists a vanishing sum $\alpha_1 + \cdots + \alpha_n = 0$ where each α_i is an m th root of unity in $\overline{\mathbb{F}}_p$, the algebraic closure of the prime field \mathbb{F}_p . Note that, if $m = p^t m'$ where gcd(p, m') = 1, we have $x^m = 1$ in $\overline{\mathbb{F}}_p$ iff $x^{m'} = 1$; in particular, $W_p(m) = W_p(m')$. Therefore, we may assume throughout that gcd(p, m) = 1, i.e. p is not among the prime divisors p_i of m. As in the case of characteristic 0, we have

Lam was supported in part by NSF. Research at MSRI is supported in part by NSF grant DMS-9022140.

Leung's research was carried out while he was on sabbatical leave at U.C. Berkeley from the National University of Singapore. The hospitality of the former institution is gratefully acknowledged.

 $p_i \in W_p(m)$ for all *i*. But in characteristic *p*, we also have $p \in W_p(m)$ (due to the vanishing sum $p \cdot 1 = 0$), so now

(1.2)
$$W_p(m) \supseteq \mathbb{N} \, p + \mathbb{N} \, p_1 + \dots + \mathbb{N} \, p_r.$$

Easy examples (see (2.1)) show that this need not be an equality in general, so we are left with no viable conjecture on the structure of the weight set $W_p(m)$ in characteristic p. However, (1.2) does show that, if m > 1, all sufficiently large integers n (in fact all $n \ge (p-1)(p_i-1)$) belong to $W_p(m)$. A more tractable problem will then be the determination of more accurate bounds n_0 such that all integers $n \ge n_0$ belong to $W_p(m)$.

In this paper, we will show how such an integer n_0 can be determined. Our work is divided into three cases, depending on whether gcd(p-1, m) is 1, 2, or bigger. The estimates on n_0 differ from case to case, and are given respectively in (5.6), (4.1), and (3.1)-(3.3). Although we have three different estimates on n_0 , there does exist a (necessarily weaker) uniform estimate for all cases. In the following, we shall try to explain what this uniform estimate is, and why is it a reasonable one.

A guiding principle for our work throughout is the fact that a finite field is a C_1 -field (see [Gr]). If $K = \mathbb{F}_{p^k}$ is a finite field containing all m th roots of unity, then, for $d := (p^k - 1)/m$, the m th roots of unity in $\overline{\mathbb{F}}_p$ comprise the group \dot{K}^d . Therefore, a vanishing sum of m th roots of unity of weight n corresponds precisely to a "good" solution of $x_1^d + \cdots + x_n^d = 0$ in K, where by a "good" solution we mean one with each $x_i \neq 0$. If n > d, the fact that K is C_1 implies that we have a solution $(x_1, \cdots, x_n) \neq (0, \cdots, 0)$. It certainly seems tempting to speculate that there exists in fact a "good" solution (in K). If this is indeed the case, then by what we said earlier in this paragraph, any integer n > d will be in the weight set $W_p(m)$.

The desired conclusion that, for n > d, $x_1^d + \cdots + x_n^d = 0$ has a "good" solution in K is, however, not true in general! For instance, if $d = p^k - 1$, then $x^d = 1$ for each $x \in \dot{K}$, so we have a "good" solution for $x_1^d + \cdots + x_n^d = 0$ in K only when n is a multiple of p. In a similar vein, if k = 1, p is odd, and d = (p-1)/2, then any nonzero d th power in K is ± 1 . For any odd integer $n \in (d, p)$, the equation $x_1^d + \cdots + x_n^d = 0$ again has no "good" solution in \mathbb{F}_p . The trouble with these cases is that $m \leq 2$, for which we don't have "enough" m th roots to play with. As it turns out, as soon as we ignore the above cases, we'll have the following uniform result for getting "good" solutions.

Theorem 1.3. Let $K = \mathbb{F}_{p^k}$ and $d = (p^k - 1)/m$ as above, and assume that $m \neq 1$, $(m, k) \neq (2, 1)$. Then, whenever n > d, the equation $x_1^d + \cdots + x_n^d = 0$ has a "good" solution in K. In other words, the weight set $W_p(m)$ contains all integers $\geq d + 1$.

The results in §§3-5 below will cover this theorem in the case $m \ge 3$. In the case m = 2, (1.3) is quickly checked as follows. Since we assume in this case that $k \ge 2$,

we have $d \ge (p^2 - 1)/2 \ge p - 1$. Given $n \ge d + 1 \ge p$, it is easy to solve the equation $\alpha_1 + \cdots + \alpha_n = 0$ with $\alpha_i = \pm 1$, by considering the parity of n. Having disposed of the trivial cases m = 1, 2, we may assume in §§3-6 of this paper that $m \ge 3$.

In the case when p is odd and d = 2, (1.3) says precisely that, for any n > 2, the quadratic form $x_1^2 + \cdots + x_n^2$ has a "good" zero over any finite field of more than five elements. This is a special case of a well-known observation of Witt for isotropic diagonal quadratic forms (see [Wi: p.39], or [BS: p.394], [La: p.25, Ex.7]). Thus, (1.3) may be thought of as a generalization of Witt's result to the higher degree diagonal forms $x_1^d + \cdots + x_n^d$ over finite fields. Note that $d \mid (p^k - 1)$ is not a really essential assumption in (1.3). In dealing with the equation $x_1^e + \cdots + x_n^e = 0$, we can replace the degree e by $d := \gcd(p^k - 1, e)$, and define m to be $(p^k - 1)/d$. Then $\dot{K}^e = \dot{K}^d$, so under the assumptions of (1.3), $x_1^e + \cdots + x_n^e = 0$ will have a "good" solution as long as n > d.

In the literature, there are many results dealing with diagonal equations over finite fields; see, for instance, [LN], [Sch], [Sm], and more recently [QY]. Conventionally, one could apply algebro-geometric methods, or alternatively the method of Gauss and Jacobi sums. As the referee of this paper pointed out, these methods can be utilized to show the existence of "good" solutions to a diagonal equation $x_1^d + \cdots + x_n^d =$ $0 \ (n > 2)$ in \mathbb{F}_q if q is suitably large compared to d (without the condition n > d). However, these conventional methods do not seem to give enough information if qis "small" in comparison to d. In our setting, working mostly with n > d and taking full advantage of the additive nature of the special equation $x_1^d + \cdots + x_n^d = 0$, we apply instead the methods of additive number theory. These methods do give fairly precise results, without reference to the size of \mathbb{F}_q . In fact, the analysis in §§3-5 will not only prove (1.3), but also show that, in various cases, the equation $x_1^d + \cdots + x_n^d = 0$ has a "good" solution in K often for much smaller values of n(than $n \ge d + 1$). Thus, the more precise results in this paper are to be found in (3.1)-(3.3), (4.1) and (5.6). Theorem 1.3 is only a common denominator of these results giving a convenient and uniform summary of the main work in this paper.

Acknowledgment. We thank the referee of this paper, whose comments enabled us to rewrite more accurately the last paragraph above comparing the use of different methods in treating diagonal equations over finite fields.

2. Some Basic Examples

We shall begin with some examples and computations of the weight sets $W_p(m)$. The first couple of examples show that various properties of weight sets in characterictic 0 are no longer valid in characteristic p. For convenience of expressing weight sets, let us use the notation $[n, \infty)_{\mathbb{Z}}$ for the set of integers $\geq n$. **Example 2.1.** Referring to (1.2), the smallest positive element in the set $W_p(m)$ may not be min $\{p, p_1, \dots, p_r\}$. For instance, when p = 11 and m = 5, the 5th roots of unity in $\overline{\mathbb{F}}_{11}$ are $\{1, 3, 9, 5, 4\}$. Observing that 1 + 1 + 9 = 0 in \mathbb{F}_{11} , we see that $W_{11}(5)$ contains 3, which is smaller than 5 and 11. By (2.3) below, we have $W_{11}(5) = \{0\} \cup [3, \infty)_{\mathbb{Z}}$. Thus, not only (1.2) fails to be an equality, but also $W_{11}(5)$ is not even of the form $\sum_i \mathbb{N} q_i$ for a set of primes q_i 's.

Example 2.2. Contrary to the characteristic 0 case, the set $W_p(m)$ may be larger than $W_p(m_0)$ where m_0 is the square-free part of m. For instance, let p = 5 and m = 4, so $m_0 = 2$. It is easy to see that $W_5(2) = \{0, 2\} \cup [4, \infty)_{\mathbb{Z}}$, but $W_5(4) = \{0\} \cup [2, \infty)_{\mathbb{Z}}$.

Example 2.3. Let $q = p^a > 5$ where p is an odd prime, and let m = (q-1)/2. Then d := (q-1)/m = 2. For any $n \ge 3$, the quadratic form $X_1^2 + \cdots + X_n^2$ is isotropic over \mathbb{F}_q , so by the theorem of Witt referenced before, it has a "good" zero in \mathbb{F}_q . Therefore, $n \in W_p(m)$. It follows that $W_p(m) = \{0\} \cup [2, \infty)_{\mathbb{Z}}$ if $q \equiv 1 \pmod{4}$, and $W_p(m) = \{0\} \cup [3, \infty)_{\mathbb{Z}}$ if $q \equiv 3 \pmod{4}$.

Example 2.4. (\mathbb{F}_p contains all *m* th roots of unity.) Let p = 31, and m = 3. The third roots of unity are $\{1, 5, 25\}$, so the equation $25 + 6 \cdot 1 = 0 \in \mathbb{F}_{31}$ shows that $7 \in W_{31}(3)$. A routine computation shows that $W_{31}(3) = \{0, 3, 6, 7\} \cup [9, \infty)_{\mathbb{Z}}$.

Example 2.5. (\mathbb{F}_p contains no *m* th roots of unity other than 1.) Let p = 2, and m = 73. We work in $K = \mathbb{F}_{2^9}$ which contains all 73 rd roots of unity. By standard tables of irreducible polynomials over finite fields, the trinomial $f(X) = X^9 + X + 1$ is irreducible over \mathbb{F}_2 , so we can take *K* to be $\mathbb{F}_2[X]/(f(X))$. Let $\alpha := \overline{X} \in K$. We have $0 = (\alpha^9 + \alpha + 1)^8 = \alpha^{72} + \alpha^8 + 1$, so $\alpha^{73} = \alpha(\alpha^8 + 1) = \alpha^9 + \alpha = 1$. Thus, the relation $\alpha^9 + \alpha + 1 = 0$ shows that $3 \in W_2(73)$, and it follows easily that $W_2(73) = \{0\} \cup [2, \infty)_{\mathbb{Z}}$.

In the balance of this section, let us consider $W_p(m)$ in the case when m is a prime power (not divisible by p). Under a special hypothesis on the cyclotomic polynomial $\Phi_m(X)$, the weight set $W_p(m)$ can be determined explicitly.

Theorem 2.6. Let $m = \ell^a$ where ℓ is a prime different from p, and assume that the cyclotomic polynomial $\Phi_m(X) \in \mathbb{Z}[X]$ remains irreducible modulo p. Then $W_p(m) = \mathbb{N} p + \mathbb{N} \ell$.

Proof. Of course, it suffices to prove the inclusion " \subseteq ". Let ζ be a primitive m th root of unity in $\overline{\mathbb{F}}_p$. Let $m' := m/\ell$, and $\alpha := \zeta^{m'}$ (a primitive ℓ th root of unity). Let $K = \mathbb{F}_p(\zeta)$, and $L = \mathbb{F}_p(\alpha)$. Since $\Phi_m(X)$ is irreducible mod p, $[K : \mathbb{F}_p] = \varphi(m) = m'(\ell - 1)$. From this, it is easy to see that [K : L] = m' and $[L : \mathbb{F}_p] = \ell - 1$.

Any vanishing sum of m th roots of unity can be written in the form $\sum_{i=0}^{m'-1} g_i \zeta^i = 0$, where each g_i is a sum of ℓ th roots of unity. Since the degree of ζ over L

is m', the elements $1, \zeta, \dots, \zeta^{m'-1}$ are linearly independent over L. Therefore, each $g_i \in L$ is itself a vanishing sum, and it suffices to show that its weight is in $\mathbb{N} p + \mathbb{N} \ell$. Starting over again, we are now down to considering a vanishing sum $\sum_{i=0}^{\ell-1} a_i \alpha^i = 0$, where each $a_i \in \mathbb{N}$. Let a_j be the smallest among the a_i 's. Since the minimal equation of α over \mathbb{F}_p is $1 + \alpha + \dots + \alpha^{\ell-1} = 0$, it follows easily that $a_0 = a_1 = \dots = a_{\ell-1} \in \mathbb{F}_p$. The weight of the vanishing sum in question is $\sum_i a_i \equiv \ell a_j \pmod{p}$. Since $\sum_i a_i \geq \ell a_j$, it follows that $\sum_i a_i = \ell a_j + bp$ for some $b \in \mathbb{N}$, as desired.

Remark 2.7. A vanishing sum of m th roots of unity is said to be minimal if no proper subsum of it is also vanishing. In general, the problem of determining the minimal vanishing sums is difficult (both in characteristic 0 and in characteristic p). Under the hypothesis of (2.6), however, this problem can be solved. In fact, the argument presented in the proof above can be used to show that, in the setting of (2.6), the minimal vanishing sums of m th roots of unity are, up to multiplication by a power of ζ : (1) $p \cdot 1 = 0$, and (2) $1 + \alpha + \cdots + \alpha^{\ell-1} = 0$. (Of course, this implies that $W_p(m) = \mathbb{N} p + \mathbb{N} \ell$.) For this conclusion, however, the assumption on the irreducibility of $\Phi_m(X)$ modulo p is essential, as the examples (2.1), (2.4) and (2.5) show. (In (2.1) and (2.4), $\Phi_m(X)$ splits completely modulo p, and in (2.5), $\Phi_m(X)$ splits into the product of eight irreducible factors of degree 9 in $\mathbb{F}_p[X]$.)

3. The Case $\operatorname{GCD}(p-1, m) \geq 3$

In dealing with $W_p(m)$, our main goal is to find good estimates for integers n_0 such that $[n_0, \infty)_{\mathbb{Z}} \subseteq W_p(m)$. We begin our analysis with the case when $gcd(p-1, m) \geq 3$. This case turns out to be fairly easy if we use the right tools from additive number theory modulo p. It will be convenient to use the following notations. For a subset A in a field, we shall write |A| for the cardinality of A, and for any integer $n \geq 1$, we write n * A for the set $A + \cdots + A$ with n summands of A.

Theorem 3.1. Assume that $m_0 := \gcd(p-1, m) \ge 3$ and let $d_0 = (p-1)/m_0$. Then $[d_0 + 1, \infty)_{\mathbb{Z}} \subseteq W_p(m_0) \subseteq W_p(m)$.

Proof. Since $m_0|(p-1)$, the group H of m_0 th roots of unity in \mathbb{F}_p is exactly $\dot{\mathbb{F}}_p^{d_0}$ and has exactly m_0 elements. We claim that $|n * H| \ge nm_0$ for $n \le d_0$, and |n * H| = p for $n \ge d_0 + 1$. It suffices to prove this for $n = 1, 2, \dots, d_0 + 1$ (for, once we show that $|(d_0 + 1) * H| = p$, then $(d_0 + 1) * H = \mathbb{F}_p$, and this implies that $(d_0 + i) * H = \mathbb{F}_p$ for any $i \ge 1$). We proceed by induction on n, the case n = 1 being clear. Assume that $|n * H| \ge nm_0$ where $n < d_0$. By the Cauchy-Davenport Theorem (see [Ma: Cor.1.2.3]), |(n + 1) * H| is either p (and hence $\ge (n + 1)m_0$), or else

$$|(n+1) * H| \ge |n * H| + |H| - 1 \ge (n+1)m_0 - 1.$$

In the latter case, $|(n+1)*H \setminus \{0\}| \ge (n+1)m_0-2$. Since H acts on $(n+1)*H \setminus \{0\}$ by multiplication, $|(n+1)*H \setminus \{0\}|$ is a multiple of m_0 . Since $m_0 \ge 3$, we must therefore have $|(n+1)*H \setminus \{0\}| \ge (n+1)m_0$, which gives what we want. This proves our claim for $n \le d_0$. In particular, $|d_0*H| \ge d_0m_0 = p-1$. By the Cauchy-Davenport Theorem again, $(d_0+1)*H$ must be \mathbb{F}_p , for otherwise we would have

$$|(d_0+1)*H| \ge |d_0*H| + |H| - 1 \ge d_0m_0 + m_0 - 1 = p + (m_0 - 1) > p_2$$

a contradiction. This completes our inductive proof. Thus, for any $n \ge d_0 + 1$, we have $0 \in n * H$. This means that $n \in W_p(m_0)$, and so $[d_0 + 1, \infty)_{\mathbb{Z}} \subseteq W_p(m_0) \subseteq W_p(m)$.

Example 3.2. In many cases Theorem 3.1 gives the best result. For instance, if $p \equiv 3 \pmod{4}$ and $m = (p-1)/2 \geq 3$, then $d_0 = 2$ and we have $d_0 \notin W_p(m)$ since m is odd. Even in the case $p \equiv 1 \pmod{4}$, the Theorem may still give the best result. For instance, if p = 13 and m = 4, then $d_0 = 3$ and $G = \{\pm 1, \pm 8\}$. By a simple calculation, $3 * G = \dot{\mathbb{F}}_{13}$, so again $d_0 = 3 \notin W_p(m)$. On the other hand, if m is divisible by two distinct primes p_1, p_2 , then the fact that $p_1, p_2 \in W_p(m)$ implies that $[n_0, \infty)_{\mathbb{Z}} \subseteq W_p(m)$ for $n_0 := (p_1 - 1)(p_2 - 1)$ (see [LeV: p.22, Ex.4]). In case the number d_0 in (3.1) is "large", $[n_0, \infty)_{\mathbb{Z}} \subseteq W_p(m)$ will of course give a better result.

We can now derive the first case of Theorem 1.3.

Corollary 3.3. Let $K = \mathbb{F}_{p^k}$ be a finite field containing all m th roots of unity, and let $d = (p^k - 1)/m$. If $m_0 := \gcd(p - 1, m) \ge 3$, then $[d+1, \infty)_{\mathbb{Z}} \subseteq W_p(m_0) \subseteq W_p(m)$.

Proof. Say $p-1 = m_0 d_0$ and $m = m_0 m_1$. Then

$$d = \frac{(p-1)(p^{k-1} + \dots + p + 1)}{m} = d_0 \cdot \frac{p^{k-1} + \dots + p + 1}{m_1}.$$

Since $gcd(d_0, m_1) = 1$, the fraction on the RHS above is an integer. Therefore, we have $d_0 \mid d$, and the desired conclusion follows from Theorem 3.1.

4. The Case GCD(p-1, m) = 2

We shall assume throughout this section that gcd(p-1, m) = 2 (and as before $m \geq 3$). In particular, p is odd and m is even. In this case, $W_p(m)$ contains $2\mathbb{N}$ and is stable under addition by 2. Thus, once we have an odd integer $n \in W_p(m)$, we will have automatically $[n-1, \infty)_{\mathbb{Z}} \subseteq W_p(m)$. This observation will be used without further mention in the following.

Let $K = \mathbb{F}_{p^k}$ be any finite field containing the group G of all m th roots of unity. The following result gives a somewhat sharper form of Theorem 1.3 in the

case gcd(p-1, m) = 2 (in that the index [K : G] itself is shown to be a weight, with a minor exception).

Theorem 4.1. Assume that gcd(p-1, m) = 2, and let $d = [K : G] = (p^k - 1)/m$. Then $[d, \infty)_{\mathbb{Z}} \subseteq W_p(m)$ unless p = 3 and $m = 3^k - 1$, in which case $[d+1, \infty)_{\mathbb{Z}} \subseteq W_3(m)$.

Proof. Let us first check the Theorem when m = 4. In this case, the assumption gcd(p-1, m) = 2 implies that G is not contained in \mathbb{F}_p , so $k \ge 2$. If p > 3, then $d \ge (p^2 - 1)/4 \ge p$, and we have $[d, \infty)_{\mathbb{Z}} \subseteq [p, \infty)_{\mathbb{Z}} \subseteq W_p(m)$. If p = 3, then $d \ge (9-1)/4 = 2$, and we have again $[d, \infty)_{\mathbb{Z}} \subseteq [2, \infty)_{\mathbb{Z}} \subseteq W_3(m)$ (since $W_3(m)$ contains both 2 and 3). In the following, we may therefore assume that $m \ge 6$.

Write m = 2m', so that

(4.2)
$$d = \frac{(p-1)(p^{k-1} + \dots + p+1)}{2m'} = \frac{p-1}{2} \cdot \frac{p^{k-1} + \dots + p+1}{m'}$$

Since gcd(m', (p-1)/2) = 1, we have $m'|(p^{k-1}+\cdots+p+1)$. If $m' < p^{k-1}+\cdots+p+1$, the second factor on the RHS in (4.2) is ≥ 2 , so $d \geq p-1$. Since $p \in W_p(m)$, we have $[d, \infty)_{\mathbb{Z}} \subseteq [p-1, \infty)_{\mathbb{Z}} \subseteq W_p(m)$, as desired. Therefore, in the following we may assume that

(4.3)
$$m' = p^{k-1} + \dots + p + 1$$
, and $d = (p-1)/2$.

In this case $\dot{K} = \dot{\mathbb{F}}_{p-1} \cdot G$, so any coset of G in \dot{K} has a "scalar" representative. We fix a generator ζ for the group G, and try to put a lower bound on the cardinality of the set $A := \mathbb{F}_p \cap 2 * G$.

Recalling that $m \ge 6$, write

$$(\zeta - 1)G = a_1G$$
, $(\zeta^2 - 1)G = a_2G$, and $(\zeta^4 - 1)G = a_3G$ where $a_i \in \dot{\mathbb{F}}_p$.

Clearly, $\pm a_i \in A$, since $-1 \in G$. First let us assume that these three G-cosets in \dot{K} are different. Since $a_i \neq -a_j$, $\{0, \pm a_i\}$ are seven different elements of A. (In particular, $p \geq 7$ here.) Applying repeatedly the Cauchy-Davenport Theorem in \mathbb{F}_p , we see that $|n * A| \geq \min\{p, 6n + 1\}$. It follows that

$$n \ge (p-1)/6 \Longrightarrow n * A = \mathbb{F}_p \Longrightarrow -1 \in n * A \Longrightarrow 2n+1 \in W_p(m).$$

This yields $2\lceil \frac{p-1}{6}\rceil + 1 \in W_p(m)$ (where $\lceil \cdot \rceil$ denotes the ceiling function). Writing p in the form $6t \pm 1$, we see easily that $2\lceil \frac{p-1}{6}\rceil = \lceil \frac{p-1}{3}\rceil$. Thus, in this case, we get the stronger conclusion that $\lfloor (p-1)/3, \infty \rangle_{\mathbb{Z}} \subseteq W_p(m)$.

From now on, we may assume that the three cosets $\{a_iG\}$ above are not all different. If $a_1G = a_2G$, then $\zeta^2 - 1 = (\zeta - 1)\zeta^i$ for some *i*, and so $\zeta^i = \zeta + 1$. Since *m* is even, this shows that $3 \in W_p(m)$, and so $[2, \infty)_{\mathbb{Z}} \subseteq W_p(m)$. We have certainly no problem in this case (except when d = 1, which occurs only when p = 3). If $a_2G = a_3G$, we can finish similarly. Now assume $a_1G = a_3G$. Here, $\zeta^4 - 1 = (\zeta - 1)\zeta^j$ for some j, so $\zeta^j = \zeta^3 + \zeta^2 + \zeta + 1$. As before, this gives $5 \in W_p(m)$. If p > 7, then $d = (p-1)/2 \ge 5$, and we have what we want. Thus we are only left with the cases p = 3, 5, 7.

If p = 3, we have d = (p-1)/2 = 1 (and $m = 3^k - 1$ by (4.3)). In this case the desired conclusion is $[2, \infty)_{\mathbb{Z}} \subseteq W_3(m)$, which is true since $2, 3 \in W_3(m)$.

If p = 5, then $d = 2 \in W_5(m)$. In this case we need to show that $3 \in W_5(m)$. If $a_1G = a_2G$, we are done as before. Otherwise, one of these cosets must be the identity coset G (since $[\dot{K}:G] = d = 2$), and this implies again that $3 \in W_5(m)$.

Finally, we treat the case p = 7. Here we must show that [K : G] = d = 3 is in the weight set $W_7(m)$. We first note that:

(4.4) If
$$(\zeta - 1)G = (\zeta^{i} - 1)G = (\zeta^{i+1} - 1)G$$
 for some $i \ge 1$, then $3 \in W_7(m)$.

Indeed, if we write $\zeta^i - 1 = (\zeta - 1)\zeta^r$ and $\zeta^{i+1} - 1 = (\zeta - 1)\zeta^s$, then $\zeta^r = \zeta^{i-1} + \cdots + \zeta + 1$ and $\zeta^s = \zeta^i + \cdots + \zeta + 1$ imply that $\zeta^s = \zeta^i + \zeta^r$, so $3 \in W_7(m)$. Now let C, C' be the two nonidentity cosets of G in K. By reasonings we have used before, we may assume that $(\zeta^2 - 1)G = C$ and $(\zeta - 1)G = (\zeta^4 - 1)G = C'$. Noting that 3 is prime to m (since $\gcd(p - 1, m) = \gcd(6, m) = 2$), we may also assume, in view of (4.4), that $(\zeta^3 - 1)G = C$. Replacing ζ by ζ^3 , we may further assume that $(\zeta^9 - 1)G = C'$. Next, note that since $m \ge 6$, it cannot divide 10, so $\zeta^{10} \ne 1$. Thus, in view of (4.4), we may assume that $(\zeta^5 - 1)G = C$, and hence that $(\zeta^{10} - 1)G = C'$. Now we have $C' = (\zeta - 1)G = (\zeta^9 - 1)G = (\zeta^{10} - 1)G$, so $3 \in W_7(m)$ once more by (4.4).

5. The CASE GCD(p-1, m) = 1

Throughout this section, we shall assume that gcd(p-1, m) = 1 (and as before, $m \ge 3$). The analysis of the weight set $W_p(m)$ in this case turns out to require the hardest work.

The assumption that gcd(p-1, m) = 1 means that the only m th root of unity in \mathbb{F}_p is 1. Therefore, upon factoring the polynomial $X^m - 1$ modulo p, we have

(5.1)
$$X^m - 1 = (X - 1)g_1(X)g_2(X)\cdots,$$

where the g_i 's are irreducible monic polynomials in $\mathbb{F}_p[X]$, each of degree ≥ 2 . Let $\ell := \min\{\deg(g_i)\}$. This integer ℓ will play an important role in finding the estimates on $W_p(m)$ in this section, so let us first note a few other characterizations of it.

Recall that the cyclotomic polynomial $\Phi_n(X) \in \mathbb{Z}[X]$ factors modulo p into a product of irreducible factors each of degree given by the order of the element p in the unit group $U(\mathbb{Z}/n\mathbb{Z})$ (see, e.g. [Gu]). Since $X^m - 1 = \prod_{n|m} \Phi_n(X)$, it follows that ℓ is the minimum of the orders of p in $U(\mathbb{Z}/n\mathbb{Z})$ for n ranging over the divisors of m greater than 1. From this, we see that ℓ is also the minimum of the orders of p in $U(\mathbb{Z}/q\mathbb{Z})$ for q ranging over the prime divisors of m. It is now an easy exercise to check the following:

(5.2)
$$\ell = \min \{ e \ge 1 : \gcd(p^e - 1, m) > 1 \}.$$

This simply means that $\mathbb{F}_{p^{\ell}}$ is the field with the smallest extension degree over \mathbb{F}_p which contains an *m* th root of unity other than 1. This can also be verified directly from the definition of ℓ .

For the rest of this section, let $L := \mathbb{F}_{p^{\ell}}$, $m' := \gcd(p^{\ell} - 1, m)$, and let H be group of m' th roots of unity in L. By (5.2), $|H| = m' \geq 2$. It will be important to work with the set $T := \operatorname{tr}(H)$ where "tr" denotes the field trace from L to \mathbb{F}_p . The next theorem gives a description of $W_p(m)$ in terms of ℓ and the cardinality t := |T| (under the standing assumption that $\gcd(p-1, m) = 1$).

Theorem 5.3. Let ℓ and t be as defined above, and let $n := \lceil \frac{p-1}{t-1} \rceil$. Then $\lfloor \ell n, \infty \rangle_{\mathbb{Z}} \subseteq W_p(m') \subseteq W_p(m)$.

Proof. Applying the Cauchy-Davenport Theorem to the subset T in \mathbb{F}_p , we have $|2 * T| \ge \min \{p, 2t - 1\}$, and inductively $|i * T| \ge \min \{p, it - (i - 1)\}$. By the definition of n, we have $n(t-1) \ge p-1$, so $nt - (n-1) \ge p$. Therefore, |n * T| = p. In particular, for every $j \ge 0$, there exists an equation $t_1 + \cdots + t_n = -j \in \mathbb{F}_p$, where all $t_i \in T$. Now each $t_i \in \operatorname{tr}(H)$ is a sum of ℓ elements of H, so $t_1 + \cdots + t_n + j \cdot 1 = 0$ is a vanishing sum of m' th roots of unity of weight $\ell n + j$. This shows that $[\ell n, \infty)_{\mathbb{Z}} \subseteq W_p(m') \subseteq W_p(m)$, as desired.

Note that the above theorem is meaningful only if we know that the trace set $T \subseteq \mathbb{F}_p$ has at least two elements. Fortuitously, this is always the case, according to the following result.

Trace Lemma 5.4. In the notations of (5.3), $t = |T| \ge 2$.

The proof of this lemma will be postponed to the last section (§6). We shall first assume this lemma and try to get to the main conclusions of this section. Note that the larger the trace set T is, the better bound on $W_p(m)$ is given by (5.3). Since $t \ge 2$ by (5.4), we have in any case:

Corollary 5.5. $[\ell(p-1), \infty)_{\mathbb{Z}} \subseteq W_p(m') \subseteq W_p(m).$

Now consider any field $K = \mathbb{F}_{p^k}$ containing the group G of all m th roots of unity. Clearly, $L \subseteq K$, and $H = \dot{L} \cap G$. Let $d = (p^k - 1)/m$ and $d' = (p^{\ell} - 1)/m'$. We now proceed to the proof of the following, which is a stronger version of Theorem 1.3 in the case treated in this section.

Theorem 5.6. Assume that gcd(p-1, m) = 1. Then $[d', \infty)_{\mathbb{Z}} \subseteq W_p(m')$ and $[d, \infty)_{\mathbb{Z}} \subseteq W_p(m)$ except in the following two special cases: (A) d' = p - 1; (B) p = 2, d' = 3, and m' = 5. In these special cases, we have $[d'+1, \infty)_{\mathbb{Z}} \subseteq W_p(m')$ and $[d+1, \infty)_{\mathbb{Z}} \subseteq W_p(m)$.

Proof. Since $d' = [\dot{L} : H] = [\dot{L}G : G]$ divides $d = [\dot{K} : G]$ and $m' \mid m$, it suffices to prove the theorem for $W_p(m')$. Let

(5.7)
$$s = (p^{\ell-1} + \dots + p + 1)/m',$$

so that d' = s(p-1). First let us treat the special case (A), where we have s = 1. Here we are supposed to prove that $[p, \infty)_{\mathbb{Z}} \subseteq W_p(m')$. Since now $[\dot{L}: H] = p - 1$ (and $H \cap \mathbb{F}_p = \{1\}$), we have $\dot{L} = H \cdot \dot{\mathbb{F}}_p$. Therefore, fixing a primitive m'-th root of unity $\alpha \in H$, we have $\alpha - 1 = b^{-1}\alpha^i$ for some integer i and some $b \in \dot{\mathbb{F}}_p$. For convenience, let us think of b as an integer in [1, p - 1]. Multiplying $b \cdot 1 + (p - b) \cdot 1 = 0 \in \mathbb{F}_p$ by α and using the relation $b\alpha = b + \alpha^i$, we get

$$0 = b\alpha + (p-b)\alpha = b \cdot 1 + \alpha^{i} + (p-b)\alpha,$$

which is a vanishing sum of weight p+1. Multiplying this by α again and repeating the argument, we get vanishing sums (of m'-th roots of unity) of weight p+i for any i > 0. Coupled with $p \in W_p(m')$, this gives $[p, \infty)_{\mathbb{Z}} \subseteq W_p(m')$, as desired. For the rest of the proof, we may assume that s > 1. We claim the following:

Lemma 5.8. s > 1 implies that $s \ge \ell$, except perhaps when p = 2 and $\ell = 4, 6, 8, 9$.

Thus, leaving aside the four special cases, we have $d' = s(p-1) \ge \ell(p-1)$, so the desired conclusion for $W_p(m')$ in (5.6) follows from (5.5). The four special cases will have to be treated later.

Proof of (5.8). We go into the following two cases.

Case 1. ℓ is prime. We claim that $\ell \leq q$ for any prime $q \mid s$ (and therefore $\ell \leq s$). In fact, from (5.7), we get $p^{\ell-1} + \cdots + p + 1 \equiv 0 \pmod{q}$, so $p^{\ell} \equiv 1 \pmod{q}$. If $p \not\equiv 1 \pmod{q}$, then p has order ℓ in $U(\mathbb{Z}/q\mathbb{Z})$ (since ℓ is prime), and so $\ell \mid (q-1)$. In this case $\ell \leq q-1 < q$. If $p \equiv 1 \pmod{q}$, then from $p^{\ell-1} + \cdots + p + 1 \equiv \ell \equiv 0 \pmod{q}$, we have in fact $\ell = q$.

Case 2. ℓ is composite. Let q be the smallest prime divisor of ℓ and write $\ell = qt$. Then $1 < t < \ell$ and (5.2) implies that $gcd(p^t - 1, m) = 1$. Since $(p^t - 1) | (p^\ell - 1)$, we see that $p^t - 1$ divides $(p^\ell - 1)/m' = s(p - 1)$. Thus, $s \ge (p^t - 1)/(p - 1)$. We shall now exploit the following elementary fact which is easy to prove using calculus:

Lemma 5.9. $p^x \ge (p-1)x^2 + 1$ for every $x \in [2, \infty)_{\mathbb{Z}}$ with the exception of p = 2and x = 2, 3, 4.

Applying this lemma to x = t, we get the desired conclusion

$$s \ge \frac{p^t - 1}{p - 1} \ge t^2 \ge qt = \ell,$$

except when p = 2 and t = 2, 3, 4. If t = 2, we have q = 2 so $\ell = 4$. If t = 3, we have q = 2, 3, so $\ell = 6$ or 9. Finally, if t = 4, we have q = 2 so $\ell = 8$. This proves (5.8), but we still have to complete the proof of (5.6) in the four special cases noted.

In these cases, $d' = (2^{\ell} - 1)/m'$, so both m', d' are odd (and > 1). We may assume that d' < m'. (For, if $d' \ge m'$, we have $[m', \infty)_{\mathbb{Z}} \subseteq W_2(m')$ since m' is odd, and hence $[d', \infty)_{\mathbb{Z}} \subseteq W_2(m')$.) We simply have to check the four outstanding cases individually.

(1) $\ell = 4$. Here $2^{\ell} - 1 = 15$, so d' = 3, m' = 5, and we are in the case (B) of (5.6). The desired conclusion in this case is $[4, \infty)_{\mathbb{Z}} \subseteq W_2(m')$, which is true since $5 = m' \in W_2(m')$. In fact, by (2.6), we have $W_2(5) = 2\mathbb{N} + 5\mathbb{N} = \{0, 2\} \cup [4, \infty)_{\mathbb{Z}}$. In particular, $3 \notin W_2(5)$, so this case is truly exceptional.

(2) $\ell = 6$. Here $2^{\ell} - 1 = 63$, so we have either d' = 7, m' = 9 or d' = 3, m' = 21. In both cases, $[2, \infty)_{\mathbb{Z}} \subseteq W_2(m')$ (since $2, 3 \in W_2(m')$), so there is no problem. (Actually, in the case d' = 7, we are in the good case $s = d' \ge \ell$ already.)

(3) $\ell = 8$. Here $2^{\ell} - 1 = 255$, so we have either d' = 5, m' = 51 or d' = 15, m' = 17. The latter case presents no problems, since we are once more in the good case $s = d' \ge \ell$. In the former case, $3 \mid m'$ implies that $W_2(m') = [2, \infty)_{\mathbb{Z}}$, so again there is no problem.

(4) $\ell = 9$. Here $2^{\ell} - 1 = 511$, so d' = 7, m' = 73. We have shown in (2.5) that $W_2(73) = \{0\} \cup [2, \infty)_{\mathbb{Z}}$, so there is no problem.

This finally completes the proof of Theorem 5.6.

6. Traces of m th Roots of Unity

In §5, we stated without proof the Trace Lemma 5.4, which was crucial for the proofs of (5.5) and (5.6). In this section, we return to the trace set T = tr(H), and offer a general analysis of T which we believe to be of independent interest. The proof of the Trace Lemma is an easy by-product of this general analysis.

The notations (and hypotheses) introduced at the beginning of §5 will remain in force. In particular, H is the group of m' th roots of unity in $L = \mathbb{F}_{p^{\ell}}$, and "tr" is the field trace from L to \mathbb{F}_p . To enumerate the elements in T, let

(6.1)
$$X^{m'} - 1 = (X - 1)h_1(X) \cdots h_r(X)$$

be the factorization of $X^{m'} - 1$ into (monic) irreducibles over \mathbb{F}_p . Then deg $h_i \geq \ell$ by the definition of ℓ (and the fact that $(X^{m'} - 1) \mid (X^m - 1)$). On the other hand, since L contains all m' th roots of unity, each $h_i(X)$ splits completely in L, so deg $h_i \leq [L : \mathbb{F}_p] = \ell$. Therefore, deg $h_i = \ell$ for all i. Let

(6.2)
$$h_i(X) = X^{\ell} - a_i X^{\ell-1} + \cdots,$$

and let $\{\alpha_{ij}\}$ be all the roots of $h_i(X)$ in L. For each h_i , we can identify the field $\mathbb{F}_p[X]/(h_i(X))$ with L by the correspondence $\overline{X} \leftrightarrow \alpha_{ij}$ (for any j). Therefore, $\operatorname{tr}(\alpha_{ij}) = \sum_k \alpha_{ik} = a_i$ for all i, j. We have thus

(6.3)
$$T = \{ \operatorname{tr}(1), a_1, \cdots, a_r \} = \{ \ell, a_1, \cdots, a_r \},\$$

with possible duplications.

It is now easy to prove the Trace Lemma 5.4, which asserted that $|T| \ge 2$. Assume, for the moment, that T is a singleton. Then, by (6.3), $a_i = \ell$ for all i. Summing all roots of the polynomial in (6.1) (and recalling that $m' \ge 2$), we get

$$0 = 1 + a_1 + \dots + a_r = 1 + r \,\ell = m' \in \mathbb{F}_p$$
,

contradicting the fact that m' is prime to p.

The equation (6.3) gives an upper bound $|T| \leq 1+r$, and this becomes an equality iff the elements listed in (6.3) are distinct. This is the case, for instance, if $\ell = 2$. To see this, note that the constant term of each $h_i(X)$ in (6.1) is 1, since it is an m th root of unity in \mathbb{F}_p , and we are assuming that gcd(p-1, m) = 1. Therefore, if $\ell = 2$, we have $h_i(X) = X^2 - a_iX + 1$. Since the h_i 's are distinct, so are the a_i 's, and of course $a_i \neq 2$ (since otherwise $h_i(X) = (X-1)^2$). Therefore, |T| = 1+r, and (5.3) gives the pretty good estimate $[2n, \infty)_{\mathbb{Z}} \subseteq W_p(m') \subseteq W_p(m)$ with $n = \lceil \frac{p-1}{r} \rceil$. For a simple example of this, let p = 5, and m = 3. Here m' = 3, $\ell = 2$, r = 1, $L = \mathbb{F}_{25}$, and |T| = 2. By (2.6), $W_5(3)$ is the set

$$3\mathbb{N} + 5\mathbb{N} = \{0, 3, 5, 6\} \cup [8, \infty)_{\mathbb{Z}}.$$

Since 2n = 8, the conclusion in (5.3) is sharp here.

In general, t := |T| may be less than 1+r, since there may be duplications among the elements of T listed in (6.3). For an example where interesting duplications occur, take p = 7 and m = 19. Here m' = 19 and $\ell = 3$, r = 6. Mathematica gives a factorization

$$X^{19} - 1 = (X - 1)(X^3 + 2X + 6)(X^3 + 4X^2 + X + 6)(X^3 + 4X^2 + 4X + 6)$$

$$\cdot (X^3 + 5X^2 + 6)(X^3 + 3X^2 + 3X + 6)(X^3 + 6X^2 + 3X + 6) \in \mathbb{F}_7[X].$$

Since -4 = 3 = tr(1) in \mathbb{F}_7 , *T* has only five (two less than 1 + r = 7) distinct elements $\{0, 1, 2, 3, 4\}$. In this case, the number *n* in (5.3) is $\lceil \frac{7-1}{5-1} \rceil = 2$, and (5.3) shows that $[6, \infty)_{\mathbb{Z}} \subseteq W_7(19)$. Note that, in spite of the trace duplications, this is still much sharper than what is given in (5.6).

In general, we cannot hope to improve upon the lower bound $|T| \ge 2$. For one thing, \mathbb{F}_p may have only two elements to begin with. Also, we may have r = 1, in which case (5.4) and (6.3) show that |T| = 2. Even if $p \ge 3$ and $r \ge 2$, there are many cases in which T is just a doubleton. Let us illustrate the situation r = 2 by taking m to be an odd prime q (so that m' = q too), and assuming that p is also odd and has order $\ell = (q - 1)/2$ in the group $U(\mathbb{Z}/q\mathbb{Z})$. In this case, r = 2, and (6.1) becomes

(6.4)
$$X^{q} - 1 = (X - 1)h_{1}(X)h_{2}(X)$$

where h_1 , h_2 are monic irreducible (over \mathbb{F}_p) of degree ℓ . Following a standard notation in number theory, let us define q^* to be q if $q \equiv 1 \pmod{4}$, and q^* to be -q if $q \equiv 3 \pmod{4}$. Then the size of the trace set T = tr(H) is determined as follows.

Proposition 6.5. Under the above assumptions |T| = 2 iff $p | (q^*-1)$ (and |T| = 3 otherwise).

Proof. Let $E = \mathbb{Q}(\zeta)$ where $\zeta = e^{2\pi i/q}$, and fix a generator t of $U(\mathbb{Z}/q\mathbb{Z})$. Then $\sigma : \zeta \mapsto \zeta^t$ is a generator for $\operatorname{Gal}(E/\mathbb{Q})$, and $\sigma^2 : \zeta \mapsto \zeta^{t^2}$ is a generator for $\operatorname{Gal}(E/F)$, where F is the fixed field E^{σ^2} . Note that $X^{q-1} + \cdots + X + 1$ factors into f(X)g(X) over F[X], where

$$f(X) = X^{\ell} - a X^{\ell-1} + \cdots$$
 and $g(X) = X^{\ell} - b X^{\ell-1} + \cdots$

are, respectively, the minimal polynomials of ζ and ζ^t over F. We have

$$a - b = \operatorname{tr}_{E/F}(\zeta) - \operatorname{tr}_{E/F}(\zeta^t) = \sum_{j=0}^{\ell-1} \zeta^{t^{2j}} - \sum_{j=0}^{\ell-1} \zeta^{t^{2j+1}} \in F.$$

This is precisely the quadratic Gauss sum (with respect to the Legendre character on \mathbb{F}_q), so by [IR: (8.2.2)], $a - b = \sqrt{q^*}$. (Gauss showed that the $\sqrt{q^*}$ here is the one taken in the upper half plane if $q \equiv 3 \pmod{4}$, but this will not be needed in the following.) Since we also have a + b = -1, it follows that

$$a = (\sqrt{q^*} - 1)/2, \quad b = -(\sqrt{q^*} + 1)/2.$$

Incidentally, this proves the well-known fact that $F = \mathbb{Q}(\sqrt{q^*})$.

Let R be the ring of algebraic integers in F. Since p is unramified in E, it is also unramified in F, so $pR = \mathfrak{p}\mathfrak{p}'$, where $\mathfrak{p}, \mathfrak{p}'$ are distinct prime ideals of R, both of residue degree 1. Identifying R/\mathfrak{p} with \mathbb{F}_p , we may take the polynomials h_1, h_2 in (6.4) to be \overline{f} and \overline{g} , where "bar" means reduction modulo \mathfrak{p} . In particular, $T = \{\ell, \overline{a}, \overline{b}\}$ by (6.3). Here the two elements $\overline{a}, \overline{b}$ are *always* different (for otherwise \mathfrak{p} would contain $(a-b)^2 = q^*$ as well as p). Therefore, T will have only two elements iff \mathfrak{p} also contains

$$4(\ell - a)(\ell - b) = (q - \sqrt{q^*})(q + \sqrt{q^*}) = q^2 - q^* = q^*(q^* - 1).$$

Since $q^* \notin \mathfrak{p}$, this happens iff $q^* - 1 \in \mathfrak{p}$, that is, iff $p \mid (q^* - 1)$, as claimed.

Corollary 6.6. Let $q = 2\ell + 1$ and $p = 2\ell' + 1$ be distinct primes such that the order of p is ℓ modulo q. If $p \mid (q^* - 1)$, then $[2\ell\ell', \infty)_{\mathbb{Z}} \subseteq W_p(q)$. Otherwise, $[\ell\ell', \infty)_{\mathbb{Z}} \subseteq W_p(q)$.

Proof. This follows from (5.3) and (6.5), since the number n in (5.3) is $2\ell'$ in the first case, and ℓ' in the second case.

Example 6.7. Let q = 11 (with $q^* = -11$). Then the primes 3 and 5 both have order $\ell = (q-1)/2 = 5$ modulo q, and according to *Mathematica*:

$$X^{11} - 1 = (X - 1)(X^5 + X^4 + 2X^3 + X^2 + 2)(X^5 + 2X^3 + X^2 + 2X + 2) \in \mathbb{F}_3[X],$$

$$X^{11} - 1 = (X - 1)(X^5 + 2X^4 + 4X^3 + X^2 + X + 4)(X^5 + 4X^4 + 4X^3 + X^2 + 3X + 4) \in \mathbb{F}_5[X].$$

Thus, for p = 3, $T = \{\ell, -1, 0\} = \{2, 0\}$ in \mathbb{F}_3 . This is consistent with (6.5) since p = 3 divides $q^* - 1 = -12$. Here, $\ell = 5$, $\ell' = 1$, so (6.6) gives $[10, \infty)_{\mathbb{Z}} \subseteq W_3(11)$. (In fact, from $0, 2 \in T$, we see easily that 5, $6 \in W_3(11)$, and so 8, $9 \in W_3(11)$ also.) On the other hand, if we choose p = 5, then $T = \{\ell, -2, -4\} = \{0, 3, 1\}$ in \mathbb{F}_5 , consistently with (6.5) since p = 5 does not divide $q^* - 1 = -12$. Here, $\ell = 5$, $\ell' = 2$, so (6.6) gives again $[10, \infty)_{\mathbb{Z}} \subseteq W_5(11)$, and 0, 1, $3 \in T$ show further that 5, 7, $9 \in W_5(11)$.

The arguments in the proof of (6.5) can be generalized. However, if the order of p modulo q is smaller than (q-1)/2 (in other words r > 2), the computations of the trace elements in T will involve Gaussian sums with (higher) character values as coefficients. We shall not go into this analysis here. We should point out, however, that if q is *fixed*, then the prime ideal method (in characteristic 0) used in the proof of (6.5) will suffice to show that the upper bound $|T| \leq 1 + r$ becomes an *equality* for sufficiently large p. Therefore, by (5.3), $\lfloor \ell n, \infty \rangle_{\mathbb{Z}} \subseteq W_p(q)$ with $n = \lceil \frac{p-1}{r} \rceil$, for sufficiently large p.

References

- [BS] Z. I. Borevich and I. R. Sharfarevich: *Number Theory*, Academic Press, New York/London, 1966.
- [Gr] M. J. Greenberg: Lectures on Forms in Many Variables, Mathematics Lecture Notes Series, W. A. Benjamin, Reading, Mass., 1969.
- [Gu] W. J. Guerrier: The factorization of the cyclotomic polynomial mod p, Amer. Math. Monthly **75**(1968), p.46.
- [IR] K. Ireland and M. Rosen: A Classical Introduction to Modern Number Theory, Graduate Texts in Mathematics, Vol. 84, Springer-Verlag, Berlin-Heidelberg-New York, 1982.
- [La] T. Y. Lam: The Algebraic Theory of Quadratic Forms, Mathematics Lecture Notes Series, W. A. Benjamin, Reading, Mass., 1973. (Second Printing with Revisions, 1980.)
- [LL] T. Y. Lam and K. H. Leung: On vanishing sums of roots of unity, preprint series, MSRI, 1995.
- [LeV] W. LeVeque: Topics in Number Theory, Vol. 1, Addison-Wesley, Reading, Mass., 1956.
- [LN] R. Lidl and H. Niedereiter: *Finite Fields*, Encyclopedia of Mathematics and Its Applications, Vol. 20, Addison-Wesley, 1983.
- [Ma] H. B. Mann: Addition Theorems, Krieger Publ. Co., Huntington, New York, 1976.
- [Sch] W. Schmidt: Equations over Finite Fields, Lecture Notes in Mathematics, Vol. 536, Springer-Verlag, Berlin-Heidelberg-New York, 1976.
- [Sm] C. Small: Arithmetic of Finite Fields, M. Dekker Inc., New York-Basel-Hong Kong, 1991.

- [QY] S. Qi and P.-Z. Yuan: On the number of solutions of diagonal equations over a finite field, Finite Fields and Their Applications 2(1996), 35-41.
- [Wi] E. Witt: Theorie der quadratischen Formen über beliebigen Körpern, J. reine angew. Math. **176**(1937), 31-44.
- T. Y. LAM, MATHEMATICS DEPARTMENT, UNIVERSITY OF CALIFORNIA, BERKELEY, CA, 94720 *E-mail address*: lam@msri.org
- K. H. LEUNG, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119260