

MASSEY AND FUKAYA PRODUCTS ON ELLIPTIC CURVES

A. POLISHCHUK

This note is a complement to [8]. Its goal is to show that some higher Massey products on an elliptic curve can be computed as higher compositions in Fukaya category of the dual symplectic torus in accordance with the homological mirror conjecture of M. Kontsevich [6]. Namely, we consider triple Massey products of very simple type which are uniquely defined, compute them in terms of theta-functions and compare the result with the series one obtains in Fukaya picture. The identity we get in this way was first discovered by Kronecker.

The interesting phenomenon is that although Massey products on an elliptic curve are partially defined and multivalued, one always has the corresponding univalued Fukaya product. The conjecture seems to predict that one can choose canonical representatives for all (not necessary well-defined) Massey products such that the A_∞ -laws are satisfied. In order to understand what kind of functions appear as such representatives we compute some higher compositions m_3 in Fukaya category of a torus corresponding to multiple-valued Massey products on an elliptic curve. The functions appearing in this way can be expressed in terms of theta-functions and the function $\kappa(y, x; \tau)$ given by the series

$$\kappa(y, x; \tau) = \sum_{n \in \mathbb{Z}} \frac{\exp(2\pi i(n^2/2\tau + nx))}{\exp(2\pi i n \tau) - \exp(2\pi i y)}$$

where τ is in the upper half-plane. The latter function was introduced (with slightly different choice of variables) by M. P. Appell in his study of doubly-periodic functions of the third kind in [1]. The A_∞ -constraints between m_2 and m_3 in Fukaya category of a torus give some non-trivial identities involving κ and theta-functions.

Our computation of univalued triple Massey products on elliptic curve can be easily generalized to the case of higher genus curves. The representation of these products as values of a rational section of some universal bundle is closely related to the cube structure of the theta line bundle on the Jacobian of a curve. We expect also that they can be compared with the Fukaya compositions on the symplectic torus,

which is mirror dual to the Jacobian of a curve. This generalization will be considered in another paper.

Throughout this paper we use the notation $e(z) = \exp(2\pi iz)$.

Acknowledgment I am grateful to M. Kontsevich for explaining to me the signs in Fukaya composition and to E. Zaslow for correcting some errors. This work was partially supported by NSF grant.

1. TRIPLE MASSEY PRODUCTS

1.1. General construction. Let $X_1 \xrightarrow{f_1} X_2 \xrightarrow{f_2} X_3 \xrightarrow{f_3} X_4$ be a sequence of morphisms in a triangulated category such that $f_2 \circ f_1 = 0$ and $f_3 \circ f_2 = 0$. Then one can construct the subset of elements $\langle f_3, f_2, f_1 \rangle$ in $\text{Hom}^{-1}(X_1, X_4)$ which is a coset by the sum of the images of the maps

$$\text{Hom}^{-1}(X_2, X_4) \rightarrow \text{Hom}^{-1}(X_1, X_4) : g \mapsto g \circ f_1,$$

$$\text{Hom}^{-1}(X_1, X_3) \rightarrow \text{Hom}^{-1}(X_1, X_4) : h \mapsto f_3 \circ h.$$

Namely, $k \in \langle f_3, f_2, f_1 \rangle$ iff the following diagram is commutative

$$\begin{array}{ccccccc} X_1 & \xrightarrow{f_1} & X_2 & & & & \\ \downarrow k & & \downarrow & \searrow f_2 & & & \\ X_4[-1] & \longrightarrow & Y & \longrightarrow & X_3 & \xrightarrow{f_3} & X_4 \end{array} \quad (1.1.1)$$

where the lower row is an exact triangle. This definition is equivalent to the one given in [3] where one includes f_2 into an exact triangle. In particular, if $\text{Hom}^i(X_1, X_3) = \text{Hom}^i(X_2, X_4) = 0$ for $i = -1, 0$ then the compositions $f_2 f_1$ and $f_3 f_2$ are always zero, the product $\langle f_3, f_2, f_1 \rangle$ contains one element, and we obtain the linear map

$$m_3 : \text{Hom}(X_1, X_2) \otimes \text{Hom}(X_2, X_3) \otimes \text{Hom}(X_3, X_4) \rightarrow \text{Hom}^{-1}(X_1, X_4).$$

Here is a more concrete description of this map. Consider the exact triangle

$$K \rightarrow \text{Hom}(X_3, X_4) \otimes X_3 \rightarrow X_4 \rightarrow K[1]$$

Then our assumptions imply that the following natural maps are isomorphisms:

$$\alpha : \text{Hom}(X_2, K) \rightarrow \text{Hom}(X_3, X_4) \otimes \text{Hom}(X_2, X_3),$$

$$\beta : \text{Hom}^{-1}(X_1, X_4) \rightarrow \text{Hom}(X_1, K).$$

Now m_3 is equal to the following composition

$$\begin{aligned} & \text{Hom}(X_3, X_4) \otimes \text{Hom}(X_2, X_3) \otimes \text{Hom}(X_1, X_2) \xrightarrow{\alpha^{-1} \otimes \text{id}} \\ & \text{Hom}(X_2, K) \otimes \text{Hom}(X_1, X_2) \rightarrow \text{Hom}(X_1, K) \xrightarrow{\beta^{-1}} \text{Hom}^{-1}(X_1, X_4) \end{aligned}$$

where the middle arrow is the natural composition map.

1.2. Some Massey products on elliptic curve. Let us consider the simplest example of triple Massey product on elliptic curve E over a field k . Namely, we want to describe the Massey product

$$\text{Hom}(\mathcal{O}, \mathcal{O}_{x_0}) \otimes \text{Hom}(\mathcal{O}_{x_0}, \mathcal{L}[1]) \otimes \text{Hom}(\mathcal{L}[1], \mathcal{O}_x[1]) \rightarrow \text{Hom}(\mathcal{O}, \mathcal{O}_x)$$

where $x \neq x_0$, $\mathcal{L} \neq \mathcal{O}$, $\deg \mathcal{L} = 0$. If $\mathcal{L}|_{x_0}$ is trivialized then using the Serre duality the source of this arrow can be identified with $(\mathcal{L}|_x \otimes \omega)^*$ where ω is the stalk of the canonical bundle of E at zero, while the target is k . The dual map gives a canonical element $s(\mathcal{L}, x) \in \mathcal{L}|_x \otimes \omega$. Note that $\mathcal{L}|_x$ is a stalk of the Poincaré line bundle \mathcal{P} on $E \times \hat{E}$ where \hat{E} is the dual elliptic curve. As we'll see below the above Massey product is the value at the point (\mathcal{L}, x) of the canonical rational section of $\mathcal{P} \otimes \omega$ (with poles at $x = x_0$ and $\mathcal{L} = \mathcal{O}$).

Proposition 1.1. *The above element $s(\mathcal{L}, x)$ corresponds to the following composition*

$$k \simeq \mathcal{L}|_{x_0} \xrightarrow{s_x(x_0)^{-1}} \mathcal{L}(-x)|_{x_0} \xrightarrow{r_{x_0}} H^0(E, \mathcal{L}^{-1}(x))^* \xrightarrow{r_x^{-1}} \mathcal{L}(-x)|_x \simeq \mathcal{L}|_x \otimes \omega \quad (1.2.1)$$

where s_x is the canonical global section of $\mathcal{O}(x)$, r_z for any $z \in E$ is the dual of the evaluation map $H^0(E, \mathcal{L}^{-1}(x)) \rightarrow \mathcal{L}^{-1}(x)|_z$ (note that r_x is invertible if $\mathcal{L} \neq \mathcal{O}$).

The proof is straightforward and is left to the reader.

Now assume that $k = \mathbb{C}$ and the elliptic curve E is \mathbb{C}/Γ where $\Gamma = \Gamma_\tau = \mathbb{Z} + \mathbb{Z}\tau$, τ is in the upper-half plane. Let us consider the line bundle L on E (equipped with a trivialization of its pull-back to \mathbb{C}/\mathbb{Z}) such that the classical theta-function

$$\theta(z, \tau) = \sum_n e(\tau n^2/2 + nz)$$

is a section of L (as a function of z), so $L \simeq \mathcal{O}(\xi)$ where $\xi = (\tau + 1)/2$. Now set $\mathcal{L} = t_y^* L \otimes L^{-1}$ where $y \in \mathbb{C}$ is not a lattice point. Then $\mathcal{L}^{-1}(x) \simeq t_{-x-y+\xi}^* L$ so that $t_{-x-y+\xi}^* \theta$ generates its space of global sections. Under the natural isomorphisms we have $s_x = \theta(-x + \xi, \tau)$,

the map r_{x_0} becomes multiplication by $\theta(-x - y + \xi, \tau)$ while $r_x^{-1} = \theta(-y + xi, \tau)^{-1}$. So the composition is given by

$$s(\mathcal{L}, x) = \theta(-x - y + (\tau + 1)/2, \tau) \theta(-x + (\tau + 1)/2, \tau)^{-1} \theta(-y + (\tau + 1)/2, \tau)^{-1}.$$

Actually the above computation only works for a fixed elliptic curve E so below we'll have to multiply the obtained expression by some constant depending only on τ to get a canonical section of $\mathcal{P} \otimes \omega$.

2. COMPARISON WITH FUKAYA COMPOSITION

2.1. Fukaya series. Let us consider the Fukaya composition of three morphisms considered in 1.2 (see [8] for notation). The corresponding four objects of the Fukaya category of a torus are: $\Lambda_1 = (t, 0)$ with trivial connection, $\Lambda_2 = (0, t)$ with trivial connection, $\Lambda_3 = (t, -\alpha_1)$ with the connection $(-2\pi i \beta_1)dx$, $\Lambda_4 = (-\alpha_2, t)$ with the connection $2\pi i \beta_2 dy$. Here $\alpha_1, \alpha_2, \beta_1, \beta_2$ are real numbers. There is an essentially unique choice of logarithms of slopes for Λ_i such that $\text{Hom}^0(\Lambda_i, \Lambda_{i+1}) \neq 0$ and with such a choice one has $\text{Hom}^{-1}(\Lambda_1, \Lambda_4) \neq 0$. Moreover, all these spaces are one-dimensional so the corresponding Fukaya composition m_3 is just a number. This composition is defined only if α_1 and α_2 are not integers. Then it is given by the following series

$$\sum_{(\alpha_1+m)(\alpha_2+n)>0} \text{sign}(\alpha_1+m) e(\tau(\alpha_1+m)(\alpha_2+n) + (\alpha_1+m)\beta_2 + (\alpha_2+n)\beta_1)$$

where we denote $\text{sign}(t) = 1$ for $t > 0$, $\text{sign}(t) = -1$ for $t < 0$, the sum is over integers m and n subject to the condition that $(\alpha_1 + m)$ has the same sign as $(\alpha_2 + n)$, The restriction on m and n is imposed by the condition for maps from the disk to be holomorphic in the definition of Fukaya composition while the sign comes from the canonical orientation of the corresponding moduli space. We can write the above expression in the form

$$e(\tau\alpha_1\alpha_2 + \alpha_1\beta_2 + \alpha_2\beta_1) f(\alpha_1\tau + \beta_1, \alpha_2\tau + \beta_2; \tau)$$

where

$$f(z_1, z_2; \tau) = \sum_{(\alpha(z_1)+m)(\alpha(z_2)+n)>0} \text{sign}(\alpha(z_1) + m) e(\tau mn + nz_1 + mz_2)$$

is a holomorphic function of z_1 and z_2 defined for $\text{Im } z_i \notin \mathbb{Z}(\text{Im } \tau)$, where $\alpha(z) = \text{Im}(z)/\text{Im}(\tau)$.

2.2. Poles. We claim that the above function $f(z_1, z_2; \tau)$ extends to a meromorphic function in z_1, z_2 with poles at the lattice points $z_1 \in \Gamma_\tau$ or $z_2 \in \Gamma_\tau$ where $\Gamma_\tau = \mathbb{Z} + \mathbb{Z}\tau$. The following identities are easy to check:

$$f(z_2, z_1; \tau) = f(z_1, z_2; \tau), \quad (2.2.1)$$

$$f(z_1 + m + n\tau, z_2; \tau) = e(-nz_2)f(z_1, z_2; \tau), \quad (2.2.2)$$

$$f(z_1, z_2 + m + n\tau; \tau) = e(-nz_1)f(z_1, z_2; \tau). \quad (2.2.3)$$

Thus, in order to prove the above claim it suffices to show that $f(z_1, z_2) = f(z_1, z_2; \tau)$ extends to a meromorphic function in the region

$$U = \{(z_1, z_2) : 0 < \text{Im}(z_2) < \text{Im}(\tau), -\text{Im}(\tau) < \text{Im}(z_1) < \text{Im}(\tau)\},$$

with poles only at $p_1^{-1}(\mathbb{Z})$ where p_1 is the projection to the first coordinate. If $(z_1, z_2) \in U$ and $\text{Im}(z_1) > 0$ then

$$\begin{aligned} f(z_1, z_2) &= \sum_{m \geq 0, n \geq 0} e(mn\tau + nz_1 + mz_2) - \sum_{m < 0, n < 0} e(mn\tau + nz_1 + mz_2) = \\ \sum_{n \geq 0} e(nz_1) + \tilde{f}(z_1, z_2) &= \frac{1}{1 - e(z_1)} + \tilde{f}(z_1, z_2) \end{aligned}$$

where

$$\tilde{f}(z_1, z_2) = \sum_{m(\alpha(z_2) + n) > 0} \text{sign}(m) e(mn\tau + nz_1 + mz_2)$$

is holomorphic on U . On the other hand, if $(z_1, z_2) \in U$ and $\text{Im}(z_1) < 0$ then

$$\begin{aligned} f(z_1, z_2) &= \sum_{m > 0, n \geq 0} e(mn\tau + nz_1 + mz_2) - \sum_{m \leq 0, n < 0} e(mn\tau + nz_1 + mz_2) = \\ - \sum_{n < 0} e(nz_1) + \tilde{f}(z_1, z_2) &= -\frac{e(-z_1)}{1 - e(-z_1)} + \tilde{f}(z_1, z_2). \end{aligned}$$

Taking into account the identity

$$\frac{1}{1 - e(z_1)} = -\frac{e(-z_1)}{1 - e(-z_1)}$$

we get the required continuation.

The above computation also shows that the poles of $f(z_1, z_2; \tau)$ at the divisors $z_1 = \lambda$ and $z_2 = \lambda$ where $\lambda \in \Gamma_\tau$ are of order one.

2.3. Identity. Consider the function

$$\phi(z_1, z_2; \tau) = \theta(z_1 - (\tau + 1)/2, \tau) \theta(z_2 - (\tau + 1)/2, \tau) f(z_1, z_2; \tau).$$

Since $\theta((\tau + 1)/2, \tau) = 0$, $\phi(z_1, z_2; \tau)$ is a holomorphic function in z_1 and z_2 . Moreover, using the transformation formula

$$\theta(z + m + n\tau, \tau) = e(-n^2\tau/2 - nz) \theta(z, \tau)$$

and the formulas (2.2.2), (2.2.3) we get

$$\phi(z_1 + m + n\tau, z_2; \tau) = \phi(z_1, z_2 + m + n\tau; \tau) = e(-n^2\tau/2 - n(z_1 + z_2 - (\tau + 1)/2)) \phi(z_1, z_2; \tau).$$

But there is only one holomorphic function of z_1, z_2 satisfying this transformation law (up to a constant), namely, $\theta(z_1 + z_2 - (\tau + 1)/2, \tau)$. Hence, we obtain the identity

$$\phi(z_1, z_2; \tau) = c_\tau \theta(z_1 + z_2 - (\tau + 1)/2, \tau) \quad (2.3.1)$$

where c_τ is a non-zero constant depending on τ . Below we determine this constant substituting particular values of z_1 and z_2 . The identity (2.3.2) was first discovered by Kronecker [7] (see also [9], ch. VIII).

Theorem 2.1. *One has the following identity*

$$f(z_1, z_2; \tau) = \prod_{n \geq 1} (1 - q^n)^3 \cdot \frac{\theta(z_1 + z_2 - (\tau + 1)/2, \tau)}{\theta(z_1 - (\tau + 1)/2, \tau) \theta(z_2 - (\tau + 1)/2, \tau)} \quad (2.3.2)$$

where $q = e(\tau)$.

Proof. In order to determine the constant c_τ let us substitute $z_1 = (\tau + 1)/2$, $z_2 = \tau/2$ in (2.3.1). Then we obtain

$$\begin{aligned} \sum_{m \geq 0, n \geq 0} e(mn\tau + \frac{m}{2} + \frac{m+n}{2}\tau) - \sum_{m < 0, n < 0} e(mn\tau + \frac{m}{2} + \frac{m+n}{2}\tau) &= c_\tau \frac{\theta(\tau/2, \tau)}{\theta(0, \tau) \theta(1/2, \tau)} \quad (2.3.3) \\ &= c_\tau \frac{\theta(\tau/2, \tau)}{\theta(0, \tau) \theta(1/2, \tau)} \end{aligned}$$

Let us rewrite the LHS in a more familiar form. First of all replacing m and n by $-m - 1$ and $-n - 1$ in the second sum one can easily see that the LHS is equal to

$$2 \sum_{m \geq 0, n \geq 0} e(mn\tau + \frac{m}{2} + \frac{m+n}{2}\tau) = 2 \sum_{m \geq 0, n \geq 0} (-1)^m q^{mn + (m+n)/2}.$$

Now we notice that

$$\begin{aligned} \sum_{m \geq 0, n \geq 0} (-1)^m q^{mn + (m+n)/2} &= \sum_{m \geq 0, n \geq 0} (-1)^m q^{(m+1/2)(n+1/2) - 1/4} = \\ &= q^{-1/4} \cdot \sum_{m \geq 0, n \geq 0} (q^{(4m+1)(4n+1)/4} - q^{(4m+3)(4n+3)/4}). \end{aligned}$$

In the last step we noticed that the part of the sum over m and n of different parity is zero and then replaced (m, n) by $(2m, 2n)$ and $(2m + 1, 2n + 1)$. The obtained series is equal to

$$q^{-1/4} \sum_{k \geq 1} \delta(k) q^{k/4}$$

where $\delta(k) = 0$ for even k and $\delta(k) = \sum_{d|k} (-1)^{(d-1)/2}$ for odd k . It is well-known that $4\delta(k)$ is equal to $r(k)$, the number of representations of k as sum of two squares. Basing on this it is easy to show that $\delta(k)$ is equal to the number of representations of $2k$ in the form $2k = (2m + 1)^2 + (2n + 1)^2$ where $m \geq 0, n \geq 0$. Hence, we have the identity

$$q^{-1/4} \sum_{k \geq 1} \delta(k) q^{k/4} = q^{-1/4} \left(\sum_{n \geq 0} q^{(2n+1)^2/8} \right)^2 = \left(\sum_{n \geq 0} q^{n(n+1)/2} \right)^2 = \frac{1}{4} \theta(\tau/2, \tau)^2.$$

Hence, we have shown that the LHS of (2.3.3) is equal to $\frac{1}{2} \theta(\tau/2, \tau)^2$. Thus, we have

$$c_\tau = \frac{1}{2} \theta(0, \tau) \theta(1/2, \tau) \theta(\tau/2, \tau).$$

Using the representation of θ as an infinite product one can easily show that the right hand side is equal to $\prod_{n \geq 1} (1 - q^n)^3$. \square

If we fix z_2 and consider residues of both parts of the identity (2.3.2) at $z_1 = 0$ we obtain

$$\text{Res}_{z_1=0} f(z_1, z_2; \tau) = \prod_{n \geq 1} (1 - q^n)^3 \theta'(-(\tau + 1)/2, \tau).$$

On the other hand, the reasoning in 2.2 implies that

$$\text{Res}_{z_1=0} f(z_1, z_2; \tau) = \text{Res}_{z_1=0} \frac{1}{1 - \exp(2\pi i z_1)} = -\frac{1}{2\pi i}.$$

Thus, we recover the identity

$$\prod_{n \geq 1} (1 - q^n)^3 = \frac{1}{2\pi i} \theta'((\tau + 1)/2, \tau) = \sum_{n \in \mathbb{Z}} (-1)^n n q^{(n^2+n)/2}.$$

Alternatively, one can use this identity to give another proof of (2.3.2).

2.4. Functional equation for $f(z_1, z_2; \tau)$. It is clear from the definition that

$$f(z_1, z_2; \tau + 1) = f(z_1, z_2; \tau).$$

Now using the identity (2.3.2) one can easily deduce from the functional equation for theta function that $f(z_1, z_2; \tau)$ satisfies the functional equation of the form

$$f(z_1/\tau, z_2/\tau; -\tau^{-1}) = \zeta \cdot \tau \cdot e(z_1 z_2 / \tau) f(z_1, z_2; \tau)$$

where ζ is a root of unity. Using the property

$$f(-z_1, -z_2; \tau) = -f(z_1, z_2; \tau)$$

we immediately conclude that $\zeta^2 = 1$, so $\zeta = \pm 1$. Finally substituting $z_1 = (\tau + 1)/2$, $z_2 = \tau/2$ and looking at the sign of both sides when $\tau = it$, $t \in \mathbb{R}$ and $t \rightarrow +\infty$ we find that $\zeta = 1$. So the functional equation for f becomes

$$f(z_1/\tau, z_2/\tau; -\tau^{-1}) = \tau \cdot e(z_1 z_2/\tau) f(z_1, z_2; \tau).$$

The functional equation can also be derived from the representation of f in the following form (cf. [9], ch.VIII):

$$f(z_1, z_2; \tau) = -\frac{e(-\alpha(z_2)z_1)}{2\pi i} \sum_e \frac{\chi(w)}{z_1 + w}$$

provided that $0 < \alpha(z_i) < 1$ for $i = 1, 2$. Here \sum_e denotes the Eisenstein summation over the lattice Γ_τ , χ is the character of Γ_τ such that $\chi(1) = e(-\alpha(z_2))$, $\chi(\tau) = e(z_2 - \alpha(z_2)\tau)$.

3. SOME HIGHER COMPOSITIONS IN FUKAYA CATEGORY

3.1. Trapezoid compositions. Now we are going to consider some compositions m_3 in Fukaya category of a torus such that the corresponding triple Massey products on elliptic curve are not well-defined. Namely, consider four lagrangians: $\Lambda_1 = (t, -t)$ and $\Lambda_4 = (t, 0)$ with trivial connections, $\Lambda_2 = (t, -\alpha_2)$ with the connection $-2\pi i \beta_2 dx$, and $\Lambda_3 = (-\alpha_1, t)$ with the connection $2\pi i \beta_1 dy$, where α_i, β_i are real numbers, $\alpha_i \notin \mathbb{Z}$. There is a natural choice of logarithms of slopes so that $\text{Hom}^0(\Lambda_i, \Lambda_{i+1}) \simeq \mathbb{C}$ and $\text{Hom}^{-1}(\Lambda_1, \Lambda_4) \simeq \mathbb{C}$. Note that $\text{Hom}^0(\Lambda_1, \Lambda_3) \neq 0$, so the corresponding triple Massey product on elliptic curve is not defined. The Fukaya composition

$$m_3 : \text{Hom}(\Lambda_1, \Lambda_2) \otimes \text{Hom}(\Lambda_2, \Lambda_3) \otimes \text{Hom}(\Lambda_3, \Lambda_4) \rightarrow \text{Hom}^{-1}(\Lambda_1, \Lambda_4)$$

is just the number given by the series

$$\sum_{(n+\alpha_1)(m+\alpha_2)>0} \text{sign}(m+\alpha_2) e((n+\alpha_1+(m+\alpha_2)/2)(m+\alpha_2)\tau + (m+\alpha_2)\beta_1 + (m+\alpha_2+n+\alpha_1)\beta_2).$$

Let us define

$$g(z_1, z_2; \tau) = \sum_{(n+\alpha(z_1))(m+\alpha(z_2))>0} \text{sign}(m+\alpha(z_2)) e((n+m/2)m\tau + mz_1 + (m+n)z_2)$$

where as before $\alpha(z) = \text{Im}(z)/\text{Im}(\tau)$, $\alpha(z_1) \notin \mathbb{Z}$, $\alpha(z_2) \notin \mathbb{Z}$. Then the above composition is equal to

$$e((\alpha_1 + \alpha_2/2)\alpha_2\tau + \alpha_2\beta_1 + (\alpha_1 + \alpha_2)\beta_2)g(\alpha_1\tau + \beta_1, \alpha_2\tau + \beta_2; \tau).$$

3.2. Properties of $g(z_1, z_2; \tau)$. The function g is holomorphic for $\alpha(z_i) \notin \mathbb{Z}$, $i = 1, 2$, and satisfies the following quasi-periodicity identities

$$g(z_1 + m + n\tau, z_2; \tau) = e(-nz_2)g(z_1, z_2; \tau), \quad (3.2.1)$$

$$g(z_1, z_2 + m + n\tau; \tau) = e(-n^2\tau/2 - n(z_1 + z_2))g(z_1, z_2; \tau). \quad (3.2.2)$$

In other words, g can be considered as a holomorphic section of a line bundle on E^2 over the open subset $(E \setminus S)^2$ where $S = \mathbb{R}/\mathbb{Z} \subset \mathbb{C}/\Gamma_\tau$. However, it doesn't extend to a meromorphic section on E^2 . On the other hand, let us denote by g_0 the restriction of g to the region $0 < \alpha(z_1) < 1$, $\alpha(z_2) \notin \mathbb{Z}$. Then we claim that g_0 extends to a meromorphic function on \mathbb{C}^2 with poles of order 1 at $z_2 \in \Gamma_\tau$. Indeed, if we sum first over n in the series defining g we get

$$g_0(z_1, z_2; \tau) = \sum_{m \in \mathbb{Z}} \frac{e(m^2\tau/2 + m(z_1 + z_2))}{1 - e(m\tau + z_2)}.$$

The latter series clearly satisfies the properties we claimed. However, g_0 lacks the quasi-periodicity of g . More precisely, it is easy to see that g and g_0 are related as follows:

$$g_0(z_1, z_2; \tau) - g(z_1, z_2; \tau) = p(z_1, \tau)\theta(z_1 + z_2; \tau)$$

where $p(z, \tau)$ is the following piecewise polynomial function of $e(z_1)$:

$$p(z, \tau) = \begin{cases} -\sum_{0 < n \leq \alpha(z)} e(-n^2\tau/2 + nz_1), & \alpha(z) \geq 0, \\ \sum_{\alpha(z) < n \leq 0} e(-n^2\tau/2 + nz_1), & \alpha(z) < 0 \end{cases}$$

Let us consider the following series

$$\kappa(y, x; \tau) = \sum_{n \in \mathbb{Z}} \frac{e(n^2/2\tau + nx)}{e(n\tau) - e(y)}. \quad (3.2.3)$$

This function is holomorphic for $y \notin \Gamma_\tau$ and satisfies the difference equation

$$\kappa(y, x + m + \tau; \tau) = e(y)\kappa(y, x; \tau) + \theta(x, \tau)$$

where $m \in \mathbb{Z}$. Then we have

$$g_0(z_1, z_2; \tau) = -e(-z_2)\kappa(-z_2, z_1 + z_2; \tau) = \kappa(z_2, \tau - z_1 - z_2; \tau). \quad (3.2.4)$$

Remark. The function κ and its derivatives were used by M. P. Appell to represent an arbitrary doubly-periodic function of the third kind as a sum of simple elements (cf. [1],[5]). On the other hand, κ can be expressed via the bilateral basic hypergeometric series. Namely, using notation of [2] we have

$$\kappa(y, x; \tau) = (1 - e(-y))^{-1} \cdot {}_1\psi_2(e(-y); 0, e(\tau - y); e(\tau), e(x + (\tau + 1)/2)).$$

3.3. Trapezoid Massey products. The geometric meaning of the function κ above is the following: the pair $(\kappa(y, x; \tau), \theta(x, \tau))$ determines a global section of a rank 2 bundle F_y on E which is a non-trivial extension of L by the line bundle of degree 0 corresponding to y . Namely, the bundle F_y is defined as follows

$$F_y = \mathbb{C}^* \times \mathbb{C}^2 / (z, v) \mapsto (z \cdot e(\tau), A_y(z)v)$$

where

$$A_y(z) = \begin{pmatrix} e(y) & 1 \\ 0 & e(-\tau/2)z^{-1} \end{pmatrix}.$$

We claim that some of triple Massey products corresponding to the trapezoid Fukaya products considered above are univalued and are also expressed via κ . Namely, consider the triple Massey product for $X_1 = \mathcal{O}$, $X_2 = L$, $X_3 = \mathcal{L}[1]$, $X_4 = \mathcal{O}_\xi[1]$ where \mathcal{L} is a non-trivial line bundle of degree 0, $\xi = (\tau + 1)/2$. Then $\text{Hom}^*(X_1, X_3) = 0$ and $\text{Hom}^0(X_2, X_4) = 0$. Furthermore, the composition map

$$\text{Hom}(X_1, X_2) \otimes \text{Hom}^{-1}(X_2, X_4) \rightarrow \text{Hom}^{-1}(X_1, X_4)$$

is zero since the unique section of L vanishes at ξ . It follows that the Massey product is well-defined and univalued in this case. To compute it one should include the non-zero morphism $X_2 \rightarrow X_3$ into an exact triangle

$$X_2 \rightarrow X_3 \rightarrow C \rightarrow X_2[1]$$

then lift the morphisms $X_1 \rightarrow X_2$ and $X_3 \rightarrow X_4$ to morphisms $X_1 \rightarrow C[-1]$ and $C \rightarrow X_4$ and compose the obtained two morphisms. In our case one starts with a global section $s : \mathcal{O} \rightarrow L$, then s can be lifted canonically to a section \tilde{s} of F_y where $\mathcal{L} \simeq t_y L^{-1} \otimes L$. Now one chooses a splitting $r : (F_y)_\xi \rightarrow \mathbb{C}$ of the embedding $\mathbb{C} \simeq \mathcal{L}_\xi \rightarrow (F_y)$ and applies r to $\tilde{s}(\xi)$. The result doesn't depend on a choice of splitting at ξ since $s(\xi) = 0$. More concretely, for $s = \theta(x, \tau)$ we have $\tilde{s} = \kappa(y, x; \tau)$, hence, the above Massey product is given by $\kappa(y, (\tau + 1)/2; \tau)$. One can easily check that this answer agrees with the corresponding Fukaya product. Note also that in fact this Massey product can still be expressed via theta functions due to the identity

$$\kappa(y, \frac{\tau + 1}{2}; \tau) = \prod_{n \geq 1} (1 - e(n\tau))^3 \cdot \frac{1}{\theta(y - \frac{\tau + 1}{2}, \tau)}$$

If we replace ξ by another point on E the Massey product will no longer be univalued. However, if one chooses a splitting of the embedding $\mathcal{L} \rightarrow F_y$ over some open subset $U \subset E$ and a trivialization

of $\mathcal{L}|_U$ then replacing r by this splitting we get a univalued operation. The function $\kappa(y, x; \tau)$ appears as such operation corresponding to the choice of a splitting over $E \setminus S$ coming from the trivialization of the pull-back of F_y to \mathbb{C}^* .

3.4. Associativity constraint. Let us consider an example of associativity constraint for Fukaya's A_∞ -category of a torus involving triple products computed above. Namely, since $m_1 = 0$ we have the following identity

$$m_3(m_2(a_1, a_2), a_3, a_4) \pm m_3(a_1, m_2(a_2, a_3), a_4) \pm m_3(a_1, a_2, m_2(a_3, a_4)) \\ \pm m_2(m_3(a_1, a_2, a_3), a_4) \pm m_2(a_1, m_3(a_2, a_3, a_4)) = 0$$

where a_1, \dots, a_4 is the sequence of composable morphisms.

Note that the proof of A_∞ -identities for the Fukaya compositions in our case is easy. Basically they follow from additivity of the area of plane figures. For example, the usual associativity of m_2 (which is equivalent to the addition formula for theta function) can be proved by considering two ways of cutting a non-convex quadrangle into two triangles and converting this into the identity for the corresponding generating series. Similarly, one can prove higher A_∞ -identities.

Let us consider the following five lagrangians in $\mathbb{R}^2/\mathbb{Z}^2$: $\Lambda_1 = (t, -t)$ with trivial connection, $\Lambda_2 = (t, \alpha_2)$ with $2\pi i \beta_2 dx$, $\Lambda_3 = (-\alpha_1 - \alpha_2, t)$ with $2\pi i(\beta_1 + \beta_2)dy$, $\Lambda_4 = (t, \alpha_1 + \alpha_2 + \alpha_3)$ with $2\pi i(\beta_1 + \beta_2 + \beta_3)dx$, $\Lambda_5 = (0, t)$ with trivial connection. Here α_i, β_i are real numbers, $\alpha_1 + \alpha_2 \notin \mathbb{Z}$, $\alpha_1 + \alpha_3 \notin \mathbb{Z}$. We choose liftings of Λ_i to objects in Fukaya category in such a way that there is a non-zero morphism a_i of degree 0 from Λ_i to Λ_{i+1} . We want to write the above A_∞ -identity for these morphisms. Note that all the Hom-spaces between our objects are either zero or one-dimensional, so the relevant compositions m_2 and m_3 are just numbers. Taking into account the fact that $\Lambda_2 \cap \Lambda_4 = \Lambda_3 \cap \Lambda_5 = \emptyset$ the identity boils down to

$$m_2(\Lambda_1, \Lambda_2, \Lambda_3)m_3(\Lambda_1, \Lambda_3, \Lambda_4, \Lambda_5) + m_2(\Lambda_1, \Lambda_4, \Lambda_5)m_3(\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4) - \\ m_2(\Lambda_1, \Lambda_2, \Lambda_5)m_3(\Lambda_2, \Lambda_3, \Lambda_4, \Lambda_5) \neq 0 \quad (3.4.1)$$

where for example $m_2(\Lambda_1, \Lambda_2, \Lambda_3)$ is the unique non-zero m_2 -composition of morphisms between $\Lambda_1, \Lambda_2, \Lambda_3$, etc.

Now we can express all ingredients of (3.4.1) in terms of theta-functions, and functions f and g introduced above. Namely, denoting

$z_i = \alpha_i \tau + \beta_i$ for $i = 1, 2, 3$ we obtain

$$\begin{aligned}
m_2(\Lambda_1, \Lambda_2, \Lambda_3) &= e(\alpha_1^2 \tau / 2 + \alpha_1 \beta_1) \theta(z_1, \tau), \\
m_2(\Lambda_1, \Lambda_4, \Lambda_5) &= e((\alpha_1 + \alpha_2 + \alpha_3)^2 \tau / 2 + (\alpha_1 + \alpha_2 + \alpha_3)(\beta_1 + \beta_2 + \beta_3)) \theta(z_1 + z_2 + z_3, \tau), \\
m_2(\Lambda_1, \Lambda_2, \Lambda_5) &= e(\alpha_2^2 \tau / 2 + \alpha_2 \beta_2) \theta(z_2, \tau), \\
m_3(\Lambda_1, \Lambda_3, \Lambda_4, \Lambda_5) &= \\
&e((\alpha_1 / 2 + \alpha_2 / 2 + \alpha_3)(\alpha_1 + \alpha_2) \tau + (\alpha_1 + \alpha_2)(\beta_1 + \beta_2 + \beta_3) + \alpha_3(\beta_1 + \beta_2)) g(z_3, z_1 + z_2; \tau), \\
m_3(\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4) &= e((\alpha_1 / 2 - \alpha_3^2 / 2) \tau + \alpha_1 \beta_1 - \alpha_3 \beta_3) g(-z_3, z_1 + z_3; \tau), \\
m_3(\Lambda_2, \Lambda_3, \Lambda_4, \Lambda_5) &= \\
&e((\alpha_1 + \alpha_2)(\alpha_1 + \alpha_3) \tau + (\alpha_1 + \alpha_2)(\beta_1 + \beta_3) + (\alpha_1 + \alpha_3)(\beta_1 + \beta_2)) f(z_1 + z_2, z_1 + z_3; \tau).
\end{aligned}$$

Substituting this into (3.4.1) and deleting similar terms we obtain the following identity

$$\theta(z_1, \tau) g(z_3, z_1 + z_2; \tau) + \theta(z_1 + z_2 + z_3, \tau) g(-z_3, z_1 + z_3; \tau) = \theta(z_2, \tau) f(z_1 + z_2, z_1 + z_3; \tau). \quad (3.4.2)$$

By (3.2.4) this implies the following identity between meromorphic functions of \mathbb{C}^3 :

$$e(y) \theta(y + z, \tau) \kappa(y, z - x; \tau) - e(-x) \theta(x - z, \tau) \kappa(-x, y + z; \tau) = \theta(z, \tau) f(x, y; \tau) \quad (3.4.3)$$

where we have put $x = z_1 + z_2$, $y = z_1 + z_3$, $z = z_2$.

We used the fact that Fukaya composition satisfies axioms of A_∞ -category to derive the above identity. However, it can be also proved in a straightforward way comparing residues of both sides at poles and using the difference equation for κ . It appears in a slightly different form in Halphen's book [5] (p.481, formula (45) and the next one).

3.5. More Fukaya products. Let us consider another example of triple Fukaya products where none of the four lines are parallel. Namely, let $\Lambda_1 = (\alpha_1 + t, -\alpha_1 + t)$, $\Lambda_2 = (-\alpha_2, t)$, $\Lambda_3 = (t, 0)$, $\Lambda_4 = (t, -t)$. Then we can choose (essentially uniquely) lifts of the corresponding lagrangian circles in $\mathbb{R}^2 / \mathbb{Z}^2$ to objects of Fukaya category in such a way that there is a non-zero morphism of degree zero from Λ_i to Λ_{i+1} . All the Hom-spaces $\text{Hom}(\Lambda_i, \Lambda_{i+1})$ are 1-dimensional, however, $\text{Hom}^{-1}(\Lambda_1, \Lambda_4)$ is 2-dimensional since the corresponding circles have 2 points of intersection. Let us consider the component of the triple product corresponding to the point $(\alpha_1, -\alpha_1) \in \Lambda_1 \cap \Lambda_4$. Then as before its value is expressed via certain holomorphic function of $z_1 = \alpha_1 \tau$ and $z_2 = \alpha_2 \tau$ (one could add some monodromies in the above picture

to get all values of complex variables z_1 and z_2). This function has the following form

$$h(z_1, z_2; \tau) = \sum_{(m+\alpha(z_1))(n+\alpha(z_2))>0} \text{sign}(m + \alpha(z_1)) e \left(\frac{\tau}{2} (2m^2 + 4mn + n^2) + 2(m+n)z_1 + (2m+n)z_2 \right).$$

It is holomorphic in the region $\alpha(z_i) \notin \mathbb{Z}$ and can be considered as a section of a line bundle on E^2 over the corresponding open subset of E^2 because of the following quasi-periodicity:

$$h(z_1 + m + \tau, z_2; \tau) = e(-\tau - 2z_1 - 2z_2)h(z_1, z_2; \tau), \quad (3.5.1)$$

$$h(z_1, z_2 + m + \tau; \tau) = e(-\tau/2 - 2z_1 - z_2)h(z_1, z_2; \tau). \quad (3.5.2)$$

Let us denote by h_0 the restriction of h to the region $0 < \alpha(z_i) < 1$ for $i = 1, 2$. Then it is easy to see that h_0 extends to a holomorphic function on \mathbb{C}^2 (the series converges absolutely).

As before we can translate certain A_∞ -associativity axiom into an identity involving $h(z_1, z_2)$. More precisely, we can consider five lines: two with slope 0, one with slope 1, one with slope -1, and one with slope ∞ . Their relative position is described by three parameters. Adding monodromies we get the following identity with three complex variables z_1, z_2, z_3 :

$$\theta(2z_1 + z_3, \tau)h(z_1 + z_3, -z_1 + z_2 - z_3; \tau) + \theta(z_1 + z_2 + z_3, \tau)h(-z_1 - z_3, z_3; \tau) = \theta(2z_2, 2\tau)g(-z_1 + z_2 - z_3, -z_1 - z_2; \tau) + \theta(2z_1, 2\tau)g(z_3, z_1 + z_2; \tau). \quad (3.5.3)$$

Using difference equations for h and g we can further transform this into the following identity involving h_0 and κ :

$$\theta(2x + y, \tau)h_0(x, z; \tau) - \theta(2x + z, \tau)h_0(x, y; \tau) = \theta(2(x + z), 2\tau)\kappa(-2x - y - z, 2x + y + \tau; \tau) - \theta(2(x + y), 2\tau)\kappa(-2x - y - z, 2x + z + \tau; \tau). \quad (3.5.4)$$

Substituting $y = -x$ in this identity we can express $h_0(x, z; \tau)$ via $h_0(x, -x; \tau)$ and functions κ and θ . We claim that $h_0(x) := h_0(x, -x; \tau)$ can also be expressed as a rational function of κ and θ . Indeed, we have

$$h_0(x) = \sum_{(m+1/2)(n+1/2)>0} \text{sign}(m + 1/2) e \left(\frac{\tau}{2} (2m^2 + 4mn + n^2) + nx \right).$$

Hence,

$$h_0(x + \tau) = e(\frac{\tau}{2} + x) \times \sum_{(m+1/2)(n+1/2) > 0} \text{sign}(m + 1/2) e\left(\frac{\tau}{2} (2(m + 1)^2 + 4(m + 1)(n - 1) + (n - 1)^2) + (n - 1)x\right) = e(\frac{\tau}{2} + x) (h_0(x) - \theta(x, \tau)) + \theta(0, 2\tau).$$

Let us denote $\psi(x) = \theta(x - (\tau + 1)/2, \tau)h_0(x)$. Then ψ satisfies the equation

$$\psi(x + \tau) = e(\frac{\tau + 1}{2})\psi(x) + e(\frac{\tau}{2})\theta(x, \tau)\theta(x - \frac{\tau + 1}{2}, \tau) + \theta(0, 2\tau)\theta(x + \frac{\tau + 1}{2}, \tau).$$

It is easy to see that this equation has the unique holomorphic solution, namely,

$$\psi(x) = \theta(\frac{\tau + 1}{2}, 2\tau) \cdot \left(e(\frac{\tau}{2})\kappa(\frac{\tau + 1}{2}, 2x - \frac{\tau + 1}{2}; 2\tau) - e(x - \frac{\tau}{2})\kappa(-\frac{\tau + 1}{2}, 2x + \frac{\tau + 1}{2}; 2\tau) \right) + \theta(0, 2\tau)\kappa(\frac{\tau + 1}{2}, x + \frac{\tau + 1}{2}; \tau).$$

3.6. General case. It is easy to see that for general four lines the triple Fukaya products are expressed via holomorphic functions of the following sort

$$F_B(x, y; \tau) = \sum_{(m+1/2)(n+1/2) > 0} \text{sign}(m + \frac{1}{2}) e(B(m, n)\tau + \Delta_1 B(m, n)x + \Delta_2 B(m, n)y)$$

where $B(m, n) = am^2 + bmn + cn^2$ is a quadratic form with rational coefficients, which is positive in the range of summation, $\Delta_1 B(m, n) = 2am + bn + a$ and $\Delta_2 B(m, n) = 2cn + bm + c$ are its partial difference derivatives. This function satisfies the following difference equations:

$$F_B(x + \tau, y; \tau) = e(-2ax - by)F_B(x, y) - e(-ax + (c - b)y)\theta(bx + 2cy, 2c\tau)$$

$$F_B(x, y + \tau; \tau) = e(-2cy - bx)F_B(x, y) - e(-cy + (a - b)x)\theta(by + 2cx, 2a\tau)$$

plus $F_B(x + N, y, \tau) = F_B(x, y + N, \tau)$ for sufficiently divisible N . From this equations it seems plausible that the function F_B can be expressed via the functions κ and θ . Also the A_∞ -axioms should produce analogues of the identities (3.4.3) and (3.5.4) involving F_B . We leave these subjects for future investigation.

REFERENCES

- [1] M. P. Appell, *Sur le fonctions doublement periodique de troisieme espece*, Annales scientifiques de l'École Normale Supérieure, 3e série, t.I, p.135, t.II, p.9, t.III, p.9 (1884–1886).

- [2] G. Gasper, M. Rahman, *Basic hypergeometric series*, Cambridge University Press, 1990.
- [3] S. Gelfand, Yu. Manin, *Methods of homological algebra*. Springer-Verlag, 1996.
- [4] K. Fukaya, *Morse Homotopy, A^∞ -Category, and Floer Homologies*, in *The Proceedings of the 1993 GARC Workshop on Geometry and Topology*, H. J. Kim, ed., Seoul National University.
- [5] G.-H. Halphen, *Traité des fonctions elliptiques*, I. Paris, 1886.
- [6] M. Kontsevich, *Homological algebra of mirror symmetry*, Proceedings of ICM (Zürich, 1994), 120–139. Birkhäuser, Basel, 1995.
- [7] L. Kronecker, *Zur theorie der elliptischen functionen* (1881), in *Leopold Kronecker's Werke*, vol. IV, 311–318. Chelsea Pub. Co., 1968.
- [8] A. Polishchuk, E. Zaslow, *Categorical mirror symmetry: the elliptic curve*, preprint math.AG/9801119.
- [9] A. Weil, *Elliptic functions according to Eisenstein and Kronecker*. Springer-Verlag, 1976.