

ON THE REPRESENTATIONS OF THE INFINITE SYMMETRIC GROUP

ANDREI OKOUNKOV

ABSTRACT. We classify all irreducible admissible representations of three Olshanski pairs connected to the infinite symmetric group $S(\infty)$. In particular, our methods yield two simple proofs of the classical Thoma's description of the characters of $S(\infty)$. Also, we discuss a certain operation called mixture of representations which provides a uniform construction of all irreducible admissible representations.

CONTENTS

- 0. Introduction
 - 0.1 Tame representations and factor-representations
 - 0.2 Olshanski pairs and admissible representations
 - 0.3 The statement of the problem and of the results
- 1. Olshanski semigroups
 - 1.1 Definition and Olshanski's theorem
 - 1.2 Parameterization of representations
 - 1.3 An example: Thoma multiplicativity
- 2. Classification of irreducible admissible representations
 - 2.1 Spectra of the operators A_i in spherical representations of the pair (G^D, K^D)
 - 2.2 Spectra of the operators A_i in admissible representations of the pair (G^D, K^D)
 - 2.3 The Thoma theorem
 - 2.4 Another proof of Thoma theorem
 - 2.5 Classification of the irreducible admissible representations of the pair (G^D, K^D)
 - 2.6 Description of $K^E \backslash G^E / K^E$
 - 2.7 Spherical representations of the pair (G^E, K^E)
 - 2.8 Classification of irreducible admissible representations of the pairs (G^O, K^O) , (G^E, K^E)
- 3. Construction of representations
 - 3.1 Mixtures of representations
 - 3.2 Mixtures and induction
 - 3.3 Elementary representations.
 - 3.4 Mixtures in the case of (G^D, K^D)
- 4. Concluding remarks

This is the English version of the author's PhD thesis (1995, Moscow State University).

0. INTRODUCTION

0.1 Tame representations and factor-representations.

Denote by $S(n)$ the group of permutations of the set $\{1, \dots, n\}$ and let

$$S(\infty) = \bigcup_n S(n)$$

be the union of groups $S(n)$ over all n . The group $S(\infty)$ is one of the simplest examples of a *wild* group (see, for example, the book [23] for the definitions of tame and wild groups, as well as for other basic notions of infinite-dimensional representation theory). This means that the study of all irreducible representations of $S(\infty)$ does not seem to be a reasonable problem.

There is probably a unique natural simple topology on the group $S(\infty)$ induced by the weak (as well as by the strong) operator topology in the representation by permutations of basis vectors

$$(0.1) \quad S(\infty) \rightarrow U(l_2),$$

where $U(l_2)$ denotes the group of unitary operators in the coordinate Hilbert space l_2 . Denote by

$$S_n(\infty), \quad n = 1, 2, 3, \dots,$$

the subgroup of $S(\infty)$ which fixes the numbers $1, 2, \dots, n$. The subgroups $S_n(\infty)$ form a fundamental neighborhood base of identity in this topology. A unitary representation

$$(0.2) \quad \pi : S(\infty) \rightarrow U(H),$$

where $U(H)$ is the group of unitary operators in a Hilbert space H , is called *tame* if it is continuous with respect to this topology on $S(\infty)$ and the weak topology on $U(H)$. All tame representations were described by Lieberman [27] (see also [39]). In particular, it is known that:

- (1) any tame representation is a direct sum of irreducible ones;
- (2) irreducible tame representations are labeled by all Young diagrams μ ;
- (3) the k th tensor power of the representation (0.1) decomposes into irreducible representations labeled by all Young diagrams μ such that $k \geq |\mu| > 0$.

These properties make the topological group $S(\infty)$ look similar to a compact group. One expects the group $S(\infty)$ to be a good combinatorial model of a big infinite-dimensional group. It is clear that the supply of tame representations is much too small for a unitary dual of anything truly infinite-dimensional.

Denote by $\overline{S(\infty)}$ the group of all bijections of the set of natural numbers. One can introduce a similar topology on this group and in this topology $S(\infty)$ is a dense subgroup. Tame representations are precisely those representations of the group $S(\infty)$ that can be extended by continuity to the entire group $\overline{S(\infty)}$.

We conclude the discussion of tame representations by one more technically useful definition of a tame representation. Denote by

$$H_n = H^{S_n(\infty)}, \quad n = 1, 2, \dots,$$

the subspace of invariants for the action of the group $S_n(\infty)$ in the representation (0.2). As shown in [39], the representation (0.2) is tame if and only if

$$H = \overline{\bigcup_n H_n}.$$

Another approach to the representation theory of the group $S(\infty)$ focuses on the study of its factor-representations of finite type (or, more generally, of semi-infinite type). A beautiful theory of these representations was developed by Thoma [49], and Vershik and Kerov [6,7,9,10].

Any finite type factor representation π is uniquely determined, up to quasi-equivalence, by its trace, that is, the restriction of the trace in the factor on the image of the group. The traces of finite type factor-representations of the group G are precisely the *characters* of the group G . A character of the the group G is, by definition, a function ϕ on the group G which is

- (1) central, that is, $\phi(g_1 g_2) = \phi(g_2 g_1)$ for all $g_1, g_2 \in G$;
- (2) positive definite, that is, for all $g_1, \dots, g_n \in G$ the matrix $(\phi(g_i g_j^{-1}))$ is Hermitian and non-negatively definite;
- (3) indecomposable, that is, it cannot be represented as a sum of two linearly independent functions satisfying (1) and (2);
- (4) normalized by $\phi(e) = 1$.

If the group G is compact then its characters are precisely the functions

$$\phi_\pi(g) = \frac{\text{tr } \pi(g)}{\dim \pi},$$

where π runs over the set of equivalence classes of irreducible representations of the group G . The character theory of compact groups is a classical chapter of representation theory. The characters of wild groups were the subject of intense recent studies, see, for example, [3, 6-11, 13-15, 25, 44, 49-52].

In the paper [49], Thoma obtained the following description of all characters of the group $S(\infty)$. The characters of the group $S(\infty)$ are labeled by a pair of sequences of real numbers $\{\alpha_i\}$, $\{\beta_i\}$, $i = 1, 2, \dots$, such that

$$\alpha_1 \geq \alpha_2 \geq \alpha_3 \geq \dots > 0, \quad \beta_1 \geq \beta_2 \geq \beta_3 \geq \dots > 0, \\ \sum \alpha_i + \sum \beta_i \leq 1.$$

The value of the corresponding character on a permutation with a single cycle of length k is

$$\sum_i \alpha_i^k + (-1)^{k-1} \sum_i \beta_i^k.$$

Its value on a permutation with several disjoint cycles equals the product of the values on each cycle. As usual, it is assumed that an empty product equals 1. In particular, the character of the regular representation of the group $S(\infty)$ corresponds to the sequences $\alpha_i \equiv 0$, $\beta_i \equiv 0$.

The heart of the Thoma's proof is the classification of so called *totally positive* sequences. Recall that a sequence of real numbers $\{a_i\}, i = 0, 1, 2, \dots$ is said to be totally positive if all minors of the following infinite Toeplitz matrix

$$\begin{bmatrix} a_0 & a_1 & a_2 & a_3 & \dots \\ 0 & a_0 & a_1 & a_2 & \ddots \\ & 0 & a_0 & a_1 & \ddots \\ & & 0 & a_0 & \ddots \\ & & & \ddots & \ddots \end{bmatrix}$$

are non-negative. Thoma obtained the description of all totally positive sequences using some deep results about entire functions. He established that a totally positive sequence

$$a_0, a_1, a_2, \dots$$

has generating function of the following form

$$\sum_i a_i t^i = e^{\gamma t} \prod_i \frac{1 + \beta_i t}{1 - \alpha_i t}$$

for some non-negative $\{\alpha_i\}, \{\beta_j\}, \gamma$ such that

$$\sum \alpha_i + \sum \beta_i < \infty.$$

We shall briefly explain below the connection between totally positive sequences and representation theory and show how simple representation theoretical considerations allow one to simplify Thoma's original argument significantly and, in particular, to avoid entire functions entirely.

As a matter of fact, Thoma's description of totally positive sequences was found earlier in the papers [1, 17]. Totally positive sequences arise in many problems of analysis (such as approximation theory or small oscillations), geometry (convex curves), and probability. An important role is played, for instance, by the following characteristic property of totally positive sequences: the convolution of an arbitrary sequence with a totally positive sequence contains no more changes of sign than the original sequence. This is a classical result by I. Schoenberg [43] which he obtained in the course of his studies of various generalizations of the Descartes rule. Totally positive sequences and their continuous analogs were studied by F. R. Gantmakher, M. G. Krein, D. Polya, I. Schoenberg and his collaborators, A. Edrei, S. Karlin, and others. See, for example, [1, 16-18, 21] and especially [42] where one can find references to the most recent applications of the theory of totally positive sequences. Among the papers discussing the relations between total positivity and representation theory we mention [6-11, 12].

An explicit construction of all corresponding factor-representations of the group $S(\infty)$ was given later by A. M. Vershik and S. V. Kerov [6]. A. M. Vershik and S. V. Kerov have also found another proof of Thoma theorem based on the so called *ergodic method*; see [5,7] and also [57].

For the group $S(\infty)$, the general ergodic method specializes to the following procedure. One starts with a sequence λ_n of Young diagrams such that $|\lambda_n| = n$. Let

$$\chi_n(g) = \frac{\text{tr } \pi_{\lambda_n}(g)}{\dim \pi_{\lambda_n}}$$

be the normalized character of the symmetric group $S(n)$ corresponding to the irreducible representation π_{λ_n} .

We say that a sequence of characters χ_n converges, as $n \rightarrow \infty$, to a function χ on the group $S(\infty)$ if

$$\chi_n(g) \rightarrow \chi(g), \quad n \rightarrow \infty$$

for every element $g \in S(\infty)$. Note that the expression $\chi_n(g)$ makes sense for all sufficiently large n .

It follows from the general approximation theorems [5, 12, 33] that every character χ of the infinite symmetric group $S(\infty)$ is a limit of a suitable sequence of characters χ_n of finite symmetric groups. A. M. Vershik and S. V. Kerov proved that a sequence χ_n has a limit if and only if the following limits exist:

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{i\text{-th row of } \lambda_n}{n} &= \alpha_i, \\ \lim_{n \rightarrow \infty} \frac{i\text{-th column of } \lambda_n}{n} &= \beta_i. \end{aligned}$$

If these limits exist then the characters χ_n converge to the Thoma character with parameters $\{\alpha_i\}, \{\beta_i\}$. In other words, the parameters $\{\alpha_i\}, \{\beta_i\}$ have the meaning of asymptotic lengths of rows and columns of a Young diagram.

0.2 Olshanski pairs and admissible representations.

In [32], G. Olshanski initiated the study of a more general class of representations of the infinite symmetric group $S(\infty)$. Before giving a definition, let us consider an example. Let

$$\pi : S(\infty) \rightarrow U(M)$$

be a finite type factor-representation of the group $S(\infty)$. Here $U(M)$ is the group of unitary operators in a finite factor M . Let the Hilbert space H be the completion of M with respect to the following Hermitian inner product

$$(A, B) = \text{tr } AB^*.$$

The group

$$(0.3) \quad G = S(\infty) \times S(\infty),$$

acts in this space by left and right multiplications and this representation is irreducible. The identity operator

$$1 \in M \subset H$$

is the unique vector invariant under the action of the diagonal subgroup

$$(0.4) \quad K = \text{diag } S(\infty) \subset S(\infty) \times S(\infty),$$

and the corresponding matrix element

$$(\pi(g)1, 1) = \text{tr } \pi(g)$$

is exactly the trace of the factor representation π .

Let us check that the irreducibility of the action of the group (0.3) and the existence of a vector invariant under the action of the subgroup (0.4) implies that the action of the subgroup (0.4) in the space H is tame [32]. Indeed, the subgroups

$$S(n) \times S(n) \quad \text{and} \quad K_n = \text{diag } S_n(\infty)$$

commute. Therefore, the subspace

$$1 \in \overline{\bigcup_n H^{K_n}}$$

is G -invariant, and also closed and non-trivial, hence, equal to H .

As this example suggests, one should study unitary representations of the group (0.3) such that their restrictions to the subgroup (0.4) are tame. Such representations are called *admissible representations of the pair* (0.4) or simply *representations of the pair* (0.4).

Olshanski's general idea was that, in the infinite-dimensional situation, it takes two groups to produce a good representation theory. Namely, one should study the unitary representations not of a single group G , but rather unitary representations of a pair

$$(0.5) \quad K \subset G,$$

where K is a subgroup of G designated to play the role of a maximal compact subgroup of G . A unitary representation

$$G \rightarrow U(H)$$

of the group G is said to be a *representation of the pair* (0.5) if its restriction on the subgroup K belongs to a given simple class of representations of the group K (for example, the class of tame representations). Recall that tame representations do resemble in many aspects representations of a compact group. In addition to the pair (0.4), two other pairs closely connected to the group $S(\infty)$ were considered by G. Olshanski in [32] and will be studied in the present paper.

Equivalently, admissible representation can be viewed as continuous unitary representations of a certain not locally-compact group \overline{G} containing G as a dense subgroup, see [32] or the next subsection.

The *spherical* representations which, by definition, are irreducible unitary representations of G with a non-zero K -fixed vector, form a distinguished subclass of admissible representations. It is known (see [32] or below) that in case $\dim H^K = 1$ and also that the above correspondence between finite factor representations of $S(\infty)$ and spherical representations of (G, K) is a bijection.

One of the advantages of the class of admissible representations is that it is closed under all natural operations on representations such as restriction to the subgroup $S_n(\infty) \times S_n(\infty)$, induction from such subgroups, taking direct sums and tensor products.

As it was shown by Olshanski, every representation of the pair (0.4) is of von Neumann type I. It follows that the study of general representations of the pair (0.4) can be essentially reduced to the study of its irreducible representations. Olshanski obtained this result using his so called *semigroup method*, which is a very powerful and beautiful tool for the study of admissible representations [34,32]. This method is a far reaching generalization of the multiplicativity property of characters of $S(\infty)$ found by Thoma (see also [15, 19-20, 31]). It can be compared to the use of Hecke algebras in the p -adic representation theory; the subgroups K_n play the role of the principal congruence subgroups. The main difference between the p -adic groups and the groups like (0.4) is that K_n are not compact. Consequently, the definition of the convolution product for K_n -biinvariant functions on G involves a certain limit transition. On the bright side, in this limit, the multiplication greatly simplifies and one obtains an actual semigroup and not just a hypergroup.

As another application of the semigroup machinery one obtains a way of labeling the irreducible admissible representations. All irreducible admissible representation of the pair (0.4) are indexed by continuous parameters (namely, parameters used in the Thoma theorem) and some discrete parameters. More precisely, to every element of the set $\{\alpha_i\} \cup \{-\beta_j\} \cup \{0\}$ one assigns two Young diagrams in such a way that all but finitely many diagrams are empty. Conversely, any such data correspond, in general, to a representation in a vector space with an invariant sesquilinear scalar product, not necessarily positive definite. The classification problem for irreducible admissible representation this way can be reformulated as finding all values of the parameters which correspond to unitary representation. One can compare this with the problem of describing the unitary highest weight modules of a Lie algebra.

We shall give a brief summary of the semigroup method and the resulting labeling of representations in Section 1.

In the same paper [32], G. Olshanski has constructed a large supply of irreducible admissible representations. His construction generalizes, on the one hand, that of factor-representations of the infinite symmetric group by A. M. Vershik and S. V. Kerov, and, on the other hand, it is an infinite-dimensional generalization of the classical Hermann Weyl's duality for representations in traceless tensors. Using this construction and the ergodic method, G. Olshanski obtained two-sided estimates for the set of parameters corresponding to irreducible admissible representations.

In the present paper we shall prove, and this is our main result, that the lower bound from [32] is actually the correct answer.

0.3 The statement of the problem and of the results.

We study irreducible representations of three similar Olshanski pairs related to the infinite symmetric group.

By definition, a bijection $g : \mathbb{Z} \rightarrow \mathbb{Z}$ is called *finite* if the set

$$\{i \in \mathbb{Z} \mid g(i) \neq i\}$$

is finite. Define a group G^O as the group of all finite bijections $\mathbb{Z} \rightarrow \mathbb{Z}$. Set

$$G^E = \{g \in G^O \mid g(0) = 0\}$$

$$G^D = \{g \in G^O \mid g(0) = 0, g(\mathbb{N}) = \mathbb{N}\}$$

The superscripts O , E , D mean “odd”, “even” and “double”. We write simply G if the formulas are valid for all three groups. Set

$$G(n) = \{g \in G \mid g(i) = i, |i| > n\}$$

$$G_n = \{g \in G \mid g(i) = i, |i| \leq n\}$$

The group G is the union of an increasing chain of its subgroups $G(n)$

$$\{e\} = G(0) \subset G(1) \subset G(2) \subset \dots$$

$$\bigcup_n G(n) = G.$$

The subgroups G_n form a decreasing chain of subgroups,

$$G_0 \supset G_1 \supset G_2 \supset \dots,$$

$$\bigcap_n G_n = \{e\}.$$

The subgroups $G(n)$ and G_n commute.

There is a natural involution $i \mapsto -i$ on the set \mathbb{Z} . Using this involution, one can define a subgroup of the group G which shall play the role of a maximal compact subgroup. Let

$$K = \{g \in G \mid g(-i) = -g(i)\}.$$

The pairs (G, K) are Gelfand pairs in the sense of [32]. Define an increasing and decreasing chains of subgroups

$$K(n) = K \cap G(n),$$

$$K_n = K \cap G_n$$

in the subgroup K .

The group G^D is isomorphic to a direct product of two copies of the group $S(\infty)$ (one permutes the positive numbers, the other — negative ones). The group K^D is isomorphic to $S(\infty)$.

The groups G^O and G^E are isomorphic to $S(\infty)$, and the subgroups K^O , K^E are isomorphic to a semi-direct product of $S(\infty)$ and the group \mathbb{Z}_2^∞ . In this case the Olshanski pairs differ by the way of embedding of the subgroup into the group (and the resulting representation theory is also different).

We have $G_n^D \cong G^D$ and $G_n^E \cong G_n^O \cong G^E$. Also $G^D(n) \cong S(n) \times S(n)$, $G^E(n) \cong S(2n)$, and $G^O(n) \cong S(2n+1)$.

Let π be a unitary representation of the group G in a Hilbert space $H(\pi)$. Denote by $H(\pi)_n$ the subspace formed by K_n -invariant vectors. This subspace is invariant under the action of the group $G(n)$. Furthermore, the subspaces $H(\pi)_n$ form an increasing sequence of subspaces

$$H(\pi)_0 \subset H(\pi)_1 \subset H(\pi)_2 \dots$$

Hence, their union $\cup_n H(\pi)_n$ is an algebraically invariant subspace in $H(\pi)$. A representation π is called (see [32]) *admissible representation of the pair (G, K)* , if

$$H(\pi) = \overline{\bigcup_n H(\pi)_n}.$$

It is known [32] that any admissible representation is of von Neumann type I.

The notion of admissible representation can also be stated in topological terms. Namely, as shown in [32], for every one of the three pairs (G, K) there exists some topological group \overline{G} which is not locally compact and contains G as dense subgroup. The admissible representations of the pair (G, K) are exactly unitary representations of G admitting an extension by continuity to the group \overline{G} . As a result, the theory of admissible representations is equivalent to the theory of unitary representations of non locally-compact groups \overline{G} . Since every admissible representation generates a von Neumann algebra of type I, non locally-compact topological groups \overline{G} are actually tame groups.

The construction of the group \overline{G} is as follows. Embed the group G into the group of all (not necessarily finite) bijections

$$g : \mathbb{Z} \rightarrow \mathbb{Z}.$$

Consider subgroups

$$\overline{K}_n = \{g | g(i) = -g(-i), g(i) = i, i > n\} \cap G, \quad n = 0, 1, \dots$$

and let

$$\overline{G} = G \cdot \overline{K}.$$

Finally, define a topology on the group \overline{G} such that a fundamental neighborhood system of unity is formed by the subgroups \overline{K}_n , $n = 0, 1, \dots$.

The viewpoint of unitary representations of the groups \overline{G} is convenient for the constructions of representations, see Section 3. Still, the viewpoint of admissible representations of (G, K) pairs is more convenient for the proof of classification theorems.

The **main result** of this paper is a complete description of all irreducible admissible representations of the three pairs (G, K) given for the pair (G^D, K^D) in Theorem 3, Section 2.5, and for the other two pairs in Theorem 5, Section 2.8.

The sufficiency of the conditions of given in these theorems was known before; it follows from Olshanski's explicit construction of irreducible admissible representations [32]. We devote the entire Section 3 to the discussion of the construction of admissible representations. However, in that section, our improvement upon [32] is

only a very modest one. It follows from our classification theorems that the representations constructed in [32] form actually an open subset of the admissible dual. Here we introduce the notion of a *mixture* of admissible representations which gives a uniform construction of all irreducible admissible representation. This mixture is a kind of an induced representation as discussed in Section 3.2.

Another known result which we discuss at length in the present paper is the Thoma's description of characters of $S(\infty)$. Traditionally, this result was considered as a very hard one. Our methods allow to give an new simple proof (see Section 2.3) and also to very much simplify Thoma's original proof (see Section 2.4).

The analog of Thoma theorem for the pair (G^E, K^E) is obtained in Theorem 4, Section 2.7.

A brief account of the most important results of this paper was published in [55].

1. OLSHANSKI SEMIGROUPS

1.1 Definition and Olshanski's theorem.

In this section we describe semigroups which shall play a prominent role in the sequel. These semigroups were introduced by G. Olshanski in [32]; they are similar to the Brauer semigroups. A more detailed exposition can be found in [32, 34].

Given two disjoint finite sets S and S' , we consider the following set. An element of $B(S, S')$ is by definition the following data


- (1) a partition of the set $S \cup S'$ into pairs together with a nonnegative real number assigned to each pair, and
- (2) a finite unordered collection of nonnegative real numbers.

Geometrically, this data can be visualized as a compact 1-dimensional manifold M such that

- (1) the boundary ∂M of M is $S \cup S'$, and
- (2) each connected component of M is equipped with a non-negative real number (which can be thought of as its length).

The connected components with boundaries give a partition of $S \cup S'$ into pairs and the lengths of the cycles of M (by which we mean the loops in M , that is, connected components homeomorphic to S^1) give an array of nonnegative numbers.


It is convenient to position the elements of S and S' on two horizontal lines, one above the other. An example of an element of $B(S, S')$ with $|S| = |S'| = 5$ is depicted in the following figure:



There is a natural map

$$B(S, S') \times B(S', S'') \rightarrow B(S, S'')$$

which glues a manifold $M_1 \in B(S, S')$ to a manifold $M_2 \in B(S', S'')$ along S' . The lengths of the two glued components, naturally, add up. Note that the resulting manifold may have more cycles than M_1 and M_2 combined. This operation makes $B(S, S)$, where the two S 's are considered as two disjoint copies of the same set, a noncommutative semigroup. For example, if $|S| = 5$, then



In more technical terms, one can think of an element of $B(S, S')$ as of a *wiring diagram* with inputs/outputs indexed by $S \cup S'$ and a nonnegative resistance (instead of length) assigned to each wire. When one multiplies (that is, connects) two such objects, the resistances of the connected wires add up. We shall occasionally refer to the elements of $B(S, S')$ as (wiring) diagrams. By analogy with a computer chip, the term *chip* was used in [22,32].

The group $\text{Aut}(S)$ is naturally a subgroup of

$$\text{Aut}(S) \subset B(S, S)$$


formed by all diagrams with no cycles and all other components of length zero. Namely, a bijection

$$g : S \rightarrow S$$

corresponds to such diagram that each element $s \in S$ in the first copy of S is connected to the element $g^{-1}(s)$ in the second copy of S by a segment of length zero. Recall that we consider the length as just a formal number assigned to a each connected component; in particular, components of length zero are still non-trivial.

Our next goal is to make some sense out of the object " $B(\mathbb{Z}, \mathbb{Z})$ ". The above multiplication rule can fail for infinite diagrams because one can get infinitely many loops, which is what we want to avoid.


However, we shall need only the semigroup generated by a certain special set of infinite diagrams; in that semigroup the multiplication will be indeed well defined. First, we take all diagrams with no cycles and no components of positive length. They form a group isomorphic to G^O . We add to them the following diagrams A_k , $k \in \mathbb{Z}$. The diagram A_k is defined in the following figure:



All but one segments of A_k have zero length and the only segment of length 1 joins $k \in \mathbb{Z}$ from the first copy of \mathbb{Z} with k in the second copy. Clearly, the semigroup

$$\langle G^O, A_1 \rangle \cong S(\infty) \ltimes \mathbb{Z}_{\geq 0}^\infty$$


generated by G^O and A_1 contains also all other A 's. Now, let C_k , $k = 1, 2, 3, \dots$, be following pure cycles. The diagram C_k is, by definition,



In our semigroup, we wish to mod out by the relation

$$C_1 = 1.$$

The purpose of doing this is to make the following diagrams P_k , $k = 0, 1, 2, \dots$, into idempotents. The diagram P_k is depicted in the following picture:



It has $2k + 1$ vertical segments of length 0 and all the remaining components are arcs of length 1/2. Clearly, $P_k^2 = P_k \pmod{C_1}$.

Definition. Denote by Γ^O the semigroup generated by the following diagrams

$$\Gamma^O = \left\langle G^O, A_1, C_j, P_k \right\rangle_{j \geq 1, k \geq 0} \Big/ \left\langle C_1 = 1 \right\rangle$$

modulo the relation $C_1 = 1$. Similarly, set

$$\Gamma^E = \left\langle G^E, A_1, C_j, P_k \right\rangle_{j \geq 1, k \geq 0} \Big/ \left\langle C_1 = 1 \right\rangle.$$

$$\Gamma^D = \left\langle G^D, A_1, C_j, P_k \right\rangle_{j \geq 1, k \geq 0} \Big/ \left\langle C_1 = 1 \right\rangle.$$

We call these semigroups the *Olshanski semigroups*. They are slightly smaller than the ones defined by Olshanski in [32], Sections 2.8 and 3.10. Those defined in [32] are topological semigroups which contain ours as dense subsemigroups.

There is a natural involution $*$ in these semigroups, namely, the reflection in the horizontal axis. This involution fixes all A 's, C 's, and P 's and takes a permutation g to the inverse permutation $g^* = g^{-1}$.

The importance of Olshanski semigroups for the theory of admissible representations lies in the following fundamental

Theorem (Olshanski, [32]). *Every admissible representation π of a pair (G, K) extends canonically to a $*$ -representation by contractions (that is, operators of norm ≤ 1) of the corresponding semigroup Γ in $H(\pi)$. In this representation, the idempotent P_n maps to the orthogonal projection onto $H(\pi)_n$.*

This canonical extension we shall denote with the same letter π .

Let us say a few words about how this canonical extension is constructed. We have to specify the action of the A 's and C 's. For simplicity, consider the pair (G^O, K^O) ; the other pairs are very similar. Consider the transposition $(i, n) \in G^O$ and consider the limit of the operators

$$\lim_{n \rightarrow \infty} \pi((i, n)),$$

which exists in the weak operator topology. To see this, it suffices to check that the limits

$$\lim_{n \rightarrow \infty} (\pi((i, n))\zeta, \eta)$$

do exist, where the vectors ζ and η belong to the dense subspace

$$\bigcup_{m=1}^{\infty} H(\pi)_m.$$

Let ζ and η belong to the subspace $H(\pi)_m$ and assume that the numbers n_1, n_2 are chosen big enough, $n_1, n_2 > m$. Then the permutation

$$(n_1, n_2)(-n_1, -n_2) \in K_m$$

belongs to the subgroup K_m and, by the definition of the subspace $H(\pi)_m$, we have

$$\begin{aligned} (\pi((i, n_1))\zeta, \eta) &= \\ &= (\pi((n_1, n_2)(-n_1, -n_2)(i, n_2)(n_1, n_2)(-n_1, -n_2))\zeta, \eta) = \\ &= (\pi((i, n_2))\pi((n_1, n_2)(-n_1, -n_2))\zeta, \pi((n_1, n_2)(-n_1, -n_2))\eta) = \\ &= (\pi((i, n_2))\zeta, \eta). \end{aligned}$$

Therefore, the number

$$(\pi((i, n))\zeta, \eta)$$

does not depend on n provided that $n > m$, hence coincides with the limit

$$(\pi((i, n))\zeta, \eta) = \lim_{n \rightarrow \infty} (\pi((i, n))\zeta, \eta), \quad n > m.$$

By definition, one sets

$$\pi(A_i) = \lim_{n \rightarrow \infty} \pi((i, n)),$$

and, similarly,

$$\pi(C_k) = \lim_{n_1, \dots, n_k \rightarrow \infty} \pi((n_1, n_2, \dots, n_k)),$$

where we assume that the numbers n_1, \dots, n_k are pairwise distinct.

Sometimes, it is convenient to replace the limits in the definition of the operators $\pi(A_i)$, $\pi(C_i)$ by the corresponding Cesaro limits

$$\pi(A_i) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \pi((i, j)),$$

which exist in the strong operator topology. This formula be interpreted as saying that A_i is the transposition of i and a “random” number j . The cycle C_k can be thought of as a “random” cyclic permutation of length k .

The operator $\pi(P_m)$ may be represented in a similar form

$$\pi(P_m) = \lim_{n \rightarrow \infty} \frac{1}{n!} \sum_{g \in K_m(n)} \pi(g),$$

where $K_m(n) = K_m \cap K(n)$. Indeed, the operator

$$\frac{1}{n!} \sum_{g \in K_m(n)} \pi(g)$$

is the projection onto the subspace of $K_m(n)$ -invariants. Denote this subspace by $H(\pi)_{m;n}$. Clearly,

$$H(\pi)_{m;0} \supset H(\pi)_{m;1} \supset H(\pi)_{m;2} \supset \dots$$

and

$$H(\pi)_m = \bigcap_n H(\pi)_{m;n}.$$

Hence, in the strong operator topology, the projection onto the subspace $H(\pi)_m$ is the limit of those onto the subspaces $H(\pi)_{m;n}$.

For the convenience of the future references we list some useful identities. Given a permutation σ , denote by $[\sigma]$ the array of numbers formed by the lengths of non-trivial cycles of σ . One easily checks the following equalities:

Proposition 1.

$$(1.1) \quad A_i A_j = A_j A_i,$$

$$(1.2) \quad g A_i g^{-1} = A_{g(i)}, \quad g \in G^O,$$

$$(1.3) \quad A_i P_n = P_n A_i, \quad |i| \leq n,$$


$$(1.4) \quad A_i P_n = A_{-i} P_n, \quad |i| > n,$$

$$(1.5) \quad P_n A_i P_n = P_n(i, k) P_n, \quad |i| \leq n, |k| > n,$$

$$(1.6) \quad P_n A_{i_1}^{k_1} A_{i_2}^{k_2} \dots A_{i_r}^{k_r} P_n = P_n \prod_{j=1}^r C_{k_j+1}, \quad n < i_j, i_m \neq i_l,$$

$$(1.7) \quad P_0 \sigma P_0 = P_0 \prod_{k \in [\sigma]} C_k, \quad \sigma \in S(\infty).$$

All these identities are straightforward to check. As a visual aid for the proof of the last equality we provide the following figure (where $P_0(123)P_0 = P_0 C_3$ is depicted):

**1.2 Parameterization of representations.**

Let π be an irreducible admissible representation of a pair (G, K) in a Hilbert space $H(\pi)$. Let $d = d(\pi)$ denote the least integer k , such that $H(\pi)_k \neq 0$; it is called the *depth* of π . Denote the subspace $H(\pi)_d$ by $R(\pi)$ and call it the *root* of π .

Let ζ be a vector in the subspace $R(\pi)$. The representation π is uniquely determined by any matrix element, in particular by the one corresponding to ζ :

$$\psi(g) = (\pi(g)\zeta, \zeta).$$

The operator $\pi(P_d)$ is the orthogonal projection onto the subspace $R(\pi)$, hence

$$(1.8) \quad (\pi(g)\zeta, \zeta) = (\pi(P_d g P_d)\zeta, \zeta).$$

The set

$$\Gamma(d) = P_d \Gamma P_d \subset \Gamma$$

is a subsemigroup which acts in $R(\pi)$. The following proposition is due to G. Olshanski:

Proposition 2. *Let π be an admissible representation of the group G . Choose n so that $H(\pi)_n \neq 0$. Denote by π_n the representation of the semigroup $\Gamma(n)$ in the subspace $H(\pi)_n$. Then*

- (1) *if the representation π is irreducible, then the representation π_n is irreducible;*
- (2) *if the representation π_n is irreducible and the subspace $H(\pi)_n$ is cyclic then π is also irreducible.*

Proof. Assume that π is irreducible. Let B denote an arbitrary bounded operator in the subspace $H(\pi)_n$. Denote by \tilde{B} the operator in the subspace $H(\pi)$ which coincides with B on the subspace $H(\pi)_n$ and equals zero on its orthogonal complement $H(\pi)_n^\perp$. Clearly, this operator is bounded. Since π is irreducible, there exists a sequence $b^{(i)}$ of elements in the group algebra $\mathbb{C}[G]$, such that

$$\pi(b^{(i)}) \rightarrow \tilde{B}, \quad i \rightarrow \infty$$

in the weak operator topology. But this implies that

$$\pi(P_n b^{(i)} P_n) \rightarrow \pi(P_n) \tilde{B} \pi(P_n), \quad i \rightarrow \infty.$$

Hence,

$$\pi_n(P_n b^{(i)} P_n) \rightarrow B, \quad i \rightarrow \infty.$$

Therefore, the representation π_n is irreducible.

In the opposite direction, let us argue by contradiction. Let W be a non-trivial closed invariant subspace. For any $\zeta \in H(\pi)_n$, its orthogonal projections onto W and W^\perp are also K_n -invariant vectors. Therefore, at least one of $\Gamma(n)$ -invariant subspaces

$$H(\pi)_n \cap W \neq 0, \quad \text{or} \quad H(\pi)_n \cap W^\perp \neq 0$$

is non-trivial. Since π_n is irreducible, we conclude that

$$H(\pi)_n \subset W, \quad \text{or} \quad H(\pi)_n \subset W^\perp.$$

Since the subspace $H(\pi)_n$ is cyclic,

$$H(\pi) = W, \quad \text{or} \quad H(\pi) = W^\perp,$$

which contradicts the non-triviality of W . \square

By virtue of (1.8), we need only to know the representation π_d of the subgroup $\Gamma(d)$ in the subspace $R(\pi)$ in order to reconstruct the representation π .

By definition of the number d , we have $H(\pi)_{d-1} = 0$. Hence,

$$\pi(P_{d-1}) = 0.$$

Set

$$\Gamma(d)^\times = \Gamma(d) \setminus \Gamma(d) P_{d-1} \Gamma(d).$$

One can check [32] that $\Gamma(d)^\times$ is the subsemigroup of $\Gamma(d)$ generated by the elements

$$\begin{aligned} gP_d, \quad g &\in G(d), \\ A_iP_d, \quad |i| &\leq d, \\ C_kP_d, \quad k &\geq 2. \end{aligned}$$

It is clear that only the elements of this subsemigroups can act in $H(\pi)_d$ by non-zero operators.

The semigroup $\Gamma(d)^\times$ is very simple and its representations can be easily described. In case of Γ° , it is isomorphic to

$$\Gamma^\circ(d)^\times \cong (S(2d+1) \ltimes \mathbb{Z}_+^{2d+1}) \times \mathbb{Z}_+^\infty.$$

Here the three factors correspond to the three sets of generators of $\Gamma^\circ(d)^\times$. To simplify notation, set

$$\Sigma(m) = S(m) \ltimes \mathbb{Z}_+^m.$$

All irreducible $*$ -representations of $\Sigma(m)$ by contraction operators are the following.

Given a point $x \in [-1, 1]$ and a partition λ of m , denote by $V_{\lambda,x}$ the $\Sigma(m)$ -module in which \mathbb{Z}_+^m acts by multiplication by x and $S(m)$ acts by the irreducible representation corresponding to λ . This is an irreducible $*$ -representation. More general representation will be induced from the subsemigroups of the form


$$\Sigma(\rho) = \prod_i \Sigma(\rho_i),$$

where ρ is a partition of m and the the product is direct.

By definition, a *Young distribution* $\Lambda(x)$ is a function from $[-1, 1]$ to the set of Young diagrams such that $\Lambda(x) = \emptyset$ for all but finitely many x . We set

$$\begin{aligned} |\Lambda| &= \sum_x |\Lambda(x)|, \\ \text{supp } \Lambda &= \{x, \Lambda(x) \neq \emptyset\}. \end{aligned}$$

One can visualize a Young distribution as a collection of Young diagrams growing out of various points of the interval $[-1, 1]$:



By ordering the numbers $|\Lambda(x)|$, $x \in \text{supp } \Lambda$, one obtains a partition of $|\Lambda|$ which we denote by $\rho(\Lambda)$. If $|\Lambda| = m$ then

$$\bigotimes_{x \in \text{supp } \Lambda} V_{\Lambda(x), x}$$

is an irreducible module over the semigroup $\Sigma(\rho(\Lambda)) \subset \Sigma(m)$. Denote by T_Λ the representation of $\Sigma(m)$ in the following induced module

$$V_\Lambda = C[\Sigma(m)] \otimes_{C[\Sigma(\rho(\Lambda))]} \left(\bigotimes_{x \in \text{supp } \Lambda} V_{\Lambda(x), x} \right).$$

This is an irreducible $*$ -representation. All irreducible $*$ -representations of $\Sigma(m)$ by contractions are precisely the representations in V_Λ .

The C 's are central and $*$ -stable, therefore

$$\pi(C_i) = c_i,$$

for some numbers $c_i \in [-1, 1]$.

Similarly, we have

$$\begin{aligned} \Gamma^E(d)^\times &\cong (S(2d) \ltimes \mathbb{Z}_+^{2d}) \times \mathbb{Z}_+^\infty, \\ \Gamma^D(d)^\times &\cong (S(d) \ltimes \mathbb{Z}_+^d) \times (S(d) \ltimes \mathbb{Z}_+^d) \times \mathbb{Z}_+^\infty. \end{aligned}$$

The representations $\Gamma^E(d)^\times$ are labeled by a Young distribution Λ such that $|\Lambda| = 2d$ and by a sequence $\{c_i\} \subset [-1, 1]$. The representations of $\Gamma^D(d)^\times$ are indexed by a pair of Young distributions Λ, M such that $|\Lambda| = |M| = d$ and numbers c_i , $i = 2, 3, \dots$

The classification problem of admissible irreducible representations of the group G can now be reformulated as the problem of description of all parameters $\Lambda, \{c_i\}$ (respectively, $\Lambda, M, \{c_i\}$) such that an admissible irreducible representation with these parameters does exist. Equivalently, one has to describe all parameters for which the matrix element (1.8) is a positive definite function on the group G .

1.3 An example: Thoma multiplicativity.

Let π be a spherical representation of the pair (G^D, K^D) , which is, by definition, an irreducible unitary representation of G^D such that $H(\pi)_0 \neq 0$. By Olshanski's theorem, the subspace $H(\pi)_0$ carries an irreducible $*$ -representation of the semigroup $\Gamma^D(0)$. This semigroup consists of the elements of the form

$$\prod_i C_i^{k_i} P_0.$$

It is commutative, and has, therefore, only 1-dimensional irreducible representations. Whence,

$$\dim H(\pi)_0 = 1.$$

In other words, there exists a unique, up to a scalar factor, K^D -invariant vector ξ . We normalize it by the condition $\|\xi\| = 1$ and call the *spherical* vector. Such a vector is determined up to multiplication by a complex number of absolute value 1. The spherical function

$$\phi_\pi(g) = (\pi(g)\xi, \xi)$$

does not depend on the choice of the spherical vector.

Since the spherical vector ξ is K^D -invariant, the spherical function is constant on the double cosets $K^D \backslash G^D / K^D$:

$$\phi_\pi(k_1 g k_2) = (\pi(k_1 g k_2)\xi, \xi) = (\pi(g)\pi(k_2)\xi, \pi(k_1^{-1})\xi) = (\pi(g)\xi, \xi) = \phi_\pi(g),$$

where k_1 and k_2 are two arbitrary permutations in K^D .

Every double coset in $K^D \backslash G^D / K^D$ intersects with the subgroup $S(\infty)$ by a conjugacy class in $S(\infty)$. If $\sigma \in S(\infty)$ and σ has just one non-trivial cycle of length k then

$$\phi_\pi(\sigma) = (\pi(\sigma)\xi, \xi) = (\pi(P_0 \sigma P_0)\xi, \xi) \stackrel{(1.7)}{=} (\pi(C_k)\xi, \xi) = (c_k \xi, \xi) = c_k.$$

For arbitrary σ , we have

$$(1.9) \quad (\sigma\xi, \xi) = (P_0 \sigma P_0 \xi, \xi) \stackrel{(1.7)}{=} \left(\prod_{k \in [\sigma]} C_k \xi, \xi \right) = \prod_{k \in [\sigma]} c_k.$$

In other words, any spherical function of (G^D, K^D) is multiplicative in the following sense: its value on any permutation σ equals the product of its values on the disjoint cycles of σ . This multiplicativity property was first established by Thoma in [49]. Olshanski's theorem is, therefore, a generalization of this Thoma multiplicativity.

2. CLASSIFICATION OF IRREDUCIBLE ADMISSIBLE REPRESENTATIONS

2.1 Spectra of the A_i 's in spherical representations of (G^D, K^D) .

Fix a spherical representation π of the pair (G^D, K^D) and write simply g instead of $\pi(g)$.

Let μ denote the spectral measure for the operator A_1 and the spherical vector ξ . Since $\|A_i\| \leq 1$, this measure is supported at the segment $[-1, 1]$. The numbers c_k are the moments of μ

$$(2.1) \quad \int t^k \mu(dt) = (A_1^k \xi, \xi) = (P_0 A_1^k P_0 \xi, \xi) \stackrel{(1.6)}{=} c_{k+1}.$$

Therefore, the spherical function of the pair (G^D, K^D) is uniquely determined by the measure μ . We denote the spherical function corresponding to the measure μ by ϕ^μ

$$\phi^\mu(\sigma) = \prod_{k \in [\sigma]} \int_{[-1, 1]} t^{k-1} \mu(dt).$$

Let λ be the measure on $[-1, 1]^\infty$ which is the spectral measure for the operators A_1, A_2, \dots and the vector ξ . The claim of the following lemma is parallel to the Thoma multiplicativity.

Lemma 1. $\lambda = \mu^{\otimes \infty}$.

Proof. It suffices to check the identity

$$\begin{aligned} \int t_1^{k_1} \dots t_l^{k_l} d\lambda &= (A_1^{k_1} \dots A_l^{k_l} \xi, \xi) \\ &= (P_0 A_1^{k_1} \dots A_l^{k_l} P_0 \xi, \xi) \stackrel{(1.6)}{=} \prod_i c_{k_i+1} = \prod_i \int t^{k_i} d\mu \end{aligned}$$

for the integrals of all monomials. \square

Now we can prove the following

Theorem 1. *The measure μ is discrete and its atoms can only accumulate to zero $0 \in [-1, 1]$.*

Proof. Denote by s the transposition $(12) \in S(\infty)$. Let E be a Borel subset in $[\varepsilon, 1]$ where $\varepsilon > 0$. Denote by χ_E its characteristic function. We claim that

$$\varepsilon\mu(E) \leq \mu(E)^2.$$

To this end, we prove two inequalities

$$\varepsilon\mu(E) \leq (s\chi_E(A_1)\xi, \chi_E(A_1)\xi) \leq \mu(E)^2.$$

The expression in the middle is real because

$$s^{-1} = s.$$

On the one hand, by (1.3) and (1.5), we have

$$\begin{aligned} (s\chi_E(A_1)\xi, \chi_E(A_1)\xi) &= (s\chi_E(A_1)P_1\xi, \chi_E(A_1)P_1\xi) \\ &\stackrel{(1.3)}{=} (\chi_E(A_1)P_1sP_1\chi_E(A_1)\xi, \xi) \\ &\stackrel{(1.5)}{=} (\chi_E(A_1)P_1A_1P_1\chi_E(A_1)\xi, \xi) \\ &\stackrel{(1.3)}{=} (A_1\chi_E(A_1)\xi, \xi) = \int_E t d\mu \geq \varepsilon\mu(E). \end{aligned}$$

On the other hand, since A_1 and A_2 are commuting projections,

$$\begin{aligned} \chi_E(A_1)s\chi_E(A_1) &= \chi_E(A_1)^2s\chi_E(A_1)^2 \stackrel{(1.2)}{=} \chi_E(A_1)^2\chi_E(A_2)s\chi_E(A_1) \\ &\stackrel{(1.1,2)}{=} \chi_E(A_1)\chi_E(A_2)s\chi_E(A_1)\chi_E(A_2). \end{aligned}$$

Therefore,

$$\begin{aligned} (s\chi_E(A_1)\xi, \chi_E(A_1)\xi) &= (s\chi_E(A_1)\chi_E(A_2)\xi, \chi_E(A_1)\chi_E(A_2)\xi) \\ &\leq (\chi_E(A_1)\chi_E(A_2)\xi, \chi_E(A_1)\chi_E(A_2)\xi) = \mu(E)^2, \end{aligned}$$

where the last step relies on Lemma 1.

It follows from the inequality $\varepsilon\mu(E) \leq \mu(E)^2$ that either $\mu(E) = 0$ or $\mu(E) \geq \varepsilon$. An similar estimate holds for $E \subset [-1, -\varepsilon]$. This implies that the measure μ is discrete. Since μ is a probability measure, there are no more than $1/\varepsilon$ of its atoms in the interval $[\varepsilon, 1]$. This implies the second claim of the theorem. \square

We denote by $\text{supp } \mu$ the set of atoms of the measure μ .

2.2 Spectra of the A_i 's in admissible representations of (G^D, K^D) .

Now let π denote an irreducible admissible representation of depth $d > 0$ corresponding to some Young distributions Λ, M and some numbers $\{c_i\}$.

Let μ be the spectral measure μ of the operator A_{d+1} with respect to some unit vector ζ in the subspace $R(\pi)$. The measure μ is independent of the choice of ζ because the numbers $\{c_i\}$ are the moments of μ . By construction, the measure μ corresponds to a spherical representation of the pair

$$(G_d^D, K_d^D) \cong (G^D, K^D).$$

Therefore, μ is discrete. Let $\text{supp } \mu$ denote the set of its atoms.

By definition of the distribution Λ , the spectrum of any of the operators A_i , $i = 1, \dots, d$ in the space $R(\pi)$ is $\text{supp } \Lambda$.

Proposition 3. $\text{supp } \Lambda \subset \text{supp } \mu \cup \{0\}$, $\text{supp } M \subset \text{supp } \mu \cup \{0\}$.

Proof. Take $x \in \text{supp } \Lambda \setminus \{0\}$. Let ζ be a vector in the subspace $R(\pi)$, such that $\|\zeta\| = 1$ and $A_1\zeta = x\zeta$. Denote by δ_x the function equal to 1 at the point x , and to 0 at all other points. Then

$$\zeta = \delta_x(A_1)\zeta.$$

Denote by s the permutation $(1, d+1) \in S(\infty)$. Then

$$\begin{aligned} 0 < |x| &= |(A_1\zeta, \zeta)| \stackrel{(1.5)}{=} |(P_d s P_d \zeta, \zeta)| = |(s\zeta, \zeta)| = |(s\delta_x(A_1)\zeta, \delta_x(A_1)\zeta)| \\ &\stackrel{(1.1,2)}{=} |(s\delta_x(A_{d+1})\zeta, \delta_x(A_{d+1})\zeta)| \leq (\delta_x(A_{d+1})\zeta, \delta_x(A_{d+1})\zeta) = \mu(x), \end{aligned}$$

i.e., $x \in \text{supp } \mu$. The argument for M is analogous. \square

This proposition was previously proved in [32, Theorem 4.6] in a more complicated way.

2.3 The Thoma theorem.

Let π be a spherical representation and μ the corresponding spectral measure. Denote by $\alpha_i, -\beta_i, \alpha_i > 0, \beta_i > 0$ the non-zero elements in $\text{supp } \mu$. For $x \neq 0$ we set $\nu(x) = \mu(x)/|x|$. By virtue of (2.1),

$$\begin{aligned} (2.2) \quad c_k &= \sum_i \alpha_i^{k-1} \mu(\alpha_i) + \sum_i (-\beta_i)^{k-1} \mu(-\beta_i) \\ &= \sum_i \alpha_i^k \nu(\alpha_i) + (-1)^{k-1} \sum_i \beta_i^k \nu(-\beta_i) \end{aligned}$$

for every $k > 1$.

Theorem 2. *The numbers $\nu(\alpha_i), \nu(-\beta_i)$ are positive integers.*

In the proof of this theorem we shall need the following lemma. Let $\sigma \in S(\infty)$ be an arbitrary permutation. Denote by \mathbb{N}/σ the set of orbits of σ on the set \mathbb{N} . For an orbit $p \in \mathbb{N}/\sigma$, denote by $|p|$ its cardinality.

Lemma 2. *Let $f_i(t), g_i(t), i = 1, 2, \dots$ be continuous functions on $[-1, 1]$, all but finitely many identically equal to 1. Then*

$$\begin{aligned} \text{a)} \quad & \left(\sigma \prod_{i=1}^{\infty} f_i(A_i) \xi, \xi \right) = \prod_{p \in \mathbb{N}/\sigma} \int t^{|p|-1} \prod_{j \in p} f_j(t) d\mu, \\ \text{b)} \quad & \left(\sigma \prod_{i=1}^{\infty} f_i(A_i) \xi, \prod_{i=1}^{\infty} g_i(A_i) \xi \right) = \prod_{p \in \mathbb{N}/\sigma} \int t^{|p|-1} \prod_{j \in p} f_j(t) \overline{g_j(t)} d\mu, \end{aligned}$$

for $\sigma \in S(\infty)$.

Proof. a) One can assume that $f_i(t) = t^{k_i}$, $i = 1, 2, \dots$, and that $k_i = 0$ for $i \gg 1$. For $p \in \mathbb{N}/\sigma$ set $\Sigma(p) = \sum_{j \in p} k_j$. Then $\prod_{j \in p} f_j(t) = t^{\Sigma(p)}$. The following identity

$$P_0 \sigma \prod_i A_i^{k_i} P_0 = P_0 \prod_{p \in \mathbb{N}/\sigma} C_{|p| + \Sigma(p)}$$

generalizes the identities (1.6) and (1.7) and is proved similarly. It follows from this identity that

$$\begin{aligned} \left(\sigma \prod_{i=1}^{\infty} f_i(A_i) \xi, \xi \right) &= \left(\sigma \prod A_i^{k_i} \xi, \xi \right) = \prod_{p \in \mathbb{N}/\sigma} c_{|p|+\Sigma(p)} \\ &= \prod_{p \in \mathbb{N}/\sigma} \int t^{|p|+\Sigma(p)-1} d\mu = \prod_{p \in \mathbb{N}/\sigma} \int t^{|p|-1} \prod_{j \in p} t^{k_j} d\mu. \end{aligned}$$

Part b) follows from a), equation (1.2), and from the obvious equality

$$\prod_{j \in p} f_j(t) = \prod_{j \in p} f_{\sigma(j)}(t). \quad \square$$

Remark. The above lemma holds for a larger class of functions, for example, for functions which are pointwise limits of uniformly bounded sequences of continuous functions. This follows from the functional calculus of operators, cf. [41, v.1, Theorem VII.2(d)]. The function $\delta_x(t)$,

$$\delta_x(t) = \begin{cases} 1, & t = x, \\ 0, & t \neq x, \end{cases}$$

belongs to this class. Alternatively, by Theorem 1, we can take instead of δ_x a continuous function equal to 1 at the point x , and to zero at other points in $\text{supp } \mu$.

Proof of Theorem 2. Fix some $\alpha = \alpha_i$ and set $\nu = \nu(\alpha_i)$. Consider the vector

$$\zeta^{(m)} = \prod_{i=1}^m \delta_{\alpha}(A_i) \xi.$$

By Lemma 2,

$$(\sigma \zeta^{(m)}, \zeta^{(m)}) = \prod_{p \in \mathbb{N}/\sigma} \alpha^{|p|-1} \mu(\alpha) = \alpha^{m-\ell(\sigma)} \mu(\alpha)^{\ell(\sigma)} = \alpha^m \nu^{\ell(\sigma)}$$

for $\sigma \in S(m)$, where $\ell(\sigma)$ denotes the number of cycles of the permutation $\sigma \in S(m)$. Effectively, what the consideration of the vectors $\zeta^{(m)}$, $m = 1, 2, \dots$, allows us is to single out just one point α from the set $\text{supp } \mu$.

Let $\text{Alt}(m)$ be the operator of antisymmetrization over the group $S(m)$. Then

$$\begin{aligned} (\text{Alt}(m) \zeta^{(m)}, \zeta^{(m)}) &= \frac{1}{m!} \sum_{\sigma \in S(m)} \text{sgn}(\sigma) (\sigma \zeta^{(m)}, \zeta^{(m)}) = \\ (2.3) \quad &= \frac{\alpha^m}{m!} \sum_{\sigma \in S(m)} \text{sgn}(\sigma) \nu^{\ell(\sigma)} = \frac{\alpha^m}{m!} \sum_{\sigma \in S(m)} (-1)^{m+\ell(\sigma)} \nu^{\ell(\sigma)} = \\ &= \frac{\alpha^m}{m!} \nu(\nu-1) \dots (\nu-m+1), \end{aligned}$$

where we have applied the equality $\text{sgn } \sigma = (-1)^{m+\ell(\sigma)}$ and the well known identity

$$(2.4) \quad \sum_{\sigma \in S(m)} x^{\ell(\sigma)} = x(x+1) \dots (x+m-1)$$

(cf. [45, Proposition 1.3.4]). Since $\text{Alt}(m)$ is a projection, the last product in (2.3) should be nonnegative for all m which is only possible if it is terminating, i.e., if $\nu = \nu(\alpha) \in \mathbb{N}$.

If we replace the representation T by its tensor product with the representation $\text{sgn} \otimes \text{sgn}$, then the measure $\mu(x)$ is replaced by the measure $\mu(-x)$. It follows that $\nu(-\beta_i) \in \mathbb{N}$. \square

We call to a discrete probability measure μ on $[-1, 1]$ satisfying

$$\frac{\mu(x)}{|x|} \in \mathbb{Z}_+, \quad x \neq 0,$$

a *Thoma measure*.

The theorem just proved provides a necessary condition for the existence of representations. It follows from the explicit construction of representations [6, 32] (see also Chapter 3), or else from a direct verification of the positive definiteness [49], that this condition is sufficient as well. Therefore the description of spherical functions for the pair $(G^\mathcal{D}, K^\mathcal{D})$ is established. In order to state it in the classical form, it is convenient to treat the set $\text{supp } \mu$ as a multiset in which every element x is repeated $\mu(x)/|x|$ times.

Theorem (Thoma, [49]). *The characters of the group $S(\infty)$ are precisely the functions of the form*

$$\phi(\sigma) = \prod_{k \in [\sigma]} \left(\sum_i \alpha_i^k + (-1)^{k-1} \sum_i \beta_i^k \right),$$

where

$$\begin{aligned} \alpha_1 \geq \alpha_2 \geq \alpha_3 \geq \dots > 0 \quad \beta_1 \geq \beta_2 \geq \beta_3 \geq \dots > 0 \\ \sum \alpha_i + \sum \beta_i \leq 1. \end{aligned}$$

2.4 Another proof of Thoma theorem.

In this Section we show how the presentation

$$c_k = \int t^{k-1} d\mu$$

simplifies the original proof of the Thoma theorem. Throughout the section, except for the very last punch-line, we closely follow Thoma's original exposition [49].

Let ϕ be a character of $S(\infty)$. Consider the restriction of ϕ to a finite symmetric group $S(n)$. The characters of finite symmetric group $S(n)$ are labeled by Young diagrams with n boxes. Let χ^λ be the (non normalized) character corresponding to an irreducible representation λ . Since the function ϕ is positive definite, its

restriction $\phi|_{S(n)}$ to the group $S(n)$ is a non-negative linear combination of the functions χ^λ

$$\phi|_{S(n)} = \sum_{\lambda, |\lambda|=n} m(\lambda) \chi^\lambda, \quad m(\lambda) \geq 0.$$

We call the numbers $m(\lambda)$ the *Fourier coefficients* of the function ϕ . One can compute them using the orthonormality of characters with respect to the Hermitian inner product

$$\langle f_1, f_2 \rangle_{S(n)} = \frac{1}{n!} \sum_{g \in S(n)} f_1(g) \overline{f_2(g)}.$$

Evidently, the numbers $m(\lambda)$ have to satisfy some coherency conditions. Namely, consider the representation of the group $S(n+1)$ determined by a Young diagram Λ , $|\Lambda| = n+1$. According to the Young branching rule

$$(2.5) \quad \chi^\Lambda|_{S(n)} = \sum_{\lambda, \Lambda \searrow \lambda} \chi^\lambda,$$

where the notation $\Lambda \searrow \lambda$ means that the diagram λ is obtained from the diagram Λ by removing a box. By virtue of (2.5), the numbers $m(\lambda)$ have to satisfy the conditions

$$m(\lambda) = \sum_{\Lambda, \Lambda \searrow \lambda} m(\Lambda).$$

Conversely, every collection of non-negative numbers $m(\lambda)$ satisfying the above coherence condition determines some positive definite function on the group $S(\infty)$. The function is normalized if and only if

$$m(\emptyset) = 1.$$

By Proposition 2 (see also Section 1.3) this function is indecomposable if and only if it is multiplicative in the cycles a of permutation. This multiplicativity imposes severe restrictions on the numbers $m(\lambda)$. Our next goal is to obtain a precise form of these restrictions.

Recall the definition of the external product of characters of symmetric groups. Let $\mathfrak{Z}(S(n))$ denote the linear space of central functions on the groups $S(n)$ and suppose $f_1 \in \mathfrak{Z}(S(n_1))$ and $f_2 \in \mathfrak{Z}(S(n_2))$ are some central functions. Consider the following function on $S(n_1) \times S(n_2)$

$$(f_1 \otimes f_2)(g_1, g_2) = f_1(g_1) f_2(g_2).$$

The external product of f_1 and f_2 is, by definition,

$$f_1 \circ f_2 = \text{Ind}_{S(n_1) \times S(n_2)}^{S(n_1+n_2)} f_1 \otimes f_2 \in \mathfrak{Z}(S(n_1+n_2)).$$

Denote by η^k the trivial character of $S(k)$ if $k \geq 0$, and zero otherwise. The Frobenius formula [23] asserts that

$$\chi^\lambda = \left| \begin{array}{cccccc} \eta^{\lambda_1} & \eta^{\lambda_1+1} & \eta^{\lambda_1+2} & \dots & \dots & \dots \\ \eta^{\lambda_2-1} & \eta^{\lambda_2} & \eta^{\lambda_2+1} & \eta^{\lambda_2+2} & \dots & \dots \\ \vdots & \eta^{\lambda_3-1} & \eta^{\lambda_3} & \eta^{\lambda_3+1} & \dots & \dots \\ \vdots & & & \vdots & & \vdots \\ \vdots & & & \vdots & & \vdots \\ \vdots & \dots & \dots & \eta^{\lambda_i-i+j} & \dots & \vdots \\ \vdots & & & \vdots & & \vdots \end{array} \right|^\circ$$

where the multiplication is to be understood as the external one. The Frobenius formula expresses the character χ^λ of the irreducible representation of the symmetric group as a linear combination of the functions η^λ

$$\eta^\lambda = \eta^{\lambda_1} \circ \eta^{\lambda_2} \circ \dots$$

The following lemma is straightforward.

Lemma 3. *A function $\phi \in \mathfrak{Z}(S(n))$ is multiplicative in the cycles of a permutation if and only if for all n_1, n_2 such that $n_1 + n_2 = n$, and for all functions $f_1 \in \mathfrak{Z}(S(n_1))$, $f_2 \in \mathfrak{Z}(S(n_2))$ we have*

$$(2.6) \quad \begin{aligned} & \langle \phi|_{S(n_1) \times S(n_2)}, f_1 \otimes f_2 \rangle_{S(n_1) \times S(n_2)} = \\ & = \langle \phi|_{S(n_1)}, f_1 \rangle_{S(n_1)} \langle \phi|_{S(n_2)}, f_2 \rangle_{S(n_2)}. \end{aligned}$$

Remark. The equality (2.6) is antilinear in f_1 and f_2 . Therefore, it sufficient to check it for f_1 and f_2 ranging independently over some linear bases of the spaces $\mathfrak{Z}(S(n_1))$ and $\mathfrak{Z}(S(n_2))$.

Proposition 4. *The coefficients $m(\lambda)$ correspond to a multiplicative central function on the group $S(\infty)$ if and only if we have*

$$(2.7) \quad m(\lambda) = \det [m(\lambda_i - i + j)]_{i,j}$$

for any Young diagram λ . Here $m(k)$, $k \geq 0$, stands for $m((k))$ and we set $m(k) = 0$ for $k < 0$.

Proof. Let $\phi \in \mathfrak{Z}(S(\infty))$ be a multiplicative function. By the Frobenius reciprocity,

$$\langle \phi|_{S(|\lambda|)}, \eta^\lambda \rangle_{S(|\lambda|)} = \langle \phi|_{S(\lambda_1) \times \dots \times S(\lambda_s)}, 1 \rangle_{S(\lambda_1) \times \dots \times S(\lambda_s)}.$$

By the multiplicativity of ϕ this equals

$$\prod_i \langle \phi|_{S(\lambda_i)}, 1 \rangle_{S(\lambda_i)} = \prod_i m(\lambda_i).$$

Therefore, by the Frobenius formula,

$$m(\lambda) = \langle \phi|_{S(|\lambda|)}, \chi^\lambda \rangle_{S(|\lambda|)} = \det [m(\lambda_i - i + j)].$$

In the opposite direction, assume (2.7). Fix some n and consider the following dominance ordering of all partitions of n

$$\mu \leq \lambda \iff \forall i \quad \mu_1 + \dots + \mu_i \leq \lambda_1 + \dots + \lambda_i.$$

Then (it is clear from Frobenius formula) the transition matrix from the basis $\{\chi^\lambda\}$ to the basis $\{\eta^\lambda\}$ is triangular. The equality

$$(2.8) \quad \langle \phi, \eta^\lambda \rangle = \prod_i m(\lambda_i)$$

is evident if $\lambda = (n)$. Assume that the equality is proved for all partitions μ of n , such that $\mu > \lambda$. Then it follows from (2.7) and Frobenius formula that it is also true for λ . Therefore, the equality (2.8) is true for all partitions λ .

Now let λ, μ be two arbitrary partitions. Set $|\lambda| = n_1$, $|\mu| = n_2$. Denote by $\lambda \cup \mu$ the union of parts of λ and μ . Then

$$\begin{aligned} \langle \phi|_{S(n_1) \times S(n_2)}, \eta^\lambda \otimes \eta^\mu \rangle_{S(n_1) \times S(n_2)} &= \langle \phi|_{S(n_1+n_2)}, \eta^{\lambda \cup \mu} \rangle_{S(n_1+n_2)} \\ &= \prod_i m(\lambda_i) \prod_j m(\mu_j) = \langle \phi|_{S(n_1)}, \eta^\lambda \rangle_{S(n_1)} \langle \phi|_{S(n_2)}, \eta^\mu \rangle_{S(n_2)}. \end{aligned}$$

Since the functions η^λ form a basis in the space of central functions, the multiplicativity of ϕ is established. \square

Recall that a sequence of real numbers $\{a_i\}$, $i = 0, 1, 2, \dots$ is said to be *totally positive* if all the minors of the following infinite Toeplitz matrix

$$\begin{bmatrix} a_0 & a_1 & a_2 & a_3 & \dots \\ 0 & a_0 & a_1 & a_2 & \ddots \\ & 0 & a_0 & a_1 & \ddots \\ & & 0 & a_0 & \ddots \\ & & & \ddots & \ddots \end{bmatrix}$$

are non-negative. In particular, all the numbers a_i are non-negative.

A shift of indices and the multiplication of the entries by a positive number preserve the total positivity, hence we can always assume that $a_0 = 1$. Assume that for some $i > 0$ we have $a_i = 0$. Then it follows from the inequality

$$\det \begin{bmatrix} a_i & a_{i+1} \\ a_0 & a_1 \end{bmatrix} \geq 0$$

that $a_j = 0$ for all $j \geq i$. Hence either all entries of a totally positive sequence are positive or there exist such N that

$$a_i = 0 \iff i \geq N.$$

One easily checks that the multiplication by a positive geometric series preserves total positivity. Therefore, if a sequence $\{a_i\}$ has at least two positive entries, we can assume that

$$a_0 = a_1 = 1.$$

We call such a sequence *normalized totally positive*.

Proposition 5. *The map*

$$\phi \rightarrow \{m(i)\}, \quad i = 0, 1, \dots$$

is a bijection of the set of the characters of the group $S(\infty)$ onto the set of normalized totally positive sequences.

Proof. Let ϕ be a character of the group $S(\infty)$. Clearly, $m(0) = m(1) = 1$. The inner product of $\phi|_{S(n)}$ with the trace of any representation of $S(n)$ is always non-negative. In particular, this is true for the trace $\chi^{\lambda/\mu}$ of the representation corresponding to the skew Young diagram λ/μ . By multiplicativity of ϕ and the analog of the Frobenius formula for $\chi^{\lambda/\mu}$ [28, §1.5, (5.4)],

$$\langle \phi|_{S(|\lambda/\mu|)}, \chi^{\lambda/\mu}|_{S(|\lambda/\mu|)} \rangle = \det [m(\lambda_i - \mu_j - i + j)]_{ij} \geq 0.$$

One can easily see that all the minors involved in the definition of total positivity can be obtained in this way.

In the opposite direction, assume that we are given a normalized totally positive sequence $\{m(i)\}$, $i = 0, 1, \dots$. Define a central function ϕ_n on every symmetric group $S(n)$ by the formula

$$\langle \phi_n, \eta^\lambda \rangle = \prod_i m(\lambda_i), \quad |\lambda| = n.$$

Because of $m(1) = 1$ we have

$$\langle \phi_{n+1}|_{S(n)}, \eta^\lambda \rangle = \langle \phi_{n+1}, \eta^{\lambda \cup 1} \rangle = m(1) \prod_i m(\lambda_i) = \langle \phi_n, \eta^\lambda \rangle.$$

This yields the coherence of $\{\phi_n\}$

$$\phi_{n+1}|_{S(n)} = \phi_n.$$

The multiplicativity of ϕ is clear, its positive definiteness follows from the Frobenius formula, and $m(0) = 1$ implies $\phi(1) = 1$. \square

It is convenient to form a generating series

$$H(t) = 1 + m(1)t + m(2)t^2 + \dots$$

for the numbers $m(i)$, $i = 0, 1, \dots$. As we already know, either $H(t)$ is a polynomial, or all the terms of $H(t)$ are non-zero. The series $H(t)$ has the following expression in terms of the numbers c_i , $i = 1, 2, \dots$. Recall that

$$c_k = \phi((12 \dots k)).$$

Suppose that $\lambda = 1^{\rho_1} 2^{\rho_2} \dots$. We have

$$\begin{aligned} m(k) &= \langle \phi|_{S(k)}, 1 \rangle_{S(k)} \\ &= \frac{1}{k!} \sum_{\lambda, |\lambda|=k} \frac{k!}{\prod_i \rho_i! i^{\rho_i}} \prod_i c_i^{\rho_i} \\ &= \sum_{\lambda, |\lambda|=k} \prod_i \frac{1}{\rho_i!} \left(\frac{c_i}{i} \right)^{\rho_i}. \end{aligned}$$

Therefore,

$$\begin{aligned}
H(t) &= \sum_{k=0}^{\infty} t^k \sum_{\rho_1+2\rho_2+3\rho_3+\dots=k} \prod_i \frac{1}{\rho_i!} \left(\frac{c_i}{i}\right)^{\rho_i} \\
&= \prod_{i=1}^{\infty} \sum_{j=0}^{\infty} \frac{1}{j!} \left(\frac{c_i t^i}{i}\right)^j \\
&= \prod_{i=1}^{\infty} \exp\left(\frac{c_i t^i}{i}\right) = \exp\left(\sum_{i=1}^{\infty} \frac{c_i}{i} t^i\right) \quad .
\end{aligned}$$

Vice versa, the numbers c_i can be easily obtained from the series $H(t)$,

$$c_1 + c_2 t + c_3 t^2 + \dots = \frac{H'(t)}{H(t)}.$$

We shall state the Thoma theorem in the following form.

Theorem (Thoma [49]). *The generating functions for Fourier coefficients $\{m(i)\}$, $i = 0, 1, \dots$ of the characters of the group $S(\infty)$ have the form*

$$H(t) = e^{\gamma t} \prod_i \frac{1 + \beta_i t}{1 - \alpha_i t}$$

where

$$\alpha_i \geq 0, \quad \beta_i \geq 0, \quad \gamma \geq 0, \quad \sum \alpha_i + \sum \beta_i + \gamma = 1.$$

This is equivalent to the following:

Theorem (Edrei [17]). *The generating functions of totally positive sequences have the form*

$$H(t) = e^{\gamma t} \prod_i \frac{1 + \beta_i t}{1 - \alpha_i t}$$

where

$$\alpha_i \geq 0, \quad \beta_i \geq 0, \quad \gamma \geq 0, \quad \sum \alpha_i + \sum \beta_i < \infty.$$

Proof of the Thoma theorem. First, assume that $H(t)$ is not a polynomial. Then it follows from the inequalities

$$\det \begin{bmatrix} m(i) & m(i+1) \\ m(i-1) & m(i) \end{bmatrix} \geq 0$$

that

$$(2.9) \quad 1 = \frac{m(1)}{m(0)} \geq \frac{m(2)}{m(1)} \geq \frac{m(3)}{m(2)} \geq \dots \geq 0.$$

Therefore, there exists a limit

$$\alpha = \lim_{n \rightarrow \infty} \frac{m(n+1)}{m(n)}, \quad n \rightarrow \infty.$$

Obviously, $0 \leq \alpha \leq 1$ and $1/\alpha$ is the convergence radius of the series $H(t)$.

If $\alpha = 1$ then by (2.9) we have $m(i) \equiv 1$ and

$$H(t) = \frac{1}{1-t}.$$

This H -series corresponds to the trivial representation. We denote it by $H_1(t)$.

Assume now that $0 < \alpha < 1$. Then we construct, starting from the character ϕ , a new character $\tilde{\phi}$ with Fourier coefficients

$$\tilde{m}(\lambda) = (1 - \alpha)^{-|\lambda|} \lim_{n \rightarrow \infty} \frac{m(n \cup \lambda)}{m(n)}, \quad n \rightarrow \infty.$$

It is easy to compute these limits explicitly. One has

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{m(n \cup \lambda)}{m(n)} &= \lim_{n \rightarrow \infty} \frac{1}{m(n)} \det \begin{bmatrix} m(n) & m(n+1) & m(n+2) & \dots \\ m(\lambda_1 - 1) & m(\lambda_1) & m(\lambda_1 + 1) & \dots \\ m(\lambda_2 - 2) & m(\lambda_2 - 1) & m(\lambda_2) & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \\ &= \lim_{n \rightarrow \infty} \det \begin{bmatrix} 1 & m(n+1)m(n)^{-1} & m(n+2)m(n)^{-1} & \dots \\ m(\lambda_1 - 1) & m(\lambda_1) & m(\lambda_1 + 1) & \dots \\ m(\lambda_2 - 2) & m(\lambda_2 - 1) & m(\lambda_2) & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \\ &= \det \begin{bmatrix} 1 & \alpha & \alpha^2 & \dots \\ m(\lambda_1 - 1) & m(\lambda_1) & m(\lambda_1 + 1) & \dots \\ m(\lambda_2 - 2) & m(\lambda_2 - 1) & m(\lambda_2) & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \\ &= \det \begin{bmatrix} 1 & 0 & 0 & \dots \\ m(\lambda_1 - 1) & m(\lambda_1) - \alpha m(\lambda_1 - 1) & m(\lambda_1 + 1) - \alpha m(\lambda_1) & \dots \\ m(\lambda_2 - 2) & m(\lambda_2 - 1) - \alpha m(\lambda_2 - 2) & m(\lambda_2) - \alpha m(\lambda_2 - 1) & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \\ &= \det \begin{bmatrix} m(\lambda_1) - \alpha m(\lambda_1 - 1) & m(\lambda_1 + 1) - \alpha m(\lambda_1) & \dots \\ m(\lambda_2 - 1) - \alpha m(\lambda_2 - 2) & m(\lambda_2) - \alpha m(\lambda_2 - 1) & \dots \\ \vdots & \vdots & \ddots \end{bmatrix}. \end{aligned}$$

In particular,

$$(2.9) \quad \tilde{m}(k) = (1 - \alpha)^{-k} (m(k) - \alpha m(k-1)).$$

Therefore,

$$\tilde{m}(\lambda) = \det [\tilde{m}(\lambda_i - i + j)]_{i,j}.$$

We see that the Fourier coefficients $\{\tilde{m}(\lambda)\}$ determine a multiplicative positive definite function on the group $S(\infty)$. Set

$$\tilde{H}(t) = \sum \tilde{m}(k) t^k.$$

Then the equality (2.9) means that

$$H(t) = H_1(\alpha t) \tilde{H}((1 - \alpha)t).$$

The representation-theoretical meaning of this factorization will be made clear in Section 3. Note that the convergence radius of the series $\tilde{H}((1 - \alpha)t)$ is not less than that of the series $H(t)$. Iterating this factorization, we obtain the following factorization

$$H(t) = H_{\text{ent}} \left(\left(1 - \sum \alpha_i \right) t \right) \prod_i H_1(\alpha_i t),$$

where

$$\alpha_i \geq 0, \quad \sum \alpha_i \leq 1,$$

and H_{ent} is an entire function which is also an H -series for a character of the group $S(\infty)$.

Now, instead of ϕ , consider the character ϕ^- ,

$$\phi^-(g) = \text{sgn}(g)\phi(g), \quad g \in S(\infty).$$

In particular,

$$c_k^- = (-1)^{k-1} c_k.$$

Therefore, the character ϕ^- is associated with the H -series

$$H^-(t) = \exp \left(\sum_{i=1}^{\infty} (-1)^{i-1} \frac{c_i}{i} t^i \right) = \exp \left(- \sum_{i=1}^{\infty} \frac{c_i}{i} (-t)^i \right) = H(-t)^{-1}.$$

Now factoring the function $H_{\text{ent}}(-t)^{-1}$ we obtain

$$H(t) = H^*(\gamma t) \prod H_1(\alpha_i t) \prod H_{\text{sgn}}(\beta_i t) \quad \sum \alpha_i + \sum \beta_i + \gamma = 1,$$

where

$$H_{\text{sgn}}(t) = 1 + t,$$

and $H^*(t)$ is an H -series for a character of the group $S(\infty)$ which is an entire function with no zeros. Hence the series

$$c_1 + c_2 t + c_3 t^2 + \cdots = \frac{H'(t)}{H(t)}$$

represents an entire function. Now, and this is the whole point of this section, we shall show that this is only possible if

$$c_1 = 1, \quad c_2 = c_3 = \cdots = 0.$$

Indeed, we have

$$c_k = \int t^{k-1} d\mu.$$

Assume that the measure μ is not concentrated at $t = 0$. Then for some $\varepsilon > 0$ we have $\mu([-1, 1] \setminus [-\varepsilon, \varepsilon]) = \text{const} > 0$, hence

$$c_{2k+1} = \int t^{2k} d\mu \geq \text{const} \varepsilon^{2k}$$

for all k , and hence the convergence radius of the series $\sum c_k z^{k-1}$ does not exceed $1/\varepsilon$. Therefore, the measure μ should be supported at zero, which implies $c_2 = c_3 = \dots = 0$. In other words,

$$H^*(t) = e^t. \quad \square$$

Remark. The case of entire function without zeros and poles was actually the most difficult part in the original proof [17,49].

2.5 Classification of the irreducible admissible representations of the pair (G^D, K^D) .

Let π be an irreducible admissible representation of depth d determined by a Thoma measure μ and Young distributions Λ, M .

Denote by s and t the permutations $(1, d+1), (-1, -d-1) \in G^D$.

Lemma 4. *Suppose $x \in \text{supp } \Lambda$ and suppose a vector $\zeta \in R(\pi)$, $\|\zeta\| = 1$, satisfies $A_1 \zeta = x \zeta$. Set $\zeta^{(x)} = \delta_x(A_{d+1}) \zeta$. Then*

- a) $(\zeta^{(x)}, \zeta^{(x)}) = \mu(x)$,
- b) $(s\zeta^{(x)}, \zeta^{(x)}) = x$,
- c) $(st\zeta^{(x)}, \zeta^{(x)}) = 0$.

If, in addition, $x \in \text{supp } M$ and $A_{-1} \zeta = x \zeta$ then

- d) $(t\zeta^{(x)}, \zeta^{(x)}) = x$.

Proof. Part a) follows from the definition of the spectral measure. By Proposition 3, $\mu(x) \neq 0$ if $x \neq 0$. Part b) has actually been already established in the proof of Proposition 3. Part d) then follows from the fact that, by virtue of (1.4),

$$A_{-d-1}^k \zeta = A_{d+1}^k \zeta, \quad k = 1, 2, \dots,$$

and hence

$$\zeta^{(x)} = \delta_x(A_{-d-1}) \zeta.$$

Let us check part c). The operator $P_d st P_d$ is conjugate to P_{d-1} , and therefore equals zero since $H(\pi)_{d-1} = 0$ by assumption. Hence,

$$\begin{aligned} 0 &= (P_d st P_d \zeta, \zeta) = (st \zeta, \zeta) = (st \delta_x(A_1) \zeta, \delta_x(A_1) \zeta) \\ &\stackrel{(1.1), (1.2)}{=} (st \delta_x(A_{d+1}) \zeta, \delta_x(A_{d+1}) \zeta) = (st \zeta^{(x)}, \zeta^{(x)}). \quad \square \end{aligned}$$

Theorem 3. *An irreducible admissible representation of depth d with the Thoma measure μ and Young distributions Λ, M , $|\Lambda| = |M| = d$ exists if and only if for all $x \in [-1, 1]$ we have*

$$\begin{aligned} \ell(\Lambda(x)) + \ell(M(x)) &\leq \mu(x)/|x|, & x > 0, \\ \ell(\Lambda'(x)) + \ell(M'(x)) &\leq \mu(x)/|x|, & x < 0, \end{aligned}$$

where the prime denotes the transposition of Young diagrams and ℓ stands for the number of rows in a Young diagram.

Proof. The sufficiency was proved in [32] using an explicit construction of the representation. We shall give the explicit construction of all corresponding representations in Section 3.

Let us prove the necessity. Assume that

$$x \in \text{supp } \Lambda \cup \text{supp } M \quad \text{and} \quad x > 0.$$

Set

$$l_1 = \ell(\Lambda(x)), \quad l_2 = \ell(M(x)).$$

We shall assume that $l_1, l_2 > 0$. The case where $l_1 = 0$ or $l_2 = 0$ is more simple, and can be dealt with analogously. Denote by $S(l_1) \times S(l_2)$ the subgroup in the group $G^D(d)$ which permutes the numbers $\{1, \dots, l_1\}$ and $\{-1, \dots, -l_2\}$.

By the branching rule for the representations of finite symmetric groups and the definition of Λ and M , there exists a vector $\zeta \in R(\pi)$ which is anti-invariant with respect to the group $S(l_1) \times S(l_2)$ and also satisfies

$$A_i \zeta = x \zeta, \quad i = 1, \dots, l_1, -1, \dots, -l_2.$$

Then the vector $\zeta^{(x)}$ is $S(l_1) \times S(l_2)$ -anti-invariant, too.

Let Alt be the operator of anti-symmetrization over the group $S(l_1 + 1) \times S(l_2 + 1)$ permuting $\{1, 2, \dots, l, d + 1\}$ and $\{-1, -2, \dots, -l, -d - 1\}$. The function

$$(\text{sgn}(g) g \zeta^{(x)}, \zeta^{(x)})$$

is invariant with respect to right and left translations by the elements of the group $S(l_1) \times S(l_2)$. The group $S(l_1 + 1) \times S(l_2 + 1)$ consists of four double cosets with respect to subgroup $S(l_1) \times S(l_2)$; their representatives are $1, s, t, st$, and their cardinalities are $l_1! l_2!, l_1 l_1! l_2!, l_2 l_1! l_2!, l_1 l_2 l_1! l_2!$. Hence, by the above lemma we have

$$\begin{aligned} (\text{Alt } \zeta^{(x)}, \zeta^{(x)}) &= \frac{l_1! l_2! \mu(x) - l_1 l_1! l_2! x - l_2 l_1! l_2! x + l_1 l_2 l_1! l_2! 0}{(l_1 + 1)! (l_2 + 1)!} \\ &= \frac{x(\mu(x)/x - l_1 - l_2)}{(l_1 + 1)(l_2 + 1)}. \end{aligned}$$

Since the operator Alt is a projection, the result is non-negative, and therefore

$$l_1 + l_2 \leq \mu(x)/x.$$

The case $x < 0$ can be reduced to that of $x > 0$ by replacing T with $T \otimes (\text{sgn} \otimes \text{sgn})$. \square

Remark. From the point of view of the figure in Section 1.2, the use of spectral projectors $\delta_x(A_i)$ allows one to get rid of all Young diagrams but those growing from the point x . In Section 3, we shall be concerned with the opposite problem: how to plant a Young diagram at a given point of the interval $[-1, 1]$.

2.6 Description of $K^E \backslash G^E / K^E$.

In this section G stands the group $G^E(n)$ and K denotes the subgroup $K^E(n)$. We shall recall basic facts about the cosets $K \backslash G / K$ which we shall need in the next section. Denote by $S(n) \subset G^E(n)$ the subgroup fixing the points $\{-1, \dots, -n\}$.

The set $K \backslash G / K$ clearly coincides with the set of orbits of the group K on the set G / K . The set G / K is naturally identified with the set Π of partitions of the set $\{\pm 1, \pm 2, \dots, \pm n\}$ into pairs (note the difference between partitions of a set and partitions of a number). The group K is the stabilizer of the partition σ

$$\sigma = \{\{\pm 1\}, \dots, \{\pm n\}\}.$$

For any two partitions τ, v let $\tau \vee v$ denote their least upper bound, that is, the finest partition consisting of whole blocks of τ and v . If τ and v were partitions into pairs (more generally, into even blocks), then the partition $\tau \vee v$ is also a partition into even blocks. Therefore the block cardinalities of $\tau \vee v$, divided by two, form a partition of n which we denote by $\tau \nabla v$. The function $f(\tau) = \tau \nabla \sigma$ with values in the set of partitions of n is an invariant of the action of the group K on the space Π . The following proposition is well known [46, 32].

Proposition. *The function $f(\tau) = \tau \nabla \sigma$ separates the orbits of the group K in Π .*

Corollary.

- (1) *The set $K \backslash G / K$ is parameterized by partitions λ of the number n .*
- (2) *The intersection of the double coset corresponding to a partition λ with the subgroup $S(n)$ consists of permutations with the cycle structure λ . In particular, this intersection is non-empty.*

Let $\lambda = 1^{m_1} 2^{m_2} \dots$ be the partition with m_i parts of size i . Denote by $\ell(\lambda)$ the number of parts in the partition λ . Set $z_\lambda = \prod i^{m_i} m_i!$. Denote by $K \lambda K$ the double coset corresponding to the partition λ . The following proposition can be established by a direct combinatorial argument [46]:

Proposition 7.

$$|K \lambda K| = 2^{2n - \ell(\lambda)} \frac{(n!)^2}{z_\lambda}$$

Corollary. *For an element $g \in G^E(n)$, let $\ell(g)$ be the number of parts in the partition corresponding to the double coset KgK . If t is a formal variable, then*

$$\sum_{g \in G^E(n)} t^{\ell(g)} = n! 2^n t(t+2)(t+4) \cdots (t+2n-2).$$

Proof. We have

$$\begin{aligned} \sum_{g \in G^E(n)} t^{\ell(g)} &= \sum_{\lambda \vdash n} 2^{2n-\ell(\lambda)} (n!)^2 z_\lambda^{-1} t^{\ell(\lambda)} \\ &= n! 2^{2n} \sum_{\lambda \vdash n} n! z_\lambda^{-1} \left(\frac{t}{2}\right)^{\ell(\lambda)} \end{aligned}$$

By the identity (2.4) for the Stirling numbers [45] already employed in the proof of Theorem 2, this is

$$\begin{aligned} &= n! 2^{2n} \frac{t}{2} \left(\frac{t}{2} + 1\right) \left(\frac{t}{2} + 2\right) \cdots \left(\frac{t}{2} + n - 1\right) \\ &= n! 2^n t(t+2)(t+4) \cdots (t+2n-2). \quad \square \end{aligned}$$

2.7 Spherical representations of the pair (G^E, K^E) .

Let π be a spherical representation of the pair (G^E, K^E) and let ϕ be the corresponding spherical function. We know from the previous section that the group G^E is the product of its subgroups $K^E S(\infty) K^E$. Hence ϕ , as a K^E -biinvariant function, is completely determined by its restriction to the subgroup $S(\infty)$. This restriction is a normalized positive definite function. Just as in the case of spherical representations of the pair (G^D, K^D) , one checks that

$$\phi(\sigma) = \prod_{k \in [\sigma]} c_k, \quad \sigma \in S(\infty).$$

That is, the function ϕ is multiplicative in the cycles of a permutation and, hence, by Proposition 2, ϕ is indecomposable. Therefore, it has the form ϕ^μ , for some Thoma measure μ . Since for any $g \in S(\infty)$ the intersection

$$S(\infty) \cap K^E g K^E$$

is the conjugacy class of g in the group $S(\infty)$, every function ϕ^μ has a unique K^E -biinvariant extension to the group G^E . We denote this extension by $\tilde{\phi}^\mu$.

To summarize, the spherical functions of the pair (G^E, K^E) are precisely those functions $\tilde{\phi}^\mu$ that are positive definite on G^E . The description of this set is given by the following

Theorem 4. *The function $\tilde{\phi}^\mu$ is a spherical function of the pair (G^E, K^E) if and only if*

$$\frac{\mu(x)}{|x|} \in 2\mathbb{Z}_+$$

for every $x \in [-1, 0)$.

Proof. The sufficiency of the condition follows from the explicit constructions of representations (see Chapter 3). Let us prove the necessity.

Let π be the spherical representation of the pair (G^E, K^E) corresponding to the spherical function $\tilde{\phi}^\mu$. Let ξ be the spherical vector of the representation π . Take some $x \in [-1, 0)$. In the same way as we already did it in the proof of Theorem 2, by replacing the vector ξ by the vectors

$$\xi^{(m)} = \prod_{i=1}^m \delta_x(A_i) \xi, \quad m = 1, 2, 3, \dots,$$

one can effectively reduce the case of a general Thoma measure μ to that of Thoma measure supported at a single point x . Thus, we can assume that

$$\text{supp } \mu = \{x\}, \quad \mu(x) = 1.$$

In this case the value of the function $\tilde{\phi}^\mu$ at the element $g \in G^E(n)$ is

$$\tilde{\phi}^\mu(g) = x^{n-\ell(g)}.$$

Consider the non-negative expression

$$\sum_{g \in G^E(n)} \tilde{\phi}^\mu(g) = \sum_{g \in G^E(n)} x^{n-\ell(g)}.$$

By the Corollary to Proposition 7 it equals

$$\begin{aligned} n! 2^n x^n \frac{1}{x} \left(\frac{1}{x} + 2 \right) \left(\frac{1}{x} + 4 \right) \cdots \left(\frac{1}{x} + 2n - 2 \right) = \\ = n! 2^n 1(1 + 2x)(1 + 4x) \cdots (1 + (2n - 2)x). \end{aligned}$$

Since $x < 0$, this product can be non-negative for all n only if it terminates, which happens if

$$x = -\frac{1}{2k}$$

for some $k \in \mathbb{N}$. That is,

$$\nu(x) = 2k. \quad \square$$

This result is parallel to the following well known result from representation theory of finite symmetric groups. Namely, an irreducible admissible representation of the group $G^E(n)$ corresponding to a Young diagram λ , $|\lambda| = 2n$, contains a $K^E(n)$ -invariant vector if and only if all the parts of λ are even [28, §1.8, Example 6].

2.8 Classification of irreducible admissible representations of the pairs (G^O, K^O) , (G^E, K^E) .

Let π be an irreducible admissible representation of depth d of the Gelfand pair (G^E, K^E) . The case of the pair (G^O, K^O) is entirely analogous. The representation π is determined by a Young distribution Λ , such that $|\Lambda| = 2d$, and a Thoma measure μ . By the results of the previous section, $\mu(x)/|x| \in 2\mathbb{Z}_+$ for all $x < 0$.

The proof of the next proposition is a word for word copy of that of Proposition 3 in Section 2.2.

Proposition 8. $\text{supp } \Lambda \subset \text{supp } \mu \cup \{0\}$.

The description of irreducible admissible representations of the pair (G^E, K^E) is provided by the following theorem.

Theorem 5. *An irreducible admissible representation of the pair (G^E, K^E) (respectively, of the pair (G^O, K^O)) with depth d , Thoma measure μ , and Young distribution Λ , where $|\Lambda| = 2d$ (resp. $2d + 1$), exists if and only if $\mu(x)/|x| \in 2\mathbb{Z}_+$ for all $x < 0$ and*

$$\begin{aligned} \Lambda'(x)_1 + \Lambda'(x)_2 &\leq \mu(x)/|x|, & x > 0, \\ \Lambda(x)_1 &\leq \mu(x)/2|x|, & x < 0 \end{aligned}$$

for all $x \in [-1, 1]$.

Proof. The sufficiency follows from the explicit construction of representations [32] to be discussed in Section 3.

Let us prove the necessity. Suppose $x > 0$ and set

$$l_1 = \Lambda'(x)_1, \quad l_2 = \Lambda'(x)_2.$$

Consider the main case when $l_1, l_2 > 0$. Other cases are similar. Denote by $S(l_1) \times S(l_2)$ the subgroup in the group $G^E(d)$ which permutes the numbers $\{d - l_1 + 1, \dots, d\}$ and $\{-d, \dots, -d + l_2 - 1\}$.

By the branching rule for representations of finite symmetric groups and the definition of Λ , there exists a vector $\zeta \in R(\pi)$ which is antiinvariant under the action of the group $S(l_1) \times S(l_2)$ and satisfies

$$A_i \zeta = x \zeta, \quad i = d - l_1 + 1, \dots, d, -d, \dots, -d + l_2 - 1.$$

Then the vector

$$\zeta^{(x)} = \delta_x(A_{d+1})\zeta$$

is also $S(l_1) \times S(l_2)$ -antiinvariant.

Let Alt be the antisymmetrization over the group $S(l_1 + 1) \times S(l_2 + 1)$ which permutes the numbers $\{d - l_1 + 1, \dots, d, d + 1\}$ and $\{-d - 1, -d, \dots, -d + l_2 - 1\}$. The function

$$(\text{sgn}(g) g \zeta^{(x)}, \zeta^{(x)})$$

is $S(l_1) \times S(l_2)$ -biinvariant. Denote by s and t the permutations $(d, d + 1), (-d, -d - 1) \in G^E$. Arguing as in the proof of Lemma 4 we obtain

$$\begin{aligned} (\zeta^{(x)}, \zeta^{(x)}) &= \mu(x) \\ (s \zeta^{(x)}, \zeta^{(x)}) &= x \\ (t \zeta^{(x)}, \zeta^{(x)}) &= x \\ (st \zeta^{(x)}, \zeta^{(x)}) &= 0. \end{aligned} \tag{2.10}$$

The group $S(l_1 + 1) \times S(l_2 + 1)$ consists of four double cosets of the subgroup $S(l_1) \times S(l_2)$; their representatives are $1, s, t, st$ and the corresponding cardinalities are $l_1!l_2!, l_1l_1!l_2!, l_2l_1!l_2!, l_1l_2l_1!l_2!$. Therefore, by virtue of (2.10),

$$\begin{aligned} (\text{Alt } \zeta^{(x)}, \zeta^{(x)}) &= \frac{l_1!l_2!\mu(x) - l_1l_1!l_2!x - l_2l_1!l_2!x + l_1l_2l_1!l_2!0}{(l_1 + 1)!(l_2 + 1)!} \\ &= \frac{x(\mu(x)/x - l_1 - l_2)}{(l_1 + 1)(l_2 + 1)}. \end{aligned}$$

Since the result should be non-negative, we conclude that

$$l_1 + l_2 \leq \mu(x)/x.$$

Now suppose that $x < 0$ and set

$$l = \Lambda(x)_1.$$

Denote by $S(l)$ the subgroup in the group $G^E(d)$ which permutes the numbers $\{d - l + 1, \dots, d\}$.

Again, there exists a vector $\zeta \in R(\pi)$ invariant under the action of the group $S(l)$ and such that

$$A_i \zeta = x \zeta, \quad i = d - l + 1, \dots, d.$$

Then the vector

$$\zeta^{(x)} = \delta_x(A_{d+1})\zeta$$

is also $S(l)$ -invariant. The permutation $r = (d + 1, -d - 1)$ belongs to the group K_d^E . By the definition of the subspace $R(\pi)$, the vector $\zeta \in R(\pi)$ will be invariant under r , hence with respect to the group $S(l) \times S(2)$ which permutes the numbers $\{d - l + 1, \dots, d\}$ and $\{d + 1, -d - 1\}$. Let Sym be the symmetrization over the group $S(l + 2)$ which permutes the numbers $\{d - l + 1, \dots, d, d + 1, -d - 1\}$. Denote by s and t the permutations $(d, d + 1), (d - l + 1, -d - 1) \in G^E$. As usual,

$$\begin{aligned} (\zeta^{(x)}, \zeta^{(x)}) &= \mu(x) \\ (s\zeta^{(x)}, \zeta^{(x)}) &= x \\ (st\zeta^{(x)}, \zeta^{(x)}) &= 0. \end{aligned}$$

The group $S(l + 2)$ consists of three double cosets of $S(l) \times S(l_2)$; their representatives are $1, s, st$ and the cardinalities are $2l!, 4l!l!, l(l - 1)l!$. Therefore,

$$\begin{aligned} (\text{Sym } \zeta^{(x)}, \zeta^{(x)}) &= \frac{2l!\mu(x) + 4l!l!x + l(l - 1)l!0}{(l + 2)!} \\ &= \frac{2|x|(\mu(x)/|x| - 2l)}{(l + 1)(l + 2)}. \end{aligned}$$

The result should be non-negative, whence

$$l \leq \mu(x)/2|x|. \quad \square$$

Remark. The cases $x > 0$ and $x < 0$ for the pairs (G^E, K^E) and (G^O, K^O) are not symmetric. This is because the function sgn on the group $S(\infty)$ cannot be extended to a K^E -invariant positive definite function on the group G^E .

3. CONSTRUCTION OF REPRESENTATIONS

The main object considered in this section is a certain operation on admissible representations which we call *mixing* the representations. This construction is very much parallel to the Olshanski's construction in [32] and only slightly more general (see also [6,56]). It yields an explicit construction of actually all irreducible admissible representations whereas the methods of [32] produce only an open subset in of the admissible dual. The mixture of representations is, essentially, a special sort of an induced representation as we shall see in Section 3.2.

In Sections 3.1–3.3 we shall deal with the pairs G^E and G^O (mainly with G^E). We shall comment briefly on the case of G^D in Section 3.4.

3.1 Mixtures of representations.

Let π_1 and π_2 be two admissible representations of the pair (G^E, K^E) . Let p_1 and p_2 be two numbers, such that $p_1 > 0$, $p_2 > 0$, $p_1 + p_2 = 1$. We shall define the *mixture* of representations π_1 and π_2 with the weights p_1 and p_2 . One can similarly define the mixture of admissible representation of the pair (G^E, K^E) with an admissible representation of the pair (G^O, K^O) or the mixture of two admissible representations of the pair (G^O, K^O) . Set $d_i = d(\pi_i)$, $i = 1, 2$.

In the set of all functions $f : \mathbb{Z} \setminus \{0\} \rightarrow \{1, 2\}$ consider the following subset X ,

$$X = \{f | f(i) = f(-i) \text{ for almost all } i\},$$

where “almost all” means “all but finitely many”. The set X is a union of an increasing sequence of subsets

$$X_0 \subset X_1 \subset X_2 \subset \dots$$

where

$$X_n = \{f \in X | f(i) = f(-i), |i| > n\}.$$

The map

$$X_n \ni f \mapsto (f(-n), \dots, f(-1), f(1), \dots) \in \{1, 2\}^\infty$$

is a bijection. We transfer the product topology from $\{1, 2\}^\infty$ to X_n via this map and endow the set X with the direct limit topology. Consider the following measure ω_{p_1, p_2} on X ,

$$\omega_{p_1, p_2}(\{f | f(i) = f_i, |i| \leq n, f(i) = f(-i), i > n\}) = \prod_{|i| < n} p_{f_i}^{1/2}.$$

On each set X_n the measure ω_{p_1, p_2} is finite. The group G^E acts on X and preserves the measure ω_{p_1, p_2} .

It follows from the definition of the space X that for every $f \in X$ the parity of the number $|f^{-1}(1) \cap \{-N, \dots, N\}|$ stabilizes as $N \rightarrow \infty$. Consider the subset $Y \subset X$,

$$Y = \{f, |f^{-1}(1) \cap \{-N, \dots, N\}| \in 2\mathbb{Z}, N \gg 0, |f^{-1}(i)| = \infty, i = 1, 2\}.$$

This subset is measurable and $\omega_{p_1, p_2}(Y) > 0$, since the condition $|f^{-1}(i)| = \infty$, $i = 1, 2$ means deletion of a countable set of zero measure. It is also clear that this subset is G^E -invariant. We set $Y_n = Y \cap X_n$.

For every $f \in Y$ there exist unique bijections $\eta_i : f^{-1}(i) \rightarrow \mathbb{Z} \setminus \{0\}$, such that $\eta_i(-a) = -\eta_i(a)$ for almost all a and $a < b$ if and only if $\eta_i(a) < \eta_i(b)$. Define a cocycle

$$c : G^E \times Y \rightarrow G^E \times G^E$$

on the generators $(i, i+1)$ of the group G^E by the formula

$$c((i, i+1), f) = \begin{cases} (e, e), & f(i) \neq f(i+1) \\ ((\eta_1(i), \eta_1(i+1)), e), & f(i) = f(i+1) = 1 \\ (e, (\eta_2(i), \eta_2(i+1))), & f(i) = f(i+1) = 2. \end{cases}$$

Denote by H the Hilbert space of maps

$$F : Y \rightarrow H(\pi_1) \otimes H(\pi_2)$$

with inner product

$$(F_1, F_2)_H = \int_Y (F_1(f), F_2(f))_{H(\pi_1) \otimes H(\pi_2)} \omega_{p_1, p_2}(df).$$

Define a representation of the group G^E in the space H by the formula

$$[g \cdot F](f) = \pi_1 \otimes \pi_2 (c(g, g^{-1} \cdot f)) F(g^{-1} \cdot f).$$

This representation is unitary.

Now, our next goal is to compute the subspaces of invariants H_n . Fix two numbers $a, b \in \mathbb{Z}_+$. Set $K = 2a + 2b$ and define

$$\begin{aligned} Y_{a,b} = \{f \in Y, & |f^{-1}(1) \cap \{-K, \dots, K\}| = 2a \\ & |f^{-1}(2) \cap \{-K, \dots, K\}| = 2b \\ & f(i) = f(-i), |i| > K\}. \end{aligned}$$

Let D range over all subsets of cardinality $2a$ of the set $\{-K, \dots, K\} \setminus \{0\}$. Set

$$Y_{a,b,D} = \{f \in Y_{a,b} | f^{-1}(1) \cap \{-K, \dots, K\} = D\}.$$

Denote by $\text{supp } F$ the complement to the largest open subset where F is equal to zero almost everywhere. Denote by $H(Y_{a,b})$ the space of maps F , such that $\text{supp } F \subset Y_{a,b}$. Denote by $H_{a,b} \subset H(Y_{a,b})$ the subspace of maps with the support in $Y_{a,b}$, which are constant on all of $Y_{a,b,D}$ and take the values in $H(\pi_1)_a \otimes H(\pi_2)_b$.

Proposition 9.

$$H_n = \bigoplus_{a+b=n} H_{a,b}.$$

Proof. The inclusion $\bigoplus_{a+b=n} H_{a,b} \subset H_n$ is obvious. Let us check the inverse inclusion. Assume that $F \in H_n$, and let the numbers r, s be such that $s \geq r > n$. Set

$$W_{r,s} = \{f \in Y \mid f(r) \neq f(-r), f(i) = f(-i), |i| > s\}.$$

The sets $W_{r,s}$ are open. Let l_1, l_2 be two distinct integers, such that $l_1, l_2 > s$. Consider the action of permutations $(r, l_1)(-r, -l_1) \in K_n^E$ on the sets $W_{r,s}$. We have

$$(r, l_1)(-r, -l_1) \cdot W_{r,s} \cap (r, l_2)(-r, -l_2) \cdot W_{r,s} = \emptyset$$

$$\omega_{p_1, p_2}((r, l_1)(-r, -l_1) \cdot W_{r,s}) = \omega_{p_1, p_2}((r, l_2)(-r, -l_2) \cdot W_{r,s}).$$

Since the map F is square summable, it should vanish almost everywhere on each set $W_{r,s}$. This means that

$$\text{supp } F \subset Y \setminus \bigcup_{n < r \leq s} W_{r,s} = \bigcup_{a+b=n} Y_{a,b}.$$

In other words, $F \in \bigoplus_{a+b=n} H(Y_{a,b})$.

All the subspaces $H(Y_{a,b})$ are invariant with respect to K_n^E . The subspaces of invariants are always consistent with decompositions in a direct sum, hence

$$F \in \bigoplus_{a+b=n} (H(Y_{a,b}))_n.$$

Let $F \in (H(Y_{a,b}))_n$ for some a, b . For every $g \in K_n^E$ the equality $[g \cdot F](f) = F(f)$ holds for almost all f with respect to the measure ω_{p_1, p_2} . Since the group K_n^E is countable, for almost all f the equality $[g \cdot F](f) = F(f)$ holds for all $g \in K_n^E$.

Consider the stabilizer $\text{Stab}(f) \subset K_n^E$ of a point $f \in Y_{a,b}$. The image of the group $\text{Stab}(f)$ under the map $g \mapsto c(g, f)$ is the subgroup $K_a^E \times K_b^E \subset K^E \times K^E$. Therefore, for almost all $f \in Y_{a,b}$ we have $F(f) \in H(\pi_1)_a \otimes H(\pi_2)_b$.

For every $\zeta \in H(\pi_1)_a \otimes H(\pi_2)_b$ the set $F^{-1}(\zeta)$ is a measurable and a K_n^E -invariant mod 0 subset. Consider an arbitrary set $Y_{a,b,D}$. The action of the group K_n^E on the set $Y_{a,b,D}$ is clearly isomorphic to the action of the symmetric group by permutations of factors on $\{1, 2\}^\infty$ with a Bernoulli measure. The ergodicity of this latter action is well known. Hence the set $F^{-1}(\zeta)$ is, up to a subset of measure zero, a union of the sets $Y_{a,b,D}$.

Therefore, $(H(Y_{a,b}))_n = H_{a,b}$. \square

Corollary. *The representation of the group G^E in the space H is admissible. Its depth is $d = d_1 + d_2$ and*

$$H_d = H_{d_1, d_2},$$

$$\dim H_d = \binom{2d}{2d_1} \dim H(\pi_1)_{d_1} \dim H(\pi_2)_{d_2}.$$

Definition. Let π be the representation of the group G^E in the cyclic span of H_{d_1, d_2} . We call the representation π the *mixture* of representations π_1, π_2 with the weights p_1, p_2 .

Proposition 10. *Suppose that the representations π_1, π_2 are irreducible and that $\mu_i, i = 1, 2$ are their Thoma measures. Then the representation of the pair (G_d^E, K_d^E) in the cyclic span of $R(\pi)$ is a multiple of the spherical representation with the Thoma measure*

$$\mu(x) = p_1 \mu_1(x/p_1) + p_2 \mu_2(x/p_2) .$$

Proof. We have to prove that

$$\pi \left(\prod_i C_i \right) = \prod_i \pi(C_i)$$

and

$$\pi(C_k) = p_1^k \pi_1(C_k) + p_2^k \pi_2(C_k) .$$

Let us prove the second equality. The proof of the first one is similar. Denote by $z_k \in G_d^E$ the permutation

$$z_k = (d+1, \dots, d+k) .$$

Choose a vector $\xi \in H(\pi_1)_{d_1} \otimes H(\pi_2)_{d_2}$, $\|\xi\| = 1$. Consider the following map $F \in R(\pi)$,

$$F(f) = \begin{cases} (\omega_{p_1, p_2}(Y_{d_1, d_2}))^{-1} \xi, & f \in Y_{d_1, d_2} \\ 0, & \text{otherwise} . \end{cases}$$

It is clear that $\|F\| = 1$. Consider two subsets

$$^{(1)}Y_{d_1, d_2} = \{f \in Y_{d_1, d_2} | f(i) = 1, |i| = d+1, \dots, d+k, \}$$

$$^{(2)}Y_{d_1, d_2} = \{f \in Y_{d_1, d_2} | f(i) = 2, |i| = d+1, \dots, d+k, \} .$$

in Y_{d_1, d_2} . Clearly, the two inclusions $f \in Y_{d_1, d_2}$ and $z_k \cdot f \in Y_{d_1, d_2}$ occur simultaneously only if $f \in ^{(1)}Y_{d_1, d_2}$ or $f \in ^{(2)}Y_{d_1, d_2}$. Therefore,

$$\begin{aligned} \pi(C_k) &= (\pi(z_k)F, F)_H \\ &= \frac{1}{\omega_{p_1, p_2}(Y_{d_1, d_2})} \left[\pi_1(C_k) \omega_{p_1, p_2} \left(^{(1)}Y_{d_1, d_2} \right) + \pi_2(C_k) \omega_{p_1, p_2} \left(^{(2)}Y_{d_1, d_2} \right) \right] \\ &= \pi_1(C_k) p_1^k + \pi_2(C_k) p_2^k . \quad \square \end{aligned}$$

Now consider the action of the operators $A_i, |i| = 1, \dots, d$ in the space $R(\pi)$. Set $m_i = \dim R(\pi_i)$, $i = 1, 2$. Let $\{\zeta_1^{(i)}, \dots, \zeta_{m_i}^{(i)}\} \in R(\pi_i)$ be the eigenbases of the operators $A_k, |k| = 1, \dots, d$. Let D run over the subsets of cardinality $2d_1$ in the set $\{-d, \dots, d\} \setminus 0$. Then the maps $F_{i, j, D}, i = 1, \dots, m_1, j = 1, \dots, m_2$, where

$$F_{i, j, D}(f) = \begin{cases} \zeta_i^{(1)} \otimes \zeta_j^{(2)}, & f \in Y_{d_1, d_2, D} \\ 0, & \text{otherwise} \end{cases}$$

form a basis in the space $R(\pi)$. This basis consists of eigenvectors of the operators A_k . Indeed, let P be the orthogonal projector of the space H onto the subspace $R(\pi)$. Then by (1.5), for every $F \in R(\pi)$ and every $k, |k| = 1, \dots, n$, we have the following equality

$$\pi(A_k) \cdot F = P\pi((k, d+1)) \cdot F.$$

Therefore

$$\pi(A_k) \cdot F_{i,j,D}(f) = \begin{cases} p_1 \pi_1(A_{\eta_1(k)}) \zeta_i^{(1)} \otimes \zeta^{(2)}, & f \in Y_{d_1, d_2, D}, k \in D \\ p_2 \zeta_i^{(1)} \otimes \pi_2(A_{\eta_2(k)}) \zeta^{(2)}, & f \in Y_{d_1, d_2, D}, k \notin D \\ 0, & \text{otherwise.} \end{cases}$$

Let Λ_1 and Λ_2 denote the Young distributions of representations π_1 and π_2 . Let T_{Λ_i} , $i = 1, 2$, denote the corresponding representations of the semigroups

$$S(d_i) \ltimes \mathbb{Z}_+^{d_i}$$

in the spaces $R(\pi_i)$. Denote by $\Lambda(\cdot/p_1)$ the Young distribution which equals $\Lambda(x/p_1)$ at the point x . Then the above formulas for the action of operators A_i imply that the representation of the semigroup $S(d) \ltimes \mathbb{Z}_+^d$ in the space $R(\pi)$ is

$$\text{Ind}_{(S(d_1) \ltimes \mathbb{Z}_+^{d_1}) \times (S(d_2) \ltimes \mathbb{Z}_+^{d_2})}^{S(d) \ltimes \mathbb{Z}_+^d} T_{\Lambda_1(\cdot/p_1)} \otimes T_{\Lambda_2(\cdot/p_2)}.$$

This representation is irreducible if and only if $\text{supp } \Lambda_1(\cdot/p_1) \cap \text{supp } \Lambda_2(\cdot/p_2) = \emptyset$. In this case the representation π is also irreducible and has the Young distribution Λ equal to $\Lambda(x) = \Lambda_1(x/p_1) \cup \Lambda_2(x/p_2)$.

We summarize this discussion as follows:

Theorem 6. *Let π_1 and π_2 be two irreducible admissible representations of the pair (G^E, K^E) with Thoma measures μ_i , $i = 1, 2$, and with Young distributions Λ_i , $i = 1, 2$. Let the representation π be the mixture of representations π_1 and π_2 with the weights p_1 and p_2 , where $p_1 > 0$, $p_2 > 0$, $p_1 + p_2 = 1$.*

The representation π is admissible. It is a sum of irreducible admissible representations with Thoma measure

$$\mu(x) = p_1 \mu_1(x/p_1) + p_2 \mu_2(x/p_2).$$

It is irreducible if and only if

$$\text{supp } \Lambda_1(\cdot/p_1) \cap \text{supp } \Lambda_2(\cdot/p_2) = \emptyset.$$

In this case it has the Young distribution Λ , where

$$\Lambda(x) = \Lambda_1(x/p_1) \cup \Lambda_2(x/p_2).$$

The mixture of any finite number of representations may be defined in the same way. Moreover, one can define a mixture of finitely many admissible representations

and countably many spherical representations π_i , $i = 1, \dots$ of the pair (G^E, K^E) . In this case $\bigotimes_1^\infty H(\pi_i)$ denotes the direct limit of Hilbert spaces

$$\lim \operatorname{ind} \bigotimes_1^N H(\pi_i)$$

with respect to the inclusions

$$\bigotimes_1^N H(\pi_i) \rightarrow \bigotimes_1^N H(\pi_i) \otimes \xi_{N+1},$$

where $\xi_i \in H(\pi_i)$ is the spherical vector of the representation π_i . In this case Theorem 6 can be generalized as follows.

Theorem 7. *Let π_i , $i = 1, 2, \dots$, be irreducible admissible representations of the pair (G^E, K^E) with Thoma measures μ_i , $i = 1, 2, \dots$, and Young distributions Λ_i , $i = 1, 2, \dots$, such that the sum*

$$\sum_i |\Lambda_i| < \infty$$

is finite. Define the representation π as the mixture of representations π_i , $i = 1, 2, \dots$, with the weights p_i , $i = 1, 2, \dots$, where $p_i > 0$, $\sum_i p_i = 1$.

The representation π is admissible. It is a sum of irreducible admissible representations with Thoma measure

$$\mu(x) = \sum_i p_i \mu_i(x/p_i).$$

It is irreducible if and only if

$$\operatorname{supp} \Lambda_i(\cdot/p_i) \cap \operatorname{supp} \Lambda_j(\cdot/p_j) = \emptyset, \quad i \neq j.$$

In this case it has the following Young distribution:

$$\Lambda(x) = \bigcup_i \Lambda_i(x/p_i).$$

3.2 Mixtures and induction.

In this Section we show that the operation of mixing the representations is intimately related to that of inducing of representations. In the group of all (not necessarily finite) bijections $g : \mathbb{Z} \setminus \{0\} \rightarrow \mathbb{Z} \setminus \{0\}$ consider the subgroups

$$\overline{K}_n^E = \{g | g(i) = -g(-i), g(i) = i, i > n\}, n = 0, 1, \dots$$

As usual, write $\overline{K}^E = \overline{K}_0^E$. Set

$$\overline{G}^E = G^E \cdot \overline{K}^E.$$

Define a topology in the group \overline{G}^E in which a fundamental neighborhood system of unity is formed by the subgroups \overline{K}_n^E , $n = 0, 1, \dots$. In this topology the group G^E is a dense subgroup of the group \overline{G}^E . The representations of the group G^E which admit a continuous extensions onto the group \overline{G}^E are exactly the admissible representations of the pair (G^E, K^E) [32].

A mixture of representations is an induced representation in the following sense. The space Y is a homogeneous space of the group \overline{G}^E . The stabilizer of a point is isomorphic to the group $\overline{G}^E \times \overline{G}^E$. The cocycle $c(\cdot, \cdot)$ is exactly the usual cocycle on the homogeneous space. We have the following

Proposition 11. *The measures ω_{p_1, p_2} are, up to a factor, exactly the measures on Y which are*

- (1) *invariant with respect to the action of the group G^E ;*
- (2) *finite on all the sets Y_n , $n = 0, 1, \dots$;*
- (3) *extreme in the class of measures with the properties (1)–(2).*

Proof. Consider the set Y_k . For every function $f \in Y_k$ there is an element $g \in G^E(k)$ such that $g \cdot f \in Y_0$. In other words, $Y_k \subset \bigcup_{g \in G^E(k)} g \cdot Y_0$. This means that a G^E -invariant measure on Y is determined by its restriction to Y_0 .

Let ν be a measure satisfying the conditions (1)–(3). If $\nu(Y_0) = 0$, then also $\nu(Y) = 0$. Therefore we can assume that $\nu(Y_0) = 1$. The set Y_0 is embedded in $X_0 \cong \{1, 2\}^\infty$. The group K^E acts on X_0 by permutations of the factors. It follows from the de Finetti theorem [53, p. 256] that the measure ν on X_0 has the form ω_{p_1, p_2} for some $p_1, p_2 \geq 0$, $p_1 + p_2 = 1$. It is also clear that $\nu(Y_0) \neq 0$ if $p_1 > 0$ and $p_2 > 0$. \square

Let us now consider the G^E -invariant measures supported on $\overline{Y} \setminus Y$, where \overline{Y} is the closure of the set Y in the space X :

$$\overline{Y} = \{f, |f^{-1}(1) \cap \{-N, \dots, N\}| \text{ is even for almost all } N\}.$$

The set \overline{Y} coincides with the orbit of the set X_0 under the action of the group G^E . The set $\overline{Y} \setminus Y$ consists of a countable number of G^E -orbits $Z_{2k}^{(i)}$, $i = 1, 2$, $k = 0, 1, \dots$

$$Z_r^{(i)} = \{f \in X, |f^{-1}(i)| = r\}.$$

It suffices to consider the sets $Z_{2k}^{(1)}$. The set $Z_{2k}^{(1)}$ is a single orbit of a countable group G^E . It supports a unique, up to a factor, G^E -invariant measure ν_{2k} which is just the counting measure. The stabilizer of a point in $Z_{2k}^{(1)}$ is isomorphic to the group $G^E(k) \times G_k^E \cong G^E(k) \times G^E$.

Let ρ be an irreducible representation of the group $G^E(d_1)$, and π_2 be an irreducible admissible representation of the group $G_{d_2}^E \cong G^E$. Let μ_2, Λ_2 be the Thoma measure and the Young distribution corresponding to the representation π_2 . The representation π ,

$$\pi = \text{Ind}_{G^E(d_1) \times G_{d_2}^E}^{G^E} \rho \otimes \pi_2,$$

can be realized in the space of maps $F : X \rightarrow H(\rho) \otimes H(\pi_2)$ which are square summable with respect to the measure ν_{2d_1} .

Assume that $\Lambda_2(0) = \emptyset$. Then, the same argument we used for mixtures yields that the representation π is admissible and irreducible. Its Thoma measure is

$$\mu = \mu_2,$$

and the Young distribution is

$$\Lambda(x) = \begin{cases} \Lambda_2(x), & x \neq 0 \\ \rho, & x = 0. \end{cases}$$

This conventional induction of representations may be considered as a limit case of mixtures. Recall the definition of the topology in the space of unitary representations of a discrete group G [23]. Let T_0 be a unitary representation. Given a finite subset $M \in G$ and an array of vectors $\xi_1, \dots, \xi_k \in H(T_0)$, we denote by $U(T_0, M, \xi_1, \dots, \xi_k; \varepsilon)$ the set of unitary representations T of the group G , for which the corresponding space contains the vectors ζ_1, \dots, ζ_n such that

$$|(T(g)\zeta_i, \zeta_j) - (T_0(g)\xi_i, \xi_j)| < \varepsilon, \quad g \in M, i, j = 1, \dots, n.$$

The sets $U(T_0, M, \xi_1, \dots, \xi_k; \varepsilon)$ form a neighborhood base of the representation T_0 . If the representation T_0 is irreducible then in order to check the convergence $T_n \rightarrow T_0$ it suffices to check that a certain matrix element $(T_0(\cdot)\xi, \xi), \xi \in H(T_0)$ can be approximated by the matrix elements of representations T_n .

Let π_1 be an irreducible admissible representation of the group G^E with the Young distribution Λ_1 such that $\text{supp } \Lambda_1$ is a one point set $\{y\}$, $y \in [-1, 1]$ and $\Lambda_1(y) = \rho$. Denote by $\pi(p_1, p_2)$ the mixture of representations π_1, π_2 with the weights p_1, p_2 .

Proposition 12.

$$\pi(p_1, p_2) \rightarrow \pi \text{ as } p_1 \rightarrow 0.$$

Proof. It follows from Theorem 6 that, as $p_1 \rightarrow 0$, the representations of the semi-group $\Gamma(d)$ in the spaces $R(\pi(p_1, p_2))$ converge to the representation of this semi-group in $R(\pi)$. Hence, the matrix coefficients of representations $\pi(p_1, p_2)$ corresponding to vectors in subspaces $R(\pi(p_1, p_2))$ converge to the matrix coefficients of the representation π . Since the representation π is irreducible, the proposition follows. \square

The indecomposable invariant measures on X supported by the set $X \setminus \overline{Y}$ have the following meaning. The restriction of a measure ω_{p_1, p_2} onto $X \setminus \overline{Y}$ corresponds to the representations of the group G^E which are the mixtures of two representations of the group G^O with the weights p_1, p_2 . The measures ν_{2k+1} correspond to representations of the group G^E which are induced from a subgroup isomorphic to $G^O(k) \times G^O$.

3.3 Elementary representations.

Let π be an irreducible admissible representation of G^E or G^O with Thoma measure μ and Young distribution Λ . Call this representation *elementary* if

$$\text{supp } \mu = \text{supp } \Lambda = \{y\}$$

for some point $y \in [-1, 1]$. It follows from the classification of irreducible representations and the results of the two previous sections that in order to construct all irreducible admissible representation it remains to give a construction of elementary ones.

We shall briefly describe the realization of elementary representations of the groups G^E and G^O obtained in [32]. One has to distinguish between three cases: $y = 0$, $y > 0$ and $y < 0$.

Suppose $y = 0$. Then the corresponding elementary representation is

$$\mathrm{Ind}_{G(d) \times K_d}^G \Lambda(0) \otimes 1.$$

Suppose $y > 0$. By Theorem 4, y has to be of the form $y = 1/n$, $n \in \mathbb{N}$. Consider the space \mathbb{C}^n with the standard basis $\{e_1, \dots, e_n\}$ and set, by definition,

$$\xi = n^{-1/2} \sum_{i=1}^n e_i \otimes e_i \in \mathbb{C}^n \otimes \mathbb{C}^n.$$

Consider the direct limit

$$H = \lim \operatorname{ind} \bigotimes_1^{2N} \mathbb{C}^n$$

of Hilbert spaces with respect to inclusions

$$\bigotimes_1^{2N} \mathbb{C}^n \rightarrow \bigotimes_1^{2N} \mathbb{C}^n \otimes \xi.$$

Using the bijection $\mathbb{Z} \setminus \{0\} \rightarrow \mathbb{N}$,

$$i \mapsto \begin{cases} 2i - 1, & i > 0 \\ -2i, & i < 0, \end{cases}$$

we can define the action of the group G^E in the space H . (In case of G^O one should consider $\lim \operatorname{ind} \bigotimes_0^{2N} \mathbb{C}^n$.)

The group of orthogonal matrices $O(n)$ preserves the vector ξ , hence its action on the space H is well defined. As explained in [54], the irreducible representations of the group $O(n)$ are labeled by Young diagrams λ such that

$$(\lambda)_1' + (\lambda)_2' \leq n.$$

One has the following:

Theorem (Olshanski, [32]).

- (1) *The representation of the group G^E in the space H is admissible.*
- (2) *The representations of the groups G^E and $O(n)$ generate the commutant of each other.*
- (3) *The space H , as an $G^E \times O(n)$ -module, decomposes into the following direct sum*

$$H = \bigoplus_{\lambda, (\lambda)_1' + (\lambda)_2' \leq n} \pi_{1/n, \lambda} \otimes T_\lambda,$$

where T_λ is the representation of the group $O(n)$ corresponding to a diagram λ , and $\pi_{1/n, \lambda}$ is the irreducible admissible representation of the group G^E , such that $\operatorname{supp} \mu = \operatorname{supp} \Lambda = \{1/n\}$, $\Lambda(1/n) = \lambda$.

There is a little inaccuracy in the paper [32] in case of $y < 0$, indicated by G. Olshanski. The correct construction of representations is as follows. By Theorem 4, in case of $y < 0$ we are forced to take $y = -1/2n$, $n \in \mathbb{N}$. Consider a basis $\{e_1, \dots, e_{2n}\}$ in the space \mathbb{C}^{2n} . Consider the vector

$$\xi = (2n)^{-1/2} \sum_{i=1}^n (e_i \otimes e_{i+n} - e_{n+i} \otimes e_i)$$

in the space $\mathbb{C}^{2n} \otimes \mathbb{C}^{2n}$ and let

$$H = \lim_{\text{ind}} \bigotimes_1^{2N} \mathbb{C}^{2n}$$

be the direct limit of Hilbert spaces with respect to inclusions

$$\bigotimes_1^{2N} \mathbb{C}^{2n} \rightarrow \bigotimes_1^{2N} \mathbb{C}^{2n} \otimes \xi.$$

Define the representation of the group G^E in the space H as the tensor product of the representation by permutations of tensor factors, and the one dimensional representation sgn .

The group $Sp(n)$ of symplectic matrices preserves the vector ξ , hence its action on the space H is well defined. Again, as explained in [54], the representations of the group $Sp(n)$ are labeled by Young diagrams λ , such that

$$\lambda_1 \leq n.$$

One has the following

Theorem (Olshanski).

- (1) *The representation of the group G^E in the space H is admissible.*
- (2) *The representations of groups G^E and $Sp(n)$ generate the commutant of each other.*
- (3) *The space H , as an $G^E \times Sp(n)$ -module, decomposes into the following direct sum*

$$H = \bigoplus_{\lambda, \lambda_1 \leq n} \pi_{-1/2n, \lambda} \otimes T_\lambda,$$

where T_λ is the representation of the group $Sp(n)$ corresponding to a diagram λ , and $\pi_{-1/2n, \lambda}$ is an irreducible admissible representation of the group G^E , such that $\text{supp } \mu = \text{supp } \Lambda = \{-1/2n\}$, $\Lambda(1/2n) = \lambda$.

3.4 Mixtures in the case of G^D .

The definition of a mixture of representations can be easily extended to the case of the pair (G^D, K^D) . In fact, it is natural to consider the following “unbalanced” groups

$$G_{m_1, m_2}^D = \{g \in G^D | g(i) = i, -m_2 \leq i \leq m_1\},$$

where m_1, m_2 are integers, and define mixtures of representations of these groups. Since the inclusion $K_n^D \subset G_{m_1, m_2}^D$ is valid, given m_1, m_2 , for all n but finitely many, the definition of admissible representation also works for the groups G_{m_1, m_2}^D . All the theory of admissible representations can be transferred word-for-word to this “unbalanced” case. In particular, such representations are labeled by a Thoma measure and a pair of Young distributions Λ, M , for which it is now possible that $|\Lambda| \neq |M|$.

The construction of elementary representations can be taken from [32].

Remark that in the language of H -series from Section 2.4 the mixture of spherical representations corresponds to the product of H -series.

4. CONCLUDING REMARKS

This paper is the English version of the author’s PhD thesis (1995, Moscow State University). I did not try to update anything in it. For some related recent results the reader is referred to [57, 58] and references therein.

I am very grateful to my advisor A. A. Kirillov, A. M. Vershik, R. S. Ismagilov, S. V. Kerov, and Yu. A. Neretin for their constant interest, encouragement, and help. My very special thanks are due to G. Olshanski, who not only laid the foundations of the whole subject, thus making this paper possible in the first place, but also was of absolutely indispensable help to me from the very beginning of my work to the proofreading stage of the present paper.

REFERENCES

1. M. Aissen, I. J. Schoenberg, A. M. Whitney, *On the generating function of totally positive sequences I.*, J. Analyse Math. **2** (1952), 93–103.
2. M. Aissen, A. Edrei, I. J. Schoenberg, A. M. Whitney, *On the generating function of totally positive sequences*, Proc. Nat. Acad. Sci. USA **37** (1951), 303.
3. R. Boyer, *Infinite traces of AF-algebras and characters of $U(\infty)$* , J. Operator Theory **9** (1983), 205–236.
4. R. Brauer, *On algebras which are connected with semisimple continuous groups*, Annals of Math. **38** (1937), 857–872.
5. A. M. Vershik, *A description of invariant measures for actions of some infinite dimensional groups*, Soviet Math. Dokl. **15** (1974), 1396–1400.
6. A. M. Vershik, S. V. Kerov, *Characters and factor-representations of the infinite symmetric group*, Soviet Math. Dokl. **23**, (1981), no. 2, 389–392.
7. A. M. Vershik, S. V. Kerov, *Asymptotic character theory of the symmetric group*, Functional analysis and its applications **15** (1981), no. 4, 246–255.
8. A. M. Vershik, S. V. Kerov, *Characters and factor-representations of the infinite unitary group*, Soviet Math. Dokl. **26** (1982), no. 3, 570 – 574.
9. S. V. Kerov, A. M. Vershik, *Characters, factor representations and K -functor of the infinite symmetric group*, Operator Algebras and Group Representations, Monographs Stud. Math., 18, vol. 2, Pitman, Boston-London, 1984, pp. 23–32.
10. S. V. Kerov, A. M. Vershik, *The Grothendieck group of infinite symmetric group and symmetric functions (with elements of the theory of K_0 -functor of AF-algebras)*, Representation of Lie groups and related topics, Adv. Stud. Contemp. Math., vol. 7, Gordon and Breach, New York, 1990, pp. 39–117.
11. S. V. Kerov, A. M. Vershik, *The characters of the infinite symmetric group and probability properties of the Robinson-Schensted-Knuth algorithm*, SIAM J. Alg. Discr. Methods **7** (1986), no. 1, 116–124.

12. G. Olshanski, A. M. Vershik, *Ergodic unitarily invariant measures on the space of infinite Hermitian matrices*, Contemporary mathematical physics, AMS Transl. Ser. 2, vol. 175, Amer. Math. Soc., Providence, 1996, pp. 137–175.
13. D. Voiculescu, *Représentations factorielles de type II_1 de $U(\infty)$* , J. Math. pures et appl. **55** (1976), 1–20.
14. D. Voiculescu, *On extremal invariant functions of positive type on certain groups*, INCREST Preprint Series Math. (1978).
15. D. Voiculescu, *Sur les représentations factorielles finies de $U(\infty)$ et autres groupes semblables*, C. Rend. Acad. Sci. Paris **279** (1974), A945–A946.
16. F. P. Gantmakher, M. G. Krein, *Oscillation matrices and kernels and small oscillations of mechanical systems*, (Russian), Gosudarstv. Isdat. Tehn.-Teor. Lit., Moscow-Leningrad.
17. A. Edrei, *On the generating function of totally positive sequences II*, J. Analyse Math. **2** (1952), 104–109.
18. A. Edrei, *On the generating function of a doubly infinite, totally positive sequence*, Trans. Amer. Math. Soc. **47** (1953), 367–383.
19. R. S. Ismagilov, *On linear representations of matrix groups with elements from a normed field*, Math. USSR Izvestija **3** (1969), no. 6, 1219–1244.
20. R. S. Ismagilov, *Spherical functions over a normed field whose residue field is infinite*, Func. analysis and its application **4** (1970), no. 1, 42–51.
21. S. Karlin, *Total positivity and applications*, Stanford University Press, 1968.
22. S. V. Kerov, *A realization of representations of Brauer semigroup*, J. Soviet Math. **47** (1989), no. 2, 2503–2507.
23. A. A. Kirillov, *Introduction to the theory of representations and noncommutative harmonic analysis*, Representation theory and noncommutative harmonic analysis, I, Encyclopedia Math. Sci., vol. 22, Springer-Verlag, Berlin, 1994, pp. 1–156.
24. A. A. Kirillov, *Elements of the theory of representations*, Grundlehren der Mathematischen Wissenschaften, vol. 220, Springer-Verlag, Berlin–New York, 1976.
25. A. A. Kirillov, *Positive definite functions on a matrix group with the elements in a discrete field*, Soviet Math. Dokl. **6** (1965), no. 3, 707–709.
26. A. A. Kirillov, *Representations of the infinite dimensional unitary group*, Soviet Math. Dokl. **14** (1973), 1355–1358.
27. A. Lieberman, *The structure of certain unitary representations of infinite symmetric group*, Trans. Amer. Math. Soc. **164** (1972), 189–198.
28. I. G. Macdonald, *Symmetric functions and Hall polynomials*, Clarendon Press, Oxford, 1979.
29. M. L. Nazarov, Yu. A. Neretin, G. Olshanski, *Semigroupes engendrés par la représentation de Weil du groupe symplectique infinie*, Comptes Rendus Acad. Sci. Paris, Sér. I **309** (1989), no. 7, 443–446.
30. M. A. Naimark, *Normed rings*, Nauka, Moscow, 1968.
31. N. I. Nessonov, *The complete classification of representations of $GL(\infty)$ containing the identity representation of the unitary subgroup*, Math. USSR Sbornik **58** (1987), no. 1, 127–147.
32. G. Olshanski, *Unitary representations of (G, K) -pairs that are connected with the infinite symmetric group $S(\infty)$* , Leningrad Math. J. **1** (1990), no. 4, 983–1014.
33. G. Olshanski, *Unitary representations of infinite-dimensional pairs (G, K) and the formalism of R. Howe*, Adv. Stud. Contemp. Math. **7** (1990), Gordon and Breach, New York, 269–463.
34. G. Olshanski, *On semigroups related to infinite-dimensional groups*, Topics in representation theory, Adv. Soviet Math., vol. 2, Amer. Math. Soc., Providence, 1991, pp. 67–101.
35. G. Olshanski, *Unitary representations of infinite dimensional (G, K) -pairs and the formalism of R. Howe*, Soviet Math. Dokl. **27** (1983), no. 2, 290 – 294.
36. G. Olshanski, *The infinite dimensional classical groups of finite R -rank: a description of representations and asymptotic theory*, Functional Anal. Appl. **18** (1984), no. 1, 22–34.
37. G. Olshanski, *Unitary representations of the group $SO_0(\infty, \infty)$ as limits of unitary representations of the groups $SO_0(n, \infty)$ when $n \rightarrow \infty$* , Functional Anal. Appl. **20** (1986), no. 4, 292–301.
38. G. Olshanski, *The method of holomorphic extensions in the theory of unitary representations of infinite-dimensional classical groups*, Functional Anal. Appl. **22** (1988), no. 4, 273–285.

39. G. Olshanski, *Unitary representations of the infinite symmetric group: a semigroup approach*, Representations of Lie groups and Lie algebras, Academiai Kiadó, Budapest, 1985, pp. 181–197.
40. G. Olshanski, *New "large" groups of type I*, J. Soviet Math. **18** (1982), 22–39.
41. M. Reed, B. Simon, *Methods of modern mathematical physics*, Academic Press, NY, London, 1972.
42. I. Schoenberg, *Selected Papers*, vol. 1–2, Birkhäuser, Boston, Basel, 1988.
43. I. Schoenberg, *Über die variationvermindernde lineare Transformationen*, Math. Zeitschr. **32** (1930), 321–328.
44. S. V. Smirnov, *Positive definite functions on algebraically nilpotent groups over a discrete field*, Soviet Math. Dokl. **7** (1966), no. 5, 1240–41.
45. R. Stanley, *Enumerative Combinatorics*, vol. 1, Wadsworth & Brooks/Cole, Monterey, California, 1986.
46. J. R. Stembridge, *On Schur's Q -functions and the primitive idempotents of a commutative Hecke algebra*, J. Alg. Comb. **1** (1992), 71–96.
47. S. Stratila, D. Voiculescu, *Representations of AF -algebras and of the group $U(\infty)$* , Lect. Notes in Math. **186** (1975), Springer.
48. S. Stratila, D. Voiculescu, *A survey on representations of the unitary group $U(\infty)$* , Spectral Theory Banach Center Publications **8** (1982), 415–434.
49. E. Thoma, *Die unzerlegbaren, positiv-definiten Klassenfunktionen der abzählbar unendlichen symmetrischen Gruppe*, Math. Zeitschr. **85** (1964), no. 1, 40–61.
50. E. Thoma, *Über unitäre Darstellungen abzählbarer diskreter Gruppen*, Math. Annalen **153** (1964), 111–138.
51. E. Thoma, *Eine Charakterisierung diskreter Gruppen von Typ I*, Invent. Math. **6** (1968), 190–196.
52. E. Thoma, *Characters of infinite groups*, Operator Algebras and Group Representations, vol. 2 Pitman, 1984, pp. 211–216.
53. W. Feller, *An Introduction to Probability Theory and its Applications*, vol. 2, John Wiley & Sons, Inc., NY, London, Sydney, 1966.
54. H. Weyl, *The classical groups, their invariants and representations* (1939), Princeton University Press.
55. A. Okounkov, *The Thoma theorem and representations of the infinite bisymmetric group*, Funct. Anal. Appl. **28** (1994), no. 2, 100–107.
56. A. J. Wassermann, *Automorphic actions of compact groups on operator algebras*, PhD thesis, University of Pennsylvania (1981).
57. S. Kerov, A. Okounkov, and G. Olshanski, *The boundary of the Young graph with Jack edge multiplicities*, q-alg/9703037.
58. A. Okounkov and G. Olshanski, *Asymptotics of Jack polynomials as the number of variables goes to infinity*, q-alg/9709011.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 UNIVERSITY AVENUE,
CHICAGO, IL 60637–1546

E-mail address: okounkov@math.uchicago.edu