A q-ANALOGUE OF THE BEREZIN QUANTIZATION OF THE UNIT DISC

D. Shklyarov \diamond S. Sinel'shchikov * L. Vaksman * [†]

*Institute for Low Temperature Physics & Engineering National Academy of Sciences of Ukraine

> [°]Kharkov State University Department of Mechanics and Mathematics

It is well known [8, 3] that the Berezin quantization can be used to construct a formal deformation of the algebra $C^{\infty}(U)$ of smooth functions in the unit disc $U \subset \mathbb{C}$.

In the well known work [7] a family of q-analogues of the unit disc $U \subset \mathbb{C}$ was introduced. The simplest, and hence the most important one among those was also considered in [9, 6, 4]. With these results as a background, we produce an explicit formula for a formal deformation of the quantum disc. To develop it, a Berezin method [2] is used.

Consider an involutive algebra $\operatorname{Pol}(\mathbb{C})_q$, determined by its generator z and the relation

$$z^* z = q^2 z z^* + 1 - q^2.$$
⁽¹⁾

This *-algebra may be treated as a q-analogue of the Weyl algebra [4]. However, we use an approach of [7, 9, 6] to the above algebra as a q-analogue of the polynomial algebra on the plane. We use a standard differential calculus [12]. Specifically, let $\Omega_q(\mathbb{C})$ be an involutive algebra given by its generators z, dz, and the relations (1),

$$dz \cdot z^* = q^{-2}z^* \cdot dz, \qquad dz \cdot z = q^2 z \cdot dz, \qquad dz \cdot dz^* = -q^{-2}dz^* \cdot dz, \qquad dz \cdot dz = 0.$$

A differential $d : \Omega_q(\mathbb{C}) \to \Omega_q(\mathbb{C})$ is defined in a standard manner on the elements z, z^* , dz, dz^* , and then extended in a unique way up to a differentiation of the graded algebra $\Omega_q(\mathbb{C})$ (deg $(z) = \deg(z^*) = 0$, deg $(dz) = \deg(dz^*) = 1$). Partial derivatives $\frac{\partial^{(l)}}{\partial z}, \frac{\partial^{(r)}}{\partial z}, \frac{\partial^{(l)}}{\partial z^*}, \frac{\partial^{(l)}}{\partial z^*}, \frac{\partial^{(l)}}{\partial z^*}$ are defined as such linear operators in $\operatorname{Pol}(\mathbb{C})_q$ that for all $f \in \operatorname{Pol}(\mathbb{C})_q$

$$df = dz \cdot \frac{\partial^{(l)} f}{\partial z} + dz^* \frac{\partial^{(l)} f}{\partial z^*} = \frac{\partial^{(r)} f}{\partial z} \cdot dz + \frac{\partial^{(r)} f}{\partial z^*} \cdot dz^*.$$

The bilinear maps $L: \operatorname{Pol}(\mathbb{C})_q \times \operatorname{Pol}(\mathbb{C})_q \to \operatorname{Pol}(\mathbb{C})_q$ of the form

$$L: f_1 \times f_2 \mapsto \sum_{i,j,k,m=0}^{N(L)} c_{ijkm} \cdot \left(\left(\frac{\partial^{(r)} f}{\partial z^*} \right)^i f_1 \right) \cdot z^{*j} z^k \cdot \left(\left(\frac{\partial^{(l)} f}{\partial z} \right)^m f_2 \right),$$

*Partially supported by the grant INTAS-RFBR-95-418

[†]Partially supported by the grant INTAS-94-4720

with $c_{ijkm} \in \mathbb{C}$, will be called q-bidifferential operators.

Our goal is to produce a formal deformation of the multiplication law in $Pol(\mathbb{C})_q$

*:
$$\operatorname{Pol}(\mathbb{C})_q \times \operatorname{Pol}(\mathbb{C})_q \to \operatorname{Pol}(\mathbb{C})_q[[t]],$$

*: $f_1 \times f_2 \mapsto f_1 \cdot f_2 + \sum_{j \in \mathbb{Z}_+} t^j \cdot C_j(f_1, f_2)$

and then to get explicit formulae for q-bidifferential operators $\{C_j\}_{j \in \mathbb{Z}_+}$. (The requirement of formal associativity [1] will be satisfied.)

It should be noted that the operators $\frac{\partial^{(l)}}{\partial z}$, $\frac{\partial^{(l)}}{\partial z^*}$, $\frac{\partial^{(r)}}{\partial z}$, $\frac{\partial^{(r)}}{\partial z^*}$ are extended "by a continuity" from the space of polynomials $\operatorname{Pol}(\mathbb{C})_q$ onto the space D(U)' of series of the form $f = \sum_{j,k \in \mathbb{Z}_+} a_{jk} z^j z^{*k}$, $a_{jk} \in \mathbb{C}$. This allows one to extend the *-product described in the sequel

from the polynomial algebra onto the "algebra of smooth functions in the quantum disc".

Everywhere below $q, t \in (0, 1)$.

We follow [7] in considering the operator of "weighted shift" \hat{z} given by $\hat{z}e_m = e_{m+1}$ with respect to an orthogonal base such that

$$||e_m||^2 = \frac{(q^2; q^2)_m}{(q^2 t; q^2)_m}.$$
(2)

A standard notation $(a;q^2)_m = (1-a)(1-q^2a) \cdot \ldots \cdot (1-q^{2(m-1)}a)$ is used here.

It was explained in [7] that the operators \hat{z} , \hat{z}^* supply a two-parameter quantization of the unit disc in the complex plane.

A covariant symbol of the operator $\hat{f} = \sum_{i,j} a_{ij} \hat{z}^i \hat{z}^{*j}$, $a_{ij} \in \mathbb{C}$ is defined to be the series $f = \sum_{ij} a_{ij} z^i z^{*j}$. The map $f \mapsto \hat{f}$ is a q-analogue of the Berezin quantization.¹ The formal

deformation * of the multiplication in $\operatorname{Pol}(\mathbb{C})_q$ is defined via this quantization:

$$\widehat{f_1 * f_2} = \widehat{f_1} \cdot \widehat{f_2}, \qquad f_1, f_2 \in \operatorname{Pol}(\mathbb{C})_q.$$

This deformation is well defined, as one can see from the following observation. (2) implies the relation

$$\hat{z}^*\hat{z} = q^2\hat{z}\hat{z}^* + 1 - q^2 + t\frac{1 - q^2}{1 - t}(1 - \hat{z}\hat{z}^*)(1 - \hat{z}^*\hat{z}).$$
(3)

Explicit formulae for the multiplication * are also accessible via multiple application of (3). However, in our opinion, the proof of those formulae presented in [10] is more appropriate.

Our principal result is For all $f_1, f_2 \in \operatorname{Pol}(\mathbb{C})_q$

$$f_1 * f_2 = (1-t) \sum_{j \in \mathbb{Z}_+} t^j m(p_j(\widetilde{\Box}) f_1 \otimes f_2),$$

with

i) $m : \operatorname{Pol}(\mathbb{C})_q \otimes \operatorname{Pol}(\mathbb{C})_q \to \operatorname{Pol}(\mathbb{C})_q$ being a multiplication in $\operatorname{Pol}(\mathbb{C})_q$,

¹In this work we consider only polynomial symbols $f \in \operatorname{Pol}(\mathbb{C})_q$. The general case was investigated in [10].

$$\begin{array}{l} ii) \stackrel{\simeq}{\square} = q^{-2}(1 - (1 + q^{-2})z^* \otimes z + q^{-2}z^{*2} \otimes z^2) \cdot \frac{\partial^{(r)}}{\partial z^*} \otimes \frac{\partial^{(l)}}{\partial z}, \\ iii) p_j(\stackrel{\simeq}{\square}) = \sum_{k=0}^j \frac{(q^{-2j};q^2)_k}{(q^2;q^2)_k^2} q^{2k} \cdot \prod_{i=0}^{k-1} \left(1 - q^{2i}\left((1 - q^2)^2 \stackrel{\simeq}{\square} + 1 + q^2\right) + q^{4i+2}\right). \end{array}$$

The proof of this theorem is presented in [10]. It involves the notions of a contravariant symbol and of a Berezin transform [11]. The contravariant symbol of the operator $\hat{f} = \sum_{i,j=1}^{N(\hat{f})} a_{ij} \hat{z}^{*i} \hat{z}^{j}$, $a_{ij} \in \mathbb{C}$, is defined to be a polynomial $\stackrel{\circ}{f} = \sum_{i,j=1}^{N(\hat{f})} a_{ij} z^{*i} z^{j} \in \text{Pol}(\mathbb{C})_q$. The Berezin transformation B_{σ} : $\text{Pol}(\mathbb{C})_{\sigma} \to \text{Pol}(\mathbb{C})_{\sigma}[[t]], B_{\sigma} : \stackrel{\circ}{f} \mapsto f$ takes the contravariant symbols of

transformation $B_q : \operatorname{Pol}(\mathbb{C})_q \to \operatorname{Pol}(\mathbb{C})_q[[t]], B_q : \stackrel{\circ}{f} \mapsto f$ takes the contravariant symbols of operators to their covariant symbols.

Just as in the case of an ordinary disc, the proof of the theorem reduces to constructing an "asymptotic expansion" of the Berezin transform. The explicit form for the coefficients of this expansion could be found via an application of the methods of the quantum group theory [5]. (*-algebra $\text{Pol}(\mathbb{C})_q$ is a $U_q\mathfrak{sl}_{1,1}$ -module algebra, and the Berezin transformation is a morphism of $U_q\mathfrak{sl}_{1,1}$ -modules.)

References

- F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowich, D. Sternheimer, *Deformation theory* and quantization, I, II, Ann. Physics, **110** (1978), 61 – 151.
- [2] F. Berezin, General concept of quantization, Commun. Math. Phys., 40 (1975), 153 174.
- [3] M. Cahen, S. Gutt, J. Rawnsley, Quantization of Kähler manifolds III, Letters in Mathematical Physics, 30, 291 305.
- [4] M. Chaichian, P. P. Kulish. Quantum group covariant systems, preprint q-alg/9512017.
- [5] V. Chari, A. Pressley. A Guide to Quantum Groups, Cambridge Univ. Press, 1995.
- [6] P. G. A. Floris. Addition Formula for q-adic Polynomials, Report W94-24, Mathematical Institute, University of Leiden, 10 p.
- S. Klimek, A. Lesniewski, A two-parameter quantum deformation of the unit disc, J. Funct. Anal. 115, (1993), 1 23.
- [8] C. Moreno, P. Ortego-Navaro, *-Products on D¹(ℂ), S², and related spectral analysis, Letters in Mathematical Physics, 7 (1983), 181 – 193.
- [9] G. Nagy, A. Nica. On the "quantum disc" and a "non-commutative circle", in: Algebraic Methods on Operator Theory, R. E. Curto, P. E. T. Jorgensen (eds.), Birkhauser, Boston, 1994, p. 276 – 290.
- [10] D. Shklyarov, S. Sinel'shchikov, L. Vaksman. On function theory in quantum disc: a q-analogue of the Berezin transformation, preprint, q-alg, 1998 (to appear).

- [11] A. Unterberger, H. Upmeier, The Berezin transform and invarian differential operators, Comm. Math. Phys., 164 (1994), 563 – 598.
- [12] Chong-Sun Chu, Pei-Ming Ho, B. Zumino. Some complex quantum manifolds and their geometry, preprint hep-th/9608188, 1996, 49 pp.