

Albanese and Picard 1-motives

Luca BARBIERI VIALE and Vasudevan SRINIVAS

Abstract – We define, in a purely algebraic way, 1-motives $\text{Alb}^+(X)$, $\text{Alb}^-(X)$, $\text{Pic}^+(X)$ and $\text{Pic}^-(X)$ associated with any algebraic scheme X over an algebraically closed field of characteristic zero. For X over $\mathbf{C}$ of dimension n the Hodge realizations are, respectively, $H^{2n-1}(X, \mathbf{Z}(n))/(\text{torsion})$, $H_1(X, \mathbf{Z})/(\text{torsion})$, $H^1(X, \mathbf{Z}(1))$ and $H_{2n-1}(X, \mathbf{Z}(1-n))/(\text{torsion})$.

1-motifs de Albanese et de Picard

Résumé – Nous définissons, de façon purement algébrique, les 1-motifs $\text{Alb}^+(X)$, $\text{Alb}^-(X)$, $\text{Pic}^+(X)$ et $\text{Pic}^-(X)$ associés au tout schéma algébrique X sur un corps de caractéristique 0. Pour X sur $\mathbf{C}$ les réalisations de Hodge sont, respectivement, $H^{2n-1}(X, \mathbf{Z}(n))/(\text{torsion})$, $H_1(X, \mathbf{Z})/(\text{torsion})$, $H^1(X, \mathbf{Z}(1))$ et $H_{2n-1}(X, \mathbf{Z}(1-n))/(\text{torsion})$.

Version française abrégée – Dans cette Note nous annonçons certains résultats dont les preuves se trouvent dans [1]. Nous donnons une construction algébrique pour les 1-motifs “d’Albanese et de Picard” en accord avec la conjecture de Deligne selon laquelle les 1-motifs associés à la structure de Hodge mixte sur la cohomologie des variétés algébriques admettent une définition algébrique (voir [3], §10.4.1). Etant donné que les structures de Hodge mixtes $H^{2n-1}(X, \mathbf{Z}(n))/(\text{torsion})$, $H_1(X, \mathbf{Z})/(\text{torsion})$, $H^1(X, \mathbf{Z}(1))$ et $H_{2n-1}(X, \mathbf{Z}(1-n))/(\text{torsion})$ pour X sur $\mathbf{C}$ de dimension n , sont de niveau ≤ 1 , elles sont conjecturellement algébrisables. Soit X un schéma algébrique sur un corps k algébriquement clos de caractéristique 0. Soit $f : \tilde{X} \rightarrow X$ une résolution des singularités, $\overline{X}$ une compactification lisse de $\tilde{X}$, Y le diviseur à croisements normaux et $Y = \cup Y_i$ la réunion des composantes lisses. Soit $\text{Pic}(\overline{X}, Y)$ le groupe des classes d’isomorphisme des paires $(\mathcal{L}, \varphi)$, telles que $\mathcal{L}$ soit un fibré en droites sur $\overline{X}$ et $\varphi : \mathcal{L}|_Y \cong \mathcal{O}_Y$ une trivialisations sur Y .

⊖ LEMME – Le foncteur de Picard rigidifié $T \leadsto \text{Pic}(\overline{X} \times_k T, Y \times_k T)$ est représentable. La composante connexe de l’identité $\text{Pic}_{(\overline{X}, Y)/k}^0$ est un schéma semi-abélien, représentable par une extension

$$0 \rightarrow T(\overline{X}, Y) \rightarrow \text{Pic}_{(\overline{X}, Y)/k}^0 \rightarrow [\ker^0 \text{Pic}_{\overline{X}/k}^0 \rightarrow \oplus \text{Pic}_{Y_i/k}^0] \rightarrow 0$$

d’un schéma abélien par un tore.

Pour chaque D disjoint de Y il y a une paire canonique $(\mathcal{O}(D), 1)$ dans $\text{Pic}(\overline{X}, Y)$. Soit S le lieu singulier de X et soit $\overline{S}$ la fermeture de $f^{-1}(S)$ dans $\overline{X}$. Alors $\text{Div}_{\overline{S}}^0(\overline{X}, Y)$ est le groupe des diviseurs engendré par les composantes propres de $f^{-1}(S)$ qui sont algébriquement nulles. Soit $\text{Div}_{\overline{S}/S}^0(\overline{X}, Y)$ le sous-groupe des diviseurs ayant image directe nulle sur S . Nous définissons le 1-motif “Picard homologique” de la façon suivante:

$$\text{Pic}^-(X) \stackrel{\text{def}}{=} [\text{Div}_{\overline{S}/S}^0(\overline{X}, Y) \rightarrow \text{Pic}^0(\overline{X}, Y)]$$

Le 1-motif “Albanese cohomologique” $\text{Alb}^+(X)$ est, par définition, le dual de Cartier de $\text{Pic}^-(X)$.

⊖ THÉORÈME – Si X est défini sur $\mathbf{C}$ et $n = \dim X$, alors la réalisation de Hodge de

$\mathrm{Pic}^-(X)$ est $H_{2n-1}(X, \mathbf{Z}(1-n))/(\text{torsion})$.

La réalisation de Hodge de $\mathrm{Alb}^+(X)$ est alors $H^{2n-1}(X, \mathbf{Z}(n))/(\text{torsion})$; par exemple, $\mathrm{Alb}^+(X) \cong J^n(X)$ lorsque X est une variété projective complexe. Soit $f_\bullet : X_\bullet \rightarrow X$ un hyperrecouvrement de X ; alors $X_\bullet$ est un schéma simplicial avec composantes lisses. Nous considérons une compactification propre et lisse $\overline{X}_\bullet$ à croisements normaux Y_i ; nous désignons par $\mathrm{Div}_{Y_\bullet}(\overline{X}_\bullet)$ le noyau de $d_0^* - d_1^* : \mathrm{Div}_{Y_0}(\overline{X}_0) \rightarrow \mathrm{Div}_{Y_1}(\overline{X}_1)$

⊕ LEMME – Le *fpqc*-faisceau $\mathbf{Pic}_{\overline{X}_\bullet/k}$ associé au foncteur de Picard simplicial $T \leadsto \mathbf{Pic}(\overline{X}_\bullet \times_k T)$ est représentable. La composante connexe de l'identité est un schéma semi-abélien, représentable par une extension

$$0 \rightarrow T(\overline{X}_\bullet) \rightarrow \mathbf{Pic}_{\overline{X}_\bullet/k}^0 \rightarrow [\ker^0 \mathrm{Pic}_{\overline{X}_0/k}^0 \rightarrow \mathrm{Pic}_{\overline{X}_1/k}^0] \rightarrow 0$$

d'un schéma abélien par un tore.

Nous désignons par $\mathrm{Div}_{Y_\bullet}^0(\overline{X}_\bullet)$ le sous-groupe des diviseurs ayant image dans $\mathbf{Pic}^0$. Nous définissons le 1-motif “Picard cohomologique” de X

$$\mathrm{Pic}^+(X) \stackrel{\mathrm{def}}{=} [\mathrm{Div}_{Y_\bullet}^0(\overline{X}_\bullet) \rightarrow \mathbf{Pic}^0(\overline{X}_\bullet)]$$

Le dual de Cartier de $\mathrm{Pic}^+(X)$ est le 1-motif “Albanese homologique” $\mathrm{Alb}^-(X)$.

⊕ THÉOREME – Si X est défini sur $\mathbf{C}$, alors la réalisation de Hodge de $\mathrm{Pic}^+(X)$ est $H^1(X, \mathbf{Z}(1))$.

La réalisation de Hodge de $\mathrm{Alb}^-(X)$ est $H_1(X, \mathbf{Z})/(\text{torsion})$. Certaines applications à l'étude des cycles algébriques ainsi que d'autres propriétés concernant les 1-motifs définis ci-dessus paraîtront dans [1].

In this Note we announce some results whose proof will appear in [1]. We give an algebraic construction for “Albanese and Picard” 1-motives, in accordance with Deligne's conjecture that the 1-motives associated to mixed Hodge structures on the cohomology groups of an algebraic variety should be algebraically defined (see [3], §10.4.1). The mixed Hodge structures $H^{2n-1}(X, \mathbf{Z}(n))/(\text{torsion})$, $H_1(X, \mathbf{Z})/(\text{torsion})$, $H^1(X, \mathbf{Z}(1))$ and $H_{2n-1}(X, \mathbf{Z}(1-n))/(\text{torsion})$ for X over $\mathbf{C}$ of dimension n are of the appropriate type, *i.e.*, are of niveau ≤ 1 , and hence the associated 1-motives conjecturally do have an algebraic description.

Let X be an algebraic scheme over an algebraically closed field k of characteristic zero.

HOMOLOGICAL PICARD AND COHOMOLOGICAL ALBANESE 1-MOTIVES – Let $f : \tilde{X} \rightarrow X$ be a resolution of singularities and let $\overline{X}$ be a smooth compactification of $\tilde{X}$ with normal crossing boundary Y , which has smooth irreducible components Y_i . Let $\pi_{\overline{X}} : \overline{X} \rightarrow \mathrm{Spec} k$, $\pi_Y : Y \rightarrow \mathrm{Spec} k$ be the structure morphisms. Then the *fpqc*-sheaves $(\pi_{\overline{X}})_* \mathbf{G}_{m,\overline{X}}$ and $(\pi_Y)_* \mathbf{G}_{m,Y}$ are represented by algebraic k -tori, as is $T(\overline{X}, Y) = \mathrm{coker} \left((\pi_{\overline{X}})_* \mathbf{G}_{m,\overline{X}} \rightarrow (\pi_Y)_* \mathbf{G}_{m,Y} \right)$. If $G \rightarrow H$ is a homomorphism of k -group schemes which are locally of finite type over k , let $\ker^0 G \rightarrow H$ denote the connected component of the identity of the k -group scheme $\ker G \rightarrow H$.

⊖ LEMMA – The rigidified Picard functor $T \leadsto \mathrm{Pic}(\overline{X} \times_k T, Y \times_k T)$ is representable by a k -group scheme which is locally of finite type, and whose group of k -points is $\mathrm{Pic}(\overline{X}, Y)$. The connected component of the identity $\mathrm{Pic}_{(\overline{X}, Y)/k}^0$ is a

semi-abelian scheme which can be represented as an extension

$$0 \rightarrow T(\overline{X}, Y) \rightarrow \mathrm{Pic}_{(\overline{X}, Y)/k}^0 \rightarrow [\ker^0 \mathrm{Pic}_{\overline{X}/k}^0 \rightarrow \oplus \mathrm{Pic}_{Y_i/k}^0] \rightarrow 0$$

of an abelian scheme by a torus.

We let $\mathrm{Pic}^0(\overline{X}, Y) \subset \mathrm{Pic}(\overline{X}, Y)$ denote the subgroup of k -points of $\mathrm{Pic}_{(\overline{X}, Y)/k}^0$. Any divisor D on $\overline{X}$ with support disjoint from Y has a class $(\mathcal{O}_{\overline{X}}(D), 1) \in \mathrm{Pic}(\overline{X}, Y)$; we say D is algebraically equivalent to 0 relative to Y if this class is in the subgroup $\mathrm{Pic}^0(\overline{X}, Y)$. Let S denote the singular locus of X , and $\overline{S}$ the closure in $\overline{X}$ of $f^{-1}(S)$. Let $\mathrm{Div}_{\overline{S}}^0(\overline{X}, Y)$ denote the group of divisors D on $\overline{X}$ such that $\mathrm{supp} D$ is disjoint from Y , and contained in $\overline{S}$, and also D is algebraically equivalent to 0 relative to Y . Let $\mathrm{Div}_{\overline{S}/S}^0(\overline{X}, Y) \subset \mathrm{Div}_{\overline{S}}^0(\overline{X}, Y)$ denote the subgroup of such divisors D which have vanishing push-forward under f . We can now define the “homological Picard” 1-motive as follows:

$$\mathrm{Pic}^-(X) \stackrel{\mathrm{def}}{=} [\mathrm{Div}_{\overline{S}/S}^0(\overline{X}, Y) \rightarrow \mathrm{Pic}^0(\overline{X}, Y).]$$

The “cohomological Albanese” 1-motive $\mathrm{Alb}^+(X)$ is defined to be the Cartier dual of $\mathrm{Pic}^-(X)$.

◻ THEOREM – *If X is defined over $\mathbf{C}$ and $n = \dim X$, then the Hodge realization of $\mathrm{Pic}^-(X)$ is $H_{2n-1}(X, \mathbf{Z}(1-n))/(\text{torsion})$.*

By duality, we have that the Hodge realization of $\mathrm{Alb}^+(X)$ is $H^{2n-1}(X, \mathbf{Z}(n))/(\text{torsion})$, e.g., $\mathrm{Alb}^+(X) \cong J^n(X)$ if X is a projective complex variety, where $J^n(X)$ is the semi-abelian variety considered in [2].

COHOMOLOGICAL PICARD AND HOMOLOGICAL ALBANESE 1-MOTIVES – We let $f_* : X_* \rightarrow X$ be a proper smooth hypercovering of X ; then X_* is a simplicial k -scheme with smooth components. Consider a proper smooth compactification $\overline{X}_*$ with normal crossing boundary Y_* , and denote by $\mathrm{Div}_{Y_*}(\overline{X}_*)$ the group of divisors on X_0 which are supported on Y_0 , and have zero pullback on X_1 , i.e., by definition

$$\mathrm{Div}_{Y_*}(\overline{X}_*) \stackrel{\mathrm{def}}{=} \ker \mathrm{Div}_{Y_0}(\overline{X}_0) \xrightarrow{d_0^* - d_1^*} \mathrm{Div}_{Y_1}(\overline{X}_1).$$

Let $\pi_i : \overline{X}_i \rightarrow \mathrm{Spec} k$ be the structure morphism. Let $T(\overline{X}_*)$ be the k -torus representing the $fpqc$ -sheaf

$$\frac{\ker[(\pi_1)_* \mathbf{G}_{m, \overline{X}_1} \rightarrow (\pi_2)_* \mathbf{G}_{m, \overline{X}_2}]}{\mathrm{im}[(\pi_0)_* \mathbf{G}_{m, \overline{X}_0} \rightarrow (\pi_1)_* \mathbf{G}_{m, \overline{X}_1}]}.$$

◻ LEMMA – *The sheaf $\mathbf{Pic}_{\overline{X}_*./k}$, with respect to $fpqc$ -topology, associated to the simplicial Picard functor $T \leadsto \mathbf{Pic}(\overline{X}_* \times_k T)$, is representable by a group scheme locally of finite type over k , with k -points $\mathrm{Pic}(\overline{X}_*)$. The connected component of the identity is a semi-abelian scheme, which can be represented as an extension*

$$0 \rightarrow T(\overline{X}_*) \rightarrow \mathbf{Pic}_{\overline{X}_*./k}^0 \rightarrow [\ker^0 \mathrm{Pic}_{\overline{X}_0/k}^0 \rightarrow \mathrm{Pic}_{\overline{X}_1/k}^0] \rightarrow 0$$

of an abelian scheme by a torus.

Let $\mathrm{Div}_{Y_*}^0(\overline{X}_*)$ denote the subgroup of $\mathrm{Div}_{Y_*}(\overline{X}_*)$ of those divisors which are

mapped to $\mathbf{Pic}^0(k)$ under the canonical mapping. We then define the “cohomological Picard” 1-motive of X as follows:

$$\mathrm{Pic}^+(X) \stackrel{\mathrm{def}}{=} [\mathrm{Div}_Y^0(\overline{X}.) \rightarrow \mathbf{Pic}^0(\overline{X}.)].$$

The “homological Albanese” 1-motive $\mathrm{Alb}^-(X)$ is defined to be the Cartier dual of $\mathrm{Pic}^+(X)$.

⊕ THEOREM – *If X is defined over $\mathbf{C}$, then the Hodge realization of $\mathrm{Pic}^+(X)$ is $H^1(X, \mathbf{Z}(1))$.*

We then also have that the Hodge realization of $\mathrm{Alb}^-(X)$ is $H_1(X, \mathbf{Z})/(\text{torsion})$.

REMARKS – We keep the same notation as above.

1. We can show that our definitions are independent of choices of resolutions, compactifications or hypercoverings.
2. If X is normal and projective over $k = \overline{k}$ then $\mathrm{Alb}^+(X)$ is an abelian variety, which may be identified with $\mathrm{Alb}(\tilde{X})$, the Albanese variety of a resolution. If X is projective, but not necessarily normal, then $\mathrm{Alb}^+(X)$ is a semi-abelian variety. If X is smooth, possibly open, then $\mathrm{Alb}^-(X)$ is a semi-abelian variety.
3. Let X be irreducible and projective of dimension n , and let X_{reg} denote the set of smooth points of X ; we may also regard X_{reg} as an open subscheme of any given resolution of singularities $\tilde{X}$. Let $x_0 \in X_{\mathrm{reg}}$ be a smooth point and let $a_{x_0} : \tilde{X} \rightarrow \mathrm{Alb}(\tilde{X})$ be the corresponding Albanese mapping; considering Alb^+ as a torus bundle over $\mathrm{Alb}(\tilde{X})$, and pulling back along a_{x_0} , one can see it has a natural trivialization on X_{reg} , yielding a section $a_{x_0}^+ : X_{\mathrm{reg}} \rightarrow \mathrm{Alb}^+(X)$. This map yields a universal regular homomorphism $CH^n(X)_{\mathrm{deg} 0} \rightarrow \mathrm{Alb}^+(X)$ from the “cohomological” Levine-Weibel Chow group of zero cycles to semi-abelian varieties, and the corresponding Roitman torsion theorem can be proved (*c.f.* [2]).
4. We clearly have a commutative square of 1-motives

$$\begin{array}{ccc} \mathrm{Pic}^-(\tilde{X}) & \longrightarrow & \mathrm{Pic}^+(\tilde{X}) \\ \downarrow & & \downarrow \\ \mathrm{Pic}^-(X) & \longrightarrow & \mathrm{Pic}^+(X_{\mathrm{reg}}) \end{array}$$

In particular, by taking Cartier duals we have a map $\mathrm{Alb}^-(X_{\mathrm{reg}}) \rightarrow \mathrm{Alb}^+(X)$. Then there is a morphism $X_{\mathrm{reg}} \rightarrow \mathrm{Alb}^-(X_{\mathrm{reg}})$ which lifts the section $a_{x_0}^+$ above when X is proper. In fact, Alb^- of a smooth variety is the universal semi-abelian variety associated to it by Serre [6], §5.

5. The claimed representabilities of “rigidified” and “simplicial” Picard functors can be obtained *via* the classical theorems due to Grothendieck [4] and Murre [5].
6. For further properties we refer to [1] where we will consider ℓ -adic and De Rham realizations; functoriality properties, algebraic Gysin maps and

Lefschetz theorems (*c.f.* [2]) are also expected. There are also analogous definitions of the 1-motives in the case when the ground field k is an arbitrary field of characteristic 0. However, the tori appearing need not be split, for example. The general case is reduced to the case of an algebraically closed field using Galois descent.

Ce travail a été réalisé lors de séjours des auteurs à Paris VII, TIFR Mumbai, DIMA Genova et ICTP Trieste, que nous remercions.

RÉFÉRENCES

- [1] L. BARBIERI VIALE et V. SRINIVAS: Albanese and Picard 1-motives, travail en préparation.
- [2] J. BISWAS et V. SRINIVAS: Roitman's theorem for singular projective varieties, preprint, 1996.
- [3] P. DELIGNE: Théorie de Hodge III, *Publ. Math. IHES* **44** (1974) 5–78.
- [4] A. GROTHENDIECK: Technique de descente et théorèmes d'existence en géométrie algébrique V–VI Les schémas de Picard, dans “Fondements de la Géométrie Algébrique” Extraits du *Séminaire BOURBAKI* 1957/62.
- [5] J. MURRE: On contravariant functors from the category of preschemes over a field into the category of abelian groups (with application to the Picard functor), *Publ. Math. IHES* **23** (1964) 581–619.
- [6] J.P. SERRE: Morphismes universels et variétés d'Albanese, dans “Variétés de Picard” *ENS Séminaire C. CHEVALLEY* 3e année: 1958/59.

*L. B. V.: Dipartimento di Matematica, Università di Genova
Via Dodecaneso, 35, I-16146 Genova, Italia;
V.S.: Tata Institute of Fundamental Research, School of Mathematics
Homi Bhabha Road, 400 005 Mumbai, India.*