Jacobian Conjecture and Nilpotent Mappings

Vik.S. Kulikov *

Abstract

We prove the equivalence of the Jacobian Conjecture (JC(n)) and the Conjecture on the cardinality of the set of fixed points of a polynomial nilpotent mapping (JN(n)) and prove a series of assertions confirming JN(n).

Introduction

where

Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be a morphism of complex affine spaces. If we choose coordinates in the image and preimage, then F is defined by n polynomials $Y_i = F_i(X_1, ..., X_n) \in \mathbb{C}[X] = \mathbb{C}[X_1, ..., X_n]$ in n variables. Put

 $F'(X) = \det JF(X),$

$$JF = \left(\frac{\partial F_i}{\partial X_j}\right)_{1 \le i,j \le n}$$

is the Jacobi martix of F.

It is well-known that F is locally invertible in a neighborhood of some point $x \in \mathbb{C}^n$ (as an analytic mapping) if and only if $F'(x) \neq 0$. Therefore, " $F'(x) \neq 0$ for all $x \in \mathbb{C}^n$ " is a necessary condition for the invertibility of F.

The famous Jacobian Conjecture (JC(n)) claims:

Conjecture JC(n). Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be a morphism such that $F'(x) \neq 0$ for all $x \in \mathbb{C}^n$ (or, equivalently, $F'(X) \in \mathbb{C}^*$), then F is an isomorphism, i.e. F possesses an inverse mapping which is also given by polynomials.

Without loss of generality we can assume that $F'(X) \equiv 1$ if $F'(X) \in \mathbb{C}^*$. The following reformulations of conjecture JC(n) are well-known

Theorem 1 (cf. [2]). Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be a morphism with $F'(X) \in \mathbb{C}^*$. The Jacobian conjecture JC(n) is equivalent to either of the following assertions:

(i) F is injective;

- (ii) The degree deg F of F is equal to 1 (i.e. $\mathbb{C}(X_1,...,X_n)) = \mathbb{C}(F_1(X),...,F_n(X)));$
- (iii) The ring $\mathbb{C}[X_1, ..., X_n]$ is a finitely generated $\mathbb{C}[F_1(X), ..., F_n(X)]$ -module;
- (iv) The ring $\mathbb{C}[X_1, ..., X_n]$ is a projective $\mathbb{C}[F_1(X), ..., F_n(X)]$ -module;

^{*}Partly supported by RFFI (No. 96-01-00614) and INTAS (No. 96-0713).

(v) The ring $\mathbb{C}[F_1(X), ..., F_n(X)]$ is integrally closed in $\mathbb{C}[X_1, ..., X_n]$; (vi) The field $C(X_1, ..., X_n)$ is a Galois extension over $\mathbb{C}(F_1(X), ..., F_n(X))$; (vii) F is proper.

The Jacobian Conjecture have been formulated by O.H.Keller in 1939 [7] in the case n = 2for the polynomials with integer coefficients. For n = 1 conjecture JC(n) is trivial since a polynomial $F(X_1) \in \mathbb{C}[X_1]$ with $F'(X_1) \in \mathbb{C}^*$ is of degree one and, therefore, defines an isomorphism $F : \mathbb{C}^1 \to \mathbb{C}^1$. The Jacobian Conjecture JC(n) remains an open problem even in the case n = 2 in spite of attempts of many mathematicians to prove it or to find a counterexample (a survey of the results on the Jacobian Conjecture one can find in [2], [4], [5]). These attempts allow to obtain many partial results connected with the Jacobian Conjecture and even several false proofs were published (for the critique of these false proofs we refer to [12] and [2]).

The aim of this short note is to attract attention of the reader to a connection of JC(n)and one of conjectures on the nilpotent mappings.

Definition 1 A morphism $N = (N_1, ..., N_n) : \mathbb{C}^n \to \mathbb{C}^n$ is called nilpotent if JN(X) is a nilpotent matrix, i.e.

$$(JN(X))^n = \left(\frac{\partial N_i}{\partial X_i}\right)^n \equiv 0$$

in the ring of matrices $\mathcal{M}(n, \mathbb{C}[X])$ of n-th order with coefficients in $\mathbb{C}[X]$.

We consider nilpotent mappings as endomorphisms of \mathbb{C}^n . A point $x \in \mathbb{C}^n$ is called a *fixed* point of a nilpotent mapping N if N(x) = x. It is well-known that if N is linear and nilpotent, then $N^n(\mathbb{C}^n)$ consists of a unique point. In particular, a linear nilpotent mapping N possesses only one fixed point.

Conjecture JN(n). A nilpotent mapping $N : \mathbb{C}^n \to \mathbb{C}^n$ possesses at most one fixed point.

The following theorem describes a connection between the conjectures JC(n) and JN(n).

Theorem 2 The conjectures JC(n) and JN(n) are equivalent, i.e. JC(n) is true for all n if and only if JN(n) is true for all n.

In section 1, we recall some well-known results relating to the Jacobian Conjecture. In section 2, we prove Theorem 2 and a series of assertions confirming JN(n).

This paper was written during my stay at the Max-Planck-Institut für Mathematik in Bonn. It is a pleasure to thank the Institut for its hospitality and financial support.

1 Stable equivalence of mappings and the Jacobian Conjecture

1.1. Let $F : \mathbb{C}^n \to \mathbb{C}^n$ and $G : \mathbb{C}^n \to \mathbb{C}^n$ be morphisms given by polynomials $F_1(X), ..., F_n(X)$ and $G_1(X), ..., G_n(X)$ respectively. We shall say that F and G are *equivalent* if there exist automorphisms $H, R : \mathbb{C}^n \to \mathbb{C}^n$ such that the following diagram is commutative

$$\begin{array}{ccc} \mathbb{C}^n & \xrightarrow{H} & \mathbb{C}^n \\ G & & & \downarrow F \\ \mathbb{C}^n & \xrightarrow{R} & \mathbb{C}^n. \end{array}$$

In other words, F and G are equivalent if

$$R_i(G_1(X), ..., G_n(X)) = F_i(H_1(X), ..., H_n(X))$$

for all *i*, where $H_1(X), ..., H_n(X)$ and $R_1(X), ..., R_n(X)$ are polynomials in $\mathbb{C}[X_1, ..., X_n]$ defining the automorphisms H and R.

Definition 2 Morphisms $F : \mathbb{C}^n \to \mathbb{C}^n$ and $G : \mathbb{C}^m \to \mathbb{C}^m$, $n \leq m$, are called stably equivalent if $F \times Id : \mathbb{C}^m = \mathbb{C}^n \times \mathbb{C}^{m-n} \to \mathbb{C}^n \times \mathbb{C}^{m-n} = \mathbb{C}^m$ and G are equivalent.

The following assertion is a consequence of the definition.

Claim 1 If F and G are stably equivalent, then

$$\deg F = \deg G.$$

In particular, if $F'(X) \in \mathbb{C}^*$ for F and F and G are stably equivalent, then F and G are invertible (or not invertible) simultaneously.

1.2. To avoid confusion with the degree of a morphism, we shall call *p*-degree or polynomial degree of a morphism $F = (F_1, ..., F_n) : \mathbb{C}^n \to \mathbb{C}^n$ the maximum of the degrees of $F_i(X)$ defining F,

$$\deg_n F = \max \deg F_i(X).$$

Of course, *p*-degree depends on a choice of affine coordinates in the image and preimage.

A usual construction to simplify the form of polynomials defining a morphism is the following: one replaces a mapping by a stably equivalent one and performs linear and triangular automorphisms of the image and the preimage.

Definition 3 An automorphism $H : \mathbb{C}^n \to \mathbb{C}^n$ is called triangular if, in some coordinate system, it can be given by polynomials $H_i(X_1, ..., X_n)$, i = 1, ..., n, of the form

$$H_i(X_1, ..., X_n) = X_i + H_i(X_1, ..., X_{i-1}),$$

where $H_i(X_1, ..., X_{i-1}) \in \mathbb{C}[X_1, ..., X_{i-1}]$ is a polynomial in variables $X_1, ..., X_{i-1}$.

The following theorem is well-known.

Theorem 3 ([2], [15]). For any morphism $F : \mathbb{C}^n \to \mathbb{C}^n$ there exists a stably equivalent morphism $G : \mathbb{C}^m \to \mathbb{C}^m$ such that $\deg_p G \leq 3$.

1.3. Let $F : \mathbb{C}^n \to \mathbb{C}^n$, given by polynomials $F_i(X)$, map the origin to the origin, i.e. F(0) = 0. A morphism $F_t : \mathbb{C}^{n+1} \to \mathbb{C}^{n+1}$ given by the polynomials

$$\widetilde{F}_{i}(X_{1},...,X_{n},T) = \sum_{j=1}^{\deg F_{i}} T^{j-1}F_{(j)i}(X_{1},...,X_{n}), \quad i = 1,...,n,$$

$$\widetilde{F}_{n+1}(X_{1},...,X_{n},T) = T,$$

is called a blow up of F, where $F_{(j)i}(X)$ is the homogeneous component of degree j of the polynomial $F_i(X)$. It is easy to check the following assertion.

Claim 2 Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be a morphism such that F(0) = 0, then (i) $F'_t(X_1, ..., X_n, T) = F'(TX_1, ..., TX_n)$; (ii) F, with $F'(X) \in \mathbb{C}^*$, is invertible if and only if F_t is invertible.

Geometrically, to obtain F_t from F, one needs to perform σ -processes with centers at the origins of the image and the preimage of the morphism $F \times Id : \mathbb{C}^{n+1} = \mathbb{C}^n \times \mathbb{C}^1 \to \mathbb{C}^n \times \mathbb{C}^1 = \mathbb{C}^{n+1}$ and to restrict the obtained morphism onto coordinates neighborhoods corresponding the last coordinate T.

In [3], theorem 3 was made more precise.

Theorem 4 Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be a morphism such that F(0) = 0 and $F'(0) \neq 0$. Then the blow up F_t of F is stably equivalent to $G : \mathbb{C}^m \to \mathbb{C}^m$ defined by polynomials $G_i(X)$ of the form

$$G_i(X_1, ..., X_m) = X_i + G_{(3)i}(X_1, ..., X_m),$$
(1)

where $G_{(3)i}(X_1, ..., X_m)$ are homogeneous forms of degree 3.

Since the space of the homogeneous polynomials of degree k is generated by k-th powers of the linear polynomials, the following theorem is a simple consequence of Theorem 4.

Theorem 5 Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be given by polynomials

$$F_i(X_1, ..., X_n) = X_i + F_{(3)i}(X_1, ..., X_n)$$

Then F is stably equivalent to $G: \mathbb{C}^m \to \mathbb{C}^m$ given by polynomials $G_i(X)$ of the form

$$G_i(X_1, ..., X_m) = X_i + (\sum a_{i,j} X_j)^3.$$
 (2)

1.4. The Jacobian conjecture was proved in [14] and [6] for the morphisms of the form (1) with $m \leq 4$ and in [6] for the morphisms of the form (2) with $m \leq 7$. Besides, in [13] the following theorem was proved

Theorem 6 The conjecture JC(n) is true for $F : \mathbb{C}^n \to \mathbb{C}^n$ of polynomial degree $\deg_p F \leq 2$.

This theorem will be obtained in the next section as a consequence of more general result (see Proposition 1).

Note also that the conjecture JC(n) was proved in [8] in the following two cases: when F is given by polynomials $F_i(X) \in X_i \mathbb{C}[X]$ and when $F_i(X) = X_i + \lambda_i M_i(X)$, where $M_i(X)$ are monomials.

1.5. In the case $F : \mathbb{C}^2 \to \mathbb{C}^2$ the Jacobian conjecture holds for all F of polynomial degree $\deg_p F \leq 100 [10]$. In [9] (see also [5]), it was proved that if the degrees of polynomials defining F with $F'(X) \in \mathbb{C}^*$ are coprime, then F is invertible and moreover in this case one of polynomials must be linear. In particular, the Jacobian conjecture holds if one of the polynomials defining F is of prime degree. In [1], this result was generalized to the case when the degree of one of the polynomials defining F is a product of two different prime integers.

In addition, in [11], it was shown that if $F : \mathbb{C}^2 \to \mathbb{C}^2$ with $F'(X) \equiv 1$ is not invertible, then deg $F \geq 4$.

1.6. A morphism $F: \mathbb{C}^n \to \mathbb{C}^n$ induces a polynomial mapping $\widetilde{F}: \mathbb{R}^{2n} \to \mathbb{R}^{2n}$ given by

$$\widetilde{F} = (Re F_1, Im F_1, ..., Re F_n, Im F_n),$$

where F_i are polynomials defining F. It is easy to see that

$$\det J\widetilde{F} = |\det JF|^2 .$$

Hence, det $JF \in \mathbb{C}^*$ if and only if det $J\widetilde{F} \in \mathbb{R}^*$.

Conjecture RJC(n). A polynomial mapping $F : \mathbb{R}^n \to \mathbb{R}^n$ with det $JF \in \mathbb{R}^*$ is injective.

It is easy to show that if RJC(n) is true, then $F : \mathbb{R}^n \to \mathbb{R}^n$ with det $JF \in \mathbb{R}^*$ is invertible and the following proposition holds.

Proposition 1 The conjecture JC(n) is true for all n if and only if the conjecture RJC(n) is true for all n.

Consider $F : \mathbb{R}^n \to \mathbb{R}^n$ with det $JF \in \mathbb{R}^*$. Without loss of generality, we can assume that F(0) = 0 and JF(0) = E, where E is the identity matrix. Write F in the form

$$F = Id + F_{(2)} + \dots + F_{(m)}, \tag{3}$$

where $F_{(i)}$ are morphisms defined by homogeneous polynomials of degree *i*.

Definition 4 A polynomial mapping $F : \mathbb{R}^n \to \mathbb{R}^n$ of the form (3) is called positive (resp. negative) if all non-zero coefficients of the $F_{(i)}$ are positive (resp. negative).

In [16], the following theorem has been proved.

Theorem 7 (i) The conjecture RJC(n) holds for all n and all negative polynomial mappings $F : \mathbb{R}^n \to \mathbb{R}^n$.

(ii) It suffices to prove RJC(n) for all positive polynomial mappings $F : \mathbb{R}^n \to \mathbb{R}^n$.

It follows from Theorem 7 (i) (passing to the blow up of F) that

The conjecture RJC(n) holds for all n and for all positive polynomials mappings $F : \mathbb{R}^n \to \mathbb{R}^n$ of the form $F = Id + F_{(2)} + F_{(4)} + ... + F_{(2m)}$, and also for all polynomial mappings F of the form $F = Id + F_{(2)} - F_{(3)} + ... + (-1)^m F_{(m)}$, where $F_{(i)}$ are morphisms defined by homogeneous polynomials of degree i with non-negative coefficients.

2 Nilpotent Mappings

2.1. It is well-known that a matrix $A \in \mathcal{M}(n, \mathbb{C})$ is nilpotent if and only if $\det(tE - A) \equiv t^n$, where *E* is the identity matrix. As a consequence we obtain the following Lemma.

Lemma 1 A morphism $N : \mathbb{C}^n \to \mathbb{C}^n$ given by polynomials $N_i(X)$ is nilpotent if and only if $\widetilde{F} = (Id - T \cdot N) \times Id : \mathbb{C}^n \times \mathbb{C}^1 \to \mathbb{C}^n \times \mathbb{C}^1$, given by the polynomials

$$\widetilde{N}_i(X_1, ..., X_n, T) = X_i - TN_i(X_1, ..., X_n) \quad for \ i = 1, ..., n,$$

$$\widetilde{F}_{n+1}(X_1, ..., X_n, T) = T,$$

has $\widetilde{F}'(X,T) \equiv 1$.

If we apply lemma 1 to the restriction of $\widetilde{F} = Id - T \cdot N$ to the hypersurface $\mathbb{C}^n = \{T = 1\}$ we obtain that for the nilpotent mapping N the corresponding morphism $F = Id - N : \mathbb{C}^n \to \mathbb{C}^n$ has $F'(X) \equiv 1$. Besides, if there exist d distinct fixed points of N, then for F = Id - N the preimage of the origin consists of d distinct points, and hence degree of F = Id - N is not less than d. In particular, if a nilpotent mapping $N : \mathbb{C}^n \to \mathbb{C}^n$ possesses d > 1 fixed points, then $F = Id - N : \mathbb{C}^n \to \mathbb{C}^n$ is not invertible.

2.2. Theorem 2 is a consequence of the following theorem.

Theorem 8 Any morphism $F : \mathbb{C}^n \to \mathbb{C}^n$ of degree deg F = d with $F'(X) \equiv 1$ is stably equivalent to \overline{F} of the form

$$\overline{F} = Id - N,$$

where N is a nilpotent morphism possessing d fixed points.

Conversely, if a nilpotent mapping $N : \mathbb{C}^n \to \mathbb{C}^n$ possesses d distinct fixed points, then $\overline{F} = Id - N$ is of degree not less than d.

Proof. Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be a morphism of degree deg F = d with $F'(X) \equiv 1$. Performing a translation and linear transformations, if necessary, we can assume that F(0) = 0, the number of the preimages of the origin $\#\{F^{-1}(0)\} = \deg F$, and F is given by polynomials

$$F_i(X_1, ..., X_n) = X_i - F_{(\geq 2)i}(X_1, ..., X_n),$$
(4)

where $F_{(\geq 2)i}(X_1, ..., X_n)$ are polynomials containing no monomials of degree zero and one.

Consider the blow up $F_t: \mathbb{C}^{n+1} \to \mathbb{C}^{n+1}$ of F. The morphism F_t is defined by polynomials

$$\begin{array}{rcl} Y_i &=& \widetilde{F}_i(X_1,...,X_n,T) &=& X_i - \sum_{j=2}^{\deg F_i} T^{j-1} F_{(j)i}(X_1,...,X_n), \quad i=1,...,n, \\ \widetilde{T} &=& \widetilde{F}_{n+1}(X_1,...,X_n,T) &=& T. \end{array}$$

Let us show that F_t is stably equivalent to $\widetilde{G} = (Id - T \cdot G) \times Id : \mathbb{C}^m \times \mathbb{C}^1 \to \mathbb{C}^m \times \mathbb{C}^1$ given by $\widetilde{Y}_i = \widetilde{G}_i(X_1, ..., X_m, T) = X_i - TG_i(X_1, ..., X_m)$ for i = 1, ..., m, and $\widetilde{G}_{m+1}(X_1, ..., X_n, T) = T$, and of degree deg $\widetilde{G} = \deg F$, and the number of the preimage of the point with coordinates (0, ..., 0, 1) is $\#\{\widetilde{G}^{-1}((0, ..., 0, 1))\} = \deg F$. For this we must get rid of all monomials of the form

 $T^k X_1^{r_1} \cdots X_n^{r_n}$ with k > 1 and then apply Lemma 1 to $G : \mathbb{C}^m \to \mathbb{C}^m$ defined by $G_i(X_1, ..., X_m)$, i = 1, ..., m. By Lemma 1, G is nilpotent and the number of the fixed points of G is equal to deg F.

Removing the monomial $T^k X_1^{r_1} \cdots X_n^{r_n}$ with k > 1 can be performed by adding new variables (passing to stably equivalent morphisms) and performing linear and triangular transformations. In fact, let for some i a polynomial \widetilde{Y}_i be of the form

$$\widetilde{Y}_i = \widetilde{X}_i - \sum_{j=1}^{k_i} T^j F_{(j)i}(\widetilde{X}_1, ..., \widetilde{X}_n)$$

(here $F_{(j)i}(\widetilde{X}_1, ..., \widetilde{X}_n)$ is not necessary homogeneous). We add a new variable $\widetilde{Y}_{n+1} = \widetilde{X}_{n+1}$ and perform a triangular transformation in the preimage:

$$\begin{aligned} X_i &= \widetilde{X}_i, & i = 1, ..., n \\ X_{n+1} &= \widetilde{X}_{n+1} + TF_{(k_i)i}(\widetilde{X}_1, ..., \widetilde{X}_n). \end{aligned}$$

Next we perform a triangular transformation in the image:

$$\begin{array}{ll} Y_l = & \widetilde{Y}_l, & l = 1,...,i-1,i+1,...,n+1 \\ Y_i = & \widetilde{Y}_i - T^{k_i-1}\widetilde{Y}_{n+1}. \end{array}$$

It is easy to see that in the coordinates $X_1, ..., X_{n+1}, T$ and $Y_1, ..., Y_{n+1}, T$ the morphism $F_t \times Id$ is given by equations:

$$\begin{array}{rcl} Y_s & = & X_s - \sum_{j=1}^{k_s} T^j F_{(j)s}(X_1, ..., X_n), & s = 1, ..., i-1, i+1, ..., n \\ Y_i & = & X_i - \sum_{j=1}^{k_i-1} T^j F_{(j)l}(X_1, ..., X_n) - T^{k_i-1} X_{n+1}, \\ Y_{n+1} & = & X_{n+1} - T F_{(k_i)i}(X_1, ..., X_n). \end{array}$$

Thus, we removed the summand of $\tilde{F}_i(X_1, ..., X_n, T)$ containing the variable T is contained with the maximal multiplicity k_i . Obviously, after a finite number of similar steps we obtain a morphism stably equivalent to F_t and satisfying the required conditions.

2.3. Let $F : \mathbb{C}^n \to \mathbb{C}^n$ be of the form $F = Id - F_{(k)}$, i.e. F is defined by

$$F_i(X) = X_i - F_{(k)i}(X_1, ..., X_n), \qquad i = 1, ..., n_i$$

where $F_{(k)i}(X_1, ..., X_n)$ are homogeneous forms of degree k. If $F'(X) \equiv 1$, then $F_{(k)}$ is nilpotent. In fact, by Theorem 2, the Jacobian $F'_t(X, T)$ of the blow up F_t is equal to

$$F'_t(X,T) = F'(TX_1, ..., TX_n) = \det(E - T^{k-1}JF_{(k)}) \equiv 1.$$

Therefore, the Jacobian of \hat{F}_t which is defined by polynomials

$$\begin{array}{rcl} Y_i &=& X_i - TF_{(k)i}(X_1,...,X_n), & i = 1,...,n, \\ T &=& T, \end{array}$$

is also equal to 1. Hence, by Lemma 1, $F_{(k)}$ is nilpotent.

Proposition 2 The conjecture JN(n) holds for nilpotent morphisms $N_{(k)} : \mathbb{C}^n \to \mathbb{C}^n$ defined by homogeneous forms of degree k.

Proof. The origin, i.e. the point o = (0, ..., 0), is a fixed point of $N = N_{(k)}$. Let $x = (x_1, ..., x_n) \neq o$ be another fixed point of $N_{(k)}$. This contradicts the nilpotency of N, since the vector $\overline{x} = (x_1, ..., x_n)$ is an eigenvector (corresponding to a non-zero eigenvalue (equals to k)) of the linear mapping $JN_{(k)}(X)$ at the point X = x. In fact, by Euler's formula

$$JN_{(k)}(x)(\overline{x}) = \left(\sum_{j=1}^{n} x_j \frac{\partial N_1}{\partial X_j}(x), \dots, \sum_{j=1}^{n} x_j \frac{\partial N_n}{\partial X_j}(x)\right)$$

= $(kN_1(x), \dots, kN_n(x)) =$
= $(kx_1, \dots, kx_n) = k \cdot \overline{x},$

where $N_i(X)$ are polynomials defining $N = N_{(k)}$.

Theorem 6 is an easy consequence of this theorem. For let F be of polynomial degree $\deg_p F = 2$. After a linear changes of variables, we can assume that F is of the form $F = Id - F_{(2)}$ and the preimage $F^{-1}(0)$ consists of deg F distinct points. Hence, the nilpotent morphism $F_{(2)}$ possesses deg F distinct fixed points. Therefore, deg F = 1, i.e. F is invertible.

As a consequence of above considerations, Lemma 1, Theorem 6, and the pool of Theorem 8 we obtain the following theorem.

Theorem 9 The conjecture JN(n) holds for $N : \mathbb{C}^n \to \mathbb{C}^n$ of polynomial degree $\deg_p N \leq 2$.

Proposition 2 may be generalized as follows.

Proposition 3 The conjecture JN(n) holds for $N : \mathbb{C}^n \to \mathbb{C}^n$ of the form $N = N_{(k_1)} + N_{(k_2)}$, where $N_{(k_i)}$ are morphisms defined by homogeneous forms of degree $k_i > 0$.

Proof. The origin is a fixed point of N. Let $x = (x_1, ..., x_n) \neq o$ be another fixed point of N. As in the proof of Proposition 2 this contradicts the nilpotency of N, since the vector $\overline{x} = (x_1, ..., x_n)$ is an eigenvector (corresponding to a non-zero eigenvalue) of the linear mapping JN(X) at the point $X = \lambda x = (\lambda x_1, ..., \lambda x_n)$, where

$$\lambda = \left(\frac{k_1}{k_2}\right)^{\frac{1}{k_2 - k_1}}$$

In fact,

$$JN(\lambda x)(\overline{x}) = \left(\sum_{j=1}^{n} x_{j} \frac{\partial N_{(k_{1})1}}{\partial X_{j}}(\lambda x), \dots, \sum_{j=1}^{n} x_{j} \frac{\partial N_{(k_{1})n}}{\partial X_{j}}(\lambda x)\right) + \\ + \left(\sum_{j=1}^{n} x_{j} \frac{\partial N_{(k_{2})1}}{\partial X_{j}}(\lambda x), \dots, \sum_{j=1}^{n} x_{j} \frac{\partial N_{(k_{2})n}}{\partial X_{j}}(\lambda x)\right) = \\ = \left(k_{1}\lambda^{k_{1}-1}N_{(k_{1})1}(x), \dots, k_{1}\lambda^{k_{1}-1}N_{(k_{1})n}(x)\right) + \\ + \left(k_{2}\lambda^{k_{2}-1}N_{(k_{2})1}(x), \dots, k_{2}\lambda^{k_{2}-1}N_{(k_{2})n}(x)\right) = \\ = k_{1}\left(\frac{k_{1}}{k_{2}}\right)^{\frac{k_{1}-1}{k_{2}-k_{1}}} \cdot N_{(k_{1})}(x) + k_{2}\left(\frac{k_{1}}{k_{2}}\right)^{\frac{k_{2}-1}{k_{2}-k_{1}}} \cdot N_{(k_{2})}(x) =$$

$$= \frac{k_1^{\frac{k_2-1}{k_2-k_1}}}{k_2^{\frac{k_2-1}{k_2-k_1}}} \cdot N_{(k_1)}(x) + \frac{k_1^{\frac{k_2-1}{k_2-k_1}}}{k_2^{\frac{k_1-1}{k_2-k_1}}} \cdot N_{(k_2)}(x)$$

$$= \frac{k_1^{\frac{k_2-1}{k_2-k_1}}}{k_2^{\frac{k_2-1}{k_2-k_1}}} \cdot (N_{(k_1)}(x) + N_{(k_2)}(x))$$

$$= \frac{k_1^{\frac{k_1-1}{k_2-k_1}}}{k_2^{\frac{k_2-1}{k_2-k_1}}} \cdot \overline{x}.$$

2.4. The following result is a consequence of Theorem 3 and the proof of Theorem 8

Theorem 10 If each nilpotent morphism $N : \mathbb{C}^n \to \mathbb{C}^n$ of polynomial degree $\deg_p N = 3$ has a unique fixed point, then JN(n) holds.

In the case $\deg_p N = 3$ the morphism N is of the form

$$N = N_{(1)} + N_{(2)} + N_{(3)}$$

where $N_{(i)}$ is a homogeneous form of degree *i*, and Proposition 3 can be reformulated as follows:

The conjecture JN(n) holds for $N : \mathbb{C}^n \to \mathbb{C}^n$ of the form $N = N_{(1)} + N_{(3)}$ and $N = N_{(2)} + N_{(3)}$.

It follows from [6] that:

The conjecture JN(3) holds for $N : \mathbb{C}^3 \to \mathbb{C}^3$ of polynomial degree $\deg_p N \leq 3$.

2.5. The following proposition is a consequence of Theorem 7.

Proposition 4 Let $F : \mathbb{R}^n \to \mathbb{R}^n$ be a polynomial mapping of the form F = Id + N, where N is a positive nilpotent morphism. Then F is invertible.

Proof. Without loss of generality, we can assume that N(0) = 0. Consider a morphism $\overline{F} : \mathbb{R}^{n+1} \to \mathbb{R}^{n+1}$ defined by $\overline{F}_i = X_i - TN_i(X)$, i = 1, ..., n, and $\overline{F}_{n+1} = T$. By Theorem 7, \overline{F} is invertible. The restriction of \overline{F} to the hyperplane T = -1 coincides with F. Therefore, F is invertible.

In particular, it follows from this that:

The conjecture JN(n) holds for $N : \mathbb{R}^n \to \mathbb{R}^n$ if N is either a positive or a negative nilpotent morphism.

It follows from Theorem 7 (if we use a blow up) that:

The conjecture JN(n) holds for the nilpotent morphisms $N: \mathbb{R}^n \to \mathbb{R}^n$ of the form

$$N = \pm (N_{(1)} - N_{(2)} + \dots + (-1)^{m+1} N_{(m)}),$$

where $N_{(i)}$ are defined by positive homogeneous forms of degree *i*. **2.6.** Let r = rk JN(X) be the rank of a nilpotent morphism $N : \mathbb{C}^n \to \mathbb{C}^n$. In [2] it was shown that if r = 1, then F = Id - N is invertible:

The conjecture JN(n) holds for nilpotent morphisms $N : \mathbb{C}^n \to \mathbb{C}^n$ of rank r = 1. In particular, the conjecture JN(2) is true.

References

- H. Appelgate, H. Onishi, The Jacobian Conjecture in two variables // J.P.A.A., 37 (1985), 215-227.
- [2] H. Bass, E. Connell, D. Wright, The Jacobian Conjecture: Reduction of degree and formal expansion of the inverse // Bulletin of the AMS., 7, No. 2 (1982), 287-330.
- [3] L. Drużkowski, An affective approach to Keller's Jacobian Conjecture // Math. Ann. 264 (1983), 303-313.
- [4] A. van den Essen, Seven lectures on polynomial automorphisms // Univ. of Nijmegen, Report No. 9442 (October 1994).
- [5] A. van den Essen, Polynomial automorphisms and the Jacobian Conjecture // Univ. of Nijmegen, Report No. 9540 (October 1995).
- [6] E. Hubberts, The Jacobian Conjecture: Cubic homogeneous maps in dimension four // Master's thesis, Univ. of Nijmegen, Febr. 1994.
- [7] O. Keller, Ganze Cremona-Transformationen // Monatsh. Math. Phys. 47 (1939), 299-306.
- [8] J. Lang, S. Maslamani, Some results on the Jacobian Conjecture in higher dimension // J. of Pure and Applied Algebra 94 (1994), 327-330.
- [9] A. Magnus, On polynomial solutions of a differential equation // Math. Scand. 3 (1955), 255-260.
- [10] T.-T. Moh, On the global Jacobian Conjecture and the configuration of roots // J. Angew. Math. 340 (1983), 140-212.
- [11] S. Yu. Orevkov, On three-sheeted polynomial mappings of C² // Math. of the USSR, Izvestiya, vol. 29 (1987), 587-596.
- [12] A.G. Vitushkin, On polynomial transformation of Cⁿ manifolds (Tokyo, 1973) // Tokyo Univ. Press, Tokyo, 1975, 415-417.
- [13] S. Wang, A Jacobian criterion for separability // J. of Algebra 65 (1980), 453-494.
- [14] D. Wright, The Jacobian Conjecture: linear triangulation for cubics in dimension three // Linear and Multilinear Algebra 34 (1993), 85-97.
- [15] A. Yagzhev, On Keller's problem // Siberian Math. J. 21 (1980), 747-754.
- [16] J. Yu, On the Jacobian Conjecture: Reduction of coefficients // J. of Algebra 171 (1995), 515-523.

Steklov Mathematical Institute victor@olya.ips.ras.ru kulikov@mpim-bonn.mpg.de