

3-FOLD LOG FLIPS ACCORDING TO V. V. SHOKUROV

HIROMICHI TAKAGI*

ABSTRACT. We review §8 of V. V. Shokurov's paper "3-fold log flips".

CONTENTS

0. Introduction
1. Preliminaries
2. Auxiliary flips and back tracking method
3. Set up and dividing into cases
4. Treatment of case (a) and (b1)
5. Preliminaries for case (b2) and (c)
6. Treatment of case (b2, δ)
7. Treatment of case (c)

0. INTRODUCTION

In his paper '3-fold log flips', V. V. Shokurov proved the existence of the flips for flipping contractions from log 3-folds with only Kawamata log terminal singularities. First we review his strategy after 3 definition needed for explanation.

Definition (0.1) (Special flip) (See [S, Definition 6.1] or [FA, Definition 18.8]). Let X be a \mathbb{Q} -factorial n -fold and S a integral Weil divisor. Let $f : X \rightarrow Y$ be a small contraction with $\rho(X/Y) = 1$. We say that f is a special flipping contraction if

- (1) $K_X + S$ is LT;
- (2) $K_X + S$ and any component of S are f -negative.

We call the flip of such an f the special flip.

Definition (0.2) (Special termination). We say that the special termination holds if for any sequence of flips

$$(X_1, B_1) \dashrightarrow \cdots \dashrightarrow (X_k, B_k) \cdots,$$

where (X_i, B_i) is a LC pair, there is $k_0 \in \mathbb{N}$ (depending on a sequence) such that flipping curves on X_{k_0} are disjoint from $\perp B_{k_0} \lrcorner$.

Key words and phrases. Extremal ray, Good divisorial extraction, Log flip.

Mathematics Subject Classification. Primary : 14E30, Secondary : 14J35.

* Research Fellow of the Japan Society for the Promotion of Science

Definition (0.3) ($\overline{\mathcal{S}}_n^0(\text{local})$). We denote by $\overline{\mathcal{S}}_n^0(\text{local})$ the set of sequences (b_1, b_2, \dots, b_m) such that there is a \mathbb{Q} -factorial variety X of dimension at most n , a subset $Z \subset X$ and a divisor $B_0 + \sum b_i B_i$ ($B_0 \neq \phi$ is reduced but possibly reducible) such that every B_i intersects Z , $Z \subset B_0$, $K_X + B_0$ is PLT and $K_X + B_0 + \sum b_i B_i$ is maximally log canonical near Z .

Shokurov proved the following Reduction Theorem:

Theorem (0.4) (Reduction Theorem). *Let X be a n -dimensional normal variety and B a boundary such that $K_X + B$ is KLT. Let $f : X \rightarrow Y$ be a small contraction such that $-(K_X + B)$ is f -ample. Then the flip of f exists if the following three hold:*

- (1) (*Existence of Special flips*)
there exists the flip for any special flipping contraction of dimension n ;
- (2) (*Special termination*)
The special termination holds in dimension n ;
- (3) (*A.C.C for $\overline{\mathcal{S}}_n^0(\text{local})$*)
The ascending chain condition holds for the set $\overline{\mathcal{S}}_n^0(\text{local})$.

Proof. See [S, Theorem 6.4] or [FA, §18]. \square

In dimension 3, the special termination holds by [S, Theorem 4.1] or [FA, Theorem 7.1]. Furthermore the A.C.C holds for $\overline{\mathcal{S}}_3^0(\text{local})$ by [S, Chicago Lemma 4.9] or [FA, Proposition 18.19 and Corollary 18.25.1]. Hence in dimension 3, it suffices to prove the existence of the special flip. So we consider only the special flipping contraction in the below and we use the notation in Definition 0.1 for explanation.

First we can easily prove the existence in case S is reducible (See [FA, Proposition 21.2].) Hence we may assume that S is irreducible in the below. To flip in this case, Shokurov introduced the concept of the complement and divided special flipping contractions into cases using it:

Definition (0.5) (Complement) (See [S, §5] or [FA, §19]). Let X be a normal variety and B a subboundary on X . Let S be the smallest effective Weil divisor on X such that $\lfloor B - S \rfloor \leq 0$, and let $B_0 := B - S$. An n -complement of $K_X + B$ is a divisor $\overline{B} \in |-nK_X - nS - \lfloor (n+1)B_0 \rfloor|$ such that $K_X + B^+$ is LC, where $B^+ := S + \frac{1}{n}(\lfloor (n+1)B_0 \rfloor + \overline{B})$. We say $K_X + B$ is n -complemented if an n -complement exists.

The definition is arranged so that the vanishing theorem can be applied. But in the explanation below, we consider only the case B is a reduced boundary. So we restate the definition for this case.

Definition (0.5') (Complement). Let X be a normal variety and S a reduced boundary. An n -complement of $K_X + S$ is a divisor $\overline{B} \in |-nK_X - nS|$ such that $K_X + S + B$ is LC, where $B := \frac{1}{n}\overline{B}$. We say $K_X + B$ is n -complemented if an n -complement exists.

Then the following theorem holds:

Theorem (0.6). *Let $f : X \rightarrow Y$ be a 3-dimensional special flipping contraction such that S is irreducible. Then $K_X + S$ is n -complemented, where $n \in \{1, 2, 3, 4, 6\}$.*

Proof. See [S, Theorem 5.12] or [FA, Theorem 19.6 and Theorem 19.8]. \square

We call the minimum of n as in Theorem 0.6 the index of f .

In case the index is 1, it is easy to show the existence of the flip. (See [S, Proposition 6.8] or [FA, Proposition 21.4].)

In case the index is 2, if $K_X + S + B$ is LT, it is easy to prove the existence of the flip. (See below Proposition (2.2) which we quot from [S] and [FA].)

In case $n \geq 2$, Shokurov used the back tracking method (see [FA, §6 and §21] or the section 2 of this paper for the back tracking method) and proved the following:

Theorem (0.7). *If index 2 special flips exist, all special flips exist.*

Proof. See [S, Reduction 7.6] or [FA, Theorem 21.10]. \square

Hence the existence of index 2 special flips are left which we explain in this paper.

By some minor reductions, it suffices to consider the flipping contraction as in Set up (3.0). After some preparations in the sections 1 and 2, we start the proof of the existence of the special flip. The case (a) or (b) in Set up (3.0) is relatively easy to treat. (But the author had trouble understanding the proof of [S, Proposition 8.6] or [FA, Lemma 22.6] which is Theorem (4.0) in this paper.) The case (c) is very hard to treat and we have to divide it into finer cases. The proof in the section 7 is almost taken from [S, Proposition-Reduction 8.8] but we separate Shokurov's argument into Lemmas (Lemma 7.6 ~ 7.9 are principal ones). Futhermore the author cannot understand the inductive argument in [S, (8.8.4) and (8.8.5)] and hence find the alternative argument (see the proof of Theorem 7.1).

Acknowledgement. I express my hearty thanks to Professor Yujiro Kawamata to recommend me to write down this paper. This is the extended version of my talk in the seminar of the doctor course (1997) (I explained the contents of the section 0~6). I am grateful to the member of the seminar, Professor Yujiro Kawamata, Professor Keiji Oguiso, Doctor Nobuyoshi Takahashi, Mr. Tatsuhiro Minagawa and Mr. Kimikazu Kato for listening patiently to my talk and giving me useful comments.

Notation and Conventions.

- (1) In this paper, we will work over \mathbb{C} , the complex number field;
- (2) The \mathbb{P}^1 -bundle $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-a))$ over \mathbb{P}^1 , a Hirzebruch surface of degree a , is denoted by \mathbb{F}_a . Its unique negative section is denoted by C_0 and a ruling is denoted by f . The projective cone obtained from \mathbb{F}_a by the contraction of C_0 is denoted by $\mathbb{F}_{a,0}$. A generating line on $\mathbb{F}_{a,0}$ is denoted by l ;
- (3) we use the following abbreviations:
ODP for ordinary double point, LT for log terminal, PLT for purely log terminal, KLT for kawamata log terminal, DLT for divisorial log terminal, LC for log canonical, NC for normal crossing and MRS for minimal resolution.

1. PRELIMINARIES

In this section, we gather results used in the proof of the main theorem. The reader had better skip this section and refer later when the results in this section are quoted.

We start by easy lemmas about log surfaces.

Definition and Proposition (1.0). *Let $(S, K_S + B, s)$ be a germ of LC surface singularity. Then there is a unique extraction $\mu : T \rightarrow S$ satisfying the following conditions:*

- (1) $K_T + B_T = \mu^*(K_S + B)$ and $K_T + B_T$ is LT, where all exceptional curves appear in B_T with coefficients 1;
- (2) if there is a μ -exceptional (-1) -curve, then $K_T + B_T$ become non LT after contracting this.

We call this $\mu : T \rightarrow S$ the minimal LT model (abbreviated MLT) of $K_S + B$.

Proposition (1.1) (Description of LC surface 1). *Let $(S, K_S + B, s)$ be a germ of LC surface singularity. Then $\lfloor B \rfloor$ consists of at most 2 components. Furthermore if $\lfloor B \rfloor$ consists of 2 components, then $B = \lfloor B \rfloor$.*

Proof. See [Ka1, Theorem 9.6]. \square

We will repeatedly use this Proposition for 3-dimensional situations as follows:

Corollary (1.2). *Let $(X, K_X + D, x)$ be a 3-dimensional LC germ. Then every irreducible curve on X is contained in at most two reduced components of D . Furthermore if an irreducible curve is contained in two reduced components of D , the curve is contained in no other component of D .*

Proof. Take a general hyperplane section of X and apply (1.1). \square

Proposition (1.3) (Description of LC surface 2). *Let $(S, K_S + B, s)$ be a germ of LC surface singularity. Then for an exceptional curve F of MRS of s , we have $d(F, B) \leq 1$.*

Proof. Easy. \square

Proposition (1.4) (Description of LC surface 3). *Let $(S, K_S + B, s)$ be a germ of LC surface singularity such that $2(K_S + B) \sim 0$. Then its MLT is described as follow: FIGURE (I)*

Assume that $(S, K_S + B, s)$ is not LT. Then all exceptional curves of MRS except the end (-2) -curves of the dual graph are log crepant for $K_S + B$.

Proof. Easy. \square

Definition (1.5) (LLC, CLC and PLC) (See [Ka4, Definition 1.3]. Let X be a normal variety and D be a \mathbb{Q} -divisor such that $K_X + D$.

Then we define

CLC $(K_X + D)$ to be the set of subvarieties W on X such that there is a birational morphism from a normal variety $\mu : Y \rightarrow X$ and a prime divisor $E \subset Y$ with the discrepancy coefficient $e \leq -1$ such that $\mu(E) = W$.

We denote the set of m -dimensional elements in CLC $(K_X + D)$ by $\text{CLC}_m(K_X + D)$.

We define LLC $(K_X + D)$ to be the union of subvarieties in CLC $(K_X + D)$.

We define PLC $(K_X + D)$ to be the set of discrete valuations E in $\mathbb{C}(X)$ such that there is a birational morphism from a normal variety $\mu : Y \rightarrow X$ such that E is realized as a prime divisor on Y with the discrepancy coefficient $e \leq -1$.

The next two results describe LLC .

Theorem (1.6) (Connectedness Lemma (a)). *Let X and Y be normal varieties and $f : X \rightarrow Y$ a proper surjective morphism with only connected fibers. Let D be a \mathbb{Q} -divisor on X such that $K_X + D$ is \mathbb{Q} -Cartier. Write $D := \sum_{i=1}^n d_i D_i$, where D_i is an irreducible component of D .*

Assume the following:

- (1) if $d_i < 0$, $\text{codim } f(D_i) \geq 2$;
- (2) $-(K_X + D)$ is f -nef and f -big;

Then for any point $y \in Y$, $\text{LLC}(K_X + D) \cap f^{-1}(y)$ is connected.

Proof. See [S, 5.7] or [FA, 17.4 Theorem] and also [Ka, Theorem 1.4]. \square

Theorem (1.7) (Connectedness Lemma (b)). *Let (S, Θ) be a proper LC surface. Assume that $K_S + \Theta \equiv 0$. Then one of the following holds:*

- (1) $\perp\Theta\perp$ is irreducible and there is no point contained in $CLC_0(K_S + \Theta)$ on $\perp\Theta\perp$;
- (2) $\perp\Theta\perp$ is connected and there is a point contained in $CLC_0(K_S + \Theta)$ on every component of $\perp\Theta\perp$. Furthermore $LLC(K_S + \Theta) = \perp\Theta\perp$;
- (3) S is a generically \mathbb{P}^1 -bundle over a smooth rational or elliptic curve and $\perp\Theta\perp = C_1 \cup C_2$, where C_1 and C_2 are two disjoint sections of above generically \mathbb{P}^1 -bundle. There is no point contained in $CLC_0(K_S + \Theta)$ on C_1 or C_2 . Furthermore $LLC(K_S + \Theta) = \perp\Theta\perp$.

Proof. See [S, Theorem 6.9] or [FA, 12.3.1 Proposition]. \square

Lemma (1.8) (Ampleness). *Let $f : S \dashrightarrow T$ be a birational map of normal projective surfaces and D an effective ample divisor on S such that D_T is irreducible. (D_T is the log birational transform, i.e., all the blown up curve are contained in it with multiplicity 1.) Then D_T is numerically ample.*

Proof. See [S, 8.10 Lemma]. \square

The following theorem is slightly weaker than [S, Lemma 8.9]. But in our application, it is sufficient.

Theorem (1.9) (CLC₀ for log surface). *Let (S, B) be a projective LC surface and C_1 and C_2 connected contractible curves. (Caution: we allow that C_i is empty. The contractibility doesn't mean C_1 and C_2 are contractible simultaneously.) We assume the following:*

- (i) $2(K_S + B) \sim 0$;
- (ii) $\perp B \perp = B_1 + B_2$, where B_i is an irreducible curve and $B_1^2 > 0$ and B_2 is ample;
- (iii) $B_1 \cap C_1 = \emptyset$;
- (iv) $P := B_1 \cap B_2$ is one point and on B_1 , P is the unique point contained in $CLC_0(K_S + B)$;
- (v) all the component of C_2 pass through P .

(See FIGURE (II).) Then $C_1 \cap B_2 \notin CLC_0(K_S + B)$.

Proof. First note that if $C_1 \cap B_2$ (resp. $C_2 \cap (B_1 \cup B_2)$) is not empty, $C_1 \cap B_2$ (resp. $C_2 \cap (B_1 \cup B_2)$) is one point by (1.6).

Let $\mu : T \rightarrow S$ be the MRS of S .

Claim 1. *Let C'_2 be any component of C_2 . Then*

- (1) $\mu^{-1}C'_2$ is a (-1) -curve;
- (2) P is singular point of S . Outside P , there are only Du Val singularities on C'_2 ;
- (3) $C'_2 \cap \text{Supp } \{B\} = \emptyset$.

Proof. By the assumption of (v), C'_2 passes through P and there are two reduced boundaries of $K_S + B$ through P . Hence $C'_2 \not\subset \text{Supp } B$. So $K_S.C'_2 = -B.C \leq -(B_1 + B_2).C'_1 < 0$, which shows (1).

If P is nonsingular on S , we obtain $K_S.C'_2 \leq -(B_1 + B_2).C'_2 \leq -2$, a contradiction to the contractibility of C'_2 . Hence P is singular.

Define B_T by $K_T + B_T = \mu^*(K_S + B)$. Then $B_T.\mu^{-1}C'_2 \geq 1$ since $\mu^{-1}C'_2$ intersects log crepant curves over P . But by (1), this must be equality. From this we can deduce the latter half of (2) and (3). \square

Let $\nu : U \rightarrow S$ be MLT for $K_S + B$.

Below we assume that $C_1 \cap B_2 \in CLC_0(K_S + B)$ to get a contradiction.

Claim 2. $C_1 \cap C_2 = \phi$ (See FIGURE (III).)

Proof. Assume there is a component C'_2 of C_2 such that $C_1 \cap C'_2 \neq \phi$. Let C'_1 be a component of C_1 such that $C'_1 \cap C'_2 \neq \phi$. Then by Claim 1, $C'_1 \not\subset \text{Supp } \{B\}$. Hence by (ii) and (iii), $K_S \cdot C'_1 \leq -B_2 \cdot C'_1 < 0$, which shows that $\mu^{-1}C'_1$ is a (-1) -curve. On the other hand, $\nu^{-1}C'_1 \cap \nu^{-1}C'_2 \neq \phi$ since by Claim 1, $C'_1 \cap C'_2 (\neq \phi)$ is at worst Du Val singularity and not contained in $\text{Supp } B$.

Furthermore we know that $C'_1 \cap B_2$ is singular as follow:

If $C'_1 \cap B_2$ is smooth, we have $K_S \cdot C'_1 \leq -B_2 \cdot C'_1 \leq -1$ by $C'_1 \not\subset \text{Supp } B$. So this must be equality since $\mu^{-1}C'_1$ is a (-1) -curve. Hence $C'_1 \cap \text{Supp } \{B\} = \phi$. But this shows that $C'_1 \cap B_2 \notin \text{CLC}_0(K_S + B)$, a contradiction.

Next we claim that $(\nu^{-1}B_1)^2 < 0$ as follows:

By the assumption (iii), $\nu^{-1}B_1 \cap \nu^{-1}C'_1 = \phi$. By Claim 1 (2), $\nu^{-1}B_1 \cap \nu^{-1}C'_2 = \phi$. Clearly ν -exceptional curves over $C'_1 \cap B_2$ and $\nu^{-1}B_1$ doesn't intersect. Now contract ν -exceptional curves over $C'_1 \cap C'_2$ and $\nu^{-1}C'_1$ then $(\nu^{-1}C'_2)^2$ becomes ≥ 0 (cf. Claim 1 (2) and the fact $\mu^{-1}C$ is a (-1) -curve). Next contract ν -exceptional curves over $C'_1 \cap B_2$ (Note that there exist such exceptional curve by Claim 1 (2) and the fact $P \in \text{CLC}_0(K_S + B)$). then $(\nu^{-1}C'_2)^2$ becomes > 0 . Even after such contractions, $\nu^{-1}B_1$ and $\nu^{-1}C'_2$ doesn't intersect since such contractions are isomorphic near $\nu^{-1}B_1$. Hence by Hodge index theorem, $(\nu^{-1}B_1)^2 < 0$.

By the way we know that $B_1 \simeq \mathbb{P}^1$ and $K_S + B|_{B_1} = K_{B_1} + \frac{1}{2}P_1 + \frac{1}{2}P_2 + P$ by the assumption (iv).

Here we claim that one of P_1 and P_2 , say P_1 is smooth and P_2 is an ODP as follow: If both P_1 and P_2 are singular, by the above subadjunction they are ODP's. Hence we can write

$$\mu^*B_1 = \mu^{-1}B_1 + \sum e_j E_j + \frac{1}{2}E^1 + \frac{1}{2}E^2,$$

where E^i is the exceptional curve resolving P_i and E_j 's are exceptional curves over P and $e_j \leq 1$. (We assume that E_1 intersects $\mu^{-1}B_1$.) By $(B_1)^2 > 0$ and $(\mu^{-1}B_1)^2 < 0$, $0 < (\mu^{-1}B_1)^2 + e_1 + 1$ and hence $\mu^{-1}B_1$ is a (-1) -curve. On the other hand since $\nu^{-1}B_1$ is also (possibly analytically) contractible, E^1 , $\mu^{-1}B_1$ and E^2 is a contractible chain. But this is absurd. By similar calculation, P_2 must be an ODP.

By the above subadjunction, we know that there is a component B_3 of $\{B\}$ which passes through P_1 .

Next we claim that $(\nu^{-1}B_3)^2 < 0$ below. First by Claim 1, $\nu^{-1}B_3 \cap \nu^{-1}C'_2 = \phi$. Furthermore we obtain also $\nu^{-1}B_3 \cap \nu^{-1}C'_1 = \phi$ as follows:

Looking the degree of $K_S + B|_{B_2}$, Only P and $B_2 \cap C'_1$ can contribute to its boundary. On the other hand by the ampleness of B_2 , $B_2 \cap B_3 \neq \phi$ and hence $B_2 \cap B_3 = B_2 \cap C'_1$. Consider the description of MLT over $B_2 \cap B_3$. If $\nu^{-1}B_3 \cap \nu^{-1}C'_1 \neq \phi$, we have $K_T \cdot \mu^{-1}C'_1 \leq -\frac{3}{2}$, a contradiction to that $\mu^{-1}C'_1$ is a (-1) -curve. Hence $\nu^{-1}B_3 \cap \nu^{-1}C'_1 = \phi$. Now contract first $\nu^{-1}C'_1$. Then $(\nu^{-1}C'_2)^2$ become ≥ 0 . Next contract ν -exceptional curve over P . Then $(\nu^{-1}C'_2)^2$ become > 0 . These contractions are isomorphic near $\nu^{-1}B_3$ since $\nu^{-1}B_3 \cap \nu^{-1}C'_1 = \phi$ and hence even after the contractions, $\nu^{-1}B_3 \cap \nu^{-1}C'_2$ remain empty. Hence by Hodge index theorem, $(\nu^{-1}B_3)^2 < 0$.

But this gives the final contradiction. For $\nu^{-1}B_3$ intersects at least one log crepant curve over $B_2 \cap B_3$ and also $\nu^{-1}B_1$ at another point. Hence applying (1.6) for the contraction of B_3 , we get a contradiction. We obtain Claim 2. \square

By this Claim 2, even after contracting C_2 , every assumption of (1.9) and assumption that $C_1 \cap B_2 \in \text{CLC}_0(K_S + B)$ hold by setting $C_2 = \phi$. Hence it is sufficient to get a contradiction assuming also $C_2 = \phi$.

Consider extremal rays for $K_S + (1 - \epsilon)(B - B_1)$, where ϵ is a sufficiently small positive rational number. Note that $K_S + (1 - \epsilon)(B - B_1)$ is KLT by (1.7).

Claim 3. *There is an extremal ray R for $K_S + (1 - \epsilon)(B - B_1)$ such that $B_1.R > 0$.*

Proof. Assume that any extremal ray R' for $K_S + (1 - \epsilon)(B - B_1)$ satisfies $B_1.R' \leq 0$. For an irreducible curve l such that $(K_S + (1 - \epsilon)(B - B_1)).l \geq 0$, we have $((1 - \epsilon)B_1 + \epsilon B).l \leq 0$. Since ϵ is sufficiently small, we have $B_1.l \leq 0$. But by the cone theorem, for every curve l , we must have $B_1.l \leq 0$, a contradiction to $(B_1)^2 > 0$. \square

Fix a extremal ray R as in Claim 3.

If the contraction of R is birational, by the choice of R and ampleness of B_2 , $B_1 \cap R \neq \phi$ and $B_2 \cap R \neq \phi$. By (1.6), $B_1 \cap R = B_2 \cap R = \{P\}$. Now set $C_2 = \text{Supp } R$. Then again all the assumptions of this theorem and the assumption $C_1 \cap B_2 \in \text{CLC}_0(K_S + B)$ hold. Hence by considering inductively, we have the maximal composition of birational extremal contractions $S \rightarrow \bar{S}$ such that all extremal ray intersect B_1 positively and for \bar{S} , all the assumptions of this theorem and the assumption $\bar{C}_1 \cap \bar{B}_2 \in \text{CLC}_0(K_{\bar{S}} + \bar{B})$ hold. (We denote all the image on \bar{S} with $\bar{\cdot}$.) For this \bar{S} , we can take an extremal ray \bar{R} as in Claim 3. By the choice of \bar{S} , the contraction of \bar{R} is not birational. By (iii), it is not a contraction to a point. Hence it is a contraction to a curve. Let l be its fiber such that $\bar{B}_2 \cap \bar{C}_1 \in l$. By $\bar{B}_1.l > 0$, $\bar{B}_1 \cap l \neq \phi$. Let $\bar{\nu} : \bar{T} \rightarrow \bar{S}$ be MLT for $K_{\bar{S}} + \bar{B}$. Then $(\bar{\nu}^{-1}l)^2 < 0$ since $\bar{B}_2 \cap \bar{C}_1$ is contained in $\text{CLC}_0(K_{\bar{S}} + \bar{B})$ and singular. Note that $\bar{B}_1 \cap l \neq \bar{B}_2 \cap \bar{C}_1$ and $l \not\subset \text{LLC } \bar{S}\bar{B}$. Furthermore $\bar{\nu}^{-1}l$ intersects $\text{LLC } \bar{\nu}^*(K_{\bar{S}} + \bar{B})$ at two point (and is not contained in it), a contradiction to (1.6). We finish the proof of this theorem. \square

The following proposition is used in the treatment of case (b) in Set up (3.0).

Proposition (1.10). *Let $(S, K_S + B, s)$ be a pointed LC surface. Assume that $C := \lrcorner B \lrcorner$ is a smooth projective curve. Let $a_i : S_i \rightarrow S$ for $i = 1, 2$ be a birational morphism satisfying the following:*

- (1) a_i ($i = 1, 2$) is log crepant for $K_S + B$ and the exceptional curve F_i for a_i is irreducible;
- (2) $\text{CLC}_0(a_1^*(K_S + B)) \cap F_1 = a_1^{-1}C \cap F_1$;
- (3) $a_2^*(K_S + B)$ is LT at $a_2^{-1}C \cap F_2$ (and hence S_2 is smooth at $a_2^{-1}C \cap F_2$).

Then $(a_2^{-1}C)^2 > 0$ (resp. $(a_2^{-1}C)^2 \geq 0$) implies that $(a_1^{-1}C)^2 > 0$ (resp. $(a_1^{-1}C)^2 \geq 0$).

Proof. Let $m : \tilde{S}_1 \rightarrow S_1$ be the MLT for $a_1^*(K_S + B)$ at $m^{-1}C \cap F_1$. By this and assumption (2), the assumption that C is irreducible and the uniqueness of MLT (1.0), $a_1 \circ m$ must be the MLT for $K_S + B$. Let $n : \tilde{S}_2 \rightarrow S_2$ be the MLT for $a_2^*(K_S + B)$ near F_2 . Possibly after contracting some $a_2 \circ n$ -exceptional curves, we obtain the MLT for $K_S + B$. Assume that $(a_2^{-1}C)^2 > 0$ (resp. $(a_2^{-1}C)^2 \geq 0$). Then by assumption (3), there is no n -exceptional curve over $a_2^{-1}C \cap F_2$ and hence $((a_2 \circ m)^{-1}C)^2 > 0$ (resp. $((a_2 \circ m)^{-1}C)^2 \geq 0$). Since the MLT for $K_S + B$ is dominated by \tilde{S}_2 , we have also $((a_1 \circ n)^{-1}C)^2 > 0$ (resp. $((a_1 \circ n)^{-1}C)^2 \geq 0$) and hence $(a_1^{-1}C)^2 > 0$ (resp. $(a_1^{-1}C)^2 \geq 0$). \square

Proposition (1.11) (Smoothness). *Let X be a normal variety and $K_X + S + B$ is LT, where S is a \mathbb{Q} -Cartier integral Weil divisor which contains no codimension 2 singular locus. Assume that S is nonsingular. Then X is nonsingular.*

Proof. See [S, 3.7, Corollary]. \square

Proposition (1.12) (Classification of LT singularity). *Let $(X, K_X + B, x)$ be a germ of a normal LT 3-fold such that $K_X + \lrcorner B \lrcorner$ is \mathbb{Q} -Cartier (and hence LT). Then*

- (1) *if B has at least 3 reduced components which are \mathbb{Q} -Cartier, then locally analytically*

$$(x \in B \subset X) \simeq (o \in (xyz = 0) \subset \mathbb{C}^3);$$

(2) if B has 2 components which are \mathbb{Q} -Cartier, then locally analytically

$$(x \in \perp B \lrcorner \subset X) \simeq (o \in (xy = 0) \subset \mathbb{C}^3 / \mathbb{Z}_n(q_1, q_2, 1),$$

where q_1 and q_2 are natural numbers such that $(q_1, q_2, 1) = 1$.

Proof. In [FA,16.10 Corollary] and [FA,16.15 Theorem], this theorem is proved under the assumption that $K_X + B$ is DLT. But as we will see below, we can prove under the above assumptions that (X, x) is analytically \mathbb{Q} -factorial and hence $K_X + \perp B \lrcorner$ is DLT.

If (X, x) is not analytically \mathbb{Q} -factorial, then we can take a small analytic \mathbb{Q} -factorialization $\pi \tilde{X} \rightarrow X$ by applying (1.13) for $K_X + B_1$. In fact everything is clear except that π is small. Since π is log crepant for $K_X + B_1$ and $K_X + B_1$ is PLT, it must be small. Let B_1 and B_2 be two of reduced components in $\perp B \lrcorner$ which are \mathbb{Q} -Cartier. Since B_i 's are \mathbb{Q} -Cartier, exceptional curves of π are contained in $\pi^{-1}B_i$'s. Hence $\pi^{-1}B_1 \cap \pi^{-1}B_2$ is not irreducible. Now we apply [FA,16.15 Theorem] for $K_{\tilde{X}} + \pi^{-1}B_1 + \pi^{-1}B_2$, a contradiction.

Hence we can directly apply [FA,16.10 Corollary] and [FA,16.15 Theorem] for $K_X + \perp B \lrcorner$. Note that in (1), $B = \perp B \lrcorner$. In fact the simple blow up of x is log crepant for $K_X + \perp B \lrcorner$ and hence there can be no component other than $\perp B \lrcorner$ since $K_X + B$ is also LT. \square

Theorem (1.13) (\mathbb{Q} -factorial LT model). *Let X be a 3-fold and B a boundary on X . Assume that B is \mathbb{Q} -Cartier and $\perp B \lrcorner$ is an LSEPD divisor, i.e., there is a principal divisor (h) near $\text{Supp } \perp B \lrcorner$ such that $\text{Supp } (h) = \text{Supp } \perp B \lrcorner$. Then in a neighborhood of $\perp B \lrcorner$, there exists a \mathbb{Q} -factorial LT model for $K_X + B$, i.e., there is a birational morphism $f : Y \rightarrow X$ such that $K_Y + B_Y$ is f -nef and over a neighborhood of $\perp B \lrcorner$, $K_Y + B_Y$ is LT, where B_Y has all f -exceptional divisors with coefficients 1.*

Proof. See [S, Corollary 5.19] or [FA, Corollary 20.9]. \square

2. AUXILIARY FLIPS AND BACK TRACKING METHOD

Proposition (2.0) (Flip of type 1). *Let X be a \mathbb{Q} -factorial 3-fold. Let $D = S + B$ be a \mathbb{Q} -divisor, where S is reduced and $\perp B \lrcorner = 0$. Let $f : X \rightarrow Y$ be a small contraction with $\rho(X/Y) = 1$ and C its exceptional curve. Assume that*

- (1) $K_X + D$ is LC and $(K_X + D).C \leq 0$;
- (2) S has at least 2 irreducible components S_1 and S_2 such that $S_1.C < 0$ and $S_2.C > 0$.

Then the flip of f exists.

Proof. See [FA, 20.7 Theorem]. \square

Proposition (2.1) (Flip of type 2). *Let $f : X \rightarrow Y$ be a small contraction and C its exceptional curve. Let S_1 and S_2 are effective divisors such that $S_1 \cap S_2 = C$. Assume that for some $m_1, m_2 \in \mathbb{N}$, $m_1 S_1 \sim m_2 S_2$ near C . Then the flip of f exists.*

Proof. See [FA, 20.11 Theorem]. \square

Proposition (2.2) (Flip of type 3). *Let X be a \mathbb{Q} -factorial 3-fold. Let $D = S + B$ be a \mathbb{Q} -divisor, where S is reduced. Let $f : X \rightarrow Y$ be a small contraction with $\rho(X/Y) = 1$ and C its exceptional curve. Assume that*

- (1) f is special with respect to $K_X + S$;
- (2) $2(K_X + D) \sim 0$ near C ;
- (3) $K_X + D$ is LT;

Then the flip of f exists.

Proof. See [FA, Proposition 21.5]. \square

Proposition (2.3) (Flip of type 4). *Let X be a \mathbb{Q} -factorial 3-fold. Let $D = S + B$ be a \mathbb{Q} -divisor, where S is irreducible and reduced. Let $f : X \rightarrow Y$ be a small contraction with $\rho(X/Y) = 1$ and C its exceptional curve. Assume that*

- (1) f is special with respect to $K_X + S$;
- (2) $2(K_X + D) \sim 0$ near C ;
- (3) $K_X + D$ is LC;
- (4) $K_X + D|_S$ is exceptional, i.e., there is exactly one divisor in $PLC(K_X + D|_S)$;

Then the flip of f exists.

Proof. See [FA, 21.1.2.1 Corollary]. \square

Theorem (2.4). *Let X be an algebraic 3-fold, S_1 and S_2 irreducible \mathbb{Q} -Cartier surfaces. Let $f : X \rightarrow Y$ be a small contraction. We assume the following:*

- (a) $K_X + S_1 + S_2$ is LC;
- (b) $C := \text{excep } f$ is an irreducible curve and $C \subset S_1 \cap S_2$;
- (c) $(K_X + S_1).C < 0$, $S_1.C < 0$ and $(K_X + S_1 + S_2).C = 0$;
- (d) $2(K_X + S_1 + S_2) \sim 0$ near C ;
- (e) Along C , S_1 and S_2 are generically normal crossing and S_1 is normal;
- (f) $K_X + S_1 + S_2|_C = K_C + \frac{1}{2}P_1 + \frac{1}{2}P_2 + P$, where P_1 , P_2 and P are distinct points.

Then the flip of f exists. Let $f^+ : X^+ \rightarrow Y$ be the flip and C^+ the exceptional curve for f^+ . We denote all the strict transforms of divisors on X with superscript $+$. Then we obtain the following descriptions;

- (1) One of P_1 and P_2 , say P_1 is an ODP on S_1 . P_2 is a smooth point on S_1 . P is smooth or a singularity with the following description on the MRS of S_1 : FIGURE (IV) of type $(3, 2, 2, 2, 2, \dots, 2)$;
- (2) Near P , $S_2|_{S_1} = C + C'$ as a Weil divisor on S_1 , where C' is a smooth curve. In particular along C' , S_1 and S_2 are also generically normal crossing and $K_X + S_1 + S_2$ is LT outside P ;
- (3) The natural birational map $S_1 \dashrightarrow (S_1)^+$ is a morphism whose exceptional locus is C . The image point of C (we call this Q) is smooth on $(S_1)^+$;
- (4) C^+ is irreducible and near Q , C^+ coincides with the nonnormal locus of S_2^+ . Let $\nu : S_2^{+\nu} \rightarrow S_2^+$ be the normalization. Then $\nu^{-1}Q$ is one point and a nonsingular point of $S_2^{+\nu}$.

(See FIGURE (VI).)

Proof. Since f is a flipping contraction of type 1, the flip exists by (2.0). We prove the descriptions (1) \sim (4) below.

First we prove (1) and (2).

Claim 1. $S_1 \cap S_2$ is not irreducible.

Proof. If $S_1 \cap S_2$ is irreducible, $C_{S_1}^2 = S_2.C > 0$ by the assumption (c) and (e). But this contradicts the contractibility of C in S_1 . \square

By LC property of $K_X + S_1 + S_2|_{S_1}$ and Claim 1, there are two components of $S_1 \cap S_2$ at P . Let C' be the component other than C . By [FA, 16.6.3] and the assumption (e), we can write $S_2|_{S_1} = C + \frac{1}{m}C'$, where m is a natural number. We prove the description (1). Let $\mu : (S_1)^\mu \rightarrow S_1$ be the MRS. Assume that both P_1 and P_2 are singular, i.e., ODP's by (f). Let

$$(\star) \quad \mu^*(C + \frac{1}{m}C') = \mu^{-1}C + \frac{1}{m}\mu^{-1}C' + \frac{1}{2}E^1 + \frac{1}{2}E^2 + \sum_{i=1}^n e_i E_i,$$

where E^i is the (-2) -curve resolving P_i and E_j 's are exceptional curves resolving P and E_1 is the nearest component to $\mu^{-1}C$. We note that $e_i \leq 1$ by the minimality of μ and LC property of $K_{S_1} + C + \frac{1}{m}C'$. Intersecting \star with $\mu^{-1}C$, we obtain

$$0 < S_2 \cdot C = (\mu^{-1}C)^2 + 1 + \begin{cases} \frac{1}{m} & \text{if } n = 0 \\ e_1 & \text{if } n \geq 1. \end{cases}$$

Hence by this and the contractibility of $\mu^{-1}C$, $(\mu^{-1}C)^2 = -1$. But we obtain a non-contractible chain consisting of E^1 , $\mu^{-1}C$ and E^2 , a contradiction.

Assume that both P_1 and P_2 are smooth. Then by similar calculations, we obtain $(\mu^{-1}C)^2 \geq 0$, a contradiction.

Hence we proved the description about P_1 and P_2 .

Furthermore by the similar calculations, we obtain $(\mu^{-1}C)^2 = -1$ and

$$\begin{cases} \frac{1}{m} > \frac{1}{2} & \text{if } n = 1 \\ e_1 > \frac{1}{2} & \text{if } n \geq 1. \end{cases}$$

If $n = 1$, we immediately obtain (1) and (2).

If $n \geq 1$, we see that $E_1^2 \leq -3$ since otherwise E^1 , $\mu^{-1}C$ and E_1 construct a non-contractible chain. Based on this and $e_1 > \frac{1}{2}$, we obtain P is a singularity as in the statement of (1) and

$$\mu^*C = \mu^{-1}C + \frac{1}{2}E^1 + \sum_{i=1}^n \frac{n-i+1}{2n+1}E_i$$

and

$$\mu^*C' = \mu^{-1}C' + \sum_{i=1}^n \frac{2i-1}{2n+1}E_i$$

by elementary calculations. Hence $e_1 = \frac{n}{2n+1} + \frac{1}{m} \frac{1}{2n+1} > \frac{1}{2}$, i.e., $m = 1$. So in this case we proved (1) and (2).

Next we prove (3). By (1) and (2), $f(C)$ is a smooth point of $f(S_1)$. In particular $f(S_1)$ is normal. Hence if we prove that no component of C^+ is contained in S_1^+ , then by the Zariski's Main Theorem, $S_1^+ \simeq f(S_1)$ and (3) follows. Note that $K_X + S_1 + S_2|_{S_1} = K_{S_1} + C + \frac{1}{2}C'' + C'$ by (f) and (1), where C'' is an irreducible curve which crosses normally with C at P_2 . Here $K_Y + f(S_1) + f(S_2)|_{f(S_1)} = K_{f(S_1)} + \frac{1}{2}f(C'') + f(C')$ is LC since $K_X + S_1 + S_2$ is f -trivial. Hence $K_Y + f(S_1)|_{f(S_1)} = K_{f(S_1)} + \frac{1}{2}f(C'')$ is canonical since $f(C)$ is smooth on $f(S_1)$. By this and the f'^+ -ampleness of $K_{X^+} + S_1^+|_{S_1^+}$ (we define that $f'^+ := f^+|_{S_1^+}$), S_1^+ contains no exceptional curve of f . Hence we finish the proof of (3).

Next we prove (4). Since S_1^+ is \mathbb{Q} -Cartier, any analytic branch of S_2^+ intersects S_1^+ along curves. On the other hand by (2) and (3), $S_1^+ \cap S_2^+ = C'^+$ and the intersection is generically NC so the number of analytic branch of S_2^+ is 1, i.e., $\nu^{-1}Q$ is one point.

Claim 2. *The index of $K_{X^+} + S_2^+$ at Q is 1.*

Proof. Assume the contrary. Let $\pi : \tilde{X}^+ \rightarrow X^+$ be the index 1 cover of $K_{X^+} + S_2^+$. On S_1^+ , X^+ has only generically ODP curve singularity (along C''^+) near Q . Hence $\pi^{-1}S_1^+ \rightarrow S_1^+$ is only ramified at Q near Q . But S_1^+ is smooth at Q so $\pi^{-1}S_1^+$ has at least 2 components intersecting only at $\pi^{-1}Q$. Since $K_{\tilde{X}^+} + \pi^{-1}S_1^+$ is PLT by (c) and (2), we can take a small \mathbb{Q} -factorialization $\rho : \bar{X}^+ \rightarrow \tilde{X}^+$ by (1.13). The exceptional curves of ρ are contained in $\rho^{-1}\pi^{-1}S_1$ and $\rho^{-1}\pi^{-1}S_2$ (they have at most 3 components in all.) But this contradicts LC property of $K_{\bar{X}^+} + \rho^{-1}\pi^{-1}S_1^+ + \rho^{-1}\pi^{-1}S_2^+$ along the exceptional curves of ρ . Hence we obtain Claim 2. \square

Claim 3. *Near $\nu^{-1}Q$, $K_{X^+} + S_1^+ + S_2^+|_{S_2^{+\nu}} = K_{S_2^{+\nu}} + C^* + C'^+$, where C^* is some curve through $\nu^{-1}Q$. (We will prove later that $C^* := \nu^{-1}C^+$.)*

Proof. Since $K_{X^+} + S_1^+ + S_2^+|_{S_2^{+\nu}}$ is LC at $\nu^{-1}Q$ and $K_{X^+} + S_2^+|_{S_2^{+\nu}}$ is LT of index 1 at $\nu^{-1}Q$ by Claim 2, the boundary near $\nu^{-1}Q$ is reduced and the number of it is at most 1. Hence we have only to prove that $K_{X^+} + S_2^+|_{S_2^{+\nu}} = K_{S_2^{+\nu}} + C^*$ for some irreducible curve C^* . Assume the contrary. Then $K_{X^+} + S_2^+|_{S_2^{+\nu}} = K_{S_2^{+\nu}}$. By this, we know that S_2^+ is normal at Q . Since $K_{X^+} + S_1^+ + S_2^+|_{S_2^+} = K_{S_2^+} + C'^+$ and $K_{S_2^+}$ is Cartier at Q , $K_{S_2^+}$ is canonical at Q .

On the other hand $K_{X^+} + S_1^+ + S_2^+|_{S_2^+}$ is not LT. In fact if it is LT, it is PLT. Hence its restriction to C'^+ is KLT and by the inversion of adjunction [FA, 17.6 Theorem], $K_{X^+} + S_1^+ + S_2^+|_{S_2^+}$ is PLT. But this contradicts the description (1) and (2) (see FIGURE (V)).

So by (1.12), Q is a canonical singularity of type D_m for some m . Let C_1^+ be any component of C^+ and $\sigma : S_2^{+\sigma} \rightarrow S_2$ the MRS. Since $K_{S_2^+}.C_1^+ = (K_{X^+} + S_2^+).C_1^+ < 0$, C_1^+ is (-1) -curve on the MRS of S_2 . But the union of the exceptional curves of MRS of a canonical singularity of type D_m and C_1^+ cannot be contracted at one time, a contradiction. \square

By Claim 2 and Claim 3, we see that $S_2^{+\nu}$ is smooth at $\nu^{-1}Q$.

In fact it is derived by the fact that $K_{S_2^{+\nu}} + C^*$ is Cartier and LT.

By Claim 3, $\nu(C^*)$ coincides with the nonnormal locus near Q . Since S_1^+ is normal outside C^+ by (e) and all the components of C^+ pass through Q , every 1 dimensional component of the nonnormal locus of S_1^+ is contained in C^+ (and passes through Q). Hence $\nu(C^*) \subset C^+$. If there is a component C_1^+ of C^+ which is not contained in $\nu(C^*)$, we have $K_{S_2^{+\nu}}.\nu^{-1}C_1^+ = -(C^* + C'^+).\nu^{-1}C_1^+ \leq -2$, a contradiction to the contractibility of $\nu^{-1}C_1^+$ in $S_2^{+\nu}$. Hence $\nu(C^*) = C^+$ and in particular C^+ is irreducible.

Now we finish the proof of (2.4). \square

Description of BTM (2.5). We review the back tracking method for introducing the notation. The material here is contained in [FA, Chapter 21]. Let X be an algebraic \mathbb{Q} -factorial normal 3-fold and $f : X \rightarrow Y$ a projective small contraction such that $\rho(X/Y) = 1$. Let S be an irreducible divisor on X , B a boundary and C the exceptional curve of f . We assume the following assumptions.

- (i) $K_X + S$ is PLT and $K_X + S + B$ is LC.
- (ii) $(K_X + S).C < 0$, $S.C < 0$ and $(K_X + S + B).C = 0$.

Then we can make the following procedure:

Choose a primitive log crepant divisorial extraction $g_1 : Z_1 \rightarrow X$ for $K_X + S + B$. Let E_1 be the exceptional divisor.

By assumption, $K_{Z_1} + S_1 + B_1 + E_1 = g_1^*(K_X + S + B)$, where S_1 and B_1 are the strict transforms of S and B . Hence by choosing a rational number b ($0 < b < 1$) appropriately, we have $K_{Z_1} + S_1 + bB_1 + e'E_1 = g_1^*(K_X + S + bB)$, where $e' \in (0, 1)$ is a rational number. Here note that $(K_X + S + bB).C < 0$ by the assumption (2). So by choosing a rational number $e \in (e', 1)$, $-(K_{Z_1} + S_1 + bB_1 + eE_1)$ become $f \circ g$ -ample. Furthermore $K_{Z_1} + S_1 + bB_1 + eE_1$ is PLT by the following argument:

By (i), $K_X + S + bB$ is PLT. Hence $K_{Z_1} + S_1 + bB_1 + e'E_1$ is also PLT. The PLT property is an open one so $K_{Z_1} + S_1 + bB_1 + eE_1$ is also PLT.

Consequently we can run the $K_{Z_1} + S_1 + bB_1 + eE_1$ -MMP over Y . Since $\rho(Z_1/Y) = 2$, the

surgeries are uniquely determined once we choose g_1 . We obtain the following diagram:

$$\begin{array}{ccccccc}
 & & Z_1 & \dashrightarrow & Z_2 & \dashrightarrow & \\
 & \swarrow g_1 & & h_1 \searrow & \swarrow g_2 & h_2 \searrow & \\
 X & & & X_1 & & X_2 \dots & \\
 & \searrow f & & & \searrow & & \\
 & & Y & \swarrow & & Y & .
 \end{array}$$

We denote the extremal ray associated to h_i by R_i . If $h_i : Z_i \rightarrow X_i$ is a flipping contraction, we denote the exceptional curve of $h_i : Z_i \rightarrow X_i$ by C_i and the exceptional curve of $g_{i+1} : Z_{i+1} \rightarrow X_i$ by C_i^+ . We also denote the strict transforms of S_1 , B_1 and E_1 on Z_i by S_i , B_i and E_i . Then we have the following important numerical properties:

Numerical properties.

- (1) For some positive rational numbers p and q , $(pS_i + qE_i).C_i < 0$;
- (2) For some positive rational numbers r and s , $(S_i + rB_i + sE_i).C_i = 0$;
- (3) For fixed i , we have $S_{i+1}.C_i^+ > 0$ or $S_{i+1}.C_{i+1} > 0$.

For the proof, see [FA, 21.6.2 Lemma, 21.7.1 Proposition and 21.7.2 Lemma].

If we encounter divisorial contraction $h_n : Z_n \rightarrow X_n$, then $X_n \rightarrow Y$ is the flip of f ([FA, 6.5.5]). Hence if all flips in BTM exist and terminate, then the flip of f exists. So we need to examine such flips in detail.

Roughly speaking there are three types of such flips as follow:

(a) beginning flip.

Let

$$j := \min \{k | h_k \text{ is flipping and } E_k.C_k \leq 0, \text{ or } h_k \text{ is divisorial.}\}.$$

We call g_i with $1 \leq i \leq j - 1$ a beginning flipping contraction.

By the numerical property (1), we have $E_i.C_i > 0$ and $S_i.C_i < 0$. Hence this is a flipping contraction of type 1 and the flip exists by (2.0) and terminates (for termination, see [FA, §7]).

After beginning flips, we encounter h_j . If it is a divisorial contraction, then we are done. Assume that h_j is a flipping contraction.

(b) middle flip.

C_j satisfies $E_j.C_j \leq 0$ by definition. If C_j satisfies also $S_j.C_j \leq 0$, we call h_j a middle flipping contraction.

We treat this flip by dividing into the following cases:

(b1) $E_j.C_j < 0$ and $S_j.C_j < 0$.

If $C_j \subsetneq S_j \cap E_j$, there is no general strategy to prove the existence of the flip of g_j . In our application, we can choose g so as to avoid this situation.

If $C_j = S_j \cap E_j$, it is a flip of type 2 and the flip of h_j exists by (2.1) and h_{j+1} is a divisorial contraction (see [FA, 21.8.2.1]). Hence in this case we are done.

(b2) $E_j.C_j = 0$ and $S_j.C_j < 0$.

There is no general strategy to prove the existence of the flip of g_j . In our application, we can choose g so as to avoid this situation.

(b3) $E_j.C_j < 0$ and $S_j.C_j = 0$.

If some component of C_j is contained in S_j , there is no general strategy to prove the existence of the flip of g_j . In our application, we can choose g so as to avoid this situation.

The left case is that any component of C_j does not intersect S_j . The heart of the Shokurov's paper is to prove the existence of the flip for such a flipping contraction by induction (see §4 and §7).

Assume that h_j satisfies (b3) and any component of C_j does not intersect S_j . Assume furthermore that the flip of h_j exists and h_{j+1} is also a flipping contraction. By the numerical property (3), we have $S_{j+1}.C_{j+1} > 0$.

(c) final flip. We can prove that if h_k is a flipping contraction such that $S_k.C_k > 0$ for some k , every C_l 's for $l \geq k$ satisfies $S_l.C_l > 0$ (see [FA, 21.8.3.1 Lemma]). We call such a h_l a final flipping contraction. By the numerical property (1), we have $E_k.C_k < 0$. Hence h_k is of type 1. So the flip exists by (2.0) and terminate (for termination, see [FA, §7]). After final flips, we obtain a divisorial contraction. By contracting the divisor we obtain the flip of f .

We remark that if $S_j \cap E_j$ is irreducible, then h_j satisfies the assumption of (b1) and that $S_j \cap E_j = C_j$ or the assumption of (b2) and that any component of C_j does not intersect S_j . If the latter case occurs, once we clear the flip of C_j final flipping contractions or a divisorial contraction appear and we are done.

3. SET UP AND DIVIDING INTO CASES

Set up (3.0). Let X be an algebraic \mathbb{Q} -factorial normal 3-fold, $f : X \rightarrow Y$ is a projective small contraction onto a normal 3-fold Y such that $\rho(X/Y) = 1$ and C the exceptional locus. Let S be an irreducible divisor on X and B a boundary on X such that $\lfloor B \rfloor = 0$. We assume all the following assumptions in the below sections:

- (1) $K_X + S$ is PLT. $K_X + S + B$ is LC and LT outside S . But $K_X + S + B$ is not LT;
- (2) $(K_X + S).C < 0$, $S.C < 0$ and $(K_X + S + B).C = 0$;
- (3) $2(K_X + S + B) \sim 0$ near C ;
- (4) there are at least 2 divisors in $\text{PLC}(K_S + \text{Diff}_S(B))$;
- (5) there is a unique component L of $\lfloor \text{Diff}_S(B) \rfloor$ such that $L \not\subset C$. For any component D of $\text{Supp } B|_S$, if $D \cap \text{LLC}(K_S + \text{Diff}_S(B)) \neq \emptyset$ then $D = L$ or $D \subset C$.

As we reviewed in the introduction, we have only to prove the existence of the flip under the assumptions

- (1)' $K_X + S$ is PLT but $K_X + S + B$ is not LT

and (2) and (3). But by [S, Proposition 6.12] or [FA, Proposition 21.9], we may assume (1), by [S, Corollary 7.3] or [FA, 21.12.1], we may assume (4) and by [S, Reduction 8.2] or [FA, 22.1], we may assume (5).

Main Theorem. *Under the assumptions in (3.0), the flip of f exists.*

(3.1) Reduction to the case that C is irreducible. Here we consider f as in (3.0) but only assume (1)~(3). Replace X by an analytic neighborhood of C and take a \mathbb{Q} -factorial LT model $\tilde{X} \rightarrow X$ for $K_X + S$ near S which exists by (1.13). Let \tilde{S} be the strict transform of S . Run the $K_{\tilde{X}} + \tilde{S}$ -LMMP. Then the intermediate flips satisfy (1)~(3) in (3.0). Hence by taking algebraization, it suffices to prove the main theorem under the additional assumption that C is irreducible (note that we can assume also (4) and (5).)

Dividing into cases (3.2). If $C \cap L$ consists of more than 1 points, by (3.1), we may assume L is not irreducible. But in this case the flip exists by (5) of (3.0). Hence we may assume that $C \cap L$ is one point and we denote it by P .

Note that if $C \not\subset \text{Supp } B$, $K_X + S + B$ is PLT at the generic point of C since $K_X + S$ is PLT. Hence the following 4 cases occur.

(a). C is irreducible. $C \not\subset \text{Supp } B$. $K_X + S + B$ is PLT at the generic point of C .

By applying (1.6) for $f|_S$ and the assumption (5) in (3.0), $\text{CLC}_0(K_S + \text{Diff}_S(B)) = \{C \cap L\}$.

(b). C is irreducible. $C \subset \text{Supp } B$. $K_X + S + B$ is not PLT at the generic point of C .

In this case we furthermore divide into 2 cases.

(b1). There is a point $Q \in C$ and $Q \notin L$ such that $Q \in \text{CLC}_0(K_S + \text{Diff}_S(B))$.

We see that such a Q is unique by considering the degree of the boundary of $K_X + S + B|_S|_C$.

(b2). The contrary case to (b1).

(c). $C \subset \text{Supp } B$. $K_X + S + B$ is PLT at the generic point of C . By applying (1.6) for $f|_S$ and the assumption (5) in (3.0), $\text{CLC}_0(K_S + \text{Diff}_S(B)) = \{C \cap L\}$.

In this case we do not assume that C is irreducible for induction. But we may assume that all the components intersect L . In fact a component which does not intersect L is analytically of type 3 (see (2.2)). Hence by (1.4), the number of irreducible components is at most 2.

Definition (3.3) (Good extraction). Consider the situation as in (3.0). Let $g : Z \rightarrow X$ be a log crepant divisorial (primitive) extraction for $K_X + S + B$ and E the exceptional divisor. We say that g is a good extraction (resp. semi-good extraction) if and only if the following hold:

- (1) $D := g^{-1}S \cap E \simeq \mathbb{P}^1$ (resp. $g^{-1}S \cap E$ is a chain of $D_1 \simeq \mathbb{P}^1$ and a smooth curve D_2);
- (2) $K_Z + g^{-1}S + E$ is LT (resp. $K_Z + g^{-1}S + E$ is LT outside P);
- (3) $K_Z + g^{-1}S + E + B|_D = K_D + P + \frac{1}{2}P_1 + \frac{1}{2}P_2$, where P, P_1 and P_2 are distinct points. (resp. $K_Z + g^{-1}S + E + B|_{D_1} = K_{D_1} + P + \frac{1}{2}P_1 + \frac{1}{2}P_2$, where $P := D_1 \cap D_2, P_1$ and P_2 are distinct points).

Corollary for the Definition (3.4). Consider the situation as in Definition (3.3). Assume that $g(E)$ is a point. Then if g is good (resp. g is semi-good),

$$\text{LLC}(K_Z + g^{-1}S + g^{-1}B + E|_E) = D \text{ or } D \cup M,$$

where M is an irreducible curve and intersects D only at P

$$\text{(resp. } \text{LLC}(K_Z + g^{-1}S + g^{-1}B + E|_E) = D_1 \cup D_2 \text{ or } D_1 \cup D_2 \cup M,$$

where M is an irreducible curve and intersects D_2 only at one point $\neq P$).

Furthermore on M there is at most one $\text{CLC}_0(K_Z + g^{-1}S + g^{-1}B + E|_E)$ except P (resp. except $D_2 \cap M$).

proof. We prove the claim only for the case g is good (the proof for another case is similar). By (1.7) and (3) of the definition of good extraction in (3.3), $\text{LLC}(K_Z + g^{-1}S + g^{-1}B + E|_E)$ is a union of curves. Assume that $\text{LLC}(K_Z + g^{-1}S + g^{-1}B + E|_E)$ has another irreducible component except D . Let M be one of it. Since $D = g^{-1}S|_E$ is ample and $\text{CLC}_0(K_Z + g^{-1}S + g^{-1}B + E|_E) \cap D = P$, M must intersect D only at P . By this, M must be unique. For the latter half, we have only to note $\deg K_Z + g^{-1}S + g^{-1}B + E|_{E|_M} = 0$. \square

4. TREATMENT OF CASE (a) AND (b1)

In this section we consider cases (a) and (b1). We use the notation as in (3.2).

Theorem (4.0). *Let $h_t : Z_t \rightarrow X$ be a \mathbb{Q} -factorial LT model for $K_X + S + B$ near S (which exists by (1.13)).*

Run the $(K_{Z_t} + S_t + B_t + E_t - \epsilon B_t)$ -MMP over X .

Then by interchanging the surgeries of MMP if necessarily, there is a nonempty sequence of divisorial extractions

$$Z_n \xrightarrow{g_n} Z_{n-1} \xrightarrow{g_{n-1}} \dots Z_1 \xrightarrow{g_1} X$$

satisfying the following properties:

- (1) *Let E_i be the exceptional divisor of g_i . Then $Q_{i-1} := g_i(E_i)$ is a point for $i \geq 2$ and $Q = g(E_1)$. Furthermore $Q_{i-1} \in E_{i-1}$ but $\notin \cup_{j \neq i-1} E_j$;*
- (2) *$K_{Z_i} + S_i + \sum_{j=1}^i E_j$ is LT, where S_i is the strict transform of S on Z_i ;*
- (3) *Let $D_i := E_i \cap E_{i-1}$ for $i \geq 2$ and $D_1 := E_1 \cap S_1$. Then for $1 \leq i \leq n-1$ (resp. $i = n$), $CLC_0(K_{Z_i} + S_i + B_i + E_i|_{E_i}) = \{Q_i, P_i\}$ (resp. $= \{P_i\}$), and $LLC(K_{Z_i} + S_i + B_i + E_i|_{E_i}) = D_i \cup L_i$, where $L_i := \text{Supp } B_i|_{E_i}$ and $P_i := L_i \cap D_i$. Furthermore L_i is irreducible and $Q_i \in L_i$. In particular g_1 is a good extraction.*

(See FIGURE (VII).)

Proof. As long as a component of E_t is left, $-\epsilon B_t$ is not nef by [FA, 2.19 Lemma] so the MMP continue till every component of E_t is contracted. After contracting every component of E_t , the morphism going back to X is at most a small contraction but since X is \mathbb{Q} -factorial, it must be isomorphism. Hence the last morphism is a divisorial contraction to X . Let it $g_1 : Z_1 \rightarrow X$ and E_1 the exceptional divisor.

Claim 1. *We can assume that $g_1(E_1) = Q$ by interchanging the surgeries of MMP if necessarily.*

Proof.

case (a).

If $g_1(E_1) \neq Q$, we must have $g_1(E_1) = L$. But we prove that it is impossible by the nonexceptional assumption. Assume that $g_1(E_1) = L$. Since $K_Z + S_1 + E_1$ is LT by the choice of MMP, $S_1 \cap E_1$ is irreducible. Hence $g'_1 := g_1|_{S_1} : S_1 \rightarrow S$ is isomorphism. By subadjunction,

$$(4.1.1) \quad K_{S_1} + \text{Diff}_{S_1}(E_1 + B_1) = g'^*_1(K_S + \text{Diff}_S(B)).$$

Let L' be the divisor corresponding to L by the isomorphism $S_1 \simeq S$. Then $L' = E_1 \cap S_1$. Hence $L' \not\subset \text{Supp } B_1$ by LC property of $K_Z + S_1 + B_1 + E_1$ along L' . On the other hand in case (a), $L = \text{Supp } B|_S$ so $\text{Supp } B_1 \cap S_1$ is at most a set of finite points. But Z_1 is also \mathbb{Q} -factorial, it must be empty. Hence $K_{Z_1} + S_1 + B_1 + E_1$ (and also $K_{Z_1} + S_1 + B_1 + E_1|_{S_1}$) is LT. Consequently by (4.1.1), $K_S + \text{Diff}_S(B)$ is also LT. But since $C \not\subset LLC(K_S + \text{Diff}_S(B))$ it must be PLT, a contradiction to the nonexceptional assumption.

Case (b1).

If $g_1(E_1) \neq Q$, we must have $g_1(E_1) = C, L$ or $C \cap L$.

We treat them separately.

Subcase 1. $g_1(E_1) = C$. We exclude this case as in Case (a).

Subcase 2. $g_1(E_1) = L$ or $C \cap L$.

If this case occurs, we prove the statement of Claim 1. Let

$$Z'_{i+1} \xrightarrow{g_{i+1}} Z'_i \cdots \dashrightarrow \dots Z_1 \xrightarrow{g_1} Z$$

be the sequence of the surgeries of MMP such that $Z'_i \dashrightarrow X$ consists of flips or divisorial extractions whose centers are the strict transform of L or $C \cap L$ and $Z'_{i+1} \rightarrow Z'_i$ is a divisorial extraction whose center is Q or the strict transform of C .

Note that $Z'_i \dashrightarrow X$ is isomorphic near Q . Then by the similar argument to Case (a), we can prove that the center of $Z'_{i+1} \rightarrow Z'_i$ is not C . Hence we can interchange $Z'_{i+1} \rightarrow Z'_i$ and $Z_1 \rightarrow X$.

□

We check (1), (2) and (3) of the statement of (4.0) for $g_1 : Z_1 \rightarrow X$ as in Claim 1. (1) is proved by Claim 1 and (2) is clear by the choice of MMP.

Let C' and L' be the strict transforms of C and L on S_1 respectively. Let $P_1 := L' \cap D_1$ in case (a) (resp. $P_1 := C' \cap D_1$ in case (b1)).

In case (a), $L = \text{Supp } B|_S$ and by LC property of $K_{Z_1} + S_1 + B_1 + E_1$ along D_1 , $D_1 \not\subset \text{Supp } B_1$. Hence $L' = \text{Supp } B_1|_{S_1}$. So $K_{Z_1} + S_1 + B_1 + E_1$ is LT at points of D_1 except P_1 , i.e., $P_1 = \text{CLC}_0(K_{Z_1} + S_1 + B_1 + E_1|_{E_1}) \cap D_1$.

In case (b1), near Q , $C = \text{Supp } B|_S$. Hence by the similar argument to case (a), $P_1 = \text{CLC}_0(K_{Z_1} + S_1 + B_1 + E_1|_{E_1}) \cap D_1$.

If there is no point in $\text{CLC}_0(K_{Z_1} + S_1 + B_1 + E_1|_{E_1})$ except P_1 , by setting $n = 1$, we finish the poof of (4.0).

Assume that there is a point Q_1 in $\text{CLC}_0(K_{Z_1} + S_1 + B_1 + E_1|_{E_1})$ other than P_1 . We first show the uniqueness of Q_1 . By (1.7) and the existence of P_1 , $\text{LLC}(K_{Z_1} + S_1 + B_1 + E_1|_{E_1})$ is connected. Hence $P_1, Q_1 \in L_1$, where $L_1 := \text{LLC}(K_{Z_1} + S_1 + B_1 + E_1|_{E_1}) - D_1$.

Claim 2. L_1 irreducible.

Proof. If L_1 is not irreducible, by $\rho(Z_1/X) = 1$ and $g_1(E_1) = Q$, D_1 is ample. So any component of L_1 intersects with D_1 . By just above, any component of L_1 passes through P_1 . But this is a contradiction since there is at least three reduced boundary of $K_{Z_1} + S_1 + B_1 + E_1|_{E_1}$ at P_1 . □

Consider the degree of $K_{Z_1} + S_1 + B_1 + E_1|_{E_1}|_{L_1}$. P_1 and Q_1 contribute degree 1 respectively to its boundary. By this, we know the uniqueness of Q_1 .

Next Claim 3 complete the proof of (3) for g_1 .

Claim 3. $L_1 = \text{Supp } B_1|_{E_1}$

Proof. Assume that there is another component L'_1 of $\text{Supp } B_1|_{E_1}$ except L_1 . Since D_1 is ample, $D_1 \cap L'_1 \neq \emptyset$. As above, L'_1 cannot pass through P_1 by LC property of $K_{Z_1} + S_1 + B_1 + E_1|_{E_1}$ at P_1 . This means that there is a component of $\text{Supp } B_1|_{S_1}$ except L' in case (a) (resp. C' in case (b1)). But this contradicts that $L = \text{Supp } B|_S$ in case (a) (resp. $C = \text{Supp } B|_S$ near Q in case (b1)). □

Next we prove the existence of the sequence by induction. Assume that we can construct the sequence satisfying the statement of (4.0) until i -th stage:

$$Z_i \rightarrow Z_{i-1} \rightarrow \cdots \rightarrow X.$$

We decompose the surgeries before Z_i as follow:

$$Z_{j+1} \xrightarrow{g_{j+1}} Z_j \dashrightarrow \dots Z_{i+1} \rightarrow Z_i,$$

where $Z_j \dashrightarrow Z_i$ consists of flips or divisorial contractions which are isomorphic near Q_i and $Z_{j+1} \dashrightarrow Z_j$ is not isomorphic near Q_i .

Claim 4. $Z_{j+1} \dashrightarrow Z_j$ is a divisorial contraction whose center is Q_i .

Proof. Assume the contrary. Then $Z_{j+1} \dashrightarrow Z_j$ is a divisorial contraction of (the strict transform of) L_i or flip.

First assume that $Z_{j+1} \dashrightarrow Z_j$ is a divisorial contraction of (the strict transform of) L_i . Here we denote the strict transform of E_i on Z_j by E_i and the strict transform of E_i on Z_{j+1} by E'_i . The argument is almost the same as one in Claim 1. Since $K_Z + E'_i + E_{j+1}$ is LT by the choice of MMP, $E'_i \cap E_{j+1}$ is irreducible. Hence $g'_{j+1} := g_{j+1}|_{E'_i} : E'_i \rightarrow E_i$ is isomorphism. By subadjunction,

$$(4.1.2) \quad K_{E'_i} + \text{Diff}_{E'_i}(S_{j+1} + B_{j+1} + E_{j+1} + E') = g'^*_1(K_{E_i} + \text{Diff}_{E_i}(S_j + B_j + E)),$$

where E is the reduced sum of exceptional divisors of $Z_j \dashrightarrow X$ except E_i and E' is the strict transform of E . Let L'_i be the divisor corresponding to L_i by the isomorphism $E'_i \simeq E_i$. Then $L'_i = E'_i \cap E_{j+1}$. Hence $L'_i \not\subset \text{Supp } B_{j+1}$. On the other hand, $L_i = \text{Supp } B_j|_{E_i}$ near Q_i by the assumption of induction (3). Hence $\text{Supp } B_{j+1} \cap E'_i$ must be empty near Q_i . So Q_i is not contained in $\text{CLC}_0(K_{E'_i} + \text{Diff}_{E'_i}(S_{j+1} + B_{j+1} + E_{j+1} + E'))$, a contradiction to the assumption of the induction (3).

Next assume that g_{j+1} is a flip. By assumption, there is a connected component of the flipped curve through Q_i and we call it l . Since $Z_j \dashrightarrow Z_i$ is isomorphic near Q_i , (the strict transform of) l on Z_i passes through Q_i . On Z_i , l is a exceptional curve of $Z_i \rightarrow X$ so on Z_j , $l \subset E_i$ and l is not contained in another exceptional divisor of $h_j : Z_j \rightarrow X$ or S_j . On the other hand we have $h_j^*B.l = 0$ and $B_j.l < 0$. Hence there exists a h_j -exceptional divisor E_k such that $E_k.l > 0$. We also have $h_j^*S.l = 0$ so there is a divisor F which is S_j or h_j -exceptional such that $F.l < 0$ Hence F must be E_i and in particular $E_i.l < 0$. Then we see that $l = L_i$ and $l = \text{Supp } B_j|_{E_i}$ near l by the following argument:

$l = L_i$ follows from the fact that near Q_i , $L_i = \text{Supp } B_j|_{E_i}$ and $l \subset \text{Supp } B_j$. For the latter half, we have only to consider the degree of $K_{E_i} + \text{Diff}_{E_i}(S_j + B_j + E)|_l$.

Here we consider the situation before the flip. We use the notation as in the FIGURE (VI). By the above numerical consideration, $E_i^-.l^- > 0$ and $E_k^-.l^- < 0$. Then any component of l^- is contained in E_i^- . For if a component l_1^- of l^- is contained in E_i^- , $l_1^- = E_i^- \cap E_k^-$ since $K_{Z_{j+1}} + E_i^- + E_k^-$ is LT and hence $E_i^- \cap E_k^-$ is irreducible. Since E_k^- is normal and l_1^- is contractible in E_k^- , we have $E_i^-.l^- = (l_1^-)_{E_k^-}^2 < 0$, a contradiction. By this, the natural map $E_i \dashrightarrow E_i^-$ is a morphism. Let this morphism be $c : E_i \rightarrow E_i^-$. Here we consider the following subadjunction:

$$(4.1.3) \quad K_{E_i} + \text{Diff}_{E_i}(S_j + B_j + E) = c^*(K_{E_i^-} + \text{Diff}_{E_i^-}(S_{j+1} + B_{j+1} + E^-)),$$

where E^- is the strict transform of E . Note that $l = L_i$ is a log crepant curve for this. Since $l = \text{Supp } B_j|_{E_i}$ near l , $\text{Supp } B_{j+1}|_{E_i^-}$ is at most a point $c(l)$ near $c(l)$. But by the \mathbb{Q} -factoriality of Z_{j+1} , $\text{Supp } B_{j+1}|_{E_i^-}$ is in fact empty near $c(l)$. Hence near $c(l)$,

$$K_{E_i^-} + \text{Diff}_{E_i^-}(S_{j+1} + B_{j+1} + E^-)$$

is LT. On the other hand there is a log crepant curve for this log divisor, so there must be F^- which is S_{j+1} or E_l^- for some $l \neq k$ such that $c(l)$ is contained in it and $c(l)$ is a smooth point of E_i^- . Hence l intersects F and E_k outside Q_i . But this gives a contradiction by considering the degree of the boundary of $K_{E_i} + \text{Diff}_{E_i}(S_j + B_j + E)|_l$. \square

By replacing $Z_{i+1} \dashrightarrow Z_i$ by $Z_{i+1} \dashrightarrow Z_i$, we can assume that $g_i : Z_{i+1} \rightarrow Z_i$ is a divisorial extraction whose center is Q_i . For this g_i , (2) is clear by the choice of MMP and the check of (3) is similar to the check for g_1 . \square

Definition (4.1) (Good sequence, Invariant λ). We call a good sequence a sequence of divisorial extractions as in (4.0). We define

$\lambda :=$

$\min\{n | n \text{ is the length of a good sequence for a } \mathbb{Q}\text{-factorial LT model for } K_X + S + B\}.$

We remark that it is a well defined finite number by (4.0).

(4.2) Conclusion of case (a, λ) and (b1, λ). *Fix a good sequence whose length is λ . We apply BTM starting by g_1 . So we will use the notation as in BTM. We used the same notation in (2.5) and (4.0) for different objects but in this conclusion (4.2), the notation in (4.0) is not used except $g_1 : Z_1 \rightarrow X$ and the notation by which we denote objects on Z_1 . So no confusion is caused because the same notation in (2.5) and (4.0) stands for the same object on Z_1 .*

Our conclusion is explained in the following flow chart:

Case (a)

Choose a good sequence with length n
and run BTM starting by g_1

↓

R_1 is a beginning flipping ray. So the flip of R_1 exists

R_2 is a divisorial ray. DONE	R_2 is a flipping ray of type 2. DONE	R_2 is a flipping ray and $S_2 \cap C_2 = \emptyset$.	R_2 is a final flipping ray. DONE
------------------------------------	--	--	--

$P \notin C_1.$	$P \in C_1.$
-----------------	--------------

$C_2 \cap \text{CLC}_0 \neq \emptyset$. Then the connected component C'_2 of C_2 containing CLC_0 is of type (a, $\lambda' < \lambda$). (Another component is of type 3 or 4. DONE)	$C_2 \cap \text{CLC}_0 = \emptyset$. Then C_2 is of type 3 or 4. DONE	A component of C_1^+ is contained in LLC. Then C_1^+ is irreducible.	No component of C_1^+ is contained in LLC. Then C_2 is of type 3. DONE		
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 5px; vertical-align: top;"> $C_2 \cap \text{CLC}_0 = \emptyset$. Then C_2 is of type 3 or 4. DONE </td> <td style="width: 50%; padding: 5px; vertical-align: top;"> $C_2 \cap \text{CLC}_0 \neq \emptyset$ </td> </tr> </table>		$C_2 \cap \text{CLC}_0 = \emptyset$. Then C_2 is of type 3 or 4. DONE	$C_2 \cap \text{CLC}_0 \neq \emptyset$		
$C_2 \cap \text{CLC}_0 = \emptyset$. Then C_2 is of type 3 or 4. DONE	$C_2 \cap \text{CLC}_0 \neq \emptyset$				

↓

Reduction to (a, $\lambda' < \lambda$)

L'_1 exists. Then the connected component C'_2 of C_2 containing e is of type (b).	L'_1 does not exist. Then the connected component C'_2 of C_2 containing e is of type (c)
--	---

↓

↓

Reduction to

Reduction to

$$\text{CLC}_0 := \text{CLC}_0 (K_{Z_2} + S_2 + B_2 + E_2|_{E_2})$$

$$\text{LLC} := \text{LLC} (K_{Z_2} + S_2 + B_2 + E_2|_{E_2})$$



$$P := L' \cap S_1 \cap E_1$$

Case (b1)

Choose a good sequence with length n
and run BTM starting by g_1






R_1 is a beginning flipping ray. So the flip of R_1 exists



R_3 is a flipping ray of type 2. DONE

A component of C_1^+ is contained in LLC.

No component of C_1^+ is contained in LLC. Then C_2 is of type 3. DONE



Reduction to case (b1, $\lambda' < \lambda$)	Reduction to case (b2)
--	---------------------------

$$\text{CLC}_0 := \text{CLC}_0 (K_{Z_2} + S_2 + B_2 + E_2|_{E_2})$$

e is the unique element in CLC_0 outside S_2

$$\text{LLC} := \text{LLC} (K_{Z_2} + S_2 + B_2 + E_2|_{E_2})$$

Case (a).

Since $g(E)$ is a point, $\text{Supp } R_1$ is the strict transform of C . Clearly we have $E.R_1 > 0$. Hence by $f^*S.R_1 < 0$, $S_1.R_1 < 0$. So R_1 is a beginning flipping ray and hence the flip exists.

Claim 1.

- (1) $S_2 \cap E_2$ is irreducible;
- (2) $K_{Z_2} + E_2$ is PLT.

Proof.

- (1) First assume that $P \notin C_1$. If $S_2 \cap E_2$ is reducible, a component $\overline{C_1^+}$ of C_1^+ is contained in $S_2 \cap E_2$. Then $\overline{C_1^+} \in \text{LLC}(K_{Z_2} + S_2 + B_2 + E_2)$, which in turn shows that C_1 contains an element of $\text{CLC}(K_{Z_1} + S_1 + B_1 + E_1)$ since $K_{Z_1} + S_1 + B_1 + E_1$ is numerically trivial over Y . But this contradicts the assumption of Subcase 1.

Next assume that $P \in C_1$. If $S_2 \cap E_2$ is reducible, a component $\overline{C_1^+}$ of C_1^+ is contained in $S_2 \cap E_2$. On the other hand by $P \in C_1$, $B_1.C_1 > 0$ and hence $B_2.C_1^+ < 0$. So $\overline{C_1^+} \subset S_2 \cap E_2 \cap \text{Supp } B_2$. But this contradicts LC property of $K_{Z_2} + S_2 + B_2 + E_2$ along $\overline{C_1^+}$.

- (2) By the proof of (1), E_2 is normal and $\text{LLC}(K_{Z_2} + S_2 + B_2 + E_2|_{E_2}) \subset \text{Supp } B_2|_{E_2} \cup \text{Supp } S_2|_{E_2}$. Hence $K_{Z_2} + E_2|_{E_2}$ is PLT, which in turn shows that $K_{Z_2} + E_2$ is PLT by the inversion of adjunction.

□

Consider the next extremal ray R_2 . R_2 may be a divisorial ray but in this case we can obtain the flip of C by contracting E_2 . Assume that R_2 is a flipping ray. Since $S_2 \cap E_2$ is irreducible and all exceptional curves over Y are contained in E_2 , R_2 is a middle flipping ray or a final flipping ray. we may consider only the case that $E_2.R_2 < 0$ and $\text{Supp } R_2 \cap S_2 = \phi$ (see the last remark in (2.5)). Note that by the numerical property (2) in (2.5), $B_2.R_2 < 0$. Hence R_2 is a special flipping ray by Claim 1 (2). Here we divide into two cases.

Subcase 1. $P \notin C_1$. See FIGURE (VIII).

Claim 2. C_1^+ is irreducible.

Proof. Easy. □

Note that $\text{LLC}(K_{Z_1} + S_1 + B_1 + E_1|_{E_1}) = S_1|_{E_1}$ or $S_1|_{E_1} \cup L_1$. By the proof of Claim 1, $\text{LLC}(K_{Z_2} + S_2 + B_2 + E_2|_{E_2}) = S_2|_{E_2}$ or $S_2|_{E_2} \cup L'_1$, where L'_1 is the strict transform of L_1 . In particular $C_2 \notin \text{LLC}(K_{Z_2} + S_2 + B_2 + E_2|_{E_2})$.

If $C_2 \cap \text{CLC}_0(K_{Z_2} + S_2 + B_2 + E_2|_{E_2}) = \phi$, $K_{Z_2} + E_2 + B_2|_{E_2}$ is PLT or exceptional near C_2 . If the former case occurs, $K_{Z_2} + E_2 + B_2$ is PLT near C_2 by the inversion of adjunction and hence C_2 is of type 3 and the flip exists by (2.2). If the latter case occurs, C_2 is of type 4 and the flip exists by (2.3).

Hence we assume below that $C_2 \cap \text{CLC}_0(K_{Z_2} + S_2 + E_2 + B_2) \neq \phi$. By the proof of Claim 1, we know that there is no $\text{CLC}_0(K_{Z_2} + S_2 + E_2 + B_2)$ on C_1^+ . Hence $\text{CLC}_0(K_{Z_2} + S_2 + E_2 + B_2) = \{Q_1, P\}$. By the assumption that $C_2 \cap S_2 = \phi$, $Q_1 \in C_2$. Let C'_2 be the connected component of C_2 containing Q_1 . We will show that the flip of C'_2 is of type $(a, \lambda' < \lambda)$. See FIGURE (IX). (Another component is of type 3 or 4. So we have only to consider C'_2 .) First we prove that C'_2 is irreducible. Denote the strict transform of L_1 on E_2 by L'_1 . Note that $L'_1.C'_2 > 0$ and $L'_1.C_1^+ = 0$. Since $\rho(Z_1/Y) = 2$ and $L'_1 = \text{Supp } B_2|_{E_2}$, only components of C_1^+ are numerically trivial for L'_1 among curves in E_2 . By similar reason, we see that $(S_2|_{E_2})^2 > 0$. Note also

that every component of C'_2 intersects C_1^+ because C_1^+ is irreducible and before the flip of C_1 , every component of C'_2 intersect S_1 . So after contracting a component of C'_2 , L'_1 becomes ample. By these properties, we can apply (1.9) after contracting one component of C'_2 by setting $B_1 = S_2 \cap E_2$, $B_2 = L'_1$, $C_1 = C'_2$ and $C_2 = \phi$ (every left side is the notation in (1.9)). Then $C'_2 \cap L'_1 \notin \text{CLC}_0(K_{Z_2} + E_2 + B_2)$. But by the assumption that $Q_1 \in C'_2$, C'_2 must be empty after the contraction of a component of it, i.e., C'_2 is irreducible.

We can easily check that (Z_2, E_2, B_2, C_2) satisfies the assumption of case (a) (in this case L'_1 corresponds to L). Note that the flip $Z_1 \dashrightarrow Z_2$ is isomorphism near Q_1 . Hence the above good sequence except g_1 is also a good sequence with length $\lambda - 1$. So we finish the reduction to the case $(a, \lambda' < \lambda)$.

Subcase 2. $P \in C_1$. See FIGURE (X).

Note that $S_2.C_2 = 0$ and $S_2.C_1^+ > 0$. By $\rho(Z_2/Y_1) = 2$, only components of C_2 is numerically trivial for S_2 among curves in E_2 . Let $a : E_2 \rightarrow E_1$ be the contraction which is the restriction of $Z_2 \dashrightarrow Z_1$. Then C_1^+ is the exceptional curve of a .

Assume that any component of C_1^+ is not contained in LLC $(K_{Z_2} + S_2 + E_2 + B_2|_{E_2})$. Since $C_1^+ \cap S_2 \ni P$, $C_1^+ \cap \text{LLC}(K_{Z_2} + S_2 + E_2 + B_2|_{E_2}) = \{P\}$ by (1.6). If L_1 exists, its strict transform L'_1 must pass through P . Hence $K_{Z_2} + S_2 + E_2 + B_2|_{E_2}$ has two reduced boundaries at P . But $C_1^+ \subset \text{Supp } B_2|_{E_2}$, a contradiction to LC property of $K_{Z_2} + S_2 + E_2 + B_2|_{E_2}$. So L_1 does not exist and $K_{Z_2} + S_2 + E_2 + B_2|_{E_2}$ is PLT near C_2 . Hence $K_{Z_2} + E_2 + B_2$ is PLT near C_2 by the inversion of adjunction and C_2 is of type 3. So the flip of C_2 exists by (2.2).

Assume that a component of C_1^+ is contained in LLC $(K_{Z_2} + S_2 + E_2 + B_2|_{E_2})$. By $S_2.C_1^+ > 0$, all the components of C_1^+ pass through P . Furthermore by $C_1^+ \subset \text{Supp } B_2$, C_1^+ must be irreducible and furthermore $\text{Supp } B_2|_{E_2} - C_1^+$ does not intersect S_2 . Then $\text{LLC}(K_{Z_2} + S_2 + E_2 + B_2|_{E_2}) = S_2|_{E_2} \cup C_1^+$ or $S_2|_{E_2} \cup C_1^+ \cup L'_1$, where L'_1 is the strict transform of L_1 (if it exists). So if $C_2 \cap \text{CLC}_0(K_{Z_2} + S_2 + B_2 + E_2|_{E_2}) = \phi$, C_2 is of type 3 or 4 as in Subcase 1. Assume below that $C_2 \cap \text{CLC}_0(K_{Z_2} + S_2 + B_2 + E_2|_{E_2}) \neq \phi$. Note that outside $S_2|_{E_2}$, elements of $\text{CLC}_0(K_{Z_2} + S_2 + B_2 + E_2|_{E_2})$ are only on C_1^+ and in fact only one point exists on C_1^+ outside $S_2|_{E_2}$. We denote this point by e . Note that if L_1 exists, $L_1 \subset C_2$ and $e \in L_1$. Hence in any case $e \in C_2$. Let C'_2 be the connected component of C_2 containing e . We will show that $\text{Supp } B_2|_{E_2} - C_1^+ = C'_2$.

By $\text{Supp } B_2|_{E_2} - C_1^+ \cap S_2 = \phi$, $\text{Supp } B_2|_{E_2} - C_1^+ \subset C_2$. After contracting only $\text{Supp } B_2|_{E_2} - C_1^+$, C_1^+ becomes ample by (1.8) and ampleness of $\text{Supp } B_1|_{E_1}$ on E_1 . Hence we can apply (1.9) by setting $B_1 = S_2 \cap E_2$, $B_2 = C_1^+$, $C_1 = C'_2$ and $C_2 = \phi$ after the contraction of $\text{Supp } B_2|_{E_2} - C_1^+$. We obtain that C'_2 cannot pass through $\text{CLC}_0(K_{Z_2} + E_2 + B_2)$ after the contraction of $\text{Supp } B_2|_{E_2} - C_1^+$ but it means C'_2 becomes empty after the contraction of $\text{Supp } B_2|_{E_2} - C_1^+$, i.e., $C'_2 = \text{Supp } B_2|_{E_2} - C_1^+$.

If L'_1 exists, then $L'_1 = \text{Supp } B_2|_{E_2} - C_1^+$ and C'_2 is of type (b). (In this case C_1^+ corresponds to L in (3.2). Furthermore if Q_1 exists, C'_2 is of type (b, $\lambda' < \lambda$) or if Q_1 does not exist, C'_2 is of type (b2).)

If L'_1 does not exist, C'_2 is of type (c). (In this case C_1^+ corresponds to L in (3.2).) See FIGURE (XI).

Case (b1). See FIGURE (XII).

We know as in case (a) that R_1 is a beginning flipping ray such that $\text{Supp } R_1$ is the strict transform of C on Z_1 . We examine the next extremal ray R_2 . It may be a divisorial ray but in this case we obtain the flip of C by contracting E_2 . We may assume that R_2 is a flipping ray. We treat by dividing into two cases.

Subcase 1. $S_2 \cap E_2$ is reducible. See FIGURE (XIII).

Let D be the strict transform of $S_1 \cap E_1$ on S_2 . First we prove that $S_2 \cap E_2 = D \cup l$, where l is an irreducible component of C_1^+ .

Let $S_2 \cap E_2 = D \cup l_1 \cup l_2 \cdots \cup l_n$, where l_i is an irreducible component of C_1^+ and they form a chain with this order. By LC property of $K_{Z_2} + S_2 + B_2 + E_2$, $l_i \not\subset \text{Supp } B_2$. On the other hand l_i 's form a chain and only l_n intersect $\text{Supp } B_2$. Hence n must be 1. We newly denote this l_1 by l . Note that $B_2.D = 0$. On the other hand $B_2.C_1^+ > 0$. Hence $B_2.C_2 \leq 0$. By the numerical properties (1) and (2) in (2.5), $E_2.C_2 < 0$ and $S_2.C_2 > 0$ or $E_2.C_2 > 0$ and $S_2.C_2 < 0$. In the former case R_2 is a final flipping ray so we are done. In the latter case $C_2 \subset S_2$ but contractible curves in S_2 are only D and l and $B_2.D = 0$ and $B_2.l > 0$. So in the latter case $B_2.C_2 = 0$. In this case we immediately see that $D = C_2$ by $B_2.C_2 = 0$, $B_2.C_1^+ > 0$ and $B_2.D = 0$. Furthermore we can easily check that D satisfies the assumption of (2.4). We will apply (1.10) by setting $a_1 = h_2|_{E_2}$, $a_2 = g_3|_{E_3} \circ \mu$ and $K_S + B = K_{X_2} + g(S_2) + g(B_2) + g(E_2)|_{g(E_2)}$, where μ is the normalization of E_3 . For this we need to check the assumptions of (1.10). The former half of the assumptions can be checked by (2.4) (1) and (2). Assumption (1) is clear. Assumption (2) follows from assumption (6) of (2.4). Assumption (3) follows from (2) of Theorem (2.4). Let l' be the strict transform of l on E_3 . Hence if $(l')^2 \geq 0$, then $l^2 \geq 0$. But l is a component of C_1^+ and hence $l^2 < 0$, a contradiction. So $(l')^2 < 0$. Since $l' = S_3 \cap E_3$, this shows that $S_3.l' < 0$. Note that $S_3.C_2^+ > 0$. So for the next extremal ray R_3 , we have $S_3.R_3 < 0$. Hence it is a flipping ray and its support is contained in S_3 . Note that $l' = S_3 \cap E_3$ is irreducible and it is the unique projective curve in S_3 . Hence R_3 is of type 3 and we are done (see (2.5)).

Subcase 2 $S_2 \cap E_2$ is irreducible.

By the irreducibility of $S_2 \cap E_2$, we may assume as in case (a) that $C_2 \cap S_2 = \phi$. By the numerical property (2) in (2.5), we have $B_2.C_2 > 0$.

Claim 3. $K_{Z_2} + E_2$ is PLT.

Proof. If $B_2.C_1^+ \leq 0$, then $C_1^+ \subset \text{Supp } B_2$ and hence we can prove the assertion as in Claim 1. If $B_2.C_1^+ > 0$, $B_2|_{E_2}$ is ample since $B_2.C_2 > 0$. We can prove similar statements to Claim 1 and Claim 2 in (5.2) and deduce from these that $K_{Z_2} + S_2 + E_2$ is LT. Hence we obtain the assertion. \square

(*) Note that some component of C_1^+ is contained in $\text{Supp } B_2$ by $L' \subset \text{Supp } B_2$,

where L' is the strict transform of L on E_2 . Furthermore by the g_2 -ampleness of S_2 , any component of C_1^+ passes through P .

Assume first that no component of C_1^+ is contained in $\text{LLC}(K_{Z_2} + S_2 + B_2 + E_2|_{E_2})$. By (1.6), $C_1^+ \cap \text{LLC}(K_{Z_2} + S_2 + B_2 + E_2) = P$. Hence if L_1 exists, its strict transform L'_1 cannot pass through P by * and (1.1). But L'_1 intersects C_1^+ , a contradiction to (1.6). Hence L_1 does not exist. Consequently $K_{Z_2} + S_2 + B_2 + E_2|_{E_2}$ is PLT near C_2 by (1.7) and C_2 is of type 3.

Assume next that some component of C_1^+ is contained in $\text{LLC}(K_{Z_2} + S_2 + B_2 + E_2|_{E_2})$. We know that such a component is unique. Denote it by $C_1'^+$. Note that $\text{LLC}(K_{Z_2} + S_2 + E_2 + B_2|_{E_2}) = S_2|_{E_2} \cup C_1'^+$ or $S_2|_{E_2} \cup C_1'^+ \cup L'_1$, where L'_1 is the strict transform of L_1 . So if $C_2 \cap \text{CLC}_0(K_{Z_2} + S_2 + B_2 + E_2|_{E_2}) = \phi$, C_2 is of type 3 or 4 as in case (a) Subcase 2. Assume below that $C_2 \cap \text{CLC}_0(K_{Z_2} + S_2 + B_2 + E_2|_{E_2}) \neq \phi$. Note that outside $S_2|_{E_2}$, elements of $\text{CLC}_0(K_{Z_2} + S_2 + B_2 + E_2|_{E_2})$ (except Q_1 if it exists) is only on $C_1'^+$ and in fact

only one point is on $C_1'^+$ outside $S_2|_{E_2}$. We denote this point by e (clearly $e \in \text{Supp } B_2|_{E_2}$). Note that if L_1 exists, $L_1 \subset C_2$ and $e \in L_1$. Hence in any case $e \in C_2$. Let C_2' be the connected component of C_2 containing e . We will show that $\text{Supp } B_2|_{E_2} - C_1'^+ = C_2'$. By $\text{Supp } B_2|_{E_2} - C_1'^+ \cap S_2 = \phi$, $\text{Supp } B_2|_{E_2} - C_1'^+ \subset C_2$. After contracting $\text{Supp } B_2|_{E_2} - C_1'^+$ and $C_1^+ - C_1'^+$, C_1^+ becomes ample by (1.8) and ampleness of $\text{Supp } B_1|_{E_1}$ on E_1 . Hence we can apply (1.9) by setting $B_1 = S_2 \cap E_2$, $B_2 = C_1'^+$, $C_1 = C_2'$ and $C_2 = \phi$ after the contraction of $\text{Supp } B_2|_{E_2} - C_1'^+$. We obtain that C_2' cannot pass through $\text{CLC}_0(K_{Z_2} + E_2 + B_2)$ after the contraction of $\text{Supp } B_2|_{E_2} - C_1'^+$ but it means C_2' becomes empty after the contraction of $\text{Supp } B_2|_{E_2} - C_1'^+$, i.e., $C_2' = \text{Supp } B_2|_{E_2} - C_1'^+$.

If L_1' exists, then $L_1' = \text{Supp } B_2|_{E_2} - C_1'^+$ and C_2' is of type (b). (In this case $C_1'^+$ corresponds to L in (3.2).) Furthermore if Q_1 exists, C_2' is of type (b1, $\lambda' < \lambda$). Hence the induction works well.

If L_1' does not exist, C_2' is of type (c). (In this case $C_1'^+$ corresponds to L in (3.2).) See FIGURE (XIV).

5. PRELIMINARIES FOR CASE (B2) AND (C)

Definition (5.0) (Invariant δ). Consider case (b2) or (c). We define

$\delta := \#\{E|E \text{ is a log crepant divisor for}$

$$K_X + S + B \text{ and } d(E, S) \leq 1\}.$$

We can prove $\delta < \infty$. See [FA, 4.12.1 Lemma].

The following lemmas is used for checking the goodness or semi-goodness of an extraction.

Lemma (5.1) (LT Lemma). *Let $(Z, K_Z + S + B + E)$ be a \mathbb{Q} -factorial LC 3-fold, where S and E is irreducible surfaces. Assume that $S \cap E$ is irreducible (resp. a union of two irreducible curves) and set $D_1 = S \cap E$ (resp. $D_1 \cup D_2 = S \cap E$).*

- (1) $D_1 \simeq \mathbb{P}^1$. S and E are generically normal crossing along D_1 (resp. $D_1 \simeq \mathbb{P}^1$. S and E are generically normal crossing along D_1 and D_2);
- (2) $\text{CLC}_0(K_Z + S + B + E|_E) \cap D_1$ consists of one point P and $P \in \text{Supp } B|_E$ (resp. $\text{CLC}_0(K_Z + S + B + E|_E) \cap D_1$ consists of one point P and $P = D_1 \cap D_2$);
- (3) S is normal.
- (4) $M := \text{LLC}(K_Z + g^{-1}S + g^{-1}B + E|_{E^\mu})$ is connected and we can write

$$M = \mu^{-1}D_1, B_1, \dots, B_m$$

(they form a chain with this order) (resp.

$$M = \mu^{-1}D_1, \mu^{-1}D_2, B_1, \dots, B_m$$

(they form a chain with this order)), where μ is the normalization of E and B_i is an irreducible curve contained in $\text{Supp } B|_E$.

Then $K_Z + S + E$ is LT (resp. $K_Z + S + E$ is LT outside P).

Proof. By assumption (4), E is normal. Also by assumption (4), $\text{CLC}_0(K_Z + S + E|_E)$ is contained in D_1 (resp. D_1 and D_2) and does not contain $D_1 \cap \text{Supp } B|_E$ (resp. $D_2 \cap \text{Supp } B$). Hence $K_Z + S + E|_E$ is LT (resp. $K_Z + S + E|_E$ is LT outside P), which in turn show that $K_Z + S + E$ is LT (resp. $K_Z + S + E$ is LT outside P) by the inversion of adjunction ([S, 5.13 proposition]). \square

Lemma (5.2) (Lemma for goodness). *Consider the situation of (b2) or (c) in (3.0). Let $g : Z \rightarrow X$ be a (primitive) log crepant divisorial extraction for $K_X + S + B$. Let E be the exceptional divisor and $\cup_{i=1}^n D_i := g^{-1}S \cap E$, where D_i 's form a chain and D_1 intersects with $g^{-1}C$. Then $n = 1$ or 2 .*

Furthermore we assume that the following hold:

If $n = 1$ (resp. $n = 2$),

- (1) $D_1 \simeq \mathbb{P}^1$. $g^{-1}S$ and E are generically normal crossing along D_1 (resp. $D_1 \simeq \mathbb{P}^1$. $g^{-1}S$ and E are generically normal crossing along D_1 and D_2);
- (2) $K_Z + g^{-1}S + g^{-1}B + E|_{D_1} = K_{D_1} + \frac{1}{2}P_1 + \frac{1}{2}P_2 + P$ (resp. we assume the same thing and also assume that $P = D_1 \cap D_2$);
- (3) $g^{-1}S$ is normal.

Then g is a good contraction (resp. a semi-good contraction).

Proof. Note that $g^{-1}B$ is g -ample. So if $n \geq 3$, D_2, \dots, D_{n-1} are contained in fibers but do not intersect $g^{-1}B$, a contradiction. Hence $n \leq 2$.

Assume that $n = 1$ (resp. $n = 2$) and above (1), (2) and (3) hold. We have only to check the (2) of the definition of good extraction (resp. semi-good extraction) in (3.3). For this, we use Lemma (5.1). We have to check assumption (4) of Lemma (5.1).

Let $\mu : E^\mu \rightarrow E$ be the normalization of E .

Claim 1. *LLC $(K_Z + g^{-1}S + g^{-1}B + E|_{E^\mu})$ is connected.*

Proof. In the case $g(E) \neq L$, E is a projective surface such that $K_Z + g^{-1}S + g^{-1}B + E|_E \equiv 0$. So by (1.7) and the assumption (2), we immediately obtain that LLC $(K_Z + g^{-1}S + g^{-1}B + E|_{E^\mu})$ is connected.

Next we treat the case $g(E) = L$. By the assumption (1), $n = 2$ in this case.

Assume that LLC $(K_Z + g^{-1}S + g^{-1}B + E|_{E^\mu})$ is not connected. Then there is a point over Q contained in LLC $(K_Z + g^{-1}S + g^{-1}B + E|_{E^\mu})$ and not in D_1 . In particular, the fiber is reducible. So we can contract bimeromorphically all the components of the fiber over Q which are not contained in LLC $(K_Z + g^{-1}S + g^{-1}B + E|_{E^\mu})$. But this contradicts (1.6). Hence we finish the proof of Claim 1. \square

Let $M := \text{LLC}(K_X + g^{-1}S + g^{-1}B + E|_{E^\mu})$. By Claim 1 and (2), this is a tree of curves and $\mu^{-1}D_1$ is an end of the tree.

Claim 2.

$$M = \begin{cases} \mu^{-1}D_1, B_1, \dots, B_m \text{ (with this order) in case } n = 1 \\ \mu^{-1}D_1, \mu^{-1}D_2, B_1, \dots, B_m \text{ (with this order) in case } n = 2 \text{ and } g(E) \neq L \\ \mu^{-1}D_1, \mu^{-1}D_2 \text{ (with this order) in case } n = 2 \text{ and } g(E) = L. \end{cases}$$

In the above, B_i is an irreducible curve contained in $\text{Supp } g^{-1}B|_{E^\mu}$.

Proof. Note that in case (b2), $g(E) = Q$ or C and in case (c), $g(E) = Q$ or L . First assume that $g(E) \neq L$. If $n = 1$ (resp. $n = 2$), $g^{-1}B|_{E^\mu}$ is ample (resp. ample in the cases $g(E) = Q$ or nef and only trivial for D_1 in the case $g(E) = C$). (For the case $g(E) = C$, see Theorem (4.0) Claim 2 in detail.) Hence $\text{Supp } g^{-1}B|_{E^\mu}$ is connected. (In the ample case, this is well known. In the case $g(E) = C$, D_1 coincides with the support of the next flipping ray (see Theorem (4.0) Claim 2). Hence after contracting $\mu^{-1}D_1$ in E^μ , $g^{-1}B|_{E^\mu}$ becomes ample. Hence we obtain the connectedness as usual.)

Let $\mu^{-1}D_1, B_1, \dots, B_n$ (resp. $\mu^{-1}D_1, C_1, \dots, C_m, \mu^{-1}D_2, B_1, \dots, B_n$) be the irreducible components of M (they form a tree with this order). We will show that B_1, \dots, B_n is exactly in

Supp $g^{-1}B|_{E^\mu}$

(resp. B_1, \dots, B_n is exactly in Supp $g^{-1}B|_{E^\mu}$ and $\{C_i\}$ is an empty set). Let Supp $g^{-1}B|_{E^\mu} = B' + B''$ be the decomposition such that B' is the union of components of Supp $g^{-1}B|_{E^\mu}$ contained in M . Let B'_k be any connected component of B' . By the connectedness of Supp $g^{-1}B|_{E^\mu}$, $B'_k \cap B'' \neq \emptyset$ so an end of B'_k is B_n since $\deg(K_X + g^{-1}S + g^{-1}B + E|_{E^\mu|_{B_j}}) = 0$ for $1 \leq j \leq n$ and for $j \neq n$ B_j intersects already two component of LLC $(K_Z + g^{-1}S + g^{-1}B + E|_{E^\mu})$. In particular we know that B' is actually connected and $B' \subset B_1 \cup \dots \cup B_n$. But by the fact $g^{-1}B|_{E^\mu}$ is ample (resp. ample in the cases $g(E) = Q$ or nef and only trivial for D_1 in the case $g(E) = C$), $\mu^{-1}D_1$ (resp. C_i and $\mu^{-1}D_2$) must intersect B' so we are done.

Next we treat the case $g(E) = L$. Let $\mu^{-1}D_1, C_1, \dots, C_m, \mu^{-1}D_2$ be the irreducible components of M , where C_i is a component of a fiber of g . No component of Supp $g^{-1}B|_{E^\mu}$ pass through P . In fact otherwise on $g^{-1}S$ (this is normal by the assumption (3)), $K_Z + g^{-1}S + g^{-1}B + E|_{g^{-1}S}$ has two reduced boundary D_1 and D_2 and another boundary, a contradiction. Hence $\{C_i\}$ must be empty since $g^{-1}B|_{E^\mu}$ is $g \circ \mu$ -ample. Now we finish the proof of Claim 2 \square

By these 2 claims, we can apply (5.1) and we are done. \square

Remark (5.3). We can prove the normality of E even if we replace (1) by the following weaker assumption in case $n = 2$:

(1)' $D_1 \simeq \mathbb{P}^1$. $g^{-1}S$ and E are generically normal crossing along D_1 .

In fact the assumption that $g^{-1}S$ and E are generically normal crossing along D_2 is used only in the last paragraph of the above proof.

6. TREATMENT OF CASE (B2, δ)

We use the notation as in (3.2).

Theorem (6.0). *There is a good extraction or semi-good extraction $g : Z \rightarrow X$ such that $g(E) = C$, where E is the exceptional divisor of g . Furthermore if g is good one of the following holds:*

- (1) *Supp B and S are not generically simply tangent along C and $d(E, S) \leq 1$ or*
- (2) *Supp B and S are generically simply tangent along C and $g^*(K_X + S + B)|_E$ is PLT over a general point of C .*

If g is semi-good, let $D_1 \cup D_2 := g^{-1}S \cap E$, where D_1 is the strict transform of C on $g^{-1}S$. Then D_1 coincides the support of a flipping ray satisfying the assumption of (2.4).

Proof.

Claim 1. *Except the case that Supp B and S are generically simply tangent along C , there is a log crepant exceptional divisor for $K_X + S + B$ with multiplicity in $S \leq 1$.*

Proof. It is sufficient to prove the statement after cutting X by a general hyperplane section at a general point of C . Let x be a general point of C and H a general hyperplane section of X through x . Let $\mu : H_t \rightarrow H$ be the MLT for $(H, K_X + S + B|_H, x)$ and F_i ($1 \leq i \leq n$) exceptional curves, where F_1 intersects $\mu^{-1}(S|_H)$. First assume that there are 2 reduced components of the boundary of $K_X + S + B|_H$. If x is smooth, then the simple blow up at x gives what we want. If x is singular, then MLT coincides with MRS, so by (1.3), we are done. Next assume that there is 1 reduced component of the boundary of $K_X + S + B|_H$. If x is singular, then MRS is dominated by MLT, so we are done by (1.3). If x is smooth and $n \geq 2$, $d(F_1, S|_H) = 1$. Hence we are done. If x is smooth and $n = 1$, then by the description of MLT, we encounter the following 3 possibilities: See FIGURE (I)

Only in the middle case, we have no log crepant curve as we want. Hence we obtain Claim 1. \square

We go back to 3-dimensional situation. Let $h_t : X^t \rightarrow X$ be a \mathbb{Q} -factorial LT model for $K_X + S + B$ near S and we denote the exceptional divisor corresponding to E_1 by E . Run $N := h_t^*(K_X + S + B) - \epsilon E$ -MMP and let $g' : Z' \rightarrow X$ be the end result. Then $N' := K_{Z'} + S' + B' + E' - \epsilon E - \delta g'^*S$ ($0 < \delta \ll 1$) is KLT and g' -nef and g' -big, its sufficient multiple is g' -free by Kawamata's base point free theorem. Let $h : Z' \rightarrow Z$ be the morphism defined by such a multiple and $g : Z \rightarrow X$ be the natural morphism. Then exceptional divisor of g is exactly E . Indeed, for a general curve in E , N is nef while MMP since $N \equiv -\epsilon E$ over X . So E is g -exceptional. On the other hand some h_t -exceptional divisor F ($\neq E$) are g -exceptional, a general curve in F , $h(N')$ is numerically nonpositive, a contradiction.

We will prove that this g satisfies the statement of Theorem. By (5.2), we can write $g^{-1}S \cap E = D_1$ or $D_1 \cup D_2$. By the choice of E , $g^{-1}S$ and E are generically normal crossing along D_1 . By the assumption of (b2), (2) of Lemma (5.2) holds. Note that $g^{-1}S$ is normal since S is normal and $g^{-1}S$ is normal along intersection curves with g -exceptional divisors. Hence by (5.2), g is good if $g^{-1}S \cap E$ is irreducible. In this case we check that (1) or (2) holds. If $\text{Supp } B$ and S are not generically simply tangent along C , (1) holds by Claim 1. If $\text{Supp } B$ and S are generically simply tangent along C , the restriction of g over a general hyperplane section of X at a general point of C coincides with MLT. Hence (2) of Theorem (6.0) holds.

Consider the case $g^{-1}S \cap E$ is reducible in the below. Run the BTM starting from g . We use the notation as in (2.5).

Claim 2. $g^{-1}B.R_1 = 0$, $E.R_1 > 0$ and $g^{-1}S.R_1 < 0$. R_1 is a flipping ray with $\text{Supp } R_1 = D_1$ which satisfies the assumption of (2.4).

Proof. It is easy to see that $g^{-1}B.D_1 = 0$. On the other hand, $g^{-1}B$ is positive for fibers of g . Hence $g^{-1}B.R_1 \leq 0$ since we play a 2-ray game. Considering g^*B and g^*S , we obtain that $E.R_1 > 0$ and $g^{-1}S.R_1 < 0$. By this $\text{Supp } R_1 \subset g^{-1}S$. Clearly $\text{Supp } R_1 \subset E$. By these, we see that $\text{Supp } R_1 = D_1$. The check that it satisfies the assumption of (2.4) is easy so we left it to readers.

By the property (2) of (2.4), (1) of Lemma (5.2) holds. Hence g is semi-good and we are done.

\square

(6.1) Conclusion of case (b2). We fix a divisorial extraction $g : Z \rightarrow X$ as in (6.0) and apply BTM starting by this g . We will use the notation as in (2.5). Our conclusion is explained in the following flow chart:

Case (b2)

Choose a primitive log crepant extraction g as in (6.0) and run BTM starting by g

<p>(A) g is semi-good. Then R_1 is of the type as in (2.4). \downarrow</p>	<p>(B) g is good.</p>
---	--

<p>R_2 is a flipping ray of type 2. DONE</p>	<p>R_1 is a flipping ray and $g^*(K_X + S + B) _{g^*H}$ is PLT</p>	<p>R_1 is a divisorial ray. DONE</p>	<p>R_1 is a flipping ray and $g^*(K_X + S + B) _{g^*H}$ is not PLT</p>
---	--	---	--

<p>$LLC = S_1 \cup E_1$. Then C_1 is of type 3. DONE</p>	<p>$LLC = S_1 \cup E_1 \cup l_1$</p>	<p>$C_1 \cap CLC_0 = \phi$. Then C_2 is of type 3 or 4. DONE</p>	<p>$C_2 \cap CLC_0 \neq \phi$. Then the connected component C'_1 of C_1 containing CLC_0 is of type (b2, $\delta' < \delta$). (Another component is of type 3 or 4. So we are done)</p>
	<p>$C_1 \cap CLC_0 = \phi$. Then C_1 is of type 3 or 4. DONE</p>	<p>$C_1 \cap CLC_0 \neq \phi$. Then the connected component C'_1 of C_1 containing CLC_0 is of type (c).</p>	

\Downarrow

\Downarrow

Reduction to case (c)

Reduction to case (b2, $\delta' < \delta$)

$$CLC_0 := CLC_0 (K_{Z_1} + S_1 + B_1 + E_1|_{E_1})$$

$$LLC := LLC (K_{Z_1} + S_1 + B_1 + E_1|_{E_1})$$

(A). First we assume that g is semi-good. See FIGURE (XVI). As in the proof of (6.0), R_1 is a flipping ray satisfying the assumption of (2.4) and $\text{Supp } R_1 = D_1$. After the flip D_1 , we have $S_2.C_1^+ > 0$ and $E_2.C_1^+ < 0$. Then we obtain the following:

Claim 1. $S_2.R_2 < 0$.

Proof. We derive a contradiction assuming that $S_2.R_2 \geq 0$. Let $\nu : E_2' \rightarrow E_2$ be the normalization. First we see that $(D_2^+)^2 > 0$ on E_2' as follow:

Since $D_2^+ = S_2 \cap E_2$, and S_2 and E_2 are generically normal crossing along D_2^+ , $(D_2^+)^2 = S_2.D_2^+$. On Z_2 , two extremal rays over Y are R_2 and the ray generated by D_1^+ . They satisfy $S_2.R_2 \geq 0$ and $S_2.D_1^+ > 0$. If $D_2^+ \subset \text{Supp } R_2$, D_2^+ is contractible in E_2 . But by $D_2^+ = S_2 \cap E_2$, we have $S_2.D_2^+ < 0$, a contradiction. Hence $D_2^+ \not\subset R_2$, which in turn show that $S_2.D_2^+ > 0$.

Next we deduce from $(D_2^+)^2 > 0$ on E_2' that $(D_2)^2 > 0$ on E_1 . (This give us a contradiction since D_2 is contained in a fiber of g .) Let Q be the point as in Theorem (2.4) (3). Note that Q is a smooth point. On the other hand on D_1 there is one non PLT point for $K_Z + g^{-1}S + g^{-1}B + E|_E$ by the assumption of (b2). By these and (1.10), we have $(D_2)^2 > 0$ on E_1 . As we saw above, this give a contradiction and prove Claim 3. \square

Hence it is a flipping ray and its support is contained in S_2 . Note that $S_2 \cap E_2$ is irreducible and it is the unique projective curve in S_2 . Hence R_2 is of type 2 and we are done (see (2.5)).

(B). Next we assume that g is good. We may assume that R_1 is a flipping ray. Since $S_1 \cap E_1$ is irreducible, we may assume that $C_2 \cap S_2 = \phi$ (see Conclusion of (a) and (b1)). Let x be a general point of C and H a general hyperplane section of X through x . Let F_1, \dots, F_n be the exceptional curves of MLT for $(H, K_X + S + B|_H, x)$. We consider by dividing into two cases.

Subcase 1. $K_Z + S_1 + B_1 + E_1|_{g^*H}$ is **PLT**. See FIGURE (XVII).

In this case g_1 coincides with MLT over H . Hence

$$\text{LLC}(K_Z + S_1 + B_1 + E_1|_{E_1}) = (S_1 \cap E_1) \text{ or } l_1 \cup \dots \cup l_m \cup (S_1 \cap E_1),$$

where l_i is a irreducible curve contained in a fiber of g . In the former case R_1 is a flipping ray of type 3 so we are done by (2.2). We assume that the latter case occurs. In this case $m = 1$ holds. For otherwise some l_i would not intersect with $S_1 \cap E_1$ (consider the degree of the boundary of $K_Z + S_1 + B_1 + E_1|_{E_1|_{S_1 \cap E_1}}$). But this contradicts the g -ampleness of S_1 . By considering the degree of the boundary of $K_Z + S_1 + B_1 + E_1|_{E_1|_{l_1}}$, we can prove that there is at most one point in $\text{CLC}_0(K_Z + S_1 + B_1 + E_1|_{E_1})$ outside $S_1 \cap E_1$.

If C_1 does not pass through a point in $\text{CLC}_0(K_Z + S_1 + B_1 + E_1|_{E_1})$ outside $S_1 \cap E_1$, R_1 is a flipping ray of type 3 or 4 so we are done by (2.2) and (2.3). We assume that C_1 passes through the point Q_1 in $\text{CLC}_0(K_Z + S_1 + B_1 + E_1|_{E_1})$ outside $S_1 \cap E_1$. Note that $(\text{Supp } B_1|_{E_1} - l_1).S_1 = 0$. Hence by $S_1.l_1 > 0$ and $S_1.C_1 = 0$ and $\rho(Z_1/Y) = 2$, we have $\text{Supp } B_1|_{E_1} - l_1 \subset C_1$. Let C_1' be the connected component of C_1 containing $\text{Supp } B_1|_{E_1} - l_1$. By $Q_1 \in C_1'$ and (1.9), we can prove as before $C_1' = \text{Supp } B_1|_{E_1} - l_1$. Since $\text{Supp } B_1|_{E_1} - l_1$ is not contained in $\text{LLC}(K_Z + S_1 + B_1 + E_1|_{E_1})$, the type of C_1' is (c) (in this case l_1 corresponds to L).

Subcase 2. Contrary to Subcase 1. See FIGURE (XVIII).

In this case, (1) holds in (6.0).

Claim 2. *Supp $B_1|_{E_1}$ is contained in LLC $(K_Z + S_1 + B_1 + E_1|_{E_1})$ and is a section or a union of a section and an irreducible fiber.*

Proof. Recall that in the construction of g , we take $E = E_1$. Hence the fiber of g over x intersects $\text{Supp } B_1|_{E_1}$ at one point and the intersection point is contained in LLC $(K_Z + S_1 + B_1 + E_1|_{E_1})$ (cf. (1.4)). So $\text{Supp } B_1|_{E_1}$ is a union of a section contained in LLC $(K_Z + S_1 + B_1 + E_1|_{E_1})$ and (possibly empty) components of fibers. Assume that the fiber-part is not empty. Then it must pass through P by (5) in Set up (3.0) and g_1 -ampleness of S_1 and hence the section-part cannot pass through P by (1.1). By (1.7) and the property (3) of a good extraction in (3.3), LLC $(K_Z + S_1 + B_1 + E_1|_{E_1})$ is connected, so some component of the fiber-part is contained in LLC $(K_Z + S_1 + B_1 + E_1|_{E_1})$. By this and (1.1), the fiber-part must be irreducible and contained in LLC $(K_Z + S_1 + B_1 + E_1|_{E_1})$. \square

We can prove as in Subcase 1 that there is at most one point in $\text{CLC}_0(K_Z + S_1 + B_1 + E_1|_{E_1})$ outside $S_1 \cap E_1$.

If C_1 does not pass through a point in $\text{CLC}_0(K_Z + S_1 + B_1 + E_1|_{E_1})$ outside $S_1 \cap E_1$, R_1 is a flipping ray of type 3 or 4 so we are done by (2.2) and (2.3).

We assume that C_1 passes through the point Q_1 in $\text{CLC}_0(K_Z + S_1 + B_1 + E_1|_{E_1})$ outside $S_1 \cap E_1$. In this case we will prove that the former case in Claim 2 cannot occur. Indeed otherwise Q_1 would sit on a section contained in LLC $(K_Z + S_1 + B_1 + E_1|_{E_1})$. Let m be the fiber containing Q_1 and m' its connected component containing Q_1 and intersecting S_1 at a time. After extracting a log crepant curve from Q_1 , m' become contractible but this contradicts (1.6).

Hence we denote the fiber-part of LLC $(K_Z + S_1 + B_1 + E_1|_{E_1})$ by l . Let C'_1 be the connected component of C_1 through Q_1 . By the similar argument to Subcase 1 using (1.9), we can prove $C'_1 = \text{Supp } B_1|_{E_1} - l$. Furthermore C'_1 is of type (b2, $\delta' < \delta$).

7. TREATMENT OF CASE (c)

(7.0) Set up for (c). If $\delta = 0$, we set

$$d := \min\{d(E, S) \mid E \text{ is a log crepant exceptional divisor for } K_X + S + B\}.$$

Let

$$I := \{i \mid E_i \text{ is a log crepant exceptional divisor for } K_X + S + B \text{ with } d_i \leq 1\}$$

if $\delta > 0$ or

$$I := \{i \mid E_i \text{ is a log crepant exceptional divisor for } K_X + S + B \text{ with } d_i = d\}$$

if $\delta = 0$, where $d_i := d(E_i, S)$. Let $h : Z_t \rightarrow X$ be a \mathbb{Q} -factorial LT model of $K_X + S + B$ near S which extracts any divisor E_i with $i \in I$. We apply $N := K_{Z_t} + S_t + B_t + E_t - \epsilon(\sum_{i \in I} d_i E_i)$ -MMP

over X , where S_t (resp. B_t) is the strict transform of S (resp. B) and E_t is the reduced exceptional divisor of h . Since any flip while MMP is a flip of type 1, we can carry out the program. Let $g' : Z' \rightarrow X$ be the end result. Since $N' := K_{Z'} + S' + B' + E' - \epsilon(\sum_{i \in I} d_i E_i) - \delta g'^* S$

($\delta \ll 0$) is KLT and g' -nef and g' -big, its sufficient multiple is g' -free by Kawamata's base point free theorem. Let $h : Z' \rightarrow Z''$ be the morphism defined by such a multiple and $g'' : Z'' \rightarrow X$ be the natural morphism. Then exceptional divisors of g'' are exactly the divisors E_i 's such that $i \in I$. Indeed, for any E_i with $i \in I$ and a general curve in E_i , N is positive while MMP since $N \equiv \epsilon(S_t + \sum_{j \notin I} d_j E_j)$ over X . So any E_i with $i \in I$ is g'' -exceptional. On the other hand some

h_t -exceptional divisors E_j 's with $j \notin I$ are g'' -exceptional, for at least one E_j and a general curve in E_j , $h'(N')$ is numerically nonpositive, a contradiction. Note that $g''^{-1}S$ is normal since S is normal and $g''^{-1}S$ is normal along intersection curves with g'' -exceptional divisors. Let the chain $\cup_{i \in I} E_i \cap g''^{-1}S$ be $\cup_{k=1}^n D_k$, where D_1 intersects $g''^{-1}C$ and D_n intersects $g''^{-1}L$ (if L is the center of some E_i with $i \in I$, D_n is equal to $g''^{-1}L$). Then we encounter the following two possibilities:

- (1) $g''^{-1}C \cap D_1$ is contained in $\text{CLC}_0(g''^*(K_X + S + B)|_{g''^{-1}S})$;
- (2) $g''^{-1}C \cap D_1$ is not contained in $\text{CLC}_0(g''^*(K_X + S + B)|_{g''^{-1}S})$.

We can express above two possibilities in a more concrete way:

- (1)
 - (1.1) $D_1 \neq g''^{-1}L$ (and hence $D_1 \simeq \mathbb{P}^1$) and $g''^*(K_X + S + B)|_{g''^{-1}S}|_{D_1} = K_{D_1} + P' + P$, where $P := D_1 \cap D_2$ if $n \geq 2$ and $P := D_1 \cap g''^{-1}L$ if $n = 1$
 - or
 - (1.2) $D_1 = g''^{-1}L$ and $P' := g''^{-1}C \cap g''^{-1}L$;
- (2) $D_1 \neq g''^{-1}L$ (and hence $D_1 \simeq \mathbb{P}^1$) and $g''^*(K_X + S + B)|_{g''^{-1}S}|_{D_1} = K_{D_1} + \frac{1}{2}P_1 + \frac{1}{2}P_2 + P$, where $P := D_1 \cap D_2$ if $n \geq 2$ and $P := D_1 \cap g''^{-1}L$ if $n = 1$.

See FIGURE (XIX). In any case, we call E the exceptional divisor which cut out D_1 and $d := d(E, S)$ (it is consistent with the above definition of d). We remark that $g''^{-1}S$ and E cross normally along D_i . In fact otherwise there is a log crepant divisor F for $K_X + S + B$ whose center is D_i with $d(F, g''^{-1}S + E) \leq 1$. If $\delta > 0$, $1 \geq d(F, g''^{-1}S + E) \geq d(F, g''^{-1}S + dE)$, i.e., $d(F, S) \leq 1$. But this contradicts the property of h . If $\delta = 0$,

$$d \geq d(F, dg''^{-1}S + dE) \geq d(F, g''^{-1}S + d(E, S)E), \text{ i.e., } d(F, S) \leq d.$$

But this again contradicts the property of h .

We remark the following:

Claim. *Above division into cases (1-1), (1-2) and (2) depends only on f .*

Proof. By running $g''^*(K_X + S + B) - \epsilon(g''^{-1}S + dE)$ -MMP and taking the LC model as in (7.0), we obtain the primitive extraction of E from g'' . We can check the above procedure is isomorphism near P' in case (1) of (7.0) (resp. near P_1 and P_2 in case (2) of (7.0).)

Assume that the case (2) occurs in the above. Note that even after making g'' a primitive extraction of E , the property (2) holds for this primitive contraction.

Assume furthermore that there is another \tilde{g}'' with the similar properties to g'' but the case (1) occurs. Then we can obtain the primitive extraction of the strict transform of E from \tilde{g}'' and after making \tilde{g}'' a primitive extraction of E , the property (1) for this primitive contraction. Hence we obtain two different primitive extraction of E . But this contradicts [FA, Lemma 6.2].

If the case (1) occurs, then by above argument, (2) does not occur for any \tilde{g}'' .

It is determined only by f whether L is a center of an exceptional divisor E_i for $i \in I$. Hence it is determined only by f whether (1-1) or (1-2) occurs. This completes the proof. \square

We treat the case (1) and (2) separately in the below. We first treat the case (1). We deduce the following from the condition of (1).

Claim. *For any log crepant divisor F for $K_X + S + B$ whose center is P' or a curve through P' , $d(F, g''^{-1}S + E) > 1$.*

Proof. Note that among E_i with $i \in I$, E is the unique divisor whose center contains P' . Hence if $\delta > 0$, $d(F, g''^{-1}S + E) \geq d(F, g''^{-1}S + dE) = d(F, S) > 1$ and if $\delta = 0$, $d(F, g''^{-1}S + E) \geq d(F, \frac{1}{d}(g''^{-1}S + dE)) = \frac{1}{d}d(F, S) > 1$. \square

Note that $K_{Z''} + g''^{-1}S + g''^{-1}B$ and $K_{Z''} + E + g''^{-1}B$ is PLT at P' by normality of $g''^{-1}S$ and E at P' , and the inversion of adjunction (cf. the subadjunction formula in (1) above). By these, we obtain the following singularity.

(*) (Z, P) is an algebraic germ of a normal 3-fold singularity, S_1 and S_2 are irreducible \mathbb{Q} -Cartier divisors and B is a \mathbb{Q} -Cartier boundary such that $\lfloor B \rfloor = \phi$ with the following properties:

- (i) $K_Z + S_1 + S_2 + B$ is LC and $K_Z + S_i + B$ is PLT for $i = 1, 2$;
- (ii) $2(K_Z + S_1 + S_2 + B) \sim 0$;
- (iii) $C := S_1 \cap S_2$ is a smooth curve and $K_Z + S_1 + S_2 + B|_{S_i}|_C = K_C + P$, i.e., $K_Z + S_1 + S_2 + B|_{S_i}$ is not PLT at P for $i = 1, 2$;
- (iv) for any log crepant divisor F with respect to $K_Z + S_1 + S_2 + B$ whose center is P or a curve through P , $d(F, S_1 + S_2) > 1$.

We examine the (*) for a while and prove

Theorem (7.1) (Classification of (*)). *We consider the object (*). Then one of the following holds:*

- (1) P is smooth on Z , S_1 and S_2 ;
- (2) locally analytically $(P \in S_1 \cup S_2 \subset Z) \simeq (o \in (xy = 0) \subset \frac{1}{3}(2, 2, 1))$. The weighted blow up $h : Z' \rightarrow Z$ with weight $\frac{1}{3}(2, 2, 1)$ is log crepant for $K_Z + S_1 + S_2 + B$. Let F' be the exceptional divisor of h . Then $h^*(K_Z + S_1 + S_2 + B)|_{F'} = K_{F'} + M_1 + M_2 + \frac{1}{2}M_3 + \frac{1}{2}M_4$, where all M_i 's are irreducible curves, $M_i := h^{-1}S_i \cap F'$ for $i = 1, 2$, $M_3 := \text{Supp } h^{-1}B \cap F'$ and M_4 is an ODP curve singularity. On F' , $h^*(K_Z + S_1 + S_2 + B)$ is LT; See FIGURE (XX).
- (3) locally analytically $(P \in S_1 \cup S_2 \subset Z) \simeq (o \in (xy = 0) \subset \frac{1}{2}(1, 0, 1))$.

We begin with some lemmas.

Lemma (7.2). *Assume that Z is not smooth at P . Let \tilde{B} be a \mathbb{Q} -Cartier boundary such that*

- (i) $K_Z + S_1 + S_2 + \tilde{B}$ is LC and $K_Z + S_i + \tilde{B}$ is PLT for $i = 1, 2$;
- (ii) $2(K_Z + S_1 + S_2 + \tilde{B}) \sim 0$.

Then $K_Z + S_1 + S_2 + \tilde{B}|_{S_i}|_C = K_C + P$, i.e., $K_Z + S_1 + S_2 + \tilde{B}|_{S_i}$ is not PLT at P for $i = 1, 2$.

Proof. Assume the contrary to what we want, i.e., $K_Z + S_1 + S_2 + \tilde{B}|_{S_i}$ is PLT at P for $i = 1, 2$. By the existence of $\tilde{B}|_{S_i}$ and the description of index 2 PLT surface singularity (see (1.4)), we know that S_i is smooth at P and $\text{Diff}_{S_i}(0) = 0$. Hence by (1.11), Z is also smooth, a contradiction. \square

Lemma (7.3). *(Z, P) is analytically \mathbb{Q} -factorial. $K_Z + S_1 + S_2 + B$ is LT outside P and curves where S_i and $\text{Supp } B$ are simply tangent. In particular $K_Z + S_1 + S_2$ is LT.*

Proof. If (Z, P) is not \mathbb{Q} -factorial, by (1.13), we can take an analytically \mathbb{Q} -factorial LT model $\pi : \tilde{Z} \rightarrow Z$ for $K_Z + S_1 + B$ near S_1 . By (i) of (*), $K_{Z_1} + S_1 + B$ is PLT so this modification

is a small morphism. Since S_1, S_2 and B is \mathbb{Q} -Cartier, the fiber over P is contained in the inverse images of S_1, S_2 and B . But this contradicts (1.2). We will prove the rest. Let $P' \neq P$ be a point on Z and Z' a general hyperplane section of Z through P' . On Z' , we express the restricted divisors with $'$. Note that the property (iv) of $(*)$ holds for $K_{Z'} + S'_1 + S'_2 + B'$ and P' . If P' is singular on Z' and $K_{Z'} + S'_1 + S'_2 + B'$ is not LT, there is a log crepant curve F for $K_{Z'} + S'_1 + S'_2 + B'$ such that $d(F, S'_1 + S'_2) \leq 1$ by (1.3), a contradiction. If P' is nonsingular on Z' and $K_{Z'} + S'_1 + S'_2 + B'$ is not LT, then by (iv) of $(*)$ and (1.4), P' is contained in only one S'_i and S'_i is simply tangent to $\text{Supp } B'$ at P' . So we obtain the result. \square

Lemma (7.4). *$K_Z + S_1 + S_2$ is 1-complemented. Assume that Z is not smooth. Let \bar{B} be an 1-complement. Then possibly after replacing B while keeping $(*)$, $a_l(K_Z + S_1 + S_2 + B) \geq a_l(K_Z + S_1 + S_2 + \bar{B})$ holds for any exceptional divisor.*

Proof. By (7.3), $K_Z + S_1 + S_2$ is LT. Hence there exists a good resolution $r : Z' \rightarrow Z$ for $K_Z + S_1 + S_2$ such that all log discrepancy for $K_Z + S_1 + S_2$ are positive and $r^{-1}S_1 \cap r^{-1}S_2 \neq \emptyset$. Set $K_{Z'} + r^{-1}S_1 + r^{-1}S_2 + G = r^*(K_Z + S_1 + S_2)$. Then all the coefficients in G are < 1 and the left hand side (we call this D) is LT (subboundary version). Let H be a very ample divisor on Z' and $r^*r_*H := H + \sum g_i G_i$, where G_i 's are exceptional divisors. Then $-\sum g_i G_i$ is r -ample. Hence for $\epsilon \ll 1$, $D' := -(K_{Z'} + r^{-1}S_1 + r^{-1}S_2 + G + \epsilon \sum g_i G_i)|_{r^{-1}S_1}$ is LT and $r|_{r^{-1}S_1}$ -ample. Furthermore $r^{-1}S_2|_{r^{-1}S_1} \neq \emptyset$ and is not contained in a fiber of r . So by [FA, 19.11, Corollary], there is a 1-complement for D' near any fiber intersecting $r^{-1}S_2|_{r^{-1}S_1}$. Hence there is a 1-complement for $K_Z + S_1 + S_2|_{S_1}$. This is lifted to a 1-complemented for $K_{Z'} + r^{-1}S_1 + r^{-1}S_2 + G$ by inversion of adjunction.

Next we assume that Z is not smooth. By (7.2), if we replace B by a more general element such that $a_l(K_Z + S_1 + S_2 + B) \geq a_l(K_Z + S_1 + S_2 + \bar{B})$ for any exceptional divisor, the conditions of $(*)$ are preserved. Hence we are done. \square

The next lemma is a key to the proof of (7.1).

Lemma (7.5). *Assume that Z is not smooth. Let F be a exceptional divisor over P (not over a curve through P) with $a_l(F, K_Z + S_1 + S_2) \leq 1$. Then F satisfies one of the following:*

- (1) $d(F, S_1 + S_2) > 1$;
- (2) *there is a log crepant divisorial extraction $Z' \rightarrow Z$ for $K_Z + S_1 + S_2 + B$ whose exceptional divisor F' contains a generically ODP curve singularity of Z' and F is extracted by the simple blow up of the curve singularity. $a_l(F, K_Z + S_1 + S_2 + B) = \frac{1}{2}$ and $d(F, S_1 + S_2) = \frac{1}{2}d(F', S_1 + S_2) > \frac{1}{2}$.*

Proof. By (ii) of $(*)$ and the assumption that the center of F is P , $a_l(F, K_Z + S_1 + S_2 + B) = 0$ or $\frac{1}{2}$. If the former case occurs, F satisfies (1) by (iv) of $(*)$. We assume that the latter case occurs. Let $h : Z_t \rightarrow Z$ be a \mathbb{Q} -factorial LT model for $K_Z + S_1 + S_2 + B$ near $S_1 \cup S_2$ and F_j 's exceptional divisors. F lands on some F_j . Let $(S_i)_t$ be the strict transform of S_i .

Case 1. F is contracted to a point z on Z_t . Then z is contained in at most two of F_j 's and $(S_i)_t$'s. In fact, if z is contained in three of F_j 's and $(S_i)_t$'s, z is smooth and is not contained in any other component of the boundaries of $h_t^*(K_Z + S_1 + S_2 + B)$ by (1.12). Hence the index of $h_t^*(K_Z + S_1 + S_2 + B)$ at z is 1. But this contradicts $a_l(F, K_Z + S_1 + S_2 + B) = \frac{1}{2}$. So we divide into two cases as follows:

Subcase 1.1. z is contained in two of F_j 's and $(S_i)_t$'s. Let such two components D_1 and D_2 . Near z , $h^*(K_Z + S_1 + S_2 + B)|_{D_i} = K_{D_i} + \text{Diff}_{D_i}(0) + D_{2-i}|_{D_i} + h^{-1}B|_{D_i}$ holds. It is LT and of index 2 at z . By (1.4) and (1.12), we easily deduce that $z \in Z$ is a $\frac{1}{2}(1, 1, 1)$

singularity if $z \notin h^{-1}B$ or z is smooth point of Z if $z \in h^{-1}B$. In any case $D_1 + D_2$ is Cartier at z . On the other hand $d(D_1 + D_2, S_1 + S_2) > 1 + 1 = 2$ by the assumption (iv) of (*). Hence $d(F, S_1 + S_2) > 1$. So F satisfies (1).

Subcase 1.2. z is contained in only F_j . We prove that this case does not occur. Take a 1-complement \overline{B} as in (7.4). Since we may assume that B is general more than $2\overline{B}$ (see (7.4)), $a_l(F, K_Z + S_1 + S_2 + \overline{B}) \leq \frac{1}{2}$, which in turn show that $a_l(F, K_Z + S_1 + S_2 + \overline{B}) = 0$ because the index of $K_Z + S_1 + S_2 + \overline{B}$ is 1. By the similar way, we know that $a_l(F_j, K_W + S_1 + S_2 + \overline{B}) = 0$. Assume that $z \notin \text{Supp } h^{-1}\overline{B}$. Then $K_W + F_j$ is PLT and of index 1 so $a_l(F, K_W + S_1 + S_2 + \overline{B}) \geq 1$, a contradiction. Assume that $z \in \text{Supp } h^{-1}\overline{B}$ and let $D := h^{-1}\overline{B} \cap F_j$. Then

$$K_W + h^{-1}S_1 + h^{-1}S_2 + h^{-1}\overline{B} + F_j|_{F_j} = K_{F_j} + D' + D \text{ near } z,$$

where D' is $\text{Diff}_{F_j}(0)$ (contribution of D). But the index of this log divisor is 1 and $K_W + F_j$ is PLT so $D' = 0$. Hence the index of $K_{F_j} + D$ is 1 and so $K_{F_j} + D$ is PLT. Hence $a_l(F, K_Z + S_1 + S_2 + \overline{B}) \geq 1$, a contradiction. Now we exclude this case.

Case 2. F is contracted to a curve l . By (1.2), l is contained in at most two of F_j 's and $(S_i)_t$. If l is contained in F_j and F_k (or $(S_i)_t$), then l is not contained in any component of $\text{Supp } B_t$. Hence the index of $h_t^*(K_Z + S_1 + S_2 + B)$ along l is 1, a contradiction to $a_l(F, K_Z + S_1 + S_2 + B) = \frac{1}{2}$. So l is contained in only one F_j . For a general point P' on l and a general hyperplane section Z'_t of Z_t through P' , $K_{Z'_t} + B'_t + F_j := h_t^*(K_Z + S_1 + S_2 + B)|_{Z'_t}$ is LT of index 2 and not of index 1 by the existence of F whose log discrepancy is $\frac{1}{2}$. So by (1.4), $P' \notin \text{Supp } B'_t$ and P' is an ODP on Z'_t or $P' \in \text{Supp } B'_t$ and P' is smooth on Z'_t . Note that

$$(7.5.1) \quad d(F, S_1 + S_2) = d(F, (S_1)_t + (S_2)_t) + d(F_j, S_1 + S_2)F_j = d(F, F_j)d(F_j, S_1 + S_2).$$

For the latter case, $d(F, F_j)$ is an integer. Hence F satisfies (1) by (7.5.1). For the former case, if $d(F, S_1 + S_2) \leq 1$ then $d(F, F_j) = \frac{1}{2}$ by (7.5.1) and hence F is the exceptional divisor of the simple blow up of l . So F satisfies (2).

Now we finish the proof of (7.5). \square

Proof of Theorem (7.1). Since $K_Z + S_1 + S_2$ is LT and S_1 and S_2 are \mathbb{Q} -Cartier, we have $(P \in S_1 \cup S_2 \subset Z) \simeq (o \in (xy = 0) \subset \frac{1}{s}(t_1, t_2, t_3))$, where t_i is a nonnegative integer less than s and $(s, t_1, t_2) = 1$ by (1.12).

First we treat the case P is an isolated singularity.

Claim 1. *If P is an isolated singularity, then P is a terminal singularity.*

Proof. We may assume that P is singular. Let F be any exceptional divisor over P . If $a_l(F, K_Z + S_1 + S_2) \leq 1$, then by (7.5), $a_l(F, K_Z) = a_l(F, K_Z + S_1 + S_2) + d(F, S_1 + S_2) > 1$. Hence for any F , we have $a_l(F, K_Z) > 1$ and we are done. \square

If P is smooth on Z , we obtain (1) by (1.12). Assume below that P is singular point of Z . Then by (1.12), the index of P is greater than 1. Hence by Kawamata's minimal discrepancy theorem [Ka 2], there is a exceptional divisor F over P with $a_l(F, K_Z) = 1 + \frac{1}{r}$, where r is the index of P . Since P is terminal, $a_l(F, K_Z + S_1 + S_2) \in \frac{\mathbb{Z}}{r}$ and $a_l(F, K_Z + S_1 + S_2) < 1 + \frac{1}{r}$ by $a_l(F, K_Z) = 1 + \frac{1}{r}$. Hence $a_l(F, K_Z + S_1 + S_2) \leq 1$ and we apply (7.5). If F satisfies (1) of Lemma (7.5), then $d(F, S_1 + S_2) \geq 1 + \frac{1}{r}$ since P is terminal. On the other hand, $a_l(F, K_Z + S_1 + S_2) > 0$ since $a_l(F, K_Z + S_1 + S_2 + B) \geq 0$ and the center of F on Z is P . But these contradict

$$(7.1.1) \quad d(F, S_1 + S_2) + a_l(F, K_Z + S_1 + S_2) = 1 + \frac{1}{r}.$$

So F satisfies (2). By $a_l(F, K_Z + S_1 + S_2 + B) = \frac{1}{2}$, $a_l(F, K_Z + S_1 + S_2) > \frac{1}{2}$. On the other hand $d(F, S_1 + S_2) > \frac{1}{2}$. By these and (7.1.1), r must be odd (we set $r = 2r' + 1$) and $a_l(F, K_Z + S_1 + S_2) = d(F, S_1 + S_2) = \frac{r'+1}{2r'+1}$. Furthermore by the theorem of Morrison-Stevens [MS, Theorem 2.4] and that $S_1 + S_2$ is not a Cartier divisor (since $d(F, S_1 + S_2)$ is not an integer), we can take $s = 2r' + 1$, $t_1 + t_3 = 2r' + 1$ and $t_2 = 1$. By the weighted blow up with a weight $\frac{1}{2r'+1}(t_1, 1, 2r'+1-t_1)$, F is extracted by the uniqueness of the exceptional divisor with minimal discrepancy for terminal quotient singularity (see [Ka3]). So $\frac{r'+1}{2r'+1} = d(F, S_1 + S_2) = \frac{t_1+1}{2r'+1}$. Hence $t_1 = r'$. We prove $r' = 1$, which in turn show that (2) of (7.1) holds with a weight change. It suffices to prove that if $r' \geq 2$, F must be log crepant for $K_Z + S_1 + S_2 + B$ (this is a contradiction). Let F'' be the curve which is extracted by the restriction of the above weighted blow up to $y = 0$. Let $F_1, F_2, \dots, F_{2r'}$ be the exceptional curve of the MRS of the singularity of the origin on $y = 0$, where F_1 intersects the strict transform of $x = y = 0$. Then F'' is $F_{r'+1}$ on the MRS by the choice of weight. Hence if $r' \geq 2$, F'' must be log crepant for $K_Z + S_1 + S_2 + B|_{S_2}$ by (1.3), which in turn show that F is log crepant for $K_Z + S_1 + S_2 + B$, a contradiction. The latter half of (2) is an easy calculation.

Next we treat the case that P' is non-isolated singularity. We will prove that P satisfies (3) of (7-1). Since no curve singularity through P' is contained in LLC ($K_Z + S_1 + S_2 + B$) by (7.3) and any curve singularity through P' is a generically ODP curve since the index of $K_Z + S_1 + S_2 + B$ is 2. Let $\tilde{Z} := \mathbf{Spec}(\mathcal{O}_Z \oplus \mathcal{O}_Z(-S_1 - S_2))$. Then $\pi : \tilde{Z} \rightarrow Z$ is a double covering such that $\tilde{P} := \pi^{-1}P$ is an isolated singularity on \tilde{Z} . It is easy to see that $K_{\tilde{Z}} + \tilde{S}_1 + \tilde{S}_2 + \tilde{B} := \pi^*(K_Z + S_1 + S_2 + B)$ satisfies the condition (*). If \tilde{P} satisfies (1), we obtain what we want. We show that \tilde{P} does not satisfies (2). Assume that \tilde{P} satisfies (2). Then it is easy to see that locally analytically $(P \in S_1 \cup S_2 \subset Z) \simeq (o \in (xy = 0) \subset \frac{1}{6}(5, 2, 1))$. Take the weighted blow up $h : W \rightarrow Z$ with the weight $\frac{1}{6}(5, 2, 1)$ and let F be the exceptional divisor. The restriction of h to S_1 is the MRS of P which extracts a (-3) -curve so h is log crepant for $K_Z + S_1 + S_2 + B$ by (1-3). Then at the intersection of F and $h^{-1}S_2$, there is a singularity Q such that $(Q \in F \cup h^{-1}S_2 \subset W) \simeq (o \in (xy = 0) \subset \frac{1}{5}(2, 1, 3))$. Since the index of $h^*(K_Z + S_1 + S_2 + B)$ is 2 at the singularity, $\text{Supp } h^{-1}B$ passes through it and neither $h^*(K_Z + S_1 + S_2 + B)|_F$ or $h^*(K_Z + S_1 + S_2 + B)|_{h^{-1}S_2}$ is PLT. Hence the weighted blow up with the weight $\frac{1}{5}(2, 1, 3)$ is log crepant for $h^*(K_Z + S_1 + S_2 + B)$ since the restriction of this weighted blow up to $h^{-1}S_2$ is log crepant for $h^*(K_Z + S_1 + S_2 + B)|_{h^{-1}S_2}$ by (1.3). Let F' be the exceptional divisor of this weighted blow up. Then $d(F', S_1 + S_2) = d(F', h^{-1}S_2 + \frac{7}{6}F) = \frac{2}{3}$. But this contradicts the property of (*).

□

Lemma (7.6) (Reducible case). *Consider the case (c). Let $g : Z \rightarrow X$ be a primitive log crepant divisorial extraction for $K_X + S + B$ and E the exceptional divisor. Assume the following conditions:*

- (1) $g^{-1}S \cap E$ is reducible (and hence by Lemma (5.2), we can write $g^{-1}S \cap E = D_1 \cup D_2$);
- (2) $D_1 \simeq \mathbb{P}^1$. $g^{-1}S$ and E are generically normal crossing along D_1 (Note that we assume nothing about D_2 here.);
- (3) $K_Z + g^{-1}S + g^{-1}B + E|_{D_1} = K_{D_1} + \frac{1}{2}P_1 + \frac{1}{2}P_2 + P$;
- (4) $g^{-1}S$ is normal.

Run the BTM starting by g .

Then D_1 becomes a flipping ray satisfying the assumptions of (2.4) after the flip of all the strict transforms of the components of C . It satisfies the following numerical properties: $B'.D_1 = 0, S'.D_1 < 0$ and $E'.D_1 > 0$, where B', S' and E' are the strict transform of B, S and E (by abuse of notation, we express the strict transform of D_1 also by D_1). In particular g is

a semi-good extraction by the property (2) of (2.4).

Proof. We use the notation as in (2.5). In particular we write S_1 , B_1 and E_1 instead of $g^{-1}S$, $g^{-1}B$ and E . By an abuse of notation, we also call C , D_1 and D_2 the strict transforms of C , D_1 and D_2 . In a neighborhood of C on Z , $C = \text{Supp } B_1|_{S_1}$ and C is contractible on S_1 . Hence there exists a component C' of C such that $B_1.C' < 0$ and $E_1.C' > 0$. On the other hand, for an exceptional curve l for g , $B_1.l > 0$ and $E_1.l < 0$, so we must have $B_1.C_1 < 0$, $E_1.C_1 > 0$ and $S_1.C_1 < 0$. Hence $C_1 \subset S_1 \cap \text{Supp } B_1$, which in turn shows that $C_1 \subset C$. By the above numerical property, C_1 is a beginning flipping ray and the flip exists.

Claim. *After the flip of C_1 , Any component of C_1^+ is not contained in S_2 .*

Proof. If some component $(C_1^+)'$ is contained in S_2 , it contained in $S_2 \cap E_2$. In particular $(C_1^+)' \subset \text{LLC}(K_{Z_2} + S_2 + B_2 + E_2)$. Hence C_1 contains an element of $\text{CLC}(K_{Z_1} + S_1 + B_1 + E_1)$ since $(K_{Z_2} + S_2 + B_2 + E_2).C_1^+ = 0$. But this contradicts the assumption of case (c) and (3) of this lemma. \square

Note that E_2 must be normal since E_1 is normal by Remark (5.3) and the proof of Claim. If some components of C left on S_2 , then as above there exists a component C'' of C such that $B_2.C'' < 0$ and $E_2.C'' > 0$. On the other hand, for the flipped curve C_1^+ , $B_2.C_1^+ > 0$ and $E_2.C_1^+ < 0$, so we must have $B_2.C_2 < 0$, $E_2.C_2 > 0$ and $S_2.C_2 < 0$. Hence as above we see that $C_2 \subset C$ and any component of C_2^+ is not contained in S_3 . By the above numerical property, we can easily see that $E_2.C_2 > 0$ and $S_2.C_2 < 0$ so the flip of C_2 is a beginning flip. (After the flip, E_3 is normal.) Since the number of the components of C on X is at most 2, no component of C lefts on the strict transform of S . After the flips of the strict transform of C , we call Z' the image of Z and S' , B' and E' the strict transform of S , B and E . Let R be the next extremal ray. We show that R is a flipping ray stated as in the statement of this lemma. We claim that first $B'.D_1 = 0$. After the flips of the strict transform of C , $\text{Supp } B'$ intersects S' at most in finite points near D_1 since the flipped curves are not contained in S' but by \mathbb{Q} -factoriality, $\text{Supp } B' \cap S'$ must be empty near D_1 . Hence we obtain $B'.D_1 = 0$. On the other hand, B' is positive for the flipped ray one before R . Hence we must have $B'.R \leq 0$.

We prove that $B'.R < 0$ does not occur. Assume that $B'.R < 0$. Then we see that neither D_1 or D_2 is not contained in $\text{Supp } R$ since neither D_1 or D_2 is contained in $\text{Supp } B'$. Hence by the numerical property (2) in (2.5), we must have $S'.R > 0$. Hence by the numerical property (1) in (2.5), $E'.R < 0$, i.e., R is a final flipping ray (it is not divisorial since $B'.R < 0$). Note that $D_1 \cap \text{Supp } R = \emptyset$ since $\text{Supp } R \subset \text{Supp } B'$ and $D_1 \cap \text{Supp } B = \emptyset$. We see inductively that after the flip of R , an extremal ray is negative for B' and E' and positive for S' as long as D_1 is not a flipping curve while BTM (hence it is not a divisorial ray and its support does not intersects D_1). Hence after finite number of such a flip, an extremal ray R'' such that $B''.R'' = 0$ appears. We show that R'' is not divisorial. If R'' is divisorial, after the contraction we obtain the flip of C . Let C^+ be the flipped curve. Then $B^+.C^+ < 0$ since $B.C > 0$. If E'' is contracted to a point, C^+ passes through the contracted point. But by $B''.R'' = 0$, C^+ cannot be contained in $\text{Supp } B^+$, a contradiction. If R'' is contracted to a curve, the image curve is not contained in $\text{Supp } B^+$ since $B''.R'' = 0$. Hence the center of E on X cannot be L and the image of E'' is contained in C^+ . But this again contradicts $B^+.C^+ < 0$.

Hence R'' is not divisorial and D_1 is a flipping curve (I don't know if $\text{Supp } R'' = D_1$.) Here we consider the flipping contraction of R'' in the analytic category. First we flip only D_1 . Since E' is normal and above procedures after the flip of the strict transform of C are isomorphic near D_1 , E'' is normal near D_1 . Hence we can check that D_1 satisfies the assumption of (2.4). Hence by (2.4), after the flip, D_1 go outside E'' . Next we flip other components of $\text{Supp } R''$. While such flips, D_1 never become a flipping curve again. In fact the numerical property (1) in

(2.5) still holds in the analytic category. While such flips, $E''.D_1$ continues to be positive and hence $S''.D_1$ continues to be negative. So $(K_{Z''} + E'').D_1$ continues to be positive. But such flipping curves (other component of $\text{Supp } R''$) are negative for $K_{Z''} + E''$. Hence we obtain the assertion.



After the flip of R'' (and possibly the flips of some final flipping rays), finally D_1 become a curve contained in C^+ . But D_1 is not contained in $\text{Supp } B^+$, a contradiction to $B^+.C^+ < 0$.

Hence we have $B'.R = 0$. we immediately see that $\text{Supp } R \subset D_1$. By the numerical property (2) in (2.5), we must have $S'.R > 0$ or $E'.R > 0$. We can show the former case cannot occurs just as above. Hence the latter case occurs and We can easily check R satisfies the assumptions of (2.4) by the assumptions of this Lemma. \square

Lemma (7.7) (Reducible case 1). *Consider the situation as in (7.6). Assume furthermore that $g(E) = Q$.*

Then we have the description as explained in the following flow chart:

D_1 become a flipping ray of the type in (2.4). Consider the next extremal ray R'



↓

- (1) $Q' := D_2 \cap g^{-1}L$ is smooth on $g^{-1}S$, E and Z . $(D_2)_E^2 = 2$. $\text{Supp } B_1|_E$ and D_2 are simply tangent at Q' ;
- (2) P is smooth on $g^{-1}S$, E and Z ;
- (3) $d = \frac{3}{-(D_2)_{g^{-1}S}^2 - 1}$;
- (4) $d(F, S) = d + 1$ and $a_l(F, S) = 0$ for the exceptional divisor F of the simple blow up along D_1 or D_2 and for any other log crepant divisor $F \neq E$ for $K_X + S + B$, $d(F, S) > d + 1$.

$$\text{LLC} := \text{LLC} (K_{Z^+} + S^+ + B^+ + E^+|_{E^+}).$$

$$\text{CLC}_0 := \text{CLC}_0 (K_{Z^+} + S^+ + B^+ + E^+|_{E^+}).$$

See also FIGURE (XXI).

Proof. When D_1 becomes a flipping ray, we use the same notation as in (7.6) (S' , E' ..., etc). After the flip of D_1 , we call the flipped curve D_1^+ and S^+ , B^+ , E^+ and D_2^+ the strict transforms of S , B , E and D_2 . Let R' be the next extremal ray. Since $S^+ \cap E^+$ is irreducible, we can assume that $\text{Supp } R' \cap S^+ = \phi$ and $E^+ \cdot R' < 0$ (see the flow chart). Furthermore we can assume that $\text{Supp } R' \cap \text{LLC} (K_{Z^+} + S^+ + B^+ + E^+|_{E^+}) \neq \phi$, where $\nu : E^{+\nu} \rightarrow E^+$ is the normalization of E^+ . Let $D^* = \mu^{-1}D_1^+$.

Claim 1. $\text{LLC} (K_{Z^+} + S^+ + B^+ + E^+|_{E^+}) = D^* \cup D_2^+$.

Proof. If $\text{LLC} (K_{Z^+} + S^+ + B^+ + E^+|_{E^+})$ has another component except D^* and D_2^+ , $\text{LLC} (K_Z + S + B + E|_E)$ has another component D' except D_1 and D_2 , and D' must intersect D_2 at a point ($\neq P$) by (3.4). Furthermore $\text{LLC} (K_{Z^+} + S^+ + B^+ + E^+|_{E^+}) = D^* \cup D_2^+ \cup D'^+$, where D'^+ is the strict transform of D' , and $D^* \cap D'^+ = \phi$ and $D_2^+ \cap D'^+ \neq \phi$. Note

that $(D_2^+)^2_{E^{+\nu}} = S^+ \cdot D_2 > 0$ since $S^+ \cdot R' = 0$ and $S^+ \cdot D_1^+ > 0$. Hence $aD_2^+ - (K_{E^{+\nu}} + \{\text{Diff}_{E^{+\nu}}(S^+ + B^+)\}) \equiv aD_2^+ + \perp \text{Diff}_{E^{+\nu}}(S^+ + B^+) \perp$ is nef and big for $a \gg 0$. Furthermore $K_{E^{+\nu}} + \{\text{Diff}_{E^{+\nu}}(S^+ + B^+)\}$ is KLT by Theorem (1.7) (2), so by Kawamata's base point free theorem, D_2^+ is semi-ample.

On the image \overline{E} of $E^{+\nu}$ by the morphism defined by a sufficient multiple of D_2^+ , take a $K_{\overline{E}} + (1 - \epsilon)(\overline{\text{Diff}_{E^{+\nu}}(S^+ + B^+) - \overline{D'^+}})$ -extremal ray A ($0 < \epsilon \ll 1$ is a sufficiently small rational number) such that $\overline{D'^+} \cdot A > 0$. We can take such a ray since $\overline{D'^+}^2 > 0$ (cf. Claim 3 in the proof of (1.9)). Note that $\overline{D_2^+}$ is ample. If A is birational, by $\overline{D'^+} \cdot A > 0$ and the ampleness of $\overline{D_2^+}$, $\text{Supp } A$ intersects $\overline{D_2^+}$ and $\overline{D'^+} \cdot A > 0$. Hence by (1.6), $\text{Supp } A \cap \overline{D_2^+} \cap \overline{D'^+}$ is one point. We take this procedure till the extremal ray is not birational. Note that the situation does not change while this procedure. So we may assume that the image of \overline{E} by the extremal contraction associated to A is a curve. Since $\overline{D'^+} \cdot A > 0$, $\overline{D^*}$ is not contained in a fiber. (Note that $\overline{D^*}$ does not intersect $\overline{D'^+}$ by applying (1.6) to the morphism defined by sufficiently multiple of D_2^+ .) Hence a general fiber of the contraction of A intersects LLC $(K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}})$ with at least three points. But this is impossible since for a general fiber r , $K_{E^{+\nu}} \cdot r = -2$.

The contraction of A cannot contract \overline{E} to a point since $\overline{D^*}$ and $\overline{D'^+}$ does not intersect.

This is a final contradiction and we finish the proof of Claim 1. \square

Subcase 1. $\text{CLC}_0(K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}}) \cap \text{Supp } R' = \phi$

Let l be any connected component of $\text{Supp } R'$. In this case we prove that after perturbing E^+ near l , the flip of l become a flip of type 3.

By assumption of Subcase 1, $K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}}$ is exceptional near l . By (2.4) (4), the nonnormal locus of E^+ is D^* . Hence l is not contained the nonnormal locus of E^+ and taking a more general element E' than E^+ in $|E^+ \supset l|$ near l , E' is normal. Since LLC $(K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}})$ is not contained in $\text{Supp } B^+|_{E^{+\nu}}$ near l , $K_{Z^+} + S^+ + B^+ + E'|_{E'}$ has at most isolated LLC near l . If $K_{Z^+} + S^+ + B^+ + E'|_{E'}$ has isolated LLC near l , LLC $(K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}})$ also has isolated LLC near l . But this contradicts the assumption of Subcase 1.

Then $K_{Z^+} + S^+ + B^+ + E'|_{E'}$ is KLT and by the inversion of adjunction, $K_{Z^+} + S^+ + B^+ + E'$ is PLT.

Subcase 2. $\text{CLC}_0(K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}}) \cap \text{Supp } R' \neq \phi$

In this case we will prove (1), (2), (3) and (4) of this Lemma. We first prove (1). Since $D^* \cap D_2^+$ is contained in $\text{CLC}_0(K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}})$, D^* has exactly one point in $\text{CLC}_0(K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}})$ other than $D^* \cap D_2^+$. We call this Q'' . Let l be the connected component of $\text{Supp } R'$ containing Q'' .

Claim 2. *There is no log crepant primitive extraction $\mu : E' \rightarrow E^{+\nu}$ for $K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}}$ over Q' such that $(\mu^{-1}D_2^+)^2 \geq 0$ and $\text{Supp } \mu^{-1}(B^+|_{E^{+\nu}}) \cap \mu^{-1}D_2^+ = \phi$.*

Proof. Assume that there exists such a μ . As in the proof of Claim 1, we can prove that $\mu^{-1}D_2^+$ is semi-ample. Let m be the fiber of the morphism defined by a sufficient multiple of $\mu^{-1}D_2^+$ containing l . There are components of $\text{Supp } \mu^{-1}(B^+|_{E^{+\nu}})$ intersecting $\mu^{-1}l$ since $B^+ \cdot l > 0$. Let b be the union of such components of $\text{Supp } \mu^{-1}(B^+|_{E^{+\nu}})$. By the assumption for μ , $\text{Supp } \mu^{-1}(B^+|_{E^{+\nu}}) \cap \mu^{-1}D_2^+ = \phi$. Hence b intersects the exceptional curve for μ and $\mu^{-1}l \cup b \subset m$. But $\mu^{-1}l$ contains Q'' and b intersects the exceptional curve for μ , so we obtain a contradiction by applying (1.6) for the birational contraction of m after extracting some log crepant curve from Q'' . \square

Assume that Q' is a singular point of $E^{+\nu}$. Since Q' is the unique singular point of $E^{+\nu}$ on D_2^+ by Theorem (2.4) (3), $(m^{-1}D_2^+)^2 > 0$, where m is the MRS along D_2^+ . We can make

m primitive and satisfying the assumption of Claim 2, a contradiction. So we see that Q' is a smooth point on $E^{+\nu}$. Furthermore we see that $(D_2^+)^2 = 1$ and $\text{Supp } B^+|_{E^{+\nu}}$ and D_2^+ are simply tangent at Q' . By the last property, E^+ contains no curve singularity near Q' . By the property of semi-good extraction, $K_{Z^+} + S^+ + E^+$ is LT near Q' . Hence by (1.11), Z^+ is also smooth at Q' . Since $K_{Z^+} + S^+ + E^+$ is LT at Q' and E^+ contains no curve singularity near Q' , S^+ is also smooth by (1.12). We finish the proof of (1).

Next we prove (2). It suffices to prove (2) on Z' since the flips before R are isomorphism near P . Let $t : \tilde{E}^{+\nu} \rightarrow E^{+\nu}$ be MLT of $K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}}$ at Q'' and F_1, \dots, F_n exceptional divisors of t , where F_1 intersects $t^{-1}D^*$. By (2.4), E' is obtained from $\tilde{E}^{+\nu}$ by contracting $t^{-1}D^*, F_1, \dots, F_{n-1}$. Then we claim the following:

Claim 3. $n = 1$ and $t^{-1}D^*$ is a (-1) -curve.

(2) follows from this claim and (1.11) immediately .

Proof of Claim 3. Let $h : E^{+\nu} \rightarrow \overline{E}^{+\nu}$ be the morphism defined by a sufficient multiple of D_2^+ . We see that $\overline{E}^{+\nu} \simeq \mathbb{P}^2$ since $h(D_2^+) \simeq \mathbb{P}^1$ is an ample Cartier divisor on $\overline{E}^{+\nu}$ and $h(D_2^+)^2 = 1$ by (1) and (2). Also we know that $h(D^*)$ is a line on $\overline{E}^{+\nu}$. We claim that $\cup_{i=1}^n F_i \cup t^{-1}D^*$ intersects strict transforms of h -exceptional curves only at points on F_n . In fact, if the strict transform F of a h -exceptional curve intersects $\cup_{i=1}^n F_i \cup t^{-1}D^*$ at points on F_i with $i \neq n$ (resp. $t^{-1}D^*$), then F is not contained in components of the boundary of $t^*(K_{Z^+} + S^+ + B^+ + E^+)|_{E^{+\nu}}$ by considering the degree of $t^*(K_{Z^+} + S^+ + B^+ + E^+)|_{F_i}$ and so F is a (-1) -curve on the MRS of Q'' . Hence also F does not intersect $\text{Supp } \text{Diff}_{E^{+\nu}}(S^+ + B^+)$ except F_i with $i \neq n$ (resp. $t^{-1}D^*$). By these, the strict transform of F on E' intersects D_1 only at P and does not intersect $\text{Supp } B'|_{E'}$. But this is a contradiction since $B^-|_{E^-}$ is nef on E' and numerically trivial only for D_1 . Hence we obtain the description as follows:

FIGURE (XXII).

Let $K_{\overline{E}^{+\nu}} + h(D^*) + h(D_2^+) + \overline{B} = h_*(K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}})$. and let $s : \tilde{E}^{+\nu} \rightarrow G$ the contraction of the strict transforms of h -exceptional curves. By Claim 1, they are not log crepant curves for $K_{\overline{E}^{+\nu}} + h(D^*) + h(D_2^+) + \overline{B}$. Hence we obtain the MLT for $K_{\overline{E}^{+\nu}} + h(D^*) + h(D_2^+) + \overline{B}$ possibly after contracting the images of some F_i 's on G . By the above, such curves intersect $\cup_{i=1}^n F_i \cup t^{-1}D^*$ at points only on F_n . Hence after contraction, the image of $\text{Supp } (\nu \circ t)^{-1}B^+|_{E^+}$ intersects the image of $\cup_{i=1}^n F_i \cup t^{-1}D^*$ at points only on the image of F_n . So if we contract the image of F_n , $s_*(t^*(K_{Z^+} + S^+ + B^+ + E^+|_{E^{+\nu}}))$ become non LT. Hence G is MLT for $K_{\overline{E}^{+\nu}} + h(D^*) + h(D_2^+) + \overline{B}$. On the other hand $\text{Supp } \overline{B}$ is tangent to $h(D_2^+)$ by (1). Since $K_{\overline{E}^{+\nu}} + h(D^*) + h(D_2^+) + \overline{B}$ is numerically trivial, $\text{Supp } \overline{B}$ is a smooth conic. Furthermore by the existence of Q'' , $\text{Supp } \overline{B}$ is also tangent to $h(D^*)$. Hence MLT for the image of Q'' is described as in (1.4). From these, we obtain Claim 3.

□

For (3), we consider $0 = (S_1 + dE).D_2 = \{(D_1 + D_2).D_2\}_E + d\{(D_1 + D_2).D_2\}_{S_1}$. By (1) and (2), $(D_1.D_2)_E = 1$, $(D_1.D_2)_{S_1} = 1$, $(D_2)_E^2 = 2$ so (3) follows. (4) is easily deduced from (1) and (2). □

Lemma (7.8) (Reducible case 2). *Consider the situation as in (7.6). Assume furthermore that $g(E) = L$.*

Then

- (1) $P := D_1 \cap D_2$ is smooth on S_1 and Z and is an ODP or a smooth point on E .
- (2) $d(F, S) = d+1$ and $a_l(F, S) = 0$ for the exceptional divisor F of the simple blow up along D_1 or D_2 and for any other log crepant divisor $F \neq E$ for $K_X + S + B$, $d(F, S) > d+1$.

Proof. When D_1 becomes a flipping ray, we use the same notation as in (7.6) (S' , E' ..., etc).

Claim. P is an ODP or a smooth point on E .

Proof. Let l be the fiber of $E \rightarrow L$ containing D_1 and l' any other component of l intersecting D_1 . Note that $B_1.l' > 0$. So if l' passes through P , after contraction of l' , $K_Z + S_1 + B_1 + E|_E$ has two reduced boundaries D_1 and D_2 and some other boundary contained in $\text{Supp } B_1$ at P , a contradiction. Hence l' does not pass through P . Hence contractions of components of l except D_1 does not change the singularity of P on E . Furthermore by the fact that any component of l except D_1 is not contained in LLC ($K_Z + S_1 + B_1 + E|_E$) and $K_Z + S_1 + B_1 + E|_E$ is numerically trivial for g , the property (3) of Lemma (7.6) does not change. so we may assume that D_1 is an irreducible fiber. By this, we easily show that P is an ODP or a smooth point on E . \square

Assume that P is smooth on E . Since $K_Z + S_1 + B_1 + E|_E$ has two reduced boundary D_1 and D_2 at P , E contains no singular curve near P . So P is smooth on Z by (1.11) and LT property of $K_Z + E$. Furthermore by (2.4), on S_1 , P is a smooth point or a singular point of index ≥ 3 . So P must be smooth on S_1 .

If P is an ODP on E , it suffices to prove that P is smooth on S_1 (by (1.11)). If P is singular on S_1 , P is a singular point of index ≥ 3 . Hence the index of $K_Z + S_1$ at P is greater than or equal to 3. Take the index 1 cover π of $K_Z + S_1$. Then the inverse image of E has more than or equal to 2 components since P is an ODP on E . But this contradicts LC property of $\pi^*(K_Z + S_1 + E)$ after taking a small \mathbb{Q} -factorization for $\pi^*(K_Z + S_1)$. Now we finish the proof of (1). (2) easily follows from (1) and the property of semi-good extraction (2) in (3.3). \square

Lemma (7.9) (Irreducible case). *Consider the case (c). Let $g : Z \rightarrow X$ be a primitive log crepant divisorial extraction for $K_X + S + B$ satisfying the following conditions:*

- (1) $D := g^{-1}S \cap E$ is irreducible and \mathbb{P}^1 .
- (2) $g^{-1}S$ and E are generically normal crossing along D ;
- (3) $K_Z + g^{-1}S + g^{-1}B + E|_D = K_D + \frac{1}{2}P_1 + \frac{1}{2}P_2 + P$;
- (4) $g^{-1}S$ is normal.

Then g is a good extraction (immediately by Lemma (5.2)). Run the BTM starting by g (we use the notation as in (2.5)). Then surgeries while BTM are described as in the following flow chart:

Flip the strict transform of $g^{-1}C$.
Consider the next extremal ray R' .

R' is a divisorial ray. DONE	R' is a flipping ray of type 2. DONE	R' is a flipping ray and $S' \cap \text{Supp } R' = \phi$.	R' is a final flipping ray. DONE
--------------------------------------	--	---	--

$\text{Supp } R' \cap \text{CLC}_0 = \phi$. Then $\text{Supp } R'$ is of type 3 or 4. DONE	$\text{Supp } R' \cap \text{CLC}_0 \neq \phi$. Then the connected compo- nent C' of $\text{Supp } R'$ contain- ing CLC_0 is of type (c)
---	--

$$\text{LLC} := \text{LLC} (K_{Z'} + S' + B' + E'|_{E'})$$

$$\text{CLC}_0 := \text{CLC}_0 (K_{Z'} + S' + B' + E'|_{E'})$$

Proof. By the similar reason to the proof of (7.6), first we must flip the strict transform of C . After the flips, we call Z' the image of Z and S' , B' and E' the strict transform of S , B and E . We can prove as Claim in the proof of (7.6) that any component of the flipped curve is not contained in S' . Let R' be the next extremal ray. Since $S' \cap E'$ is irreducible, we may assume that R' is a flipping ray such that $\text{Supp } R' \cap S' = \phi$, $E'.R' < 0$ and $B'.R' > 0$ (see the above flow chart). Furthermore we may assume that there exists the unique component M' of $\text{LLC} (K_{Z'} + S' + B' + E'|_{E'})$ and there exists a point $P' \in \text{CLC}_0 (K_{Z'} + S' + B' + E'|_{E'})$ on M . (see the flow chart). Let C' be the connected component of $\text{Supp } R'$ through P' . (Needless to say, the flip of another component of $\text{Supp } R'$ is of type 3 or 4.) We show that the flip of C' is of type (c). By the assumption of (5) in Set up (3.0) and Lemma, $\text{Supp } B'|_{E'} - M'$ does not intersect S' . Hence $\text{Supp } B'|_{E'} - M'$ is contained in $\text{Supp } R'$ since $S'.R' = 0$ and S' is positive for the flipped curve one before R' . Furthermore by the existence of P' and the connectedness of $\text{Supp } B'|_{E'}$, $P' \in \text{Supp } B'|_{E'} - M'$, i.e., $\text{Supp } B'|_{E'} - M' \subset C'$. We show that $\text{Supp } B'|_{E'} - M' = C'$. Assume that there is a component C'' of C' which is not contained in $\text{Supp } B'|_{E'} - M'$ and contains P' . After contracting $\text{Supp } B'|_{E'} - M'$, M' becomes ample since $B'.R' > 0$ and B' is positive for the flipped curve one before R' . So by (1.9), $P' \notin C''$, a contradiction. Hence $\text{Supp } B'|_{E'} - M' = C'$. Replacing C by C' , S by E' , B by B' and L by M' , we can easily check the condition of case (c). (Note that $K_{Z'} + E' + S'$ is LT since LT property of $K_Z + S + E$ is preserved by the flip of the strict transform of C .) \square

Lemma (7.10). *Consider the case (c). Let $g : Z \rightarrow X$ be a primitive log crepant divisorial extraction for $K_X + S + B$ and E the exceptional divisor. Let D_1 be the component of $g^{-1}S \cap E$ which intersects $g^{-1}C$. Let C' be a component of C . Assume that $g^{-1}S \cap E$ is generically normal crossing along D_1 and $g^{-1}C' \cap D_1$ is a smooth point on $g^{-1}S$. Then $g^{-1}C'$ is a (-1) -curve or a (-2) -curve on the MRS of $g^{-1}S$. More precisely, $g^{-1}C'$ is a (-2) -curve on the MRS of $g^{-1}S$ if and only if $g^{-1}S$ is nonsingular on $g^{-1}C'$ and there is no component of $\text{Diff}_{g^{-1}S}(0)$ and $g^{-1}B|_{g^{-1}S} - g^{-1}C'$ intersecting $g^{-1}C'$.*

Assume in addition that $K_Z + g^{-1}S + g^{-1}B + E|_{g^{-1}S}$ is PLT at $g^{-1}C' \cap D_1$ and $g^{-1}C'$ is a (-1) -curve on the MRS of $g^{-1}S$. Then one of the following holds:

- (1) there is only one singularity $x \neq g^{-1}C' \cap D_1$ on $g^{-1}C'$ resolved by a (-3) -curve. There is no component of $\text{Diff}_{g^{-1}S}(0)$ and $g^{-1}B|_{g^{-1}S} - g^{-1}C'$ intersecting $g^{-1}C'$;
- (2) there is only one singularity $x \neq g^{-1}C' \cap D_1$ which is canonical of type A_m (we allow $m = 0$). There is no component of $g^{-1}B|_{g^{-1}S} - g^{-1}C'$ intersecting $g^{-1}C'$ and there is only one component of $\text{Diff}_{g^{-1}S}(0)$ (an ODP curve singularity of Z) or $\text{Supp } g^{-1}B|_{g^{-1}S}$ intersecting $g^{-1}C'$ which pass through x .

proof. By $(K_Z + g^{-1}S + g^{-1}B + E).g^{-1}C' = 0$, we have

$$(K_{g^{-1}S} + \text{Diff}_{g^{-1}S}(0) + g^{-1}B|_{g^{-1}S} + D_1).g^{-1}C' = 0.$$

Note that $g^{-1}C'$ is contained in $g^{-1}B|_{g^{-1}S}$ with coefficient $\frac{1}{2}$. Also note that P' is smooth on $g^{-1}S$. By these, we obtain the equality $K_{g^{-1}S}.g^{-1}C' + \frac{1}{2}(g^{-1}C')^2 + 1 + \alpha = 0$, where α is a non-negative rational number. On the other hand, we have the equality $K_{g^{-1}S}.g^{-1}C' + (g^{-1}C')^2 = -2 + \beta$, where β is a nonnegative rational number. By these, we obtain $K_{g^{-1}S}.g^{-1}C' = -2\alpha - \beta$ and $(g^{-1}C')^2 = -2 + 2\alpha + 2\beta$. Hence we obtained the first part of the lemma.

Assume in addition that $K_Z + g^{-1}S + g^{-1}B + E|_{g^{-1}S}$ is PLT at $g^{-1}C' \cap D_1$ and $g^{-1}C'$ is a (-1) -curve on the MRS of $g^{-1}S$. Note that after contraction of C , $K_X + S + B|_S$ becomes LC of index 2 at the image of C . $g^{-1}S$ becomes its MLT after contracting $g^{-1}C$. By the description of the MLT of a LC point of index 2, the image of $g^{-1}C$ by its contraction is an ODP or is smooth and there is one of an ODP singular curve of Z or $\text{Supp } g^{-1}B|_{g^{-1}S}$ containing it. If the former case occurs, we have (1). If the latter case occurs, we have (2). \square

Lemma (7.11) (Complement). *Let $(X, K_X + S)$ is a PLT d -fold, where $d \geq 3$ and S is a \mathbb{Q} -Cartier irreducible divisor. Let $f : X \rightarrow Y$ be a projective contraction such that $-(K_X + S)$ is f -nef and f -big. Assume that the index of $K_X + S$ is n except finite points. Then if there is a KLT index n complement \overline{B} for $K_X + S|_S$, it lifts to a PLT index n complement for $K_X + S$, i.e., there is a PLT index n complement B for $K_X + S$ such that $B|_S = \overline{B}$.*

Proof. Let X_0 be the maximum locus where the index of $K_X + S$ is n . and $S_0 := S|_{X_0}$. Consider the local cohomology exact sequence:

$$H^1(X, \mathcal{O}_X(-n(K_X + S) - S)) \rightarrow H^1(X_0, \mathcal{O}_{X_0}(-n(K_X + S) - S)) \rightarrow H^2_{\{X - X_0\}}(X, \mathcal{O}_X(-n(K_X + S) - S)). \blacksquare$$

The first term vanishes by Kawamata-Viehweg vanishing theorem. The last term vanishes by Cohen-Macaulay property of $\mathcal{O}_X(-n(K_X + S) - S)$ and $d \geq 3$. Hence we obtain $H^1(X_0, \mathcal{O}_{X_0}(-n(K_X + S) - S)) = 0$. Next consider the exact sequence:

$$0 \rightarrow \mathcal{O}_{X_0}(-n(K_X + S) - S) \rightarrow \mathcal{O}_{X_0}(-n(K_X + S)) \rightarrow \mathcal{O}_{S_0}(-n(K_X + S)) \rightarrow 0.$$

Note that the first part of the sequence is exact since $-n(K_X + S)$ is Cartier on X_0 . By the above vanishing, we obtain the surjection

$$H^0(X, \mathcal{O}_X(-n(K_X + S))) \rightarrow H^0(S, \mathcal{O}_S(-n(K_X + S))).$$

Note that

$$H^0(X_0, \mathcal{O}_{X_0}(-n(K_X + S))) \simeq H^0(X, \mathcal{O}_X(-n(K_X + S)))$$

and

$$H^0(S_0, \mathcal{O}_{S_0}(-n(K_X + S))) \simeq H^0(S, \mathcal{O}_S(-n(K_X + S)))$$

by reflexivity. Hence there is a member $B' \in |-n(K_X + S)|$ such that $B'|_S = n\overline{B}$. Let $B := \frac{1}{n}B'$. Then $K_X + S + B|_S = K_S + \text{Diff}_S(0) + \overline{B}$. By the assumption, the right side is KLT. Hence by the inversion of adjunction, $K_X + S + B$ is PLT. This completes the proof. \square

Lemma (7.12) (Case $\delta = 0$). *Consider the case (c). Assume that $\delta = 0$. Then $\text{Supp } B$ and S are generically simply tangent along L . In particular $1 < d \leq 2$.*

Proof. If $\text{Supp } B$ and S are not simply tangent along L , δ must be positive by the proof of Claim 1 in the proof of (6.0). For the latter half, note that there is a log crepant divisor F over L for $K_X + S + B$ with $d(F, S) = 2$. \square

Conclusion of case (c, δ).

By running $g''^*(K_X + S + B) - \epsilon(g''^{-1}S + dE)$ -MMP and taking the LC model as in (7.0), we obtain the primitive extraction of E . We can check the above procedure is isomorphism near P' in case (1) of (7.0)(resp. near P_1 and P_2 in case (2) of (7.0)). Hence in case (2), we can check that g is good or semi-good.

First we exclude some cases.

Proposition (7.13). *If the case (1) of (7.0) occurs, P' is not smooth on Z .*

Proof. Assume that P' is smooth on Z . Note first that all the components of $g^{-1}C$ pass through P' . Let C' be a component of C . If there is another component of C , then $g^{-1}C'$ is (-1) -curve on the MRS of $g^{-1}S$ by (7.10). The same thing holds for another component of $g^{-1}C$. But this is a contradiction since $g^{-1}C$ is contractible and P' is smooth on $g^{-1}S$. Hence $g^{-1}C$ is irreducible. In particular, near P' , $K_Z + g^{-1}S + g^{-1}B + E|_{g^{-1}S} = K_{g^{-1}S} + \frac{1}{2}g^{-1}C + D_1$, where $g^{-1}C$ and D_1 are normal crossing at P' . But this contradicts the assumption of (1) in (7.0). So we exclude this case. \square

Proposition (7.14). *If the case (1) of (7.0) occurs and $(P' \in g^{-1}S \cup E \subset Z) \simeq (o \in (xy = 0) \subset \frac{1}{2}(1, 0, 1))$, then one of the following holds:*

- (1) *after the flip of the strict transform of C , D_1 become a final flipping ray or the strict transform of E becomes a exceptional divisor of a divisorial contraction. In particular the flip of f exists by (2.5);*
- (2) *there is a index 2 PLT complement near C . In particular the flip of f exists by (2.2).*

Proof.

Claim. *P' is a smooth point on $g^{-1}S$ and an ODP on E .*

Proof. Assume the contrary, i.e., P' is an ODP on $g^{-1}S$. If $g^{-1}C$ is irreducible, $K_Z + g^{-1}S + g^{-1}B + E|_{g^{-1}S} = K_{g^{-1}S} + D_1 + \frac{1}{2}g^{-1}C$ near P' . But then $K_Z + g^{-1}S + g^{-1}B + E|_{g^{-1}S}$ must be PLT at P' , a contradiction to the assumption of (1) in (7.0). Assume that $g^{-1}C$ is reducible, i.e., it has two components $g^{-1}C'$ and $g^{-1}C''$. By (7.10), $g^{-1}C'$ (resp. $g^{-1}C''$) is a (-1) -curve or (-2) -curve on the MRS of $g^{-1}S$.

If $D_1 = g^{-1}L$, $g^{-1}S \simeq S$ and hence Q is an ODP on S . On the other hand, $K_S.C' < 0$ (resp. $K_S.C'' < 0$) by $(K_X + S).C' < 0$ (resp. $(K_X + S).C'' < 0$). So C' (resp. C'') is a (-1) -curve on the MRS of singularities of $g^{-1}S$. But on it we have the non-contractible chain consisting of the strict transform of C' , (-2) -curve resolving Q and the strict transform of C'' , a contradiction.

If $D_1 \neq g^{-1}L$, D_1 is not a (-1) -curve on the MRS of singularities of $g^{-1}S$. For otherwise we have the non-contractible tree consisting of the strict transform of $g^{-1}C'$, (-2) -curve resolving P' and the strict transform of $g^{-1}C''$ and the strict transform of D_1 , a contradiction. In particular, $K_{g^{-1}S}.D_1 \geq 0$ and hence $(K_Z + g^{-1}S).D_1 \geq 0$. So we can write $g^*(K_X + S) = K_Z + g^{-1}S + aE$, where a is a nonnegative rational number. Intersecting this with $g^{-1}C'$ (resp. $g^{-1}C''$), we obtain $K_{g^{-1}S}.g^{-1}C' < 0$ (resp. $K_{g^{-1}S}.g^{-1}C'' < 0$). Hence $g^{-1}C'$ (resp. $g^{-1}C''$) is a (-1) -curve on the MRS of singularities of $g^{-1}S$. But this is a contradiction by the same reason as in case $D_1 = g^{-1}L$. \square

By this Claim, there is an ODP curve singularity D_0 on $g^{-1}S$ through P' . Hence $g^{-1}C$ must be irreducible by (1.4). By the argument of (7.10), $g^{-1}C$ is (-1) -curve on the MRS of $g^{-1}S$ and there is no component of $\text{Diff}_{g^{-1}S}(0)$ intersecting $g^{-1}C$ except D_0 and there is at most one singular point on $g^{-1}C$ which is of type A_m . (See FIGURE (XXIII).)

Assume that $g^{-1}S \cap E$ is reducible. We prove that (1) holds in this case.

We run the BTM starting from g . We use the notation as in (2.5). We see as in (7.6) that C_1 is the strict transform of C . We prove that C_1^+ is not contained in S_2 . Note that $h_1(C_1)$ is a smooth point of $h_1(S_1)$ by the above description of $g^{-1}C$. So $K_{h_1(S_1)} + \frac{1}{2}h_1(D_0)$ is canonical. On the other hand $K_{S_2} + \frac{1}{2}D'_0 + B_2|_{S_2}$ is g_2 -ample, where D'_0 is the strict transform of D_0 . Hence no component of C_1^+ is contained in S_2 .

By this, $S_1 \dashrightarrow S_2$ is a morphism contracting C_1 . We denote the image point of C_1 by z . We can easily show that $z \notin \text{Supp } B_2$. By this we have $B_2.D_1 = 0$, where D_1 is the strict transform of D_1 (by abuse of notation). We can deduce from this that D_1 is contained in $\text{Supp } R_2$. By the numerical properties (1) and (2), we have $S_2.R_2 < 0$ and $E_2.R_2 > 0$, or $S_2.R_2 > 0$ and $E_2.R_2 < 0$. If the latter case occurs, we are done. Hence it suffices to exclude the former case. Assume that the former case occurs. As we proved above, z is smooth on S_2 . Let $\mu : \tilde{S}_2 \rightarrow S_2$ be the MRS of the point $D_1 \cap D_2$. Let $E_2|_{S_2} = D_1 + \alpha D_2$ and $\mu^*(D_1 + \alpha D_2) = \mu^{-1}D_1 + \alpha\mu^{-1}D_2 + \sum_{i=1}^n f_i F_i$, where F_i is exceptional curves on $D_1 \cap D_2$. Note that $E_2.D_1 > 0$. So we obtain $(\mu^{-1}D_1)^2 + \alpha > 0$ if $D_1 \cap D_2$ is smooth on $g^{-1}S$ or $(\mu^{-1}D_1)^2 + f_1 > 0$ if $D_1 \cap D_2$ is singular on $g^{-1}S$. In any case $(\mu^{-1}D_1)^2 > 0$ by $\alpha \leq 1$ and $f_i \leq 1$, a contradiction to the contractibility of $\mu^{-1}D_1$. So we are done.

Next we assume that $g^{-1}S \cap E$ is irreducible. We have two cases, one is that $g(E) = L$ and another is that $g(E) = Q$. In these cases, we show (2) holds.

If $g(E) = L$, $g^{-1}S \simeq S$. Let $K_X + S|_S = K_S + \frac{1}{2}g(D_0) + \frac{n-1}{n}L$, where n is a natural number. Since $g^*(K_X + S).g^{-1}C < 0$, we have $-1 + \frac{1}{2} + \frac{n-1}{n} < 0$. Hence $n = 1$. Near Q , let $\tilde{X} := \mathbf{Spec}(\mathcal{O}_X \oplus \mathcal{O}_X(-(K_X + S)))$. The natural morphism $\tilde{X} \rightarrow X$ is a double cover of X ramified along $g(D_0)$. By $n = 1$ and (1.11), \tilde{X} is smooth. Hence the index of $K_X + S$ is 2 at Q and hence on C . Take a general member \tilde{B} of $-2(K_X + S)$. Then near C , $K_X + S + \frac{1}{2}\tilde{B}$ is PLT and of index 2. So the flip exists by (2.2).

Assume that $g(E) = Q$. Let $g^*(K_X + S)|_{g^{-1}S} = K_{g^{-1}S} + \frac{1}{2}D_0 + \frac{n-1}{n}g^{-1}L + a'D_1$, where n is a natural number and a' is a positive natural number. Intersecting this with $g^{-1}C$, we obtain $(g^*(K_X + S)).g^{-1}C = a' - \frac{1}{2} < 0$ and so $a' < \frac{1}{2}$. Let $\mu : (g^{-1}S)^\mu \rightarrow g^{-1}S$ be the MRS of $g^{-1}S$ and F_1, \dots, F_m exceptional curves over $D_1 \cap g^{-1}L$. Note that $D_1 \cap g^{-1}L$ is the unique possibility of singularity of $g^{-1}S$ on D_1 . Write

$$\begin{aligned} \mu^*(K_{g^{-1}S} + \frac{n-1}{n}g^{-1}L + \frac{1}{2}D_0 + a'D_1) = \\ K_{g^{-1}S}^\mu + \frac{n-1}{n}\mu^{-1}g^{-1}L + \frac{1}{2}\mu^{-1}D_0 + a'\mu^{-1}D_1 + \sum_{i=1}^m f_i F_i. \end{aligned}$$

Intersecting this successively with $\mu^{-1}D_1, F_1, \dots$ and F_m noting $a' < \frac{1}{2}$, we know that $\mu^{-1}D_1, F_1, \dots$ and F_m are (-2) -curves, i.e., Q is a canonical singularity of type A_{m+1} on S and $n = 1$. In this case, we can easily construct a index 2 KLT complement for $K_X + S|_S$ and hence by (7.11), it lifts to a index 2 PLT complement of $K_X + S$. This completes the proof of this proposition. \square

Hence we need to treat the following 6 cases:

- (i) Treatment of case (1) in (7.0)- $\delta > 0$ and P' is a $\frac{1}{3}(2, 2, 1)$ -singularity

- (ii) Treatment of (2) in Set up (7.0)- $\delta > 0$ and g is good
- (iii) Treatment of (2) in Set up (7.0)- $\delta > 0$ and g is semi-good
- (iv) Treatment of (1) in Set up (7.0)- $\delta = 0$ and P' is a $\frac{1}{3}(2, 2, 1)$ -singularity
- (v) Treatment of (2) in Set up (7.0)- $\delta = 0$ and g is good
- (vi) Treatment of (2) in Set up (7.0)- $\delta = 0$ and g is semi-good

Our conclusion is explained in the following flow chart:

$$\delta > 0$$

Choose a good primitive log crepant extraction g such that $d(E, S) \leq 1$ and run BTM starting by g

We fall into Case (1) of (7.0) and P' satisfies (3) of (7.1). DONE

We fall into Case (1) of (7.0) and P' satisfies (2) of (7.1)

We fall into (2) of (7.0).

↓

Let \tilde{g} be the primitive extraction of \tilde{E} . Then we fall into Case (2) of (7.0). Run the BTM starting by \tilde{g} .

$g^{-1}S \cap E$ is reducible. Let \tilde{g} be the primitive extraction of \tilde{E} . Then we fall into Case (2) of (7.0). Run the BTM starting by \tilde{g} .

$g^{-1}S \cap E$ is irreducible. Apply (7.9). We may assume that \tilde{R} is of type (c). Then this is of type (c, $\delta' < \delta$)

⇓

(B) $\tilde{g}^{-1}S \cap \tilde{E}$ is reducible. Apply (7.6). Then (1)~(4) of (7.6) do not hold. DONE

(A) $\tilde{g}^{-1}S \cap \tilde{E}$ is irreducible. Apply (7.9). We may assume that \tilde{R} is of type (c).

(D) $\tilde{g}^{-1}S \cap \tilde{E}$ is reducible. Apply (7.6). Then (1)~(4) of (7.6) do not hold. DONE

(C) $\tilde{g}^{-1}S \cap \tilde{E}$ is irreducible. Apply (7.9). We may assume that \tilde{R} is of type (c, δ').


Reduction to case (c, $\delta' < \delta$)

(A-2) \tilde{R} is of type (2)

(A-1) \tilde{R} is of the type


(C-1) $g(E) = Q$

(C-2) $g(E) = L$. Then $\delta' = 0$




↓

↓



↓



↓

↓



Case (c) and $\delta = 0$

Choose a good primitive log crepant extraction g such that $d(E, S) = d$ and run BTM starting by g

We fall into Case (1) of (7.0) and P' satisfies (3) of (7.1). DONE	We fall into Case (1) of (7.0). Then P' satisfies (2) of (7.1)	We fall into Case (2) of (7.0).
--	--	---------------------------------

↓

Let \tilde{g} be the primitive extraction of \tilde{E} . Then we fall into Case (2) of (7.0). Run the BTM starting by \tilde{g} .

$g^{-1}S \cap E$ is reducible.	$g^{-1}S \cap E$ is irreducible. Then there is an exceptional complement of index 2 for $K_X + S$ and hence C become of type 4. DONE
--------------------------------	--

(B) $\tilde{g}^{-1}S \cap \tilde{E}$ is reducible. Apply (7.6). Then (1)~(4) of	(A) $\tilde{g}^{-1}S \cap \tilde{E}$ is irreducible. Apply (7.9). We may assume	$g(E) = Q$. Then there is a PLT complement of	$g(E) = L$. Then there is a PLT complement of
---	---	--	--

(A-1) \tilde{R} is of the type in (7.14). DONE	(A-2) \tilde{R} is of type (2) in (7.0) and δ for \tilde{R} is 0
---	---

↓

Reduction to case $(c, \delta' = 0)$ and the case of (2) in (7.0)
--

Treatment of (1) in Set up (7.0)- P' is a $\frac{1}{3}(2, 2, 1)$ -singularity. We treat the case $\delta > 0$ and $\delta = 0$ simultaneously. Recall there is a exceptional divisor F' over Z as in Theorem (7.1) (2). We change the notation. We write \tilde{E} instead of F' . Note that $d(\tilde{E}, S) > 1$. Running $(g \circ h)^*(K_X + S + B) - \epsilon((g \circ h)^{-1}S + \tilde{d}\tilde{E})$ -MMP over X , we obtain a primitive extraction of \tilde{E} , which we call $\tilde{g} : \tilde{Z} \rightarrow X$. Let $\tilde{D} := \tilde{g}^{-1}S \cap \tilde{E}$ if $\tilde{g}^{-1}S \cap \tilde{E}$ is irreducible or $\tilde{D}_1 \cup \tilde{D}_2 := \tilde{g}^{-1}S \cap \tilde{E}$ if $\tilde{g}^{-1}S \cap \tilde{E}$ is reducible, where only \tilde{D}_1 intersects $\tilde{g}^{-1}C$. (By (5.2), there are such two possibilities.) Note that outside E , the MMP is isomorphism. Hence if $\tilde{g}^{-1}S \cap \tilde{E}$ is irreducible (resp. reducible), \tilde{D} (resp. \tilde{D}_1) is the strict transform of $\tilde{E} \cap (g \circ h)^{-1}S$. Furthermore by the description of (7.1), (7.0) (2) holds for \tilde{g} . By Theorem (7.1) (2), the assumptions of (7.9) (resp. (7.6)) holds. Hence \tilde{g} is good (resp. semi-good).

(A). If \tilde{g} is good, by applying (7.9), we have only consider the case (c) appears while BTM. We use the notation as in the proof of (7.9) (replacing $g, E, D_1, C', M, M', \dots$ etc by $\tilde{g}, \tilde{E}, \tilde{D}_1, \tilde{C}', \tilde{M}, \tilde{M}', \dots$ etc). We claim that \tilde{C}' satisfies the assumption of Proposition (7.14) (and there exists the flip of \tilde{C}' , hence of C) or \tilde{C}' is of type (2) in (7.0), $\delta = 0$. Do the same procedure as in (7.0) for \tilde{C}' . Recall $K_{\tilde{Z}} + \tilde{g}^{-1}S + \tilde{g}^{-1}B + \tilde{E}|_{\tilde{E}} = K_{\tilde{E}} + \tilde{M}_1 + \tilde{M} + \frac{1}{2}\tilde{M}_3 + \frac{1}{2}\tilde{M}_4$, where $\tilde{M}_1 := \tilde{g}^{-1}S|_{\tilde{E}}$, $\tilde{M}_3 := \text{Supp } \tilde{g}^{-1}B|_{\tilde{E}} - \tilde{M}$ and \tilde{M}_4 is a generically ODP curve of \tilde{Z} . Since \tilde{g} is good, $\tilde{M}_1 \cap \tilde{M}_3 \neq \tilde{M}_1 \cap \tilde{M}_4$. See FIGURE (XXIV).

(A-1). If we fall into case (1) of (7.0), the flip of \tilde{C}' satisfies the assumption of Proposition (7.14) by the existence of the generically ODP curve along \tilde{M}_4 .

(A-2). If we fall into case (2) of (7.0), assume that there exists a log crepant divisor F for $K_{\tilde{Z}'} + \tilde{S}' + \tilde{B}' + \tilde{E}'$ whose center contains \tilde{P}' (i.e., is equal to \tilde{P}' or \tilde{M}') and $d(\tilde{E}', F) \leq 1$ (to get a contradiction). Let $l : W \rightarrow \tilde{Z}'$ be a (primitive) log crepant extraction of F . Here we can take F the divisor corresponding to E in (7.0). Since the flips before \tilde{C}' do not touch with \tilde{M} , the center of F on \tilde{Z} is \tilde{P} or \tilde{M} and we can consider l is defined over \tilde{Z} . Run the $(\tilde{g} \circ l)^*(K_X + S + B) - \epsilon((\tilde{g} \circ l)^{-1}S + d(F, S)F)$ -MMP over X . We show possibly after the flip of $l^{-1}\tilde{M}$ that $l^{-1}\tilde{E}$ is contracted to a curve.

Let A be the first extremal ray. By the assumption that we fall into case (2) of (7.0), $l^{-1}\tilde{M}_3 \cap F|_{l^{-1}\tilde{E}} \neq l^{-1}\tilde{M}_4 \cap F|_{l^{-1}\tilde{E}}$. Hence $(\tilde{g} \circ l)^{-1}B.l^{-1}\tilde{M}_4 = 0$. Since for a l -exceptional curve, $(\tilde{g} \circ l)^{-1}B$ is positive, we know that $l^{-1}\tilde{B}.A \leq 0$.

If $(\tilde{g} \circ l)^{-1}B.A = 0$, $l^{-1}\tilde{M}_4 \subset \text{Supp } A$ since the extremal ray for F is positive for $(\tilde{g} \circ l)^{-1}B$ and we play 2 ray game. By (1.6), $l^{-1}\tilde{M}_4$ cannot be birationally contracted. Hence A is a divisorial ray and $l^{-1}\tilde{E}$ is contracted to a curve.

If $(\tilde{g} \circ l)^{-1}B.A < 0$, A is a flipping ray and $\text{Supp } A \subset (\tilde{g} \circ l)^{-1}B$. But $l^{-1}\tilde{M}_3$ cannot be birationally contracted by (1.6) since $l^{-1}\tilde{M}_3$ intersects $F|_{l^{-1}\tilde{E}}$ and $(\tilde{g} \circ l)^{-1}S|_{l^{-1}\tilde{E}}$ at two points. So $\text{Supp } A = l^{-1}\tilde{M}_2$ and $l(F) = \tilde{P}$. Since A is $(\tilde{g} \circ l)^*(K_X + S + B) - \epsilon((\tilde{g} \circ l)^{-1}S + d(F, S)F)$ -extremal ray, $l^{-1}\tilde{E}.A < 0$ and $F.A > 0$. So this is a flip of type 1 and the flip exists by (2.0). Let A' be the next extremal ray. We express all the strict transforms with superscript $+$. Since $(\tilde{g} \circ l)^{-1}B^+.l^{-1}\tilde{M}_4^+ = 0$ and for the flipped curve, $(\tilde{g} \circ l)^{-1}B^+$ is positive, we know that $(\tilde{g} \circ l)^{-1}B^+.A' \leq 0$. But we see that $(\tilde{g} \circ l)^{-1}B^+.A' = 0$ and $l^{-1}\tilde{E}^+$ is contracted to a curve. Hence we obtain what we want.

Furthermore by the above argument, we also see that when \tilde{E} is contracted to a curve, $l^{-1}\tilde{E}$ and F is generically normal crossing along the irreducible component of $l^{-1}\tilde{E} \cap F$ intersecting $l^{-1}\tilde{C}'$. The same thing also holds for $l^{-1}\tilde{E}$ and $(\tilde{g} \circ l)^{-1}S$. Since $l^{-1}\tilde{M}_4^+.(l^{-1}\tilde{E})^{-1}B^+ = 0$ and it intersects F and $(\tilde{g} \circ l)^{-1}S$ at distinct two points, $l^{-1}\tilde{M}_4^+$ is an irreducible fiber by (1.6). Furthermore it intersects F and $(\tilde{g} \circ l)^{-1}S$ at smooth points normally. Hence

$(\tilde{g} \circ l)^{-1} S.l^{-1}\tilde{M}_4 = F.l^{-1}\tilde{M}_4 = 1$. Since $l^{-1}\tilde{M}_4$ does not intersect singular curve of \tilde{Z} and neither does a general fiber, $n := -l^{-1}\tilde{E}.l^{-1}\tilde{M}_4$ is a positive integer. Since $((\tilde{g} \circ l)^{-1}S + d(\tilde{E}, S)l^{-1}\tilde{E} + d(F, S)F).l^{-1}\tilde{M}_4 = 0$ and $d(F, S) = d(F, \tilde{E})d(\tilde{E}, S)$, we obtain the equality $1 - nd(\tilde{E}, S) + d(F, \tilde{E})d(\tilde{E}, S) = 0$. From now on, we treat separately the case $\delta > 0$ and the case $\delta = 0$.

First we treat the case $\delta > 0$.

If $n \geq 2$ in the above equality, then

$$d(\tilde{E}, S) \leq \frac{1}{2}(d(\tilde{E}, S)d(F, \tilde{E}) + 1) \leq \frac{1}{2}(d(\tilde{E}, S) + 1)$$

by the assumption $d(F, \tilde{E}) \leq 1$. So $d(\tilde{E}, S) \leq 1$, a contradiction to the choice of \tilde{E} .

If $n = 1$, $d(F, S) = d(\tilde{E}, S) - 1 \leq \frac{4}{3} - 1 < 1$. Hence after contracting \tilde{E} , we fall into case (2) of (7.0) with $\delta > 0$ since after contracting $l^{-1}\tilde{E}$, F and $(\tilde{g} \circ l)^{-1}S$ cross normally along the image of $l^{-1}\tilde{E}$ by $l^{-1}\tilde{E}.l^{-1}\tilde{M}_4 = -1$, a contradiction by Claim in (7.0).

Next we treat the case $\delta = 0$.

If $n \geq 2$ in the above equality, we get a contradiction by the same way as the case $\delta > 0$.

If $n = 1$, then $d(F, S) = d(\tilde{E}, S) - 1 \leq 2 - 1 = 1$, a contradiction to the assumption $\delta = 0$.

Now we finish the treatment of the case \tilde{g} is good.

(B). If \tilde{g} is semi-good, we can apply (7.6) and (7.7) for \tilde{g} . Note that $d(\tilde{E}, S) > 1$ so if (4) of Lemma (7.7) holds, then for all log crepant divisors their multiplicities in S is greater than 1 (resp. d) in case $\delta > 0$ (resp. $\delta = 0$), a contradiction. Hence in this case, we can construct the flip of f as explained in the flow chart. \square

Treatment of (2) in Set up (7.0), $\delta > 0$ and g is good.

In this case, we can apply (7.9). We have only to consider the case (c) appears while BTM. We use the notation as in the proof of (7.9). Then we can check the flip of C' is of type $(c, \delta' < \delta)$. So we reduce the existence of the flip of C to the flip of type $(c, \delta' < \delta)$. \square

Treatment of (2) in Set up (7.0), $\delta > 0$ and g is semi-good. (See FIGURE (XXV).)

Let $h : W \rightarrow Z$ be the simple blow up of D_1 and \tilde{E} the exceptional divisor of h . Running $(g \circ h)^*(K_X + S + B) - \epsilon((g \circ h)^{-1}S + \tilde{d}\tilde{E})$ -MMP over X , we obtain an primitive extraction of \tilde{E} , which we call $\tilde{g} : \tilde{Z} \rightarrow X$. Let $\tilde{D} := \tilde{g}^{-1}S \cap \tilde{E}$ if $\tilde{g}^{-1}S \cap \tilde{E}$ is irreducible or $\tilde{D}_1 \cup \tilde{D}_2 := \tilde{g}^{-1}S \cap \tilde{E}$ if $\tilde{g}^{-1}S \cap \tilde{E}$ is reducible, where only \tilde{D}_1 intersects $\tilde{g}^{-1}C$. (By (5.2), there are such two possibilities.) Since $h|_{(g \circ h)^{-1}S}$ is isomorphism over D_1 and the MMP is isomorphism outside E , the conditions (2) and (3) of a semi-good extraction in (3.3) are preserved (replacing E by \tilde{E}). Hence for \tilde{g} , the assumptions of (7.9) or (7.6) are satisfied.

(C). If $\tilde{g}^{-1}S \cap \tilde{E}$ is irreducible, we can apply (7.9) for \tilde{g} and we may assume that a flip of type (c) appears while BTM. We use the notation as in the lemma (replacing g, E, D_1, C', M', \dots etc by $\tilde{g}, \tilde{E}, \tilde{D}_1, \tilde{C}', \tilde{M}', \dots$ etc). We prove that δ for the flip of \tilde{C}' is 0.

First consider the case $g(E) = Q$.

(C-1-1). Assume that \tilde{Z}' is nonsingular along \tilde{M}' . Then the following claim holds:

Claim. *Assume that \tilde{Z}' is nonsingular generically along \tilde{M}' . Then there is a log crepant divisor F for $K_X + S + B$ over $\tilde{Q}' := \tilde{M}' \cap (\tilde{S}'|_{\tilde{E}'})$ whose multiplicity in $\tilde{S}' + \tilde{E}'$ is less than or equal to 1.*

Proof. Note that $S_1 \rightarrow \tilde{S}'$ is the contraction of D_2 . By (1) and (2) of Lemma (7.7), S_1 is smooth along D_2 and by (3) of the same Lemma, $d(E, S) = \frac{3}{-(D_2)_{S_1}^2 - 1} \leq 1$. Let $n := -(D_2)_{S_1}^2$.

Then by these, \tilde{Q} is $\frac{1}{n}(1, 1)$ singularity on \tilde{S} and $n \geq 4$. It is easy to see that the index of $K_{\tilde{Z}} + \tilde{S}$ at \tilde{Q} is n and the index 1 cover for this is smooth by (1.11). Hence locally analytically $(\tilde{Q} \in \tilde{S} \cup \tilde{E} \subset \tilde{Z}) \simeq (o \in (xy = 0) \subset \frac{1}{n}(k, 1, 1))$, where $1 \leq k < n$ is a natural number. The weighted blow up with the weight $\frac{1}{n}(k, 1, 1)$ extracts a log crepant divisor for $K_{\tilde{Z}} + \tilde{S} + \tilde{E} + \tilde{B}$ since the restriction of the weighted blow up to \tilde{S} coincides with the MRS and it must be log crepant for $K_{\tilde{Z}} + \tilde{S} + \tilde{E} + \tilde{B}|_{\tilde{S}}$ by $n \geq 4$. The multiplicity of the exceptional divisor in $\tilde{S} + \tilde{E}$ is equal to $\frac{k+1}{n}$ and this is less than or equal to 1. So we are done. \square

By this claim we complete the reduction if \tilde{Z}' is nonsingular generically along \tilde{M}' . In fact, let $d(F, \tilde{S}') = a$ and $d(F, \tilde{E}') = b$. Since $a + b \leq 1$, $a < 1$ and $b < 1$,

$$d(F, S) = d(F, \tilde{S}' + d(\tilde{E}', S)\tilde{E}') = a + d(\tilde{E}', S)b = a + (d(E, S) + 1)b < 1 + d(E, S).$$

Hence by (4) of Lemma (7.7) for g , it must be E . Furthermore we know by this that on \tilde{Z} , the center of the exceptional divisor of the simple blow up of D_2 is also \tilde{Q}' . Hence for any log crepant divisor for $K_{\tilde{Z}} + \tilde{S}' + \tilde{B}' + \tilde{E}'$ whose center contains \tilde{P}' , its multiplicity in $\tilde{S}' + (d(E, S) + 1)\tilde{E}'$ is greater than $d(E, S) + 1$, which in turn show that its multiplicity in \tilde{E}' is greater than 1, which we want.

(C-1-2). If \tilde{Z}' is singular along \tilde{M}' , then there exists a log crepant divisor for $K_{\tilde{Z}'} + \tilde{S}' + \tilde{E}' + \tilde{B}'$ over \tilde{M}' such that its multiplicity in \tilde{E}' is less than 1 (take a general hyperplane section at a general point of \tilde{M} and take the MRS). By (4) of Lemma (7.7), it must be E . Furthermore the exceptional divisor of simple blow up of D_2 lands on \tilde{Q}' . Since E is the unique log crepant divisor for $K_{\tilde{Z}} + \tilde{E} + \tilde{B}$ such that its multiplicity in \tilde{E} is less than 1, the flip of \tilde{C}' is the flip of the type (1) in (7.0) or the type treated next in (C-2). (Note that \tilde{M}' corresponds to L in (3.2).) So we obtain the reduction as explained in the flow chart.

(C-2). Next consider the case $g(E) = L$. By (2) of Lemma (7.8), only E and the exceptional divisor of the simple blow up of D_1 or D_2 are the divisors such that its multiplicity in S is less than or equal to $d(E, S) + 1$. Note that they cannot land on \tilde{P}' or \tilde{M}' since they are not projective. Hence for any log crepant divisor for $K_{\tilde{Z}} + \tilde{S}' + \tilde{B}' + \tilde{E}'$ whose center contains \tilde{P}' , its multiplicity in $\tilde{S}' + (d(E, S) + 1)\tilde{E}'$ is greater than $d(E, S) + 1$, which in turn show that its multiplicity in \tilde{E}' is greater than 1, which we want.

(D). If $\tilde{g}^{-1}S \cap \tilde{E}$ is reducible, we can apply (7.6) for \tilde{g} and assume that (1), (2), (3) and (4) of Lemma (7.7) hold. By (4), $d(\tilde{F}, S) = d(\tilde{E}, S) + 1$ and $a_l(\tilde{F}, K_X + S + B) = 0$ for the exceptional divisor \tilde{F} of the simple blow up along \tilde{D}_1 or \tilde{D}_2 and for any other log crepant divisor $\tilde{F} \neq \tilde{E}$ for $K_X + S + B$, $d(\tilde{F}, S) > d(\tilde{E}, S) + 1$. In particular, for any log crepant divisor $\tilde{F} \neq \tilde{E}$ for $K_X + S + B$, $d(\tilde{F}, S) > 1$. But this contradicts the existence of E . Hence in this case we can construct the flip of f as explained in the flow chart. \square

Treatment of (2) in Set up (7.0), $\delta = 0$ and g is good.

By (7.9), we can assume that there is a curve M of LLC $(K_Z + g^{-1}S + g^{-1}B + E|_E)$ through Q' .

Claim. Q' is smooth on $g^{-1}S$, E and Z .

Proof. By (1.12), there is no curve singularity outside $g^{-1}S \cup E$. If there is a curve singularity through Q' contained in $g^{-1}S \cup E$, it must be M (note that Z is smooth generically along L' by (7.12)). Hence there is a log crepant divisor F for $K_Z + g^{-1}S + g^{-1}B + E$ such that $d(F, g^{-1}S + E) \leq 1$ over that curve singularity. Hence

$$d(F, S) = d(F, g^{-1}S) + d(E, S)d(F, E) < d(d(F, g^{-1}S) + d(F, E)) \leq d,$$

a contradiction to the choice of d . By the same way, we can prove that there is no log crepant divisor for $K_Z + g^{-1}S + g^{-1}B + E$ such that $d(F, g^{-1}S + E) \leq 1$ over Q' or a curve through Q' . So $K_Z + g^{-1}S + g^{-1}B + E$ satisfies (*) at Q' . Then (1) or (2) of Theorem (7.1) holds since both $g^{-1}S$ and E contain no curve singularity through Q' . If (2) holds, let $h : Z' \rightarrow Z$ be the extraction of F' taken in (7.1). The existence of $g^{-1}L$ and M contradicts the description of (2). \square

By the above claim, there is a index 2 exceptional complement for $K_X + S|_S$. In fact, we have only to replace $g^{-1}L$ by two smooth curve intersecting D at smooth points transversely with coefficient $\frac{1}{2}$ and consider the images of them. By the argument of (7.11), it lifts to a index 2 exceptional complement for $K_X + S$. Hence the flip of C exists (type 4). \square

Treatment of (2) in Set up (7.0), $\delta = 0$ and g is semi-good.

First we treat the case $g(E) = Q$. By (7.7), we may assume that (1), (2), (3) and (4) hold. On the other hand, $2 \geq d(E, S) > 1$ by (7.12), Hence $-(D_2)_{g^{-1}S}^2 = 3$ and $d(E, S) = \frac{3}{2}$. We prove that there is a PLT index 2 complement near C , which in turn show that the flip of C exists (type 3).

Claim 1.

- (1) Let $\mu : E^\mu \rightarrow E$ be the MRS of E . Then $(\mu^{-1}D_1)^2 \leq 0$.
- (2) One of P_1 and P_2 , say P_1 is smooth on E and $g^{-1}S$ and $g^{-1}C$ passes through it. P_2 is an ODP on E and is smooth on $g^{-1}S$. $(D_1)_E^2 = \frac{1}{2}$ and $(D_1)_{g^{-1}S}^2 = -2$.
- (3) C is irreducible. On $g^{-1}C$, $g^{-1}S$ has one singular point resolved by one (-3) -curve.
- (4) $E \simeq \mathbb{F}_{2,0}$. The index of $K_X + S$ is 2 at Q .

See FIGURE (XXVI).

Proof.

- (1) Assume that $(\mu^{-1}D_1)^2 > 0$. Since $\mu^{-1}D_1$ and $\mu^{-1}D_2$ are nef and big, the inequality $(\mu^{-1}D_1)^2 \cdot (\mu^{-1}D_2)^2 \leq (\mu^{-1}D_1 \cdot \mu^{-1}D_2)^2$ holds. On the other hand, $(\mu^{-1}D_2)^2 = 2$ and $\mu^{-1}D_1 \cdot \mu^{-1}D_2 = 1$ since E is smooth on D_2 , $(D_2)_E^2 = 2$ and $D_1 \cdot D_2 = 1$. But this contradicts the inequality.
- (2) By the ampleness of $g^{-1}S|_E$, we have

$$0 < g^{-1}S \cdot D_1 = ((D_1 + D_2) \cdot D_1)_E = (D_1)_E^2 + 1, i.e.,$$

$$(*) \quad (D_1)_E^2 > -1.$$

On the other hand, by $0 = g^*S \cdot D_1 = g^{-1}S \cdot D_1 + \frac{3}{2}E \cdot D_1$, we obtain

$$(**) \quad (D_1)_{g^{-1}S}^2 = -\frac{2}{3}(D_1)_E^2 - \frac{5}{3}.$$

We may assume that an irreducible component of $g^{-1}C$ passes through P_1 . Then P_1 is smooth since $K_Z + g^{-1}S + g^{-1}B + E|_{g^{-1}S}$ is LT at P_1 of index 2 and $g^{-1}C \subset \text{Supp } g^{-1}B|_{g^{-1}S}$. P_1 is smooth also on E by the same reason. P_2 is a smooth point or an ODP on $g^{-1}S$ and E . By (*) and (1), $(D_1)_E^2 = 0$ (resp. $(D_1)_E^2 = -\frac{1}{2}$ or $\frac{1}{2}$) if P_2 is smooth on E (resp. if P_2 is an ODP on E). Hence by (**), $(D_1)_{g^{-1}S}^2 = -\frac{5}{3}$ (resp. $(D_1)_{g^{-1}S}^2 = -\frac{4}{3}$ or -2) if P_2 is smooth on E (resp. if P_2 is an ODP on E). If P_2 is an ODP on $g^{-1}S$, $(D_1)_{g^{-1}S}^2 \in \frac{\mathbb{Z}}{2}$ and $\notin \mathbb{Z}$, a contradiction. So P_2 is smooth on

$g^{-1}S$ and hence $(D_1)_{g^{-1}S}^2$ is an integer. Consequently P_2 must be an ODP on E and $(D_1)_{g^{-1}S}^2 = -2$.

- (3) By (2), there is a ODP curve singularity D_0 on $g^{-1}S$ through P_2 . So no irreducible component of $g^{-1}C$ passes through P_2 . This prove that C is irreducible. By (2), $(K_Z + g^{-1}S).D_1 = (K_{g^{-1}S} + \frac{1}{2}D_0).D_1 = \frac{1}{2}$. On the other hand, by (2), $E.D_1 = -1$. Hence

$$(***) \quad K_Z + g^{-1}S + \frac{1}{2}E = g^*(K_X + S).$$

By (7.10), $g^{-1}C$ is a (-1) -curve or (-2) -curve on the MRS of $g^{-1}S$. But if $g^{-1}C$ is a (-2) -curve, by $(***)$, $(K_X + S).C = \frac{1}{2} > 0$, a contradiction to the speciality of the flipping contraction f . Hence $g^{-1}C$ is a (-1) -curve on the MRS of $g^{-1}S$ and (1) or (2) of Lemma (7.10) holds. If (2) holds and $m = 0$, by $(***)$, we have $(K_X + S).C = -1 + \frac{1}{2} + \frac{1}{2} = 0$, a contradiction to the speciality of the flipping contraction f . If (2) holds and $m > 0$, we obtain on the MRS of $g^{-1}S$ a noncontractible chain consisting of a (-2) -curve resolving of x , the strict transforms of $g^{-1}C$ and D_1 , a contradiction. Hence (1) holds and we are done.

- (4) Let $\nu : E \rightarrow \bar{E}$ be the morphism defined by sufficient multiple of D_2 . Since E is smooth on D_2 and $(D_2)_E^2 = 2$, $\bar{E} \simeq \mathbb{F}_{2,0}$ or $\mathbb{P}^1 \times \mathbb{P}^1$. Since $(D_1)_E^2 = \frac{1}{2}$, $\nu(D_1)^2 \geq \frac{1}{2}$. But on \bar{E} , $\nu(D_1)$ and $\nu(D_2)$ intersect at one point. Hence $\nu(D_1)^2 = \frac{1}{2}$, which in turn show that $\tilde{E} \simeq \mathbb{F}_{2,0}$ and ν is isomorphism near D_1 . But since $D_1 + D_2$ is ample on E , the exceptional curve of ν must intersect D_1 . So actually ν is isomorphism, i.e., $E \simeq \mathbb{F}_{2,0}$. There is no curve singularity of Z on E . In fact, if there exists a curve singularity of Z on E , it intersects D_1 or D_2 by the ampleness of $D_1 + D_2$ on E . But this is impossible by (2) and Lemma (7.7) (1). Hence E is a Cartier divisor since $E \simeq \mathbb{F}_{2,0}$ (cf. proof of (1.11)). By this and $(***)$, the index of $K_X + S$ is 2 at Q .

□

By Claim 1 (2) and (4), $(K_X + S).C = -\frac{1}{6}$. There exists a Weil divisor \bar{B} on S through x (the singularity on C) such that $K_S + \frac{1}{2}\bar{B}'$ is PLT and has index 2 and $\bar{B}'.C = -\frac{1}{3}$. By (7.11), it lifts to an index 2 PLT complement for $K_X + S$.

Next we treat the case $g(E) = L$. By (7.8), (1) and (2) of the lemma hold.

Claim 2.

- (1) One of P_1 and P_2 , say P_1 is smooth on E and $g^{-1}S$ and $g^{-1}C$ passes through it. P_2 is smooth on E and is an ODP on $g^{-1}S$. $(D_1)_E^2 = 0$ and $(D_1)_{g^{-1}S}^2 = -\frac{3}{2}$. P is smooth on E .
- (2) C is irreducible. On $g^{-1}C$, $g^{-1}S$ is smooth and there is a ODP curve singularity $D_{-1} \subset g^{-1}S$ of Z or a component of $\text{Supp } g^{-1}B$ with coefficient $\frac{1}{2}$ in $g^{-1}B|_{g^{-1}S}$ normally crossing with $g^{-1}C$.
- (3) The index of $K_X + S$ is 2 near C .

See FIGURE (XXVII).

Proof.

- (1) Take P_1 as in Claim 1. The statement about P_1 is proved as in Claim 1. Let l be a general fiber of $E \rightarrow L$. Then $g^{-1}B.l = \frac{1}{2}$ by (7.12). On the other hand $g^{-1}B.D_1 \geq \frac{1}{2}$ since D_1 intersect $g^{-1}B|_E$ at a smooth point P_1 . Since the fiber containing D_1 deforms to l , D_1 must be numerically equivalent to l in Z . In particular, D_1 is an irreducible and reduced

fiber. Hence P and P_2 are smooth on E . Furthermore $-\frac{1}{2} = E.l = E.D_1 = (D_1)_{g^{-1}S}^2 + 1$, i.e., $(D_1)_{g^{-1}S}^2 = -\frac{3}{2}$. By this, P_2 is an ODP on $g^{-1}S$.

- (2) The proof is similar to Claim 1 (3). By (1), P_2 is an ODP on $g^{-1}S$. So no irreducible component of $g^{-1}C$ passes through P_2 . This prove that C is irreducible. By $(K_Z + g^{-1}S).D_1 = (K_{g^{-1}S}).D_1 = 0$, we have

$$(*) \quad (K_Z + g^{-1}S) = g^*(K_X + S).$$

By (7.10), $g^{-1}C$ is a (-1) -curve or (-2) -curve on the MRS of $g^{-1}S$. If $g^{-1}C$ is a (-2) -curve, by (*), $(K_X + S).C = 0$, a contradiction. Hence $g^{-1}C$ is a (-1) -curve and (1) or (2) of the lemma holds. If (1) holds, then we obtain a non-contractible chain on the MRS of $g^{-1}S$ consisting of the (-3) -curve resolving the singularity on $g^{-1}C$, the strict transforms of $g^{-1}C$ and D_1 and (-2) -curve resolving P_2 , a contradiction. Hence (2) holds. But if $m > 0$, we have again a noncontractible chain, a contradiction. Hence we obtain what we want.

- (3) By (2), the index of $K_Z + g^{-1}S$ near $g^{-1}C$ is 2. Hence by (*), the index of $K_X + S$ is also 2 near C .

□

Let \overline{B} is a general smooth Cartier divisor near C such that $\overline{B}.C = 1$. By Claim 2 (3), $(K_X + S).C = -\frac{1}{2}$. Hence $(K_X + S + \frac{1}{2}\overline{B}).C = 0$. By Claim 2 (3), the index of $K_X + S + \frac{1}{2}\overline{B}$ is 2. Taking \tilde{B} generally, this is PLT so we are done. □

REFERENCES

- [S] V. V. Shokurov, *3-fold log flips*, Math. USSR. Izv. **40**, No. 1 (1993), 95–202.
 [FA] J. Kollár et.al., *Flips and abundance for algebraic threefolds*, Astérisque **211** (1992).
 [KMM] Y. Kawamata, K. Matsuda and K. Matsuki, *Introduction to the minimal model problem*, Adv. St. Pure Math. **10** (1987), 287–360.
 [Ka2] ———, *The minimal discrepancy of a 3-fold terminal singularity - appendix to [S]*.
 [Ka3] ———, *Divisorial contractions to 3-dimensional terminal quotient singularity*, Higher dimensional complex varieties, Walter de Gruyter, 1996, p. 241–246.
 [Ka4] ———, *On Fujita's freeness conjecture for 3-folds and 4-folds*, Math. Ann. **308** (1997), 491–505.
 [MS] D. Morrison and G. Stevens, *Terminal quotient singularities in dimensions three and four*, Proc. Amer. Math. Soc **90** (1984), 15–20.

DEPARTMENT OF MATHEMATICAL SCIENCE, UNIVERSITY OF TOKYO, KOMABA, MEGURO-KU, TOKYO 153, JAPAN

E-mail address: htakagi@ms.u-tokyo.ac.jp