Z-Pencils

J. J. McDonald * $^{\ddagger \ddagger}$ D.

D. D. Olesky ^{† ‡‡} H. Schneider [‡] P. van den Driessche ^{§ ‡‡} M. J. Tsat
someros * ‡‡

September 7, 2017

Abstract

The matrix pencil $(A, B) = \{tB - A \mid t \in \mathbf{C}\}$ is considered under the assumptions that A is entrywise nonnegative and B - A is a nonsingular M-matrix. As t varies in [0, 1], the Z-matrices tB - A are partitioned into the sets L_s introduced by Fiedler and Markham. As no combinatorial structure of B is assumed here, this partition generalizes some of their work where B = I. Based on the union of the directed graphs of A and B, the combinatorial structure of nonnegative eigenvectors associated with the largest eigenvalue of (A, B) in [0, 1) is considered.

Key words: Z-matrix, matrix pencil, M-matrix, eigenspace, reduced graph.

AMS subject classifications: 15A22, 15A48, 05C50.

1 Introduction

The generalized eigenvalue problem $Ax = \lambda Bx$ for $A = [a_{ij}], B = [b_{ij}] \in \mathbf{R}^{n,n}$, with inequality conditions motivated by certain economics models, was studied by Bapat et al. [1]. In keeping with this work, we consider the matrix pencil $(A, B) = \{tB - A \mid t \in \mathbf{C}\}$ under the conditions

- (1) A is entrywise nonnegative, denoted by $A \ge 0$
- (2) $b_{ij} \le a_{ij}$ for all $i \ne j$
- (3) there exists a positive vector u such that (B A)u is positive.

Note that in [1] A is also assumed to be irreducible, but that is not imposed here. When $Ax = \lambda Bx$ for some nonzero x, the scalar λ is an *eigenvalue* and x is the corresponding

 $^{^*\}mathrm{Dept}$ of Mathematics and Statistics, Univ. of Regina, Regina, Saskatchewan S4S 0A2.

[†]Dept of Computer Science, Univ. of Victoria, Victoria, British Columbia V8W 3P6.

[‡]Dept of Mathematics, Univ. of Wisconsin, Madison, Wisconsin 53706.

[§]Dept of Mathematics and Statistics, Univ. of Victoria, Victoria, British Columbia V8W 3P4.

^{‡‡}Research partially supported by NSERC research grant.

eigenvector of (A, B). The eigenspace of (A, B) associated with an eigenvalue λ is the nullspace of $\lambda B - A$.

A matrix $X \in \mathbf{R}^{n,n}$ is a Z-matrix if X = qI - P, where $P \ge 0$ and $q \in \mathbf{R}$. If, in addition, $q \ge \rho(P)$, where $\rho(P)$ is the spectral radius of P, then X is an M-matrix, and is singular if and only if $q = \rho(P)$. It follows from (1) and (2) that when $t \in [0, 1]$, tB - A is a Z-matrix. Henceforth the term Z-pencil (A, B) refers to the circumstance that tB - A is a Z-matrix for all $t \in [0, 1]$.

Let $\langle n \rangle = \{1, 2, ..., n\}$. If $J \subseteq \langle n \rangle$, then X_J denotes the principal submatrix of X in rows and columns of J. As in [3], given a nonnegative $P \in \mathbf{R}^{n,n}$ and an $s \in \langle n \rangle$, define

$$\rho_s(P) = \max_{|J|=s} \{\rho(P_J)\}$$

and set $\rho_{n+1}(P) = \infty$. Let L_s denote the set of Z-matrices in $\mathbb{R}^{n,n}$ of the form qI - P, where $\rho_s(P) \leq q < \rho_{s+1}(P)$ for $s \in \langle n \rangle$, and $-\infty < q < \rho_1(P)$ when s = 0. This gives a partition of all Z-matrices of order n. Note that $qI - P \in L_0$ if and only if $q < p_{ii}$ for some i. Also, $\rho_n(P) = \rho(P)$, and L_n is the set of all (singular and nonsingular) M-matrices.

We consider the Z-pencil (A, B) subject to conditions (1)-(3) and partition its matrices into the sets L_s . Viewed as a partition of the Z-matrices tB - A for $t \in [0, 1]$, our result provides a generalization of some of the work in [3] (where B = I). Indeed, since no combinatorial structure of B is assumed, our Z-pencil partition is a consequence of a more complicated connection between the Perron-Frobenius theory for A and the spectra of tB - A and its submatrices.

Conditions (2) and (3) imply that B - A is a nonsingular M-matrix and thus its inverse is entrywise nonnegative (see [2, N₃₈, p. 137]). This, together with (1), gives $(B - A)^{-1}A \ge 0$. Perron-Frobenius theory is used in [1] to identify an eigenvalue $\rho(A, B)$ of the pencil (A, B), defined as

$$\rho(A,B) = \frac{\rho((B-A)^{-1}A)}{1+\rho((B-A)^{-1}A)}.$$

Our partition involves $\rho(A, B)$ and the eigenvalues of the subpencils (A_J, B_J) . Our Z-pencil partition result, Theorem 2.4, is followed by examples where as t varies in [0, 1], tB - A ranges through some or all of the sets L_s for $0 \le s \le n$. In Section 3 we turn to a consideration of the combinatorial structure of nonnegative eigenvectors associated with $\rho(A, B)$. This involves some digraph terminology, which we introduce at the beginning of that section.

In [3], [7] and [5], interesting results on the spectra of matrices in L_s , and a classification in terms of the inverse of a Z-matrix, are established. These results are of course applicable to the matrices of a Z-pencil, however, as they do not directly depend on the form tB - A of the Z-matrix, we do not consider them here.

2 Partition of Z-pencils

We begin with two observations and a lemma used to prove our result on the Z-pencil partition.

Observation 2.1 Let (A, B) be a pencil with B - A nonsingular. Given a real $\mu \neq -1$, let $\lambda = \frac{\mu}{1+\mu}$. Then the following hold:

(i) $\lambda \neq 1$ is an eigenvalue of (A, B) if and only if $\mu \neq -1$ is an eigenvalue of $(B - A)^{-1}A$.

- (ii) λ is a strictly increasing function of $\mu \neq -1$.
- (iii) $\lambda \in [0,1)$ if and only if $\mu \ge 0$.

Proof. If μ is an eigenvalue of $(B - A)^{-1}A$, then there exists nonzero $x \in \mathbb{R}^n$ such that $(B - A)^{-1}Ax = \mu x$. It follows that $Ax = \mu (B - A)x$ and if $\mu \neq -1$, then $Ax = \frac{\mu}{1+\mu}Bx = \lambda Bx$. Notice that λ cannot be 1 for any choice of μ . The reverse argument shows that the converse is also true. The last statement of (i) is obvious. Statements (ii) and (iii) follow easily from the definition of λ .

Note that $\lambda = 1$ is an eigenvalue of (A, B) if and only if B - A is singular.

Observation 2.2 Let (A, B) be a pencil satisfying (2), (3). Then the following hold:

- (i) For any nonempty $J \subseteq \langle n \rangle$, $B_J A_J$ is a nonsingular M-matrix.
- (ii) If in addition (1) holds, the largest real eigenvalue of (A, B) in [0, 1) is $\rho(A, B)$.

Proof. (i) This follows since (2) and (3) imply that B - A is a nonsingular M-matrix (see [2, I₂₇, p. 136]) and since every principal submatrix of a nonsingular M-matrix is also a nonsingular M-matrix (see [2, p. 138]).

(ii) This follows from Observation 2.1, since $\mu = \rho((B - A)^{-1}A)$ is the maximal positive eigenvalue of $(B - A)^{-1}A$.

Lemma 2.3 Let (A, B) be a pencil satisfying (1)-(3). Let $\mu = \rho((B - A)^{-1}A)$ and $\rho(A, B) = \frac{\mu}{1+\mu}$. Then the following hold:

(i) For all $t \in (\rho(A, B), 1]$, tB - A is a nonsingular M-matrix.

(ii) The matrix $\rho(A, B)B - A$ is a singular M-matrix.

(iii) For all $t \in (0, \rho(A, B))$, tB - A is not an M-matrix.

(iv) For t = 0, either tB - A is a singular M-matrix or is not an M-matrix.

Proof. Recall that (1) and (2) imply that tB - A is a Z-matrix for all $0 < t \le 1$. As noted in Observation 2.2 (i), B - A is a nonsingular M-matrix and thus its eigenvalues have positive real parts [2, G₂₀, p. 135], and the eigenvalue with minimal real part is real ([2, Exercise 5.4, p. 159]. Since the eigenvalues are continuous functions of the entries of a matrix, as t decreases from t = 1, tB - A is a nonsingular M-matrix for all t until a value of t is encountered for which tB - A is singular. Results (i) and (ii) now follow by Observation 2.2 (ii).

To prove (iii), consider $t \in (0, \rho(A, B))$. Since $(B-A)^{-1}A \ge 0$, there exists an eigenvector $x \ge 0$ such that $(B-A)^{-1}Ax = \mu x$. Then $Ax = \rho(A, B)Bx$ and $(tB-A)x = (t-\rho(A, B))Bx \le 0$ since $Bx = \frac{1}{\rho(A,B)}Ax \ge 0$. By [2, A₅, p. 134], tB - A is not a nonsingular M-matrix. To complete the proof (by contradiction), suppose $\alpha B - A$ is a singular M-matrix for some $\alpha \in$ $(0, \rho(A, B))$. Since there are finitely many values of t for which tB - A is singular, we can choose $\beta \in (\alpha, \rho(A, B))$ such that $\beta B - A$ is nonsingular. Let $\epsilon = \frac{\beta - \alpha}{\alpha}$. Then $(1 + \epsilon)(\alpha B - A)$ is a singular M-matrix and

$$(1+\epsilon)(\alpha B - A) + \gamma I = \beta B - A - \epsilon A + \gamma I \le \beta B - A + \gamma I$$

since $A \ge 0$ by (1). By [2, C₉, p. 150], $\beta B - A - \epsilon A + \gamma I$ is a nonsingular M-matrix for all $\gamma > 0$, and hence $\beta B - A + \gamma I$ is a nonsingular M-matrix for all $\gamma > 0$ by [4, 2.5.4, p. 117]. This implies that $\beta B - A$ is also a (nonsingular) M-matrix ([2, C₉, p. 150]), contradicting the above. Thus we can also conclude that $\alpha B - A$ cannot be a singular M-matrix for any choice of $\alpha \in (0, \rho(A, B))$, establishing (iii). For (iv), -A is a singular M-matrix if and only if it is, up to a permutation similarity, strictly triangular. Otherwise, -A is not an M-matrix.

Theorem 2.4 Let (A, B) be a pencil satisfying (1)-(3). For s = 1, 2, ..., n let

$$\sigma_s = \max_{|J|=s} \{ \rho \left((B_J - A_J)^{-1} A_J \right) \}, \quad \tau_s = \frac{\sigma_s}{1 + \sigma_s}$$

and $\tau_0 = 0$. Then for $s = 0, 1, \ldots, n-1$ and $\tau_s \leq t < \tau_{s+1}$, the matrix $tB - A \in L_s$. For s = nand $\tau_n \leq t \leq 1$, the matrix $tB - A \in L_n$.

Proof. Fiedler and Markham [3, Theorem 1.3] show that for $1 \leq s \leq n-1$, $X \in L_s$ if and only if all principal submatrices of X of order s are M-matrices, and there exists a principal submatrix of order s+1 that is not an M-matrix. Consider any nonempty $J \subseteq \langle n \rangle$ and $t \in [0, 1]$. Conditions (1) and (2) imply that $tB_J - A_J$ is a Z-matrix. By Observation 2.2 (i), $B_J - A_J$ is a nonsingular M-matrix. Let $\mu_J = \rho \left((B_J - A_J)^{-1} A_J \right)$. Then by Observation 2.2 (ii), $\tau_J = \frac{\mu_J}{1 + \mu_J}$ is the largest eigenvalue in [0,1) of the pencil (A_J, B_J) . Combining this with Observation 2.2 (i) and Lemma 2.3, the matrix $tB_J - A_J$ is an M-matrix for all $\tau_J \leq t \leq 1$, and $tB_J - A_J$ is not an M-matrix for all $0 < t < \tau_J$. If $1 \le s \le n-1$ and |J| = s, then $tB_J - A_J$ is an M-matrix for all $\tau_s \leq t \leq 1$. Suppose $\tau_s < \tau_{s+1}$. Then there exists $K \subseteq \langle n \rangle$ such that |K| = s + 1 and $tB_K - A_K$ is not an M-matrix for $0 < t < \tau_{s+1}$. Thus by [3, Theorem 1.3] $tB - A \in L_s$ for all $\tau_s \leq t < \tau_{s+1}$. When s = n, since B - A is a nonsingular M-matrix, $tB - A \in L_n$ for all t such that $\rho(A, B) = \tau_n \leq t \leq 1$ by Lemma 2.3 (i). For the case s = 0, if $0 < t < \tau_1$, then tB - A has a negative diagonal entry and thus $tB - A \in L_0$. For t = 0, tB - A = -A. If $a_{ii} \neq 0$ for some $i \in \langle n \rangle$, then $-A \in L_0$; if $a_{ii} = 0$ for all $i \in \langle n \rangle$, then $\tau_1 = \tau_0 = 0$, namely, $-A \in L_s$ for some $s \geq 1.$

We continue with illustrative examples.

Example 2.5 Consider

$$A = \begin{pmatrix} 1 & 2 \\ 1 & 0 \end{pmatrix} \text{ and } B = \begin{pmatrix} 2 & 2 \\ 1 & 1 \end{pmatrix},$$

for which $\tau_2 = 2/3$ and $\tau_1 = 1/2$. It follows that

$$tB - A \in \begin{cases} L_0 & \text{if } 0 \le t < 1/2 \\ L_1 & \text{if } 1/2 \le t < 2/3 \\ L_2 & \text{if } 2/3 \le t \le 1. \end{cases}$$

That is, as t increases from 0 to 1, tB - A belongs to all the possible Z-matrix classes L_s .

Example 2.6 Consider the matrices in [1, Example 5.3], that is,

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix} \text{ and } B = \begin{pmatrix} 4 & 0 & -2 & 0 \\ 0 & 3 & 0 & -1 \\ -2 & 0 & 4 & 0 \\ 0 & -2 & 0 & 4 \end{pmatrix}$$

Referring to Theorem 2.4, $\tau_4 = \rho(A, B) = \frac{4+\sqrt{6}}{10} = \tau_3 = \tau_2$ and $\tau_1 = 1/3$. It follows that

$$tB - A \in \begin{cases} L_0 & \text{if } 0 \le t < 1/3 \\\\ L_1 & \text{if } 1/3 \le t < \frac{4+\sqrt{6}}{10} \\\\ L_4 & \text{if } \frac{4+\sqrt{6}}{10} \le t \le 1. \end{cases}$$

Notice that for $t \in [0, 1]$, tB - A ranges through only L_0 , L_1 and L_4 .

Example 2.7 Now let

$$A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \text{ and } B = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

In contrast to the above two examples, $tB - A \in L_2$ for all $t \in [0, 1]$. Note that, in general, $tB - A \in L_n$ for all $t \in [0, 1]$ if and only if $\rho(A, B) = 0$.

3 Combinatorial Structure of the Eigenspace of $\rho(A, B)$

Let $\Gamma = (V, E)$ be a digraph, where V is a finite vertex set and $E \subseteq V \times V$ is the edge set. If $\Gamma' = (V, E')$, then $\Gamma \cup \Gamma' = (V, E \cup E')$. Also write $\Gamma' \subseteq \Gamma$ when $E' \subseteq E$. For $j \neq k$, a path of length $m \geq 1$ from j to k in Γ is a sequence of vertices $j = r_1, r_2, \ldots, r_{m+1} = k$ such that $(r_s, r_{s+1}) \in E$ for $s = 1, \ldots, m$. As in [2, Ch. 2], if j = k or if there is a path from vertex j to vertex k in Γ , then j has access to k (or k is accessed from j). If j has access to k and k has

access to j, then j and k communicate. The communication relation is an equivalence relation, hence V can be partitioned into equivalence classes, which are referred to as the classes of Γ .

The digraph of $X = [x_{ij}] \in \mathbb{R}^{n,n}$, denoted by $\mathcal{G}(X) = (V, E)$, consists of the vertex set $V = \langle n \rangle$ and the set of directed edges $E = \{(j,k) \mid x_{jk} \neq 0\}$. If j has access to k for all distinct $j, k \in V$, then X is *irreducible* (otherwise, *reducible*). It is well known that the rows and columns of X can be simultaneously reordered so that X is in block lower triangular *Frobenius normal form*, with each diagonal block irreducible. The irreducible blocks in the Frobenius normal form of X correspond to the classes of $\mathcal{G}(X)$.

In terminology similar to that of [6], given a digraph Γ , the reduced graph of Γ , $\mathcal{R}(\Gamma) = (V', E')$, is the digraph derived from Γ by taking

$$V' = \{J \mid J \text{ is a class of } \Gamma\}$$

and

$$E' = \{(J, K) \mid \text{there exist } j \in J \text{ and } k \in K \text{ such that } j \text{ has access to } k \text{ in } \Gamma \}.$$

When $\Gamma = \mathcal{G}(X)$ for some $X \in \mathbf{R}^{n,n}$, we denote $\mathcal{R}(\Gamma)$ by $\mathcal{R}(X)$.

Suppose now that X = qI - P is a singular M-matrix, where $P \ge 0$ and $q = \rho(P)$. If an irreducible block X_J in the Frobenius normal form of X is singular, then $\rho(P_J) = q$ and we refer to the corresponding class J as a singular class (otherwise, a nonsingular class). A singular class J of $\mathcal{G}(X)$ is called distinguished if when J is accessed from a class $K \ne J$ in $\mathcal{R}(X)$, then $\rho(P_K) < \rho(P_J)$. That is, a singular class J of $\mathcal{G}(X)$ is distinguished if and only if J is accessed only from itself and nonsingular classes in $\mathcal{R}(X)$.

We paraphrase now Theorem 3.1 of [6] as follows.

Theorem 3.1 Let $X \in \mathbf{R}^{n,n}$ be an M-matrix and let J_1, \ldots, J_p denote the distinguished singular classes of $\mathcal{G}(X)$. Then there exist unique (up to scalar multiples) nonnegative vectors x^1, \ldots, x^p in the nullspace of X such that

 $x_j^i \begin{cases} = 0 \text{ if } j \text{ does not have access to a vertex in } J_i \text{ in } \mathcal{G}(X) \\ > 0 \text{ if } j \text{ has access to a vertex in } J_i \text{ in } \mathcal{G}(X) \end{cases}$

for all i = 1, 2, ..., p and j = 1, 2, ..., n. Moreover, every nonnegative vector in the nullspace of X is a linear combination with nonnegative coefficients of $x^1, ..., x^p$.

We apply the above theorem to a Z-pencil, using the following lemma.

Lemma 3.2 Let (A, B) be a pencil satisfying (1) and (2). Then the classes of $\mathcal{G}(tB - A)$ coincide with the classes of $\mathcal{G}(A) \cup \mathcal{G}(B)$ for all $t \in (0, 1)$.

Proof. Clearly $\mathcal{G}(tB - A) \subseteq \mathcal{G}(A) \cup \mathcal{G}(B)$ for all scalars t. For any $i \neq j$, if either $b_{ij} \neq 0$ or $a_{ij} \neq 0$, and if $t \in (0, 1)$, conditions (1) and (2) imply that $tb_{ij} < a_{ij}$ and hence $tb_{ij} - a_{ij} \neq 0$. This means that apart from vertex loops, the edge sets of $\mathcal{G}(tB - A)$ and $\mathcal{G}(A) \cup \mathcal{G}(B)$ coincide for all $t \in (0, 1)$.

Theorem 3.3 Let (A, B) be a pencil satisfying (1)-(3) and let

$$\Gamma = \begin{cases} \mathcal{G}(A) \cup \mathcal{G}(B) & \text{if } \rho(A, B) \neq 0 \\ \\ \mathcal{G}(A) & \text{if } \rho(A, B) = 0. \end{cases}$$

Let J_1, \ldots, J_p denote the classes of Γ such that for each $i = 1, 2, \ldots, p$,

- (i) $(\rho(A, B)B A)_{J_i}$ is singular, and
- (ii) if J_i is accessed from a class $K \neq J_i$ in $\mathcal{R}(\Gamma)$, then $(\rho(A, B)B A)_K$ is nonsingular.

Then there exist unique (up to scalar multiples) nonnegative vectors x^1, \ldots, x^p in the eigenspace associated with the eigenvalue $\rho(A, B)$ of (A, B) such that

$$x_{j}^{i} \begin{cases} = 0 \text{ if } j \text{ does not have access to a vertex in } J_{i} \text{ in } \Gamma \\ > 0 \text{ if } j \text{ has access to a vertex in } J_{i} \text{ in } \Gamma \end{cases}$$

for all i = 1, 2, ..., p and j = 1, 2, ..., n. Moreover, every nonnegative vector in the eigenspace associated with the eigenvalue $\rho(A, B)$ is a linear combination with nonnegative coefficients of $x^1, ..., x^p$.

Proof. By Lemma 2.3 (ii), $\rho(A, B)B - A$ is a singular M-matrix. Thus

$$\rho(A, B)B - A = qI - P = X,$$

where $P \ge 0$ and $q = \rho(P)$. When $\rho(A, B) = 0$, the result follows from Theorem 3.1 applied to X = -A. When $\rho(A, B) > 0$, by Lemma 3.2, $\Gamma = \mathcal{G}(X)$. Class J of Γ is singular if and only if $\rho(P_J) = q$, which is equivalent to $(\rho(A, B)B - A)_J$ being singular. Also a singular class J is distinguished if and only if for all classes $K \ne J$ that access J in $\mathcal{R}(X)$, $\rho(P_K) < \rho(P_J)$, or equivalently $(\rho(A, B)B - A)_K$ is nonsingular. Applying Theorem 3.1 gives the result.

We conclude with a generalization of Theorem 1.7 of [3] to Z-pencils. Note that the class J in the following result is a singular class of $\mathcal{G}(A) \cup \mathcal{G}(B)$.

Theorem 3.4 Let (A, B) be a pencil satisfying (1)-(3) and let $t \in (0, \rho(A, B))$. Suppose that J is a class of $\mathcal{G}(tB - A)$ such that $\rho(A, B) = \frac{\mu}{1+\mu}$, where $\mu = \rho((B_J - A_J)^{-1}A_J)$. Let m = |J|. Then $tB - A \in L_s$ with

$$s \begin{cases} \leq n-1 & \text{if } m = n \\ < m & \text{if } m < n \end{cases}$$

Proof. That $tB - A \in L_s$ for some $s \in \{0, 1, ..., n\}$ follows from Theorem 2.4. By Lemma 2.3 (iii), if $t \in (0, \rho(A, B))$, then $tB - A \notin L_n$ since $\rho(A, B) = \tau_n$. Thus $s \leq n - 1$. When m < n, under the assumptions of the theorem, we have $\tau_n = \rho(A, B) = \frac{\mu}{1+\mu} \leq \tau_m$ and hence $\tau_m = \tau_{m+1} = \ldots = \tau_n$. It follows that s < m.

We now apply the results of this section to Example 2.6, which has two classes. Class $J = \{2, 4\}$ is the only class of $\mathcal{G}(A) \cup \mathcal{G}(B)$ such that $(\rho(A, B)B - A)_J$ is singular, and J is accessed by no other class. By Theorem 3.3, there exists an eigenvector x of (A, B) associated with $\rho(A, B)$ with $x_1 = x_3 = 0$, $x_2 > 0$ and $x_4 > 0$. Since |J| = 2, by Theorem 3.4, $tB - A \in L_0 \cup L_1$ for all $t \in (0, \rho(A, B))$, agreeing with the exact partition given in Example 2.6.

References

- R. B. Bapat, D. D. Olesky, and P. van den Driessche. Perron-Frobenius theory for a generalized eigenproblem. *Linear and Multilinear Algebra*, 40:141-152, 1995.
- [2] A. Berman and R. J. Plemmons. Nonnegative Matrices in the Mathematical Sciences, Academic Press, New York, 1979.
- [3] M. Fiedler and T. Markham. A classification of matrices of class Z. Linear Algebra and Its Applications, 173:115-124, 1992.
- [4] Roger A. Horn and Charles R. Johnson. *Topics in Matrix Analysis*, Cambridge University Press, 1991.
- [5] Reinhard Nabben. Z-matrices and inverse Z-matrices. *Linear Algebra and Its Applications*, 256:31-48, 1997.
- [6] Hans Schneider. The influence of the marked reduced graph of a nonnegative matrix on the Jordan Form and on related properties: A survey. *Linear Algebra and Its Applications*, 84:161-189, 1986.
- [7] Ronald S. Smith. Some results on a partition of Z-matrices. Linear Algebra and Its Applications, 223/224:619-629, 1995.