# SMOOTH MODULES OVER LIE ALGEBROIDS I

ROLF KÄLLSTRÖM

## Contents

1. Introduction	2
2. Lie algebroids	6
2.1. Definitions	6
2.2. Linear Lie algebroids	9
2.3. Differential operators	11
2.4. Homomorphisms	13
2.4.1. Remark on the tangent mapping	13
2.4.2. Homomorphisms of Lie algebroids	14
2.4.5. Pull-back	15
2.5. Picard Lie algebroids	17
3. Modules	22
3.1. Basic facts	22
3.2. Interchanging left and right modules	25
3.3. Modules over Picard Lie algebroids	26
3.3.3. Twisted sheaves	27
3.4. Inverse images	28
3.5. Direct images	30
4. Smooth modules	36
4.1. Elementary properties	36
4.2. Smooth complexes	39
4.2.2. Inverse images	40
4.2.6. Direct images	41
4.3. Completion	43
4.4. GAGA	44
5. Prolonging over $\operatorname{codim} \ge 2$	46
5.1. Coherent extensions	46
5.2. Prolongation of smoothness	52
5.3. Proof of Theorem 5.2.1	54
6. Regular singularities	58
6.1. Torsion free modules	58
6.2. Proof of Theorem 6.1.2	61
6.3. Regular singularities	64
6.4. Completely regular complexes	75
References	79

Date: May 30, 2022.

1991 Mathematics Subject Classification. Primary 14A10, 32C38; Secondary 17B99.

### 1. INTRODUCTION

Let  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha)$  be a Lie algebroid (*Def.* 2.1.2) on a space X where  $\alpha$ :  $\mathfrak{g}_X \to T_X$  is a homomorphism to the tangent sheaf; a space is either a scheme locally of finite type over an algebraically closed field k of characteristic 0 or a complex analytic space  $(k = \mathbb{C})$ . Let M be a coherent module over  $\mathfrak{g}_X$ (*Def.* 3.1.1),  $\mathfrak{g}'_X$  a sub-algebroid of  $\mathfrak{g}_X$ , and x a point in X. We say that M is smooth along  $\mathfrak{g}'_X$  at x if its stalk  $M_x$  contains an  $\mathcal{O}_x$ -module  $M_x^0$  of finite type which generates  $M_x$  over  $\mathfrak{g}_x$  and satisfies  $\mathfrak{g}'_X \cdot M_x^0 \subseteq M_x^0$ . In this paper and its continuation [Käl98] we shall study this notion of smoothness, treating the algebraic and complex analytic theory simultaneously. Our main results will be described a little later.

Let  $S \subset X$  be a subspace and  $T_X(I_S) \subset T_X$  be the sub-algebroid of sections that preserve the ideal  $I_S$  of S. Particularly interesting torsion free (over  $\mathcal{O}_X$ )  $T_X$ -modules are those that are smooth along  $T_X(I)$ . When S is a divisor and X is normal then M has this property if and only if it has regular singularities along S in the sense of P. Deligne [Del70]; there are similar results for torsion modules.

Deligne's modestly titled but very influential work [loc. cit.] contains the first treatment of regular singularities of connections on complex spaces. Together with M. Kashiwara's work [Kas] it forms a starting point for the theory of  $\mathcal{D}_X$ -modules (=  $T_X$ -modules), leading to a more uniform treatment of singularities. Some important architects of the analytic theory are Kashiwara, M. Sato, T. Kawai, and Z. Mebkhout, while the algebraic theory is mainly due to J. Bernstein; see the text-books [Bjö93], [BGK<sup>+</sup>87], and [BB93]. Lie algebroids also play a prominent role in differential geometry, where one of the first uses is due to M. Atiyah [Ati57]; see the survey [Mac95].

The intrinsic interest in Lie algebroids should be rather obvious, but since the paper does not deal much with specific applications we briefly mention some connections with other areas.

Quantum gauge theory: The dictionary from the terminology of physics to mathematics is that gauge fields, field strengths, and the Yang-Mills equations (the field equations) correspond to sections, sections of the curvature, and the Bianchi identity for the Hodge dual of the curvature of Lie algebroids; and quantum gauge fields act on modules over Lie algebroids. Gauge theories first attracted differential geometers and analysts, considering the geometry of principal bundles and solutions of the Yang-Mills equations. Now an interesting case of gauge theory on curves occurs in conformal field theory, studying Lie algebroids on curves and Lie algebroids on moduli spaces of curves. The subject has therefore attracted the attention of a great many algebraic geometers; see [BS88], [TUY89], and [Uen97].

Cohomology: Let M be a regular holonomic  $T_X$ -module on a non-singular space X. Then the local cohomology sheaves  $H^i_{[S]}(M)$  also are regular holonomic  $T_X$ -modules, even when they are not coherent over  $\mathcal{O}_X$ ; see e.g. [Lyu93] for one use of the  $T_X$ -module structure on  $H^i_{[S]}(M)$ . As for the cohomology of S itself a nice version is intersection cohomology, occurring as the homology of certain objects in the category of perverse sheaves, which by the Riemann-Hilbert correspondence is equivalent to the category of regular holonomic  $T_X$ -modules whose support belong to S; see [KK81], [Meb84], [BGK<sup>+</sup>87] and [BBD82]. Similar facts hold for modules over any transitive Lie algebroid.

Localisation of representations: Often one is concerned with homomorphisms of Lie algebras over k of the type  $\alpha : \mathfrak{g} \to \Gamma(X, T_X)$ : The action of a Lie group on a variety  $G \times X \to X$  is one source of such  $\alpha$ ; other interesting sources occur in moduli problems related to curves, where  $\alpha$  is the inverse of a Kodaira-Spencer mapping; see [loc. cit.]. The homomorphism  $\alpha$  gives the  $\mathcal{O}_X$ -module  $\mathfrak{g}^0_X := \mathcal{O}_X \otimes_k \mathfrak{g}$  a structure of Lie algebroid, and letting  $\mathfrak{g}_X$  be the push-out of  $\mathfrak{g}^0_X$  by a character of  $\operatorname{Ker}(\mathfrak{g}^0_X \to T_X)$  one has a homomorphism  $\alpha : \mathcal{D}(\mathfrak{g}) \to \mathcal{D}(\mathfrak{g}_X)$  from the enveloping algebra of  $\mathfrak{g}$  to the enveloping ring of differential operators of  $\mathfrak{g}_X$ . Now the  $\mathfrak{g}_X$ -module  $M_X := \mathcal{D}(\mathfrak{g}_X) \otimes_{\mathcal{D}(\mathfrak{g})} M$  serves as a bridge to a geometric description of a  $\mathfrak{g}$ -module M using the Riemann-Hilbert correspondence. For example, when X is a flag variety the localisation functor  $M \mapsto M_X$  is essential in the proof of the Kazhdan-Lusztig algorithm for the Jordan-Hölder multiplicities of Verma modules over a complex semi-simple Lie algebra; see [BB81], [BK81], [BB93], [Kas89].

Families: Coherent sheaves of Lie algebras  $\mathfrak{g}_X$  over  $\mathcal{O}_X$  (the Lie bracket being  $\mathcal{O}_X$ -bilinear) are obvious examples of Lie algebroids. We think of these as families of Lie algebras  $\mathfrak{g}(x) := \mathfrak{g}_x/\mathfrak{m}_x\mathfrak{g}_x$  over the fields  $k(x) = \mathcal{O}_x/\mathfrak{m}_x$ , where a trivial family is one which is isomorphic to an extension of scalars  $\mathfrak{g}_X = \mathcal{O}_X \otimes_k \mathfrak{g}$  of a Lie algebra  $\mathfrak{g}$  over a field k; more interesting cases occur when the Lie bracket  $[\cdot, \cdot]_x$  for  $\mathfrak{g}(x)$  degenerates at special points  $x \in X$ (contraction). Similarly, a sheaf of modules M over a sheaf of Lie algebras  $\mathfrak{g}_X$  we regard as a family of modules, resulting in a fibre module M(x) := $M/\mathfrak{m}_x M$  over the k(x)-Lie algebra  $\mathfrak{g}(x)$  at each point x. An interesting decreasing filtration of M(x) by  $\mathfrak{g}(x)$ -modules  $\cdots \subset M_n(x) \subset M_{n-1}(x) \subset$  $\cdots \subset M_0(x) = M(x)$  is defined using an ideal  $I \subseteq \mathfrak{m}_x$  and a section  $S \in$  $Hom_{\mathfrak{g}_X}(M \otimes_{\mathcal{O}_X} M, \mathcal{O}_X)$  (where usually the  $\mathfrak{g}_X$ -module  $\mathcal{O}_X$  is trivial; S is a "Shapovalov form"), setting  $M_n(x) := \{m \in M : S(m, M) \subseteq I^n\} \mod \mathfrak{m}_x$ . Jantzen's filtration of Verma modules is an important example; see [Jan79], [BB93].

The content of the sections is as follows:

Section 2: is mostly a survey containing general definitions and basic facts about Lie algebroids, complementing and recalling parts of [BB93]. There are no really new results and we include more material than is absolutely needed for our main results, but it should be helpful to have this information collected.

Section 3: Proposition 3.1.2 states that it is equivalent that a space X is non-singular, that the  $T_X$ -module  $\mathcal{O}_X$  contains no coherent proper submodules, and that the Lie algebroid  $T_X$  contains no coherent ideals; this should be well-known. The main part of this section has to do with operations on Lie algebroids and their derived categories of complexes of modules, generalizing standard operations in  $\mathcal{D}$ -module theory.

Section 4: Propositions 4.1.5 and 4.1.7 are relations between global and local smoothness. Limiting ourselves to morphisms  $\pi: Y \to X$  of non-singular

spaces we first show that the derived pull-back functor  $\pi^+$  preserves the derived category  $D_s^b$  of bounded complexes of  $\mathfrak{g}$ -modules with smooth homology modules (*Prop.* 4.2.3), and we also give a partial converse (*Prop.* 4.2.4). Theorem 4.2.7 gives a condition when the direct image functor preserves  $D_s^b$ ; it will later be used to prove that the direct image of a module with regular singularities again has regular singularities. We record the important fact implied by the Artin-Rees lemma, that smoothness of a module is well reflected in its completion (*Prop.* 4.3.1). Theorem 4.4.1 contains the expected equivalences implied by GAGA; its Corollary 4.4.2 is a statement about cohomology groups of certain sheaves which are not  $\mathcal{O}$ -modules.

Section 5: We say that an  $\mathcal{O}_X$ -module M is pure if all its non-zero coherent sub-modules have the same dimension of their support; M is torsion free if it is pure and dim supp  $M = \dim X$ . Let M be a coherent  $\mathcal{O}_X$ -module with support  $V \subset X$ , and  $j : \Omega \to X$  an open inclusion of spaces such that  $\operatorname{codim}_V(V \cap (X \setminus \Omega)) \geq 2$ . Then if  $j^*(M)$  is pure we have  $j_*j^*(M) \in$  $\operatorname{coh}(\mathcal{O}_X)$  (Cor. 5.1.7). In the algebraic case Corollary 5.1.7 follows from A. Grothendieck's finiteness theorem [Gro62, Exp. VIII, p. 13, Th. 3.1] while in the complex analytic case it follows from a corresponding result by Y.-T. Siu and G. Trautmann [Siu70],[Tra69], as stated in Theorem 5.1.3. We include a proof of this theorem in the algebraic case which perhaps is somewhat more direct than that of Grothendieck.

Using Corollary 5.1.7 we can prove our first main result in Theorem 5.2.1, stating: Let  $\pi : Y \to X$  be a proper morphism of spaces,  $S \subset X$  a (closed) subspace, and  $\mathfrak{g}'_Y \subset \mathfrak{g}_Y$  an inclusion of Lie algebroids on Y. Suppose that  $M \in \operatorname{coh}(\mathfrak{g}_Y)$  contains, locally in X, an  $\mathcal{O}_Y$ -lattice  $M^0$ , i.e.  $M = \mathcal{D}(\mathfrak{g}_Y)M^0$ and  $M^0 \in \operatorname{coh}(\mathcal{O}_X)$  (such  $M^0$  always exist when X is a noetherian variety), and let  $\cup_{i=1}^k V_i$  be an irreducible decomposition of the support of M. Assume that  $V_i \cap V_j \cap \pi^{-1}(S) = \emptyset$  for all pairs  $i \neq j$  and that dim  $\pi(V_i) \geq \dim S + 2$ when  $V_i \cap \pi^{-1}(S) \neq \emptyset$ . Assume moreover that the maximal coherent  $\mathfrak{g}_X$ submodule of M with support in  $V_i$  is pure,  $i = 1, \ldots, k$ , and that the canonical homomorphisms

$$\pi^* \pi_*(M)_y \to M_y \quad \text{and} \quad \pi^* \pi_*(\mathfrak{g}'_Y)_y \to \mathfrak{g}'_y$$

are surjective when  $y \in \pi^{-1}(S)$ . Then if M is point-wise smooth along  $\mathfrak{g}'_Y$  in  $Y \setminus \pi^{-1}(S)$ , it follows that it is point-wise smooth in Y.

Letting Y = X and  $\pi = \text{id}$  we obtain a non-relative version: Let M be a coherent  $\mathfrak{g}_X$ -module, torsion free over  $\mathcal{O}_X$ , and  $\Gamma$  a closed subset such that  $\operatorname{codim}_X \Gamma \geq 2$ . Then if M is point-wise smooth along a sub-algebroid  $\mathfrak{g}'_X \subseteq \mathfrak{g}_X$  at each point in  $X \setminus \Gamma$  it follows that M is smooth along  $\mathfrak{g}'_X$  in all of X (Cor. 5.2.2); Corollary 5.2.3 is a similar result for pure modules. Moreover, if M is point-wise smooth along  $\mathfrak{g}'_X$  in the complement of a divisor  $S \subset X$ , then the points in S where M is smooth form a closed set, so one gets as an added dividend that it suffices to check smoothness in one point in every irreducible component of the non-singular locus of S (Cor. 5.2.4). This implies Deligne's theorem that a  $T_X$ -module, coherent and torsion free over  $\mathcal{O}_X(*S)$ , has regular singularities along S (in the sense of pull-backs to curves) if this holds at all points in a dense subset of the non-singular locus of S. Section 6: Theorem 6.1.1 is a useful smoothness criterion, and follows from a straightforward application of Gabber's integrability theorem. Theorem 6.1.2 is a curve test applicable to torsion free modules for any pair of Lie algebroids  $\mathfrak{g}'_X \subset \mathfrak{g}_X$ . The remaining part of the section is concerned with  $\mathfrak{g}_X$ -modules with regular singularities, where  $\mathfrak{g}'_X$  is determined by a sub-space in X. When M is a pure coherent  $\mathfrak{g}_X$ -module with support Vwe say that it has regular singularities along a divisor  $S \subset V \subset X$  if it is smooth along  $\mathfrak{g}_X(I) \cap \mathfrak{g}_X(J)$ ; here I and J are the ideals of S and V, and  $\mathfrak{g}_X(I) = \alpha^{-1}(T_X(I))$ . We refer to Section 6.3 for the general case when the module is not pure. From now assume that the spaces X and Y are *non*singular, and the Lie algebroids  $\mathfrak{g}_X$  and  $\mathfrak{g}_Y$  are locally free and transitive. Let  $D^b_{rs}(\mathfrak{g}_X)$  be the derived category of bounded complexes of  $\mathfrak{g}_X$ -modules  $M^{\bullet}$  whose homology  $H^{\bullet}(M^{\bullet})$  is a  $\mathfrak{g}_X$ -module with regular singularities.

We have  $D_{rs}^{b}(\mathfrak{g}_{X}) \subset D_{hol}^{b}(\mathfrak{g}_{X})(Prop. 6.3.10)$  where  $D_{hol}^{b}(\mathfrak{g}_{X})$  is the category of complexes with holonomic homology, and if  $S \subset X$  is a subspace, then  $R\Gamma_{[S]}(M^{\bullet}), M(*S)^{\bullet} \in D_{rs}^{b}(\mathfrak{g}_{X})$  if and only if  $M \in D_{rs}^{b}(\mathfrak{g}_{X})$ (*Th.* 6.3.11); this corresponds to a theorem by Bernstein.

Let  $\pi: Y \to X$  be a morphism of spaces. The inverse image functor preserves the property of having regular singularities,  $\pi^!(D^b_{rs}(\mathfrak{g}_X)) \subset D^b_{rs}(\mathfrak{g}_Y)$ (*Th.* 6.3.12), and if  $\pi$  is proper, the direct image functor has the same good behaviour (*Th.* 6.3.13)

$$\pi_+(D^b_{rs}(\mathfrak{g}_Y)) \subset D^b_{rs}(\mathfrak{g}_Y).$$

Let  $D^b_{crs}(\mathfrak{g}_X) \subset D^b_{coh}(\mathfrak{g}_X)$  be the sub-category of complexes  $M^{\bullet}$  such that the pull back  $\pi^+(M^{\bullet}) \in D^b_{rs}(\mathfrak{g}_C)$  when  $\pi : C \to X$  is an embedding of a curve (curve regular complexes). We have (*Th.* 6.3.15)

$$D_{rs}^{o}(\mathfrak{g}_X) = D_{crs}^{o}(\mathfrak{g}_X).$$

As a consequence any sub-quotient of the homology of a complex in  $D^b_{crs}(\mathfrak{g}_X)$  is curve regular (*Cor.* 6.3.18). That  $D^b_{crs}(T_X)$  has this property is known but the previous proofs are quite hard; see [BGK<sup>+</sup>87] and [Bjö93].

In the remaining part we study the sub-category  $\overline{D}_{rs}^{b}(\mathfrak{g}_{X}) \subset D_{rs}^{b}(\mathfrak{g}_{X})$  of completely regular complexes on a quasi-projective algebraic manifold X.

A locally free and transitive Lie algebroid  $(\bar{X}, \mathfrak{g}_{\bar{X}}, \bar{\alpha})$  is a *completion* of a Lie algebroid  $(X, \mathfrak{g}_X, \alpha)$  when there exists an open embedding  $j : X \to \bar{X}$ , where  $\bar{X}$  is projective, such that  $\mathfrak{g}_X = j^*(\mathfrak{g}_{\bar{X}})$  and  $\alpha = j^*(\bar{\alpha}) : \mathfrak{g}_X \to T_X$ . We say that  $M^{\bullet} \in \bar{D}^b_{rs}(\mathfrak{g}_X)$  if there exists such a completion so that  $j_+(M^{\bullet}) \in \bar{D}^b_{rs}(\mathfrak{g}_{\bar{X}})$ . This definition is intrinsic, independent of the choice of completion (*Lem.* 6.4.4), and  $\bar{D}^b_{rs}(T_X)$  coincides with the category of regular holonomic  $T_X$ -modules as defined in [BGK<sup>+</sup>87] (*Cor.* 6.4.7). We have  $\pi^!(\bar{D}^b_{rs}(\mathfrak{g}_X)) \subset \bar{D}^b_{rs}(\mathfrak{g}_Y)$  (*Cor.* 6.4.5). Conversely, if  $\pi(Y)$  contains the support of  $M \in \bar{D}^b_{coh}(\mathfrak{g}_X)$  and  $\pi^!(M) \in \bar{D}^b_{rs}(\mathfrak{g}_Y)$ , then  $M \in \bar{D}^b_{rs}(\mathfrak{g}_X)$ (*Prop.* 6.4.8), a result that can be used to prove that certain equivariant  $\mathfrak{g}_X$ -modules are completely regular. That

$$\pi_+(\bar{D}^b_{rs}(\mathfrak{g}_Y)) \subset \bar{D}^b_{rs}(\mathfrak{g}_X)$$

follows from Theorem 6.3.13 when  $\pi$  is proper, and it holds more generally when  $\pi$  can be factorized  $\pi = p \circ j$  where j is a completion and p is proper (*Prop.* 6.4.6). Let us compare to the proof of [BGK+87, VII, Th. 12.2], that  $\pi_+(\bar{D}^b_{crs}(T_Y)) \subset \bar{D}^b_{crs}(T_Y)$ . This proof is inspired by Deligne's proof [Del70] that the Gauss-Manin connection has regular singularities; one first has to prepare by constructing a generating class of objects, the standard modules, and one considers only completions  $Y \to \bar{Y}$  such that  $\bar{Y} \setminus Y$  is a divisor with normal crossing singularities (regular completions), which exist by Hironaka's theorem. Our proof is also based on Deligne's proof, but otherwise quite different. We do not need that fact that standard modules form a generating class, and our use of Hironaka's theorem is instead for taking care of modules whose support is not non-singular in codimension 1; one may also note that it is unnecessary to only consider regular completions.

I want to thank Rikard Bögvad for useful suggestions to improve the disposition of this work.

#### 2. Lie Algebroids

2.1. **Definitions.** By a *variety* we shall mean a separated scheme locally of finite type over an algebraically closed field k of characteristic 0. A *complex analytic space* is a separated complex analytic space  $(k = \mathbf{C})$ . By *space* we mean either a variety or a complex analytic space.

Remark 2.1.1. Our terminology is a little unconventional since a variety usually is an integral separated scheme of finite type over k. Note also that by Lefshetz' principle one can assume that all varieties of finite type are defined over  $k = \mathbf{C}$ , but not invoking transcendental methods to prove algebraic results we prefer also in such cases to work over a specific algebraically closed field k of characteristic 0.

Let  $\operatorname{coh}(\mathcal{O}_X) \subset \operatorname{Mod}(\mathcal{O}_X)$  be the subcategory of coherent sheaf of modules over the structure sheaf  $\mathcal{O}_X$  on a space X; we abbreviate sheaf of rings and sheaf of modules by ring and module. The ring  $\mathcal{O}_X$  and its  $\mathcal{O}_X$ -module of k-linear derivations  $T_X$  both belong to  $\operatorname{coh}(\mathcal{O}_X)$ . Let M be an  $\mathcal{O}_X$ -module. Then  $M_x$  is the stalk at a point x in X, and if  $\Omega$  is an open subset of X the vector space of sections over  $\Omega$  is  $\Gamma(\Omega, M)$  or  $M(\Omega)$ ; the restriction of M to  $\Omega$  is  $M_{\Omega}$ . When we say that m is a vector in M we mean that m is a section that is defined in some open subset of X. Recall that a space  $(\mathcal{O}_X, X)$  is integral when the rings  $\mathcal{O}_X(\Omega)$  are integral domains, and this is the same as  $(\mathcal{O}_X, X)$  being irreducible and reduced. For a locally principal (Weil) divisor  $S \subset X, \mathcal{O}_X(*S)$  is the coherent ring defined locally by  $\mathcal{O}_X[1/f]$  when f is a local generator of S, and we put  $M(*S) = \mathcal{O}_X(*S) \otimes_{\mathcal{O}_X} M$ . Let  $S \subset X$ be a closed subspace (algebraic or complex analytic) and  $j: X \setminus S \to X$  the canonical open immersion. We let  $\Gamma_S(M) \subseteq M$  be the maximal submodule whose support belongs to S and  $\Gamma_{[S]}(M) = H^0_{[S]}(M) = \{m \in M : I^n_S \cdot m = 0\},\$ where  $I_S$  is the ideal of S;  $R\Gamma_S(\cdot)$  and  $R\Gamma_{[S]}(\cdot)$  are the derived versions of  $\Gamma_{S}(\cdot)$  and  $\Gamma_{[S]}(\cdot)$ , defining functors on the derived category of complexes of  $\mathcal{O}_X$ -modules  $D(\mathcal{O}_X) := D(\operatorname{Mod}(\mathcal{O}_X))$ . The local cohomology modules of  $M^{\bullet} \in D(\mathcal{O}_X)$  are  $H^i_S(M^{\bullet}) = R^i \Gamma_S(M^{\bullet})$  and  $H^i_{[S]}(M^{\bullet}) = R^i \Gamma_{[S]}(M^{\bullet})$ , and there exist distinguished triangles triangles<sup>1</sup> in  $D(\mathcal{O}_X)$ 

(2.1.1) 
$$R\Gamma_{S}(M^{\bullet}) \to M^{\bullet} \to Rj_{*}j^{*}(M^{\bullet}) \to,$$
$$R\Gamma_{[S]}(M^{\bullet}) \to M^{\bullet} \to M^{\bullet}(*S) \to,$$

which coincide when X is a scheme.

The work of Beilinson and Bernstein [BB93] contains a succinct treatment of Lie algebroids and serves as an important source for this section; this section can be regarded as a complement to some of the most basic material in *loc. cit.*.

**Definition 2.1.2.** A Lie algebroid is a triple  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha)$ , where:

- (1)  $\mathfrak{g}_X$  is a Lie algebra over k on a space  $(\mathcal{O}_X, X)$ ;
- (2)  $\mathfrak{g}_X$  is a coherent  $\mathcal{O}_X$ -module;
- (3)  $\alpha$  is a homomorphism of sheaves

$$\alpha:\mathfrak{g}_X\to T_X$$

such that  $\alpha([\delta_1, \delta_2]) = [\alpha(\delta_1), \alpha(\delta_2)]$  and  $\alpha(f\delta) = f\alpha(\delta)$  for  $f \in \mathcal{O}_X$ ,  $\delta_1, \delta_2 \in \mathfrak{g}_X$ . ( $\alpha$  is a homomorphism for the structures in 1 and 2). We require the compatibility  $[\delta_1, f\delta_2] = \alpha(\delta_1)(f)\delta_2 + f[\delta_1, \delta_2], f \in \mathcal{O}_X$ ,  $\delta_1, \delta_2 \in \mathfrak{g}_X$ .

Let  $\operatorname{Lie}_X$  be the category of Lie algebroids on X. We say that  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha) \in \operatorname{Lie}_X$  is *transitive* if  $\alpha$  is surjective; if  $\alpha = 0$  then  $\mathfrak{g}_X$  is a sheaf of Lie algebras over  $\mathcal{O}_X$ .

A homomorphism of Lie algebroids  $\phi : (\mathfrak{g}_X, \alpha) \to (\mathfrak{a}_X, \beta)$  (on the same space X) is a homomorphism of  $\mathcal{O}_X$ -modules and k-Lie algebras  $\phi : \mathfrak{g}_X \to \mathfrak{a}_X$  such that  $\alpha = \beta \circ \phi$ . Of course, we often abbreviate a Lie algebroid  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha)$  to  $\mathfrak{g}_X$ . It should be obvious what is meant by an ideal of a Lie algebroid; for instance  $\operatorname{Ker}(\phi)$  is an ideal if  $\phi$  is a homomorphism of Lie algebroids and the sub-sheaf  $\mathfrak{b} = \operatorname{Ker}(\alpha)$  is an ideal which is moreover a coherent  $\mathcal{O}_X$ -Lie algebra.

Let  $\mathfrak{a}_X$  be a subalgebroid of a Lie algebroid  $\mathfrak{g}_X$ . The *normalizer* of  $\mathfrak{a}_X$  in  $\mathfrak{g}_X$  is defined by the presheaf  $\mathfrak{n}_X(U)$  that is formed by the  $\mathcal{O}_X(U)$ -submodule of  $\mathfrak{g}_X(U)$  that is generated by the k-subspace

$$\{\delta \in \mathfrak{g}_X(U) : [\delta, \mathfrak{a}_X(U)] \subset \mathfrak{a}_X(U)\}$$

where  $U \subset X$  is an open subset. Clearly, the presheaf  $\mathfrak{n}_X$  is a sheaf satisfying all conditions in Definition 2.1.2 except possibly (2). For example, let  $X = \mathbf{C}^1$  be the complex line with coordinate function  $t \in \mathcal{O}_{\mathbf{C}^1}(\mathbf{C}^1)$ ,  $\mathfrak{a}_X = \mathcal{O}_{\mathbf{C}^1}t\partial_t$ , and  $\mathfrak{g}_X = T_{\mathbf{C}^1} = \mathcal{O}_{\mathbf{C}^1}\partial_t$ . Then  $\mathfrak{n}_{\mathbf{C}^1} \cong j_!(T_{\mathbf{C}^1\setminus 0}) \notin \operatorname{coh}(\mathcal{O}_X)$ , the extension by 0 of  $T_{\mathbf{C}^1\setminus 0}$  from  $\mathbf{C}^1\setminus 0$  to  $\mathbf{C}^1$ .

We put a condition on the  $\mathfrak{a}_X$ -module  $N := \mathfrak{g}_X/\mathfrak{a}_X$  that ensures that  $\mathfrak{n}_X$  be coherent over  $\mathcal{O}_X$ .

**Lemma 2.1.3.** (1) If the action of  $\mathfrak{a}_X$  on  $\mathfrak{g}_X$  is  $\mathcal{O}_X$ -linear, i.e.  $[a, \phi\delta] = \phi[a, \delta]$  when  $a \in \mathfrak{a}_X, \delta \in \mathfrak{g}_X, \phi \in \mathcal{O}_X$ , then  $\mathfrak{n}_X$  is a sub-Lie algebroid.

(2) Assume that each point in X has a neighbourhood  $U \subseteq X$  such that  $\mathfrak{a}_U$  is generated by sections  $\{a_i\}_{i=1}^r \subset \mathfrak{a}_U(U)$  with the property: the

<sup>&</sup>lt;sup>1</sup>See [BBD82] for an account of triangulated categories.

 $\mathcal{O}_U$ -module  $\mathcal{O}_U N^{a_i}$  that is generated by the subsheaves of k-vector spaces

$$N^{a_i} := \{n \in N : [a_i, n] = 0\} \subset N_U$$

is locally generated by the sections of  $N^{a_i}$ . Then  $\mathfrak{n}_X$  is a sub-Lie algebroid of  $\mathfrak{g}_X$ .

(1) implies, not surprisingly, that the normalizer is always coherent when  $\mathfrak{g}_X$  is a coherent  $\mathcal{O}_X$ -Lie algebra.

*Proof.* (2): We need to prove that  $\mathfrak{n}_X$  is coherent and since this is a local problem we may assume that U = X, so the  $a_i$  are global generators of  $\mathfrak{a}_X$  and  $\mathcal{O}_X N^{a_i}$  is generated by  $N^{a_i}(X)$ . Since

$$\{\delta \in \mathfrak{g}_X : [\delta, a_i] \in \mathfrak{a}_X, i = 1, \dots, k\} = \bigcap_{i=1}^k \{\delta \in \mathfrak{g}_X : [\delta, a_i] \subset \mathfrak{a}_X\}$$

it follows that  $\mathfrak{n}_X = \bigcap_{i=1}^k \mathfrak{n}_X^{(i)}$  where  $\mathfrak{n}_X^{(i)}$  is the  $\mathcal{O}_X$ -submodule of  $\mathfrak{g}_X$  that is generated by  $\{\delta \in \mathfrak{g}_X : [\delta, a_i] \subset \mathfrak{a}_X\}$ . Hence, by [GD71, 0. §5, Cor. 5.3.6] it suffices to see that  $\mathfrak{n}_X^{(i)}$  is coherent. As  $\mathfrak{a}_X, \mathfrak{g}_X \in \operatorname{coh}(\mathcal{O}_X)$  [loc. cit., Prop. 5.3.2] implies that  $N = \mathfrak{g}_X/\mathfrak{a}_X \in \operatorname{coh}(\mathcal{O}_X)$  and if  $\overline{\mathfrak{n}}_X^{(i)}$  is the image of  $\mathfrak{n}_X^{(i)}$  in the canonical morphism  $\mathfrak{g}_X \to N$ , since  $\operatorname{Ker}(\mathfrak{g}_X \to N) = \mathfrak{a}_X \in \operatorname{coh}(\mathcal{O}_X)$ , it suffices to see that  $\overline{\mathfrak{n}}_X^{(i)} \in \operatorname{coh}(\mathcal{O}_X)$ . But since  $\overline{\mathfrak{n}}_X^{(i)} = \mathcal{O}_X N^{a_i}$  is generated by its sections  $N^{a_i}(X) \subset \overline{\mathfrak{n}}^{(i)}(X)$  and  $\mathcal{O}_X$  is noetherian (*Th.* 2.3.2) this implies that  $\overline{\mathfrak{n}}_X^{(i)} \in \operatorname{coh}(\mathcal{O}_X)$ .

(1): Again it suffices to prove that  $\bar{\mathfrak{n}}_X^{(i)} \in \operatorname{coh}(\mathcal{O}_X)$ . Since the problem is local on X and  $\mathfrak{g}_X$  is coherent we may assume that  $\mathfrak{g}_X$  has a finite set of global generators  $\{\delta_i\} \subset \Gamma(X, \mathfrak{g}_X)$ . Let  $\mathfrak{b}_X$  be the coherent submodule of N that is generated by the image of  $[\delta_i, a]$  in N(X). We have

$$\mathfrak{n}_x^{(i)} = \{\sum_i \phi_i \delta_i \in \mathfrak{g}_x : \sum \phi_i[a, \delta_i] \in \mathfrak{a}_x\}$$

showing that the image  $\bar{\mathfrak{n}}_X^{(i)} = \operatorname{Ann}(\mathfrak{b}_X) \in \operatorname{coh}(\mathcal{O}_X)$  [GD71, 0. 5.3.10].  $\Box$ 

A Lie algebroid  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha)$  is affine if X is an affine scheme over a field k; alternatively we denote it by  $(A, \mathfrak{g}_A, \alpha)$  when  $X = \operatorname{Spec} A$ .

A connection on  $\mathfrak{g}_X$  is an  $\mathcal{O}_X$ -homomorphism  $\nabla : T_X \to \mathfrak{g}_X$  such that  $\alpha \circ \nabla = \operatorname{id}_{T_X}$ . The curvature of  $\nabla$  is the  $\mathcal{O}_X$ -linear homomorphism  $R_{\nabla} :$   $T_X \wedge T_X \to \mathfrak{b}, \, \delta \wedge \eta \mapsto [\nabla(\delta), \nabla(\eta)] - \nabla([\delta, \eta])$ . The connection is integrable if  $R_{\nabla} = 0$ , hence an integrable connection is the same as a homomorphism  $T_X \to \mathfrak{g}_X$ . The connections on  $\mathfrak{g}_X$  form a sheaf in an obvious way, defining a torsor over the group  $Hom_{\mathcal{O}_X}(T_X, \mathfrak{b})$ .

*Example* 2.1.4. Assume that a Lie algebra  $\mathfrak{g}$  over k acts on a space X by a Lie homomorphism

$$\alpha_0:\mathfrak{g}\to T_X.$$

Then  $\mathfrak{g}_X = \mathcal{O}_X \otimes_k \mathfrak{g}$  forms a Lie algebroid  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha)$  defining  $\alpha : \mathfrak{g}_X \to T_X$  by  $\alpha = \mathrm{id}_{\mathcal{O}_X} \otimes \alpha_0$  and the bracket

 $[f \otimes_k \delta_1, g \otimes_k \delta_2] = fg[\delta_1, \delta_2] + f\alpha_0(\delta_1)(g)\delta_2 - g\alpha_0(\delta_2)(g)\delta_1.$ 

If  $\alpha_0$  induces surjective mappings on the tangent spaces  $T_x/\mathfrak{m}_x T_x$ , where  $\mathfrak{m}_x$  is the maximal ideal of a point  $x \in X$ , then  $\mathfrak{g}_X$  locally has a connection.

If  $T_X$  is generated by its sections  $T_X(X)$ , then  $\mathfrak{g}_X$  has a globally defined connection.

Suppose that  $\mathfrak{g}_X$  is a Lie algebroid which moreover is a right  $\mathcal{O}_X$ -module such that  $\delta \cdot \phi = \alpha(\delta)(\phi)\delta + \phi\delta$ , when  $\phi \in \mathcal{O}_X$  and  $\delta \in \mathfrak{g}_X$ ; next section contains an example. One can then define the *opposite* algebroid  $\mathfrak{g}_X^o$  of  $\mathfrak{g}_X$ . As sheaf of k-algebras  $\mathfrak{g}_X^o$  is the same as  $\mathfrak{g}_X$ . But if  $[\cdot, \cdot]^\circ$  is the Lie bracket on  $\mathfrak{g}_X^o$  and  $\circ$  is the multiplication between  $\mathcal{O}_X$  and  $\mathfrak{g}_X^o$  one has

$$[\delta, \eta]^{\circ} = [\eta, \delta],$$
  
$$\phi \circ \delta = \delta \phi, \quad \delta \circ \phi = \phi \delta,$$

for  $\delta, \eta \in \mathfrak{g}_X$  and  $\phi \in \mathcal{O}_X$ . This is again a Lie algebroid  $(\mathcal{O}_X, \mathfrak{g}_X^o, \alpha^o)$  if one defines  $\alpha^o : \mathfrak{g}_X^o \to T_X$  by  $\delta \mapsto -\alpha(\delta)$ .

2.2. Linear Lie algebroids. Let M be an  $\mathcal{O}_X$ -module and  $\mathfrak{gl}_k(M) := End_k(M)$  be its sheaf of k-Lie algebras. We have an obvious k-linear mapping

$$i: \mathcal{O}_X \to \mathfrak{gl}_k(M),$$

defining a structure of  $\mathcal{O}_X$ -bi-module on  $\mathfrak{gl}_k(M)$ , by  $(f \cdot \phi)(m) = f\phi(m)$  and  $(\phi \cdot f)(m) = \phi(fm), f \in \mathcal{O}_X, m \in M$ . Let  $\mathfrak{gl}_{\mathcal{O}_X}(M) \subseteq \mathfrak{gl}_k(M)$  be the  $\mathcal{O}_X$ -submodule of  $\mathcal{O}_X$ -linear homomorphisms  $M \to M$ . Clearly  $\mathfrak{gl}_{\mathcal{O}_X}(M)$  is a Lie sub-algebra which commutes with the commutative sub-algebra  $i(\mathcal{O}_X)$ . Define  $\mathcal{D}^1_X(M) \subseteq \mathfrak{gl}_k(M)$  the first order matrix differential operators on M as the sub-sheaf of k-linear mappings  $\delta$  such that  $[\delta, \mathfrak{gl}_{\mathcal{O}_X}(M)] \subseteq \mathfrak{gl}_{\mathcal{O}_X}(M)$ . Evidently  $\mathcal{D}^1_X(M)$  is a Lie algebra over k.

We temporarily forget the condition (2) in Definition 2.1.2, so Lie algebroids are not necessarily coherent. Now the Lie algebra  $\mathcal{D}^1_X(M)$  is not provided with a natural mapping  $\alpha$  to  $T_X$ ; its  $\mathcal{O}_X$ -submodule

(2.2.1) 
$$\mathfrak{c}_X(M) := \{ \delta \in \mathcal{D}^1_X(M) : [\delta, i(\mathcal{O}_X)] \subseteq i(\mathcal{O}_X) \}$$

also has a Lie bracket, but again there is no mapping  $\alpha$ . However, if M is a faithful  $\mathcal{O}_X$ -module, i.e. the mapping i is injective, we identify  $\mathcal{O}_X$  with a sub-ring of  $\mathfrak{gl}_{\mathcal{O}_X}(M)$ , and since  $[\delta, \phi] \cdot m = \delta \cdot (\phi m) - (\phi \delta) \cdot m$ , for  $\delta \in \mathfrak{c}_X(M)$ ,  $\phi \in \mathcal{O}_X, m \in M$ , we get a well-defined homomorphism  $\alpha : \mathfrak{c}_X(M) \to T_X$ ,  $\delta \mapsto [\delta, \phi]$ . Therefore  $(\mathcal{O}_X, \mathfrak{c}_X(M), \alpha) \in \text{Lie}_X$ . It is called the *linear Lie* algebroid of the faithful  $\mathcal{O}_X$ -module M.  $\mathfrak{c}_X(M)$  can also be described as the set of pairs  $(\delta, \partial)$  where  $\partial \in T_X$  and  $\delta$  is a lift of  $\partial$  to an action on Msuch that  $\delta(\phi m) = \partial(\phi)m + \phi\delta(m)$ . The homomorphism  $\alpha : \mathfrak{c}_X(M) \to T_X$  is  $(\delta, \partial) \mapsto \partial$ , and the bracket is  $[(\delta, \partial), (\delta', \partial')] = ([\delta, \delta'], [\partial, \partial'])$ . If M is locally free, then  $\mathfrak{c}_X(M)$  is locally isomorphic as Lie algebroid to  $\mathfrak{gl}_{\mathcal{O}_X}(M) \oplus T_X$ where the action of  $T_X$  is defined using a local isomorphism  $M_\Omega \cong \mathcal{O}_\Omega^n$  over an open subset  $\Omega \subset X$ . If X is non-singular and M is quasi-coherent with presentation

$$\oplus_I \mathcal{O}_X \xrightarrow{\phi} \oplus_J \mathcal{O}_X \twoheadrightarrow M,$$

then  $\mathfrak{c}_X(M)$  is locally isomorphic to

$$\frac{\{\delta \in \mathfrak{c}_X(\oplus_J \mathcal{O}_X) : \delta\phi(\oplus_I \mathcal{O}_X) \subseteq \phi(\oplus_I \mathcal{O}_X)\}}{\{\delta \in \mathfrak{c}_X(\oplus_J \mathcal{O}_X) : \delta(\oplus_J \mathcal{O}_X) \subseteq \phi(\oplus_I \mathcal{O}_X)\}}$$

Let I be an ideal of  $\mathcal{O}_X$ . Then  $\mathfrak{c}_X(I) = \mathcal{O}_X \oplus T_X(I)$ , where  $T_X(I)$  is the sheaf of derivations that preserve the ideal I, and if  $\mathfrak{g}_X$  is a Lie algebroid we let  $\mathfrak{g}_X(I)$  be the sub-algebroid of vectors  $\delta$  such that  $\alpha(\delta) \in T_X(I)$ .

The Lie algebroid  $\mathfrak{c}_X(M)$  is also a right  $\mathcal{O}_X$ -module, where  $\phi \in \mathcal{O}_X$  acts on  $\delta \in \mathfrak{c}_X(M)$  as  $\delta \cdot \phi = [\delta, \phi_M] + \phi_M \delta$ , and  $\phi_M \in \mathfrak{gl}_k(M)$  is the multiplication by  $\phi$  on M.

Remark 2.2.1. Let M be locally free  $\mathcal{O}_X$ -module of rank  $r < \infty$ . One may construct 'classical' Lie algebroids  $\mathfrak{g}_X$  from an alternating or symmetric bilinear form  $\langle \cdot, \cdot \rangle : M \otimes_{\mathcal{O}_X} M \to \mathcal{O}_X$ 

$$\mathfrak{g}_X = \{ \delta \in \mathfrak{c}_X(M) : \alpha(\delta) < m_1, m_2 > = < \delta \cdot m_1, m_2 > + < m_1, \delta \cdot m_2 > \}.$$

Letting det  $M = \wedge^r M$  be the determinant bundle of M there exists a homomorphism of Lie algebroids tr :  $\mathfrak{c}_X(M) \to \mathfrak{c}_X(\det M)$  defined by

$$\operatorname{tr}(\delta) \cdot m_1 \wedge \cdots \wedge m_d = (\delta \cdot m_1) \wedge \cdots \wedge m_r + \cdots + m_1 \wedge \cdots \wedge (\delta \cdot m_r),$$

where  $m_i \in M$  and  $\delta \in \mathfrak{c}_X(M)$ , and one may define the 'special' Lie algebroid by  $\mathfrak{sl}(M) := \operatorname{Ker}(\operatorname{tr}) \subseteq \operatorname{Ker}(\alpha : \mathfrak{c}_X(M) \to T_X)$ . Note that  $[\mathfrak{c}_X(M), \mathfrak{c}_X(M)] \nsubseteq \mathfrak{sl}(M)$ , which can be compared to  $[\mathfrak{gl}_{\mathcal{O}_X}(M), \mathfrak{gl}_{\mathcal{O}_X}(M)] \subset \mathfrak{sl}_{\mathcal{O}_X}(M)$ .

For applications of GAGA and checking coherence it is useful to identify  $\mathcal{D}_X^1(M)$  with an  $\mathcal{O}_X$ -module of  $\mathcal{O}_X$ -linear mappings. We recall this identification (see [Gro67, §16, Prop. 16.8.8]): Let  $\Delta : X \to X \times_k X$  be the diagonal morphism. One identifies sheaves on X with sheaves on  $X \times_k X$  whose support belongs to  $\Delta(X)$ . Let I be the ideal of  $\Delta(X)$  and put  $P_X^1 = \mathcal{O}_{X \times_k X}/I^2$ , which is regarded as an  $(\mathcal{O}_X, \mathcal{O}_X)$ -bimodule in an obvious way; similarly  $\Omega_X := I/I^2$  and  $\mathcal{O}_X$  are regarded as bimodules. One has the multiplication homomorphism  $P_X^1 \to \mathcal{O}_X$ , defined by  $\phi \otimes_k \phi' \mapsto \phi \cdot \phi', \phi, \phi' \in \mathcal{O}_X$ , and a short exact sequence of bimodules  $0 \to \Omega_X \to P_X^1 \to \mathcal{O}_X \to 0$ . Since this sequence has a canonical left  $\mathcal{O}_X$ -linear split  $d : P_X^1 \to \Omega_X, \phi \otimes \phi' \mapsto \phi \cdot d\phi'$ , putting  $\Omega_{X/k}(M) = \Omega_{X/k} \otimes_{\mathcal{O}_X} M$  and  $P_X^1(M) = P_X^1 \otimes_{\mathcal{O}_X} M$  one gets a short exact sequence of left  $\mathcal{O}_X$ -modules

(2.2.2) 
$$0 \to \Omega_{X/k}(M) \to P_X^1(M) \to M \to 0.$$

Then

$$\mathcal{D}^1_X(M) = Hom_{\mathcal{O}_X}(P^1_X(M), M),$$

where  $\delta \in \mathcal{D}^1_X(M)$  is identified with the mapping  $P^1_X(M) \to M$  defined by  $\phi \otimes \phi' \otimes m \mapsto \phi \delta(\phi' \cdot m)$ .

**Lemma 2.2.2.** (1) (X is a space)  $P_{X/k}^1, \Omega_{X/k} \in \operatorname{coh}(\mathcal{O}_X)$ . (2) If  $M, \Omega_X \in \operatorname{coh}(\mathcal{O}_X)$ , then  $\mathfrak{c}_X(M) \in \operatorname{coh}(\mathcal{O}_X)$ .

*Proof.* (1) is well-known. (2):  $\mathfrak{c}_X(M)$  is the normalizer of  $i(\mathcal{O}_X)$  in  $\mathcal{D}^1_X(M) = Hom_{\mathcal{O}_X}(P^1_X, M) \in \operatorname{coh}(\mathcal{O}_X)$ . Since the action of  $i(\mathcal{O}_X)$  on  $\mathcal{D}^1_X(M)$  is  $\mathcal{O}_X$ -linear the assertion follows from (1) in Lemma 2.1.3.  $\Box$ 

A connection on M is a k-linear mapping  $\nabla : M \to \Omega_{X/k}(M)$  such that  $\nabla(\phi m) = d(\phi)m + \phi \nabla(m)$ . Let us denote the (sheaf of) connections on M by  $\operatorname{Der}_k(M, \Omega_{X/k}(M))$ .

Lemma 2.2.3.  $\operatorname{Der}_k(M, \Omega_{X/k}(M)) = \operatorname{Hom}_{\mathcal{O}_X}(T_X, \mathfrak{c}_X(M))$ 

*Proof.* The mappings

$$\rho: \operatorname{Der}_{k}(M, \Omega(M)) \to \operatorname{Hom}_{\mathcal{O}_{X}}(T_{X}, \mathfrak{c}_{X}(M)),$$
  

$$\rho(\nabla)(\partial) \in \mathfrak{c}_{X}(M) : m \mapsto < \nabla(m), \partial >;$$
  

$$\psi: \operatorname{Hom}_{\mathcal{O}_{X}}(T_{X}, \mathfrak{c}_{X}(M)) \to \operatorname{Der}_{k}(M, \Omega(M)),$$
  

$$\psi(\phi)(m) \in \Omega(M) : \partial \mapsto \psi(\phi)(m)(\partial) = \phi(\partial)(m)$$

are mutually inverse isomorphisms.

Thus if M is a faithful  $\mathcal{O}_X$ -module then a connection on  $\mathfrak{c}_X(M)$  is the same as a connection on M.

The short exact sequence (2.2.2) defines an element

$$c(M) \in \operatorname{Ext}^{1}_{\mathcal{O}_{X}}(M, \Omega_{X}(M)),$$

called the Atiyah class of the  $\mathcal{O}_X$ -module M. Thus c(M) = 0 when there exists a globally defined connection  $\nabla \in \operatorname{Der}_k(M, \Omega_X(M))$  and if M is locally free, so connections exist locally, then  $c(M) \in \operatorname{Ext}^1_{\mathcal{O}_X}(M, \Omega_X(M)) =$  $H^1(X, \Omega_X \otimes_{\mathcal{O}_X} \mathfrak{gl}_{\mathcal{O}_X}(M))$ . Any vector in  $T_X$  defines locally a homomorphism  $\Omega_X(M) \to M$  and thus a push-out of (2.2.2), defining a homomorphism of  $\mathcal{O}_X$ -modules  $\psi: T_X \to \operatorname{Ext}^1_{\mathcal{O}_X}(M, M)$ . Applying  $\operatorname{Hom}_{\mathcal{O}_X}(\cdot, M)$  to the short exact sequence (2.2.2) we get long exact sequences

In particular:

$$\operatorname{Ker}(\psi) = \alpha(\mathfrak{c}_X(M)).$$

There is one obstruction class in  $\operatorname{Ext}^{1}_{\mathcal{O}_{X}}(\alpha(\mathfrak{c}_{X}(M)), \mathfrak{gl}_{\mathcal{O}_{X}}(M))$  for the existence of a split of the short exact sequence

$$0 \to \mathfrak{gl}_{\mathcal{O}_X}(M) \to \mathfrak{c}_X(M) \to \alpha(\mathfrak{c}_X(M)) \to 0$$

and one obstruction class in  $\operatorname{Ext}^{1}_{\mathcal{O}_{X}}(\frac{T_{X}}{\alpha(\mathfrak{c}_{X}(M))}, \alpha(\mathfrak{c}_{X}(M)))$  that  $\alpha(\mathfrak{c}_{X}(M))$  be a direct summand of  $T_{X}$ ; these two obstruction classes vanish if and only if  $\mathfrak{c}_{X}(M)$  has a connection.

2.3. Differential operators. We define differential operators as follows.

**Definition 2.3.1.** A ring of differential operators  $(\mathcal{O}_X, \mathcal{D}_X)$  is a k-algebra  $\mathcal{D}_X$  on a space  $(X, \mathcal{O}_X)$  that has:

- (1) a filtration,  $0 = \mathcal{D}^{-1} \subseteq \mathcal{D}_X^0 \subseteq \mathcal{D}_X^1 \subseteq \cdots \subseteq \mathcal{D}_X$  such that  $\mathcal{D}_X = \bigcup_n \mathcal{D}_X^n$  and the associated graded ring gr $\mathcal{D}_X := \bigoplus_{n \ge 0} \mathcal{D}_X^n / \mathcal{D}_X^{n-1}$  is commutative.
- (2) an inclusion  $i : \mathcal{O}_X \hookrightarrow \mathcal{D}_X^0$  such that  $[\mathcal{D}_X^n, i(\mathcal{O}_X)] \subseteq \mathcal{D}_X^{n-1}$ .

Let  $\text{Diff}_X$  be the category whose objects are rings of differential operators on X and the homomorphisms are required to be compatible with the filtrations. Any ring of differential operators defines a Lie algebroid

$$\mathfrak{g}_X = \{ \delta \in \mathcal{D}^1_X : [\delta, i(\mathcal{O}_X)] \subset i(\mathcal{O}_X) \}$$

(compare to (2.2.1) in Subs. 2.2) defining a functor  $Lie : \text{Diff}_X \to \text{Lie}_X$ . The functor Lie has a left adjoint  $\mathcal{D}_X(\cdot) : \text{Lie}_X \to \text{Diff}_X$ , where  $\mathcal{D}_X := \mathcal{D}_X(\mathfrak{g}_X)$  is called the enveloping ring of differential operators of  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha) \in \text{Lie}_X$ ; it is constructed similarly to the enveloping algebra of a Lie algebra.

Let  $S_X = S_X(\mathfrak{g}_X)$  be the symmetric algebra of the  $\mathcal{O}_X$ -module  $\mathfrak{g}_X$ .

In a ringed space  $(X, R_X)$  we recall that  $R_X$  is *noetherian* if the following holds: (i)  $R_X$  is coherent as left  $R_X$ -module; (ii) for each point x in X the stalk  $R_x$  is a noetherian ring; (iii) for any open subset  $\Omega$  of X any increasing family of coherent sub-modules of a coherent  $R_\Omega$ -module is locally stationary.

**Theorem 2.3.2.** (X is a space) The rings  $\mathcal{O}_X$ ,  $S_X$ ,  $gr(\mathcal{D}_X)$  and  $\mathcal{D}_X$  are noetherian.

*Proof.*  $\mathcal{O}_X$  is noetherian: When X is a variety see [GD71, Cor. 1.5.3] (it suffices that  $(X, \mathcal{O}_X)$  be locally noetherian). When X is a complex analytic space (i) is Oka's theorem, see [Gun90]; a proof of (i) and (ii) can be found in [Gun90]. (iii) is due to Serre [Ser66] and Grauert [Gra60, Th 8].

 $S_X$ ,  $\operatorname{gr}(\mathcal{D}_X)$ , and  $\mathcal{D}_X$  are noetherian: Follow the proofs in B. Kaup's article in [BGK<sup>+</sup>87]. Note that [loc. cit., Prop. 3.5] should be changed to: the canonical homomorphism  $S_X \to \operatorname{gr}(\mathcal{D}_X)$  is surjective. See also [Bjö93].

The image of the canonical inclusion  $\sigma_s : \mathfrak{g}_X \to S_X$  generates the algebra  $S_X$ , making it possible to define a k-linear Lie product

$$\{\cdot, \cdot\}: \mathcal{S}_X \otimes_k \mathcal{S}_X \to \mathcal{S}_X$$

by  $\{\sigma_s(\partial), \sigma_s(\eta)\} = \sigma_s([\partial, \eta]), \{\phi, \psi\} = 0, \{\sigma_s(\partial), \phi\} = \alpha(\partial)(\phi)$ , and inductively by the rules

- (1)  $\{a,b\} = -\{b,a\}$
- (2)  $\{a, bc\} = \{a, b\}c + \{a, c\}b$

for all  $a, b, c \in S_X$ . This defines a Poisson algebra  $(S_X, \cdot, \{\cdot, \cdot\})$ . In the same way one defines a bracket  $\{\cdot, \cdot\}$  on gr  $\mathcal{D}_X$  resulting in a Poisson algebra  $(\operatorname{gr} \mathcal{D}_X, \cdot, \{\cdot, \cdot\})$ . The canonical mapping  $\mathfrak{g}_X \to \operatorname{gr}(\mathcal{D}_X)$  lifts to a surjective homomorphism of Poisson algebras  $p: S_X \to \operatorname{gr}(\mathcal{D}_X)$ . We have the following generalization of the Poincaré-Birkhoff-Witt theorem:

**Theorem 2.3.3** ([Rin63]). If  $\mathfrak{g}_X$  is locally free over  $\mathcal{O}_X$  then p is an isomorphism.

Let M be a coherent  $\mathfrak{g}_X$ -module (see Definition 3.1.1 below),  $\Omega \subset X$ an open set and  $M_{\Omega}^0$  a coherent  $\mathcal{O}_{\Omega}$ -module generating  $M_{\Omega}$  over  $\mathfrak{g}_{\Omega}$ . Set  $M_{\Omega}^n = \mathfrak{g}_{\Omega} M_{\Omega}^{n-1} = \mathcal{D}_{\Omega}^n \cdot M_{\Omega}^0$  for  $n \geq 1$ , and

$$G(M_{\Omega}) = \oplus M_{\Omega}^n / M_{\Omega}^{n-1},$$

defining a coherent  $S_{\Omega}$ -module. When X is a variety we let  $\mathbf{Spec} S_X$  be the spectrum of  $S_X$ , and if X is a complex analytic space we use the same notation  $\mathbf{Spec} S_X$  to denote the analytic space that can be associated to  $S_X$ ; note that the  $\mathcal{O}_X$ -algebra  $S_X$  is locally finitely generated. There is a morphism of spaces  $p : \mathbf{Spec} S_X \to X$  such that  $p^*(S_X) = \mathcal{O}_{\mathbf{Spec} S_X}$ . The support  $\mathrm{SS}(M_{\Omega}) \subseteq \mathbf{Spec} S_{\Omega}$  of  $p^*(G(M_{\Omega}))$  is independent of the choice of  $M_{\Omega}^0$  (see e.g. [Bjö93]), and therefore the construction globalises defining a conic set  $SS(M_X) \subset Spec S_X$ , which is a subspace since  $p^*(G(M_\Omega))$  is coherent over  $\mathcal{O}_{Spec S_X}$ . The set  $SS(M_X)$  is the *singular support* of M, and the ideal J(M) of  $SS(M_X)$  is the characteristic ideal of M.

**Theorem 2.3.4** ([Gab81, Th. I]).

$$\{J(M), J(M)\} \subseteq J(M).$$

Let  $p: \operatorname{\mathbf{Spec}} S_X \to X$  be the canonical projection and  $i: X \to \operatorname{\mathbf{Spec}} S_X$ the canonical section. It is easy to see that  $M_x$  is of finite type over  $\mathcal{O}_x$ if and only if  $\operatorname{SS}(M_x) \subset i(x)$ , i.e. the ideal  $\operatorname{J}(M_x) = \operatorname{J}(M)_x$  of  $\operatorname{SS}(M_x)$  is  $(\sigma(\mathfrak{g}_x)) \subseteq \operatorname{S}(\mathfrak{g}_x)$ . Set  $\operatorname{SS}(M) = \operatorname{SS}(M) \setminus i(X)$ .

**Definition 2.3.5.** The strong support of a  $\mathfrak{g}_X$ -module M is the set

s-supp  $M = \{x \in X : M_x \text{ is not of finite type over } \mathcal{O}_x\} = p(\dot{SS}(M)).$ 

We shall need the following lemma, whose proof is well-known.

**Lemma 2.3.6.** (X is a space) Let M be a coherent  $\mathfrak{g}_X$ -module. Then s-supp M is a subspace of X.

*Proof.* We prove this only when X is a complex analytic space. The set SS(M) = supp G(M) is a conic analytic set, hence  $\dot{SS}(M)$  is a conic analytic set. Let  $\mathbf{Proj}(S_X)$  be the projective space of  $\mathbf{Spec}(S_X)$  (the analytic space that is associated with the graded  $\mathcal{O}_X$ -algebra  $S_X$ ), with canonical projection  $p_1 : \mathbf{Proj}(S_X) \to X$ . So we have an open inclusion mapping  $\phi : \mathbf{Spec}(S_X) \setminus i(X) \to \mathbf{Proj}(S_X)$ , satisfying  $p_1 \circ \phi = p'$ , where p' is the restriction of p to  $\mathbf{Spec}(S_X) \setminus i(X)$ . That  $\dot{SS}$  is conic means that  $\dot{SS}(M) = \phi^{-1}(V)$  where V is a complex analytic space in  $\mathbf{Proj}(S_X)$ . Hence

$$s-\operatorname{supp}(M) = p'(SS(M)) = p_1(V),$$

so the result follows from Remmert's proper mapping theorem [Gun90, Ch. N Th. 1].  $\Box$ 

## 2.4. Homomorphisms.

2.4.1. Remark on the tangent mapping. Let  $\pi : Y \to X$  be a morphism of spaces. There is then the canonical exact sequence  $\pi^*(\Omega_{X/k}) \to \Omega_Y \to \Omega_{Y/X} \to 0$ , inducing a homomorphism

$$T_Y = Hom_{\mathcal{O}_Y}(\Omega_{Y/k}, \mathcal{O}_Y) \to Hom_{\mathcal{O}_Y}(\pi^*(\Omega_{X/k}), \mathcal{O}_Y).$$

Consider also the canonical homomorphism

$$\psi: \pi^*(T_X) = \pi^*(Hom_{\mathcal{O}_X}(\Omega_{X/k}, \mathcal{O}_X)) \to Hom_{\mathcal{O}_Y}(\pi^*(\Omega_{X/k}), \mathcal{O}_Y).$$

If the induced homomorphism between stalks at any point  $x \in X$  is an isomorphism, then  $\psi$  is an isomorphism. As  $\Omega_{X/k} \in \operatorname{coh}(\mathcal{O}_X)$  we have

$$Hom_{\mathcal{O}_X}(\Omega_{X/k}, \mathcal{O}_X)_x = Hom_{\mathcal{O}_x}(\Omega_{\mathcal{O}_x/k}, \mathcal{O}_x),$$

so the condition is that the canonical morphism

 $\mathcal{O}_y \otimes_{\mathcal{O}_x} \operatorname{Hom}_{\mathcal{O}_x/k}, \mathcal{O}_x) \to \operatorname{Hom}_{\mathcal{O}_y}(\mathcal{O}_y \otimes_{\mathcal{O}_x} \Omega_{\mathcal{O}_x/k}, \mathcal{O}_y)$ 

is an isomorphism. Here  $\mathcal{O}_x \to \mathcal{O}_y$  is the homomorphism of local rings defined by  $\pi$ . Thus  $\psi$  is an isomorphism if either X is non-singular, so  $\Omega_{\mathcal{O}_x/k}$  is free of finite rank over  $\mathcal{O}_x$ , or if  $\pi$  is flat (to see this use also the fact that  $\Omega_{\mathcal{O}_x/k}$  is finitely presented over  $\mathcal{O}_x$ ). When  $\psi$  is an isomorphism there exists a canonical homomorphism

$$d\pi: T_Y \to \pi^*(T_X).$$

Also when  $\psi$  is not an isomorphism we shall abuse the notation in the following by writing  $\pi^*(T_X)$  when we actually mean  $Hom_{\mathcal{O}_Y}(\pi^*(\Omega_{X/k}), \mathcal{O}_Y)$ . Thus the homomorphism  $d\pi$  is always defined.

2.4.2. Homomorphisms of Lie algebroids. The definition of homomorphisms  $\mathfrak{g}_Y \to \mathfrak{g}_X$  between Lie algebroids on different spaces is slightly involved, because  $\pi^*(\mathfrak{g}_X) = \mathcal{O}_Y \otimes_{\pi^{-1}(\mathcal{O}_X)} \pi^{-1}(\mathfrak{g}_X)$  does not in general have a natural structure of Lie algebroid.

Remark 2.4.3. The  $\mathcal{O}_Y$ -module  $\pi^*(\mathfrak{g}_X)$  has a structure of Lie algebroid when there exists a homomorphism of Lie algebroids over  $\pi^{-1}(\mathcal{O}_X), \psi : \pi^{-1}(T_X) \to T_Y$ , defining the Lie bracket by

$$\begin{bmatrix} \phi_1 \otimes \partial_1, \phi_2 \otimes \partial_2 \end{bmatrix} := \phi_1 \phi_2 \otimes \begin{bmatrix} \partial_1, \partial_2 \end{bmatrix} + \phi_1 \psi(\alpha(\partial_1)(\phi_2)) \otimes \partial_2 - \\ - \phi_2 \psi(\alpha(\partial_2)(\phi_1)) \otimes \partial_1.$$

Here are two instances of this situation:

- (1)  $\pi$  is a projection, for then  $\pi^*(T_X) \subset T_Y$ .
- (2) Let  $\pi : X' \to X$  be the normalization morphism of an integral space. By [Sei66] every section of  $T_X$  has a unique lift to a section of  $T_{X'}$ , i.e. if  $U \subset X'$  is an open subset and  $\delta \in \pi^{-1}(T_X)(U)$ , then there exists a unique section  $\delta' \in T_{X'}(U)$  such that its restriction to  $\pi^{-1}(\mathcal{O}_X)(U) \subset \mathcal{O}_{X'}(U)$  coincides with  $\delta$ ; see *loc. cit.* for a more general and precise condition.

**Definition 2.4.4.** Let  $(Y, \mathfrak{g}_Y, \alpha) \in \text{Lie}_Y, (X, \mathfrak{g}_X, \beta) \in \text{Lie}_X$ . A homomorphism  $(\pi', \pi) : (Y, \mathfrak{g}_Y, \alpha) \to (X, \mathfrak{g}_X, \beta)$  is:

- (1) a morphism  $\pi: Y \to X$  of spaces;
- (2) a homomorphism of  $\mathcal{O}_Y$ -modules

$$\pi':\mathfrak{g}_Y\to\pi^*(\mathfrak{g}_X)$$

such that the following diagram commutes

and  $\pi'$  satisfies:

$$\tau'([\delta_1, \delta_2]) = \sum a_i b_j \otimes [\eta_i, \eta_j] + \sum (\alpha(\delta_1)(b_i) - \alpha(\delta_2)(a_i)) \otimes \eta_i$$

if  $\pi'(\delta_1) = \sum a_i \otimes \eta_i$ ,  $\pi'(\delta_2) = \sum b_j \otimes \eta_j$ , where  $\delta_1, \delta_2 \in \mathfrak{g}_Y$ ,  $a_i, b_i \in \mathcal{O}_Y$ ,  $\eta_i \in \mathfrak{g}_X$ . (Recall our abuse of notation in defining  $d\pi$ .)

If  $\pi = \text{id then } \pi'$  is a homomorphism of Lie algebroids on X. If  $\alpha = 0$ then  $\mathfrak{g}_Y$  is a sheaf of Lie algebras over  $\mathcal{O}_Y$  and a homomorphism  $\mathfrak{g}_Y \to \mathfrak{g}_X$ is a homomorphism of the  $\mathcal{O}_Y$ -Lie algebra  $\mathfrak{g}_Y$  onto a Lie algebra in the  $\mathcal{O}_Y$ -module  $\pi^*(\mathfrak{g}_X)$ . 2.4.5. *Pull-back.* Let  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha) \in \text{Lie}_X$ . The fibre product with respect to the  $\mathcal{O}_Y$ -linear homomorphisms  $\pi^*(\alpha) : \pi^*(\mathfrak{g}_X) \to \pi^*(T_X)$  and  $d\pi : T_Y \to \pi^*(T_X)$  is the  $\mathcal{O}_Y$ -module

$$\pi^*(\mathfrak{g}_X) \times_{\pi^*(T_X)} T_Y = \{(\delta, \partial) \in \pi^*(\mathfrak{g}_X) \times T_Y : \pi^*(\alpha)(\delta) = d\pi(\partial)\}.$$

We define a bracket on  $\pi^*(\mathfrak{g}_X) \times_{\pi^*(T_X)} T_Y$  by

$$[(\phi_1 \otimes \delta_1, \partial_1), (\phi_2 \otimes \delta_2, \partial_2)]$$
  
=  $(\partial_1(\phi_2) \otimes \delta_2 - \partial_2(\phi_1) \otimes \delta_1 + (\phi_1\phi_2) \otimes [\delta_1, \delta_2], [\partial_1, \partial_2]),$ 

for  $(\phi_i \otimes \delta_i, \partial_i) \in \pi^*(\mathfrak{g}_X) \times_{\pi^*(T_X)} T_Y$ , and letting  $\beta : \pi^*(\mathfrak{g}_X) \times_{\pi^*(T_X)} T_Y \to T_Y$ be the projection on the second factor we get a structure of Lie algebroid on  $\pi^*(\mathfrak{g}_X) \times_{\pi^*(T_X)} T_Y$ ; in that capacity we denote it by  $\pi^+(\mathfrak{g}_X)$ . Clearly, we get a functor  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha) \mapsto (\mathcal{O}_Y, \pi^+(\mathfrak{g}_X), \beta)$  from Lie<sub>X</sub> to Lie<sub>Y</sub>; being a composition of a right exact and left exact functor,  $\pi^+$  is in general neither left nor right exact. Let  $g : \pi^+(\mathfrak{g}_X) \to \pi^*(\mathfrak{g}_X)$  be the projection on the first factor. Then if  $\phi : \mathfrak{g}_Y \to \mathfrak{g}_X$  is a homomorphism of Lie algebroids there exists a unique homomorphism  $f : \mathfrak{g}_Y \to \pi^+(\mathfrak{g}_X)$  such that  $\phi = g \circ f$ ; hence the functor  $\mathfrak{g}_X \mapsto \pi^+(\mathfrak{g}_X)$  represents the functor Lie<sub>Y</sub>  $\to$  Set,  $\mathfrak{g}_Y \mapsto$  $Hom_{\text{Lie}}(\mathfrak{g}_Y, \mathfrak{g}_X)$ . We call  $\pi^+(\mathfrak{g}_X)$  the pull-back of  $\mathfrak{g}_X$  by  $\pi$ .

If  $Y \xrightarrow{f} Z \xrightarrow{g} X$  are morphisms of spaces, we have a canonical homomorphism

$$(2.4.1) \qquad c_{fg}: f^+(g^+(\mathfrak{g}_X)) \to (g \circ f)^+(\mathfrak{g}_X), \\ (h(y) \otimes_{f^{-1}(\mathcal{O}_Z)} (\phi(z) \otimes_{g^{-1}(\mathcal{O}_X)} \delta(x), \partial_z), \partial_y) \mapsto (h(y)\bar{\phi}(y) \otimes_{(g \circ f)^{-1}(\mathcal{O}_X)} \delta(x), \partial_y)$$

where  $h(y) \in \mathcal{O}_Y$ ,  $\phi(z) \in \mathcal{O}_Z$ ,  $\delta(x) \in \mathfrak{g}_X$ ,  $\partial_z \in T_Z$ ,  $\partial_y \in T_Y$ , and  $\bar{\phi}(y)$  is the image of  $\phi(z)$  in  $\mathcal{O}_Y$ . Unfortunately,  $c_{fg}$  is not always an isomorphism, so the projection functor Lie  $\rightarrow$  Sch makes Lie only into a prefibred category over the category of schemes/k (or over the category of complex analytic spaces); see [Gro71, Exp. VI] for this terminology. The following lemma is slightly more general than [BB93, Lemma 1.5.1].

Lemma 2.4.6. If either of the conditions

- (1) f is flat, or
- (2) g is smooth and Z, X are non-singular

are satisfied, then  $c_{fq}$  is an isomorphism.

*Proof.* The problem is local on Y, Z, X so we may instead consider a composition of local rings  $A \xrightarrow{g} B \xrightarrow{f} C$ , and an affine Lie algebroid  $(A, \mathfrak{g}_A)$ . Put  $\mathfrak{g}_B = g^+(\mathfrak{g}_A) = B \otimes_A \mathfrak{g}_A \times_{B \otimes_A T_A} T_B$  and  $\mathfrak{g}_C = (g \circ f)^+(\mathfrak{g}_A) = C \otimes_A \mathfrak{g}_A \times_{C \otimes_A T_A} T_C$ . Then  $f^+(g^+(\mathfrak{g}_A)) = f^+(\mathfrak{g}_B) = C \otimes_B \mathfrak{g}_B \times_{C \otimes_B T_B} T_C$ , and we have a canonical homomorphism

$$C \otimes_B \mathfrak{g}_B \times_{C \otimes_B T_B} T_C \to C \otimes_A \mathfrak{g}_A \times_{C \otimes_A T_A} T_C.$$

When necessary we here abuse the notation as in (2.4.1).

Assume (1). Then the exact sequence  $0 \to \mathfrak{g}_B \to B \otimes_A \mathfrak{g}_A \times T_B$  gives the exact sequence  $0 \to C \otimes_B \mathfrak{g}_B \to C \otimes_A \mathfrak{g}_A \times C \otimes_B T_B$ , implying that  $C \otimes_B \mathfrak{g}_B \cong C \otimes_A \mathfrak{g}_A \times_{C \otimes_A T_A} C \otimes_B T_B$ . Therefore

$$f^{+}(\mathfrak{g}_{B}) = C \otimes_{B} \mathfrak{g}_{B} \times_{C \otimes_{B} T_{B}} T_{C}$$
  

$$\cong (C \otimes_{A} \mathfrak{g}_{A} \times_{C \otimes_{A} T_{A}} C \otimes_{B} T_{B}) \times_{C \otimes_{B} T_{B}} T_{C}$$
  

$$\cong C \otimes_{A} \mathfrak{g}_{A} \times_{C \otimes_{A} T_{A}} T_{C}$$
  

$$= (g \circ f)^{+}(\mathfrak{g}_{A}).$$

Assume (2). Then we have a split short exact sequence

$$0 \to T_{B/A} \to T_B \to B \otimes_A T_A \to 0.$$

Choose a splitting, so  $T_B = T_{B/A} \oplus B \otimes_A T_A$ . Then, using the fact that  $N \times_L (M + L) = N \times M$  when N, L, M are *B*-modules with morphisms  $N \to L, M + L \to L$  and *M* is the kernel of the last morphism, one gets

$$\mathfrak{g}_B = B \otimes_A \mathfrak{g}_A \times_{B \otimes_A T_A} (T_{B/A} \oplus B \otimes AT_A) \cong T_{B/A} \oplus B \otimes_A \mathfrak{g}_A$$

where the morphism  $T_{B/A} \oplus B \otimes_A \mathfrak{g}_A \to T_B$  is  $\delta + b \otimes \eta \mapsto \delta + b \otimes \alpha(\eta) \in T_{B/A} \oplus B \otimes_A T_A = T_B$ . Therefore

$$f^{+}(\mathfrak{g}_{B}) = C \otimes_{B} (T_{B/A} \oplus B \otimes_{A} \mathfrak{g}_{A}) \times_{C \otimes_{B} T_{B}} T_{C}$$
  
$$= (C \otimes_{B} T_{B/A} \oplus C \otimes_{A} \mathfrak{g}_{A}) \times_{C \otimes_{B} T_{B/A} \oplus C \otimes_{A} T_{A}} T_{C}$$
  
$$= C \otimes_{A} \mathfrak{g}_{A} \times_{C \otimes_{A} T_{A}} T_{C}$$
  
$$= (g \circ f)^{+}(\mathfrak{g}_{A}).$$

Examples 2.4.7. (1)  $\pi^+(T_X) = T_Y$ . When  $\alpha = 0$ , then  $\pi^+(\mathfrak{g}_X) = \pi^*(\mathfrak{g}_X)$ . (2) If  $\pi: Y \to X$  is an open embedding, then  $\pi^+(\mathfrak{g}_X) = \pi^*(\mathfrak{g}_X)$ .

(3) Let  $\pi: Y \to X$  be a closed embedding to a non-singular space X, and  $I_Y$  be the ideal of  $\pi(Y)$ . Then

$$\pi^+(\mathfrak{g}_X) = \pi^+(\mathfrak{g}_X(I_Y)) = \pi^{-1}(\frac{\mathfrak{g}_X(I_Y)}{I_Y\mathfrak{g}_X}),$$

since  $T_Y = \pi^{-1}(T_X(I_Y))/I_YT_X)$ . Note that  $\pi^+(\mathfrak{g}_X)$  need not be locally free when  $\mathfrak{g}_X$  is locally free even if Y is non-singular.

- (4) If  $\alpha(\mathfrak{g}_X)$  is flat over  $\mathcal{O}_X$ , and  $i_x : \{x\} \to X$  is an inclusion of a point in X, then  $i_x^+(\mathfrak{g}_X) = i^*(\mathfrak{b}_X) = \mathfrak{b}_x/\mathfrak{m}_x\mathfrak{b}_x$ .
- (5) If X is non-singular and  $\pi$  is a submersion, i.e.  $d\pi : T_Y \to \mathcal{O}_Y \otimes_{\pi^{-1}(\mathcal{O}_X)} \pi^{-1}(T_X)$  is surjective, then there exists a canonical injection  $\pi^*(\mathfrak{g}_X) \hookrightarrow \pi^+(\mathfrak{g}_X)$ .

Clearly, if  $\mathfrak{g}_X$  is transitive then  $\pi^+(\mathfrak{g}_X)$  again is transitive, and if  $\alpha = 0$  then  $\pi^+(\mathfrak{g}_X) = \pi^*(\mathfrak{g}_X)$  is locally free. We will later need the following lemma.

**Lemma 2.4.8.** Let  $\pi : Y \to X$  be morphism of non-singular spaces and  $\mathfrak{g}_X$  a locally free and transitive Lie algebroid. Then  $\pi^+(\mathfrak{g}_X)$  is locally free and transitive.

*Proof.* Factorize  $\pi$  as  $\pi = p \circ i$  where  $i : Y \to Y \times X$  is the graph morphism and  $p : Z = Y \times X \to X$  is the projection on the second factor. Since p is smooth and Z and X are non-singular  $\pi^+(\mathfrak{g}_X) = i^+(p^+(\mathfrak{g}_X))$  (Lem. 2.4.6). One has  $\mathfrak{g}_Z := p^+(\mathfrak{g}_X) = p^*(\mathfrak{g}_X) \times_{p^*(T_X)} T_Z = p^*(\mathfrak{g}_X) \oplus q^*(T_Y)$  where  $q: Z \to Y$  is the projection to the first factor, so this is locally free and transitive.

Letting  $i : Y \to Z$  be a closed embedding and  $\mathfrak{g}_Z$  a locally free and transitive Lie algebroid it remains to see that  $i^+(\mathfrak{g}_Z)$  is locally free. There exists a short exact sequence

$$0 \to i^+(\mathfrak{g}_Z) \to i^*(\mathfrak{g}_Z) \to \frac{i^*(\mathfrak{g}_Z)}{i^+(\mathfrak{g}_Z)} \to 0$$

where  $i^*(\mathfrak{g}_Z)$  is locally free. Thus it suffices to prove that  $\frac{i^*(\mathfrak{g}_Z)}{i^+(\mathfrak{g}_Z)}$  is locally free, for then  $i^+(\mathfrak{g}_Z)$  is locally a direct summand of a free module; hence  $i^+(\mathfrak{g}_Z)$  is locally free. Y is non-singular so there exists a split short exact sequence

$$0 \to T_Y \to i^*(T_Z) \to \frac{i^*(T_Z)}{T_Y} \to 0;$$

hence  $\frac{i^*(T_Z)}{T_Y}$  is locally free (the normal bundle of Y in Z). Now by transitivity  $\frac{i^*(\mathfrak{g}_Z)}{i^+(\mathfrak{g}_Z)} \cong \frac{i^*(T_Z)}{T_Y}$ , implying the assertion.  $\Box$ 

**Lemma 2.4.9.** (X is a non-singular space) Let M be a locally free  $\mathcal{O}_X$ -module. Then  $\pi^+(\mathfrak{c}_X(M)) \cong \mathfrak{c}_Y(\pi^*(M))$ .

Proof. In Section 3.4 we give a canonical homomorphism  $\rho : \pi^+(\mathfrak{c}_X(M)) \to \mathfrak{c}_Y(\pi^*(M))$ , which we now contend is an isomorphism. This is a local question on X so we may assume that M is free, and hence that  $\mathfrak{c}_X(M) = Hom_{\mathcal{O}_X}(M, M) \oplus T_X$ . Then

$$\pi^{+}(\mathfrak{c}_{X}(M)) = \pi^{+}(Hom_{\mathcal{O}_{X}}(M, M) \oplus T_{X})$$

$$= \pi^{*}(Hom_{\mathcal{O}_{X}}(M, M) \oplus T_{X}) \times_{\pi^{*}(T_{X})} T_{Y}$$

$$= \pi^{*}(Hom_{\mathcal{O}_{X}}(M, M)) \oplus T_{Y}$$

$$= Hom_{\mathcal{O}_{Y}}(\pi^{*}(M), \pi^{*}(M)) \oplus T_{Y} \quad (M \text{ is locally free})$$

$$= \mathfrak{c}_{Y}(\pi^{*}(M)).$$

It remains to the check that these isomorphisms compose to  $\rho$ , but we omit the details.  $\Box$ 

2.5. **Picard Lie algebroids.** Let  $(\mathcal{O}_X, \mathfrak{g}'_X, \alpha^1) \in \text{Lie}_X$ , put  $\mathfrak{b} = \text{Ker}(\alpha^1)$ and let  $\chi : \mathfrak{b} \to \mathcal{O}_X$  be a homomorphism of Lie algebroids (a character of  $\mathfrak{b}$ ). The *push-out* of  $\mathfrak{g}'_X$  by  $\chi$  is the Lie algebroid  $\mathfrak{g}^{\chi}_X = \{(\phi, \delta) \in \mathcal{O}_X \oplus \mathfrak{g}'_X\}/J$ , where  $J = \{(\chi(b), -b) :\in \mathcal{O}_X \oplus \mathfrak{b}\}$ . A  $\mathfrak{g}'_X$ -module M such that the action of  $\mathfrak{b}$  is determined by  $b \cdot m = \chi(b)m, \ b \in \mathfrak{b}, \ m \in M$ , corresponds to a  $\mathfrak{g}^{\chi}_X$ -module.

**Definition 2.5.1.** A Lie algebroid  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha)$  is a Picard Lie algebroid if  $\alpha$  is surjective and  $\operatorname{Ker}(\alpha) \cong \mathcal{O}_X$ . We *identify*  $\operatorname{Ker}(\alpha) = \mathcal{O}_X$  by choosing a global central section  $1_{\mathfrak{g}_X} \in \mathfrak{g}_X$ . Let  $\operatorname{LPic}_X$  be the category of Picard Lie algebroids  $\mathfrak{g}_X$  which are locally isomorphic to  $T_X \oplus \mathcal{O}_X$  in the category of  $\mathcal{O}_X$ -modules (this is of course automatic when X is non-singular).

Thus if  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha) \in \operatorname{Lie}_X$  is transitive and  $\chi$  is a character of  $\operatorname{Ker}(\alpha)$  then the push-out  $\mathfrak{g}_X^{\chi} \in \operatorname{LPic}_X$ .

Homomorphisms of Picard Lie algebroids  $\phi : \mathfrak{g}_1 \to \mathfrak{g}_2$  are required to satisfy

$$\phi(1_{\mathfrak{g}_1}) = 1_{\mathfrak{g}_2} = 1_{\mathcal{O}_X},$$

so they are isomorphisms, i.e.  $\operatorname{LPic}_X$  is a groupoid. Note that in the category  $\operatorname{Lie}_X$  we do not require  $\phi(1_{\mathfrak{g}_1}) = 1_{g_2}$ , so  $\operatorname{LPic}_X$  is not a full sub-category of  $\operatorname{Lie}_X$ .

We shall determine the cohomology group classifying  $LPic_X$ . In general we have a short exact sequence

$$0 \to \mathcal{O}_X \to \mathfrak{g}_X \xrightarrow{\alpha} T_X \to 0$$

which is locally split as  $\mathcal{O}_X$ -modules, locally identifying  $\mathfrak{g}_X$  with  $\mathcal{O}_X \oplus T_X$ .

Local structure : Suppose that  $X_0 \subset X$  is an affine subset over which the exact sequence splits; thus putting  $\mathfrak{g} = \Gamma(X_0, \mathfrak{g}_{X_0})$  one has a short split exact sequence in the category of A-modules

$$0 \to A \to \mathfrak{g} \to \operatorname{Der}_k(A) \to 0.$$

A connection  $\nabla$  :  $\operatorname{Der}_k(A) \to \mathfrak{g}$ , determines an isomorphism  $\mathfrak{g} \cong A \oplus \operatorname{Der}_k(A)$ (as A-modules). Its curvature is the A-linear mapping  $R_{\nabla}$  :  $\operatorname{Der}_k(A) \wedge$  $\operatorname{Der}_k(A) \to A, \partial_1 \wedge \partial_2 \mapsto \eta(\partial_1 \wedge \partial_2) = [\nabla(\partial_1), \nabla(\partial_2)] - \nabla([\partial_1, \partial_2])$ , defining an element  $\eta$  in the vector space of closed 2-forms  $\Omega^{2,cl}(A)$ ; the condition that  $\eta$  be closed follows from the Jacobi identity. Hence, identifying the A-module  $\mathfrak{g}$  with  $A \oplus \operatorname{Der}_k(A)$ , the Lie bracket is expressed by

$$[(a_1,\partial_1),(a_2,\partial_2)] = (\partial_1(a_2) - \partial_2(a_1) + \eta(\partial_1 \wedge \partial_2), [\partial_1,\partial_2])$$

Conversely, any closed 2-form  $\eta$  defines a structure of Lie algebroid on  $A \oplus$ Der<sub>k</sub>(A) whose curvature 2-form is  $\eta$ . Denote by  $\mathfrak{g}^{\eta} (\cong \mathfrak{g})$  the object of LPic which corresponds to  $\eta$ .

Automorphisms of  $\mathfrak{g}$  are determined by A-linear homomorphisms  $(a, \partial) \mapsto (a+\omega(\partial), \partial)$  where  $\omega \in \Omega^{1,cl}(A)$  is a closed 1-form, hence  $\operatorname{Aut}(\mathfrak{g}) \cong \Omega^{1,cl}(A)$ . For  $\omega \in \Omega^1(A)$ , the mapping  $(a, \partial) \mapsto (a + \omega(\partial), \partial)$  defines an isomorphism of Lie algebroids  $\mathfrak{g}^{\eta} \to \mathfrak{g}^{\eta+d\omega}$  (identifying both sides with  $A \oplus \operatorname{Der}_k(A)$  as A-modules). There are no non-trivial automorphisms  $\phi : \mathfrak{g} \to \mathfrak{g}$  such that  $\phi \circ \nabla = \nabla$ , so therefore the group  $\Omega^{2,cl}(A)/d\Omega^1(A)$  classifies A-split affine Lie algebroids. Note that in the Zariski topology a closed 2-form need not be locally the differential of a 1-form; here is the typical example:

Example 2.5.2. Let  $X = \{(x, y) \in \mathbb{C}^2 : xy \neq 0\}$ , and  $A = \mathbb{C}[x, y, x^{-1}, y^{-1}]$ be its affine  $\mathbb{C}$ -algebra,  $\mathcal{O}_{X_h}$  the sheaf of holomorphic functions in the Hausdorff topology on X and  $\mathcal{O}_X$  the sheaf of regular rational functions in the Zariski topology. The 2-form  $\eta = 1/(xy)dx \wedge dy$  in X is evidently closed. Put  $\mathfrak{g}^{\eta} = A \oplus A\partial_x \oplus A\partial_y$  and define the bracket  $[\partial_x, \partial_y] = \eta(\partial_x \wedge \partial_y) = 1/(xy)$ . The canonical projection defines a Lie homomorphism  $\alpha : \mathfrak{g} \to \text{Der}_{\mathbb{C}}(A) =$  $A\partial_x + B\partial_y$ . Thus  $\mathfrak{g}$  is an affine Lie algebroid over A. Making a cut in the x-plane in  $\mathbb{C}^2$  we have  $\eta^{an} = d(\log(x)/ydy)$ , so  $\eta$  is locally exact in the Hausdorff topology. On the other hand, if  $\phi$  is a rational 1-form the residue  $\operatorname{res}_{y=0}(d\phi) = 0$  (for fixed x), so  $d\phi \neq \frac{1}{xy}$ . Therefore  $\mathcal{O}_{X_h} \otimes_A \mathfrak{g}$  is locally trivial whereas  $\mathcal{O}_X \otimes_A \mathfrak{g}$  is not locally trivial. Define the truncated complex  $\Omega^{\geq 1}(A) := \Omega^1(A) \to \Omega^{2,cl}(A)$ . An  $\Omega^{\geq 1}(A)$ -torsor is a pair  $(C, \eta)$  where C is an  $\Omega^1(A)$ -torsor and  $\eta : C \to \Omega^{2,cl}(A)$  is a mapping such that  $\eta(c + \phi) = \eta(c) + d\phi$ .

The set of connections  $\operatorname{Con}(\mathfrak{g})$  on a Lie algebroid  $(A, \mathfrak{g}, \alpha)$  is an  $\Omega^1(A)$ torsor, where the  $\Omega^1(A)$ -action is defined by  $\nabla \mapsto \nabla + \omega$ ,  $\omega \in \Omega^1(A)$ . The mapping  $\eta : \operatorname{Con}(\mathfrak{g}) \to \Omega^{2,cl}(A)$ ,  $\nabla \mapsto R_{\nabla}$ , satisfies  $\eta(\nabla + \phi) = \eta(\nabla) + d\phi$ . Therefore the pair  $(\operatorname{Con}(\mathfrak{g}), \eta)$  forms an  $\Omega^{\geq 1}(A)$ -torsor, and we have a functor  $\mathcal{C} : \operatorname{LPic}_A \to \Omega^{\geq 1}(A)$ -tors,  $\mathfrak{g} \mapsto (\operatorname{Con}(\mathfrak{g}), \eta)$ . The next lemma is taken from [BB93].

**Lemma 2.5.3.** The functor C :  $LPic_A \rightarrow \Omega^{\geq 1}$ -tors is an equivalence of categories.

Global structure : Set  $\Omega_X^{\bullet} = \Omega^{\bullet}(T_X, \mathcal{O}_X)$  (see 3.1.1). When X is nonsingular this is just the ordinary de Rham complex. Let  $\Omega_X^{\geq 1} = \Omega_X^1 \to \Omega_X^{2,cl}$  be its truncation at degree 1. If  $\Omega \subset X$  is an open subset, the automorphism group  $\operatorname{Aut}(\mathfrak{g}_{\Omega}) \cong \Gamma(\Omega, \Omega_X^{1,cl})$ , and if  $\mathfrak{g}_X, \mathfrak{g}'_X \in \operatorname{LPic}_X$  have connections  $\nabla, \nabla'$ defined over  $\Omega$ , then  $\mathfrak{g}_U \cong \mathfrak{g}'_\Omega$  precisely when  $R_{\nabla} - R_{\nabla'} \in d\Omega_X^1(\Omega)$ . It follows that the category  $\operatorname{LPic}_X$  is a sheaf forming a torsor over the sheaf of abelian groups  $\Omega_X^{\geq 1} = \Omega_X^1 \to \Omega_X^{2,cl}$ , and that the isomorphism classes of  $\operatorname{LPic}_X$  are classified by the group  $H^2(X, \Omega_X^{\geq 1})$ . Čech-representatives for vectors in  $H^2(X, \Omega_X^{\geq 1})$  are, in evident notation, pairs  $(\eta_i, \phi_{ij})$  where  $\eta_i$  are closed 2-forms defined in open sets  $U_i \subset X$  and  $\phi_{ij}$  are 1-forms defined in  $U_i \cap U_j$  such that  $\eta_i - \eta_j = d\phi_{ij}$ .

The connections on a locally  $\mathcal{O}_X$ -split Picard Lie algebroid  $\mathfrak{g}_X$  form a torsor over  $\Omega^1_X$ , so the sheaf of connections on  $\mathfrak{g}_X$  gives rise to a class in  $H^1(X, \Omega^1_X)$  (the Atiyah class of  $\mathfrak{g}_X$ ). A Picard Lie algebroid is *locally integrable* if it is locally isomorphic to  $\mathcal{O}_X \oplus T_X$ , i.e. if locally the curvature 2-form of its connections are exact. These are the same as  $\Omega^{1,cl}_X$ -torsors, hence they are classified by  $H^1(X, \Omega^{1,cl}_X) \subseteq H^2(X, \Omega^{\geq 1}_X)$ . If X is a complex manifold, then by Poincaré's lemma  $\Omega^{\geq 1}_X$  is quasi-isomorphic to  $\Omega^{1,cl}$ , so any Picard Lie algebroid is locally integrable. If X is a compact complex algebraic manifold then  $H^2(X, \Omega^{\geq 1}_X) = F^1 H^2_{DR}$ , the Hodge filtration subspace, and the locally integrable Picard Lie algebroids correspond to those classes that vanish on some Zariski open subset of X, i.e., precisely to **C**-linear combinations of the algebraic cycle classes.

The category  $\operatorname{LPic}_X$  is a k-vector space in categories, where the k-vector space structure of  $H^2(X, \Omega_X^{\geq 1})$  is defined directly on  $\operatorname{LPic}_X$  using the Baer sum construction. Namely, if  $(\mathfrak{g}_X^1, \alpha_1), (\mathfrak{g}_X^2, \alpha_2) \in \operatorname{LPic}(X), a, b \in k$ , then  $a\mathfrak{g}_X^1 = \mathcal{O}_X \oplus \mathfrak{g}_X^1/\{(a\phi, -\phi); \phi \in \mathcal{O}_X = \operatorname{Ker}(\alpha_1)\}$  and the linear combination  $a\mathfrak{g}_X^1 + b\mathfrak{g}_X^2$  is a Picard algebroid  $\mathfrak{g}_X$  that is equipped with a homomorphism of Lie algebroids  $s_{a,b} : \mathfrak{g}_X^1 \times_{T_X} \mathfrak{g}_X^2 \to \mathfrak{g}_X$  such that  $s_{a,b}(\delta_1, \delta_2) = (a\delta_1 + b\delta_2)$ .

Relation with the Picard group : Letting  $\operatorname{Pic}_X$  be the groupoid of invertible sheaves on X there exists a functor

$$g: \operatorname{Pic}_X \to \operatorname{LPic}_X$$
$$\lambda \mapsto \mathfrak{c}_X(\lambda) \quad (\text{see } (2.2)).$$

If  $\phi : \lambda_1 \to \lambda_2$  is an isomorphism, then the isomorphism  $g(\phi) : \mathfrak{c}_X(\lambda_1) \to \mathfrak{c}_X(\lambda_2)$  is defined by transport of structure, i.e.  $g(\phi)(\delta_1) = \phi \delta_1 \phi^{-1} \in \mathfrak{c}_X(\lambda_2)$ .

Also,  $\operatorname{Pic}_X$  is a group in categories with composition  $\otimes$  and q is a homomorphism of abelian groups in categories, expressed by

$$\mathfrak{c}_X(\lambda_1 \otimes_{\mathcal{O}_X} \lambda_2) \cong \mathfrak{c}_X(\lambda_1) + \mathfrak{c}_X(\lambda_2) \cong \lambda_1 \otimes_{\mathcal{O}_X} \mathfrak{c}_X(\lambda_2) \otimes_{\mathcal{O}_X} \lambda_1^{-1}.$$

Here all isomorphisms are unique to unique automorphism of either  $\mathfrak{c}_X(\lambda_1 \otimes_{\mathcal{O}_X}$  $\lambda_2$ ),  $\mathfrak{c}_X(\lambda_1) + \mathfrak{c}_X(\lambda_2)$ , or  $\lambda_1 \otimes_{\mathcal{O}_X} \mathfrak{c}_X(\lambda_2) \otimes_{\mathcal{O}_X} \lambda_1^{-1}$ . On the level of isomorphism classes  $[\lambda]$  and  $[\mathfrak{g}_X]$ , parametrized by Čech cohomology classes for an acyclic covering  $X = \bigcup U_{\alpha}$ , g defines a homomorphism of groups dlog:  $H^{1}(X, \mathcal{O}_{X}^{*}) \to H^{1}(X, \Omega^{1, cl}), \quad [(\xi_{\alpha, \beta})] \mapsto [(dlog(\xi_{\alpha, \beta}))], \text{ so that } \hat{dlog}[\lambda] =$  $[\mathfrak{c}_X(\lambda)]$  (using obvious notation).

Remark 2.5.4. For any integer n and  $\mathfrak{g}_X \in \operatorname{LPic}_X$  and  $\lambda \in \operatorname{Pic}_X$  we get  $\lambda^n \otimes_{\mathcal{O}_X} \mathfrak{g}_X \otimes_{\mathcal{O}_X} \lambda^{-n} = \mathfrak{g}_X + \mathfrak{c}_X(\lambda^n) \in \operatorname{LPic}_X$  and there exists a push-out diagram

where  $\phi$  is an isomorphism in Lie<sub>X</sub> (not in LPic<sub>X</sub>); see above Definition 2.5.1.

Assume that X is non-singular of pure dimension n and put  $\omega_X = \Omega_X^n \in$  $\operatorname{Pic}_X$ . Since  $\mathfrak{g}_X$  is also a right  $\mathcal{O}_X$ -module (2.2) we may consider the opposite Lie algebroid  $\mathfrak{g}_X^o$  of  $\mathfrak{g}_X \in \operatorname{LPic}_X(2.1)$ . This is again a Picard Lie algebroid and we have

$$\mathfrak{g}_X^o \cong \mathfrak{c}_X(\omega_X) - \mathfrak{g}_X \cong \omega_X \otimes_{\mathcal{O}_X} (-\mathfrak{g}_X) \otimes_{\mathcal{O}_X} \omega_X^{-1} \cong -\omega_X^{-1} \otimes_{\mathcal{O}_X} \mathfrak{g}_X \otimes_{\mathcal{O}_X} \omega_X.$$

An isomorphism is given by

$$\mu: \mathfrak{g}_X^o \to -\omega_X^{-1} \otimes_{\mathcal{O}_X} \mathfrak{g}_X \otimes_{\mathcal{O}_X} \omega_X$$

$$= \frac{\mathcal{O}_X \oplus (\omega_X^{-1} \otimes_{\mathcal{O}_X} \mathfrak{g}_X \otimes_{\mathcal{O}_X} \omega_X)}{\{(\phi, \phi): \phi \in \mathcal{O}_X = \operatorname{Ker}(\omega_X^{-1} \otimes_{\mathcal{O}_X} \mathfrak{g}_X \otimes_{\mathcal{O}_X} \omega_X \to T_X)\}}$$

$$\delta \mapsto (0, -h \otimes \delta \otimes h^{-1} - h \otimes \frac{\alpha(\delta) \cdot h}{h} \otimes h^{-1})$$

$$= (-h \otimes \frac{\alpha(\delta) \cdot h}{h} \otimes h^{-1}, h \otimes \delta \otimes h^{-1}),$$

where  $h \in \omega_X$ ,  $h \neq 0$ , and  $\alpha(\delta) \in T_X$  acts on h by the negative of the Lie derivative, see (3.2). In particular,  $\mathfrak{c}_X(\lambda)^o \cong \omega_X \otimes_{\mathcal{O}_X} \mathfrak{c}_X(\lambda^{-1}) \otimes_{\mathcal{O}_X} \omega_X^{-1} \cong \mathfrak{c}_X(\omega_X \otimes_{\mathcal{O}_X} \lambda^{-1})$ . Here one may either regard  $\omega_X \otimes_{\mathcal{O}_X} \lambda^{-1}$  as a left module over  $\mathfrak{c}_X(\lambda)^o$  or a right module over  $\mathfrak{c}_X(\lambda)$ ; compare to (3.2). Let us check that the isomorphisms are consistent with the fact that the operation o is an involution:

$$\mathfrak{g}_{X}^{oo} \cong (\omega_{X} \otimes_{\mathcal{O}_{X}} -\mathfrak{g}_{X} \otimes_{\mathcal{O}_{X}} \omega_{X}^{-1})^{o} \\
\cong (-\omega^{-1} \otimes_{\mathcal{O}_{X}} \mathfrak{g}_{X} \otimes_{\mathcal{O}_{X}} \omega_{X})^{o} \\
\cong \omega_{X} \otimes_{\mathcal{O}_{X}} (\omega^{-1} \otimes_{\mathcal{O}_{X}} \mathfrak{g}_{X} \otimes_{\mathcal{O}_{X}} \omega_{X}) \otimes_{\mathcal{O}_{X}} \omega_{X}^{-1} \cong \mathfrak{g}_{X},$$

and more particularly  $\mathfrak{c}(\lambda)_X^{oo} = \mathfrak{c}_X(\lambda^{-1} \otimes \omega_X)^o = \mathfrak{c}_X(\lambda \otimes \omega_X^{-1} \otimes \omega_X) = \mathfrak{c}_X(\lambda).$ 

Now let  $X_h$  be a complex manifold. We then have 3 horizontal short exact sequences, forming a commutative diagram where the vertical sequences also form short exact sequences:



inducing a commutative diagram where the vertical sequences are exact

$$\begin{array}{cccc} H^{0}(X_{h}, \mathcal{O}_{X_{h}}^{*}) & \stackrel{dlog}{\longrightarrow} H^{0}(X_{h}, \Omega_{X_{h}}^{1,cl}) \\ & \downarrow & \downarrow \\ H^{1}(X_{h}, 2\pi i \mathbf{Z}) & \stackrel{1}{\longrightarrow} H^{1}(X_{h}, \mathbf{C}^{*}) & \stackrel{m}{\longrightarrow} H^{1}(X_{h}, \mathbf{C}^{*}) \\ & \downarrow & \downarrow \\ H^{1}(X_{h}, \mathcal{O}_{X_{h}}) & \stackrel{\exp}{\longrightarrow} H^{1}(X_{h}, \mathcal{O}_{X_{h}}^{*}) & \stackrel{c_{1}}{\longrightarrow} H^{2}(X_{h}, 2\pi i \mathbf{Z}) \\ & \downarrow & \downarrow \\ H^{1}(X_{h}, \mathcal{O}_{X_{h}}^{*}) & \stackrel{dlog}{\longrightarrow} H^{1}(X_{h}, \Omega_{X_{h}}^{1,cl}) & \stackrel{c}{\longrightarrow} H^{2}(X_{h}, \mathbf{C}) \\ & \downarrow^{c_{1}} & \downarrow \\ H^{2}(X_{h}, 2\pi i \mathbf{Z}) & \stackrel{1}{\longrightarrow} H^{2}(X_{h}, \mathbf{C}^{*}) & \stackrel{m}{\longrightarrow} H^{2}(X_{h}, \mathbf{C}^{*}) \end{array}$$

The following proposition is due to A. Weil [Wei58, Ch. V, no, 4, Lemme 2]. Put  $\overline{H}^2(X_h, 2\pi i \mathbf{Z}) = \text{Im}(f)$ .

**Proposition 2.5.5.** ( $X_h$  is a complex manifold) A Picard Lie algebroid  $\mathfrak{g}_{X_h}$  is isomorphic to  $\mathfrak{c}_{X_h}(\lambda)$  for some invertible sheaf  $\lambda$  if and only if  $c([\mathfrak{g}_{X_h}]) \in \overline{H^2}(X_h, 2\pi i \mathbf{Z})$ .

In other words,  $\operatorname{Im}(dlog) = c^{-1}(\overline{H}^2(X_h, 2\pi i \mathbf{Z})).$ 

Proof. If  $\{\xi_{\alpha\beta}\}$  is a Čech representative of  $[\mathfrak{g}_{X_h}] \in H^1(X_h, \Omega^{1,cl})$ , then, since  $\{\partial\xi_{\alpha\beta}\} \in \overline{H}^2(X_h, 2\pi i \mathbf{Z})$ , using the holomorphic Poincaré lemma (assuming the Čech covering is sufficiently fine), the chain  $\{\phi_{\alpha\beta}\} = \{\exp^{\int \xi_{\alpha\beta}}\}$ is a cocycle representing a class in  $H^1(X_h, \mathcal{O}^*_{X_h})$  such that  $dlog(\phi_{\alpha\beta}) = \xi_{\alpha\beta}$ . Choosing  $\lambda \in \operatorname{Pic}_{X_h}$  such that  $[\lambda] = \{\phi_{\alpha\beta}\}$  we have  $\mathfrak{c}_{X_h}(\lambda) \cong \mathfrak{g}_{X_h}$ .  $\Box$  Remark 2.5.6. Poincaré's lemma holds more generally when  $X_h$  is a locally holomorphically contractible complex analytic space [Rei67], so Proposition 2.5.5 holds for such spaces.

Assume that X is a non-singular complex projective variety and  $X_h$  its associated compact complex manifold. Put  $\operatorname{LPic}_{X_h}^0 = \operatorname{Ker}(c)$ ; see the horizontal sequence (non-exact) above containing the mapping c. Consider the long exact sequence in homology of the second horizontal short exact sequence in (2.5.1) and let  $\overline{H}^1(X_h, \mathbb{C})$  be the image of  $H^1(X_h, \mathbb{C})$  in  $H^1(X_h, \mathcal{O}_{X_h}) \cong H^1(X, \mathcal{O}_X)$  (GAGA). In analogy with the ordinary Picard variety one may call

$$\operatorname{LPic}_X^0 \cong \frac{H^1(X, \mathcal{O}_X)}{\overline{H}^1(X_h, \mathbf{C})}$$

the Lie Picard variety of X. The functor g above induces a proper surjective morphism from the Picard variety of X to its Lie Picard variety,  $\operatorname{Pic}_X^0 \to \operatorname{LPic}_X^0$ , which is simply the canonical mapping

$$\frac{H^1(X,\mathcal{O}_X)}{H^1(X_h,2\pi i \mathbf{Z})} \to \frac{H^1(X,\mathcal{O}_X)}{\bar{H}^1(X_h,\mathbf{C})}.$$

When  $X_h$  is a curve one has  $\operatorname{LPic}_{X_h}^0 = 0$ , hence  $\operatorname{LPic}_{X_h} \subseteq H^2(X_h, \mathbb{C})$ ; in particular,  $\mathfrak{c}_{X_h}(\lambda_1) \cong \mathfrak{c}_{X_h}(\lambda_2)$  precisely when  $\lambda_1, \lambda_2 \in \operatorname{Pic}_{X_h}$  have the same degree. See also (3.3).

If  $X_h$  is a Stein manifold it follows from the first horizontal short exact sequence and Cartan's theorem B that  $H^1(X_h, \mathcal{O}^*_{X_h}) = H^2(X_h, 2\pi i \mathbf{Z})$  and  $H^2(X, \mathcal{O}^*_{X_h}) = H^3(X_h, 2\pi i \mathbf{Z})$ , so using the long exact sequence in cohomology of the third horizontal short exact sequence we get an exact sequence

$$H^1(X_h, \mathbf{C}^*) \to H^2(X_h, 2\pi i \mathbf{Z}) \to H^1(X_h, \Omega^{1, cl}_{X_h}) \to H^2(X_h, \mathbf{C}^*) \to H^3(X_h, 2\pi i \mathbf{Z})$$

#### 3. Modules

3.1. **Basic facts.** Let  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha)$  be a Lie algebroid.

**Definition 3.1.1.** A *left*  $\mathfrak{g}_X$ -module M is an  $\mathcal{O}_X$ -module M and a homomorphism of k-Lie algebras

$$\rho : \mathfrak{g}_X \to \mathfrak{c}_X(M) \quad (\text{see } (2.2.1))$$

satisfying  $\rho(f\delta) \cdot m = f\rho(\delta) \cdot m$ , and  $\rho(\delta) \cdot fm = \alpha(\delta)(f)m + f\rho(\delta) \cdot m$ , for all  $f \in \mathcal{O}_X$ ,  $\delta \in \mathfrak{g}_X$ ,  $m \in M$ . When M is faithful over  $\mathcal{O}_X$  these conditions express that  $\rho$  is a homomorphism of Lie algebroids.

A right  $\mathfrak{g}_X$ -module is defined by a k-linear homomorphism

$$\nu:\mathfrak{g}_X\to\mathfrak{gl}_k(M)$$

satisfying  $\nu([\delta, \eta]) = -[\nu(\delta), \nu(\eta)]$ , and  $\nu(f\delta) \cdot m = f\nu(\delta) \cdot m - \alpha(\delta)(f)m = \nu(\delta) \cdot (fm)$ , where  $\delta, \eta \in \mathfrak{g}_X$ .

When we say module without specifying left or right we will always mean left module. A  $\mathfrak{g}_X$ -module M is the same as a  $\mathcal{D}(\mathfrak{g}_X)$ -module and it is *coherent* (quasi-coherent) if it is coherent (quasi-coherent) over the ring  $\mathcal{D}(\mathfrak{g}_X)$ . M is torsion free if for every non-zero coherent  $\mathcal{O}_X$ -submodule  $N \subset M$  one has dim supp  $N = \dim X$ ; this coincides with the usual notion of torsion free modules when X is integral. Note that if  $M = \mathcal{D}(\mathfrak{g}_X)M^0$ and  $M^0$  is torsion free it does not follow that M is torsion free, but if  $M \in \operatorname{coh}(\mathfrak{g}_X)$  there exists locally  $M^0 \in \operatorname{coh}(\mathcal{O}_X)$  such that  $M = \mathcal{D}(\mathfrak{g}_X)M^0$ so  $\operatorname{supp} M = \operatorname{supp} M^0$ , implying that  $\operatorname{supp} M$  is an algebraic (analytic) subspace of X.

We let  $\operatorname{Mod}(\mathfrak{g}_X) = \operatorname{Mod}(\mathcal{D}(\mathfrak{g}_X))$  be the category of  $\mathfrak{g}_X$ -modules and  $\operatorname{coh}(\mathfrak{g}_X) \subset \operatorname{qcoh}(\mathfrak{g}_X)$  its sub-categories of coherent and quasi-coherent  $\mathfrak{g}_X$ modules. The triangulated derived category of complexes of  $\mathfrak{g}_X$ -modules  $M^{\bullet}$ with coherent homology  $H^{\bullet}(M^{\bullet}) \in \operatorname{coh}(\mathfrak{g}_X)$  is denoted by  $D_{\operatorname{coh}}(\mathfrak{g}_X)$  and the subcategories of complexes that are bounded above and below are  $D^-_{\operatorname{coh}}(\mathfrak{g}_X)$ and  $D^+_{\operatorname{coh}}(\mathfrak{g}_X)$ . We shall mostly deal with the subcategory of bounded complexes  $D^b_{\operatorname{coh}}(\mathfrak{g}_X) = D^+_{\operatorname{coh}}(\mathfrak{g}_X) \cap D^-_{\operatorname{coh}}(\mathfrak{g}_X)$  mainly because it is generated by its homology objects, i.e., every object  $M^{\bullet} \in D^b_{\operatorname{coh}}(\mathfrak{g}_X)$  is isomorphic to a complex of coherent  $\mathfrak{g}_X$ -modules; see the proof of [BGK<sup>+</sup>87, I, Prop. 12.8].

For completeness we quote results of Deligne and Bernstein [§VI, Th. 2.10, Prop. 2.11, loc. cit] about complexes on a noetherian variety X. Let  $D^b(\operatorname{coh}(\mathfrak{g}_X))$   $(D^b(\operatorname{qcoh}(\mathfrak{g}_X)))$  be the category of bounded complexes of coherent (quasi-coherent)  $\mathfrak{g}_X$ -modules, and  $D^b_{\operatorname{qcoh}}(\mathfrak{g}_X)$  the category of bounded complexes of  $\mathfrak{g}_X$ -modules  $M^{\bullet}$  such that  $H^{\bullet}(M^{\bullet}) \in \operatorname{qcoh}(\mathfrak{g}_X)$ . Then the natural inclusion functors

$$D^{b}(\operatorname{coh}(\mathfrak{g}_{X})) \to D^{b}_{\operatorname{coh}}(\operatorname{qcoh}(\mathfrak{g}_{X}))$$

and

$$D^{b}(\operatorname{qcoh}(\mathfrak{g}_{X})) \to D^{b}_{\operatorname{qcoh}}(\operatorname{Mod}(\mathfrak{g}_{X}))$$

are equivalences of categories. Hence we have also a third equivalence

 $D^b(\operatorname{coh}(\mathfrak{g}_X)) \to D^b_{\operatorname{coh}}(\operatorname{Mod}(\mathfrak{g}_X)).$ 

A  $\mathfrak{g}_X$ -connection on M is a homomorphism of  $\mathcal{O}_X$ -modules  $\rho : \mathfrak{g}_X \to \mathfrak{c}_X(M)$ . When  $\mathfrak{g}_X = T_X$  this is just a connection on the Lie algebroid  $\mathfrak{c}_X(M)$ (assuming that M is faithful over  $\mathcal{O}_X$ ). The k-linear homomorphism  $\nabla : M \to Hom_{\mathcal{O}_X}(\mathfrak{g}_X, M), \nabla(m)(\delta) = \rho(\delta)(m)$ , is the first step in a sequence of k-linear homomorphisms of  $\mathcal{O}_X$ -modules. Put  $l = \operatorname{rk} \mathfrak{g}_X$  and let  $\{\delta_1, \ldots, \delta_l\}$  be generators of the  $\mathcal{O}_X$ -module  $\mathfrak{g}_X(l$  and the generators are defined locally). Put  $\Omega^p(\mathfrak{g}_X, M) = Hom_{\mathcal{O}_X}(\wedge^p \mathfrak{g}_X, M)$  and consider

(3.1.1) 
$$M \xrightarrow{\nabla} \Omega^1(\mathfrak{g}_X, M) \xrightarrow{\nabla} \cdots \xrightarrow{\nabla} \Omega^l(\mathfrak{g}_X, M)$$

where the mapping  $\nabla : \Omega^p(\mathfrak{g}_X, M) \to \Omega^{p+1}(\mathfrak{g}_X, M)$  is defined in the usual way

$$(\nabla\omega)(\delta_{i_{1}}\wedge\cdots\wedge\delta_{i_{p+1}}) = \sum_{1\leq s< t\leq p+1} (-1)^{s+t} \omega([\delta_{i_{s}},\delta_{i_{t}}]\wedge\delta_{i_{1}}\wedge\cdots\hat{\delta}_{i_{s}}\cdots\hat{\delta}_{i_{t}}\cdots\delta_{i_{p+1}}) + \sum_{1\leq s\leq p+1} (-1)^{s+1}\delta_{i_{s}}\cdot\omega(\delta_{i_{1}}\wedge\cdots\hat{\delta}_{i_{s}}\cdots\wedge\delta_{i_{p+1}}).$$

If  $\rho$  gives M the structure of  $\mathfrak{g}_X$ -module this is a complex, i.e.  $\nabla^2 = 0$ , and we call  $(\Omega^{\bullet}(\mathfrak{g}_X, M), \nabla)$  the *de Rham* (-Chevalley-Hochschild) complex of the  $\mathfrak{g}_X$ -module M; see also [Käl98, Lem. 2.3]. The sheaf  $Hom_{\mathcal{O}_X}(\mathfrak{g}_X,\mathfrak{c}_X(M))$  forms a torsor over the sheaf of abelian groups  $N_X := Hom_{\mathcal{O}_X}(\mathfrak{g}_X,\mathfrak{gl}_{\mathcal{O}_X}(M))$ . Hence supposing that local  $\mathfrak{g}_X$ -connections exist we get an obstruction class in  $H^1(X, N_X)$  for the existence of a global  $\mathfrak{g}_X$ -connection.

Some basic facts about the Lie algebroid  $T_X = Hom_{\mathcal{O}_X}(\Omega_{X/k}, \mathcal{O}_X)$  are summarized as follows:

**Proposition 3.1.2.** (X is a space which in the algebraic case is assumed to be locally a sub-scheme of a regular scheme of finite type over k) The following are equivalent:

- (1) X/k is non-singular;
- (2)  $\Omega_{X/k}$  is locally free; hence  $T_X$  is locally free;
- (3)  $T_X$  is a simple Lie algebroid, i.e.  $T_X$  contains no proper coherent ideal;
- (4)  $\mathcal{O}_X$  is a simple  $T_X$ -module, i.e.  $\mathcal{O}_X$  contains no proper coherent  $T_X$ -submodule.

*Proof.* (1)  $\Leftrightarrow$  (2): See [Mat86] (k is algebraically closed of characteristic 0).

 $(3) \Rightarrow (1), (4) \Rightarrow (1)$ : The assertion is local in X so one may assume that  $X \subset Y$  where, in the complex analytic case  $Y = \mathbb{C}^n$ , and in the algebraic case  $Y = \mathbb{A}^n$ . Let  $j: X \to Y$  be the associated closed immersion of spaces and I be the kernel of the surjection  $\mathcal{O}_Y \to j_*(\mathcal{O}_X)$ . Put  $t = \dim X$  (the maximal value of  $\dim \mathcal{O}_x$  when  $x \in X$ ). Then if  $D_1, \ldots, D_t \in T_Y$  and  $a_1, \ldots, a_t \in I$  one can form the elements  $\det(D_i(a_j)) \in \mathcal{O}_Y$  and we let J be the ideal they generate. We assert that the ideal  $I^s := j^*(J) \subset \mathcal{O}_X$  is stable under the action of  $T_X$ . Since  $T_X = j^{-1}(T_Y(I)/IT_Y)$  it suffices to see that  $\delta \cdot (\det(D_i(a_j))) \in J$  when  $\delta \in T_Y(I)$ . This holds more generally, namely for any integer  $l \geq 1$  letting  $J_l$  be the ideal generated by elements  $\det(D_i(a_j))$  where  $D_1, \ldots, D_l \in T_Y$  and  $a_1, \ldots, a_l \in I$ , we have  $T_Y(I) \cdot J_l \subset J_l$ . To see this consider a term of  $\det(D_i(a_j))$ , which has the form

$$\pm D_1(a_{i_1})D_2(a_{i_2})\cdots D_l(a_{i_l}),$$

where  $(i_1, \ldots, i_l)$  is a permutation of  $(1, \ldots, l)$ . Applying  $\delta$  to such a term one gets 2*l* terms occuring either by replacing one of the  $D_i$  by  $D'_i = [\delta, D_i]$ or one of the  $a_{i_s}$  by  $a'_{i_s} := \delta(a_{i_s}) \in I$ . This implies that  $\delta \cdot \det(D_i(a_j))$  is a sum of 2*l* terms each belonging to  $J_l$ .

By the Jacobian criterion of regularity for spaces of this type the singular locus of X is the zero locus of  $I^s$ . Assuming that X is singular it follows that  $I^s \subset \mathcal{O}_X$  is a proper ideal which is moreover a  $T_X$ -submodule. Also,  $I^sT_X \subset$  $T_X$  is an ideal of the Lie algebroid  $T_X$ , which is proper by Nakayama's lemma. Hence  $\mathcal{O}_X$  is not a simple  $T_X$ -module and  $T_X$  is not a simple Lie algebroid.

 $(4) \Rightarrow (3)$ : Let  $\mathfrak{a} \subset T_X$  be a proper coherent ideal of  $T_X$ . It suffices to prove that  $1 \notin \mathfrak{a} \cdot \mathcal{O}_X$ , for then  $\mathfrak{a} \cdot \mathcal{O}_X \subset \mathcal{O}_X$  is a proper ideal, and since clearly  $\mathfrak{a} \cdot \mathcal{O}_X$  is a coherent  $T_X$ -module it follows that  $\mathcal{O}_X$  is not simple. Thus suppose that  $\partial \in \mathfrak{a}, \eta \in T_X$  and  $\partial(a) = 1$ , then  $\eta = \partial(a)\eta = [\partial, a\eta] - a[\partial, \eta] \in \mathfrak{a}$ , implying that  $T_X \subseteq \mathfrak{a}$ , contradicting the assumption that  $\mathfrak{a}$  is a proper ideal. (2)  $\Rightarrow$  (4): This is of course well-known. Let  $I \subset \mathcal{O}_X$  be a non-zero coherent  $T_X$ -submodule. It suffices to see that  $I_x = \mathcal{O}_x$  when x is a closed point in X. By (2)  $\Rightarrow$  (1) there exists a regular system of parameters  $(x_1, \ldots, x_n)$  of the regular local ring  $\mathcal{O}_x$ , so letting  $\hat{\mathcal{O}}_x$  denote the completion of  $\mathcal{O}_x$  by Artin-Rees lemma and Cohen's structure theorem one has  $\mathcal{O}_x \subset \hat{\mathcal{O}}_x \cong k[[x_1, \ldots, x_n]]$ . (2) implies that  $(\Omega_{X/k})_x$  is free over  $dx_i$ , hence there exist derivations  $\partial_{x_i} \in \text{Der}_k(\mathcal{O}_x)$  such that  $\partial_{x_i}(x_j) = \delta_{ij}$ . Then considering a non-zero  $f \in I_x \subset k[[x_1, \ldots, x_n]]$ , one checks that there exists a multi-index  $\alpha$  such that  $\partial^{\alpha}(f) \in I_x$  is invertible in  $\mathcal{O}_x$ ; therefore  $I_x = \mathcal{O}_x$ .  $\Box$ 

- Remarks 3.1.3. (1) When  $\alpha \neq 0$  the mapping  $\operatorname{ad} : \mathfrak{g}_X \to \mathfrak{c}_X(\mathfrak{g}_X), \operatorname{ad}(\partial)(\delta) = [\partial, \delta]$  is not  $\mathcal{O}_X$ -linear, and so does not define a module structure on  $\mathfrak{g}_X$ .
  - (2) Recall Zariski-Lipman's conjecture: If  $T_X$  is locally free, then the equivalent conditions in Proposition 3.1.2 hold, and  $\Omega_X \cong \Omega^1_X(T_X, \mathcal{O}_X)$  (3.1.1). [Sai80] contains conditions ensuring that  $T_X(I)$  be free when I is the ideal of a divisor on a complex manifold.
  - (3) Let  $\pi : X' \to X$  be the normalization morphism of an integral space. Let  $J \subset \mathcal{O}_X$  be the conductor ideal, that is  $J = \{\phi \in \mathcal{O}_X : \phi \pi_*(\mathcal{O}_{X'}) \subset \mathcal{O}_X\}$ . By Seidenberg's result in *Remark* 2.4.3 it easily follows that J is  $T_X$ -submodule of  $\mathcal{O}_X$ . Therefore, by Proposition 3.1.2 we get a curious proof of the well-known fact that if X is non-singular, then X is normal.
  - (4) A noetherian k-algebra R of characteristic p > 0 is a simple  $\text{Der}_k(R)$ module if and only if it has the form  $R = k[t_1, \ldots, t_n]/(t_1^p, \ldots, t_n^p)$ , where k is a field of characteristic p [Har61].

3.2. Interchanging left and right modules. We make a slight extension of the discussion in [BGK<sup>+</sup>87, VI]. Let  $M_1, M_2$  be left  $\mathfrak{g}_X$ -modules and  $N_1, N_2$  be right  $\mathfrak{g}_X$  modules (*Def.* 3.1.1) defined by homomorphisms

$$l_i : \mathfrak{g}_X \to \mathfrak{c}_X(M_i)$$
  
$$r_i : \mathfrak{g}_X \to \mathfrak{gl}_k(N_i).$$

Then  $M_1 \otimes_{\mathcal{O}_X} M_2$ ,  $Hom_{\mathcal{O}_X}(M_1, M_2)$  and  $Hom_{\mathcal{O}_X}(N_1, N_2)$  are left  $\mathfrak{g}_X$ -modules defined by the homomorphisms  $\rho_1, \rho_2, \rho_2$ 

- (3.2.1)  $\rho_1 : \mathfrak{g}_X \to \mathfrak{c}_X(M_1 \otimes_{\mathcal{O}_X} M_2)$
- $\delta \mapsto l_1(\delta) \otimes \mathrm{id}_{M_1} + \mathrm{id}_{M_1} \otimes l_2(\delta)$
- (3.2.2)  $\rho_2 : \mathfrak{g}_X \to \mathfrak{c}_X(Hom_{\mathcal{O}_X}(M_1, M_2))$
- $(3.2.3) \qquad \begin{aligned} \delta &\mapsto \rho_2(\delta) : f \mapsto l_i(\delta) \cdot f f \cdot l_2(\delta) \\ \rho_3 &: \mathfrak{g}_X \to \mathfrak{c}_X(Hom_{\mathcal{O}_X}(N_1, N_2)) \end{aligned}$
- $\delta \mapsto \rho_3(\delta) : f \mapsto f \cdot l_1(\delta) l_2(\delta) \cdot f.$

The  $\mathcal{O}_X$ -modules  $M_1 \otimes N_1$  and  $Hom_{\mathcal{O}_X}(M_1, N_1)$  are right  $\mathfrak{g}_X$ -modules defined by the homomorphisms  $\nu_1, \nu_2$ 

(3.2.4) 
$$\nu_1 \quad : \quad \mathfrak{g}_X \to \mathfrak{gl}_k(M_1 \otimes N_1)$$

$$\delta \mapsto -l_1(\delta) \otimes \mathrm{id}_{N_1} + \mathrm{id}_{M_1} \otimes r_1(\delta),$$

(3.2.5) 
$$\nu_2 : \mathfrak{g}_X \mapsto \mathfrak{gl}_k(Hom(M_1, N_1))$$
$$\delta \mapsto \nu_2(\delta) : f \mapsto f \cdot l_1(\delta) + r_1(\delta) \cdot f$$

*Remark* 3.2.1. The  $\mathcal{O}_X$ -modules  $Hom_{\mathcal{O}_X}(N_1, M_1)$  and  $N_1 \otimes_{\mathcal{O}_X} N_2$  are in general neither right nor left  $\mathfrak{g}_X$ -modules [Oda83].

Assume now that X is non-singular of pure dimension n, so that  $T_X$  is a locally free  $\mathcal{O}_X$ -module of rank n. Then

$$\omega_X := \Omega^n(T_X, \mathcal{O}_X)$$

is an invertible module which moreover is a  $right^2 \mathfrak{g}_X$ -module. The homomorphism  $\nu : \mathfrak{g}_X \to \omega_X$  is defined by

$$(\phi \cdot \delta)(d) = \phi(\delta \cdot d) - \alpha(\delta)(\phi(d)), \text{ where}$$
  
 $\delta \cdot d = [\alpha(\delta), \partial_1] \wedge \partial_2 \wedge \cdots \partial_n + \cdots + \partial_1 \wedge \cdots \wedge [\alpha(\delta), \partial_n] \in \bigwedge^n T_X,$ 

 $\partial_1, \ldots, \partial_r \in \mathfrak{g}_X$  form a local basis of  $\mathfrak{g}_X$ , and  $d = \partial_1 \wedge \cdots \wedge \partial_n$ . Note that  $\operatorname{Ker}(\mathfrak{g}_X \to T_X)$  acts trivially on  $\omega_X$ .<sup>3</sup> Then if M is a left  $\mathfrak{g}_X$ -module it follows that

$$M_r = \omega_X \otimes_{\mathcal{O}_X} M$$

is a right  $\mathfrak{g}_X$ -module (3.2.5), and starting with a right  $\mathfrak{g}_X$ -module N,

$$N_l = Hom_{\mathcal{O}_X}(\omega_X, N) = \omega_X^{-1} \otimes_{\mathcal{O}_X} N$$

is a left  $\mathfrak{g}_X$ -module (3.2.3). Since  $\omega_X$  is invertible the pair  $(\cdot \otimes_{\mathcal{O}_X} \omega_X, \omega_X^{-1} \otimes_{\mathcal{O}_X} \cdot)$  defines an equivalence between the categories of left and right  $\mathfrak{g}_X$ -modules.

3.3. Modules over Picard Lie algebroids. When  $\mathfrak{g}_X$  is a locally  $\mathcal{O}_X$ -split Picard Lie algebroid we *identify*  $\operatorname{Ker}(\alpha)$  with  $\mathcal{O}_X$ , so in this case a  $\mathfrak{g}_X$ -module structure is locally defined by the (non-Lie) action of  $T_X$  with respect to a connection. But if  $\rho : T_X \to \mathfrak{c}_X(M)$  is a  $T_X$ -module, then the composition  $\rho \circ \alpha : \mathfrak{g}_X \to \mathfrak{c}_X(M)$  does not satisfy  $\rho \circ \alpha(\phi) = i(\phi)$ , where  $\phi \in \mathcal{O}_X \subseteq \mathfrak{g}_X$  and  $i : \mathcal{O}_X \to \mathfrak{gl}_k(M)$ . Hence  $T_X$ -modules are considered as  $\mathfrak{g}_X$ -modules only when  $\mathfrak{g}_X = \mathcal{O}_X \oplus T_X$ .

Picard Lie algebroids do not always have modules that are coherent over  $\mathcal{O}_X$ . If  $r \in k$  we let  $\mathfrak{g}_X^r$  denote the push-out of  $\mathfrak{g}_X$  by the character  $\mathcal{O}_X \to \mathcal{O}_X$ ,  $\phi \mapsto r \cdot \phi$  (see above 2.5.1).

**Proposition 3.3.1.** (X is a non-singular space) Let  $\mathfrak{g}_X$  be a Picard algebroid. Any  $\mathfrak{g}_X$ -module which is coherent over  $\mathcal{O}_X$  is locally free. The following are equivalent:

<sup>3</sup>Of course, in the same way the  $\mathcal{O}_X$ -module  $\Omega^r(\mathfrak{g}_X, \mathcal{O}_X) = Hom_{\mathcal{O}_X}(\bigwedge^r \mathfrak{g}_X, \mathcal{O}_X)$ , if  $\operatorname{rk} \mathfrak{g}_X = r$ , is a right  $\mathfrak{g}_X$ -module for which  $\operatorname{Ker}(\mathfrak{g}_X \to T_X)$  acts non-trivially.

<sup>&</sup>lt;sup>2</sup>See Section 2.5 for the case of Picard Lie algebroids.

- (1)  $\mathfrak{g}_X$  has a non-zero module M which is coherent over  $\mathcal{O}_X$ , of some rank r.
- (2)  $\mathfrak{g}_X^r = \mathfrak{c}_X(\lambda)$  for some invertible sheaf  $\lambda$ , where  $\lambda = \det M$ , the determinant bundle of M.

We have  $\mathfrak{c}_X(\det M) = \mathfrak{c}_X^{\mathrm{tr}}(M)$ , the push-out of the linear algebroid  $\mathfrak{c}_X(M)$ by the trace homomorphism  $\mathrm{tr} : \mathfrak{gl}_{\mathcal{O}_X}(M) \to \mathcal{O}_X$  (see Remark 2.2.1).

 $(1) \Rightarrow (2)$  is proven in [BB93, 2.3.1]; the remaining parts are also well-known.

*Proof.* The first assertion follows since  $\mathfrak{g}_X$  is transitive and X is nonsingular; see, e.g. [BGK<sup>+</sup>87] for a corresponding statement for  $\mathcal{D}$ -modules. Thus if  $M \in \operatorname{coh}(\mathcal{O}_X)$  is a  $\mathfrak{g}_X$ -module, then M is locally free, so  $\lambda :=$ det  $M = \wedge_{\mathcal{O}_X}^r M$  is an invertible sheaf. Define a homomorphism of Lie algebroids  $\mathfrak{g}_X \to \mathfrak{c}(\lambda)$ , by  $\delta \cdot (m_1 \wedge m_2 \wedge \cdots \wedge m_r) = (\delta \cdot m_1) \wedge m_2 \wedge \cdots \wedge m_d +$  $\cdots + m_1 \wedge \cdots \wedge (\delta \cdot m_r)$  (Leibniz' rule), for  $\delta \in \mathfrak{g}_X$  and  $m_i \in M$ ; in particular  $\mathfrak{l}_{\mathfrak{g}_X}$  is mapped to  $r\mathfrak{l}_{\mathfrak{c}(\lambda)}$ . This is an isomorphism. The last statement should now also be evident.  $\Box$ 

Remark 3.3.2. Let  $\mathfrak{g}_X \in \operatorname{LPic}_X$ . Then  $\omega_X$  is not a right  $\mathfrak{g}_X$ -module (requiering that  $1_{\mathfrak{g}_X} = 1_{\mathcal{O}_X}$ ) unless  $\mathfrak{g}_X = \mathcal{O}_X \oplus T_X$ . Moreover, the formula (3.2.5) does not give  $\omega_X \otimes M$  a structure of right module in this sense, but we will later (3.5) regard it as a right  $\mathfrak{g}_X$ -module such that  $1_{\mathfrak{g}_X}$  acts by -1.

3.3.3. Twisted sheaves. Let X be a complex analytic manifold. The third horizontal short exact sequence in the diagram before Proposition 2.5.5 gives a long exact sequence in homology

$$\cdots \to H^1(X, \mathbf{C}^*) \to H^1(X, \mathcal{O}_X^*) \xrightarrow{dlog} H^1(X, \Omega_X^{1, cl}) \xrightarrow{t} H^2(X, \mathbf{C}_X^*) \to \cdots$$

So if  $\mathfrak{g}_X \in \operatorname{LPic}_X$  and  $\phi := t([\mathfrak{g}_X]) \neq 1 \in H^2(X, \mathbb{C}_X^*)$ , then  $\operatorname{Mod}(\mathfrak{g}_X)$ contains no coherent  $\mathcal{O}_X$ -modules (*Prop.* 3.3.1). One may then instead of the category of sheaves consider the category of  $\phi$ -twisted sheaves and its subcategory of twisted  $\mathfrak{g}_X$ -modules  $\operatorname{Mod}(\mathfrak{g}_X, \phi)$ . We will make some remarks about the possibility to use twisted sheaves; see also [Kas89, §3]. Objects  $M \in \operatorname{Mod}(\mathfrak{g}_X, \phi)$  are given by a descent datum with respect to a covering of X, so that the co-cycle condition for sheaves  $[\partial \psi_{ij}] = [\psi_{ik}\psi_{jk}^{-1}\psi_{ij}^{-1}] =$  $1 \in H^2(X, \mathcal{O}_X^*)$  (in Čech-notation) is broken to  $[\partial \psi_{ij}] = \phi$  id. Morphisms of  $\phi$ -twisted sheaves are still global sections of sheaves of morphisms, and  $\operatorname{Mod}(\mathfrak{g}_X, \phi)$  is an abelian category.

For example, let  $c \in \mathbf{C}^*$  and  $\lambda \in \operatorname{Pic}_X$ , and suppose that open sets  $U_i$  are chosen so that  $\lambda_{U_i}$  is trivial, giving isomorphisms  $\psi_{ij} : (\lambda_{U_j})_{U_i} \to (\lambda_{U_i})_{U_j}$ where  $\psi_{ij} \in \mathcal{O}^*_{U_i \cap U_j}$ . Choosing for each pair (i, j) a *c*th root  $\psi^c_{ij}$  of  $\psi_{ij}$ , we get a 2-cocycle  $\phi_{ijk} = \psi^c_{ik} \psi^{-c}_{jk} \psi^{-c}_{ij}$  defining a class  $[\phi_{ijk}] \in H^2(X, \mathbf{C}^*_X)$ . The descent datum  $\{f_{ij}, \mathcal{O}_{U_i}\}$  then defines an object  $\lambda^c \in \operatorname{Mod}(\mathfrak{c}_X(\lambda^c), \phi)$ . Note that the linear algebroid  $\mathfrak{c}_X(\lambda^c) = c \cdot \mathfrak{c}_X(\lambda)$  is on ordinary sheaf.

Let  $\{\psi_{ij}\}$  be a Čech representative of  $[\mathfrak{g}_X] \in H^1(X, \Omega^{1,cl})$ , choose  $\int \psi_{ij} \in \mathcal{O}_{U_{ij}}$ , using the holomorphic Poincaré lemma, such that  $\psi_{ij} = d \int \psi_{ij}$ , and put  $f_{ij} = \exp^{\int \psi_{ij}}$ , defining a 1-chain with values in  $\mathcal{O}_X^*$ . So  $\{f_{ij}\}$  is determined up to a 1-cocycle with values in  $\mathbb{C}^*$ , but let us make a *choice* of this

1-chain. The isomorphisms  $f_{ij} : \mathcal{O}_{U_{ij}} \to \mathcal{O}_{U_{ji}}$  then gives a descent datum  $\{f_{ij,\mathcal{O}_{U_i}}\}$ , defining a  $\phi$ -twisted sheaf of  $\mathfrak{g}_X$ -modules when  $\phi = t([\mathfrak{g}_X])$ . We denote this  $\phi$ -twisted sheaf by  $\lambda = \lambda(\mathfrak{g}_X)^4$ .

Let  $\mathfrak{g}_X^1, \mathfrak{g}_X^2 \in \operatorname{LPic}_X$  and  $M \in \operatorname{Mod}(\mathfrak{g}_X^1, \phi_1), N \in \operatorname{Mod}(\mathfrak{g}_X^2, \phi_2)$ .

Tensor products. We have  $M \otimes_{\mathcal{O}_X} N \in \text{Mod}(\mathfrak{g}_X^1 + \mathfrak{g}_X^2, \phi_1 \cdot \phi_2)$  where the sum  $\mathfrak{g}_X^1 + \mathfrak{g}_X^2$  is taken in  $\text{LPic}_X$ . When N locally is invertible as  $\mathcal{O}_X$ -module, then

$$M \to M \otimes_{\mathcal{O}_X} N$$
$$\operatorname{Mod}(\mathfrak{g}^1_X, \phi_1) \to \operatorname{Mod}(\mathfrak{g}^1_X + \mathfrak{g}^2_X, \phi_1 \cdot \phi_2)$$

defines an equivalence of categories. This equivalence is not unique, for one may change N to  $N \otimes_{\mathbf{C}} L$  where L is any  $\phi_2$ -twisted invertible  $\mathbf{C}^*$ -module.

In particular, if  $\mathfrak{g}_X^1 = \mathfrak{g}_X$  one can choose  $\phi = t([-\mathfrak{g}_X]) = t([\mathfrak{g}_X])^{-1}$ ,  $\mathfrak{g}_X^2 = -\mathfrak{g}_X$ , and  $N = \lambda(\phi)$ , resulting in an equivalence

$$\operatorname{Mod}(\mathfrak{g}_X) \cong \operatorname{Mod}(T_X, \phi),$$

between a category of sheaves and a category of  $\phi$ -twisted sheaves.

Homomorphisms. We have  $Hom_{\mathcal{O}_X}(M, N) \in \operatorname{Mod}(\mathfrak{g}_X^2 - \mathfrak{g}_X^1, \phi_2 \cdot \phi_1^{-1})$ . Thus if  $H^2(X, \mathbb{C}^*)$  has torsion, there may exist "untwisted" morphisms between M and N even if  $\phi_1 \neq \phi_2$ . When  $\phi_1 = \phi_2 = \phi$  then  $Hom_{\mathcal{O}_X}(M, N)$  is an ordinary sheaf of  $T_X$ -modules. Suppose that  $X = \bigcup X_i$  where  $X_i \subset X$  are open subsets, and  $M_i \in \operatorname{Mod}(\mathfrak{g}_{X_i}, \phi_i)$  are  $\phi_i$ -twisted sheaves of  $\mathfrak{g}_{X_i}^i$ -modules. Restricting  $M_i, \mathfrak{g}_{X_i}^i$ , and  $\phi_i$  to  $X_{ij} = X_i \cap X_j$  results in a  $\phi_{ij}$ -twisted sheaf of  $\mathfrak{g}_{X_{ij}}^1$ -modules, where possibly  $[\phi_{ij}] = 1 \in H^2(X, \mathbb{C}^*)$  so the  $M_{ij} := (M_i)_{X_j}$ may be ordinary sheaves. Let  $\{\psi_{ij}\}$  be a Čech-cocycle of isomorphisms of Lie algebroids  $\psi_{ij} : \mathfrak{g}_{X_{ij}}^i \to \mathfrak{g}_{X_{ji}}^j$ . Thus the Lie algebroids  $\mathfrak{g}_{X_i}^i$  on  $X_i$  can be glued to a global Lie algebroid  $\mathfrak{g}_X \in \operatorname{LPic}_X$  such that  $\mathfrak{g}_{X_i}^i \cong \mathfrak{g}_{X_i}$ . Now consider  $M_{ji}$  as a  $\mathfrak{g}_{X_{ij}}^i$ -module using  $\psi_{ij}$ . Let  $f_{ij} : M_{ij} \to M_{ji}$  be isomorphisms of  $\phi_{ij}$ -twisted sheaves of  $\mathfrak{g}_{X_{ij}}^i$ -modules. Then if  $\{f_{ij}\}$  satisfies the condition  $[\partial f_{ij}] = \phi \in H^2(X, \mathbb{C}^*)$  there exists  $M \in \operatorname{Mod}(\mathfrak{g}_X, \phi)$ , unique to unique isomorphism, such that  $M_{X_i} \cong M_i$ .

The above discussion shows that the category of  $\phi$ -twisted sheaves on X form a stack over the category of schemes/k (or over the category of complex analytic spaces)[Gir71].

3.4. Inverse images. Let  $\rho : \mathfrak{g}_X \to \mathfrak{c}_X(M)$  be a  $\mathfrak{g}_X$ -module and set  $\mathfrak{g}_Y = \pi^+(\mathfrak{g}_X)$  (2.4.5). The pull-back functor  $\pi^! : \operatorname{Mod}(\mathfrak{g}_X) \to \operatorname{Mod}(\mathfrak{g}_Y)$  is defined by  $\pi^!(M) = \mathcal{O}_Y \otimes_{\pi^{-1}(\mathcal{O}_X)} \pi^{-1}(M)$  and the homomorphism

$$\pi^{!}(\rho):\mathfrak{g}_{Y} \to \mathfrak{c}_{Y}(\pi^{*}(M)),$$
  
$$\pi^{!}(\rho)(\phi \otimes \delta(x), \partial_{y}) = \partial \otimes \mathrm{id} + \phi \otimes \rho(\delta),$$

where  $(\phi \otimes \delta(x), \partial_y) \in \mathfrak{g}_Y = \pi^*(\mathfrak{g}_X) \times_{\pi^*(T_X)} T_Y$ . We check that  $\pi^!(\rho)$  is well-defined. Let  $\psi \in \pi^{-1}(\mathcal{O}_X), m \in \pi^{-1}(M)$ , and  $\bar{\psi}$  be the image of  $\psi$  in

<sup>&</sup>lt;sup>4</sup>We make one choice of  $\lambda$  for each  $\mathfrak{g}_X \in \operatorname{LPic}_X$ .

 $\mathcal{O}_Y$ . Since  $\partial_y(\bar{\psi}) = \phi \alpha(\delta(\bar{x}))(\psi)$  one gets

$$\begin{aligned} (\phi \otimes \delta(x), \partial_y)(\psi \otimes m) &= \partial_y(\psi) \otimes m + \psi \phi \otimes \delta(x) \cdot m \\ &= \phi \alpha(\delta(\bar{x}))(\psi) \otimes m + \bar{\psi} \phi \otimes \delta(x) \cdot m \\ &= \phi \otimes \alpha(\delta(x))(\psi)m + \bar{\psi} \phi \otimes \delta(x) \cdot m \\ &= \phi \otimes \delta(x) \cdot (\psi m) \\ &= (\phi \otimes \delta(x), \partial_y)(1 \otimes \psi m). \end{aligned}$$

Set

$$\mathcal{D}_{Y\to X} = \pi^!(\mathcal{D}(\mathfrak{g}_X)) = \mathcal{O}_Y \otimes_{\pi^{-1}(\mathcal{O}_X)} \pi^{-1}(\mathcal{D}(\mathfrak{g}_X)).$$

This is a  $(\mathfrak{g}_Y, \pi^{-1}(\mathfrak{g}_X))$ -bimodule in an obvious way, and one easily sees that  $\pi^!(\cdot)$  is isomorphic to  $\mathcal{D}_{Y \to X} \otimes_{\pi^{-1}(\mathcal{D}(\mathfrak{g}_X))} \pi^{-1}(\cdot)$ .

When  $M \in \operatorname{coh}(\mathfrak{g}_X)$ , then  $\pi^!(M)$  is a countable union of its coherent submodules, and if

$$\pi^*(\operatorname{Im}(\alpha:\mathfrak{g}_X\to T_X))\subseteq \operatorname{Im}(d\pi:T_Y\to\pi^*(T_X)),$$

then  $\pi^! : \operatorname{coh}(\mathfrak{g}_X) \to \operatorname{coh}(\mathfrak{g}_Y)$ . This is easily checked by showing that the stalks  $\pi^!(M)_y, y \in Y$ , are  $\mathfrak{g}_y$ -modules of finite type.

- Examples 3.4.1. (1) Let  $\pi : Y = \mathbf{A}^1 \to X = \mathbf{A}^2$  be the embedding of affine spaces  $y \mapsto (0, y)$  and  $M = \mathcal{D}(T_X) \in \operatorname{coh}(T_X)$ . Then  $\pi^+(T_X) = T_Y$  and  $\pi^!(M) = \mathcal{O}_Y \otimes_{\pi^{-1}(\mathcal{O}_X)} \pi^{-1}(M) = \mathcal{D}_Y[\partial_x]$ , (polynomials in  $\partial_x$ with coefficients in the ring of differential operators on Y). Clearly  $\pi^!(M) \notin \operatorname{coh}(T_Y)$ .
  - (2) Let  $\pi: Y = \mathbf{A}^2 \to X = \mathbf{A}^1$  be the projection on the first coordinate  $(x, y) \mapsto x$ . Then  $\pi^+(T_X) = T_Y$  and  $\pi^!(M) = \mathcal{O}_Y \otimes_{\pi^{-1}(\mathcal{O}_X)} \pi^{-1}(M) \in \operatorname{coh}(T_Y)$ .

Let  $Y \xrightarrow{f} Z \xrightarrow{g} X$  be a composition of morphisms. Choosing pullbacks  $\mathfrak{g}_Z = g^+(\mathfrak{g}_X)$ ,  $\mathfrak{g}_Y = (g \circ f)^+(\mathfrak{g}_X)$  and  $f^+(\mathfrak{g}_Z)$ , there is a canonical homomorphism  $c_{fg} : f^+(\mathfrak{g}_Z) \to \mathfrak{g}_Y$  which in general is not an isomorphism (see Lemma 2.4.6). Hence if N is a  $\mathfrak{g}_Z$ -module, its inverse image  $f^!(N)$  is in general not a  $\mathfrak{g}_Y$ -module. This will nevertheless not cause problems for composed inverse images.

**Lemma 3.4.2.** Let M be a  $\mathfrak{g}_X$ -module. There exists a canonical isomorphism of  $\mathfrak{g}_Y$ -modules

$$(g \circ f)^!(M) \cong f^!(g^!(M)).$$

Proof. Considered as  $\mathcal{O}_Y$ -modules the isomorphism is the canonical one  $\mathcal{O}_Y \otimes_{(g \circ f)^{-1}(\mathcal{O}_X)} (g \circ f)^{-1}(M) \cong \mathcal{O}_Y \otimes_{f^{-1}(\mathcal{O}_Z)} f^{-1}(\mathcal{O}_Z \otimes_{g^{-1}(\mathcal{O}_X)} g^{-1}(M))$ , and this also defines the action of  $\mathfrak{g}_Y$  on the right side. One should verify that the  $f^+(\mathfrak{g}_Z)$ -action on the right side is compatible with this  $\mathfrak{g}_Y$ -action and the morphism  $c_{fg} : f^+(\mathfrak{g}_Z) \to \mathfrak{g}_Y$ , but we omit these straightforward details.  $\Box$ 

We thus can and will always consider  $f^!(g^!(M))$  as a  $\mathfrak{g}_Y$ -module. The derived version of  $\pi^!$  is the functor

$$L\pi^{!}: D^{-}_{\mathrm{coh}}(\mathfrak{g}_{X}) \to D^{-}_{\mathrm{qcoh}}(\mathfrak{g}_{Y}), \quad M^{\bullet} \mapsto \mathcal{D}_{Y \to X} \otimes^{L}_{\pi^{-1}(\mathcal{D}(\mathfrak{g}_{X}))} \pi^{-1}(M^{\bullet})$$
29

and if X is non-singular  $L\pi^!$  defines a functor  $D^b_{\rm coh}(\mathfrak{g}_X) \to D^b_{\rm qcoh}(\mathfrak{g}_Y)$ . One can compute  $L\pi^!(M^{\bullet})$  either by taking a resolution of the  $\mathfrak{g}_X$ -module M which is flat as  $\mathcal{O}_X$ -module or a resolution of  $\mathcal{D}_{Y\to X}$  in the category of  $(\mathfrak{g}_Y, \pi^{-1}(\mathfrak{g}_X))$ -bimodules consisting of modules which are flat as right  $\pi^{-1}(\mathfrak{g}_X)$ -module. Abusing the notation slightly we again use  $\pi^!$  for  $L\pi^!$ , so to define  $\pi^!$  completely one thus has to choose one resolution for each  $M^{\bullet} \in D^-(\mathfrak{g}_X)$ . If  $\mathfrak{g}_X$  is locally free, then by the Poincaré-Birkhoff-Witt theorem for Lie algebroids (*Th.* 2.3.3)  $\mathcal{D}(\mathfrak{g}_X)$  is locally free over  $\mathcal{O}_X$ , so a resolution  $F^{\bullet} \to M^{\bullet}$  which is flat as  $\mathfrak{g}_X$ -module is also flat as  $\mathcal{O}_X$ -module; hence

$$\mathcal{D}_{Y \to X} \otimes_{\pi^{-1}(\mathcal{D}(\mathfrak{g}_X))}^{L} \pi^{-1}(M^{\bullet}) = \mathcal{D}_{Y \to X} \otimes_{\pi^{-1}(\mathcal{D}(\mathfrak{g}_X))} \pi^{-1}(F^{\bullet})$$
$$= \mathcal{O}_Y \otimes_{\pi^{-1}(\mathcal{O}_X)} \pi^{-1}(F^{\bullet})$$
$$= \mathcal{O}_Y \otimes_{\pi^{-1}(\mathcal{O}_X)}^{L} M^{\bullet},$$

and  $H^{-i}(\pi^!(M^{\bullet})) = Tor_i^{\pi^{-1}(\mathcal{O}_X)}(\mathcal{O}_Y, \pi^{-1}(M^{\bullet})).$ 

**Proposition 3.4.3.**  $(\mathfrak{g}_X \text{ is locally free over } \mathcal{O}_X)$  Let  $Y \xrightarrow{f} Z \xrightarrow{g} X$  be a composition of morphisms. Then there exists a canonical isomorphism of functors  $D^-(\mathfrak{g}_X) \to D^-(\mathfrak{g}_Y)$ 

$$(g \circ f)^!(\cdot) \to f^! \circ g^!(\cdot).$$

See [Bjö93, Ch. II, Th. 2.3.21] and [BGK<sup>+</sup>87, VI, Prop 4.3] for proofs when  $\mathfrak{g}_X = T_X$  and all spaces are non-singular. The proof below is very similar.

*Proof.* By the above discussion, since  $\mathfrak{g}_X$  is locally free, there exists a resolution  $F^{\bullet}$  of  $M^{\bullet} \in D^-(\mathfrak{g}_X)$  such that the terms  $F^i$  are flat as  $\mathcal{O}_X$ -modules. Since flatness is preserved under base change and  $f^!(\cdot) \cong \mathcal{O}_Y \otimes_{f^{-1}(\mathcal{O}_X)}^L(\cdot)$  (restricted to  $D^-(\mathfrak{g}_Z) \subset D^-(\mathcal{O}_Z)$ ) this gives:

$$f^{!}(g^{!}(M^{\bullet})) = \mathcal{D}_{Y \to Z} \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_{Z}))}^{L} f^{-1}(\mathcal{D}_{Z \to X} \otimes_{g^{-1}(\mathcal{D}(\mathfrak{g}_{X}))}^{L} g^{-1}(M^{\bullet}))$$

$$= \mathcal{O}_{Y} \otimes_{f^{-1}(\mathcal{O}_{Z})}^{L} f^{-1}(\mathcal{O}_{Z} \otimes_{g^{-1}(\mathcal{O}_{X})} g^{-1}(F^{\bullet}))$$

$$= \mathcal{O}_{Y} \otimes_{f^{-1}(\mathcal{O}_{Z})}^{L} f^{-1}(\mathcal{O}_{Z} \otimes_{g^{-1}(\mathcal{O}_{X})} g^{-1}(F^{\bullet}))$$

$$= \mathcal{O}_{Y} \otimes_{(g \circ f)^{-1}(\mathcal{O}_{X})}^{L} (g \circ f)^{-1}(F^{\bullet})$$

$$= \mathcal{D}_{Y \to X} \otimes_{(g \circ f)^{-1}(\mathcal{D}(\mathfrak{g}_{X}))}^{L} (g \circ f)^{-1}(F^{\bullet})$$

$$= (g \circ f)^{!}(M^{\bullet}).$$

3.5. Direct images. In this section all spaces are non-singular, and all morphism  $\pi : Y \to X$  of spaces are such that the functor  $\pi_*$  on the category of sheaves on Y has finite cohomological dimension.

Let  $\pi : Y \to X$  be a morphism of spaces and set  $\mathfrak{g}_Y = \pi^+(\mathfrak{g}_X)$ . We will alway assume that  $\mathfrak{g}_Y$  is *locally free*. Following Kashiwara's procedure for  $T_X$ -modules we will define direct images of  $\mathfrak{g}_Y$ -modules, using the

 $(\mathfrak{g}_Y, \pi^{-1}(\mathfrak{g}_X))$ -bimodule  $\mathcal{D}_{Y \to X}$ . The direct image of a bounded complex of right  $\mathfrak{g}_Y$ -modules  $N^{\bullet}$  is defined by

$$\pi^r_+(N) = R\pi_*(N^\bullet \otimes^L_{\mathcal{D}(\mathfrak{g}_Y)} \mathcal{D}_{Y \to X}).$$

To define the direct image of a left module one first notices that by (3.2.3) and (3.2.5)

$$\mathcal{D}_{X\leftarrow Y} = \omega_Y \otimes_{\mathcal{O}_Y} \pi^*(\omega_X^{-1}) \otimes_{\mathcal{O}_Y} \mathcal{D}_{Y\to X}.$$

is a  $(\pi^{-1}(\mathfrak{g}_X),\mathfrak{g}_Y)$ -bimodule. Then define  $\pi_+: D^b(\mathfrak{g}_Y) \to D^b(\mathfrak{g}_X)$  by

$$\pi_+(M^{\bullet}) = R\pi_*(\mathcal{D}_{X\leftarrow Y} \otimes^L_{\mathcal{D}(\mathfrak{g}_Y)} M^{\bullet}) \in D^b(\mathfrak{g}_X)$$

when  $M^{\bullet} \in D^{b}(\mathfrak{g}_{X})$ . Since  $\mathfrak{g}_{Y}$  is locally free and Y is non-singular it follows that the functor  $\mathcal{D}_{X \leftarrow Y} \otimes_{\mathcal{D}(\mathfrak{g}_{Y})}^{L}$  has finite homological dimension (see for instance [Käl96]), so  $\pi_{+}(M^{\bullet})$  does indeed belong to  $D^{b}(\mathfrak{g}_{X})$ . Note that by the projection formula:

(3.5.1) 
$$\pi_+(M^{\bullet}) = \omega_X^{-1} \otimes_{\mathcal{O}_X} \pi_+^r(\omega_Y \otimes_{\mathcal{O}_Y} M^{\bullet}).$$

We now discuss the basic functorial property of the direct image functor. Consider a composition of morphims of spaces

 $Y \xrightarrow{f} Z \xrightarrow{g} X$ 

and put  $\mathfrak{g}_Z = g^+(\mathfrak{g}_X)$ , and  $\mathfrak{g}_Y = f^+(\mathfrak{g}_Z)$ . Then in general

$$(g \circ f)_*(\mathcal{D}_{X \leftarrow Y} \otimes_{\mathcal{D}(\mathfrak{g}_Y)} M) \neq g_*(\mathcal{D}_{X \leftarrow Z} \otimes_{\mathcal{D}(\mathfrak{g}_Z)} f_*(\mathcal{D}_{Z \leftarrow Y} \otimes_{\mathcal{D}(\mathfrak{g}_Y)} M)).$$

So the non-derived direct image does not compose well, but we will see that the situation is more satisfying when working with derived categories.

**Proposition 3.5.1.** ( $\mathfrak{g}_X$  and  $\mathfrak{g}_Z$  are locally free) As functors  $D^b(\mathfrak{g}_Y) \to D^b(\mathfrak{g}_X)$  we have

$$(g \circ f)_+(\cdot) \cong g_+f_+(\cdot).$$

The assumption that  $\mathfrak{g}_X$  and  $\mathfrak{g}_Z$  are locally free are satisfied for instance if  $\mathfrak{g}_X$  is locally free and g is submersive, or in the situation of Lemma 2.4.8.

*Proof.* The proof is similar to [Bjö93, Ch. II Th. 2.3.21].

a) We first prove

$$(3.5.2) \qquad \mathcal{D}_{Y \to X} \cong \mathcal{D}_{Y \to Z} \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))} f^{-1}(\mathcal{D}_{Z \to X}) \\ \cong \mathcal{D}_{Y \to Z} \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))}^L f^{-1}(\mathcal{D}_{Z \to X})$$

where the isomorphism holds in the category of (complexes of)  $(\mathfrak{g}_Y, (g \circ f)^{-1}(\mathfrak{g}_X))$ -bimodules.

Begin with the first isomorphism. Note that the right side need not be a  $\mathfrak{g}_Y$ -module when  $c_{fg}: f^+(\mathfrak{g}_Z) \to \mathfrak{g}_Y$  (2.4.1) is not an isomorphism, but both sides are  $f^+(\mathfrak{g}_Z)$ -modules, so if we prove the isomorphisms in the category of  $(f^+(\mathfrak{g}_Z), (g \circ f)^{-1}(\mathfrak{g}_X))$ -modules, then since the  $f^+(\mathfrak{g}_Z)$ -action in the left side is determined by the  $\mathfrak{g}_Y$ -action (using the homomorphism  $c_{fg}$ ), we may regard the isomorphisms in the category of  $(\mathfrak{g}_Y, (g \circ f)^{-1}(\mathfrak{g}_X))$ -modules. The morphism

$$G: \mathcal{D}_{Y \to X} \to \mathcal{D}_{Y \to Z} \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))} f^{-1}(\mathcal{D}_{Z \to X})$$
31

defined by

$$k(y) \otimes_{(g \circ f)^{-1}(\mathcal{O})} (g \circ f)^{-1}(P(x))$$
  
 
$$\mapsto k(y) \otimes_{f^{-1}(\mathcal{O}_Z)} f^{-1}(1_Z) \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))} f^{-1}(1_Z \otimes_{g^{-1}(\mathcal{O}_X)} g^{-1}(P(x)))$$

where  $k(y) \in \mathcal{O}_Y$  and  $P(x) \in \mathcal{D}(\mathfrak{g}_X)$ , obviously is an isomorphism in the category of  $(\mathcal{O}_Y, (g \circ f)^{-1}(\mathfrak{g}_X))$ -bimodules. It remains to see that it is  $f^+(\mathfrak{g}_Z)$ -linear (see (3.4)). In the notation of (2.4.1), letting r(y) = $(h(y) \otimes_{f^{-1}(\mathcal{O}_Z)} f^{-1}(\phi(z) \otimes_{g^{-1}(\mathcal{O}_X)} g^{-1}(\delta(x)), \partial_z), \partial_y) \in f^+(\mathfrak{g}_Z)$ , we have

$$c_{fg}(r(y)) = (h(y)\bar{\phi}(y) \otimes_{(g \circ f)^{-1}(\mathcal{O}_X)} (g \circ f)^{-1}(\delta(x)), \partial_y),$$

where  $\bar{\phi}(y)$  is the image of  $\phi(z) \in \mathcal{O}_Z$  in  $\mathcal{O}_Y$ . Therefore

$$\begin{split} & G(r(y) \cdot (k(y) \otimes_{(g \circ f)^{-1}(\mathcal{D}(\mathfrak{g}_X))} (g \circ f)^{-1}(P))) \\ &= G(\partial_y(k) \otimes_{(g \circ f)^{-1}(\mathcal{D}(\mathfrak{g}_X))} \otimes (g \circ f)^{-1}(P(x)) \\ &+ k(y)h(y)\bar{\phi}(y) \otimes_{(g \circ f)^{-1}(\mathcal{D}(\mathfrak{g}_X))} \cdot (g \circ f)^{-1}(\delta(x) \cdot P)) \\ &= \partial_y(k) \otimes_{f^{-1}(\mathcal{O}_Z)} f^{-1}(1_Z) \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))} f^{-1}(1_Z \otimes_{g^{-1}(\mathcal{O}_X)} g^{-1}(P(x))) \\ &+ k(y)h(y)\bar{\phi}(y) \otimes_{f^{-1}(\mathcal{O}_Z)} f^{-1}(1_Z) \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))} f^{-1}(1_Z \otimes_{g^{-1}(\mathcal{O}_X)} g^{-1}(\delta(x)P(x))) \\ &= \partial_y(k) \otimes_{f^{-1}(\mathcal{O}_Z)} f^{-1}(1_Z) \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))} f^{-1}(1_Z \otimes_{g^{-1}(\mathcal{O}_X)} g^{-1}(\delta(x)P(x))) \\ &+ k(y)h(y) \otimes_{f^{-1}(\mathcal{O}_Z)} f^{-1}(1_Z) \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))} f^{-1}(\phi(z) \otimes_{g^{-1}(\mathcal{O}_X)} g^{-1}(\delta(x)P(x))) \\ &= r(y) \cdot k(y) \otimes_{f^{-1}(\mathcal{O}_Z)} f^{-1}(1_Z) \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))} f^{-1}(1_Z \otimes_{g^{-1}(\mathcal{O}_X)} g^{-1}(P(x))) \\ &= r(y) \cdot G(k(y) \otimes_{(g \circ f)^{-1}(\mathcal{D}(\mathfrak{g}_X))} (g \circ f)^{-1}(P)). \end{split}$$

This proves the first isomorphism in (3.5.2). To see the second isomorphism, first note that the  $\mathcal{O}_Z$ -module  $\mathcal{D}(\mathfrak{g}_Z)$  is locally free since  $\mathfrak{g}_Z$  is locally free (*Th.* 2.3.3), so flat  $\mathfrak{g}_Z$ -modules are flat as  $\mathcal{O}_Z$ -modules, and then since  $\mathcal{D}(\mathfrak{g}_X)$ is also locally free it follows that  $\mathcal{D}_{Z\to X} = g^*(\mathcal{D}(\mathfrak{g}_X))$  is flat as  $\mathcal{O}_Z$ -module. This gives the following isomorphisms in  $D^b(\mathfrak{g}_Y)$ , where  $F^{\bullet}$  is a flat resolution of the  $\mathfrak{g}_Z$ -module  $\mathcal{D}_{Z\to X}$ :

$$\mathcal{D}_{Y \to Z} \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))}^{L} f^{-1}(\mathcal{D}_{Z \to X}) \cong \mathcal{D}_{Y \to Z} \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))} f^{-1}(F^{\bullet})$$
$$\cong \mathcal{O}_Y \otimes_{f^{-1}(\mathcal{O}_Z)} f^{-1}(F^{\bullet})$$
$$\cong \mathcal{O}_Y \otimes_{f^{-1}(\mathcal{O}_Z)} f^{-1}(\mathcal{D}_{Z \to X})$$
$$\cong \mathcal{D}_{Y \to Z} \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_Z))} f^{-1}(\mathcal{D}_{Z \to X}).$$

b) Let first N be a right  $\mathfrak{g}_Y$ -module. Applying the projection formula and (3.5.2) one gets

$$g_{+}^{r}(f_{+}^{r}(N)) = Rg_{*}(Rf_{*}(N \otimes_{\mathcal{D}(\mathfrak{g}_{Y})}^{L} \mathcal{D}_{Y \to Z}) \otimes_{\mathcal{D}(\mathfrak{g}_{Z})}^{L} \mathcal{D}_{Z \to X})$$

$$= Rg_{*}(Rf_{*}(N \otimes_{\mathcal{D}(\mathfrak{g}_{Y})}^{L} \mathcal{D}_{Y \to Z} \otimes_{f^{-1}(\mathcal{D}(\mathfrak{g}_{Z}))}^{L} f^{-1}(\mathcal{D}_{Z \to X})))$$

$$= Rg_{*} \circ Rf_{*}(N \otimes_{\mathcal{D}(\mathfrak{g}_{Y})}^{L} \mathcal{D}_{Y \to X})$$

$$= R(g \circ f)_{*}(N \otimes_{\mathcal{D}(\mathfrak{g}_{Y})}^{L} \mathcal{D}_{Y \to X}) = (g \circ f)_{+}^{r}(N),$$

proving the composition formula in the proposition for right  $\mathfrak{g}_Y$ -modules. Let now M be a left  $\mathfrak{g}_Y$ -module. By (3.5.1)

$$g_{+}(f_{+}(M)) = \omega_{X}^{-1} \otimes_{\mathcal{O}_{X}} g_{+}^{r}(\omega_{Z} \otimes_{\mathcal{O}_{Z}} \omega_{Z}^{-1} \otimes_{\mathcal{O}_{Z}} f_{+}^{r}(\omega_{Y} \otimes_{\mathcal{O}_{Y}} M))$$
  
$$= \omega_{X}^{-1} \otimes_{\mathcal{O}_{X}} g_{+}^{r}(f_{+}^{r}(\omega_{Y} \otimes_{\mathcal{O}_{Y}} M)))$$
  
$$= \omega_{X}^{-1} \otimes_{\mathcal{O}_{X}} (g \circ f)_{+}^{r}(\omega_{Y} \otimes_{\mathcal{O}_{Y}} M))$$
  
$$= (g \circ f)_{+}(M).$$

Let  $D^b_{\operatorname{coh},l}(\mathfrak{g}_Y) \subset D^b_{\operatorname{coh}}(\mathfrak{g}_Y)$  be the subcategory of bounded complexes  $M^{\bullet}$ whose homology  $H^{\bullet}(M^{\bullet})$ , locally in X, is generated by an  $\mathcal{O}_Y$ -coherent submodule. When X is a noetherian space, then  $D^b_{\operatorname{coh},l}(\mathfrak{g}_X) = D^b_{\operatorname{coh}}(\mathfrak{g}_X)$ .

**Proposition 3.5.2.** ( $\mathfrak{g}_Y$  and  $\mathfrak{g}_X$  are locally free) Let  $\pi : Y \to X$  be a proper morphism of spaces. We then have

$$\pi_+(D^b_{\operatorname{coh},l}(\mathfrak{g}_Y)) \subset D^b_{\operatorname{coh},l}(\mathfrak{g}_X).$$

*Proof.* The proof in [Bjö93, Th. 2.8.1] works fine so we make only a few points for comparison.

1) The theorem holds for right  $\mathfrak{g}_Y$ -modules of the form  $K \otimes_{\mathcal{O}_Y} \mathcal{D}(\mathfrak{g}_Y)$  when  $K \in \operatorname{coh}(\mathcal{O}_Y)$ : If  $F^{\bullet} \to K$  is a flat resolution of K, then  $F^{\bullet} \otimes_{\mathcal{O}_Y} \mathcal{D}(\mathfrak{g}_Y)$  is a flat resolution of  $K \otimes_{\mathcal{O}_Y} \mathcal{D}(\mathfrak{g}_Y)$ , so

$$\pi^{r}_{+}(K \otimes_{\mathcal{O}_{Y}} \mathcal{D}(\mathfrak{g}_{Y})) = R\pi_{*}(K \otimes_{\mathcal{O}_{Y}} \mathcal{D}(\mathfrak{g}_{Y}) \otimes_{\mathcal{D}(\mathfrak{g}_{Y})}^{L} \mathcal{D}_{Y \to X})$$

$$\cong R\pi_{*}(F^{\bullet} \otimes_{\pi^{-1}(\mathcal{O}_{X})} \pi^{-1}(\mathcal{D}(\mathfrak{g}_{X})))$$

$$\cong R\pi_{*}(F^{\bullet}) \otimes_{\mathcal{O}_{X}} \mathcal{D}(\mathfrak{g}_{X})$$

$$\cong R\pi_{*}(K) \otimes_{\mathcal{O}_{X}} \mathcal{D}(\mathfrak{g}_{X}),$$

where the third line follows from the projection formula since  $\mathcal{D}(\mathfrak{g}_X)$  is locally free over  $\mathcal{O}_X$ . By properness, Grauert's or Grothendieck's theorem implies that  $R\pi_*(K) \in D^b_{\mathrm{coh}}(\mathcal{O}_X)$ , hence  $\pi^r_+(K \otimes_{\mathcal{O}_Y} \mathcal{D}(\mathfrak{g}_Y)) \in D^b_{r,\mathrm{coh}}(\mathfrak{g}_X)$ (complexes of right  $\mathfrak{g}_X$ -modules).

2) Let M be a left  $\mathfrak{g}_Y$ -module and put  $l = \operatorname{rk} \mathfrak{g}_Y$ . Consider the induced  $(\mathfrak{g}_Y, \mathfrak{g}_Y)$ -bimodule  $M_{\operatorname{ind}} = M \otimes_{\mathcal{O}_Y} \mathcal{D}(\mathfrak{g}_Y)$  (the left action is (3.2.1) and the right action is multiplication in  $\mathcal{D}(\mathfrak{g}_Y)$ ). Put  $\Omega_Y^i(\mathfrak{g}_Y) = Hom_{\mathcal{O}_Y}(\bigwedge^i \mathfrak{g}_Y, \mathcal{O}_Y)$ ; since  $\mathfrak{g}_Y$  is locally free the de Rham complex  $\Omega^{\bullet}(M_{\operatorname{ind}}) = \Omega^i(\mathfrak{g}_Y) \otimes_{\mathcal{O}_Y} M \otimes_{\mathcal{O}_Y} \mathcal{D}(\mathfrak{g}_Y)$  (3.1.1). Then  $\Omega_Y^{\bullet}(\mathfrak{g}_Y)$  is an acyclic left resolution of the right  $\mathfrak{g}_Y$ -module  $\Omega^l(\mathfrak{g}_X, M) = \Omega^l(\mathfrak{g}_Y) \otimes_{\mathcal{O}_Y} M$  (3.2.5). This is proven as follows: First filter the complex  $\Omega^{\bullet}(M_{\operatorname{ind}})$  by the subcomplexes  $F_k^{\bullet}$ , where  $F_k^i = \Omega^i(\mathfrak{g}_Y) \otimes_{\mathcal{O}_Y} M \otimes_{\mathcal{O}_Y} \mathcal{D}^{i+k}(\mathfrak{g}_Y)$ . Then the associated graded complex  $\operatorname{gr} F^{\bullet} \cong M \otimes_{\mathcal{O}_Y} K^{\bullet}$ , where  $K^{\bullet}$  is the Koszul complex of the symmetric algebra  $S(\mathfrak{g}_Y)$  of the locally free  $\mathcal{O}_Y$ -module  $\mathfrak{g}_Y (K^i = S(\mathfrak{g}_Y) \otimes_{\mathcal{O}_Y} \bigwedge^{-i} \mathfrak{g}_Y)$ , so it is acyclic in degree  $\leq -1$ . But this implies that  $F^{\bullet}$  is acyclic in degrees  $\leq -1$ , and one sees that its homology in degree 0 is  $\Omega^l(\mathfrak{g}_X, M)$ .

3) The module  $M = \bigcup_{k\geq 0} M_k$  is filtered by  $\mathcal{O}_Y$ -coherent submodules  $M_k$  so that  $\mathfrak{g}_Y \cdot M_k \subset M_k$ . Then for each integer  $k \geq 0$  the de Rham complex  $\Omega^{\bullet}(M_{\mathrm{ind}})$  contains a subcomplex  $S_k^{\bullet}(M)$  (the Spencer complex of the filtered module M) where

$$S_k^i(M) = \Omega^i(\mathfrak{g}_Y) \otimes_{\mathcal{O}_Y} M_{k+i} \otimes_{\mathcal{O}_Y} \mathcal{D}(\mathfrak{g}_Y).$$

One proves that locally in Y when  $k \gg 1$  then  $S_k^{\bullet}(M)$  is quasi-isomorphic to  $\Omega^{\bullet}(M_{\text{ind}})$ ; but as  $\pi$  is proper this quasi-isomorphism holds locally in X when  $k \gg 1$ , and since the assertion to be proven is local in X we may assume that k is choosen so that, by 2),

$$S_k^{ullet}(M) \cong \Omega^l(\mathfrak{g}_Y) \otimes_{\mathcal{O}_Y} M$$

in  $D^b_r(\mathfrak{g}_Y)$ .

4) If a right  $\mathfrak{g}_Y$ -module N has a bounded acyclic resolution

$$F^l \to F^{l-1} \to \dots \to F^0 \to N$$

such that  $\pi^r_+(F^i) \in D^b_{r,\mathrm{coh}}(\mathfrak{g}_X)$ , then  $\pi^r_+(N) \in D^b_{r,\mathrm{coh}}(\mathfrak{g}_X)$ . This is proven by a straighforward induction in l. From this it follows by 1) and 3) that  $\pi^r_+(\Omega^l(\mathfrak{g}_Y) \otimes_{\mathcal{O}_Y} M) \in D^b_{r,\mathrm{coh}}(\mathfrak{g}_X)$ . Then  $M_1 := \omega_Y \otimes_{\mathcal{O}_Y} \Omega^l(\mathfrak{g}_Y)^{-1} \otimes_{\mathcal{O}_Y} M$  is a coherent left  $\mathfrak{g}_Y$ -module and (see (3.2))

$$\pi_{+}(M) = \omega_{X}^{-1} \otimes_{\mathcal{O}_{X}} \pi_{+}^{r}(\omega_{Y} \otimes_{\mathcal{O}_{Y}} M)$$
  
$$= \omega_{X}^{-1} \otimes_{\mathcal{O}_{X}} \pi_{+}^{r}(\Omega^{l}(\mathfrak{g}_{Y}) \otimes_{\mathcal{O}_{Y}} M_{1}) \in D^{b}_{\mathrm{coh}}(\mathfrak{g}_{X}).$$

Let  $M^{\bullet}[d]$  be the translation of a complex to the left d steps, and  $d_{Y,X} = \dim Y - \dim X$  be the relative dimension of a morphism of pure dimensional spaces  $\pi: Y \to X$ .

**Theorem 3.5.3.** ( $\mathfrak{g}_Y$  and  $\mathfrak{g}_X$  are locally free) Let  $\pi : Y \to X$  be a proper morphism of non-singular spaces. Then

$$R\pi_*RHom_{\mathfrak{g}_Y}(M^{\bullet},\pi^!(N^{\bullet})[\mathrm{d}_{Y,X}]) = RHom_{\mathfrak{g}_X}(\pi_+(M^{\bullet}),N^{\bullet})$$

when  $M^{\bullet} \in D^{b}_{\operatorname{coh}}(\mathfrak{g}_{Y})$  and  $N^{\bullet} \in D^{b}_{\operatorname{coh}}(\mathfrak{g}_{X})$ .

This adjointness property is proven in a similar way as for  $\mathcal{D}$ -modules [BGK<sup>+</sup>87, VIII, Prop. 9.10], so we omit the quite long proof. Let us only remark that it suffices to construct the trace morphism  $\operatorname{tr} : \pi_+\pi^!(M^{\bullet}[d_{Y,X}]) \to M^{\bullet}$  and for this it suffices to consider closed embeddings and projections separately. To see this, factorize  $\pi = p \circ i$ , where  $i : Y \to Y \times X$  is the graph embedding and  $p : Y \times X \to X$  is the projection on X. Then if  $\operatorname{tr}_i : i_+ i^!(N^{\bullet})[-\dim X] \to N^{\bullet}$  and  $\operatorname{tr}_p : p_+ p^!(M^{\bullet})[\dim Y] \to M^{\bullet}$  are defined for any  $N^{\bullet} \in D^b_{\operatorname{coh}}(\mathfrak{g}_{Y \times X}), M^{\bullet} \in D^b_{\operatorname{coh}}(\mathfrak{g}_X)$ , we first get  $\operatorname{tr}_i : i_+ i^! p^!(M^{\bullet}) \to p^!(M^{\bullet})$ , and applying  $p_+$ 

$$\pi_{+}\pi^{!}(M^{\bullet}) = p_{+}i_{+}i^{!}p^{!}(M^{\bullet}) \xrightarrow{p_{+}(\operatorname{tr}_{i})} p_{+}p^{!}(M^{\bullet}) \xrightarrow{\operatorname{tr}_{p}} M^{\bullet},$$

where the equality is because  $\pi^{!} = i^{!}p^{!}$  and  $\pi_{+} = p_{+}i_{+}$  (Props. 3.5.1 and 3.5.1). Note that in the situation of Theorem 3.5.4 below tr<sub>i</sub> is an isomorphism.

Letting  $M^{\bullet} \in D^{b}(\mathfrak{g}_{X})$  and  $Y \subset X$  be a closed algebraic (analytic) subset one has a distinguished triangle in  $D^{b}(\mathcal{O}_{X})$  (2.1.1)

(3.5.3) 
$$R\Gamma_{[Y]}(M^{\bullet}) \to M^{\bullet} \to M^{\bullet}(*S) \to .$$

Since  $\mathcal{D}(\mathfrak{g}_X)$  is locally free over  $\mathcal{O}_X$ , injective  $\mathfrak{g}_X$ -modules are injective as  $\mathcal{O}_X$ -modules, so an injective resolution of  $M^{\bullet}$  is acyclic for the functor  $\Gamma_{[Y]}(\cdot)$ , hence (3.5.3) is a distinguished triangle in  $D^b(\mathfrak{g}_X)$ .

Let now  $\pi : Y \to X$  be a closed immersion of non-singular spaces and  $D_Y^b(\mathfrak{g}_X) \subset D^b(\mathfrak{g}_X)$  be the subcategory of complexes  $M^{\bullet}$  such that  $\operatorname{supp} H^{\bullet}(M^{\bullet}) \subseteq i(Y)$ . Next result is basically due to Kashiwara.

**Theorem 3.5.4.** Let  $Y \to X$  be a closed immersion of non-singular spaces and  $\mathfrak{g}_X$  a locally free and transitive Lie algebroid. Then we have a distinguished triangle

(3.5.4) 
$$\pi_+\pi^!(M^{\bullet})[\mathrm{d}_{Y,X}] \to M^{\bullet} \to M^{\bullet}(*Y) \to$$

The functor  $\pi_+$  defines an equivalence of categories  $D^b(\mathfrak{g}_Y) \to D^b_Y(\mathfrak{g}_X)$ with quasi-inverse  $\pi^!(\cdot)[d_{Y,X}]$ ;  $\pi_+$  preserves coherence and the restriction of  $\pi^!(\cdot)[d_{Y,X}]$  to  $D^b_Y(\mathfrak{g}_X)$  also preserves coherence.

Because of Theorems 3.5.3 and 3.5.4 one may find it convenient to include the translation in the definition of the inverse image functor, redefining it to  $M \mapsto \pi^! (M^{\bullet})[\mathrm{d}_{Y,X}]^{.5}$ 

*Proof.* When  $\mathfrak{g}_X = T_X$  the proof can be found in [Bjö93] or [BGK<sup>+</sup>87, VI, Th. 7.13] so we only make a sketch to see how to get (3.5.4), following the proof in [loc. cit.].

a) Put  $d = -d_{Y,X}$  and let  $\{f_1, \ldots, f_d\}$  be a regular sequence locally generating the ideal I of Y. Let  $\partial_1, \ldots, \partial_d$  be a dual basis of the k-vector space  $\sum_{i=1} k^d df_i$ . Locally we have  $\mathcal{O}_Y \otimes_{\pi^{-1}(\mathcal{O}_X)} \pi^{-1}(\mathcal{O}_X df_1 \wedge \cdots \wedge df_d) = \omega_Y^{-1} \otimes_{\pi^{-1}(\mathcal{O}_X)} \omega_X$ .

b) Using the Koszul complex based on the regular sequence  $\{f_1, \ldots, f_d\}$  to resolve the  $\pi^{-1}(\mathcal{O}_X)$ -module  $\mathcal{O}_Y$  one gets, by a),

$$H^0(\pi^!(M)[d_{Y,X}]) = M^I \otimes_{\mathcal{O}_Y} \omega_Y^{-1} \otimes_{\pi^{-1}(\mathcal{O}_X)} \pi^{-1}(\omega_X),$$

where  $M^I = \{m \in M : I \cdot m = 0\} \subset M$ .

c) There are isomorphisms of  $\mathfrak{g}_Y$ -modules

$$\mathcal{D}_{Y \to X} = \pi^*(\mathcal{D}(\mathfrak{g}_X)) \cong \mathcal{D}(\mathfrak{g}_X) / I\mathcal{D}(\mathfrak{g}_X) \cong \mathcal{D}(\mathfrak{g}_Y) \otimes_k \mathbf{C}[\partial_1, \dots, \partial_d].$$

In particular  $\mathcal{D}_{Y \to X}$  is locally free over  $\mathfrak{g}_Y$ , and so  $\pi_+$  is exact.

d) Since  $\pi_*$  is exact, c) implies

$$\pi_{+}H^{0}(\pi^{!}(M)[d_{Y,X}])$$

$$= \pi_{*}(\pi^{-1}(\omega_{X}^{-1}) \otimes_{\pi^{-1}(\mathcal{O}_{X})} \mathcal{D}_{Y \to X} \otimes_{\mathcal{O}_{Y}} \omega_{Y} \otimes_{\mathcal{D}(\mathfrak{g}_{Y})} H^{0}(\pi^{!}(M)[d_{Y,X}]))$$

$$= \omega_{X}^{-1} \otimes_{\mathcal{O}_{X}} \pi_{*}(\frac{\mathcal{D}(\mathfrak{g}_{X})}{I\mathcal{D}(\mathfrak{g}_{X})} \otimes_{\pi^{-1}(\mathcal{O}_{X})} \omega_{Y} \otimes_{\mathcal{D}(\mathfrak{g}_{Y})} (M^{I} \otimes_{\mathcal{O}_{Y}} \omega_{Y}^{-1} \otimes_{\pi^{-1}(\mathcal{O}_{X})} \pi^{-1}(\omega_{X})))$$

Then the mapping  $\mathcal{D}(\mathfrak{g}_X) \times M^I \to M$ ,  $(P, m) \mapsto P \cdot M$  induces a homomorphism of  $\mathfrak{g}_X$ -modules

$$\mu : \pi_+ H^0(\pi^!(M)[d_{Y,X}]) \to \Gamma_{[Y]}(M)$$

Using the transitivity of  $\mathfrak{g}_X$  one proves that  $\mu$  is an isomorphism in the same way as [Prop. 7.10, loc. cit.].

e) By d) there exists an isomorphism of functors

$$\Gamma_{[Y]}(\cdot) \cong \pi_+ H^0(M^!(\cdot)[d_{Y,X}])$$

<sup>&</sup>lt;sup>5</sup>Which is the convention in [loc. cit.].

on the category  $\operatorname{Mod}(\mathfrak{g}_X)$ . Since  $\mathcal{D}(\mathfrak{g}_X)$  is locally free over  $\mathcal{O}_X$ , injective  $\mathfrak{g}_X$ -modules are injective as  $\mathcal{O}_X$ -modules; hence, as  $\pi_+$  is exact, injective resolutions are acyclic both for  $\Gamma_{[Y]}(\cdot)$  and  $\pi_+H^0(\pi^!(\cdot)) = \pi_+(Hom_{\mathcal{O}_X}(\mathcal{O}_X/I, \cdot))$ . Therefore  $\mu$  induces an isomorphism of functors

$$R\Gamma_{[Y]}(\cdot) \cong \pi_+ \pi^!(\cdot)[d_{Y,X}]$$

on  $D^b(\mathfrak{g}_X)$ . Hence the distinguished triangle (3.5.3) can be identified with (3.5.4).  $\Box$ 

## 4. Smooth modules

4.1. Elementary properties. Let  $(\mathcal{O}_X, \mathfrak{g}_X, \alpha)$  be a Lie algebroid on a space X. We are interested in  $\mathfrak{g}_X$ -modules which are locally of finite type with respect to some distinguished sub-algebroid  $\mathfrak{g}'_X$  of  $\mathfrak{g}_X$ . One example to keep in mind is  $\mathfrak{g}_X(I) \subseteq \mathfrak{g}_X$  for some ideal I of  $\mathcal{O}_X$  (see 2.2).

**Definition 4.1.1.** Let  $M \in \operatorname{coh}(\mathfrak{g}_X)$ . Then M is smooth along  $\mathfrak{g}'_X$  at a point  $x \in X$  if there exists a  $\mathfrak{g}'_x$ -submodule  $M^0_x \subseteq M_x$  of finite type over  $\mathcal{O}_x$ , which generates the  $\mathfrak{g}_x$ -module  $M_x$ . Let  $\Omega \subset X$  be an open subset. M is point-wise smooth in  $\Omega$  if it is smooth at all points x in  $\Omega$ , and globally smooth along  $\mathfrak{g}'_X$  in  $\Omega$  if  $M_\Omega$  contains a coherent  $\mathcal{O}_\Omega$ -submodule  $M^0_\Omega$  which generates the  $\mathfrak{g}_\Omega$ -module  $M_\Omega$  and satisfies  $\mathfrak{g}'_\Omega \cdot M^0_\Omega \subseteq M^0_\Omega$ .

It often follows from the context which sub-algebroid  $\mathfrak{g}'_X \subseteq \mathfrak{g}_X$  is intended, and we then abbreviate by simply saying that M is (point-wise) smooth. We then also let  $\operatorname{coh}_s(\mathfrak{g}_X) \subset \operatorname{coh}(\mathfrak{g}_X)$  denote the sub-category of smooth modules.

Obviously, a globally smooth module is point-wise smooth. When X is a quasi-compact variety, so X is a noetherian scheme, the converse also holds (*Prop.* 4.1.7), so there is no distinction between globally smooth and point-wise smooth modules, and we say only smooth. This is not true when X is a quasi-compact complex manifold; see 4.4.

Notice that  $\mathfrak{g}'_x M^0_x \subseteq M^0_x$  implies that M satisfies the following weaker condition, which occasionally is useful.

**Definition 4.1.2.** Let  $\delta$  be a vector in  $\mathfrak{g}_X(U)$ , where U is an open subset of X. We say that M is point-wise smooth along  $\delta$  in  $\mathfrak{g}'_U$  when for any point  $x \in U$  there exists an integer n(=n(x)) such that

$$\delta^n_x M^0_x \subseteq \mathcal{D}^{n-1}_X(\mathfrak{g}'_x) M^0_x$$

where  $M_x^0 \subset M_x$  is a submodule of finite type over  $\mathcal{O}_x$  such that  $\mathcal{D}(\mathfrak{g}_x)M_x^0 = M_x$ .

The first part of the following lemma implies that the above definitions do not depend on the choice of generating  $\mathcal{O}_X$ -module  $M^0 \subseteq M$ .

**Lemma 4.1.3.** Let  $(A, \mathfrak{g}, \alpha)$  be a Lie algebroid over a k-algebra  $A, \mathfrak{b} \subseteq \mathfrak{a} \subseteq \mathfrak{g}$  be sub-algebroids, and  $M \in \operatorname{coh}(\mathfrak{g})$ . Then:

- (1) *M* is smooth along  $\mathfrak{a}$  if and only if every *A*-submodule of finite type  $M^1 \subseteq M$  generates an  $\mathfrak{a}$ -module which remains of finite type over *A*;
- (2) If M is smooth along  $\mathfrak{a}$ , then M is smooth along  $\mathfrak{b}$ .

*Proof.* Trivial.  $\Box$
## Proposition 4.1.4. Let

$$0 \to M_1 \to M \to M_2 \to 0$$

be a short exact sequence of coherent  $\mathfrak{g}_X$ -modules. Then  $M \in \operatorname{coh}_s(\mathfrak{g}_X)$  if and only if  $M_1, M_2 \in \operatorname{coh}_s(\mathfrak{g}_X)$ .

Proof. It is evident that  $M \in \operatorname{coh}_s(\mathfrak{g}_X)$  implies  $M_1, M_2 \in \operatorname{coh}_s(\mathfrak{g}_X)$ . To prove the converse, let x be a point in X and  $M_x^0$  be an  $\mathcal{O}_x$ -submodule of  $M_x$  of finite type, and put  $N = \mathcal{D}(\mathfrak{g}'_x)M_x^0$ . Since  $M_2 \in \operatorname{coh}_s(\mathfrak{g}_X)$  the image  $\overline{N} \subset (M_2)_x$  is of finite type over  $\mathcal{O}_x$ , so N is of finite type if  $N_1 := N \cap (M_1)_x$ is of finite type. We have  $N_1 \subset N$ ; N is of finite type over  $\mathfrak{g}'_x$  and  $\mathcal{D}(\mathfrak{g}'_x)$ is noetherian; hence  $N_1$  is of finite type over  $\mathfrak{g}'_x$ . Since  $N_1 \subset (M_1)_x$  and  $M_1 \in \operatorname{coh}_s(\mathfrak{g}_X)$  it follows that  $N_1$  is of finite type over  $\mathcal{O}_x$  (Lem. 4.1.3).  $\Box$ 

**Proposition 4.1.5.** Let  $\mathfrak{g}'_X$  be a Lie sub-algebroid of  $\mathfrak{g}_X$  and  $M \in \operatorname{coh}(\mathfrak{g}_X)$ .

- (1) (X is a space) The subset  $\Omega = \{x \in X : M_x \text{ is smooth}\}$  is open. If M is smooth at a point x, then there exists a neighbourhood  $\Omega$  of x such that M is globally smooth over  $\Omega$ .
- (2) (X is a quasi-compact space) Let  $S \subset X$  be a divisor whose sheaf of ideals is I, and assume that  $M(*S) \in \operatorname{coh}(\mathcal{O}_X(*S))$ . Then there exists an integer n such that M is point-wise smooth along  $I^n\mathfrak{g}_X$ .

By (1) it follows that if  $M \in \operatorname{coh}_s(\mathfrak{g}_X)$  and  $M_0 \subset M$  is a coherent  $\mathcal{O}_X$ -submodule, then  $\mathcal{D}(\mathfrak{g}'_X)M_0 \in \operatorname{coh}(\mathcal{O}_X)$ . It also implies that M is point-wise smooth on a variety if it is smooth at all closed points.

Remark 4.1.6. We leave out the proofs of the following assertions: (i) Let  $x \in S$  and e(M, x) be the smallest integer n such that  $M_x$  is smooth along  $I_x^{n+1}\mathfrak{g}_x$ . Then for a short exact sequence of  $\mathfrak{g}_X$ -modules

$$0 \to M_1 \to M \to M_2 \to 0$$

one has

$$e(M, x) = \max\{e(M_1, x), e(M_2, x)\}.$$

(ii) Let k be a field of characteristic 0 and R a k-algebra which is a complete discrete valuation ring with residue field  $k = R/\mathfrak{m}$ , and K be the fraction field of R. Let M be an l-dimensional K-vector space which is a module over the Lie algebroid  $T_K := \operatorname{Der}_k(K)$ ;  $T_K$  contains the subalgebroids  $T_R =$  $\operatorname{Der}_k(R)$  and  $\mathfrak{g} := \{\partial \in T_R : \partial(\mathfrak{m}) \subset \mathfrak{m}\}$ . N. Katz [Kat87] has defined certain non-negative rational numbers  $\lambda_1, \lambda_2, \ldots, \lambda_l$  (the slopes of M) and an irregularity index  $\operatorname{Irr}(M) = \sum_{i=1}^n \lambda_i$ ; put  $\lambda_{max} = \max\{\lambda_1, \ldots, \lambda_l\}$ . Define e(M) = e(M, 0) as in (i) with  $\mathfrak{g}_x = \mathfrak{g}$ . Then

$$e(M) = r\lambda_{max}$$

where r is the multiplicity of  $\lambda_{\max}$ . Hence  $\operatorname{Irr}(M) = 0$  if and only if e(M) = 0.

*Proof.* (1): Assume that  $M_x$  is smooth and let  $M_x^0 = \sum_{i=1}^l \mathcal{O}_x \bar{m}_i$  be an  $\mathcal{O}_x$ -module of finite type which generates the  $\mathfrak{g}_x$ -module  $M_x$ , satisfying  $\mathfrak{g}'_x \cdot M_x^0 \subseteq M_x^0$ . The vectors  $\bar{m}_i$  are germs of sections  $m_i$  which are defined in some open set  $\Omega$ ; set  $M_\Omega^0 = \sum \mathcal{O}_\Omega m_i$ , defining a coherent submodule of  $M_\Omega$ . The  $\mathcal{O}_X$ -module  $\mathfrak{g}'_X$  is locally of finite type, so choose sections  $\delta_1, \ldots, \delta_n$  defined in an open subset  $\Omega' \subset \Omega$ ,  $x \in \Omega'$ , such that  $\mathfrak{g}'_{\Omega'} = \sum \mathcal{O}_{\Omega'} \delta_j$ . After further shrinking  $\Omega'$  we can assume that  $\delta_j m_i \subset M^0_{\Omega'}$ , for every i, j; hence  $M^0_{\Omega'}$  is a  $\mathfrak{g}'_{\Omega'}$ -module which is coherent over  $\mathcal{O}_{\Omega'}$  and  $M^0_x = (M^0_{\Omega'})_x$  generates the  $\mathfrak{g}_x$ -module  $M_x$ . Since M is coherent over  $\mathfrak{g}_X$  there exists an open subset  $\Omega_1 \subset \Omega', x \in \Omega_1$ , such that  $(M^0_{\Omega'})_{\Omega_1} = M^0_{\Omega_1}$  generates the  $\mathfrak{g}_{\Omega_1}$ -module  $M_{\Omega_1}$ , and therefore  $M^0_y$  generates  $M_y$  for all  $y \in \Omega_1$ .

(2): By quasi-compactness the assertion follows from (1) if we show that there exists an integer n such that  $M_x$  is smooth along  $I_x^n \mathfrak{g}$  for a point  $x \in X$ . Let  $j : M_x \to M(*S)_x$  be the natural mapping and put  $M_1 = \text{Ker}(j)$ ,  $M_2 = \text{Im}(j)$ . It suffices to prove that  $M_1$  and  $M_2$  are smooth (*Prop.* 4.1.4). Let  $m_1, \ldots, m_r$  be generators of the  $\mathfrak{g}_x$ -module  $M_1$  and  $\delta_1, \cdots, \delta_s$  generators of the  $\mathcal{O}_x$ -module  $\mathfrak{g}_x$ . Now there exists an integer n such that  $I_x^n \delta_j m_i = 0$ , for all i, j, which implies that  $M_1$  is smooth along  $I_x^n \mathfrak{g}_x$ .

Let instead the  $m_i$  denote vectors that generate the  $\mathcal{O}(*T)_x$ -module  $M(*S)_x$ , such that  $M_2 \subseteq M_3 := \sum \mathcal{D}(\mathfrak{g}_x)m_i$ ; by Proposition 4.1.4 it suffices to see that  $M_3$  is smooth. But since  $M_3 \subseteq M(*S)_x$ , and  $M(*S)_x$  is of finite type over  $\mathcal{O}(*T)_x$ , it is evident that there exists an integer n such that  $I_x^n \delta_i m_j \subseteq \sum \mathcal{O}_x m_l \subseteq M_3$ , for all i, j, implying that  $M_3$  is smooth along  $I_x^n \mathfrak{g}_x$ .  $\Box$ 

**Proposition 4.1.7.** (X is a quasi-compact variety) Let M be a coherent  $\mathfrak{g}_X$ -module. If M is point-wise smooth along a sub-algebroid  $\mathfrak{g}'_X \subseteq \mathfrak{g}_X$ , then it is globally smooth along  $\mathfrak{g}'_X$ .

Proof. Since X is quasi-compact and M is coherent over  $\mathfrak{g}_X$  it follows from (1) in Proposition 4.1.5 that there exist a finite number of open sets  $\Omega_i$  such that  $\cup \Omega_i = X$  and coherent  $\mathcal{O}_{\Omega_i}$ -submodules  $M_{\Omega_i}^0 \subseteq M_\Omega$ , satisfying  $\mathfrak{g}'_{\Omega_i} \cdot M_{\Omega_i}^0 \subseteq M_{\Omega_i}^0$  and  $\mathcal{D}(\mathfrak{g}'_{\Omega_i}) \cdot M_{\Omega_i}^0 = M_{\Omega_i}$ . By [BS58, Prop. 2] there exist coherent  $\mathcal{O}_X$ -sub-modules  $M^i \subseteq M$  such that  $M_{\Omega_i}^i = M_{\Omega_i}^0$ . Now put  $M^{00} =$  $\sum \mathcal{D}(\mathfrak{g}'_X) \cdot M^i$ . Since each  $\mathcal{O}_X$ -module  $\mathcal{D}(\mathfrak{g}'_X) \cdot M^i$  is coherent (Lem. 4.1.3) and the sum is finite it follows that  $M^{00} \subseteq M$  is a globally defined coherent  $\mathcal{O}_X$ -submodule such that  $\mathfrak{g}'_X \cdot M^{00} \subseteq M^{00}$  and  $\mathcal{D}(\mathfrak{g}_X) \cdot M^{00} = M$ .  $\square$ 

**Proposition 4.1.8.** Let  $M \in \operatorname{coh}(\mathfrak{g}_X)$ . There exists a maximal Lie subalgebroid  $\mathfrak{g}_X^M \subset \mathfrak{g}_X$  such that M is smooth along  $\mathfrak{g}_X^M$ .

*Proof.* Let  $\mathfrak{g}_X^M \subseteq \mathfrak{g}_X$  be the Lie subalgebroid that is generated by the sum of all sub-algebroids of  $\mathfrak{g}_X$  along which M is smooth; it is coherent since it is locally generated by its sections and  $\mathcal{O}_X$  is noetherian. By Theorem 2.3.4 M is smooth along  $\mathfrak{g}_X^M$ .  $\Box$ 

Let  $M \in \operatorname{coh}(\mathfrak{g}_X)$  and I be the coherent ideal whose zero locus is supp M.

**Proposition 4.1.9.** (X is a space)

 $I\mathfrak{g}_X \subseteq \mathfrak{g}_X^M \subseteq \mathfrak{g}_X(I) \quad (see \ 4.1.8)$ 

*Proof.*  $I\mathfrak{g}_X \subseteq \mathfrak{g}_X^M$ : If  $x \notin Z$  there is nothing to prove so let  $x \in Z$ ,  $\phi \in I_x, \nabla = \phi \delta \in I_x\mathfrak{g}_x$ , and  $m_x \in M_x$ . One proves by induction that for any positive integer n one can rewrite  $\nabla^n \in \mathcal{D}(I_x\mathfrak{g}_x)$  as

$$\nabla^n = \delta^n \phi^n + a_{n-1} \nabla^{n-1} + \dots + a_0, \quad \text{where} \quad a_i \in \mathcal{O}_x.$$

Therefore, if  $\phi^n m_x = 0$ , then  $\nabla^n \cdot m = (a_{n-1}\nabla^{n-1} + \cdots + a_0)m$ , whence the assertion.

 $\mathfrak{g}_X^M \subseteq \mathfrak{g}_X(I)$ : The assertion being local in X we can assume that all sections below are globally defined.

If  $\delta \in \mathfrak{g}_X^M$  it suffices to prove that there exists a Zariski open dense subset  $Z_0 \subseteq Z$  such that  $\delta_x \in \mathfrak{g}_x(I)$  when  $x \in Z_0$ , for then if  $\phi \in I$ ,  $\alpha(\delta)(\phi) \in I(Z_0 \cup (X \setminus Z)) = I(X)$  (Lem. 5.1.1).

Supposing that  $\delta \notin \mathfrak{g}(I)$  we prove that there exists a Zariski open dense subset  $Z_0 \subset Z$  such that  $\delta_x \notin \mathfrak{g}_x^M$  when  $x \in Z_0$ : There exists a function  $\phi \in I$  such that  $\alpha(\delta)(\phi) \notin I$ . Set  $Z_0 = \{x \in Z : \alpha(\delta)(\phi)_x \notin \mathfrak{m}_x\}$ ; this is a Zariski open dense subset of Z since  $\alpha(\delta)(\phi)_x \notin I_x$ . Let  $x \in Z_0$  and let  $M_x^{\phi_x}$ the  $\phi_x$ -invariants of  $M_x$ . Clearly  $M_x^{\phi_x} \neq 0$  so we get a non-zero  $\mathfrak{g}_x$ -module

$$M_1 = \mathcal{D}(\mathfrak{g}_x) M_x^{\phi_x}$$

Let  $\mathfrak{g}_x^1$  be the Lie algebroid that is generated by  $\mathfrak{g}_x^M$  and  $\delta_x$ . By Proposition 4.1.4 it suffices to prove that  $M_1$  is not smooth along the sub-algebroid  $\mathfrak{g}_x^1 \subset \mathfrak{g}_x$ . As  $\mathcal{D}(\mathfrak{g}_x)$  is noetherian it follows that  $M_1 \subset M$  is of finite type, hence there exists a sub-module  $M_1^0 \subseteq M_x^{\phi_x}$  of finite type over  $\mathcal{O}_x$  such that  $\mathcal{D}(\mathfrak{g}_x) \cdot M_1^0 = M_1$ . Clearly it suffices to prove that  $\delta_x^n \cdot M_1^0 \notin \mathcal{D}^{n-1}(\mathfrak{g}_x^1) \cdot M_1^0$ . Since  $\phi_x \delta \cdot M_1^0 = -\alpha(\delta)(\phi_x) \cdot M_1^0 \neq 0$ ,  $(\alpha(\phi_x) \notin \mathfrak{m}_x)$ , so  $\delta \cdot M_1^0 \notin M_1^0$ , this takes care of the n = 1 case. Assume that  $n \geq 2$ , and suppose the contrary that

$$\delta^n \cdot M_1^0 \subseteq \mathcal{D}^{n-1}(\mathfrak{g}_x^1) \cdot M_1^0.$$

Then

$$\phi \cdot \delta^n \cdot M_1^0 \subseteq \phi \cdot \mathcal{D}^{n-1}(\mathfrak{g}^1_x) \cdot M_1^0 \subseteq \mathcal{D}^{n-2}(\mathfrak{g}^1_x) \cdot M_1^0,$$

and

$$\phi \cdot \delta^n \cdot M_1^0 = [\phi, \delta^n] \cdot M_1^0 = \delta \cdot [\phi, \delta^{n-1}] \cdot M_1^0 - \alpha(\delta)(\phi) \cdot \delta^{n-1} \cdot M_1^0;$$

hence

$$\alpha(\delta)(\phi) \cdot \delta^{n-1} \cdot M_1^0 \subseteq \delta \cdot [\phi, \delta^{n-1}] \cdot M_1^0 + \mathcal{D}^{n-2}(\mathfrak{g}_x^1) \cdot M_1^0 = \mathcal{D}^{n-2}(\mathfrak{g}_x^1) \cdot M_1^0.$$

Since  $\alpha(\delta)(\phi)$  is invertible an induction gives a contradiction.  $\Box$ 

4.2. Smooth complexes. Throughout this subsection we let  $\pi : Y \to X$  be a morphism of non-singular spaces, and if  $\mathfrak{g}'_X \subset \mathfrak{g}_X$  is an inclusion of Lie algebroids we let  $\mathfrak{g}'_Y$  be the image of  $\pi^+(\mathfrak{g}'_X)$  in  $\mathfrak{g}_Y := \pi^+(\mathfrak{g}_X)$ .

Let  $D_s^b(\mathfrak{g}_X)$  be the sub-category of bounded complexes  $M^{\bullet}$  whose homology  $H^{\bullet}(M^{\bullet}) \in \operatorname{coh}_s(\mathfrak{g}_X)$ .

Lemma 4.2.1. Let

$$M_1^\bullet \to M^\bullet \to M_2^\bullet \to$$

be a distinguished triangle in  $D^b(\mathfrak{g}_X)$ . If two vertices of this triangle belong to  $D^b_s(\mathfrak{g}_X)$ , then the third vertex also belongs to  $D^b_s(\mathfrak{g}_X)$ .

*Proof.* This follows immediately from Proposition 4.1.4, using the long exact sequence in homology.  $\Box$ 

4.2.2. Inverse images. When  $M^{\bullet} \in D^{b}_{\mathrm{coh}}(\mathfrak{g}_{X})$  then any submodule of finite type of the homology module  $H^{\bullet}(\pi^{!}(M^{\bullet}))$  is coherent. Let us therefore agree that a  $\mathfrak{g}_{Y}$ -module is smooth if all its coherent submodules are smooth.

The category  $\operatorname{coh}_s(\mathfrak{g}_x), x \in X$ , does in general not contain sufficiently many projectives (or flats), so the next result is not quite immediate.

**Proposition 4.2.3.** ( $\mathfrak{g}_X$  is locally free and transitive) If  $M^{\bullet} \in D^b_s(\mathfrak{g}_X)$ , then the homology of  $\pi^!(M^{\bullet})$  is smooth along  $\mathfrak{g}'_Y$ .

*Proof.* We will be brief since standard arguments for  $\mathcal{D}$ -modules are used. It suffices to prove the assertion for a generating class in  $D_s^b(\mathfrak{g}_X)$ , so if M is a smooth  $\mathfrak{g}_X$ -module and we prove that the complex  $\pi^!(M)$  has smooth homology we are done.

Factorize  $\pi$  as  $\pi = p \circ i$  where  $i: Y \to Z = Y \times X$  is the graph morphism and  $p: Z \to X$  is the projection on the second factor. As  $\mathfrak{g}_X$  is locally free and p is a smooth morphism of non-singular spaces we have  $\pi^!(M) \cong i!p!(M)$ (*Prop.* 3.4.3) and  $\mathfrak{g}'_Y = i^+(p^+(\mathfrak{g}'_X))$  (*Lem.* 2.4.6), so it suffices to prove the assertion for i and p separately.

p: Since p is flat and  $\mathfrak{g}_X$  is locally free, p'(M) is the single degree complex

$$\mathcal{D}_{Y \to X} \otimes_{p^{-1}(\mathcal{D}(\mathfrak{g}_X))} p^{-1}(M) = \mathcal{O}_Y \otimes_{p^{-1}(\mathcal{O}_X)} p^{-1}(M);$$

see the proof of Proposition 3.4.3. This module clearly is smooth along  $\mathfrak{g}'_Z = p^+(\mathfrak{g}'_X)$  (see (3.4)).

*i*: Letting  $N \in \operatorname{coh}_s(\mathfrak{g}_Z)$  we have to prove that  $i^!(N) \in D_s^b(\mathfrak{g}_Y)$ . First note that  $\mathfrak{g}_Z = p^!(\mathfrak{g}_X)$  is locally free since p is submersive and  $\mathfrak{g}_X$  is locally free. Now the closed embedding i can be factorized into a sequence of closed embeddings of the type  $\pi : Y \to Z$  where  $\operatorname{codim}_Z \pi(Y) = 1$ , and the pullback of a locally free transitive Lie algebroid on a non-singular space to a non-singular subspace again is locally free and transitive (*Lem.* 2.4.8), so by Proposition 3.4.3 we may assume that  $\operatorname{codim}_Z i(Y) = 1$ . Let  $I_Y$  be the ideal of Y. Then since  $\mathfrak{g}_Z = p^!(\mathfrak{g}_X)$  is locally free and Y is non-singular one can use the Koszul complex to compute  $i^!(M)$ ; see above Proposition 3.4.3. The only non-zero cohomology  $\mathfrak{g}_Y$ -modules are

$$M/I_Y M$$
 and  $M^{I_Y}$ .

That any coherent  $\mathcal{O}_Y$ -submodule of these modules generates a  $\mathfrak{g}'_Y$ -module  $(i_*(\mathfrak{g}'_Y) = \mathfrak{g}'_X(I_Y)/I_Y\mathfrak{g}'_X)$  which remains coherent over  $\mathcal{O}_Y$  follows immediately since any coherent  $\mathcal{O}_X$ -submodule of M generates a  $\mathfrak{g}'_X$ -module that remains coherent over  $\mathcal{O}_X$ .  $\Box$ 

We will give a partial converse of Proposition 4.2.3. Let  $y \in Y$  and  $x = \pi(y) \in X$ . A morphism of Lie algebroids  $\pi' : \mathfrak{g}_Y \to \mathfrak{g}_X$  is submersive at y if the induced morphism  $\mathfrak{g}_Y \to k_y \otimes_{\mathcal{O}_y} \pi^*(\mathfrak{g})_y$  is surjective; this is, by Nakayama's lemma, the same as to say that the morphism  $\mathfrak{g}_y \to \pi^*(\mathfrak{g})_y$  is surjective.

**Proposition 4.2.4.** ( $\mathfrak{g}_X$  is locally free) Let  $M^{\bullet} \in D^b_{\mathrm{coh}}(\mathfrak{g}_X)$ . Assume that  $\pi$  is flat on  $\mathrm{supp} M^{\bullet} \subset X$ , and that for each point  $x \in \mathrm{supp} M^{\bullet}$  there exists a point  $y \in \pi^{-1}(x)$  at which  $\pi' : \mathfrak{g}'_Y \to \mathfrak{g}'_X$  is submersive. Then the following are equivalent:

(1)  $M^{\bullet} \in D^b_s(\mathfrak{g}_X);$ 

(2)  $\pi^!(M^{\bullet}) \in D^b_s(\mathfrak{g}_Y).$ 

*Remark* 4.2.5. When considering completely regular complexes one can improve Proposition 4.2.4, not demanding that  $\pi$  be flat (*Prop.* 6.4.8).

*Proof.* (1)  $\Rightarrow$  (2): Apply Proposition 4.2.3. (2)  $\Rightarrow$  (1): It suffices to prove this for a generating class of  $D^b(\mathfrak{g}_X)$ , so letting  $M \in \operatorname{coh}(\mathfrak{g}_X)$  satisfy  $\pi^!(M) \in D^b_s(\mathfrak{g}_Y)$ , we need to prove that  $M \in \operatorname{coh}_s(\mathfrak{g}_X)$ .

Let  $x \in \operatorname{supp} M$  and put  $A = \mathcal{O}_x$ . If  $M_x^0 \subset M_x$  an A-submodule of finite type, we have to check that  $\mathcal{D}(\mathfrak{g}'_x)M_x^0$  remains of finite type over A. There exists a point  $y \in Y$  such that  $x = \pi(y)$  and the canonical mapping  $\mathfrak{g}'_y \to B \otimes_A \mathfrak{g}'_x$  is surjective, where we have put  $B = \mathcal{O}_y$ . Then

$$\pi^{!}(M)_{y} \cong \pi^{*}(\mathcal{D}_{x}) \otimes^{L}_{\mathcal{D}(\mathfrak{g}_{x})} M_{x}$$
  
$$\cong B \otimes^{L}_{A} M_{x} \quad (\mathfrak{g}_{x} \text{ is free over } \mathcal{O}_{x})$$
  
$$\cong B \otimes_{A} M_{x} \qquad (A \to B \text{ is flat}).$$

As the restriction  $\mathfrak{g}'_y \to \mathfrak{g}'_x$  is surjective, so is its extension to a mapping  $\mathcal{D}(\mathfrak{g}'_y) \to B \otimes_A \mathcal{D}(\mathfrak{g}'_x)$ ; hence, by flatness,

$$B\otimes_A \mathcal{D}(\mathfrak{g}'_x)M^0_x \subset \mathcal{D}(\mathfrak{g}'_y)B\otimes_A M^0_x.$$

By (2) the right side is of finite type over B, and since B is noetherian the left side is also of finite type. Then by faithful flatness  $\mathcal{D}(\mathfrak{g}'_x)M^0_x$  is of finite type over A.  $\Box$ 

4.2.6. Direct images. As in Proposition 3.5.2 we consider the category  $D_{sl}^b(\mathfrak{g}_Y)$  of complexes  $M^{\bullet}$  whose homology  $H^{\bullet}(M^{\bullet}) \in \operatorname{coh}_s(\mathfrak{g}_Y)$  locally in X is generated by an  $\mathcal{O}_Y$ -coherent submodule. When X is a quasi-compact variety we have  $D_{sl}^b(\mathfrak{g}_Y) = D_s^b(\mathfrak{g}_Y)$ .

Put  $T_{Y/X} = \operatorname{Ker}(T_Y \to \pi^*(T_X))$  and  $\mathfrak{g}_{Y/X} = \alpha^{-1}(T_{Y/X})$ . For simplicity we shall assume that  $\mathfrak{b}_Y := \operatorname{Ker}(\mathfrak{g}_Y \to T_Y) \subset \mathfrak{g}'_Y$ .

**Theorem 4.2.7.**  $(\mathfrak{g}_X \text{ is locally free and transitive})$  Let  $\mathfrak{g}'_Y \subset \mathfrak{g}_Y$  and  $\mathfrak{g}'_X \subset \mathfrak{g}_X$  be sub-algebroids and  $\mathfrak{a}_Y$  be the Lie algebroid that is generated by  $\mathfrak{g}'_Y + \mathfrak{g}_{Y/X} \subseteq \mathfrak{g}_Y$ . Assume that  $\pi$  is proper and that

(4.2.1) 
$$\operatorname{Im}(\pi^*(\mathfrak{g}'_X) \to \pi^*(\mathfrak{g}_X)) \subseteq \operatorname{Im}(\mathfrak{a}_Y \to \pi^*(\mathfrak{g}_X))$$

(canonical mappings). Then  $\pi_+$  defines a functor

$$D^b_{sl}(\mathfrak{g}_Y) \to D^b_s(\mathfrak{g}_X).$$

The condition (4.2.1) means that, locally in Y and X, sections of  $\mathfrak{g}'_X$  should be liftable to sections of  $\mathfrak{a}_Y$ .

*Proof.* Letting  $M^{\bullet} \in D^b_s(\mathfrak{g}_Y)$  we have to prove that

$$R\pi_*(\mathcal{D}_{X\leftarrow Y}\otimes^L_{\mathcal{D}(\mathfrak{g}_Y)}M^\bullet)\in D^b_s(\mathfrak{g}_X).$$

a) Let  $\mathcal{D}_{X\leftarrow Y}^n$  be the right  $\mathcal{O}_Y$ -module that is generated by the image of  $\pi^{-1}(\mathcal{D}^n(\mathfrak{g}_X))$  in the  $(\pi^{-1}(\mathfrak{g}_X),\mathfrak{g}_Y)$ -bimodule  $\mathcal{D}_{X\leftarrow Y}$ . As  $\mathcal{D}_{X\leftarrow Y}^n\mathfrak{g}_{Y/X} \subseteq \mathcal{D}_{X\leftarrow Y}^n\mathfrak{b}_Y$  and  $\mathfrak{b}_Y \subseteq \mathfrak{g}'_Y$ , (4.2.1) implies

(4.2.2) 
$$\pi^{-1}(\mathfrak{g}'_X)\mathcal{D}^n_{X\leftarrow Y}\subseteq \mathcal{D}^n_{X\leftarrow Y}\mathfrak{a}_Y\subseteq \mathcal{D}^n_{X\leftarrow Y}\mathfrak{g}'_Y$$

b) It suffices to prove the assertion for a generating class for the considered category, and since the assertion is local in X, it suffices to prove that  $\pi_+(M) \in D^b_s(\mathfrak{g}_X)$  when  $M \in \operatorname{coh}_s(\mathfrak{g}_Y)$  is generated by a coherent  $\mathcal{O}_Y$ -submodule  $M^0$ .

The morphism  $\pi$  can be factorized  $\pi = p \circ i$  where  $p: Z = Y \times X \to X$ is the projection on the second factor and  $i: Y \to Z$  is the graph morphism, which is a closed embedding. We contend that it suffices to prove the theorem for i and p separately. Since  $\pi^! = i^! p^!$  (Prop. 3.4.3) we then have to check that the conditions in the theorem are satisfied for i and p: Put  $\mathfrak{g}'_Z = p^+(\mathfrak{g}'_X) = p^*(\mathfrak{g}'_X)$  and  $\mathfrak{g}_Z = p^+(\mathfrak{g}_X) = p^*(\mathfrak{g}_X)$ . Since pis a projection, by Lemma 2.4.6  $\pi^+(\mathfrak{g}'_X) = i^+(\mathfrak{g}'_Z), \mathfrak{g}_Y = i^+(\mathfrak{g}_Z),$  and  $\mathfrak{g}_{Y/X} = \mathfrak{g}_{Y/Z} + i^+(\mathfrak{g}_{Z/X})$ . Let  $\mathfrak{a}_Z$  be the Lie algebroid that is generated by  $\mathfrak{g}'_Z + \mathfrak{g}_{Z/X}$  and  $\mathfrak{a}^1_Y$  the Lie algebroid that is generated by  $\pi_1^+(\mathfrak{g}'_Z) + \mathfrak{g}_{Y/Z};$ now since  $\mathfrak{g}_{Z/X} \subset \mathfrak{g}'_Z$  and  $\mathfrak{g}_{Y/Z} \subset \pi^+(\mathfrak{g}'_X)$  we actually have  $\mathfrak{a}_Z = \mathfrak{g}'_Z$  and  $\mathfrak{a}^1_Y = \pi^+(\mathfrak{g}'_X) \subset \mathfrak{a}_Y$ . It should then be clear that (4.2.1) implies

$$\operatorname{Im}(i^*(\mathfrak{g}'_Z) \to i^*(\mathfrak{g}_Z)) = \operatorname{Im}(\mathfrak{a}^1_Y \to i^*(\mathfrak{g}_Z)) \subseteq \operatorname{Im}(\mathfrak{a}_Y \to i^*(\mathfrak{g}_Z))$$

and

$$\operatorname{Im}(p^*(\mathfrak{g}'_X) \to p^*(\mathfrak{g}_X)) = \operatorname{Im}(\mathfrak{a}_Z \to p^*(\mathfrak{g}_X)).$$

so (4.2.1), and hence (4.2.2) holds for i and p separately. Moreover, by Lemma 2.4.8 the pull-back of a locally free and transitive Lie algebroid again is locally free and transitive. In the complex analytic case we also remark that if M is generated by a coherent  $\mathcal{O}_Y$ -module, then  $H^{\bullet}(i_+(M^{\bullet}))$ is generated by a coherent  $\mathcal{O}_Z$ -module, i.e.  $i_+(D^b_{sl}(\mathfrak{g}_Y)) \subset D^b_{sl}(\mathfrak{g}_Z)$ . This proves that the conditions are satisfied for i and p separately.

 $i_+(D_s^b(\mathfrak{g}_Y)) \subset D_s^b(\mathfrak{g}_Z)$ : First, since  $\mathfrak{g}_Z$  is locally free and transitive one can prove as in [BGK<sup>+</sup>87, VI, 7.8], see also c) in the proof of Theorem 3.5.4, that the right  $\mathcal{D}(\mathfrak{g}_Y)$ -module  $\mathcal{D}_{Z\leftarrow Y}$  is locally free. Therefore, since  $i_*$  is exact one has

$$i_+(M) = i_*(\mathcal{D}_{Z\leftarrow Y} \otimes_{\mathcal{D}(\mathfrak{g}_Y)} M).$$

For a coherent  $\mathcal{O}_Y$ -submodule  $M^0 \subset M$  we let

$$i_*(\mathcal{D}^n_{Z\leftarrow Y}\cdot M^0)\subset i_*(\mathcal{D}_{Z\leftarrow Y}\otimes_{\mathcal{D}(\mathfrak{g}_Y)}M)$$

be the image of the canonical morphism  $i_*(\mathcal{D}_{Z\leftarrow Y}\otimes_{\mathcal{O}_Y} M^0) \to i_*(\mathcal{D}_{Z\leftarrow Y}\otimes_{\mathcal{D}(\mathfrak{g}_Y)} M)$ ; this is a coherent  $\mathcal{O}_Z$ -module. If  $N \subset i_*(\mathcal{D}_{Z\leftarrow Y}\otimes_{\mathcal{D}(\mathfrak{g}_Y)} M)$  is a coherent  $\mathcal{O}_Z$ -submodule there exists such a coherent submodule  $M^0$  and an integer n such that

$$N \subseteq i_*(\mathcal{D}^n_{Z \leftarrow Y} \cdot M^0).$$

Therefore by (4.2.2)

$$\mathcal{D}(\mathfrak{g}'_Z)N \subseteq \mathcal{D}(\mathfrak{g}'_Z)i_*(\mathcal{D}^n_{Z\leftarrow Y}\cdot M^0) \subseteq i_*(\mathcal{D}^n_{Z\leftarrow Y}\cdot \mathcal{D}(\mathfrak{g}'_Y)M^0).$$

Since  $\mathcal{D}(\mathfrak{g}'_Y)M^0 \in \operatorname{coh}(\mathcal{O}_Y)$  and  $\mathcal{D}^n_{Z \leftarrow Y}$  is coherent both as right  $\mathcal{O}_Y$ -module and left  $\mathcal{O}_Z$ -module the right side is coherent; the left side is locally generated by its sections and  $\mathcal{O}_Z$  is noetherian, hence  $\mathcal{D}(\mathfrak{g}'_Z)N \in \operatorname{coh}(\mathcal{O}_Z)$ .

It remains to prove: If  $M \in D^b_{sl}(\mathfrak{g}_Z)$  and the restriction of  $p: Z \to X$  to supp M is proper, then  $p_+(M) \in D^b_s(\mathfrak{g}_X)$ : Here  $\mathfrak{g}_Z = p^*(\mathfrak{g}_X) \oplus q^*(T_Z)$ , where q is the projection on Y, so  $p^*(\mathfrak{g}_X)$  is a Lie algebroid on Z in a natural way. The  $\mathcal{O}_Z$ -module  $p^*(\mathcal{D}(\mathfrak{g}_X))$  is a right  $p^{-1}(\mathfrak{g}_X)$ -module and left  $\mathfrak{g}_Y$ -module, but since the action of  $p^{-1}(\mathfrak{g}_X)$  commutes with that of  $q^{-1}(\mathcal{O}_Y)$  it follows that  $p^*(\mathcal{D}(\mathfrak{g}_X))$  is a right module over the Lie algebroid  $p^*(\mathfrak{g}_X)$ . Changing right to left we have a  $(p^*(\mathfrak{g}_X), \mathfrak{g}_Z)$ -bimodule

$$\mathcal{D}_{X\leftarrow Z} = \omega_Z \otimes_{\mathcal{O}_Z} p^*(\omega_X^{-1}) \otimes_{\mathcal{O}_Z} p^*(\mathcal{D}_X)$$

which is coherent over  $p^*(\mathfrak{g}_X)$ ; hence, since  $M \in \operatorname{coh}(\mathfrak{g}_Z)$ , one has

$$\mathcal{D}_{X\leftarrow Z}\otimes^{L}_{\mathcal{D}(\mathfrak{g}_Z)} M \in D^{b}_{\mathrm{coh}}(p^*(\mathfrak{g}_X)).$$

Now  $\Omega_{Z/X}^{\bullet} \otimes_{\mathcal{O}_Z} \mathcal{D}(\mathfrak{g}_Z) = q^*(\Omega_Y^{\bullet}) \otimes_{\mathcal{O}_Z} \mathcal{D}(\mathfrak{g}_Z)$ , where  $\Omega_Y^{\bullet}$  is the de Rham complex on Y, is a left resolution of the bimodule  $\mathcal{D}_{X \leftarrow Z}$  which is free as right  $\mathfrak{g}_Z$ -module (see [BGK<sup>+</sup>87, VI, 5.3]), so

$$\mathcal{D}_{X\leftarrow Z}\otimes^L_{\mathcal{D}(\mathfrak{g}_Z)}M=\Omega^{\bullet}_{Z/X}\otimes_{\mathcal{O}_Z}M.$$

Then (4.2.2) implies that  $\mathcal{D}_{X\leftarrow Z} \otimes_{\mathcal{D}(\mathfrak{g}_Z)}^L M \in D^b_s(p^*(\mathfrak{g}_X))$  where the index s denotes smoothness along  $p^*(\mathfrak{g}'_X) \subset p^*(\mathfrak{g}_X)$ . Since  $D^b_s(p^*(\mathfrak{g}_X))$  is generated by its homology objects it now suffices to prove the following: If  $N \in \operatorname{coh}_s(p^*(\mathfrak{g}_X))$ , then

$$Rp_*(N) \in D^b_s(\mathfrak{g}_X).$$

As the restriction of p to supp M is proper the proof in Proposition 3.5.2 implies that each homology module  $R^i p_*(N) \in \operatorname{coh}(\mathfrak{g}_X)$ . Now the problem is local on X so it remains to see that if  $L_i \subset R^i p_*(N)$  is an  $\mathcal{O}_X$ -coherent submodule then  $\mathcal{D}(\mathfrak{g}'_X)L_i \in \operatorname{coh}(\mathcal{O}_X)$ . To see this first note that N contains a  $p^*(\mathfrak{g}'_X)$  module  $N_i$  which is coherent over  $\mathcal{O}_Z$ , and such that

$$L_i \subset K_i := \operatorname{Im} \{ R^i p_*(N_i) \to R^i p_*(N) \}.$$

Then since  $K_i \in \operatorname{coh}(\mathcal{O}_X)$  is a  $\mathfrak{g}'_X$ -module the assertion follows.  $\Box$ 

Remark 4.2.8. Is Theorem 4.2.7 true for all locally free Lie algebroids? The problem occurs when  $\pi: Y \to X$  is a closed embedding and  $\mathcal{D}_{X \leftarrow Y}$  is not free as  $\mathfrak{g}_Y$ -module. We need a resolution  $F^{\bullet} \to \mathcal{D}_{X \leftarrow Y}$  of the  $(\pi^{-1}(\mathfrak{g}_X), \mathfrak{g}_Y)$ -bimodule  $\mathcal{D}_{X \leftarrow Y}$  such that each term  $F^i$  is a  $(\pi^{-1}(\mathfrak{g}_X), \mathfrak{g}_Y)$ -bimodule which is flat as  $\mathfrak{g}_Y$ -module and provided with a filtration by  $\pi^{-1}(\mathcal{O}_X)$ -coherent submodules  $F_n^i$  such that  $\cup_{n\geq 0} F_n^i = F^i$  and  $\pi^{-1}(\mathfrak{g}'_X)F_n^i \subset F_n^i\mathfrak{g}'_Y$ .

4.3. Completion. Let  $(A, \mathfrak{m}, k)$  be a noetherian local k-algebra and  $\mathfrak{g}' \subseteq \mathfrak{g}$ Lie algebroids over A. Let  $\hat{M}$  be the  $\mathfrak{m}$ -adic completion of an A-module M. The Artin-Rees lemma implies that if M is of finite type, then  $\hat{M} = \hat{A} \otimes_A M$ and  $M \subseteq \hat{M}$ , in particular  $\hat{\mathfrak{g}}' = \hat{A} \otimes_A \mathfrak{g}'$ ,  $\hat{\mathfrak{g}} = \hat{A} \otimes_A \mathfrak{g}$ . Since  $A \subseteq \hat{A}$ and derivations of A are continuous in the  $\mathfrak{m}$ -adic topology we have that  $\operatorname{Der}_k(A) \subseteq \operatorname{Der}_k(\hat{A})$ . More generally, if M is a  $\mathfrak{g}$ -module of finite type, its defining homomorphism  $\rho : \mathfrak{g} \to \mathfrak{c}(M)$  lifts to homomorphisms  $\hat{\rho} : \hat{\mathfrak{g}} \to \mathfrak{c}(\hat{M})$ and  $\tilde{\rho} : \hat{\mathfrak{g}} \to \hat{A} \otimes_A \mathfrak{c}(M)$ . Note that if M is not of finite type over A, then the natural injective mapping  $\hat{A} \otimes M \to \hat{M}$  is not surjective, so the  $\hat{\mathfrak{g}}$ -module of finite type  $\tilde{M} := \hat{A} \otimes_A M$  is not  $\mathfrak{m}$ -adically complete, and  $\hat{M}$  is not of finite type over  $\hat{\mathfrak{g}}$ . For instance,  $\mathcal{D}(\hat{\mathfrak{g}}) \cong \hat{A} \otimes_A \mathcal{D}(\mathfrak{g}) \ncong \hat{\mathcal{D}}(\mathfrak{g})$ . Still, the following result is very useful for checking if M is smooth because of the big supply of invariants in  $\tilde{M}$  (for an application, see [Käl98, Sec. 4]). **Proposition 4.3.1.** The following are equivalent:

(1) M is smooth along  $\mathfrak{g}'$ .

(2) M is smooth along  $\hat{\mathfrak{g}}'$ .

*Proof.* This is similar to the proof of Proposition 4.2.4. If  $M^0$  is a  $\mathfrak{g}'$ -module of finite type over A, then  $\hat{M}^0 \cong \hat{A} \otimes_A M^0$  is a  $\hat{\mathfrak{g}}' = \hat{A} \otimes_A \mathfrak{g}'$ -module of finite type over  $\hat{A}$ . It is also clear that  $\mathcal{D}(\hat{\mathfrak{g}}) \cdot \hat{M}^0 = \hat{A} \otimes_A M = \tilde{M}$ . This proves that (1) implies (2).

Assume (2). Let  $M^0$  be an A-submodule of finite type that generates Mover  $\mathfrak{g}$ , and set  $M_1 = \mathcal{D}(\mathfrak{g}')M^0$ . We have  $\hat{M}^0 \cong \hat{A} \otimes_A M^0$ , and since  $\mathcal{D}(\hat{\mathfrak{g}}') = \hat{A} \otimes_A \mathcal{D}(\mathfrak{g}')$  it follows that  $\tilde{M}_1 = \hat{A} \otimes_A M^1 = \hat{A} \otimes_A \mathcal{D}(\mathfrak{g}')M^0 = \mathcal{D}(\hat{\mathfrak{g}}')\hat{M}^0$ . (2) and Lemma 4.1.3 implies that  $\tilde{M}_1$  is of finite type over  $\hat{A}$ . Hence there exists a surjection  $\hat{f} : \bigoplus^n \hat{A} \to \hat{A} \otimes M_1$  for some integer n where  $\hat{f} = \mathrm{id} \otimes f$ for some homomorphism  $f : \bigoplus^n A \to M_1$ . Now since the functor  $\hat{A} \otimes_A \cdot \mathrm{is}$ faithfully flat it follows that f is surjective. This implies (1).  $\Box$ 

Remark 4.3.2. Analogous results hold for holomorphic localisation. Let  $\mathcal{O}_x$  be the local ring at a closed point x of a complex variety and  $\mathcal{O}_x^{an}$  be the stalk of holomorphic functions at x. Then  $\mathcal{O}_x \to \mathcal{O}_x^{an}$  is a faithfully flat ring extension, so  $M_x$  is smooth if and only if  $M_x^{an} := \mathcal{O}_x^{an} \otimes_{\mathcal{O}_x} M_x$  is smooth along  $\mathcal{O}_x^{an} \otimes_{\mathcal{O}_x} \mathfrak{g}'_x$ .

4.4. **GAGA.** Let X be a projective complex variety,  $X_h$  the associated compact complex analytic space, and

$$\pi: (\mathcal{O}_{X_h}, X_h) \to (\mathcal{O}_X, X)$$

the canonical continuous homomorphism of ringed spaces. If  $\mathfrak{g}_{X_h} \in \operatorname{Lie}_{X_h}$ we let  $\operatorname{coh}^l(\mathfrak{g}_{X_h}) \subset \operatorname{coh}(\mathfrak{g}_{X_h})$  be the sub-category of coherent  $\mathfrak{g}_{X_h}$ -modules  $M^h$  which contain a coherent  $\mathcal{O}_{X_h}$ -module  $M_0^h$  such that  $\mathcal{D}(\mathfrak{g}_{X_h}) \cdot M_0^h = M^h$  $(M_0^h$  is a lattice in  $M^h$ ). The following implication from GAGA [Ser56] is well-known in the case of  $T_X$ -modules.

**Theorem 4.4.1.** (X is a projective complex variety) The functor

 $\pi^* : \operatorname{Mod}(\mathcal{O}_X) \to \operatorname{Mod}(\mathcal{O}_{X_h}), \quad M \mapsto \mathcal{O}_{X_h} \otimes_{\pi^{-1}(\mathcal{O}_X)} \pi^{-1}(M)$ 

induces equivalences of categories

$$\operatorname{Lie}_X \cong \operatorname{Lie}_{X_h}, and$$
  
 $\operatorname{coh}(\mathfrak{g}_X) \cong \operatorname{coh}^l(\pi^*(\mathfrak{g}_X)).$ 

*Proof.*  $\pi^*$ : Lie<sub>X</sub>  $\to$  Lie<sub>X<sub>h</sub></sub> is fully faithful: Let  $\mathfrak{g}_X^1, \mathfrak{g}_X^2 \in$  Lie<sub>X</sub>. Then clearly  $\pi^*(\mathfrak{g}_X^1), \pi^*(\mathfrak{g}_X^2) \in$  Lie<sub>X<sub>h</sub></sub> and if  $\phi : \mathfrak{g}_X^1 \to \mathfrak{g}_X^2$  is a homomorphism the mapping  $\pi^*(\phi) : \pi^*(\mathfrak{g}_X^1) \to \pi^*(\mathfrak{g}_X^2)$  is a homomorphism of Lie algebroids on  $X_h$ . We have a canonical mapping

$$\operatorname{Hom}_{\operatorname{Lie}_{X}}(\mathfrak{g}_{X}^{1},\mathfrak{g}_{X}^{2}) \xrightarrow{\pi^{*}} \operatorname{Hom}_{\operatorname{Lie}_{X_{h}}}(\pi^{*}(\mathfrak{g}_{X}^{1}),\pi^{*}(\mathfrak{g}_{X}^{2}))$$
$$\subseteq \operatorname{Hom}_{\mathcal{O}_{X_{h}}}(\pi^{*}(\mathfrak{g}_{X}^{1}),\pi^{*}(\mathfrak{g}_{X}^{2})) \cong \operatorname{Hom}_{\mathcal{O}_{X}}(\mathfrak{g}_{X}^{1},\mathfrak{g}_{X}^{2}).$$

Here GAGA implies the last isomorphism and that  $\pi^*$  is injective. Then if  $\phi_h = \mathrm{id} \otimes \phi : \pi^*(\mathfrak{g}^1_X) \to \pi^*(\mathfrak{g}^2_X)$  is a Lie homomorphism it is obvious that  $\phi : \mathfrak{g}^1_X \to \mathfrak{g}^2_X$  also is a Lie homomorphism, so  $\pi^*$  is surjective.

 $\pi^*: \operatorname{Lie}_X \to \operatorname{Lie}_{X_h}$  is essentially surjective: By GAGA if  $\mathfrak{g}_{X_h} \in \operatorname{Lie}_{X_h} \subset \operatorname{coh}(\mathcal{O}_{X_h})$  then there exists a unique  $\mathcal{O}_X$ -module  $\mathfrak{g}_X$  such that  $\mathfrak{g}_{X_h} = \pi^*(\mathfrak{g}_X)$ and  $\pi^{-1}(\mathfrak{g}_X) \subset \mathfrak{g}_{X_h}$ . We need to prove that  $[\pi^{-1}(\mathfrak{g}_X), \pi^{-1}(\mathfrak{g}_X)] \subseteq \pi^{-1}(\mathfrak{g}_X)$ . The mapping  $\operatorname{ad}_h: \mathfrak{g}_{X_h} \to \mathfrak{c}_{X_h}(\mathfrak{g}_{X_h})$  is a 1st order differential operator on the  $\mathcal{O}_{X_h}$ -module  $\mathfrak{g}_{X_h}$  with values in  $\mathfrak{c}_{X_h}(\mathfrak{g}_{X_h})$ , hence it can be identified with an  $\mathcal{O}_{X_h}$ -linear homomorphism  $\operatorname{ad}_h: P_{X_h}^1(\mathfrak{g}_{X_h}) \to \mathfrak{c}_{X_h}(\mathfrak{g}_{X_h})$  (Sec. 2.2), hence by GAGA:  $P_{X_h}^1(\mathfrak{g}_{X_h}) = P_{X_h}^1(\pi^*(\mathfrak{g}_X)) = \pi^*(P_X^1(\mathfrak{g}_X))$ , and

$$\mathfrak{c}_{X_h}(\mathfrak{g}_{X_h}) \subset Hom_{\mathcal{O}_{X_h}}(P^1_{X_h}(\pi^*(\mathfrak{g}_X)), \pi^*(\mathfrak{g}_X)) = \pi^*(Hom_{\mathcal{O}_X}(P^1_X(\mathfrak{g}_X), \mathfrak{g}_X)),$$

implying that  $\mathfrak{c}_{X_h}(\mathfrak{g}_{X_h}) = \pi^*(\mathfrak{c}_X(\mathfrak{g}_X))$ ; and  $\mathrm{ad}_h$  is induced by a unique homomorphism  $\mathrm{ad} : P_X^1(\mathfrak{g}_X) \to \mathfrak{c}_X(\mathfrak{g}_X)$ ; therefore  $[\pi^{-1}(\mathfrak{g}_X), \pi^{-1}(\mathfrak{g}_X)] = \pi^{-1}(\mathrm{ad})(\pi^{-1}(\mathfrak{g}_X))(\pi^{-1}(\mathfrak{g}_X)) \subset \pi^{-1}(\mathfrak{g}_X).$ 

 $\pi^* : \operatorname{coh}(\mathfrak{g}_X) \to \operatorname{coh}^l(\pi^*(\mathfrak{g}_X))$  is an equivalence of categories: Let  $M^h$ ,  $N^h \in \operatorname{coh}^l(\mathfrak{g}_{X_h}), \ \phi^h \in \operatorname{Hom}_{\operatorname{Lie}_{X_h}}(M, N)$ , and  $M_0^h \subset M^h, \ N_0^h \subset N^h$  be coherent  $\mathcal{O}_{X_h}$ -submodules such that

$$\mathcal{D}_{X_h} \cdot M_0^h = M^h$$
, and  $\mathcal{D}_{X_h} \cdot N_0^h = N^h$ .

Since  $X_h$  is compact there exists an integer l so that  $\phi^h(M_0^h) \subset \mathcal{D}_{X_h}^l N_0^h$ ; let  $\phi_0^h$  denote the restriction of  $\phi^h$  to  $M_0^h$ . By GAGA there exist unique  $\mathcal{O}_X$ -modules  $M_0, N_0^l$  and a homomorphism of  $\mathcal{O}_X$ -modules  $\phi_0 : M_0 \to N_0^l$ such that  $\phi^h = \pi^*(\phi_0) : M_0^h = \pi^*(M_0) \to \mathcal{D}_{X_h} \cdot N_0^h = \pi^*(N_0^l)$ . Since  $\mathcal{D}_{X_h} = \mathcal{D}_{X_h}(\pi^*(\mathfrak{g}_X)) = \pi^*(\mathcal{D}_X)$  it follows that there exist coherent  $\mathfrak{g}_X$ modules M, N such that  $M_0 \subset M, N_0 \subset N, \pi^*(M) = M^h, \pi^*(N) = N^h$ , and a homomorphism  $\phi : M \to N$ , extending  $\phi_0$  and satisfying  $\pi^*(\phi) = \phi^h$ ; since by GAGA  $\pi^*(\phi_0) = 0$  implies  $\phi_0 = 0$ , this proves that  $\pi^*$  is essentially surjective and fully faithful.  $\Box$ 

Let us express the parallel for Lie algebroids the fact that on a projective complex variety  $H^1(X, \mathcal{O}_X^*) = H^1(X_h, \mathcal{O}_{X_h}^*)$  because  $\operatorname{Pic}_X = \operatorname{Pic}_{X_h}$ :

**Corollary 4.4.2.** Let X be a projective complex variety. Then there is an isomorphism of complex vector spaces

$$\pi^{-1}: H^2(X, \Omega_X^{\geq 1}) \cong H^1(X_h, \Omega_{X_h}^{1,cl}).$$

The proof is immediate from Theorem 4.4.1; see also 2.5.

Letting  $X_h$  be the holomorphic localisation of a projective algebraic manifold X we now say a few words about non-algebraic  $\mathfrak{g}_{X_h}$ -modules. Let  $S \subset X_h$  be a hyper-surface, put  $X_h^0 = X \setminus S$ , and let M be a coherent  $\mathfrak{g}_{X_h}$ -module such that  $M_{X_h^0} \in \operatorname{coh}^l(\mathfrak{g}_{X_h^0})$ . It is in general a difficult problem to see when  $M \in \operatorname{coh}^l(\mathfrak{g}_{X_h})$ , but we note the following rather trivial fact:

**Lemma 4.4.3.** Let  $S \subset X_h$  be a discrete subset,  $X_h^0 = X_h \setminus S$ ,  $M \in \operatorname{coh}(\mathfrak{g}_{X_h})$ , and  $M_{X_h^0} \in \operatorname{coh}(\mathcal{O}_{X_h^0})$ . Then  $M \in \operatorname{coh}^l(\mathfrak{g}_{X_h})$ .

*Proof.* Let  $x \in S$ . Since  $M \in \operatorname{coh}(\mathfrak{g}_{X_h})$  and S is discrete there exists a neighbourhood  $\Omega$  of x such that  $\Omega \cap S = \{x\}$  and an  $\mathcal{O}_{\Omega}$ -coherent submodule  $M^1_{\Omega} \subset M_{\Omega}$ , generating  $M_{\Omega}$  over  $\mathcal{D}(\mathfrak{g}_{\Omega})$ . As  $M_{\Omega \cap X^0} \in \operatorname{coh}(\mathcal{O}_{X^0 \cap \Omega})$ and the ring  $\mathcal{D}(\mathfrak{g}_{\Omega})$  is noetherian 2.3.2, there exists an integer k such that  $(\mathcal{D}^k(\mathfrak{g}_{X_h})_{\Omega}M^1)_{\Omega \cap X_h^0} = M_{\Omega \cap X_h^0}$ . Now define an  $\mathcal{O}_{X_h}$ -coherent extension of  $M_{X_0^h}$  over  $\Omega$  by  $M_{\Omega}^0 = \mathcal{D}^k M_{\Omega}^1$ . Doing this for each point in S we get a coherent  $\mathcal{O}_{X_h}$ -submodule  $M^0 \subset M$  such that  $\mathcal{D}(\mathfrak{g}_{X_h})M^0 = M$ .  $\Box$ 

Assume that M = M(\*S) and that M is coherent over  $\mathcal{O}_{X_h}(*S)$ . A nice example by Deligne in [Mal96] shows that M need not contain a coherent  $\mathcal{O}_{X_h}$ -module  $M^0$ , generating M over  $\mathcal{O}_{X_h}(*S)$ ; in particular one must expect  $\operatorname{coh}(\mathfrak{g}_{X_h}) \notin \operatorname{coh}^l(\mathfrak{g}_{X_h})$ . On the other hand, B. Malgrange [loc. cit.] proves

$$\operatorname{coh}(\mathcal{O}_{X_h}(*S)) \cap \operatorname{coh}(T_{X_h}) \subset \operatorname{coh}^l(T_{X_h}),$$

generalizing Deligne's result  $\operatorname{coh}(\mathcal{O}_{X_h}(*S))\cap \operatorname{coh}_{rs}^{tf}(T_{X_h}) \subset \operatorname{coh}_{rs}^{l}(T_{X_h})$ , where  $\operatorname{coh}_{rs}^{tf}(T_{X_h})$  is the category of torsion free coherent  $T_{X_h}$ -modules with (curve) regular singularities (*Def.* 6.3.3). Therefore, if  $\mathfrak{g}_{X_h}$  is *transitive*, by Theorem 4.4.1 we have

$$\operatorname{coh}(\mathcal{O}_{X_h}(*S)) \cap \operatorname{coh}(\mathfrak{g}_{X_h}) \subset \operatorname{coh}(\mathfrak{g}_X)$$
 (algebraic modules).

The coherent  $\mathcal{O}_X$ -submodule  $M^0 \subset M \in \operatorname{coh}(\mathcal{O}_{X_h}(*S)) \cap \operatorname{coh}(T_{X_h})$  is defined by Deligne and Malgrange using a certain spectral conditions for the action of  $t\partial_t$ , where t = 0 is a local equation at a non-singular point of S, giving 'normal Jordan generators' for M as in [Lev75]. One may speculate whether it is possible apply this idea to modules M over a non-transitive Lie algebroid  $\mathfrak{g}_{X_h}$ , defining  $M^0 \subset M \in \operatorname{coh}(\mathcal{O}_{X_h}(*S)) \cap \operatorname{coh}(\mathfrak{g}_X)$  by a spectral condition for the action of some distinguished sub-algebroid in  $\mathfrak{g}_{X_h}$ .

### 5. PROLONGING OVER codim $\geq 2$

5.1. Coherent extensions. When  $S \subset X$  is a closed subset in a (complex analytic) space we say that  $\operatorname{codim}_X S \geq 2$  if every point in S is contained in a subspace S', and there exists an open set  $\Omega$  such that  $S \cap \Omega \subset S' \cap \Omega$  and  $\operatorname{codim}_{\Omega} S' \geq 2$ .

Consider the normalization morphism of a reduced space or a general reduced scheme X,

$$f: X' \to X.$$

So X' is the disjoint union of the normalization of all the irreducible components of X. This is a finite morphism when X is a space. Also, f is almost always finite when X is a locally noetherian scheme; for instance, it suffices that  $\mathcal{O}_x$  is a G-ring at each point  $x \in X$ .

**Lemma 5.1.1.** (X is either a reduced complex analytic space or a locally noetherian reduced scheme such that the normalization morphism f is finite) Let  $S \subset X$  be a closed subset such that  $\operatorname{codim}_X S \ge 2$ . Put  $\Omega = X \setminus S$  and let  $j: \Omega \to X$  be the inclusion. Then:

- (1)  $j_*(\mathcal{O}_{\Omega}) \in \operatorname{coh}(\mathcal{O}_X)$ ; when X is normal, then the canonical mapping  $\mathcal{O}_X \to j_*(\mathcal{O}_{\Omega})$  is an isomorphism;
- (2) (X is a reduced noetherian scheme) Let  $N_{\Omega}$  be a coherent torsion free  $\mathcal{O}_{\Omega}$ -module (see Section 3.1 for the definition of torsion freeness). Then  $j_*(N_{\Omega})$  is coherent;
- (3) (X is a reduced complex analytic space) Assume that  $N_X$  is an  $\mathcal{O}_X$ -module which is locally generated by its sections and that  $N_\Omega$  is a coherent torsion free  $\mathcal{O}_\Omega$ -module. Then  $j_*(N_\Omega)$  is coherent.

Here (2) is essentially covered by a result of Grothendieck [Gro62, Exp. VIII, p. 6, Prop. 2.3], and (3) is a very special case of a result due to Serre [Ser66], but we provide simple direct proofs. When X is non-reduced  $j_*(\mathcal{O}_{\Omega})$  need not be coherent.

*Proof.* (1): The second part is in the complex analytic case Riemann's "Hebbarkeitssatz", see [Sch64]; in the algebraic case we recall the proof: One can assume that X is affine; then  $\operatorname{codim}_X(X \setminus \Omega) \geq 2$  implies that  $\mathcal{O}_X(\Omega) \subseteq \mathcal{O}_X$  when  $\operatorname{ht}(x) = 1$ ; by normality  $\mathcal{O}_X(X) = \bigcap_{\operatorname{ht}(x)=1} \mathcal{O}_X$ ; hence  $\mathcal{O}_X(\Omega) \subseteq \mathcal{O}_X(X)$ .

For the first part, consider the Cartesian square

$$\begin{array}{ccc} \Omega' & \stackrel{i}{\longrightarrow} & X' \\ g & & & \downarrow f \\ \Omega & \stackrel{j}{\longrightarrow} & X. \end{array}$$

One checks that  $\mathcal{O}_X \subset f_*(\mathcal{O}'_X)$  since X is reduced (the normalization is the union of all irreducible components of X, and these are integral). Then since formation of direct image is local on the base, so  $j^*f_* = g_*i^*$ ; hence, using  $i_*i^*(\mathcal{O}_{X'}) = \mathcal{O}_{X'}$ ,

$$j_*(\mathcal{O}_{\Omega}) = j_*j^*(\mathcal{O}_X) \hookrightarrow j_*j^*f_*(\mathcal{O}_{X'}) = j_*g_*i^*(\mathcal{O}_{X'}) = f_*i_*i^*(\mathcal{O}_{X'}) = f_*(\mathcal{O}_{X'}).$$

As f is finite,  $f_*(\mathcal{O}_{X'}) \in \operatorname{coh}(\mathcal{O}_X)$ ; from  $j_*(\mathcal{O}_\Omega) = j_*j^*(\mathcal{O}_X)$  it follows that  $j_*(\mathcal{O}_\Omega)$  is locally generated by its sections; hence since  $\mathcal{O}_X$  is noetherian (2.3.2),  $j_*(\mathcal{O}_\Omega) \in \operatorname{coh}(\mathcal{O}_X)$ .

(2): Put  $N_{\Omega}^{\vee} = Hom_{\mathcal{O}_{\Omega}}(N, \mathcal{O}_{\Omega})$ . Since X is noetherian, by Gabriel's theorem there exist  $N_e, N_X^{\vee} \in \operatorname{coh}(\mathcal{O}_X)$  such that  $j^*(N_e) = N_{\Omega}, j^*(N_X^{\vee}) = N_{\Omega}^{\vee}$  [BS58, Prop 2.]. As  $N_{\Omega}$  is coherent and torsion free there exists a nonempty open subset  $\Omega_1 \subseteq \Omega$ , with dense intersection with each irreducible component of X, such that  $N_{\Omega_1}$  is locally free; therefore, if K is the kernel of the canonical morphism  $N_{\Omega} \to N_{\Omega}^{\vee \vee}$ , then dim supp  $K < \dim \Omega$ , but  $N_{\Omega}$ is torsion free so K = 0. Hence,  $j_*$  being left exact, the canonical mapping

$$j_*(N_\Omega) \to j_*(N_\Omega^{\vee\vee})$$

is injective. Since  $N_{\Omega}$  has some coherent extension  $N_e$  to X, so  $j_*(N_{\Omega}) = j_*j^*(N_e)$ , implying that  $j_*(N_{\Omega})$  is locally generated by its sections. Therefore,  $\mathcal{O}_X$  being noetherian,  $j_*(N_{\Omega}) \in \operatorname{coh}(\mathcal{O}_X)$  will follow if  $j_*(N_{\Omega}^{\vee\vee}) \in \operatorname{coh}(\mathcal{O}_X)$ . By adjunction

$$j_*(N^{\vee\vee}) = j_*Hom_{\mathcal{O}_\Omega}(j^*(N_X^{\vee}), \mathcal{O}_\Omega) = Hom_{\mathcal{O}_X}(N_X^{\vee}, j_*(\mathcal{O}_\Omega));$$

hence  $j_*(N^{\vee\vee}) \in \operatorname{coh}(\mathcal{O}_X)$  by (1).

(3): By assumption there exist  $N_e = N, N_X^{\vee} = N^{\vee} \in \operatorname{coh}(\mathcal{O}_X)$  such that  $j^*(N_e) = N_{\Omega}$  and  $j^*(N_X^{\vee}) = N_{\Omega}^{\vee}$ . Then proceed as in (2).  $\Box$ 

Remarks 5.1.2. (1) Serre has proven a stronger result than (3) above; see [Ser66, Th.1]. Namely, the following statements are equivalent on a normal complex analytic space X for a coherent and torsion free  $\mathcal{O}_{\Omega}$ -module  $M_{\Omega}$ : (i) for each point  $s \in S$  there exists an open set  $U \subset X, x \in U$ , such that  $M_{\Omega \cap U}$  is generated by its sections; (ii)  $j_*(M_{\Omega}) \in \operatorname{coh}(\mathcal{O}_X)$ . Some other facts pertaining to the complex analytic case are exhibited in *loc. cit.*: (i) Let  $X = \mathbb{C}^2$  and  $S = \{0\}$ . There exist invertible sheaves on  $X \setminus S$  that are not extendible by a coherent sheaf on X. (ii) Let  $X = \mathbb{C}^n$ ,  $n \geq 3$ ,  $S = \{0\}$ , and let  $i : X \setminus S \to X$  be the inclusion. If M is a reflexive  $\mathcal{O}_{X \setminus S}$ -module, then if  $i_*(M)$  is coherent it is also reflexive, but it need not be locally free when M is locally free.

- (2) Assume that  $j : \Omega \to X$  is an open inclusion of complex manifolds, and put  $S = X \setminus \Omega$ , where  $\operatorname{codim}_X S \ge 2$ . Deligne proves the following result for the Lie algebroid  $T_{\Omega}$  [Del70, Cor. 5.8]: If a  $T_{\Omega}$ -module  $M_{\Omega}$ is coherent over  $\mathcal{O}_{\Omega}$ , then  $j_*(M_{\Omega})$  is coherent over  $\mathcal{O}_X$ . The proof is based on Hironaka's resolution of singularities and Grauert's finiteness theorem. Comparing to Serre's results we get: if  $U \subset X$  is a sufficiently small open neighbourhood of a point in S, then  $M_{U \setminus (U \cap S)}$ is generated by its global sections. Deligne's result cannot be generalised to general Picard Lie algebroids, which can be seen by taking an invertible sheaf  $M_{\Omega} = \lambda_{\Omega}$  on  $\Omega = \mathbb{C}^2 \setminus \{0\}$  that is not extendible to coherent  $\mathcal{O}_{\mathbb{C}^2}$ -module and put  $\mathfrak{g}_{\Omega} = \mathfrak{c}(\lambda_{\Omega})$ .
- (3) (1) in Lemma 5.1.1 implies that the canonical mapping  $T_X(X) \to T_X(\Omega)$  is a bijection when X is normal: if  $\delta \in T_X(\Omega)$  and  $f \in \mathcal{O}_X(X)$ , then  $\delta(f) \in \mathcal{O}_X(\Omega) = \mathcal{O}_X(X)$ . In particular,  $j_*(T_\Omega) = T_X$ .

The generalization to torsion modules is quite hard and we need some preparation.

We let prof  $M_x$  be the depth of an  $\mathcal{O}_x$ -module  $M_x$  of finite type.

**Theorem 5.1.3.** (X is a scheme which is locally a closed sub-scheme of a non-singular scheme, or a complex analytic space) Let  $Y \subset X$  be a closed subset and  $j : U \to X$  be the open immersion of  $U := X \setminus Y$ . Let  $M \in \operatorname{coh}(\mathcal{O}_X)$ . When X is a scheme assume:

(1) prof  $M_x \ge s - 1$  for all  $x \in U$  such that  $\operatorname{codim}_Y(\{x\}^- \cap Y) = 1$ . When X is a complex analytic space assume:

(1') Each point  $y \in Y$  has an open neighbourhood  $U_y \subset X$  such that prof  $M_x \ge s + \dim_y Y$  when  $x \in U_y$ .

Then  $R^p j_* j^*(M) \in \operatorname{coh}(\mathcal{O}_X)$  when  $p \leq s - 2$ .

Theorem 5.1.3 was proven for schemes by Grothendieck [Gro62, Exp. VIII, p. 13, Th.3.1] and when X is a complex analytic space it is due to Siu and Trautmann [Siu70], [Tra69]. We include a proof below when X is a scheme which is a little more direct than Grothendieck's proof.

Let p. d. N be the projective dimension of a module N over a regular ring R; let d(R) be the Krull dimension of R so prof R = d(R). We recall the Auslander-Buschsbaum relation [Mat86, Th. 19.1]:

(5.1.1) 
$$p. d. N + prof N = d(R).$$

That  $(1') \Rightarrow (1)$  can be seen using the next lemma, but we will not need this fact; the opposite implication is false.

**Lemma 5.1.4.** Let A be a local ring which is a quotient of a regular ring, let  $P \subset A$  be a prime ideal, and let q be an integer. Let N be an A-module of finite type. Then if prof  $N \ge q + \operatorname{coht} P$  it follows that prof  $N_P \ge q$ . Here coht P = d(A/P) is the coheight of a prime ideal. A stronger result is true without assuming that A is a quotient of a regular ring, namely that prof  $N \leq \text{prof } N_P + \text{coht } P$  [BS98, Lem. 9.3.2].

*Proof.* We consider M as a module over a regular ring R and P as prime ideal in R. Then p. d.  $M \leq d(R) - (q + \operatorname{coht} P) = d(R_P) - q$ . Therefore p. d.  $M_P = \sup\{i : Ext^i_{R_P}(M_P, R_P) \neq 0\} \leq p. d. M \leq d(R_P) - q$ . Hence by (5.1.1) prof  $M_x \geq q$ .  $\Box$ 

The following fact will be needed:

**Lemma 5.1.5** ([Gro62, Exp. III, p. 10, Prop. 3.3]). Let X be a locally noetherian scheme. The following are equivalent:

- (1)  $H_Z^i(M) = 0$ , when  $i \le n 1$ ;
- (2) prof  $M_z \ge n$  for all  $z \in Z$ .

We begin the proof of Theorem 5.1.3, following Grothendieck, by showing that one may assume that X is non-singular: The assertion of the theorem is of local nature, so we can assume that X can be embedded in a non-singular scheme. Let  $i : X \to X_{ns}$  be a closed immersion where  $X_{ns}$  is a regular scheme and put  $V_1 = i(V), Y_1 = i(Y), U_1 = X_{ns} \setminus Y_1$ , so  $\operatorname{codim}_V Y_1 \ge 2$ . Let  $j_1 : U_1 = X_{ns} \setminus Y_1 \to X_{ns}$  be the open inclusion and  $i_1 : U \to U_1 = i(j(U))$ be the restriction of i to  $U \subset X$ . Then  $j_1 \circ i_1 = i \circ j$ , so there exists an isomorphism of functors on  $D(\mathcal{O}_X), i_*Rj_* \cong Rj_{1*}i_{1*}$ , since  $i_*$  and  $i_{1*}$  are exact. In particular, using the fact that the direct image is local on the base, so  $i_{1*}j^* = j_1^*i_*$ , we get

$$i_*R^s j_*j^*(M) = R^s j_{1*}i_{1*}j^*(M) = R^s j_{1*}j_1^*i_*(M)$$

for each integer s. Here  $i_*(M)$  is a module such that  $\operatorname{prof} j_1^*(i_*(M))_x \geq s + \dim_y Y_1$ . So if one proves the theorem when X is non-singular then  $i_*(R^l j_* j^*(M)) \in \operatorname{coh}(\mathcal{O}_{X_{ns}})$  when  $l \leq s-2$ ; hence, since  $i_*$  is fully faithful,  $R^l j_* j^*(M) \cong i^* i_*(R^l j_* j^*(M)) \in \operatorname{coh}(\mathcal{O}_X)$ .

Until the end of the proof we will now asume that X is a non-singular scheme.

**Lemma 5.1.6.** Let  $N \in \operatorname{coh}(\mathcal{O}_X)$ ,  $S = \operatorname{supp} N$ , and  $Z = \overline{S \cap U} \cap Y$ . Let  $r \in \mathbb{Z}_{\geq 0}$  and assume that  $\operatorname{d}(\mathcal{O}_y) \geq r$  when  $y \in Z$ . Then

$$Hom_{\mathcal{O}_X}(N, R^s j_* j^*(\mathcal{O}_X)) \in \operatorname{coh}(\mathcal{O}_X)$$

when  $s \leq r - 2$ .

*Proof.* The long exact sequence in homology of the distinguished triangle

$$R\Gamma_Y(\mathcal{O}_X) \to \mathcal{O}_X \to Rj_*j^*(\mathcal{O}_X) \to$$

shows that we may instead prove

$$Hom_{\mathcal{O}_X}(N, H_Y^{s+1}(\mathcal{O}_X)) \in \operatorname{coh}(\mathcal{O}_X)$$

when  $s \leq r - 2$ .

Put  $N^0 = \Gamma_Y(N)$  and  $N^1 = N/N^0$ . Applying the contravariant left exact functor

$$T: N \mapsto Hom_{\mathcal{O}_X}(N, H^{s+1}_Y(\mathcal{O}_X))$$

to the short exact sequence

$$0 \to N^0 \to N \to N^1 \to 0$$

results in a short exact sequence

$$0 \to T(N^1) \to T(N) \to T(N)' \to 0$$

where  $T(N)' := \operatorname{Im}(T(N) \to T(N^0)) \subset T(N^0)$  is locally generated by its sections. Since supp  $N^0 \subset Y$ , clearly  $T(N^0) \in \operatorname{coh}(\mathcal{O}_X)$ ; therefore  $T(N)' \in$  $\operatorname{coh}(\mathcal{O}_X)$  because  $\mathcal{O}_X$  is noetherian. Now it will follow that  $T(N) \in \operatorname{coh}(\mathcal{O}_X)$ if  $T(N^1) = 0$ . First we have

$$T(N^1) = Hom_{\mathcal{O}_X}(N^1, H_Y^{s+1}(\mathcal{O}_X)) = Hom_{\mathcal{O}_X}(N^1, \Gamma_Z(H_Y^{s+1}(\mathcal{O}_X))).$$

Second, since  $R\Gamma_Z(\mathcal{O}_X) = R\Gamma_Z R\Gamma_Y(\mathcal{O}_X)$  there exists a spectral sequence  $H^p_Z(H^q_Y(\mathcal{O}_X)) \Rightarrow H^{p+q}_Z(\mathcal{O}_X);$  in particular  $\Gamma_Z(H^{s+1}_Y(\mathcal{O}_X))$  is a subquotient of  $H_Z^{s+1}(\mathcal{O}_X)$ ; since prof  $\mathcal{O}_y = d(\mathcal{O}_y) \ge r$  when  $y \in Z$  it follows that  $H_Z^{s+1}(\mathcal{O}_X) = 0$ , and hence  $\Gamma_Z(H_Y^{s+1}(\mathcal{O}_X)) = 0$ , when  $s+1 \leq r-1$ (Lem. 5.1.5). Therefore  $T(N^1) = 0$  when  $s \leq r - 2$ .

Proof of Theorem 5.1.3 when X is non-singular. Since M is coherent and X is non-singular there exists a biduality isomorphism

$$M \cong RHom_{\mathcal{O}_X}(RHom_{\mathcal{O}_X}(M,\mathcal{O}_X),\mathcal{O}_X).$$

Then adjunction gives

$$Rj_{*}j^{*}(M) = Rj_{*}j^{*}RHom_{\mathcal{O}_{X}}(RHom_{\mathcal{O}_{X}}(M,\mathcal{O}_{X}),\mathcal{O}_{X})$$
  
$$= Rj_{*}RHom_{\mathcal{O}_{U}}(j^{*}RHom_{\mathcal{O}_{X}}(M,\mathcal{O}_{X}),\mathcal{O}_{U})$$
  
$$= RHom_{\mathcal{O}_{X}}(RHom_{\mathcal{O}_{X}}(M,\mathcal{O}_{X}),Rj_{*}(\mathcal{O}_{U})).$$

giving a spectral sequence

$$Hom_{\mathcal{O}_X}(Ext^q_{\mathcal{O}_X}(M,\mathcal{O}_X), R^{q+p}j_*j^*(\mathcal{O}_X)) \Rightarrow R^pj_*j^*(M).$$

It now suffices to prove that the left side is coherent when  $p \leq s - 2$ .

Put  $S^q = \operatorname{supp} Ext^q_{\mathcal{O}_X}(M, \mathcal{O}_X)$  and  $Z^q = \overline{(S^q \cap U)} \cap Y$ .

b)  $d(\mathcal{O}_y) \ge q + s$  when  $y \in Z^q$ : For any  $y \in Y$  there exists  $x \in S^q \setminus Y$ such that  $y \in \{x\}^- \cap Y$  and  $d(\mathcal{O}_y) = d(\mathcal{O}_x) + 1$ . Then prof  $M_x \ge s - 1$  by (1), so by (5.1.1)

$$d(\mathcal{O}_y) = d(\mathcal{O}_x) + 1 = p. d. M_x + prof M_x + 1 \ge q + s - 1 + 1 = s + q,$$

where the inequality is because  $x \in S^q$ .

c) By b), prof  $\mathcal{O}_y = d(\mathcal{O}_y) \ge q + s$  when  $y \in Z^q$ . Hence by Lemma 5.1.6, with  $N := Ext_{\mathcal{O}_X}^q(M, \mathcal{O}_X),$ 

$$Hom_{\mathcal{O}_X}(Ext^q_{\mathcal{O}_X}(M,\mathcal{O}_X), R^{q+p}j_*j^*(\mathcal{O}_X)) \in \operatorname{coh}(\mathcal{O}_X)$$

when  $q + p \leq q + s - 2$ , i.e. when  $p \leq s - 2$ .  $\Box$ 

Let  $M \in \operatorname{coh}(\mathcal{O}_X)$  and put  $V = \operatorname{supp} M$ ,  $k = \dim V$ . We say that M is (k-) pure if every coherent  $\mathcal{O}_X$ -submodule  $N \subset M$  satisfies dim supp N = k. When M is not pure we will consider its filtration by supports (= the Krull filtration), which we recall. Let S(M) be the maximal coherent submodule  $N \subset M$  such that dim supp  $N < \dim \operatorname{supp} M$ . Then setting  $M_0 = M$  and inductively  $M_{i+1} = S(M_i)$  one has a decreasing filtration of length  $\leq \dim X$ 

$$0 = M_n \subset M_{n-1} \subset \dots \subset M_0 = M$$

where successive quotients  $L_i = M_i/M_{i+1}$  are pure.

**Corollary 5.1.7.** (X is of the same type as in Theorem 5.1.3) Let M be a pure coherent  $\mathcal{O}_X$ -module with support  $V \subset X$ . Let  $Y \subset V$  be a closed subset such that  $\operatorname{codim}_V Y \ge 2$  and let  $j : U \to X$  be the associated open immersion where  $U = X \setminus Y$ . Then  $j_*j^*(M) \in \operatorname{coh}(\mathcal{O}_X)$ .

*Proof.* When X is a scheme one may consider M as a module on V over the ring  $\mathcal{O}_V := \mathcal{O}_X / \operatorname{Ann}(M)$ , so one may assume that V = X. Then since M is pure with support X we have prof  $M_x \ge 1$  when  $\operatorname{codim}_Y(\{x\}^- \cap Y)$ , satisfying (1) in Theorem 5.1.3, so we may take s = 2.

To prove the assertion in the complex analytic case we employ (1'), which turns out a bit more complicated in spite of  $(1') \Rightarrow (1)!$  Put  $d = \dim V$  and let

$$S = \{x \in \operatorname{prof} M_x < d(M_x)\} \subset V$$

be the locus of points where M is not Cohen-Macaulay. We assert that this is a closed (analytic) subset such that  $\operatorname{codim}_V S \ge 2$ . Assuming this for a while we put  $Y_1 = S \cup Y$ , so  $\operatorname{codim}_V Y_1 \ge 2$  and  $M_{U_1}$  is Cohen-Macaulay, where  $U_1 := X \setminus Y_1$ . Letting  $j_1 : U_1 \to X$  be the open immersion Theorem 5.1.3 implies that  $(j_1)_*(j_1)^*(M) \in \operatorname{coh}(\mathcal{O}_X)$ ; since  $j_*j^*(M) \subset (j_1)_*(j_1)^*(M)$  is a submodule which is locally generated by its sections and  $\mathcal{O}_X$  is noetherian, it follows that  $j_*j^*(M) \in \operatorname{coh}(\mathcal{O}_X)$ .

It remains to prove  $\operatorname{codim}_V S \geq 2$ . Replacing the structure sheaf on  $V = \operatorname{supp} M$  we may assume that X = V and n = d, and since the assertion is local in X we may assume that  $X \subset X_{ns}$  where  $X_{ns}$  is either a complex manifold or a non-singular scheme; we thus consider M as an  $\mathcal{O}_{X_{ns}}$ -module with support on X. Let  $m \geq 1$  be an integer and put

$$D_m := \{x \in X : \operatorname{prof} M_x \le \operatorname{d}(M_x) - m\}$$
  
= 
$$\{x \in X : \operatorname{p.d.} M_x \ge \operatorname{d}(\mathcal{O}_x) - \operatorname{d}(M_x) + m\}$$
  
= 
$$\{x \in X : \operatorname{p.d.} M_x \ge n - d + m\}$$

where the second line follows from (5.1.1). Hence

$$D_m = \bigcup_{r \ge n-d+m} \operatorname{supp} Ext^r_{\mathcal{O}_{X_{ns}}}(M, \mathcal{O}_{X_{ns}}).$$

Since  $M \in \operatorname{coh}(\mathcal{O}_{X_ns})$  this shows that  $S = D_1$  is closed in X (see also [Mat86, exercise 24.2]), and a complex analytic subspace when X is a complex analytic space. Now  $\operatorname{codim}_{X_{ns}} Ext^r_{\mathcal{O}_{X_{ns}}}(M, \mathcal{O}_{X_{ns}}) \geq r \geq n-d+1$ , so  $\operatorname{codim}_X S \geq 2$  will follow if the  $\mathcal{O}_x$ -module  $Ext^{n-d+1}_{\mathcal{O}_{X_{ns}}}(M, \mathcal{O}_{X_{ns}})_x \cong$  $Ext^{n-d+1}_{\mathcal{O}_x}(M, \mathcal{O}_x)$  has dimension  $\leq d-2$  (M is coherent). Put  $A = \mathcal{O}_x$ ,  $N = M_x$ , and let  $p \subset A$  be a prime ideal of height n-d+1. Then since N is a pure A-module of finite type, prof  $N \geq 1$  and  $Ext^{n-d+1}_A(N, A)_p =$  $Ext^{n-d+1}_{A_p}(N_p, A_p)$ ; hence by (5.1.1), p. d.  $N_p \leq n-d+1-1 = n-d$ , and therefore  $Ext^{n-d+1}_A(N, A)_p = 0$ ; whence

$$\dim Ext_{\mathcal{O}_{X_{ns}}}^{n-d+1}(M,\mathcal{O}_{X_{ns}})_y \le d-2.$$

*Remark* 5.1.8. Note that in our definition a variety is always locally a closed subscheme of a non-singular scheme, so Theorem 5.1.3 and Corollary 5.1.7 are applicable to spaces.

5.2. Prolongation of smoothness. Let  $\pi : Y \to X$  be a proper homomorphism of spaces, S a closed subset of X, and put  $X_1 = X \setminus S$ . The mapping  $j : X_1 \to X$  is the canonical inclusion and  $\pi_1 : Y_1 \to X_1$  the corresponding base change of  $\pi$ . So the projection on the other coordinate  $i: Y_1 \to Y$  is an open inclusion and we have a Cartesian diagram:

(5.2.1) 
$$\begin{array}{ccc} Y_1 & \stackrel{i}{\longrightarrow} & Y \\ \pi_1 & & & \downarrow^{\pi} \\ X_1 & \stackrel{j}{\longrightarrow} & X. \end{array}$$

If  $\mathfrak{g}_Y \in \text{Lie}_Y$  we put  $\mathfrak{g}_{Y_1} = i^+(\mathfrak{g}_Y)$  (see 2.4.5). For a sub-space  $V \subset X$  the *V*-component of a coherent  $\mathfrak{g}_X$ -module *M* is the maximal coherent sub-module of *M* whose support is contained in *V*.

**Theorem 5.2.1.** Let  $\mathfrak{g}'_Y \subset \mathfrak{g}_Y$  be a sub-algebroid and M a coherent  $\mathfrak{g}_Y$ module containing, locally in X, a coherent  $\mathcal{O}_Y$ -submodule  $M^0 \subset M$  such that  $M = \mathcal{D}(\mathfrak{g}_Y)M^0$ ; this always holds in the algebraic case. Put V =supp M and let  $V = \bigcup_i^k V_i$  be an irreducible decomposition of V. Assume that  $V_i \cap V_j \cap \pi^{-1}(S) = \emptyset$  when  $i \neq j$ , and if  $V_i \cap \pi^{-1}(S) \neq \emptyset$ , then the  $V_i$ -component of M is pure and  $\dim \pi(V_i) \geq \dim S + 2$ . Assume also:

- (1) The canonical homomorphism  $\pi^*\pi_*(M)_y \to M_y$  is surjective when  $y \in \pi^{-1}(S)$ ;
- (2) The canonical homomorphism  $\pi^*\pi_*(\mathfrak{g}'_Y)_y \to \mathfrak{g}'_y$  is surjective when  $y \in \pi^{-1}(S)$ .

Then if  $i^!(M)$  is point-wise smooth along  $\mathfrak{g}'_{Y_1}$  it follows that M is pointwise smooth along  $\mathfrak{g}'_{Y_2}$ .

We shall consider non-relative versions of Theorem 5.2.1. First the case when M is torsion free is singled out, which may be of special interest, and we then need only the more elementary Lemma 5.1.1.

**Corollary 5.2.2.** Let  $\mathfrak{g}'_X$  be a sub-algebroid of a Lie algebroid  $(X, \mathcal{O}_X, \mathfrak{g}_X)$ and M a torsion free  $\mathfrak{g}_X$ -module.

- (1) (X is a variety) The following are equivalent:
  - (a) M is smooth at all points.
  - (b) M is smooth at all points of height 1.
- (2) (X is a complex analytic space) The following are equivalent:
  - (a) M is smooth at all points.
  - (b) There exists an open set  $\Omega \subset X$  with  $\operatorname{codim}_X(X \setminus \Omega) \ge 2$  such that M is smooth at all points  $p \in \Omega$ .

*Proof.* We need to prove  $(b) \Rightarrow (a)$ , and this follows from Theorem 5.2.1 by letting  $\pi : X \to X$  be the identity mapping. Here follows a direct proof: Let p be a point in X. Then there exists an affine (Stein) neighbourhood  $\Omega$  of p and a coherent  $\mathcal{O}_{\Omega}$ -submodule  $M_{\Omega}^0 \subset M_{\Omega}$  such that  $\mathcal{D}(\mathfrak{g}_{\Omega})M_{\Omega}^0 = M_{\Omega}$ . Put  $M'_{\Omega} := \mathcal{D}(\mathfrak{g}'_{\Omega})M_{\Omega}^0$ . As  $\Omega$  is affine (Stein) and  $\mathcal{D}(\mathfrak{g}'_{\Omega})$  is a union of coherent  $\mathcal{O}_{\Omega}$ -modules, it follows that  $\mathcal{D}(\mathfrak{g}'_{\Omega})$  generated by its sections (Cartan A in the analytic case) and since  $M'_{\Omega}$  is generated by its sections over  $\mathcal{D}(\mathfrak{g}'_{\Omega})$  it is also generated by its sections over  $\mathcal{O}_{\Omega}$ . Now, since the strong support of the  $\mathfrak{g}'_{\Omega}$ -module  $M'_{\Omega}$  is closed (*Prop.* 4.1.5), (b) implies both in the algebraic and the analytic case that there exists an algebraic (analytic) subset  $Z \subset \Omega$ ,  $\operatorname{codim}_{\Omega} Z \geq 2$ , such that  $M'_{\Omega \setminus Z} \in \operatorname{coh}(\mathcal{O}_{\Omega \setminus Z})$ . Letting  $j : \Omega \setminus Z \to \Omega$  be the inclusion mapping we have a canonical mapping

$$M'_p \to j_*(M'_{\Omega \setminus Z})_p$$

which is injective because  $M'_{\Omega}$  is pure and supp  $M'_{\Omega} = \Omega$ . Then Lemma 5.1.1 implies that  $j_*(M'_{\Omega \setminus Z})_p$  is the stalk of a coherent module, thus it is of finite type over  $\mathcal{O}_p$ ; since  $\mathcal{O}_p$  is noetherian it follows that  $M'_p$  is of finite type.  $\Box$ 

**Corollary 5.2.3.** Let  $\mathfrak{g}'_X \subset \mathfrak{g}_X$  be a sub-algebroid,  $L \in \operatorname{coh}(\mathfrak{g}_X)$  be pure,  $V = \operatorname{supp} L$ , and  $Y \subset V$  a subspace such that  $\operatorname{codim}_V Y \ge 2$ ; put  $U = X \setminus Y$ . Then if  $L_U$  is point-wise smooth along  $\mathfrak{g}'_U$  it follows that L is point-wise smooth along  $\mathfrak{g}'_X$ .

*Proof.* The proof is similar to that of Corollary 5.2.2. Namely, the problem is local at a point  $p \in Y$  so one may assume that L contains a coherent  $\mathcal{O}_X$ submodule  $L^0 \subset L$  such that  $L = \mathcal{D}(\mathfrak{g}_X)L^0$ . Put  $L' = \mathcal{D}(\mathfrak{g}'_X)L^0$ . By assumption  $L'_U \in \operatorname{coh}(\mathcal{O}_U)$  and since L' is pure

 $L' \subset j_*(L'_U)$ 

The right side is coherent by Corollary 5.1.7;  $\mathcal{O}_X$  is noetherian (2.3.2) and L' is locally generated by its sections; hence  $L' \in \operatorname{coh}(\mathcal{O}_X)$ .  $\Box$ 

**Corollary 5.2.4.** (X is a space) Let  $M \in \operatorname{coh}(\mathfrak{g}_X)$  be torsion free, and  $S \subset X$  a divisor such that M is smooth along  $\mathfrak{g}'_X \subseteq \mathfrak{g}_X$  at all points in  $X \setminus S$ . Then if  $M_x$  is smooth at one point x in each connected component of the non-singular locus of S, it follows that M is point-wise smooth in X.

Using Corollary 5.2.3 it is straightforward to modify the proof below to find a corresponding result for any pure module; we omit this generalisation.

Remark 5.2.5. Let X be a complex analytic manifold,  $S \subset X$  a divisor,  $\mathfrak{g}_X = T_X$  and  $\mathfrak{g}'_X = T_X(I_S)$  its sub-sheaf of derivations that preserve the ideal  $I_S$  of S. Let M be a  $T_X$ -module, coherent and torsion free over  $\mathcal{O}_X(*S)$ . In [Del70, Théorème 4.1(i) $\Rightarrow$ (ii)] it is proved that  $\phi^!(M)$  is smooth along  $\phi^+(T_X(I_S)) = T_C(I_{\phi^{-1}(S)})$  for each curve  $\phi : C \to X$  if this holds for each curve  $\phi : C \to \Omega$  where  $\Omega \subset X$  is a subset with  $\operatorname{codim}_X(X \setminus \Omega) \geq 2$ . By Corollary 6.1.3 this curve test is equivalent to M being smooth along  $T_X(I_S)$  (this also follows from [loc. cit]); hence Deligne's result follows from Corollary 5.2.4.

*Proof.* The assertion being local in X we can clearly assume that M contains a coherent  $\mathcal{O}_X$ -submodule  $M^0$  such that  $M = \mathcal{D}(\mathfrak{g}_X)M^0$ . Put  $M^1 = \mathcal{D}(\mathfrak{g}'_X)M^0 \in \operatorname{coh}(\mathfrak{g}'_X)$  and set

$$S_s = \{x \in S : M_x \text{ is smooth along } \mathfrak{g}'_x\} \\ = \{x \in S : M_x^1 \text{ is of finite type over } \mathcal{O}_x\}$$

Now s-supp  $M^1$  is a union of irreducible spaces (*Lem.* 2.3.6), s-supp  $M^1 \subset S$ , and  $S_s = S \setminus \text{s-supp } M^1 \subset S$  is an open subset containing a point from each component of S, implying that

$$\operatorname{codim}_X(\operatorname{s-supp} M^1) \ge 2.$$

Thus if  $\Omega = X \setminus \text{s-supp } M^1$ , then  $M_{\Omega}$  is point-wise smooth, so the assertion follows from Corollary 5.2.2.  $\Box$ 

The following is a sharpening of Theorem 4.2.7 for pure modules.

**Corollary 5.2.6.** Make the same assumptions and use the same notation as in Theorem 4.2.7. Let M be a pure  $\mathfrak{g}_Y$ -module which is smooth along  $\mathfrak{g}'_Y$ and put  $V = \operatorname{supp} M$ . Let  $V_1$  be a subset of V so that  $\operatorname{codim}_V V_1 \ge 2$ . Then if

(5.2.2) 
$$\operatorname{Im}(\pi^*(\mathfrak{g}'_X)_y \to \pi^*(\mathfrak{g}_X)_y) \subseteq \operatorname{Im}(\mathfrak{a}_y \to \pi^*(\mathfrak{g}_X)_y)$$

when  $y \in V \setminus V_1$ , it follows that  $\pi_+(M) \in D^b_s(\mathfrak{g}_X)$ .

*Proof.* The proof is close to that of Theorem 4.2.7 so we will avoid too much repetition. This time (5.2.2) implies

(5.2.3) 
$$(\pi^{-1}(\mathfrak{g}'_X)\mathcal{D}^n_{X\leftarrow Y})_y \subseteq (\mathcal{D}^n_{X\leftarrow Y}\mathfrak{a}_Y)_y \subseteq (\mathcal{D}^n_{X\leftarrow Y}\mathfrak{g}'_Y)_y,$$

when  $y \in Y - V_1$ . We factorize the proper morphism  $\pi : Y \to X$  into a closed embedding  $i : Y \to Z = X \times Y$  and a projection  $p : Z \to X$ where the restriction of p to the support of  $i_+(M)$  is proper. The right  $\mathcal{D}(\mathfrak{g}_Y)$ -module  $\mathcal{D}_{Z\leftarrow Y}$  is locally free, hence  $i_+(M) = i_*(\mathcal{D}_{Z\leftarrow Y} \otimes_{\mathcal{D}(\mathfrak{g}_Y)} M)$ and this is a pure  $\mathcal{O}_Z$ -module. Also, (5.2.2) holds for i and p separately where  $\mathfrak{g}'_Z = p^+(\mathfrak{g}'_X)$ . We may therefore prove the assertion when either  $\pi$ is a closed immersion or  $\pi$  is a projection such that the restriction of  $\pi$  to supp M is proper (*Prop.* 3.4.3).

 $\pi$  is a closed embedding: Put  $W = \pi(V)$  and  $W_1 = \pi(V_1)$ , so  $\pi_+(M)$  is pure with support W. Using the same argument as in the proof of Theorem 4.2.7 by (5.2.3)  $\pi_+(M) = \pi_*(\mathcal{D}_{X \leftarrow Y} \otimes_{\mathcal{D}(\mathfrak{g}_Y)} M)$  is smooth along  $\mathfrak{g}'_X$  in  $X \setminus W_1$ ; hence it is smooth everywhere (*Cor.* 5.2.3).

 $\pi: Y = Z \times X \to X$  is the projection on the second coordinate and M is a pure  $\mathfrak{g}_Y$ -module such that the restriction  $\pi': V \to X$  of  $\pi$  to the support  $V = \operatorname{supp} M$  is proper: Again the proof is similar to that of Theorem 4.2.7. We have

$$\mathcal{D}_{X \leftarrow Y} \otimes^L_{\mathcal{D}(\mathfrak{g}_Y)} M = \Omega^{\bullet}_{Y/X} \otimes_{\mathcal{O}_Y} M$$

so this is a complex of pure coherent  $\pi^*(\mathfrak{g}_X)$ -modules. Now (5.2.3) implies that this complex is smooth along  $\pi^*(\mathfrak{g}'_X)$  at all points in  $Y \setminus V_1$ , hence it is smooth everywhere (*Cor.* 5.2.3). Now the proof can be finished as in Theorem 4.2.7.  $\Box$ 

5.3. **Proof of Theorem 5.2.1.** In the proof, which is divided into lemmas, we will use the notation in the Cartesian diagram (5.2.1).

**Lemma 5.3.1.** Let  $M^i \in coh(\mathcal{O}_Y)$ ,  $i \in I$ , be an inductive system over a directed partial ordered set I. Then

(1)  $\lim M^i$  is locally generated by its sections.

(2)  $\pi_*(\lim_{\longrightarrow} M^i) = \lim_{\longrightarrow} \pi_*(M^i).$ 

*Proof.* (1): Let  $y \in Y$ . In the algebraic case let  $\Omega$  be an affine neighbourhood of y, in the complex analytic case y is contained in a Stein neighbourhood  $\Omega \subset Y$  and 'Theorem A' holds for Stein spaces (see [GR79]). Hence in either case  $M_y^i \cong \mathcal{O}_y \otimes_{\mathcal{O}_Y(\Omega)} M^i(\Omega)$ . Since inductive limits commute with  $\mathcal{O}_y \otimes \cdot$  (see [God73, II, 1.11]), we get

$$\mathcal{O}_y \otimes_{\mathcal{O}_Y(\Omega)} (\lim_{\to} M)(\Omega) \cong \lim_{\to} (\mathcal{O}_y \otimes_{\mathcal{O}_Y(\Omega)} M^i(\Omega)) \cong \lim_{\to} M^i_y.$$

(2): This follows by formal reasons if one admits that by properness the derived functor  $R\pi_*$  has a right adjoint  $\pi^!(\cdot)$ ; see [Har66], [BS76].

Here follows a more self-contained proof. We have a canonical homomorphism  $\phi$ : lim  $\pi_*(M^i) \to \pi_*(\lim M^i)$ . This is an isomorphism if it induces an isomorphism between stalks over any point  $x \in X$ . Therefore, by the last line in the proof of (1), it suffices to prove that the canonical mapping

$$\phi_x: \lim \pi_*(M^i)_x \to \pi_*(\lim M^i)_x$$

is an isomorphism. Let  $m \in \pi_*(\lim_{\to} M^i)_x$  be represented by the section

$$m_{\Omega} \in \pi_*(\lim_{\to} M^i)(\Omega) = (\lim_{\to} M^i)(\pi^{-1}(\Omega))$$

where  $\Omega \subset X$  is an open neighbourhood of x. If U is an open subset in  $\pi^{-1}(\Omega)$  we denote by  $m_U$  the restriction of  $m_\Omega$  to U. Every point  $y \in \pi^{-1}(\Omega)$ has an open neighbourhood  $U_y \subset Y$  and an integer  $n_y$  such that  $m_{U_y}$  is represented by a section  $m_{U_y}^{n_y} \in M^{n_y}(U_y)$ . Since  $\pi$  is proper it suffices with a finite number of the  $U_y$  to cover  $\pi^{-1}(x)$ ; let U be their union and n be the maximum of the corresponding integers  $n_y$ . Since I is directed it follows that  $m_U$  is represented by a section  $m_U^n \in M^n(U)$ . Now  $\pi : Y \to X$  is a continuous proper mapping between locally compact topological spaces, hence it is closed; hence the open sets  $\pi^{-1}(\Omega')$ , where  $\Omega'$  is an open subset of X, form a basis for the open neighbourhoods of  $\pi^{-1}(x)$ . So there exists an open neighbourhood  $\Omega'$  of x such that  $\Omega' \subset \pi(U) \subset \Omega$ , and therefore  $m_{\pi^{-1}(\Omega')}^n \in \pi_*(M^n)(\Omega')$ ; let  $m_x^n \in \pi_*(M^n)_x$  be its germ at x. Letting  $i_n$ :  $\pi_*(M^n)_x \to \lim_{x \to \infty} \pi_*(M^i)_x$  be the canonical mapping we have a mapping  $\psi_x : \pi_*(\lim_{d \to i} M^i)_x \to \lim_{d \to i} \pi_*(M^i)_x$ , well-defined by  $m_x \mapsto i_n(m_x^n)$ . One checks that  $\phi_x \circ \psi_x = \operatorname{id} \operatorname{and} \psi_x \circ \phi_x = \operatorname{id}$ .  $\Box$ 

**Lemma 5.3.2.** Let  $M^i \in \operatorname{coh}(\mathcal{O}_Y)$ ,  $i \in I$ , be as above,  $\phi_i : M^i \to M :=$ lim  $M^i$  the canonical homomorphism, and  $\psi_i : \pi^* \pi_*(M^i) \to M$  be the composition of the canonical homomorphisms  $\pi^*\pi_*(M^i) \to M^i \to M$ . Assume that the canonical homomorphism  $\pi^*\pi_*(M) \to M$  is surjective. Then for any  $i \in I$  there exists, locally in X, an index  $n = n(i) \in I$  such that

$$\operatorname{Im}(\phi_i) \subset \operatorname{Im}(\psi_{n(i)}).$$

*Proof.* Lemma 5.3.1 and the fact that  $\pi^*$  commutes with lim [God73, II, 1.11] implies that we have a surjection

$$\lim_{\to} \pi^* \pi_*(M^i) \cong \pi^* \pi_* \lim_{\to} M^i \to \lim_{\to} M^i.$$

It is obvious that locally near a point  $y \in Y$  one can find such an index  $n_y = n(i, y)$ . Then, since  $\pi$  is proper, for any  $x \in X$  one can reason as in the proof of Lemma 5.3.1 to see that there exists an index  $n(i, x) \in I$  and a neighbourhood  $\Omega$  of x such that

$$\operatorname{Im}(\phi_i)_{\pi^{-1}(\Omega)} \subset \operatorname{Im}(\psi_{n(i,x)})_{\pi^{-1}(\Omega)}.$$

Of course, a proper and surjective morphism may blow up subsets of codimension 2 to subsets of codimension 1. On the other hand, in the complex analytic case Hartogs' theorem states (see e.g. [BS76, p. 42]): If Y is a reduced complex Stein analytic space, K a compact subset such that  $Y \setminus K$  has no relatively compact irreducible components in Y, and the depth of  $\mathcal{O}_y$  is  $\geq 2$  for all  $y \in K$ , then  $\mathcal{O}_Y(Y) = \mathcal{O}_Y(Y \setminus K)$ . The main ingredient in the proof of Theorem 5.2.1 is the following coherence result which maybe can be thought of as Hartogs' type along the fibre and Riemann's Hebbarkeitssatz on the base.

**Lemma 5.3.3.** Let  $M^i \in \operatorname{coh}(\mathcal{O}_Y)$ ,  $i \in I$ , be an inductive system of modules such that, with  $M_1 = \lim_{\to} M^i$ , one has  $i^*(M_1) \in \operatorname{coh}(\mathcal{O}_{Y_1})$ . Assume that  $V = \operatorname{supp} M_1 \subset X$  is a subspace and let  $V = \bigcup_i^k V_i$  be an irreducible decomposition. Assume that  $V_i \cap V_j \cap \pi^{-1}(S) = \emptyset$  when  $i \neq j$ , and if  $V_i \cap \pi^{-1}(S) \neq \emptyset$ then the  $V_i$ -component  $M_1^{V_i} \subseteq M_1$  is pure and dim  $\pi(V_i) \geq \dim S + 2$ . Put

$$N = \operatorname{Im}(\pi^* \pi_*(M_1) \to M_1).$$

Then if M' is an  $\mathcal{O}_Y$ -submodule of N which is locally generated by its sections, it follows that  $M' \in \operatorname{coh}(\mathcal{O}_Y)$ .

Note that  $\pi(V_i)$  will be a subspace since  $\pi$  is proper, by Remmert's theorem in the complex analytic case.

*Proof.* Let  $y \in \pi^{-1}(S)$ . Since the problem is local near  $\pi(y)$  we can and will shrink X to any neighbourhood of  $\pi(y)$  when necessary.

a) It suffices to prove the assertion when  $M_1$  is pure and V is irreducible: As the  $\mathcal{O}_X$ -submodules  $M_1^{V_i} \subset M$  are pure we have  $\bigoplus_{i=1}^k M_1^{V_i} \subset M_1$ , and since  $V_i \cap V_i \cap \pi^{-1}(S) = \emptyset$  when  $i \neq j$ , one may shrink X so that  $V_i \cap V_j = \emptyset$ when  $i \neq j$ ; hence  $\bigoplus_i^k M_1^{V_i} = M_1$ . Therefore, if  $M' \cap M_1^{V_i} \in \operatorname{coh}(\mathcal{O}_Y)$ ,  $i = 1, \ldots k$ , it will follow that  $M' \in \operatorname{coh}(\mathcal{O}_Y)$ .

b) If  $M_1$  is pure, then  $\pi_*(M_1)$  is pure: Factorize  $\pi = p \circ i$  where  $i: Y \to Y \times X$  is the graph morphism and  $p: Z = Y \times X \to X$  the projection on the second factor. Clearly  $N = i_*(M)$  is pure with irreducible support  $V_1 = i(V)$ . Since  $\pi_*(M_1) = p_*i_*(M)$  it therefore suffices to prove: If N is a coherent  $\mathcal{O}_Z$ -module such that the restriction of p to a morphism supp  $N \to X$  is proper, then  $p_*(N)$  is pure.

By properness  $W := \operatorname{supp} p_*(N) \subset X$  is a subspace. Suppose the contrary, that there exists a non-zero pure submodule  $L \subset p_*(N)$  such that dim supp  $L < \dim W$ . Let  $\phi : p^*(L) \to N$  be the composed morphism  $p^*(L) \to p^*p_*(N) \to N$ . Then dim supp  $\operatorname{Im} \phi < \dim \operatorname{supp} N$ ; N is pure; hence  $\phi = 0$ ; therefore, projections being flat, the canonical morphism  $p^*p_*(N) \to N$  is 0; hence  $p_*(N) = 0$ , contradicting the assumption that  $L \neq 0$ . c) If  $M_1$  is pure and V is irreducible, then  $H^0_{[T]}(M_1) = 0$  when  $T = \pi^{-1}(S)$ : Suppose the contrary, that  $H^0_{[T]}(M_1) \neq 0$ . Since  $M_1$  is pure with irreducible support V the submodule  $H^0_{[T]}(M_1) \subset M_1$  also has support V; hence  $V \subset T$ , contradicting the relation dim  $\pi(V) \geq \dim \pi(T) + 2$ .

d)  $\pi_*(M_1) \in \operatorname{coh}(\mathcal{O}_X)$ : By c) and since  $\pi_*$  is left exact we have an inclusion

$$\pi_*(M_1) \subseteq \pi_* i_* i^*(M_1) = (\pi \circ i)_* i^*(M_1) = (j \circ \pi^1)_* i^*(M_1) = j_*(\pi_*^1(i^*(M_1))).$$

By Lemma 5.3.1  $\pi_*(M_1)$  is locally generated by its sections; then since  $\mathcal{O}_X$  is noetherian (2.3.2) it will follow that  $\pi_*(M_1) \in \operatorname{coh}(\mathcal{O}_X)$  if we prove

$$j_*(\pi^1_*(i^*(M_1))) \in \operatorname{coh}(\mathcal{O}_X).$$

Since  $i^*(M_1) \in \operatorname{coh}(\mathcal{O}_{Y_1})$  and properness is preserved under base change, by Grauert's theorem [Gra60] or Grothendieck's theorem [GD61, 3.2.1]

$$\pi^1_*(i^*(M_1)) \in \operatorname{coh}(\mathcal{O}_{X_1}).$$

Also by properness and since  $i^*(M_1) \in \operatorname{coh}(\mathcal{O}_{Y_1})$ , there exists a coherent  $\mathcal{O}_Y$ -submodule  $M_1^0 \subset M_1$  in a neighbourhood of  $\pi^{-1}(S)$  such that  $i^*(M_1^0) = i^*(M_1)$ . As the formation of direct image is local on the base, so  $j^*\pi_* = \pi_*^1 i^*$ , it follows that  $M := \pi_*(M_1^0)$  is a coherent extension of  $\pi_1^*(i^*(M_1))$ . We thus have a coherent  $\mathcal{O}_X$ -module M, such that  $j^*(M) = \pi_*^1(i^*(M_1))$  is pure by b), whose support  $\pi(V)$  contains S and  $\operatorname{codim}_{\pi(V)} S \ge 2$ ; hence by Corollary 5.1.7  $j_*(\pi_*^1(i^*(M_1))) \in \operatorname{coh}(\mathcal{O}_X)$ .

e)  $M' \in \operatorname{coh}(\mathcal{O}_Y)$ : Let  $L^i$  and  $N^i$  be the kernel and image of the canonical homomorphism  $\pi^* \pi_*(M^i) \to M^i$ , so we have a short exact sequence

$$0 \to L^i \to \pi^* \pi_*(M^i) \to N^i \to 0.$$

Now  $M^i \in \operatorname{coh}(\mathcal{O}_Y)$  and  $\pi$  is proper so  $\pi_*(M^i) \in \operatorname{coh}(\mathcal{O}_X)$  ([Gra60], resp. [GD61, 3.2.1]),  $\pi^*$  is right exact and  $\mathcal{O}_Y$  is coherent, hence  $\pi^*\pi_*(M^i) \in \operatorname{coh}(\mathcal{O}_Y)$ ; therefore  $L^i, N^i \in \operatorname{coh}(\mathcal{O}_Y)$  ([GD71, 0. Cor. 5.3.4]). Clearly we have canonical homomorphisms  $N^i \to N^j$ , and  $L^i \to L^j$ ,  $i \leq j$ , so put  $L = \lim_{\to} L^i$  and  $N = \lim_{\to} N^i$  (the sheaves associated to the presheaves). Now

$$\pi^*\pi_*(M_1) = \pi^* \lim_{\to} \pi_*(M^i) = \lim_{\to} \pi^*\pi_*(M^i),$$

where the first step follows from Lemma 5.3.1 and the second from [God73, II, 1.11], and since the index set **N** is directed the functor  $\lim_{\to}$  is exact, we get a short exact sequence

$$0 \to L \to \pi^* \pi_*(M_1) \to N \to 0.$$

By d)  $\pi^*\pi_*(M_1) \in \operatorname{coh}(\mathcal{O}_Y)$  (for the same reason above that the  $\pi^*\pi_*(M^i) \in \operatorname{coh}(\mathcal{O}_Y)$ ) and  $L^i \in \operatorname{coh}(\mathcal{O}_Y)$ , so L is locally generated by its sections (*Lem.* 5.3.1); as  $\mathcal{O}_Y$  is noetherian this implies that  $L \in \operatorname{coh}(\mathcal{O}_Y)$ ; hence  $N \in \operatorname{coh}(\mathcal{O}_Y)$  [GD71, 0. Prop 5.4.2], and again since  $\mathcal{O}_Y$  is noetherian it follows that  $M' \in \operatorname{coh}(\mathcal{O}_Y)$ .  $\Box$ 

Proof of Theorem 5.2.1. Let  $y \in \pi^{-1}(S)$ . Since the problem is local near  $x = \pi(y)$  we can and will shrink X to any small neighbourhood of x when necessary and we can assume that  $M = \mathcal{D}(\mathfrak{g}_Y)M^0$ . Arguing as in a) in the proof of Lemma 5.3.3 it suffices to prove that  $M' := \mathcal{D}(\mathfrak{g}'_Y)M^0 \in \operatorname{coh}(\mathcal{O}_Y)$ 

when M is pure and V is irreducible, and we then know as in b) that  $\pi_*(M)$ is pure.

The modules  $M^i = \mathcal{D}^i(\mathfrak{g}_Y) M^0 \in \operatorname{coh}(\mathcal{O}_Y), i \in \mathbf{N}$ , form an inductive system by inclusion  $M^i \subset M^j$  when  $i \leq j$ , and it follows from Lemma 5.3.2, possibly after shrinking X around x, that there exists an integer  $n_0$  such that  $M^0$  is contained in the image of the canonical homomorphism  $\pi^*\pi_*(M^{n_0}) \rightarrow$  $M^{n_0}$ ; set  $M_1 = \mathcal{D}(\mathfrak{g}'_V) M^{n_0}$ .

Now note that  $\pi^*\pi_*(\mathfrak{g}'_Y)$  is a Lie algebroid on Y in a natural way and that (2) is a homomorphism of Lie algebroids; since  $N := \text{Im}(\pi^* \pi_*(M_1) \to M_1)$ clearly is a  $\pi^*\pi_*(\mathfrak{g}'_V)$ -module (2) implies that

$$\mathfrak{g}'_Y \cdot N \subset N,$$

and since  $M^0 \subset N$  it follows that

 $M' \subset N.$ 

The  $\mathcal{O}_Y$ -coherent submodules  $M_1^i = \mathcal{D}^i(\mathfrak{g}'_Y)M^{n_0} \subset M_1$  form an inductive system by inclusion  $M_1^i \subset M_1^j$ ,  $j \ge i$ , its limit  $M_1 = \lim M_1^i$  is pure with irreducible support V, and dim  $\pi(V) \ge \dim S + 2$ . By assumption  $i^*(M_1) \in \operatorname{coh}(\mathcal{O}_{Y_1})$ ; therefore since  $M' = \lim \mathcal{D}^i(\mathfrak{g}'_Y)M^0$  is locally generated by its sections (Lem. 5.3.1) it follows by Lemma 5.3.3 that  $M' \in \operatorname{coh}(\mathcal{O}_Y)$ .  $\Box$ 

### 6. Regular singularities

6.1. Torsion free modules. Say that a set of vectors  $\delta_i$  in a Lie algebroid  $(A, \mathfrak{g}, \alpha)$  generates  $\mathfrak{g}$  if the subspace  $\sum A\delta_i \subseteq \mathfrak{g}$  is not contained in any proper sub-algebroid of  $\mathfrak{g}$ .

The following theorem follows from a straightforward application of the integrability theorem (Th. 2.3.4).

**Theorem 6.1.1.** (X is a space) Let  $\mathfrak{g}'_X$  be a sub-algebroid in a Lie algebroid  $\mathfrak{g}_X$  and M a  $\mathfrak{g}_X$ -module. Let x be a point in X,  $M_x^0$  an  $\mathcal{O}_x$ -submodule of finite type that generates  $M_x$  over  $\mathfrak{g}_x$ , and  $M'_x \subseteq M'_x$  the  $\mathfrak{g}'_x$ -module that is generated by  $M_x^0$ . Let  $\delta_i$  be generators of the Lie algebroid  $\mathfrak{g}'_x$ . Then the following are equivalent:

- (1) M is smooth at x;
- (2) Each  $\delta_i$  generates an  $\mathcal{O}_x$ -module of finite type

$$\sum_k \delta_i^k \cdot M_x^0;$$

- (3)  $M'_x$  is smooth along all the vectors  $\delta_i$  in  $\mathfrak{g}'_x$  (Def. 4.1.2); (4) Put inductively  ${M'}_x^n = \mathfrak{g}'_x {M'}_x^{n-1} + {M'}_x^{n-1}$  for  $n \ge 1$ . The increasing sequence of mappings

$$k_x \otimes_{\mathcal{O}_x} M'^n_x \to k_x \otimes_{\mathcal{O}_x} M'^{n+1}_x \to \cdots$$

is eventually surjective (here  $k_x = \mathcal{O}_y/\mathfrak{m}_y$ , while  $k_x = \mathbf{C}$  when X is a complex analytic space).

*Proof.* Clearly  $M'_x = \bigcup_{n \ge 0} M'^n_x$ , and M is smooth at x if and only if  $M'_x$ is of finite type over  $\mathcal{O}_x$ .

The implications  $(1) \Rightarrow (2), (2) \iff (3), (1) \Rightarrow (4)$  are evident.  $(4) \Rightarrow (1)$ : It suffices to note that by Nakayama's lemma, if the mapping  $k \otimes_{\mathcal{O}_x} M'_x^n \rightarrow k \otimes_{\mathcal{O}_x} M'_x^{n+1}$  is surjective then  $M'_x = M'_x^{n+1}$ .

(2)  $\Rightarrow$  (1): See Section 2.3 for the notation. Let  $\mathcal{D}_x = \mathcal{D}_x(\mathfrak{g}'_X)$  be the enveloping ring of differential operators of  $\mathfrak{g}'_x$ ,  $(S_x, \cdot, \{\cdot, \cdot\})$  the Poisson algebra that is defined on  $S_x := \operatorname{gr} \mathcal{D}_x$ , and let  $\sigma : \mathfrak{g}'_x \to S_x$  be the canonical morphism. Let  $G(M'_x)$  be the graded module over  $S_x$  that is associated to the canonical filtration  $M'^n_x = \mathcal{D}^n_x M'^0_x \subseteq M'^{n+1}_x, \cup M'^n_x = M'_x$ . Then  $G(M'_x)$  is a module of finite type over the noetherian algebra  $S_x$ , and the characteristic ideal  $J(M')_x \subseteq S_x$  is involutive (Th. 2.3.4); by assumption  $\sigma(\delta_i) \in J(M')_x$ , hence  $\sigma([\delta_i, \delta_j]) = \{\sigma(\delta_i), \sigma(\delta_j)\} \in J(M')_x$ ; the  $\delta_i$  generate  $\mathfrak{g}'_x$ ; hence  $\sigma(\mathfrak{g}'_x) \subseteq J(M')_x$ . Therefore there exists an integer n such that the  $S_x$ -module of finite type  $G(M'_x)$  is a module of finite type over  $R_x := S_x / (\sigma(\mathfrak{g}'_x))^n$ ; since clearly  $R_x$  is of finite type over  $\mathcal{O}_x$  it follows that  $G(M'_x)$  is of finite type over  $\mathcal{O}_x$ . This implies, by an easy argument using the short exact sequences  $0 \to M'^{n-1}_x \to M'^n_x \to G^n(M'_x) \to 0$ , that  $M'_x$  is of finite type over  $\mathcal{O}_x$ .  $\Box$ 

We shall now work out curve tests for smoothness. All curves C are assumed to be irreducible and non-singular, and morphisms  $\phi: C \to X$  are assumed to be locally closed embeddings of spaces; when we say that C is a curve in X we mean such a morphisms  $\phi$ . An analytic curve in a variety Xis a morphism of schemes  $\phi: C \to X$  where  $C = \operatorname{Spec} k[[x]]$ . If  $\phi: C \to X$ is a curve we set (Sec. 2.4.5)

$$\mathfrak{g}_C := \phi^+(\mathfrak{g}_X), \text{ and } \mathfrak{g}'_C := \phi^+(\mathfrak{g}'_X).$$

We thus have homomorphisms of Lie algebroids  $(\phi, \phi') : \mathfrak{g}_C \to \mathfrak{g}_X$ , and  $\mathfrak{g}'_C \to \mathfrak{g}'_X$ .

Let  $\delta \in \mathfrak{g}'_X(X_{reg})$  be a section defined over the non-singular locus of X. Locally one can assume that  $\delta = h\delta^1$ , where  $\alpha(\delta^1) \in T_X(\Omega)$  is non-zero outside an algebraic (analytic) subset  $X_{crit} \subset X$  such that  $\operatorname{codim}_X X_{crit} \geq$ 2. Now if X is a complex analytic space and x is a point in the nonsingular locus of  $\Omega$  there exists locally integral curves for  $\delta$ . More precisely, by Frobenius' theorem there exists a locally closed embedding of a curve  $\phi : C \to \Omega \subset X$  such that  $\alpha(\delta)(I_{C_x}) \subseteq I_{C_x}$ , where  $I_{C_x}$  is the ideal of a germ  $\phi(C)_x$  of the curve at a point  $x \in X$ . When X is algebraic there exist analytic (formal) solutions  $\phi$ , i.e. there exist ideals  $\hat{I}(x) \subset \hat{\mathcal{O}}_x$  so that  $\alpha(\delta)(\hat{I}(x)) \subseteq \hat{I}(x)$  and  $\hat{\mathcal{O}}_x/\hat{I}(x) = k[[x]]$ , but algebraic integral curves in general do not exist. If one insists that everything be algebraic we say that  $\mathfrak{g}'_X \in \operatorname{Lie}_X$  has good generators at a point  $x \in X$  if:

(G): There exist an open neighbourhood U of x and generators  $\delta_1, \ldots, \delta_r \in \mathfrak{g}'_X(U)$  of  $\mathfrak{g}'_U$  (as Lie algebroid) such that for every point  $y \in U$  there exists a locally closed embedding of an algebraic curve  $\phi^i : C^i \to U$ ,  $y \in \phi^i(C^i)$ , satisfying  $\alpha(\delta_i(x))(I^i(x)) \subset I^i(x)$  ( $I^i(x)$ ) is the ideal of  $\phi^i(C^i)_x$ ).

For example, when X is an algebraic manifold and S a non-singular divisor whose ideal is  $I_S$ , then  $T_X(I_S)$  has good generators at each point  $x \in X$ .

We say that a vector  $\delta \in \mathfrak{g}_x$  is *transversal* to a divisor S at  $x \in S$ , if there exists an ideal  $I_x \subset \mathcal{O}_x$  so that  $I_S + I_x = \mathfrak{m}_x$ ,  $I_S \nsubseteq I_x$ , and  $\alpha(\delta)_x \in T_x(I_x)$ 

 $(I_S \text{ is the ideal of } S)$ ; it is transversal to S if it is transversal at each point  $y \in S$ .

Say that a space X is non-singular in codimension 1 (or non-singular outside codim  $\geq 2$ ) if there exists a closed subset  $C \subset X$  such that  $\operatorname{codim}_X S \geq 2$  such that  $X_0 = X \setminus S$  is a non-singular space. When X is a scheme this means that  $\mathcal{O}_x$  is a regular ring when x is a point of height 1. This holds for example when X is normal.

Below we let  $\phi^!(M)$  denote the non-derived inverse image (3.4). In the notation above we have:

**Theorem 6.1.2.** (X is a space which is non-singular in codimension 1; when X is a variety we assume that the field k is uncountable) Let M be a coherent torsion free  $\mathfrak{g}_X$ -module which is point-wise smooth along  $\mathfrak{g}'_X$  outside a divisor S in X. Let  $S_0$  be an open subset of the non-singular locus  $S_{reg}$  of S, intersecting each component of  $S_{reg}$ . Assume that  $\mathfrak{g}'_X$  locally has generators which are transversal to  $S_0$ . Then the following are equivalent:

- (1) M is point-wise smooth in X;
- (2)  $\phi^!(M)$  is smooth along  $\mathfrak{g}'_C$  for any analytic curve  $\phi: C \to X$  such that  $\phi(C) \cap S \subseteq S_0$ ;
- (3)  $\phi^!(M)$  is smooth along  $\mathfrak{g}'_C$  for any analytic curve  $\phi: C \to X$ .

Assume that X is a variety and that  $\mathfrak{g}'_X$  has good generators at each point  $x \in S_0$ . Then (1-3) are equivalent to:

(4)  $\phi^!(M)$  is smooth along  $\mathfrak{g}'_C$  for any curve  $\phi: C \to X$  such that  $\phi(C) \cap S \subseteq S_0$ .

The annoying condition that  $\mathfrak{g}'_X$  be generated by vectors that preserve curves that are transversal to  $S_0$  can be removed in the following case:

**Corollary 6.1.3.** Make the same assumptions as in Theorem 6.1.2 except that  $\mathfrak{g}'_X$  need not have transversal generators. Assume instead that

$$T_X(I) \subseteq \alpha(\mathfrak{g}'_X)$$

(I is the ideal of S). Then the conclusion of Theorem 6.1.2 holds.

*Proof.* By Theorem 6.1.2 we first conclude that M is smooth along the sub-algebroid that is generated by all transversal vectors. Then the assertion follows from [Käl98, Th.4.1] (the proof is by eliminating the transitive directions in  $\mathfrak{g}_X$ ).  $\Box$ 

Remark 6.1.4. Let  $S \subset X$  be a divisor in a complex manifold and M a  $T_X$ module which is coherent as  $\mathcal{O}_X(*S)$ -module; let I be the ideal of S. By [Del70, Th. 4.1] M is smooth along  $T_X(I)$  if for any curve  $\phi : C \to X$  which is transversal to S the inverse image  $\phi^!(M)$  is smooth along  $T_C(I_{\phi^{-1}(S)})$ . The proof in *loc. cit.* is based on complex analysis in two ways: (i) The connection in  $X \setminus S$  is determined by its local system of horizontal sections; (ii) One proofs that S is a countable union of certain *closed* subsets, so one of these will have an interior point by Baire's theorem. Our use of category argument below is more complicated. 6.2. **Proof of Theorem 6.1.2.** As in Section 4.2 we put  $\tilde{M} = \hat{A} \otimes_A M$  for an *A*-module *M*, where  $\hat{A}$  is the **m**-adic completion of a local ring  $(A, \mathbf{m})$ . The proof of Theorem 6.1.2 is based on the following lemma. We use the notation in Theorem 6.1.2.

**Lemma 6.2.1.** Let  $\Omega \subseteq X$  be an open subset such that  $S_0 = S \cap \Omega$  is a nonempty open subset of S defined by the zero locus of a function  $h \in \mathcal{O}_X(\Omega)$ , and let  $\delta \in \mathfrak{g}'_X(\Omega)$ . Assume that for every point  $x \in S_0$  that there exists an ideal  $\hat{I}(x) \subset \hat{\mathcal{O}}_x$  such that:

- (1)  $h \notin \hat{I}(x);$
- (2)  $\delta_x \in \mathfrak{g}'_x(\hat{I}(x));$
- (3)  $\tilde{M}_x/\hat{I}(x)\tilde{M}_x$  is smooth along  $\delta_x$  in  $\mathfrak{g}'_x(\hat{I}(x))$  (see 4.1.2).

Then there exists a (closed) point  $y \in S$  such that  $M_y$  is smooth along  $\delta_y$ in  $\mathfrak{g}'_y$ .

As a first preparation for the proof of Lemma 6.2.1 we recall how one may reduce certain local questions about analytic sheaves to questions about modules over noetherian rings. Let X be a complex analytic space, put  $n = \dim X$ , and let x be a point in X. Then there exists a neighbourhood X' of x and a finite morphism  $f: X' \to \mathbb{C}^n$ . Letting  $K_0 \subset \mathbb{C}^n$  be a closed polydisc such that  $x \in K := f^{-1}(K_0)$ , then the ring  $\mathcal{O}_{\mathbb{C}^n}(K_0)$  of germs of holomorphic functions in neighbourhoods of  $K_0$  is noetherian [Fri67]; hence  $A := \mathcal{O}_X(K)$  is a noetherian ring since it is of finite type over  $\mathcal{O}_{\mathbb{C}^n}(K_0)$ [Gra60]. Moreover, K is a compact Stein set [GR79, Ch IV, §1, Th. 3], hence by Cartan's theorems A and B the functor

$$\Gamma : \operatorname{coh}(\mathcal{O}_K) \to A\operatorname{-mod}_f, \quad M \mapsto M(K)$$

is an equivalence between the category of coherent  $\mathcal{O}_X$ -modules on K and the category of A-modules of finite type, with quasi-inverse  $M(K) \mapsto \mathcal{O}_K \otimes_A M(K)$ ; in particular, the stalk  $M_x \cong \mathcal{O}_x \otimes_A M(K)$ .

Set  $A = \mathcal{O}_X(K)$ , where K is either an affine variety or a compact Stein set  $K \subset X$  such that A is noetherian (as above),  $X_0 := \text{mSpec } A$ , and X := Spec A. In the complex analytic case  $X_0 = K$ . The Nullstellensatz for either category implies that X is an Jacobson scheme (for this notion, see [GD71, Ch. 0, §2]), i.e. the canonical inclusion  $i : X_0 \to X$  is a quasihomeomorphism, meaning that the mapping  $i^{-1}$  defines a bijection between the set of open subsets of X and the set of open subsets of  $X_0$ , and letting U be an open subset of X, there is an isomorphism  $i^{\#} : \mathcal{O}_X(U) \to \mathcal{O}_{X_0}(i^{-1}(U))$ with the k-algebra of regular algebraic functions on  $i^{-1}(U) \subset X_0$ , defined by  $\phi \mapsto i^{\#}(\phi) : x \mapsto \phi \mod \mathfrak{m}_x = k$  (k is algebraically closed). In particular, if  $S_0 \subset X_0$  is dense then  $\cap_{x \in S_0} \mathfrak{m}_x = 0$ .

**Lemma 6.2.2.** Let  $\mathfrak{p} = (h) \subset \operatorname{Spec} A$  be a principal prime ideal, and S be a dense subset of  $\operatorname{mSpec} A/\mathfrak{p}$ . For each  $\mathfrak{m}_x \in S$  choose an ideal  $I(x) \in \operatorname{Spec} A$  such that  $I(x) + \mathfrak{p} \subset \mathfrak{m}_x$  and  $\mathfrak{p} \not\subseteq I(x)$ . Then if G is an A-module of finite type, we have

$$\mathfrak{p} \notin \operatorname{supp}(\cap_{x \in S} I(x)G).$$

*Proof.* The A-module G is noetherian so there exists an element  $f \in A$  such that  $G_f \cong A_f^r$  for some integer r, or  $G_f = 0$ . Let  $\overline{G}$  and  $G^{(f)}$  be the

image and kernel of the canonical mapping  $G \to G_f$ . Assume that  $\overline{G} \neq 0$ . Since  $h \notin I(x)$  the set  $S_1 = \{x \in \operatorname{mSpec} A_f : \cap I(x)_f \subset \mathfrak{m}_x\}$  is dense in the Jacobson scheme  $\operatorname{Spec} A_f$ , so

$$\bigcap_{x \in S} (I(x)\overline{G}) \subset \bigcap_{x \in S} (I(x)G_f) \cong \bigoplus (\bigcap_{x \in S} I(x))_f = 0;$$

hence

$$\cap_{x \in S}(I(x)G) \subset G^{(f)}$$

Since any non-zero  $\phi \in I(x)$  acts by an injective mapping on  $G \subset A_f^r$  it follows that

$$\bigcap_{x \in S} (I(x)G) = \bigcap_{x \in S} (I(x)G^{(f)}).$$

One may therefore assume that  $G = G^{(f)}$ . If  $f \notin \mathfrak{p}$ , then

$$(\cap_{x\in S}(I(x)G^{(f)}))_{\mathfrak{p}}\subset G_{\mathfrak{p}}^{(f)}=0,$$

so  $\mathfrak{p} \notin \operatorname{supp} \cap_{x \in S}(I(x)G)$ . It remains to consider the case  $f \in (h)$ . As  $G = G^{(f)}$  there exists an element  $g \notin \mathfrak{p} = (h)$  such that  $G_g = G_g^{(h)}$ , and since  $G_g$  is a noetherian  $A_g$ -module it has a finite filtration with successive quotients of the form  $A_g/(h)$ . Now  $\mathfrak{p} + I(x) \subset \mathfrak{m}_x \in \operatorname{mSpec} A/(h)$ , and since X is an Jacobson scheme, implying

$$\bigcap_{x\in S}(I+\mathfrak{p})\subseteq \bigcap_{\mathfrak{p}\subset\mathfrak{m}}\mathfrak{m}=\mathfrak{p},$$

one gets  $\bigcap_{x \in S} (I(x) \frac{A_g}{(h)}) = 0$ . Let now

$$0 \to \frac{A_g}{(h)} \to N \to \frac{A_g}{(h)} \to 0$$

be a short exact sequence of  $A_g$ -modules. As  $h \notin I(x)$  any non-zero element  $\phi \in I(x)$  acts as an injective mapping on  $A_g/(h)$  (the right copy in the above exact sequence), implying

$$\bigcap_{x \in S} (I(x)N) = \frac{A_g}{(h)} \cap \bigcap_{x \in S} (I(x)N) = \bigcap_{x \in S} (I(x)\frac{A_g}{(h)}) = 0.$$

An induction over the length of the filtration of  $G_g$  shows that

$$\bigcap_{x \in S} (I(x)G_g) = 0,$$

and since  $G_{\mathfrak{p}}^{(g)} = 0$  when  $g \notin \mathfrak{p}$  it follows that  $(\cap_{x \in S}(I(x)G))_{\mathfrak{p}} = 0$ .  $\Box$ 

**Lemma 6.2.3.** Let  $S \subset V(\mathfrak{p})$  be a dense subset. For each  $x \in S$  choose an ideal  $\hat{I}(x) \subset \hat{A}_x$  such that  $\hat{I}(x) + \hat{\mathfrak{p}} \subset \hat{\mathfrak{m}}_x$  and  $\mathfrak{p} \not\subseteq \hat{I}(x)$ . Let K and N be submodules of finite type in an A-module of finite type F. Then if  $K \subset \hat{N} + \hat{I}(x)\hat{F}$  when  $x \in S$ , we have

$$K_{\mathfrak{p}} \subseteq N_{\mathfrak{p}}.$$

*Proof.* Letting  $\bar{K}$  be the image of K in G := F/N, we have to prove that  $\bar{K}_{\mathfrak{p}} = 0$ .

a) There exists an element  $g \in A$  with the following properties:

(1)  $g \notin \mathfrak{p} = (h);$ 

(2) For any point  $x \in S \setminus g^{-1}(0)$  there exists an ideal  $I(x) \subset A$  such that  $h \notin I(x), I(x) + (h) = \mathfrak{m}_x$ , and  $G_g \cap (\hat{I}(x)G_g) \subseteq I(x)G_g$ .

The proof is similar to the proof of the previous lemma and we use a similar notation. There exists  $f \in A$  such that  $G_f$  is free over  $A_f$ . Setting  $I(x)_f = A_f \cap \hat{I}(x)_f \subset A_f$ , then  $G_f \cap \hat{I}(x)G_f = I(x)_fG_f$ . If  $f \notin (h)$  we are done letting g = f, so now suppose that  $f \in (h)$ . Then there exists  $g \in A \setminus (h)$  such that  $G_g^{(f)} = \operatorname{Ker}(G_g \to G_{fg})$  is a successive extension of modules of the type A/h. Now choose any ideal  $I(x) \subset A$  such that  $\hat{I}(x) \cap A \subset I(x)$ ,  $I(x) + (h) = \mathfrak{m}_x$ , and  $h \notin I(x)$ . Then

$$\begin{split} G_g^{(f)} &\cap \hat{I}(x) G_g^{(f)} \subset I(x) G_g^{(f)}, \\ G_{fg} &\cap \hat{I}(x) G_{fg} \subset I(x) G_{fg} \end{split}$$

implying  $(\hat{I}(x)G_g) \cap G_g \subset I(x)G_g$ .

b) We have by Artin-Rees' lemma  $G = F/N \subset \hat{F}/\hat{N}$ . Since  $\bar{K} \subseteq \hat{I}(x)\hat{F}/\hat{N}$ , by a) there exists  $g \in A$  satisfying (1-2) above, hence

 $\bar{K}_g \subset G_g \bigcap \hat{I}(x) G \subset I(x) G, \quad x \in S,$ 

hence  $\bar{K}_g \subset \bigcap_{x \in S} I(x) G_g$ . Since  $\bar{K}_p = (\bar{K}_g)_p$ , the proof can be finished with Lemma 6.2.2.  $\Box$ 

Proof of Lemma 6.2.1. Clearly one can assume that  $\Omega = X$  and that there exists an  $\mathcal{O}_X$ -coherent submodule  $M_X^0 \subset M_X$  such that  $M_X = \mathcal{D}(\mathfrak{g}_X)M_X^0$ . We need to prove that there exists a (closed) point  $y \in S$  and an integer n such that  $\delta^n M_y^0 \subseteq \mathcal{D}^{n-1}(\mathfrak{g}'_y)M_y^0$ .

a) S is not a countable union of nowhere dense subsets: When X is a variety the normalization theorem shows that we can reduce to either S being discrete or affine k-space and then the assertion is obvious as the field k is uncountable. When S is a complex analytic space it is a locally compact Hausdorff space, so the statement follows from Baire's theorem.

b) Let x be a point in S. By Artin-Rees lemma  $\mathcal{O}_x$  is faithfully flat over  $\mathcal{O}_x$ , so

$$\mathcal{D}^{n_x-1}(\hat{\mathfrak{g}}'_x)\hat{M}^0_x = \hat{\mathcal{O}}_x \otimes_{\mathcal{O}_x} \mathcal{D}^{n_x-1}(\mathfrak{g}'_x)M^0_x, \\ \hat{I}(x)\hat{M}^{m_x}_x = \hat{I}(x) \otimes_{\mathcal{O}_x} M^{m_x}_x,$$

and by (3) there exist positive integers  $n_x, m_x$  such that

$$\delta^{n_x} M_x \subset \mathcal{D}^{n_x - 1}(\hat{\mathfrak{g}}'_x) \hat{M}^0_x + \hat{I}(x) \hat{M}^{m_x}_x$$

Setting

$$S_{n,m} = \{ x \in S : \delta^n M_x \subset \mathcal{D}^{n-1}(\hat{\mathfrak{g}}'_x) M_x^0 + \hat{I}(x) \hat{M}_x^m \}$$

we thus have  $S = \bigcup_{n,m} S_{n,m}$ . By a) there exist integers  $n_0, m_0$  such that the closure of  $S_0 := S_{n_0,m_0}$  contains an interior point. Shrinking X if necessary we can therefore assume that X is either a compact Stein set or affine, so that  $A = \mathcal{O}_X(X)$  is noetherian, and that  $S_0$  is dense in S.

Put  $M^0 = M^0(X)$ ,  $K = \delta^{n_0} M^0$ ,  $N = \mathcal{D}^{n_0-1}(\mathfrak{g}')M^0$  and  $F = \mathcal{D}^{m_0}(\mathfrak{g})M^0(X)$ ; these are A-modules of finite type where N and F generate the coherent  $\mathcal{O}_X$ -modules  $\mathcal{D}^{n_0-1}(\mathfrak{g}')M^0_X$  and  $M^{m_0}_X$ . Since  $A_x$  is faithfully flat over A (Artin-Rees) we have

$$K \subset \hat{A}_x \otimes_A N + \hat{I}(x) \otimes_A F, \quad x \in S_0,$$

so by Lemma 6.2.3  $K_{\mathfrak{p}} \subset N_{\mathfrak{p}}$ . By coherence this inclusion holds in a neighbourhood of  $\mathfrak{p} \in \operatorname{Spec} A$ , and since the closed points are dense in  $S = V(\mathfrak{p})$  there exists a closed point  $y \in S$  such that  $K_y \subset N_y$ .  $\Box$ 

Proof of Theorem 6.1.2.  $(1) \Rightarrow (3), (1) \Rightarrow (4)$ : This follows from the nonderived version of Proposition 4.2.3; it is not necessary that  $\mathfrak{g}_X$  be locally free.  $(3) \Rightarrow (2)$ : o.k. It remains to prove  $(2) \Rightarrow (1)$  and  $(4) \Rightarrow (1)$ : When dim X = 1 it follows that  $X = X_0$ , hence it is a non-singular curve, so we can clearly assume that dim  $X \ge 2$ .

a) Since the assertion is local in X it will be clear that all locally defined sections can be assumed to be globally defined on X. One can clearly assume that S and hence  $S_0$  are connected. As M is torsion free it suffices to prove that  $M_x$  is smooth along  $\mathfrak{g}'_x$  at one point x in  $S_0$  (Cor. 5.2.4), and by Theorem 6.1.1 it suffices to prove that  $M_x$  is smooth along each element in a set of generators of  $\mathfrak{g}'_x$ . One can thus, by shrinking X if necessary, assume that  $S = S_0$  and that the ideal of S is a principal prime ideal  $\mathfrak{p} \subset \mathcal{O}_X$ generated by a function  $h \in \mathcal{O}_X(X)$ . Let  $(\hat{h})_y \subset \hat{\mathcal{O}}_y$  denote the principal ideal of  $h \in \mathcal{O}_y$  in the completion  $\hat{\mathcal{O}}_y$  at a point  $y \in X$ .

b) The assumption is that  $\mathfrak{g}'_X$  has globally defined generators

$$\delta(1), \delta(2), \ldots, \delta(k),$$

which in the case (4) satisfy the condition (G). By removing subsets of  $\operatorname{codim} \geq 2$  from X (Cor. 5.2.2) we on the one hand ensure that  $X = X_0$ , so X is non-singular, and on the other ensure that  $\alpha(\delta(i)) \in T_X$  is of the form  $g_i \partial_i$  where  $\partial_{iy} \notin \mathfrak{m}_y T_y$  for every point  $y \in X$  (it is non-critical). Moreover we have assumed that we can choose the  $\delta(i)$  such that  $\alpha(\delta(i)_y)(h) \notin (\hat{h})_y$ . Therefore, by a formal version of Frobenius' theorem, each generator  $\delta(i)$  has for each point  $y \in S$  an analytic integral curve passing through y, defined by an ideal  $\hat{I}^{(i)}(y) \subset \hat{\mathcal{O}}_y$ , satisfying:

$$\begin{aligned} \alpha(\delta(i)_y)(\hat{I}^{(i)}(y)) &\subseteq \hat{I}^{(i)}(y); \\ h \notin \hat{I}^{(i)}(y); \\ \hat{I}^{(i)}(y) + (\hat{h})_y &\subset \hat{\mathfrak{m}}_y. \end{aligned}$$

Therefore (2) or (4) implies, by Lemma 6.2.1, that there exists a point  $x_i \in S$  such that  $M_{x_i}$  is smooth along  $\delta(i)_{x_i}$  in  $\mathfrak{g}'_{x_i}$ .

c) There exists an open neighbourhood  $U_1$  of  $x_1$  such that for every point  $y \in U_1$  the stalk  $M_y$  is smooth along  $\delta(1)_y$  in  $\mathfrak{g}'_y$  (Prop. 4.1.5). By b), with  $X = U_1$ , there exist a point  $x_2 \in U_1 \cap S_0$  such that  $M_{x_2}$  is smooth along  $\delta(2)_{x_j}$  in  $\mathfrak{g}'_{x_2}$ . Again there exists open neighbourhood  $U_2 \subset U_1$  of  $x_2$  such that  $y \in U_2$  then  $M_y$  is smooth along  $\delta(2)_y$  in  $\mathfrak{g}'_y$ . Iterating we get an open subset  $U = U_k \subseteq U_{k-1} \subseteq \cdots \subseteq U_1$  such that  $M_x$  is smooth along each generator  $\delta(i)_x \in \mathfrak{g}'_x$  when  $x \in U \cap S_0$ .  $\Box$ 

6.3. **Regular singularities.** Unless otherwise stated we assume that X is a complex analytic or algebraic manifold and  $\mathfrak{g}_X$  is a locally free transitive

Lie algebroid. When  $\pi: Y \to X$  is a morphism of spaces we tacitly assume that  $\mathfrak{g}_Y$  denotes  $\pi^+(\mathfrak{g}_X)$ .

First let X be any space,  $\mathfrak{g}_X \in \operatorname{Lie}_X$ ,  $M \in \operatorname{coh}(\mathfrak{g}_X)$  be pure with support V, and let J be the ideal of V. Let  $S \subset X$  be a subspace whose ideal is I. We say that a pure  $\mathfrak{g}_X$ -module M has regular singularities along S if M is point-wise smooth along the sub-algebroid

$$\mathfrak{g}_X(J) \cap \mathfrak{g}_X(I) \subset \mathfrak{g}_X.$$

ļ

Consider a decomposition

$$S = D \cup D'$$

where D is the union of the irreducible components  $S_i \subseteq S$  such that  $S_i \cap V$  has pure codimension 1 in V, and D' is the union of the irreducible components  $S_j$  such that  $S_j \cap V$  has codimension  $\geq 2$  in V; let  $I_1$  be the ideal of D and  $I_2$  the ideal of D'. Now since  $\mathfrak{g}_X(I)_{|X\setminus D'} = \mathfrak{g}_X(I_1)_{|X\setminus D'}$  it follows by Corollary 5.2.3 that M is smooth along  $\mathfrak{g}_X(I_1) \cap \mathfrak{g}_X(J)$  if and only if M is smooth along  $\mathfrak{g}_X(I) \cap \mathfrak{g}_X(J)$ . We may therefore in practice assume that S = D when checking if M has regular singularities along S.

If S is a subspace of a space S' it may not be obvious that M has regular singularities along S' when it has regular singularities along S. That this is so follows from the following lemma, which also implies that M has regular singularities along S if it has regular singularities along each component  $S_i$ in an irreducible decomposition  $S = \bigcup S_i$ .

**Lemma 6.3.1.** (X is a space) Let  $I \subset \mathcal{O}_X$  be a proper ideal and  $\sqrt{I}$  its radical. Then:

(1) If  $I = I_1 \cap I_2 \cdots \cap I_r$  is a minimal primary decomposition we have  $T_X(I) = T_X(I_1) \cap T_X(I_2) \cap \cdots \cap T_X(I_r);$ (2)  $T_X(I) \subset T_X(\sqrt{I}).$ 

Remark 6.3.2. One also has  $T_X(\sqrt{I}) = T_X((\sqrt{I})^n)$  for any positive integer n, but in general  $T_X(\sqrt{I}) \notin T_X(I)$ . For example, let  $I = (x+y, y^2) \subset k[x,y]$ , then  $T(\sqrt{I}) = T((x,y)) = k[x,y]y\partial_y + k[x,y]y\partial_y$ , but  $x\partial_x(x+y) = x \notin I$ .

*Proof.* (1): It is evident that  $T_X(I_1) \cap \cdots \cap T_X(I_r) \subset T_X(I)$ . Now let  $\partial \in T_X(I)$  and  $x_1 \in I_1$ . Since we are considering a minimal primary decomposition there exists an element  $y \in I_2 \cap \cdots \setminus I_r \setminus \sqrt{I_1}$ . Then

$$y\partial(x) = \partial(yx) - x\partial(y) \in I_1$$

and since  $I_1$  is  $\sqrt{I_1}$ -primary,  $\partial(x) \in I_1$ . In the same way  $\partial(I_i) \subset I_i$ ,  $i = 1, \ldots, r$ , implying the assertion.

(2): First assume that  $P = \sqrt{I}$  is prime and let  $\partial \in T(I)$ ,  $x \in P$ , and  $x^n \in I$ . If n = 1 we have  $\partial(x) \in I \subset P$  so assume that  $x \notin I$ . Then  $nx^{n-1}\partial(x) \in I$ . Supposing that  $\partial(x) \notin P$ , since I is P-primary we get  $nx^{n-1} \in I$ , and iterating gives  $n!x \in I$ . Since the characteristic is 0 we reach a contradiction. Thus  $\partial(x) \in P$ . This shows that  $T(I) \subseteq T(P)$ . Hence  $T_X(I_i) \subset T_X(\sqrt{I_i})$ . By (1)

$$T_X(I) = T_X(I_1) \cap \dots \cap T_X(I_r)$$
$$T_X(\sqrt{I}) = T_X(\sqrt{I_1}) \cap \dots \cap T_X(\sqrt{I_r});$$

hence  $T_X(I) \subset T_X(\sqrt{I})$  for any ideal I.  $\Box$ 

Now let M be any coherent  $\mathfrak{g}_X$ -module and

$$(6.3.1) 0 = M_n \subset M_{n-1} \subset \cdots \subset M_0 = M$$

be its Krull filtration (see Section 5.1), where successive quotients  $L_i = M_i/M_{i+1}$  are pure.

**Definition 6.3.3.** A coherent  $\mathfrak{g}_X$ -module M has regular singularities if each pure component  $L_i$  has regular singularities as above. Let  $\operatorname{coh}_{rs}(\mathfrak{g}_X)$  be the category of such modules.

- Remarks 6.3.4. (1) If  $M \in \operatorname{coh}(\mathfrak{g}_X)$  and  $V = \operatorname{supp} M$  is discrete, then  $M \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ . To see this, let  $x \in \operatorname{supp} M$  and  $M_x^0 \subset M_x$  be an  $\mathcal{O}_x$ -submodule of finite type. Then  $\mathfrak{m}_x^n M_x^0 = 0$  for some big integer n. If  $\phi \in \mathfrak{m}_x$ ,  $\mu \in M_x^0$ , and  $\delta \in \mathfrak{g}(\mathfrak{m}_x)$ , then since  $\alpha(\delta)(\phi) \in \mathfrak{m}_x$ , so  $\phi^n \delta \mu = 0$ , it follows that  $\mathcal{D}(\mathfrak{g}(\mathfrak{m}_x)) M_x^0$  is of finite type over  $\mathcal{O}_x$ . Thus  $M_x$  is smooth along  $\mathfrak{g}_x(\mathfrak{m}_x)$ .
  - (2) By Proposition 6.3.7 below a torsion free coherent  $T_X$ -module M on a space X has regular singularities if and only if M has regular singularities in the sense of [Del70, Th 4.1].

**Proposition 6.3.5.** Consider a short exact sequence of coherent  $\mathfrak{g}_X$ -modules

$$0 \to K \to M \to N \to 0.$$

Then M has regular singularities if and only if K and N have regular singularities.

*Proof.* Consider the Krull filtration in (6.3.1).

a) If  $M \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$  then  $K, N \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ : We have a short exact sequence

(6.3.2) 
$$0 \to \frac{K \cap M_i}{K \cap M_{i+1}} \to \frac{M_i}{M_{i+1}} \to \frac{M_i}{\bar{M}_{i+1}} \to 0$$

where  $\overline{M}_i$  is the image of  $M_i$  in N. Here  $K \cap M_i/K \cap M_{i+1}$  is 0 or a maximal pure sub-quotient of K, and all maximal pure sub-quotients of K occur in this way, but  $\overline{M}_i/\overline{M}_{i+1}$  need not be pure. Now if  $J_i \subset I_i$  are radical ideals such that  $M_i/M_{i+1}$  is smooth along  $\mathfrak{g}_X^i = \mathfrak{g}_X(J_i) \cap \mathfrak{g}_X(I_i)$ ,  $i = 0, \ldots, m$ , then  $K \cap M_i/K \cap M_{i+1}$  and  $\overline{M}_i/\overline{M}_{i+1}$  also are smooth along  $\mathfrak{g}_X^i$  (Prop. 4.1.4). Hence  $K \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ , and every pure subquotient of  $\overline{M}_i/\overline{M}_{i+1}$  is smooth along  $\mathfrak{g}_X^i$  (Prop. 4.1.4), so  $N \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ .

b) If  $K, N \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$  then  $M \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ : Consider the short exact sequence (6.3.2). As  $\overline{M}_i \subset N$  and  $M_i \cap K \subset K$ , by a)  $\overline{M}_i, M_i \cap K \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ , and again by a)  $(K \cap M_i)/(K \cap M_{i+1}), \overline{M}_i/\overline{M}_{i+1} \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ . It therefore suffices to prove that  $M \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$  when M is pure. Then K is also pure, and by Corollary 5.2.3 we can assume also that N is pure. Put  $V_K = \operatorname{supp} K, V_M = \operatorname{supp} M$  and  $V_N = \operatorname{supp} N$ , and let  $J_K, J_M, J_N \subset \mathcal{O}_X$ be their ideals. Let  $I_K, I_N \subset \mathcal{O}_X$  be ideals such that K is smooth along  $\mathfrak{g}_X(J_K) \cap \mathfrak{g}_X(I_K)$  and N is smooth along  $\mathfrak{g}_X(J_N) \cap \mathfrak{g}_X(I_N)$ ; let S and T be the spaces of  $I_K$  and  $I_N$ . As M is pure we have dim  $V_K = \dim V_M$ , and we can assume that dim  $V_M - 1 \leq \dim V_N \leq \dim V_M$  (Cor. 5.2.3). So we have two cases. Suppose first that dim  $V_N = \dim V_M - 1$ . Then  $\operatorname{codim}_{V_M} T \ge 2$ , hence we may then assume that N is smooth along  $\mathfrak{g}_X(J_N)$  (Cor. 5.2.3). Moreover, Corollary 5.2.3 also implies that one may assume that all irreducible components of S and  $V_N$  have the same dimension. Letting  $I_M$  be the ideal of  $S \cup V_N$  we claim that M is smooth along

$$\mathfrak{g}'_X := \mathfrak{g}_X(J_M) \cap \mathfrak{g}_X(I_M),$$

so  $M \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ . By Proposition 4.1.4 this follows if N and K are smooth along  $\mathfrak{g}'_X$ . We have assumed that all irreducible components of S and  $V_N$ have the same dimension; hence the prime components of  $I_K \cap J_N$  is the union of the prime components of  $I_K$  and  $J_N$ ; hence Lemma 6.3.1 implies

$$\mathfrak{g}'_X \subset \mathfrak{g}_X(I_M) = \mathfrak{g}_X(I_K \cap J_N) \subset \mathfrak{g}_X(J_N).$$

Thus N is smooth along  $\mathfrak{g}'_X$ . Since  $V_K \subset V_M$  and  $\dim V_K = \dim V_M$  we may assume that the prime components of  $J_K$  is a part of the prime components of  $J_M$  (Cor. 5.2.3) so  $\mathfrak{g}_X(J_M) \subset \mathfrak{g}_X(J_K)$  (Lem. 6.3.1); in the same way we get  $\mathfrak{g}_X(I_M) \subset \mathfrak{g}_X(I_K)$ . Hence

$$\mathfrak{g}'_X \subseteq \mathfrak{g}_X(J_K) \cap \mathfrak{g}_X(I_K \cap J_N) \subseteq \mathfrak{g}_X(J_K) \cap \mathfrak{g}_X(I_K);$$

and therefore K is smooth along  $\mathfrak{g}'_X$ .

Now suppose that dim  $V_N = \dim V_M$ . Then we may assume that S and T have pure codimension 1 in  $V_M$  (Cor. 5.2.3); let  $I_M$  be the ideal of  $S \cup T$ . Using a very similar argument as above, based on Lemma 6.3.1, one sees that K and N are smooth along  $\mathfrak{g}_X(J_M) \cap \mathfrak{g}_X(I_M)$ ; hence M is smooth along  $\mathfrak{g}_X(J_M) \cap \mathfrak{g}_X(I_M)$  (Prop. 4.1.4).  $\Box$ 

The subcategories of  $D^b_{\rm coh}(\mathfrak{g}_X)$  that are of concern in this section are denoted  $D^b_{rs}(\mathfrak{g}_X)$ ,  $D^b_{crs}(\mathfrak{g}_X)$ , and  $D^b_{hol}(\mathfrak{g}_X)$ . They are defined as follows:  $D^b_{rs}(\mathfrak{g}_X)$  is the derived category of bounded complexes  $M^{\bullet}$  of  $\mathfrak{g}_X$ -modules whose homology  $H^{\bullet}(M^{\bullet}) \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ . It follows from Proposition 6.3.5 that  $D^b_{rs}(\mathfrak{g}_X)$  is a triangulated sub-category of  $D^b_{\mathrm{coh}}(\mathfrak{g}_X)$ ; i.e. if two vertices of a distinguished triangle in  $D^b_{\rm coh}(\mathfrak{g}_X)$  belong to  $D^b_{rs}(\mathfrak{g}_X)$ , then all three vertices belong to  $D^b_{rs}(\mathfrak{g}_X)$ .  $D^b_{crs}(\mathfrak{g}_X)$  is the sub-category of complexes  $M^{\bullet}$ such that for any curve  $\pi: C \to X$  one has  $\pi^!(M^{\bullet}) \in D^b_{rs}(\mathfrak{g}_C)$ ; we then say that  $M^{\bullet}$  is curve regular. Since  $D^b_{rs}(\mathfrak{g}_C)$  is a triangulated category and  $\pi^{!}$  preserves triangles, it follows that  $D^{b}_{crs}(\mathfrak{g}_{X})$  is a triangulated category. A coherent  $\mathfrak{g}_X$ -module M is holonomic when dim  $SS(M) = \dim X$  (2.3).  $D^b_{\text{hol}}(\mathfrak{g}_X)$  is the derived category of bounded complexes with holonomic homology modules; it is easy to see that sub-quotients of holonomic modules are holonomic, implying that  $D^b_{hol}(\mathfrak{g}_X)$  is a triangulated category. We refer to [BGK<sup>+</sup>87] and [Bjö93] for the theory of holonomic  $T_X$ -modules on a non-singular space X.

Remark 6.3.6. Let  $\phi : (Y, \mathfrak{g}_Y) \to (X, \mathfrak{g}_X)$  be a morphism of Lie algebroids on non-singular spaces, where  $\mathfrak{g}_Y = \phi^+(\mathfrak{g}_X)$ . Letting  $I \subset \mathcal{O}_X$  be the ideal of a divisor S on X and  $J \subset \mathcal{O}_Y$  the ideal of  $\pi^{-1}(S)$  we have a canonical morphism

$$\phi^+(\mathfrak{g}_X(I)) \to \mathfrak{g}_Y(J),$$

but this need not be surjective when S is singular and  $d\phi: T_Y \to \phi^*(T_X)$  is not surjective. For instance, let  $I = (x_1^3 + x_2^3 + x_3^3) \subset A := \mathbb{C}\{x_1, x_2, x_3\}$  be the ideal of the germ of the cubic cone in  $(\mathbf{C}^3, 0)$ . Then the A-module of derivations that preserve I is  $T_{(\mathbf{C}^3,0)}(I) = AE$  where  $E = x_1\partial_{x_1} + x_2\partial_{x_2} + x_3\partial_{x_3}$  [BGG72]. Therefore, if  $Y \subset (\mathbf{C}^3, 0)$  is a germ of an irreducible curve in  $(\mathbf{C}^3, 0)$  with ideal  $L \subset A$  such that  $E \notin T_{\mathbf{C}^3,0}(L)$ , then one easily sees that  $T_{\mathbf{C}^3,0}(I) \cap T_{\mathbf{C}^3,0}(L) = LE$ . Hence, if  $\phi : Y \to (\mathbf{C}^3, 0)$ is the inclusion morphism then the induced homomorphism  $\phi^+(T_{\mathbf{C}^3,0}(I)) = \phi^*(T_{\mathbf{C}^3,0}(I)) \times_{\phi^*(T_{\mathbf{C}^3,0})} T_Y \to T_Y$  is 0.

When X is non-singular we will generalize the proposition below considerably in Theorem 6.3.15.

**Proposition 6.3.7.** (X is a space of the same type as in Theorem 6.1.2) Let M be a holonomic  $\mathcal{D}_X$ -module without  $\mathcal{O}_X$ -torsion and I be the ideal of its strong support (see Lemma 2.3.6). The following are equivalent:

- (1)  $\phi^!(M)$  is regular along  $\phi^+(T_X(I))$  when  $\phi: C \to X$  is a curve (when X is singular we only consider the non-derived inverse image);
- (2) M is smooth along  $T_X(I)$ .

*Remark* 6.3.8. [BGK<sup>+</sup>87, VII, Props. 11.6-7 and Cor. 11.8] are attained by employing GAGA, but they also follow from Proposition 6.3.7 and Theorem 6.1.2 without GAGA.

*Proof.* (2)  $\Rightarrow$  (1): Let  $\phi : C \to X$  be a curve. By Theorem 6.1.2 we know that the non-derived inverse image  $\phi^{!}(M) \in \operatorname{coh}_{s}(T_{C})$ . In the non-singular case the assertion follows from Proposition 4.2.3, but we will see below that we actually have  $\phi^{!}(M) \in D^{b}_{rs}(T_{C})$  (Th. 6.3.12).

(1)  $\Rightarrow$  (2): Since M is holonomic, SS  $M \subset T^*X$  is a conic algebraic (analytic) Lagrangian subset; hence, M being torsion free, S := s-supp  $M \subset X$  is a divisor (*Def.* 2.3.5). As the space X is non-singular in codimension 1 we may assume that S and X are non-singular spaces (*Cor.* 5.2.2). Let x be a point in S and  $(t = x_1, x_2, \ldots, x_n)$  be a regular system of parameters of the local ring  $\mathcal{O}_x$  such that t = 0 is a local equation for S at x. Since  $\mathcal{O}_x$  is either isomorphic to the ring of convergent power series in the given parameters, or a localisation of a ring of finite type, there exist derivations  $\partial_t = \partial_{x_1}, \partial_{x_2}, \ldots, \partial_{x_n} \in \text{Der}_k(\mathcal{O}_x)$  such that  $\partial_{x_i}(x_j) = \delta_{ij}$  (the Kronecker delta). As SS M is an algebraic (analytic) Lagrangian set, each irreducible component of the nonsingular locus of SSM (see 2.3) belongs to the conormal set of S; therefore  $M_x$  is smooth along  $\partial_{x_2}, \ldots, \partial_{x_n}$ . Now (1) and Theorem 6.1.2 implies that  $M_x$  is smooth along  $t\partial_t$ ; hence  $M_x$  is point-wise smooth along  $\mathcal{O}_x t\partial_t + \mathcal{O}_x\partial_{x_2} + \cdots + \mathcal{O}_x\partial_{x_n} = T_x(I_S)$ , proving (2).  $\Box$ 

The following result is due to Bernstein.

**Proposition 6.3.9.** (X is a non-singular space) Let  $j : Y \to X$  be an open embedding of spaces such that  $S := X \setminus j(Y)$  is a subspace. Let  $M^{\bullet} \in D^{b}_{\mathrm{coh}}(\mathfrak{g}_{X})$  and consider the distinguished triangle

(6.3.3) 
$$R\Gamma_{[S]}(M^{\bullet}) \to M^{\bullet} \to M^{\bullet}(*S) \to .$$

Then the following are equivalent:

(1)  $M \in D^b_{\text{hol}}(\mathfrak{g}_X);$ 

(2)  $R\Gamma_{[S]}(M^{\bullet})$  and  $M^{\bullet}(*S)$  belong to  $D^{b}_{hol}(\mathfrak{g}_{X})$ .

Using a Mayer-Vietoris argument, the main case in the proof is  $(1) \Rightarrow (2)$  when S is a divisor and this follows from the existence of functional equations containing a Bernstein-Sato polynomial; see e.g. [Bjö93, Th. 3.2.13].

# **Proposition 6.3.10.** $D^b_{rs}(\mathfrak{g}_X) \subset D^b_{hol}(\mathfrak{g}_X).$

Proof.  $D_{rs}^{b}(\mathfrak{g}_{X})$  and  $D_{hol}^{b}(\mathfrak{g}_{X})$  are generated by its homology objects, so it suffices to see that a pure  $\mathfrak{g}_{X}$ -module M with regular singularities is holonomic. Let J be the ideal of the support V of M and  $I \subset \mathcal{O}_{X}$  be an ideal such that M is smooth along  $\mathfrak{g}_{X}(J) \cap \mathfrak{g}_{X}(I)$ . Let S be the space of I. Since  $\mathcal{D}(\mathfrak{g}_{X})$  is noetherian (2.3.2) the union N of all holonomic submodules of M is holonomic. Then clearly  $N|_{X\setminus S} = M|_{X\setminus S}$ , so the quotient K = M/N is a coherent  $\mathfrak{g}_{X}$ -module whose support belongs to  $S \subset V$ ; hence  $H^{0}(K(*S)^{\bullet}) = 0$ , so by the first part of the long exact sequence induced by the distinguished triangle  $N(*S)^{\bullet} \to M(*S)^{\bullet} \to K(*S)^{\bullet} \to$  one gets  $H^{0}(N(*S)) = H^{0}(M(*S))$ ; by Proposition 6.3.9  $H^{0}(N(*S)^{\bullet})$  is holonomic; hence  $M \subset H^{0}(M(*S)^{\bullet})$  is holonomic.  $\Box$ 

We next have a version of Bernstein's theorem for regular holonomic  $\mathcal{D}$ -modules; see [Bjö93, Th. 5.4.1].

**Theorem 6.3.11.** Let  $j : Y \to X$  be an open embedding such that  $S := X \setminus j(Y)$  is a subspace. Let  $M^{\bullet} \in D^b_{coh}(\mathfrak{g}_X)$  and consider the distinguished triangle (6.3.3). Then the following are equivalent:

- (1)  $M^{\bullet} \in D^b_{rs}(\mathfrak{g}_X).$
- (2)  $R\Gamma_{[S]}(M^{\bullet})$  and  $M(*S)^{\bullet}$  belong to  $D^b_{rs}(\mathfrak{g}_X)$ .

*Proof.* (2)  $\Rightarrow$  (1) follows since  $D^b_{rs}(\mathfrak{g}_X)$  is a triangulated category.

To prove the converse one may assume that  $M^{\bullet}$  is a single degree complex, consisting of a pure  $\mathfrak{g}_X$ -module M with regular singularities; let K be the ideal of  $V = \operatorname{supp} M$  and I an ideal such that M is smooth along  $\mathfrak{g}_X(K) \cap \mathfrak{g}_X(I)$ . Letting T be the space of I, by the discussion before Lemma 6.3.1 we can assume that every irreducible component of  $V \cap T$  has pure codimension 1 in V.

By Propositions 6.3.9 and 6.3.10

$$R\Gamma_{[S]}(M)^{\bullet}, M(*S)^{\bullet} \in D^{b}_{\mathrm{coh}}(\mathfrak{g}_{Y})$$

so it remains to prove that  $R\Gamma_{[S]}(M^{\bullet})^{\bullet}$  and  $M(*S)^{\bullet}$  have regular singularities.

a) We first prove that  $M(*S) \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$  when S is a divisor, locally defined by a function  $f \in \mathcal{O}_X$ . So in this case  $M(*S)^{\bullet}$  is a single degree complex given by a coherent  $\mathfrak{g}_X$ -module M(\*S). If  $S \cap V = \emptyset$  then M(\*S) =0, so we assume  $S \cap V \neq \emptyset$ , and then the discussion before Lemma 6.3.1 shows that one may assume that  $S \cap V$  is of pure codimension 1 in V. Then M(\*S) is again a pure  $\mathfrak{g}_X$ -module, and it suffices to see that it is smooth along

$$\mathfrak{g}'_X = \mathfrak{g}_X(K) \cap \mathfrak{g}_X(I \cap J),$$

where J = (f) is the ideal of S. Now there exists (locally) a coherent  $\mathcal{O}_X$ submodule  $M_0 \subset M$  such that  $\mathcal{D}(\mathfrak{g}_X)M_0 = M$  and  $\mathfrak{g}_X(K) \cap \mathfrak{g}_X(I) \cdot M_0 \subset M_0$  and an integer k such that  $\mathcal{D}(\mathfrak{g}_X)M_0f^{-k} = M(*S)$ . It suffices now to see that  $\mathfrak{g}'_X \cdot M_0f^{-k} \subseteq M_0f^{-k}$ . Since the prime components of  $I \cap J$  is the union of the prime components of I and the prime components of J, Lemma 6.3.1 implies

$$\mathfrak{g}'_X = \mathfrak{g}_X(K) \cap \mathfrak{g}_X(I) \cap \mathfrak{g}_X(J) \subset \mathfrak{g}_X(J).$$

Therefore, if  $\delta \in \mathfrak{g}'_X$ , we have

$$\delta \cdot M_0 f^{-k} \subseteq (\delta \cdot M_0) f^{-k} - k\alpha(f) M_0 f^{-k-1}$$
$$\subseteq (\delta \cdot M_0) f^{-k} - kJ M_0 f^{-k-1} \subseteq M_0 f^{-k},$$

implying the assertion.

b) The general case. Let  $(f_1, \ldots, f_l)$  be local generators of the ideal I of S. Letting  $S_i$  be the locally defined divisor that is defined by  $f_i$ , by a)  $\oplus_i M(*S_i) \in D^b_{rs}(\mathfrak{g}_X)$ , and inductively  $M(*\cup_i S_i) \in D^b_{rs}(\mathfrak{g}_X)$ . Therefore  $\oplus_i R\Gamma_{[S_i]}(M) \in D^b_{rs}(\mathfrak{g}_X)$  and  $R\Gamma_{[\cup_i S_i]}(M) \in D^b_{rs}(\mathfrak{g}_X)$  (6.3.3), so using the distinguished triangle of Mayer-Vietoris type

$$R\Gamma_{[S]}(M) \to \oplus R\Gamma_{[S_i]}(M) \to R\Gamma_{[\cup_i S_i]}(M) \to$$

this implies that the third vertex  $R\Gamma_{[S]}(M) \in D^b_{rs}(\mathfrak{g}_X)$ ; hence the third vertex  $M(*S)^{\bullet}$  in the distinguished triangle (6.3.3) also belongs to  $D^b_{rs}(M)$ .  $\Box$ 

**Theorem 6.3.12.** Let  $\pi : Y \to X$  be a morphism of non-singular spaces. Then  $\pi^{!}$  is a functor from  $D^{b}_{rs}(\mathfrak{g}_{X})$  to  $D^{b}_{rs}(\mathfrak{g}_{Y})$ .

Note that Proposition 4.2.3 does not directly imply Theorem 6.3.12, in the light of *Remark* 6.3.6.

Proof. The triangulated category  $D_{rs}^{b}(\mathfrak{g}_{X})$  is generated by  $\operatorname{coh}_{rs}(\mathfrak{g}_{X})$  and  $D_{rs}^{b}(\mathfrak{g}_{Y})$  is stable under extensions, so it suffices to prove that  $\pi^{!}(M) \in D_{rs}^{b}(\mathfrak{g}_{X})$  when  $M \in \operatorname{coh}_{rs}(\mathfrak{g}_{X})$ . Moreover, by induction over the length of the filtration by supports of M one can assume that M is pure. Let I be the ideal of  $\operatorname{supp} M$  and J an ideal such that M is smooth along  $\mathfrak{g}'_{X} := \mathfrak{g}_{X}(J) \cap \mathfrak{g}_{X}(I)$ . Now  $\pi$  can be factorized into a closed embedding and a projection, and by Proposition 3.4.3 it suffices to treat these cases separately.

A projection  $\pi$  is flat, hence

$$\pi^{!}(M) = \pi^{*}(\mathcal{D}(\mathfrak{g}_{X})) \otimes_{\pi^{-1}(\mathcal{D}(\mathfrak{g}_{X}))}^{L} \pi^{-1}(M)$$
$$= \mathcal{O}_{Y} \otimes_{\pi^{-1}(\mathcal{O}_{X})}^{L} \pi^{-1}(M)$$
$$= \mathcal{O}_{Y} \otimes_{\pi^{-1}(\mathcal{O}_{X})} \pi^{-1}(M)$$

and this is a pure coherent  $\mathfrak{g}_Y$ -module. Letting  $I_1, J_1 \subset \mathcal{O}_Y$  be the ideals that are generated by I and J one sees that  $\pi^!(M)$  is smooth along  $\mathfrak{g}'_Y = \pi^+(\mathfrak{g}'_X) = \mathfrak{g}_Y(\pi^*(I)) \cap \mathfrak{g}_Y(\pi^*(J)).$ 

Now assume that  $\pi$  is a closed embedding. By Theorems 3.5.4 and 6.3.11  $\pi_+\pi^!(M) \in D^b_{rs}(\mathfrak{g}_X)$ . So the proof will be complete if one proves: if  $N \in \operatorname{coh}(\mathfrak{g}_Y)$  and  $\pi_+(N) \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ , then  $N \in \operatorname{coh}_{rs}(\mathfrak{g}_Y)$ . To prove this one may again assume that N is pure; then  $\pi_+(N)$  is also pure with support in  $Y \subset X$ . Suppose that  $\pi_+(N)$  is smooth along  $\mathfrak{g}_X(I) \cap \mathfrak{g}_X(J)$  where J is the

ideal of  $V := \operatorname{supp} \pi_+(M)$  and I is the a ideal of a subspace  $S \subset V \subset X$ . Let  $L \subset \mathcal{O}_X$  be the ideal of  $\pi(Y) \subset X$ . By Theorem 3.5.4

$$N = \pi^! \pi_+(N)[\mathrm{d}_{Y,X}] \cong \pi_+(N)^L \subset \pi_+(N),$$

where  $\pi_+(L)^L$  is the  $\mathcal{O}_X$ -submodule of L-invariants of  $\pi_+(N)$ . Letting  $I_1$ and  $J_1$  be the ideals of  $S \subset V \subset Y$  in  $\mathcal{O}_Y = \mathcal{O}_X/L$  we need to prove that Nis smooth along  $\mathfrak{g}_Y(I_1) \cap \mathfrak{g}_Y(J_1)$ , for then  $N \in \operatorname{coh}_{rs}(\mathfrak{g}_Y)$ . Since  $L \subset J \subset I$ and, Y being non-singular,  $T_Y = T_X(L)/LT_X$ , it follows that the canonical mapping  $T_X(I) \cap T_X(J) \cap T_X(L) \to T_Y(I_1) \cap T_Y(J_1)$  is surjective. Hence the canonical mapping

$$\mathfrak{g}_X(I) \cap \mathfrak{g}_X(J) \cap \mathfrak{g}_X(L) \to \mathfrak{g}_Y(I_1) \cap \mathfrak{g}_Y(J_1)$$

is surjective. Now  $N = \pi_+(N)^L$  is a  $\mathfrak{g}_X(L)$ -module which is smooth along the sub-algebroid  $\mathfrak{g}_X(I) \cap \mathfrak{g}_X(J) \cap \mathfrak{g}_X(L)$ . This readily implies that N is smooth along  $\mathfrak{g}_Y(I_1) \cap \mathfrak{g}_Y(J_1)$ .  $\Box$ 

Let  $(\mathfrak{g}_X, \alpha)$  be a transitive locally free Lie algebroid on X; then  $\mathfrak{g}_Y := \pi^+(\mathfrak{g}_X)$  is also locally free and transitive on Y (Lem. 2.4.8).

**Theorem 6.3.13.** Let  $\pi: Y \to X$  be a proper morphism of spaces. Then

$$\pi_+(D^b_{rs}(\mathfrak{g}_Y)) \subset D^b_{rs}(\mathfrak{g}_X).$$

*Proof.* We will use the following criterion for pure modules.

(C): Assume that M is a pure  $\mathfrak{g}_Y$ -module. Let J be the ideal of Vand I the ideal of a subspace  $S \subset V \subset Y$  such that M is smooth along  $\mathfrak{g}'_Y = \mathfrak{g}_Y(J) \cap \mathfrak{g}_Y(I)$ ; then clearly  $\operatorname{Ker}(\mathfrak{g}_Y \to T_Y) \subset \mathfrak{g}'_Y$ . Putting  $\mathfrak{g}_{Y/X} = \alpha^{-1}(T_Y/X)$  we let  $\mathfrak{a}_Y$  be the Lie algebroid that is generated by  $\mathfrak{g}'_Y + \mathfrak{g}_{Y/X}$ . If there exist a subspace  $V_1 \subset V$ , so that  $\operatorname{codim}_V V_1 \ge 2$ , and radical ideals  $J_1, I_1 \subset \mathcal{O}_X$  such that

(6.3.4) 
$$\operatorname{Im}(\pi^*(\mathfrak{g}_X(J_1) \cap \mathfrak{g}_X(I_1))_y \to \pi^*(\mathfrak{g}_X)_y) \subseteq \operatorname{Im}(\mathfrak{a}_y \to \pi^*(\mathfrak{g}_X)_y)$$

when  $y \in V \setminus V_1$ , then  $\pi_+(M) \in D^b_{rs}(\mathfrak{g}_X)$  (Cor. 5.2.6).

We now start the proof. Let  $M^{\bullet} \in D^b_{rs}(\mathfrak{g}_Y)$  and set  $V = \operatorname{supp} M^{\bullet}$ . To prove that  $\pi_+(M^{\bullet}) \in D^b_{rs}(\mathfrak{g}_X)$  we may assume that  $M^{\bullet} = M$  is a single degree complex consisting of a coherent  $\mathfrak{g}_Y$ -module with regular singularities. We factorize  $\pi$  as  $\pi = p \circ i$ , where  $i : Y \to Y \times X$  is the graph embedding and  $p : Y \times X \to X$  is the projection on the second factor.

a)  $i_+(M) \in D^b_{rs}(\mathfrak{g}_{Y \times X})$ : To see this one may assume that M is pure and apply (C), letting  $J_1$  and  $I_1$  be the ideals of i(V) and i(S).

b) Assume that M is torsion free. Let  $S \subset Y$  be a divisor such that M is smooth along  $\mathfrak{g}_Y(I')$ , where I' is the ideal of S. Let T be the critical locus of  $\pi$ , i.e. the locus of points  $y \in Y$  such that  $\pi$  is not submersive; this also is a divisor on Y. Let I be the ideal of  $S \cup T$ . Then M is smooth along  $\mathfrak{g}_Y(I)$ . Letting  $V_1$  be the singular locus of  $S \cup T$  and V = Y we have  $\operatorname{codim}_V V_1 \geq 2$ , and it is straightforward to check that (6.3.4) is satisfied; hence  $\pi_+(M) \in D^b_{rs}(\mathfrak{g}_X)$ .

c) The proof now is by induction over  $n = \dim Y$ . When n = 0 the assertion follows from *Remark* 1, so assume that the theorem holds for all proper morphisms  $\pi_1 : Y_1 \to X_1$  when  $\dim Y_1 < n$ .

d) First assuming that dim  $V \leq n-1$  we prove that  $\pi_+(M) \in D^b_{rs}(\mathfrak{g}_X)$  by induction over dim V. If dim V = 0 the assertion follows from Remark 1, so assume that dim  $V \geq 1$ . Let  $r: V' \to Y \times X$  be a resolution of supp  $i_+(M) = i(V)$ . We then have a distinguished triangle (Th. 3.5.3)

(6.3.5) 
$$r_+r'(i_+(M)) \to i_+(M) \to C^{\bullet} \to$$

and applying  $p_+$ , using  $p_+r_+ = (p \circ r)_+$  and  $\pi_+ = p_+i_+$  (Prop. 3.5.1), we also have the distinguished triangle

(6.3.6) 
$$(p \circ r)_+ r^! (i_+(M)) \to \pi_+(M) \to p_+(C^{\bullet}) \to .$$

By a)  $i_+(M) \in D^b_{rs}(\mathfrak{g}_{Y \times X})$ , hence  $r^!(i_+(M)) \in D^b_{rs}(\mathfrak{g}_{V'})$  (*Th.* 6.3.12). Since  $\dim V' \leq n-1$ , by induction  $(p \circ r)_+ r^!(i_+(M)) \in D^b_{rs}(\mathfrak{g}_X)$  and  $r_+ r^!(i_+(M)) \in D^b_{rs}(\mathfrak{g}_{Y \times X})$ . It will now follow that  $\pi_+(M) \in D^b_{rs}(\mathfrak{g}_X)$  if we prove that  $p_+(C^{\bullet}) \in D^b_{rs}(\mathfrak{g}_X)$ .

First,  $C^{\bullet} \in D^{b}_{rs}(\mathfrak{g}_{Y \times X})$  since  $r_{+}r^{!}(i_{+}(M)), i_{+}(M) \in D^{b}_{rs}(\mathfrak{g}_{Y \times X})$ . Second, *i* is a closed embedding and  $\operatorname{supp} C^{\bullet} \subset i(Y)$  so  $C^{\bullet} = i_{+}i^{!}(C^{\bullet})[-\dim X]$ (*Th.* 3.5.4). Hence

$$p_{+}(C^{\bullet}) = p_{+}i_{+}i^{!}(C^{\bullet})[-\dim X] = \pi_{+}(i^{!}(C^{\bullet})[-\dim X]) \quad (Prop. \ 3.5.1).$$

Since r is birational to the support of  $i_+(M)$ 

$$\dim \operatorname{supp} i^!(C^{\bullet}) = \dim \operatorname{supp} C^{\bullet} \le \dim V - 1;$$

hence by induction and (6.3.7) it follows that  $p_+(C^{\bullet}) \in D^b_{rs}(\mathfrak{g}_X)$ .

e) Assume that  $\dim V=n.$  There exists a divisor  $S\subset Y$  so that in the distinguished triangle

$$R\Gamma_{[S]}(M) \to M \to M(*S)$$

the  $\mathfrak{g}_Y$ -module M(\*S) is torsion free. Applying  $\pi_+$  we get the distinguished triangle

$$\pi_+ R\Gamma_{[S]}(M) \to \pi_+(M) \to \pi_+(M(*S)) \to .$$

Now  $R\Gamma_{[S]}(M), M(*S) \in D^b_{rs}(\mathfrak{g}_Y)$  (Th. 6.3.11), hence by b)  $\pi_+(M(*S)) \in D^b_{rs}(\mathfrak{g}_X)$  and by d)  $\pi_+(R\Gamma_{[S]}(M)) \in D^b_{rs}(\mathfrak{g}_X)$ . Therefore the third vertex  $\pi_+(M) \in D^b_{rs}(\mathfrak{g}_X)$ .  $\Box$ 

Remark 6.3.14. Using an argument similar to b) in the proof one can prove the theorem for pure modules whose support V is regular in codimension 1 (for example when V is normal) without using Hironaka's resolution of singularities. Still, I was unable to avoid the general resolution of singularities altogether by instead in d) let  $r: V' \to Y \times X$  be the normalization of V.

## Theorem 6.3.15.

$$D^b_{crs}(\mathfrak{g}_X) = D^b_{rs}(\mathfrak{g}_X).$$

The proof of Theorem 6.3.15 is based on two lemmas, where the first is the counterpart for  $D_{crs}^b$  of Theorem 6.3.11.

**Lemma 6.3.16.** Let  $S \subset X$  be a non-singular divisor, where  $i: S \to X$  is the inclusion. Consider the distinguished triangle (Th. 3.5.4)

$$i_*i^!(M^{\bullet})[\mathrm{d}_{S,X}] \to M^{\bullet} \to M^{\bullet}(*S) \to .$$

Then the following are equivalent:

- (1)  $M \in D^b_{crs}(\mathfrak{g}_X);$
- (2)  $R\Gamma_{[S]}(M^{\bullet})$  and  $M^{\bullet}(*S)$  belong to  $D^{b}_{crs}(\mathfrak{g}_{X})$ .

*Proof.* (2)  $\Rightarrow$  (1): Apply Theorem 6.3.12 and Proposition 6.3.5. (1)  $\Rightarrow$  (2): Pull back the distinguished triangle (6.3.3) to a curve  $\phi : C \rightarrow X$ ,

$$\phi^!(R\Gamma_{[S]}(M^{\bullet})) \to \phi^!(M^{\bullet}) \to \phi^!(M^{\bullet}(*S)) \to$$

Then either  $\phi(C) \subset S$ , or the support of the homology of  $\phi^!(R\Gamma_{[S]}(M^{\bullet}))$ is discrete. In the latter case,  $\phi^!(R\Gamma_{[S]}(M^{\bullet})) \in D^b_{rs}(\mathfrak{g}_C)$  by Remark 1, and since by assumption  $\phi^!(M^{\bullet}) \in D^b_{rs}(\mathfrak{g}_C)$ , all vertices in the above distinguished triangle belong to  $D^b_{rs}(\mathfrak{g}_C)$ . In the first, the assertion follows if  $\phi^!(M^{\bullet}(*S)) \cong 0$ , and this follows if  $R\Gamma_{[C]}(M^{\bullet}(*S)) \cong 0$  (Th. 3.5.4). The last assertion is well-known, but not finding an accurate reference we include the argument.

First note that if  $C \subset S$  is an inclusion of closed subsets of X, then the functor  $R\Gamma_{[C]}(\cdot)$  is equivalent to  $R\Gamma_{[C]} \circ R\Gamma_{[S]}(\cdot)$ . To see this, the nonderived functors satisfy for any  $\mathcal{O}_X$ -module M the composition property  $H^0_{[C]}(H^0_{[S]}(M)) = H^0_{[C]}(M)$ , where  $H^0_{[S]}(M) = \{m \in M : I^n_S \cdot m = 0, n \gg 1\}$  $(I_S$  is the ideal of S);  $H^0_{[C]}(\cdot)$  is defined similarly. Now the assertion follows since  $H^0_{[S]}(\cdot)$  takes injective  $\mathcal{O}_X$ -modules to injective modules.

Second, applying  $R\Gamma_{[C]}$  to the distinguished triangle

$$R\Gamma_{[S]}(M^{\bullet}) \to M^{\bullet} \to M^{\bullet}(*S) \to$$

one gets  $R\Gamma_{[C]}(M^{\bullet}(*S)) \cong 0.$ 

**Lemma 6.3.17.** Let  $M^{\bullet} \in D^{b}(\mathfrak{g}_{X})$  and  $x \in X$ . There exists an open neighbourhood  $X_{0}$  of x and a divisor  $S \subset X_{0}$  such that  $M^{\bullet}_{X_{0}}(*S) \cong N^{\bullet}$  (in  $D^{b}(\mathfrak{g}_{X_{0}})$ ) where  $N^{\bullet}$  is a complex of free  $\mathcal{O}_{X_{0}}(*S)$ -modules and  $H^{\bullet}(M^{\bullet}(*S))$  is a free  $\mathcal{O}_{X_{0}}(*S)$ -module.

*Proof.* This follows from [BGK+87, VII, Lem. 9.3].  $\Box$ 

Proof of Theorem 6.3.15. The inclusion  $D^b_{rs}(\mathfrak{g}_X) \subseteq D^b_{crs}(\mathfrak{g}_X)$  follows from Theorem 6.3.12. Now letting  $M^{\bullet} \in D^b_{crs}(\mathfrak{g}_X)$  we have to prove that  $M^{\bullet} \in D^b_{rs}(\mathfrak{g}_X)$ .

The proof uses an induction over  $d = \dim X$ . The assertion is obvious when d = 1, so assume that  $d \ge 2$  and that the theorem is true for all non-singular spaces Y with  $\dim Y < d$ .

a) The assertion is true when dim supp  $M^{\bullet} < d$ : The proof is by induction over the dimension k < d of  $Y_1 = \text{supp } M^{\bullet}$ .

If dim  $Y_1 = 0$  the assertion follows from Remark 1. Assume that the assertion is true for all  $M^{\bullet}$  such that dim supp  $M^{\bullet} < k$ . Let  $\pi : Y \to X$  be

a desingularisation of  $Y_1$  and consider the distinguished triangle in  $D^b(\mathfrak{g}_X)$ (Th. 3.5.3)

$$\pi_+\pi^!(M^{ullet})[\mathrm{d}_{Y,X}] \to M^{ullet} \to N^{ullet} \to$$

Propositions 3.4.3 and 4.2.3 imply

$$\pi^!(M^{\bullet}) \in D^b_{crs}(\mathfrak{g}_Y).$$

As dim Y < d, by induction  $\pi^!(M^{\bullet}) \in D^b_{rs}(\mathfrak{g}_Y)$ , so Theorem 6.3.13 implies

$$\pi_+\pi^!(M^\bullet) \in D^b_{rs}(\mathfrak{g}_X) \subseteq D^b_{crs}(\mathfrak{g}_X).$$

By assumption  $M^{\bullet} \in D^b_{crs}(\mathfrak{g}_X)$ , so the third vertex  $N^{\bullet} \in D^b_{crs}(\mathfrak{g}_X)$ ; since dim supp  $N^{\bullet} < k$  (*Th.* 3.5.4), by induction we get  $N^{\bullet} \in D^{b}_{rs}(\mathfrak{g}_{X})$ . Then since  $\pi_+\pi^!(M^{\bullet}) \in D^b_{rs}(\mathfrak{g}_X)$  the third vertex  $M^{\bullet} \in D^b_{rs}(\mathfrak{g}_X)$ .

b) The assertion holds when dim supp  $M^{\bullet} = d$ : The statement being local in X one can, by Lemma 6.3.17, assume that  $S \subset X$  is a divisor such that  $M^{\bullet}(*S)$  and  $H^{\bullet}(M^{\bullet}(*S)) = H^{\bullet}(M^{\bullet})(*S)$  are locally free over  $\mathcal{O}_X(*S)$ .

 $M^{\bullet}(*S) \in D^{b}_{rs}(\mathfrak{g}_{X})$ : We have to prove that  $H^{\bullet}(M^{\bullet}(*S)) \in \operatorname{coh}_{rs}(\mathfrak{g}_{X})$ , and for this, since  $H^{\bullet}(M^{\bullet}(*S))$  is torsion free, one can assume that S is non-singular (Cor. 5.2.2).

As  $\mathcal{O}_X(*S)$  is locally flat over  $\mathcal{O}_X$  it follows that  $M^{\bullet}(*S)$  is a complex of flat  $\mathcal{O}_X$ -modules. Therefore, if  $\phi: C \to X$  is a curve,

$$H^{\bullet}(\phi^{!}(M^{\bullet}(*S))) \cong \phi^{!}(H^{\bullet}(M^{\bullet}(*S))).$$

Since  $M^{\bullet}(*S) \in D^{b}_{crs}(\mathfrak{g}_X)$  it follows that  $H^{\bullet}(M^{\bullet}(*S)) \in D^{b}_{crs}; H^{\bullet}(M^{\bullet}(*S))$ being torsion free Corollary 6.1.3 implies that  $H^{\bullet}(M^{\bullet}(*S)) \in \operatorname{coh}_{rs}(\mathfrak{g}_X)$ .

 $M^{\bullet} \in D^b_{rs}(\mathfrak{g}_X)$ : Since  $R\Gamma_{[S]}(M^{\bullet}) \in D^b_{crs}(\mathfrak{g}_X)$  (Lem. 6.3.16), and

$$\dim \operatorname{supp} R\Gamma_{[S]}(M^{\bullet}) \leq \dim S < d,$$

by induction  $R\Gamma_{[S]}(M^{\bullet}) \in D^b_{rs}(\mathfrak{g}_X)$ . Therefore the third vertex  $M^{\bullet} \in$  $D^b_{rs}(\mathfrak{g}_X)$ .  $\Box$ 

When  $M^{\bullet} \in D^b_{rs}(\mathfrak{g}_X)$  Proposition 6.3.5 implies that any sub-quotient of  $H^{\bullet}(M^{\bullet})$  belongs to  $\operatorname{coh}_{rs}(\mathfrak{g}_X)$  (full regularity). If we had known that the category  $D^b_{crs}(\mathfrak{g}_X)$  has the same property then Lemma 6.3.17 would be unnecessary in the above proof.

**Corollary 6.3.18.** The following are equivalent:

- (1)  $M^{\bullet} \in D^{b}_{crs}(\mathfrak{g}_{X});$ (2)  $H^{\bullet}(M^{\bullet}) \in D^{b}_{crs}(\mathfrak{g}_{X}).$

If these conditions are satisfied, then any sub-quotient of  $H^{\bullet}(M^{\bullet})$  belongs to  $D^b_{crs}(\mathfrak{g}_X).$ 

Remark 6.3.19. When  $\mathfrak{g}_X = T_X$  we get [BGK+87, VII, Cor. 12.8] and [Bjö93, Th. 5.3.4].

6.4. Completely regular complexes. In this section all spaces X are quasi-projective complex algebraic manifold, and  $X_h$  is the associated complex analytic manifold. All Lie algebroids are locally free and transitive.

Let  $(j', j) : (X, \mathfrak{g}_X, \alpha) \to (\bar{X}, \mathfrak{g}_{\bar{X}}, \bar{\alpha})$  be a morphism of locally free transitive Lie algebroids (*Def.* 2.4.4), where the underlying morphism  $j : X \to \bar{X}$ is an open embedding of algebraic manifolds. We say that (j', j) is a *completion* of  $(X, \mathfrak{g}_X, \alpha)$  if  $\bar{X}$  is projective,  $\bar{\alpha}$  is surjective, and  $j^+(\mathfrak{g}_{\bar{X}}) = \mathfrak{g}_X$ . Completions do not always exist.

Example 6.4.1. Let  $(\mathbf{C}^*, \mathfrak{g}_{\mathbf{C}^*}, \alpha)$  be a transitive Lie algebroid on  $\mathbf{C}^* = \mathbf{C}^1 \setminus \{0\}$  of the form  $\mathfrak{g}_{\mathbf{C}^*} = \mathfrak{b}_{\mathbf{C}^*} \oplus T_{\mathbf{C}^*}$  where  $\mathfrak{b}_{\mathbf{C}^*}$  is a commutative  $\mathcal{O}_{\mathbf{C}^*}$ -Lie algebra, invertible as  $\mathcal{O}_{\mathbf{C}^*}$ -module, and moreover a  $T_{\mathbf{C}^*}$ -module with the global generator  $\mu$  satisfying the relation  $(t^l\partial_t - \gamma)\mu = 0$ , where l is a non-negative integer. The structure of Lie algebroid on  $\mathfrak{b}_{\mathbf{C}^*} \oplus T_{\mathbf{C}^*}$  is  $[(b_1, \partial_1), (b_2, \partial_2)] = (\partial_1(b_2) - \partial_2(b_1), [\partial_1, \partial_2]), (b_i, \partial_i) \in \mathfrak{b}_{\mathbf{C}^*} \oplus T_{\mathbf{C}^*}$ . The unique non-singular completion of  $\mathbf{C}^*$  is the open embedding  $j : \mathbf{C}^* \to \mathbf{P}^1$ . Now if  $l > 1, \gamma \neq 0$ , or  $l = 1, \gamma \notin \mathbf{Z}$ , then  $\mathbf{T}_{\mathbf{C}^*}$ -module  $\mathcal{O}_{\mathbf{C}^*}\mu$  is not the restriction of a  $T_{\mathbf{P}^1}$ -module which is coherent over  $\mathcal{O}_{\mathbf{P}^1}$ ; hence  $(\mathbf{C}^*, \mathfrak{g}_{\mathbf{C}^*}, \alpha)$  does not have a (transitive) completion.

Remark 6.4.2. There always exist non-transitive completions of a Lie algebroid  $(X, \mathfrak{g}_X, \alpha)$ . Let I be the ideal of  $\overline{X} \setminus X$ .  $\overline{X}$  is a noetherian scheme so there exists a coherent  $\mathcal{O}_{\overline{X}}$ -submodule  $\mathfrak{g}_{\overline{X}}^1 \subseteq j_*(\mathfrak{g}_X)$  such that  $j^*(\mathfrak{g}_{\overline{X}}^1) = \mathfrak{g}_X$ . For integers  $n \gg 1$  consider the submodule  $I^n \mathfrak{g}_{\overline{X}}^1 \subset \mathfrak{g}_{\overline{X}}^1$  and let  $\overline{\alpha}$  denote the restriction of  $j_*(\mathfrak{g}_X) \to j_*(T_X)$  to  $\mathfrak{g}_{\overline{X}}^1$ . When  $n \gg 1$  one checks that  $[I^n \mathfrak{g}_{\overline{X}}^1, I^n \mathfrak{g}_{\overline{X}}^1] \subset I^n \mathfrak{g}_{\overline{X}}^1$  and  $\alpha(I^n \mathfrak{g}_{\overline{X}}^1) \subset T_{\overline{X}}^1$ . Thus for sufficiently big n the  $\mathcal{O}_{\overline{X}}$ -module  $\mathfrak{g}_{\overline{X}} = I^n \mathfrak{g}_{\overline{X}}^1$  is a completion of  $\mathfrak{g}_X$  except that the morphism  $\overline{\alpha}$  is not surjective.

We will assume that the Lie algebroids under consideration do have a completion. For example, for the Lie algebroid  $T_X$  any open embedding  $j: X \to \overline{X}$  to a proper algebraic manifold  $\overline{X}$  will do, and more generally if M is a locally free  $\mathcal{O}_X$ -module which has a locally free extension to  $\overline{X}$ , then the linear Lie algebroid  $\mathfrak{c}_X(M)$  has a completion.

Let (j', j) be a completion.

**Definition 6.4.3.** An object  $M^{\bullet} \in D^b_{rs}(\mathfrak{g}_X)$  is completely regular if  $M^{\bullet} = j^!(N^{\bullet})$  for some  $N^{\bullet} \in D^b_{rs}(\mathfrak{g}_{\bar{X}})$ , where  $(\bar{X}, \mathfrak{g}_{\bar{X}}, \alpha)$  is a completion of  $(X, \mathfrak{g}_X, \alpha)$ .

Let  $\bar{D}_{rs}^b(\mathfrak{g}_X) \subset D_{rs}^b(\mathfrak{g}_X)$  be the sub-category of completely regular complexes.

We also say that  $N^{\bullet}$  is a (regular) completion of  $M^{\bullet}$ . By Theorem 6.3.11 there exist such  $N^{\bullet}$  if and only if  $j_{+}(M^{\bullet}) = N^{\bullet}(*S) \in D^{b}_{rs}(\mathfrak{g}_{\bar{X}})$ , where  $S = \bar{X} \setminus j(X)$ , so it is natural to check if  $j_{+}(M^{\bullet}) \in D^{b}_{rs}(\mathfrak{g}_{\bar{X}})$ .

Next lemma shows that the definition of  $\bar{D}_{rs}^{b}(\mathfrak{g}_{X})$  is intrinsic.

Lemma 6.4.4. Let

$$(j_1, j'_1) : (X, \mathfrak{g}_X, \alpha) \to (X_1, \mathfrak{g}_{X_1}, \alpha_1)$$

and

$$(j_2, j'_2): (X, \mathfrak{g}_X, \alpha) \to (X_2, \mathfrak{g}_{X_2}, \alpha_2)$$
<sup>75</sup>

be completions. Put  $S_2 = X_2 \setminus j_2(X)$  and let  $M^{\bullet} \in D^b_{rs}(\mathfrak{g}_X)$ . Suppose that there exist  $N^{\bullet}_i \in D^b_{rs}(\mathfrak{g}_{X_i})$ , i = 1, 2, satisfying  $j^!(N^{\bullet}_i) \cong M^{\bullet}$ . Then if  $N^{\bullet}_1 \in D^b_{rs}(\mathfrak{g}_{X_1})$  and  $R\Gamma_{[S_2]}(N^{\bullet}_2) \in D^b_{rs}(\mathfrak{g}_{X_2})$  it follows that  $N^{\bullet}_2 \in D^b_{rs}(\mathfrak{g}_{X_2})$ .

*Proof.* By Theorem 6.3.11 it suffices to prove that  $N_2^{\bullet}(*S) \in D_{rs}^b(\mathfrak{g}_{X_2})$ .

Let  $X_3 \to X_1 \times X_2$  be a desingularisation of the closure of the image of the morphism  $(j_1, j_2) : X \to X_1 \times X_2$ , so there exist canonical proper birational morphisms  $p : X_3 \to X_1$ ,  $q : X_3 \to X_2$ , and letting  $j : X \to X_3$  be the inclusion morphism of X in  $X_3$  we have  $q \circ j = p \circ j = \operatorname{id}_X$ . Let  $S_3, S_2$ be the algebraic sets  $X_3 \setminus j(X)$  and  $X_2 \setminus j_2(X)$ .

We have  $N_3^{\bullet} := p!(N_1^{\bullet}) \in D^b_{rs}(\mathfrak{g}_{X_3})$  (Th. 6.3.12) and  $j!(N_3^{\bullet}) = j_2!(N_2^{\bullet}) = M^{\bullet}$ , so

$$N_{2}^{\bullet}(*S_{2}) = (j_{2})_{+}j_{2}^{!}(N_{2}^{\bullet})$$
  
=  $(q \circ j)_{+}j^{!}(N_{3}^{\bullet})$   
=  $q_{+}(j_{+}j^{!}(N_{3}^{\bullet}))$  (Prop. 3.5.1)  
=  $q_{+}(N_{3}^{\bullet}(*S_{3})).$ 

By Theorem 6.3.11  $N_3(*S_3) \in D^b_{rs}(\mathfrak{g}_{X_3})$ , noting that the pull-back of a transitive Lie algebroid is transitive, hence by Theorem 6.3.13  $N^{\bullet}_2(*S_2) \in D^b_{rs}(\mathfrak{g}_{X_2})$ .  $\Box$ 

In the following two corollaries to Section 6.3  $\pi: Y \to X$  is a morphism of quasi-projective algebraic manifolds.

We first note that  $\mathfrak{g}_Y$  has a completion if  $\mathfrak{g}_X$  has one. By assumption there exists a completion  $(i', i) : (X, \mathfrak{g}_X, \alpha) \to (\bar{X}, \mathfrak{g}_{\bar{X}}, \bar{\alpha})$ . Let  $Y \subseteq Y_1$  where  $Y_1$  is projective, define  $g: Y \to X \times Y_1$ ,  $g(y) = (\pi(y), y)$  and let  $\phi: \bar{Y} \to X \times Y_1$ be a desingularisation of the closure of g(Y). Now there exists an open embedding  $j: Y \to \bar{Y}$  such that  $\phi \circ j = g$  and a Cartesian diagram:

(6.4.1) 
$$\begin{array}{ccc} Y & \stackrel{j}{\longrightarrow} & \bar{Y} \\ \pi & & & \downarrow \\ X & \stackrel{i}{\longrightarrow} & \bar{X}. \end{array}$$

Since  $\mathfrak{g}_{\bar{X}}$  is locally free and transitive it follows that  $\bar{\pi}^+(\mathfrak{g}_{\bar{X}})$  is locally free and transitive (*Lem.* 2.4.8); hence

$$(\pi^+(i'),j):(Y,\mathfrak{g}_Y=\pi^+(\mathfrak{g}_X),\pi^+(\alpha))\to(\bar{Y},\pi^+(\mathfrak{g}_{\bar{X}}),\pi^+(\bar{\alpha}))$$

is a completion.

We have for  $M^{\bullet} \in D^b(\mathfrak{g}_Y)$ 

(6.4.2) 
$$j_+\pi^!(M^{\bullet}) = \bar{\pi}^! i_+(M^{\bullet})$$

since i and j are open embeddings.

Corollary 6.4.5.  $\pi^!(\bar{D}^b_{rs}(\mathfrak{g}_X)) \subset \bar{D}^b_{rs}(\mathfrak{g}_Y).$ 

*Proof.* As  $i_+(M^{\bullet}) \in D^b_{rs}(\mathfrak{g}_{\bar{X}})$  the result follows from (6.4.2) and Theorem 6.3.12.  $\Box$ 

**Corollary 6.4.6.** Assume that  $\pi$  can be factorized into an open embedding which is a completion of  $(Y, \mathfrak{g}_Y, \alpha)$  and a proper morphism. Then

$$\pi_+(\bar{D}^b_{rs}(\mathfrak{g}_Y))\subset \bar{D}^b_{rs}(\mathfrak{g}_X).$$

Proof. Let  $M^{\bullet} \in \overline{D}_{rs}^{b}(\mathfrak{g}_{Y})$ . If  $\pi = p \circ j$  where  $j : Y \to \overline{Y}$  is a completion we know that  $j_{+}(M^{\bullet}) \in D_{rs}^{b}(Y)$ , and since p is proper  $\pi_{+}(M^{\bullet}) = p_{+}(j_{+}(M^{\bullet})) \in$  $D_{rs}^{b}(\mathfrak{g}_{X})$  (Th. 6.3.13). Since  $p(\overline{Y})$  is a closed subspace of  $\overline{X}$  the restriction of i to  $p(\overline{Y})$  is proper. The proof of Theorem 6.3.13 is now applicable to prove that  $i_{+}\pi_{+}(M^{\bullet}) \in D_{rs}^{b}(\mathfrak{g}_{\overline{X}})$  only using that the restriction of i to the support of  $p_{+}(j_{+}(M^{\bullet}))$  is proper.  $\Box$ 

When  $\mathfrak{g}_X = T_X$  all open embeddings are completions so Corollary 6.4.6 holds for any morphism  $\pi$ ; in this case it is proven for algebraic  $\mathcal{D}_X$ -modules by Borel [BGK<sup>+</sup>87, VII, Th. 12.2] using curve regularity as the definition of complete regularity, cf. Corollary 6.4.7 below. The proof in [loc. cit.] is, as is ours, inspired by Deligne's proof that the Gauss-Manin connection has regular singularities [Del70, II. §7]. In [Bjö93, Ch. V, 5.5.28] the theorem is proven for analytic  $\mathcal{D}_X$ -modules, by proving the comparison condition

$$R\Gamma_{[x]}(\Omega(T_X, \pi_+(M^{\bullet}))) \cong R\Gamma_x(\Omega(T_X, \pi_+(M^{\bullet}))),$$

where the left side is the tempered local cohomology of the de Rham complex of  $\pi_+(M^{\bullet})$ , and the right side is the full local cohomology at any point  $x \in X$ , when the analogous comparison condition holds for M at all points of Y.

Let  $\overline{D}^b_{crs}(\mathfrak{g}_X) \subset D^b_{crs}(\mathfrak{g}_X)$  be the sub-category of complexes  $M^{\bullet}$  such that  $\phi^!(M^{\bullet}) \in \overline{D}^b_{rs}(\mathfrak{g}_C)$  whenever  $\phi: C \to X$  is a curve in X.

## Corollary 6.4.7.

$$\bar{D}^b_{crs}(\mathfrak{g}_X) = \bar{D}^b_{rs}(\mathfrak{g}_X).$$

*Proof.* Let  $(j, j') : (X, \mathfrak{g}_X, \alpha) \to (\overline{X}, \mathfrak{g}_{\overline{X}}, \overline{\alpha})$  be a completion, and S be the algebraic set  $\overline{X} \setminus j(X)$ .

 $\overline{D}_{rs}^{b}(\mathfrak{g}_{X}) \subseteq \overline{D}_{crs}^{b}(\mathfrak{g}_{X})$ : A morphism  $\phi: C \to X$ , where C is curve, can be completed to a morphism  $\overline{\phi}: \overline{C} \to \overline{X}$ , where  $\overline{C}$  is a proper curve, by the valuative criterion of properness. Let  $i: C \to \overline{C}$  be the corresponding open inclusion of (non-singular) curves, so  $\overline{\phi} \circ i = j \circ \phi$ . Then if  $N^{\bullet} \in D_{rs}^{b}(\mathfrak{g}_{\overline{X}})$ , we have  $\overline{\phi}^{!}(N^{\bullet}) \in D_{rs}^{b}(\mathfrak{g}_{\overline{C}})$  (Cor. 6.4.5) and  $i^{!}\overline{\phi}^{!}(N^{\bullet}) = \phi^{!}j^{!}(N^{\bullet})$  (Prop. 3.4.3); hence  $\phi^{!} \circ j^{!}(N^{\bullet}) \in \overline{D}_{crs}^{b}(\mathfrak{g}_{C})$ ; hence  $j^{!}(N^{\bullet}) \in \overline{D}_{crs}^{b}(\mathfrak{g}_{X})$ .  $\overline{D}_{crs}^{b}(\mathfrak{g}_{X}) \subseteq \overline{D}_{rs}^{b}(\mathfrak{g}_{X})$ : Let  $M^{\bullet} \in \overline{D}_{crs}^{b}(\mathfrak{g}_{X})$  and  $N^{\bullet} \in D_{coh}^{b}(\mathfrak{g}_{\overline{X}})$  be such that  $j^{!}(N^{\bullet}) = M^{\bullet}$  and  $N^{\bullet} = N^{\bullet}(*S)$ . By Theorem 6.3.15 we have to prove

 $\bar{D}^{b}_{crs}(\mathfrak{g}_{X}) \subseteq \bar{D}^{b}_{rs}(\mathfrak{g}_{X})$ : Let  $M^{\bullet} \in \bar{D}^{b}_{crs}(\mathfrak{g}_{X})$  and  $N^{\bullet} \in D^{b}_{coh}(\mathfrak{g}_{\bar{X}})$  be such that  $j^{!}(N^{\bullet}) = M^{\bullet}$  and  $N^{\bullet} = N^{\bullet}(*S)$ . By Theorem 6.3.15 we have to prove that  $N^{\bullet} \in D^{b}_{crs}(\mathfrak{g}_{\bar{X}})$ . Let  $\bar{\phi}: \bar{C} \to \bar{X}$  be a curve. Then either  $\bar{\phi}(\bar{C})$  is contained in S, or  $C := \bar{\phi}^{-1}(j(X)) \subset \bar{C}$  is Zariski dense. In the first case,  $\bar{\phi}^{!}(N^{\bullet}(*S)) = 0$ , see the proof of Lemma 6.3.16 or [BGK<sup>+</sup>87, VI, Cor. 8.5]; hence we can assume that  $i: C \to \bar{C}$  is an inclusion of a Zariski open dense subset of C. Now, by assumption,  $i^{!}\bar{\phi}^{!}(N^{\bullet}) = \phi^{!}j^{!}(N^{\bullet}) \in \bar{D}^{b}_{rs}(\mathfrak{g}_{C})$ ; hence  $\bar{\phi}^{!}(N^{\bullet}) \in D^{b}_{rs}(\mathfrak{g}_{\bar{C}})$ .  $\Box$ 

One can improve Proposition 4.2.4 when considering  $\bar{D}_{rs}^{b}(\mathfrak{g}_{X})$ :

**Proposition 6.4.8.** Let  $M^{\bullet} \in D^{b}_{\mathrm{coh}}(\mathfrak{g}_{X})$ . Assume that  $\pi$  is surjective on  $\mathrm{supp} M^{\bullet} \subset X$ . Then the following are equivalent:

(1)  $M^{\bullet} \in \overline{D}_{rs}^{b}(\mathfrak{g}_{X}).$ 

(2)  $\pi^!(M^{\bullet}) \in \overline{D}^b_{rs}(\mathfrak{g}_Y).$ 

When  $\mathfrak{g}_X = T_X$  we recover [Bjö93, Th. 5.6.5] and [BGK<sup>+</sup>87, VII, Prop. 12.9]. Proposition 6.4.8 can be used to prove the complete regularity of certain equivariant  $\mathfrak{g}_X$ -modules occurring as localizations of Harish-Chandra representations of Lie algebras.

*Proof.*  $(1) \Rightarrow (2)$  follows from Corollary 6.4.5, so we need to prove  $(2) \Rightarrow (1)$ .

Letting  $\alpha: C \to X$  be a curve one has to prove that  $\pi^!(M^{\bullet}) \in \overline{D}^b_{rs}(\mathfrak{g}_Y)$ implies  $\alpha^{!}(M^{\bullet}) \in \overline{D}^{b}_{rs}(\mathfrak{g}_{C})$  (Cor. 6.4.7). Let  $p: Y_{1} \to C$  be a desingularisation of the base change  $\alpha$  of  $\pi$ , and  $\phi: Y_1 \to Y$  be the second projection. Since  $p' \circ \alpha' \cong \phi' \circ \pi'$  (Prop. 3.4.3) we have that  $p' \alpha'(M^{\bullet}) \in \overline{D}_{rs}^{b}$ . It therefore suffices to prove the proposition when  $\pi$  is a morphism to a curve X. Moreover, there exists a curve  $\alpha_1 : C_1 \to Y$  such that the composed morphism  $\pi \circ \alpha_1 : C_1 \to X$  is dominant, which moreover can be assumed to be etale after restricting to an open subset of  $C_1$ . Since  $\alpha_1^{!}$  preserves  $D_{rs}^{b}(\mathfrak{g}_Y)$ (Cor. 6.4.5), by Lemma 6.4.4 we can then assume that  $\pi: Y \to X$  is an etale morphism of non-singular curves. Then as complex of  $\mathcal{O}_Y$ -modules the inverse image  $\pi^!(M^{\bullet})$  coincides with  $\pi^*(M^{\bullet})$ , the inverse image in the category of  $\mathcal{O}_Y$ -modules, and if  $N \in \overline{D}_{rs}^b(\mathfrak{g}_Y)$ , then  $\pi_+(N^{\bullet})$  coincides with  $\pi_*(N^{\bullet})$ , the direct image in the category of  $\mathcal{O}_X$ -modules. Moreover, that  $\pi$  is etale also implies that the canonical morphism  $M^{\bullet} \to \pi_+ \pi^! (M^{\bullet}) = \pi_* \pi^* (M^{\bullet})$  is a monomorphism in a triangulated category, hence it is split. Consider a completions of X and Y forming a Cartesian diagram of the type (6.4.1). Then  $\bar{\pi}_+ j_+ \pi^!(M^{\bullet}) \cong i_+ \pi_+ \pi^!(M^{\bullet}) \cong i_* \pi_* \pi^*(M^{\bullet})$ . Hence we have a split distinguished triangle

$$i_+(M^{\bullet}) \to \bar{\pi}_+ j_+ \pi^!(M^{\bullet}) \to C^{\bullet} \xrightarrow{0},$$

so  $H^{\bullet}(i_{+}(M^{\bullet})) \subset H^{\bullet}(\bar{\pi}_{+}j_{+}\pi^{!}(M^{\bullet}))$ ; by assumption and Theorem 6.3.13  $H^{\bullet}(\bar{\pi}_{+}j_{+}\pi^{!}(M^{\bullet})) \in \operatorname{coh}_{rs}(\mathfrak{g}_{\bar{X}})$ ; hence  $H^{\bullet}(i_{+}(M^{\bullet})) \in \operatorname{coh}_{rs}(\mathfrak{g}_{\bar{X}})$  (*Prop.* 6.3.5).  $\Box$ 

Let us make a comment about the role of  $\bar{D}_{rs}^{b}(\mathfrak{g}_{X})$  in the Riemann-Hilbert correspondence; compare also to the rather different argument in [BGK<sup>+</sup>87, VIII]. Fix a completion  $(X,\mathfrak{g}_{X},\alpha) \to (\bar{X},\mathfrak{g}_{\bar{X}},\bar{\alpha})$ . Put  $S = \bar{X} \setminus j(X)$  and let  $D_{rs}^{b}(S,\mathfrak{g}_{\bar{X}}) \subset D_{rs}^{b}(\mathfrak{g}_{\bar{X}})$  be the full sub-category of objects  $N^{\bullet} \in D_{rs}^{b}(\mathfrak{g}_{\bar{X}})$ such that  $N^{\bullet} = N^{\bullet}(*S) = Rj_{*}j^{*}(N^{\bullet})$ . Then the functor

$$j_+: \overline{D}^b_{rs}(\mathfrak{g}_X) \to D^b_{rs}(S, \mathfrak{g}_{\overline{X}}), \quad M^{\bullet} \mapsto j_+(M^{\bullet})$$

is an equivalence of categories, and using GAGA (4.4)

$$g: D^b_{rs}(S, \mathfrak{g}_{\bar{X}}) \cong D^b_{rs}(S, \mathfrak{g}_{\bar{X}_h}).$$

A main step is to prove that the restriction functor defines an equivalence

$$r: D^b_{rs}(S, \mathfrak{g}_{\bar{X}_h}) \cong D^b_{rs}(\mathfrak{g}_{X_h}).$$

Since this is a local problem in  $X_h$  the well-known proof when  $\mathfrak{g}_{X_h} = T_{X_h}$  works, based on Hironaka's resolution of singularities; see Malgrange's account in [BGK<sup>+</sup>87, IV] of Deligne's proof for the case of connections. Assume now that  $\mathfrak{g}_X \in \operatorname{LPic}_X$  and recall the discussion in (3.3.3). Put

 $\phi = t([\mathfrak{g}_{X_h}]) \in H^2(X_h, \mathbb{C}^*)$ , so one has the invertible  $\phi$ -twisted sheaf  $\lambda(\phi) \in Mod(\mathfrak{g}_{X_h}, \phi)$ , and the solution functor

$$Sol_{\phi}: D^{b}(\mathfrak{g}_{X_{h}}) \to D^{b}(\mathbf{C}_{X_{h}}, \phi), \quad M \to RHom_{\mathfrak{g}_{X_{h}}}(M, \lambda(\phi)).$$

The usual Riemann-Hilbert correspondence states that  $Sol_0$  is an equivalence of categories between  $D_{rs}^b(T_{X_h})$  and the derived category  $D^b(\mathbf{C}_{X_h})$ of complexes of sheaves whose homology sheaves are constructible, while the abelian sub-category  $\operatorname{coh}_{rs}(T_{X_h})$  is equivalent to the category of perverse sheaves on  $X_h$  ([Kas80], [Kas84], [Meb80], [Meb82], [Meb89]). Now  $Sol_{\phi}$  defines an equivalence between  $D_{rs}^b(\mathfrak{g}_{X_h})$  and the derived category of complexes of  $\phi$ -twisted sheaves whose homology  $\phi$ -twisted sheaves are constructible; see also [Kas89]. Hence by the previous equivalences there exists an equivalence between an algebraic and a topological category:

$$Sol_{\phi} \circ r \circ g \circ j_{+} : \overline{D}^{b}_{rs}(\mathfrak{g}_{X}) \cong D^{b}(\mathbf{C}_{X_{h}}, \phi).$$

## References

- [Ati57] Michael Atiyah. Complex analytic connections in fibre bundles. Trans. Amer. Math. Soc., 85:181–207, 1957.
- [BB81] Alexander Beilinson and Joseph Bernstein. Localisation de g-modules. C.R. Acad. Sci. (Paris), 292(Ser. I.), 1981.
- [BB93] Alexander Beilinson and Joseph Bernstein. A proof of Jantzen conjectures. Adv. Sov. Math., 16(1):1–50, 1993.
- [BBD82] Alexander Beilinson, Joseph Bernstein, and Pierre Deligne. Fasciuex pervers. Asterisque), 100:1–100, 1982.
- [BGG72] I. N. Bernšteĭn, I. M. Gelfand, and S. I. Gelfand. Differential operators on a cubic cone. Uspehi Mat. Nauk, 27(1(163)):185–190, 1972.
- [BGK<sup>+</sup>87] A. Borel, P.-P. Grivel, B. Kaup, A. Haefliger, B. Malgrange, and F. Ehlers. Algebraic D-modules, volume 2 of Perspectives in Mathematics. Academic Press Inc., Boston, MA, 1987.
- [Bjö93] Jan-Erik Björk. Analytic D-modules and applications. Kluwer Academic Publishers, 1993.
- [BK81] Jean-Luc Brylinski and Masaki Kashiwara. Kazhdan-Lusztig conjecture and holonomic systems. *Invent. Math.*, 64(387), 1981.
- [BS58] Armand Borel and Jean-Pierre Serre. Le théorème de Riemann-Roch (d'après des résultats inédits de A. Grothendieck). Bull. Soc. Math France, 86:97–136, 1958.
- [BS76] Constantin Bănică and Octavian Stănășilă. Algebraic methods in the global theory of complex spaces. Editura Academiei, Bucharest, 1976. Translated from the Romanian.
- [BS88] Alexander Beilinson and Vadim Schechtman. Determinant bundles and virasoro algebras. Commun. Math. Phys., 118:651 – 701, 1988.
- [BS98] M. P. Brodmann and R. Y. Sharp. Local cohomology: an algebraic introduction with geometric applications, volume 60 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1998.
- [Del70] Pierre Deligne. Équations différentielles à points singuliers réguliers, volume 163. Springer Math. L.N, 1970.
- [Fri67] Jacques Frisch. Points de platitude d'un morphisme d'espaces analytiques complexes. Invent. Math., 4:118–138, 1967.
- [Gab81] Ofer Gabber. The integrability of the characteristic variety. Am Jour Math, 103(3):445–468, 1981.
- [GD61] Alexander Grothendieck and Jean Dieudonne. étude cohomolique des faisceaux cohérent. *Publ. IHES*, (4), 1961.
- [GD71] Alexander Grothendieck and Jean Dieudonne. Eléments de géométrie algebrique. I. Berlin-Heidelberg-New York: Springer-Verlag, 1971.

- [Gir71] Jean Giraud. *Cohomologie non abélienne*. Springer-Verlag, Berlin, 1971. Die Grundlehren der mathematischen Wissenschaften, Band 179.
- [God73] Roger Godement. Topologie algébrique et théorie des faisceaux. Hermann, Paris, 1973. Troisième édition revue et corrigée, Publications de l'Institut de Mathématique de l'Université de Strasbourg, XIII, Actualités Scientifiques et Industrielles, No. 1252.
- [GR79] Hans Grauert and Reinhold Remmert. Theory of Stein spaces, volume 236 of Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, 1979.
- [Gra60] Hans Grauert. Ein theorem der analytischen garbentheorie und die modulräume komplexer strukturen. *Publ. IHES*, (5), 1960.
- [Gro62] Alexander Grothendieck. Cohomologie locale de faisceaux cohérents et Théorèmes de Lefshetz locaux et globaux (SGA 2), volume 2. North-Holland, advanced studies in pure mathematics edition, 1962.
- [Gro67] Alexander Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV. Inst. Hautes Études Sci. Publ. Math. No., 32:361, 1967.
- [Gro71] Alexander Grothendieck, et al. Revêtements étales et groupe fondamental. Springer-Verlag, Berlin, 1971. Séminaire de Géométrie Algébrique du Bois Marie 1960–1961 (SGA 1), Dirigé par Alexandre Grothendieck. Augmenté de deux exposés de M. Raynaud, Lecture Notes in Mathematics, Vol. 224.
- [Gun90] Robert C. Gunning. Introduction to holomorphic functions of several variables. Vol. II. The Wadsworth & Brooks/Cole Mathematics Series. Wadsworth & Brooks/Cole Advanced Books & Software, Monterey, CA, 1990.
- [Har61] R. Jr. Harper, Laurence. On differentiably simple algebras. Trans. Amer. Math. Soc., 100:63–72, 1961.
- [Har66] Robin Hartshorne. *Residues and duality*. Springer-Verlag, Berlin, 1966. Lecture notes of a seminar on the work of A. Grothendieck, given at Harvard 1963/64. With an appendix by P. Deligne. Lecture Notes in Mathematics, No. 20.
- [Jan79] Jens Carsten Jantzen. Moduln mit einem höchsten Gewicht, volume 750 of Lect. Notes Math. Springer-Verlag, 1979.
- [Käl96] Rolf Källström. Homological dimensions of Lie algebroids. Manuscript, 1996.
- [Käl98] Rolf Källström. Smooth modules over lie algebroids, II. Manuscript, 1998.
- [Kas] Masaki Kashiwara. Algebraic study of systems of partial differential equations. Mém. Soc. Math. France (N.S.), (63).
- [Kas80] Masaki Kashiwara. Faisceaux constructibles et systèmes holonômes d'équations aux dérivées partielles linéaires à points singuliers réguliers. In Séminaire Goulaouic-Schwartz, 1979–1980 (French), pages Exp. No. 19, 7. École Polytech., Palaiseau, 1980.
- [Kas84] Masaki Kashiwara. The Riemann- Hilbert problem for holonomic systems. Publ. Res. Inst. Math. Sci., 20(2):319–365, 1984.
- [Kas89] Masaki Kashiwara. Representation theory and D-modules on flag varieties. Société Math. de France, Astérisque, pages 55–190, 1989.
- [Kat87] Nicholas M. Katz. On the calculation of some differential Galois groups. Invent. Math., 87(1):13–61, 1987.
- [KK81] Masaki Kashiwara and Takahiro Kawai. On holonomic systems of microdifferential equations. III. Systems with regular singularities. *Publ. Res. Inst. Math. Sci.*, 17(3):813–979, 1981.
- [Lev75] A. H. M. Levelt. Jordan decomposition for a class of singular differential operators. Ark. Mat., 13:1–27, 1975.
- [Lyu93] Gennady Lyubeznik. Finiteness properties of local cohomology modules (an application of D-modules to commutative algebra). Invent. Math., 113(1):41– 55, 1993.
- [Mac95] Kirill C. H. Mackenzie. Lie algebroids and Lie pseudoalgebras. Bull. London Math. Soc., 27(2):97–147, 1995.
- [Mal96] Bernard Malgrange. Connexions méromorphes 2. *Invent. Math*, 124:367–387, 1996.

[Mat86]	Hideyuki Matsumura. <i>Commutative Ring Theory</i> . Cambridge University Press, 1986.
[Meb80]	Zoghman Mebkhout. Sur le problème de Hilbert-Riemann. In <i>Complex analysis, microlocal calculus and relativistic quantum theory (Proc. Internat. Colloq., Centre Phys., Les Houches, 1979)</i> , volume 126 of <i>Lecture Notes in Phys.</i> , pages 90–110. Springer, Berlin, 1980.
$[\mathrm{Meb82}]$	Zoghman Mebkhout. Théorèmes de bidualité locale pour les $\mathcal{D}_X$ -modules holonomes. Ark. Mat., 20(1):111–124, 1982.
[Meb84]	Zoghman Mebkhout. Une équivalence de catégories, and une autre équivalence de catégories. <i>Compositio Math.</i> , 51(1):51–62, 63–88, 1984.
[Meb89]	Zoghman Mebkhout. Le formalisme des six opérations de Grothendieck pour les $\mathcal{D}_X$ -modules cohérents, volume 35 of Travaux en Cours [Works in Progress]. Hermann, Paris, 1989. With supplementary material by the author and L. Narváez Macarro.
[Oda83]	Tadao Oda. Introduction to algebraic analysis on complex manifolds. In Al- gebraic varieties and analytic varieties (Tokyo, 1981), volume 1 of Adv. Stud. Pure Math., pages 29–48. North-Holland, Amsterdam, 1983.
[Rei67]	Hans-Jörg Reiffen. Das Lemma von Poincaré für holomorphe Differential- formen auf komplexen Räumen. <i>Math. Z.</i> , 101:269–284, 1967.
[Rin63]	George S. Rinehart. Differential forms on general commutative algebras. <i>Trans. Amer. Math. Soc.</i> , 108:195–222, 1963.
[Sai80]	Kyoji Saito. Theory of logarithmic differential forms and logarithmic vector fields. J. Fac. Sci. Univ. Tokyo Sect. IA Math., 27(2):265–291, 1980.
[Sch64]	Günter Scheja. Fortsetzungssätze der komplex-analytischen Cohomologie und ihre algebraische Charakterisierung. <i>Math. Ann.</i> , 157:75–94, 1964.
[Sei66]	Abraham Seidenberg. Derivations and integral closure. <i>Pacific J. of Math.</i> , 16:167–73, 1966.
[Ser56]	Jean-Pierre Serre. Géométrie algébrique et géométrie analytique. Ann. Inst. Fourier, Grenoble, 6:1–42, 1955–1956.
[Ser66]	Jean-Pierre Serre. Prolongement de faisceaux analytiques coherents. Ann. Inst. Fourier, 16(1):363–374, 1966.
[Siu70]	Yum-Tong Siu. Analytic sheaves of local cohomology. Trans. Amer. Math. Soc., 148:347–366, 1970.
[Tra69]	Günther Trautmann. Ein Endlichkeitssatz in der analytischen Geometrie. Invent. Math., 8:143–174, 1969.
[TUY89]	Akihiro Tsuchiya, Kenji Ueno, and Yasuhiko Yamada. Conformal field the- ory on universal family of stable curves with gauge symmetries. In <i>Integrable</i> systems in quantum field theory and statistical mechanics, volume 19 of Adv. Stud. Pure Math., pages 459–566. Academic Press, Boston, MA, 1989.
[Uen97]	Kenji Ueno. Introduction to conformal field theory with gauge symmetries. In Geometry and physics (Aarhus, 1995), volume 184 of Lecture Notes in Pure
[Wei58]	and Appl. Math., pages 603–745. Dekker, New York, 1997. André Weil. Introduction à l'étude des variétés kählériennes. Hermann, Paris, 1958. Publications de l'Institut de Mathématique de l'Université de Nancago, VI. Actualités Sci. Ind. no. 1267.

Department of Mathematics, Royal Institute of Technology, 100 44 Stockholm, Sweden

*E-mail address*: rok@math.kth.se