

**CENTER AND UNIVERSAL R-MATRIX
FOR
QUANTIZED BORCHERDS SUPERALGEBRAS**

JIN HONG

ABSTRACT. We construct a nondegenerate symmetric bilinear form on quantized enveloping algebras associated to Borchers superalgebras. With this, we study its center and its universal R-matrix.

1. INTRODUCTION

Quantized enveloping algebras for Kac-Moody algebras were introduced independently by Drinfel'd([5]) and Jimbo([10]) in their studies of the Yang-Baxter equation. The Kac-Moody algebras were generalized([3]) by Borchers to accommodate for his study of the monstrous moonshine([4]). And quantized version of the enveloping algebras for Borchers algebras([14]) was soon studied. There are also superalgebra versions of these algebras([2]). We shall study the structure of the center and find the R-matrix for the quantized Borchers superalgebras.

Much work has been done on the center of quantized enveloping algebras for finite dimensional semisimple Lie algebras([1, 7, 11, 18–21]), and there are Kac-Moody([8]) and Borchers([15]) versions also. We will mainly follow [21] and [15] to find the center for quantized Borchers superalgebras.

As for the universal R-matrix, the quantum double construction by Drinfel'd([6]) gives its existence for any Hopf algebra satisfying some conditions. Even though there is a quantum double construction for \mathbf{Z}_2 -graded Hopf algebras([9]), we do not use it in this paper. Instead, we explicitly construct a universal R-matrix and show that it satisfies the Yang-Baxter equation.

The paper is organized as follows. In Section 2, we define the quantized Borchers superalgebras and give it a Hopf algebra structure. The triangular decomposition will also be mentioned. In Section 3, the character formula for highest weight representations will be given and we prove a lemma that will be used in later sections. The next section is devoted to providing the quantized Borchers superalgebras with a bilinear form and proving its nondegeneracy. In Section 5, we define the Harish-Chandra homomorphism, show its injectivity, and prove some properties concerning its image. Information on the center of the quantized Borchers superalgebra will be obtained in Section 6. The last section will give the universal R-matrix and show that it satisfies the Yang-Baxter equation.

2. QUANTUM DEFORMATION OF BORCHERDS SUPERALGEBRAS

In this section, we define the quantized Borchers superalgebras and give it a Hopf algebra structure.

Supported in part by GARC-KOSEF at Seoul National University.

Let I be a countable index set. A matrix $A = (a_{i,j})_{i,j \in I}$ with entries in the real numbers is a *Borcherds-Cartan* matrix if

- $a_{i,i} = 2$ or $a_{i,i} \leq 0$ for all $i \in I$,
- $a_{i,j} \leq 0$ if $i \neq j$ and $a_{i,j} \in \mathbf{Z}$ if $a_{i,i} = 2$,
- $a_{i,j} = 0$ if and only if $a_{j,i} = 0$.

If there exists a diagonal matrix $D = \text{diag}(s_i | i \in I, s_i > 0)$ such that DA is symmetric, then A is said to be *symmetrizable*. If a symmetrizable Borcherds-Cartan matrix A further satisfies the constraints,

- $a_{i,j} \in \mathbf{Z}$,
- $a_{i,i} \in 2\mathbf{Z}$,
- $s_i \in \mathbf{Z}_{>0}$,

for all $i, j \in I$, then it is said to be *integral*.

A complex matrix $C = (\theta_{i,j})_{i,j \in I}$ is a *coloring matrix* if $\theta_{i,j}\theta_{j,i} = 1$ for all $i, j \in I$. Necessarily, $\theta_{i,i} = \pm 1$ and we say i is *even* when $\theta_{i,i} = 1$, *odd* when $\theta_{i,i} = -1$. A Borcherds-Cartan matrix A is *colored by* C if for every $i \in I$ such that $a_{i,i} = 2$ and $\theta_{i,i} = -1$ we have $a_{i,j} \in 2\mathbf{Z}$ for all $j \in I$.

Throughout this paper, we shall assume that A is a symmetrizable integral Borcherds-Cartan matrix which is colored by a coloring matrix C .

Let $I^e = \{i \in I | a_{i,i} = 2\}$ and $I^{\text{im}} = \{i \in I | a_{i,i} \leq 0\}$. Also let $\underline{m} = (m_i | i \in I)$ be a collection of positive integers such that $m_i = 1$ for all $i \in I^e$. We call \underline{m} the *charge* of the Borcherds-Cartan matrix A .

For a symmetrizable integral Borcherds-Cartan matrix A , which is colored by a coloring matrix C , we denote by $\mathfrak{g}(A, \underline{m}, C)$ the Borcherds superalgebra of charge \underline{m} . (See [2].)

We set $P^\vee = (\bigoplus_{i \in I} \mathbf{Z}h_i) \oplus (\bigoplus_{i \in I} \mathbf{Z}d_i)$ and let $\mathfrak{h} = \mathbf{C} \otimes_{\mathbf{Z}} P^\vee$ be the complex vector space with basis $\{h_i, d_i | i \in I\}$. For $i \in I$, we define α_i in the dual space \mathfrak{h}^* of \mathfrak{h} by setting $\alpha_i(h_j) = a_{j,i}$ and $\alpha_i(d_j) = \delta_{i,j}$. Since A is assumed to be symmetrizable, there exists a nondegenerate symmetric bilinear form $(\ |)$ on \mathfrak{h} given by $(s_i h_i | h) = \alpha_i(h)$ and $(d_i | d_j) = 0$ for $i, j \in I, h \in \mathfrak{h}$.

The free abelian group $Q = \bigoplus_{i \in I} \mathbf{Z}\alpha_i$ generated by the α_i ($i \in I$) is called the *root lattice* associated to A . Let $Q^+ = \sum_{i \in I} \mathbf{Z}_{\geq 0}\alpha_i$ and $Q^- = -Q^+$. The coloring matrix $C = (\theta_{i,j})$ gives rise to a complex valued mapping $\theta : Q \times Q \rightarrow \mathbf{C}^\times$ satisfying

- $\theta(\alpha_i, \alpha_j) = \theta_{i,j}$,
- $\theta(\alpha, \beta + \gamma) = \theta(\alpha, \beta)\theta(\alpha, \gamma)$,
- $\theta(\alpha + \beta, \gamma) = \theta(\alpha, \gamma)\theta(\beta, \gamma)$,

for all $\alpha, \beta, \gamma \in Q$.

We define the binomial coefficients by:

$$\{n\}_{q_i} = \frac{\theta_{i,i}^n q_i^n - q_i^{-n}}{\theta_{i,i} q_i - q_i^{-1}}, \quad \{n\}_{q_i}! = \prod_{t=1}^n \{t\}_{q_i}, \quad \text{and} \quad \{m\}_{q_i} = \frac{\{m\}_{q_i}!}{\{n\}_{q_i}! \{m-n\}_{q_i}!},$$

where $\{0\}_{q_i}! = 1$ and $q_i = q^{s_i}$.

We let $\xi_i = q_i - q_i^{-1}$ and $K_i = q^{s_i h_i}$.

Definition 2.1 ([2]). Suppose $\mathfrak{g} = \mathfrak{g}(A, \underline{m}, C)$ is the Borcherds superalgebra of charge \underline{m} determined by the symmetrizable integral Borcherds-Cartan matrix A which is colored by a coloring matrix C . Let q be an indeterminate. Then the *quantized Borcherds superalgebra* $U_q(\mathfrak{g})$ associated to \mathfrak{g} is the associative algebra

over $\mathbf{C}(q)$ with 1, generated by the elements q^h ($h \in P^\vee$), $e_{i,k}$, $f_{i,k}$ ($i \in I, k = 1, 2, \dots, m_i$) with the defining relations:

- (R1) $q^0 = 1, q^h q^{h'} = q^{h+h'}$ for $h, h' \in P^\vee$,
- (R2) $q^h e_{i,k} q^{-h} = q^{\alpha_i(h)} e_{i,k}$ for $h \in P^\vee, i \in I, k = 1, 2, \dots, m_i$,
- (R3) $q^h f_{i,k} q^{-h} = q^{-\alpha_i(h)} f_{i,k}$ for $h \in P^\vee, i \in I, k = 1, 2, \dots, m_i$,
- (R4) $e_{i,k} f_{j,l} - \theta_{j,i} f_{j,l} e_{i,k} = \delta_{i,j} \delta_{k,l} \frac{1}{\xi_i} (K_i - K_i^{-1})$,

for $i, j \in I, k = 1, 2, \dots, m_i, l = 1, 2, \dots, m_j$,

$$(R5) \sum_{n=0}^{1-a_{i,j}} (-1)^n \theta_{i,j}^n \theta_{i,i}^{n(n-1)/2} \left\{ \begin{matrix} 1-a_{i,j} \\ n \end{matrix} \right\}_{q_i} e_{i,k}^{1-a_{i,j}-n} e_{j,l} e_{i,k}^n = 0$$

if $a_{i,i} = 2$ and $i \neq j$,

$$(R6) \sum_{n=0}^{1-a_{i,j}} (-1)^n \theta_{i,j}^n \theta_{i,i}^{n(n-1)/2} \left\{ \begin{matrix} 1-a_{i,j} \\ n \end{matrix} \right\}_{q_i} f_{i,k}^{1-a_{i,j}-n} f_{j,l} f_{i,k}^n = 0$$

if $a_{i,i} = 2$ and $i \neq j$,

$$(R7) e_{i,k} e_{j,l} - \theta_{i,j} e_{j,l} e_{i,k} = 0 \quad \text{if } a_{i,j} = 0,$$

$$(R8) f_{i,k} f_{j,l} - \theta_{i,j} f_{j,l} f_{i,k} = 0 \quad \text{if } a_{i,j} = 0.$$

Proposition 2.2 ([2]). *The algebra $U_q(\mathfrak{g})$ has a Hopf algebra structure with comultiplication Δ , counit ε , and antipode S defined by:*

$$(2.1) \quad \Delta(q^h) = q^h \otimes q^h,$$

$$(2.2) \quad \Delta(e_{i,k}) = e_{i,k} \otimes 1 + K_i \otimes e_{i,k},$$

$$(2.3) \quad \Delta(f_{i,k}) = f_{i,k} \otimes K_i^{-1} + 1 \otimes f_{i,k},$$

$$(2.4) \quad \varepsilon(q^h) = 1,$$

$$(2.5) \quad \varepsilon(e_{i,k}) = 0,$$

$$(2.6) \quad \varepsilon(f_{i,k}) = 0,$$

$$(2.7) \quad S(q^h) = q^{-h},$$

$$(2.8) \quad S(e_{i,k}) = -K_i^{-1} e_{i,k},$$

$$(2.9) \quad S(f_{i,k}) = -f_{i,k} K_i,$$

for $h \in P^\vee, i \in I, k = 1, 2, \dots, m_i$.

We denote by U^0 the subalgebra of $U = U_q(\mathfrak{g})$ generated by q^h for $h \in P^\vee$ and U^+ (respectively, U^-) the subalgebra of U generated by the elements $e_{i,k}$ (respectively, $f_{i,k}$) for $i \in I, k = 1, 2, \dots, m_i$. We also denote by $U^{\geq 0}$ (respectively, $U^{\leq 0}$) the subalgebra of U generated by the elements q^h and $e_{i,k}$ (respectively, $f_{i,k}$) for $h \in P^\vee, i \in I, k = 1, 2, \dots, m_i$. For each $\beta \in Q$, let

$$(2.10) \quad U_\beta = \{x \in U \mid q^h x q^{-h} = q^{\beta(h)} x \text{ for all } h \in P^\vee\}.$$

We similarly define $U_{\pm\beta}^\pm, U_{\pm\beta}^{\geq 0}$, and $U_{\pm\beta}^{\leq 0}$ for $\beta \in Q^+$. We then have:

Proposition 2.3 ([2]).

- (a) $U \cong U^- \otimes U^0 \otimes U^+$.
- (b) $U^0 = \bigoplus_{h \in P^\vee} \mathbf{C} q^h$.
- (c) $U^\pm = \bigoplus_{\beta \in Q^+} U_{\pm\beta}^\pm$.

(d) (R5) and (R7) (respectively, (R6) and (R8)) are the fundamental relations for U^+ (respectively, U^-).

We give Q^+ a partial ordering by setting $\lambda \geq \mu$ if and only if $\lambda - \mu \in Q^+$. We will also use the notation $K_\gamma = \prod K_i^{n_i}$ for $\gamma = \sum n_i \alpha_i \in Q$.

3. REPRESENTATIONS OF $U_q(\mathfrak{g})$

For $i \in I$ define the \mathbf{C} -linear functionals $\Lambda_i \in \mathfrak{h}^*$ by:

$$(3.1) \quad \Lambda_i(h_j) = \delta_{i,j}, \quad \Lambda_i(d_j) = 0, \quad \text{for all } j \in I.$$

Define the lattices:

$$(3.2) \quad P = \{\lambda \in \mathfrak{h}^* \mid \lambda(h_i), \lambda(d_i) \in \mathbf{Z}, \forall i \in I\},$$

$$(3.3) \quad \bar{P} = \left(\bigoplus_{i \in I} \mathbf{Z} \alpha_i \right) \oplus \left(\bigoplus_{i \in I} \mathbf{Z} \Lambda_i \right).$$

P is called the *weight lattice* of \mathfrak{g} . An element $\lambda \in P$ is said to be a *dominant integral weight* if

$$(3.4) \quad \lambda(h_i) \in \mathbf{Z}_{\geq 0} \text{ for all } i \in I^{\text{re}},$$

$$(3.5) \quad \lambda(h_i) \in 2\mathbf{Z}_{\geq 0} \text{ for all } i \in I^{\text{re}} \cap I^{\text{odd}},$$

where I^{odd} denotes the set of $i \in I$ such that $\theta_{i,i} = -1$. Let P^+ denote the set of all dominant integral weights.

Set $\bar{\mathfrak{h}}^* = \mathbf{C} \otimes_{\mathbf{Z}} \bar{P}$. Then the nondegenerate symmetric bilinear form on \mathfrak{h} gives an isomorphism between \mathfrak{h} and $\bar{\mathfrak{h}}^*$ hence also induces a bilinear form on $\bar{\mathfrak{h}}^*$. We may extend this bilinear form to a symmetric bilinear form on \mathfrak{h}^* . We extend it so that it satisfies $(\lambda \mid \alpha_i) = \lambda(s_i h_i)$ and $(\lambda \mid \Lambda_i) = \lambda(s_i d_i)$ for every $\lambda \in \mathfrak{h}^*$. Write $\lambda \perp \mu$ if $(\lambda \mid \mu) = 0$.

For each $i \in I$ such that $a_{i,i} \neq 0$, we define the *simple reflection* $r_i \in \text{GL}(\mathfrak{h}^*)$ on \mathfrak{h}^* by

$$(3.6) \quad r_i(\lambda) = \lambda - \frac{2}{a_{i,i}} \lambda(h_i) \alpha_i.$$

The subgroup W of $\text{GL}(\mathfrak{h}^*)$ generated by r_i ($i \in I^{\text{re}}$) is called the *Weyl group* of $\mathfrak{g}(A, \underline{m}, C)$. We denote by $l : W \rightarrow \mathbf{Z}_{\geq 0}$ the natural length function.

Let R be the family of all imaginary simple roots, each root occurring as many times as its multiplicity, i.e. m_i times for α_i . For $\lambda \in P^+$, define $R(\lambda)$ to be the set of all $\mu = \sum_{j=1}^r \alpha_{i_j} + \sum_{k=1}^s l_{i_k} \beta_{i_k} \in Q^+$, where α_{i_j} (resp. β_{i_k}) are distinct even (resp. odd) roots in R , satisfying

- $\alpha_{i_j} \perp \lambda, \beta_{i_k} \perp \lambda$ for all j, k ,
- $\alpha_{i_j} \perp \beta_{i_k}$ for all j, k ,
- $\alpha_{i_j} \perp \alpha_{i_k}, \beta_{i_j} \perp \beta_{i_k}$ for $j \neq k$,
- $\beta_{i_k} \perp \beta_{i_k}$ if $l_{i_k} \geq 2$.

In particular, $0 \in R(\lambda)$. Suppose $\rho \in \mathfrak{h}^*$ satisfies $\rho(h_i) = \frac{1}{2} a_{i,i}$ for all $i \in I$.

Proposition 3.1 ([16, 17]). *Let $\lambda \in P^+$. Denote by $M^q(\lambda)$ the Verma module for $U_q(\mathfrak{g})$ with highest weight λ and let $V^q(\lambda)$ be the irreducible highest weight module over $U_q(\mathfrak{g})$ with highest weight λ . Then,*

$$(3.7) \quad \text{ch } M^q(\lambda) = \frac{e^\lambda}{\prod_{\alpha \in \Phi^-} (1 - \theta(\alpha, \alpha) e^{\alpha})^{\theta(\alpha, \alpha) \dim \mathfrak{g}_\alpha}} = e^\lambda \sum_{\beta \in Q^+} (\dim U_{-\beta}^-) e^{-\beta},$$

$$(3.8) \quad \text{ch } V^q(\lambda) = \frac{\sum_{w \in W, \mu \in R(\lambda)} (-1)^{l(w) + \text{ht}(\mu)} e^{w(\lambda + \rho - \mu) - \rho}}{\prod_{\alpha \in \Phi^-} (1 - \theta(\alpha, \alpha) e^\alpha)^{\theta(\alpha, \alpha) \dim \mathfrak{g}_\alpha}}.$$

In this formula, Φ^- is the set of all negative roots.

The following is a Corollary to this proposition.

Lemma 3.2. *Let $\gamma = \sum_{i \in I} n_i \alpha_i \in Q^+$. Suppose $\lambda \in P^+$, $\lambda(h_i) > 0$ for all $i \in I^{\text{im}}$ and $\lambda(h_i) \geq n_i$ for all $i \in I^{\text{re}}$. Then we have a linear isomorphism $U_{-\gamma}^- \xrightarrow{\sim} V^q(\lambda)_{\lambda-\gamma}$ given by $u \mapsto uv_\lambda$.*

Proof. $U_{-\gamma}^- \rightarrow M^q(\lambda)_{\lambda-\gamma}$ is surjective, so $U_{-\gamma}^- \rightarrow V^q(\lambda)_{\lambda-\gamma}$ is also surjective. Hence it suffices to show $\dim U_{-\gamma}^- = \dim V^q(\lambda)_{\lambda-\gamma}$. Since $(\alpha_i | \lambda) = \lambda(s_i h_i) = s_i \lambda(h_i) > 0$ for all $i \in I^{\text{im}}$, no nonempty subset F of R satisfies $F \perp \lambda$, and so

$$(3.9) \quad \text{ch } V^q(\lambda) = \frac{\sum_{w \in W} (-1)^{l(w)} e^{w(\lambda + \rho) - \rho}}{\prod_{\alpha \in \Phi^-} (1 - \theta(\alpha, \alpha) e^\alpha)^{\theta(\alpha, \alpha) \dim \mathfrak{g}_\alpha}}$$

$$(3.10) \quad = \left(\sum_{w \in W} (-1)^{l(w)} e^{w(\lambda + \rho) - \rho} \right) \left(\sum_{\beta \in Q^+} (\dim U_{-\beta}^-) e^{-\beta} \right)$$

Therefore, it suffices to show that if $w(\lambda + \rho) - \rho - \beta = \lambda - \gamma$ for some $w \in W$, $\beta \in Q^+$, then $w = 1$.

We will show that if $w \neq 1$, then $\gamma + w(\lambda + \rho) - (\lambda + \rho) \notin Q^+$ by using induction on the length of w .

If $w = r_i$ ($i \in I^{\text{re}}$), then

$$\begin{aligned} & \gamma + r_i(\lambda + \rho) - (\lambda + \rho) \\ &= \gamma + \lambda + \rho - (\lambda(h_i) + \rho(h_i))\alpha_i - (\lambda + \rho) \\ &= \gamma - (\lambda(h_i) + 1)\alpha_i \notin Q^+. \end{aligned}$$

If $w = w' r_i$ ($i \in I^{\text{re}}$) with $l(w) = l(w') + 1$, then

$$\begin{aligned} & \gamma + w(\lambda + \rho) - (\lambda + \rho) \\ &= \gamma + w' r_i(\lambda + \rho) - (\lambda + \rho) \\ &= \gamma + w'(\lambda + \rho - (\lambda(h_i) + \rho(h_i))\alpha_i) - (\lambda + \rho) \\ &= (\gamma + w'(\lambda + \rho) - (\lambda + \rho)) - (\lambda(h_i) + 1)w'\alpha_i \notin Q^+. \end{aligned}$$

This completes the proof. \square

4. THE BILINEAR FORM ON $U_q(\mathfrak{g})$

4.1. The bilinear form on $U^{\geq 0} \times U^{\leq 0}$. In this section, we define a bilinear form on $U^{\geq 0} \times U^{\leq 0}$ which is nondegenerate when restricted to $U_\beta^+ \times U_{-\beta}^-$, $\beta \in Q^+$.

For $\phi \in (U_\beta)^*$, $\psi \in (U_\gamma)^*$, $x \in U_\beta$, and $y \in U_\gamma$, we define $(\phi \otimes \psi)(x \otimes y) = \theta(-\gamma, \beta)\phi(x)\psi(y)$. With this, and the Hopf algebra structure on $U_q(\mathfrak{g})$, we can give an algebra structure to $\bigoplus_{\alpha \in Q^+} (U_\alpha^{\geq 0})^*$ by setting $(\phi_1 \phi_2)(x) = (\phi_1 \otimes \phi_2)(\Delta(x))$ for $\phi_1, \phi_2 \in \bigoplus_{\alpha \in Q^+} (U_\alpha^{\geq 0})^*$ and $x \in U^{\geq 0}$. For $h \in P^\vee$ and $i \in I$, $k = 1, 2, \dots, m_i$, we

define the linear functionals $\phi_h, \psi_{i,k} \in \bigoplus_{\alpha \in Q^+} (U^{\geq 0}_\alpha)^*$ by

$$(4.1) \quad \phi_h(xq^{h'}) = \varepsilon(x)q^{-(h|h')} \quad (x \in U^+, h' \in P^\vee),$$

$$(4.2) \quad \psi_{i,k}(xq^h) = 0 \quad (x \in U_\beta^+, \beta \in Q^+ \setminus \{\alpha_i\}),$$

$$(4.3) \quad \psi_{i,k}(e_{i,l}q^h) = \delta_{k,l}.$$

Proposition 4.1. *There exists an algebra homomorphism*

$$(4.4) \quad \zeta : U^{\leq 0} \longrightarrow \bigoplus_{\alpha \in Q^+} (U^{\geq 0}_\alpha)^*$$

given by

$$(4.5) \quad \zeta(q^h) = \phi_h \quad (h \in P^\vee),$$

$$(4.6) \quad \zeta(f_{i,k}) = -\frac{1}{\xi_i} \psi_{i,k} \quad (i \in I, k = 1, 2, \dots, m_i).$$

Proof. By Proposition 2.3, we have only to check that the relations (R1), (R3), (R6), and (R8) are preserved under the map ζ . Other cases being easy, we just sketch the (R6) part.

Define $e_{i,k}^{(n)} = e_{i,k}^n / \{n\}_{q_i}!$. We may check by induction on n that

$$\Delta(e_{i,k}^{(n)}) = \sum_{s+t=n} q_i^{st} e_{i,k}^{(s)} K_i^t \otimes e_{i,k}^{(t)}.$$

This show

$$((\Delta \otimes 1) \circ \Delta)(e_{i,k}^{(n)}) = \sum_{r+s+t=n} q_i^{r s + s t + t r} e_{i,k}^{(r)} K_i^{s+t} \otimes e_{i,k}^{(s)} K_i^t \otimes e_{i,k}^{(t)}.$$

We again use induction to prove

$$\psi_{i,k}^n(e_{i,k}^{(n)}) = (\theta_{i,i} q_i)^{\frac{n(n-1)}{2}}$$

With this, it is possible to show

$$\psi_{i,k}^{N-n} \psi_{j,l} \psi_{i,k}^n (e_{i,k}^{(N-m)} e_{j,l} e_{i,k}^{(m)}) = \sum \theta_{i,i}^{A'} \theta_{i,j}^{B'} q_i^{C'} \left\{ \begin{matrix} N-n \\ \alpha \end{matrix} \right\}_{q_i} \left\{ \begin{matrix} n \\ \beta \end{matrix} \right\}_{q_i},$$

with the summation over nonnegative integers $\alpha, \beta, \gamma, \delta$ such that $\alpha + \beta = N - m$, $\gamma + \delta = m$, $\alpha + \gamma = N - n$, and $\beta + \delta = n$ and where

$$A' = \beta\gamma + (N - n)n + \frac{1}{2}(N - n)(N - n - 1) + \frac{1}{2}n(n - 1),$$

$$B' = \beta - \gamma + (N - n) - n,$$

$$C' = \alpha\beta + \gamma\delta + 2\beta\gamma + (\beta + \gamma)a_{i,j} + \frac{1}{2}(N - n)(N - n - 1) + \frac{1}{2}n(n - 1).$$

Noting

$$\left\{ \begin{matrix} N \\ n \end{matrix} \right\}_{q_i} \left\{ \begin{matrix} N-n \\ \alpha \end{matrix} \right\}_{q_i} \left\{ \begin{matrix} n \\ N-m-\alpha \end{matrix} \right\}_{q_i} = \left\{ \begin{matrix} N \\ m \end{matrix} \right\}_{q_i} \left\{ \begin{matrix} N-m \\ \beta \end{matrix} \right\}_{q_i} \left\{ \begin{matrix} m \\ \delta \end{matrix} \right\}_{q_i},$$

we can calculate

$$\begin{aligned} & \left(\sum_{n=0}^{1-a_{i,j}} (-1)^n \theta_{i,j}^n \theta_{i,i}^{n(n-1)/2} \begin{Bmatrix} 1-a_{i,j} \\ n \end{Bmatrix}_{q_i} \psi_{i,k}^{1-a_{i,j}-n} \psi_{j,l} \psi_{i,k}^n \right) (e_{i,k}^{(N-m)} e_{j,l} e_{i,k}^{(m)}) \\ &= \sum_{n=0}^N \sum \theta_{i,i}^A \theta_{i,j}^B q_i^C \begin{Bmatrix} N \\ m \end{Bmatrix}_{q_i} \begin{Bmatrix} N-m \\ \beta \end{Bmatrix}_{q_i} \begin{Bmatrix} m \\ \delta \end{Bmatrix}_{q_i} \end{aligned}$$

with the second summation over nonnegative integers satisfying the same conditions as before and where

$$\begin{aligned} A &= \frac{1}{2}N(N-1) + m\beta + \frac{1}{2}\beta(\beta-1) + \frac{1}{2}\delta(\delta-1), \\ B &= N-m, \\ C &= (m-mN + \frac{1}{2}N(N-1)) + (m+1-N)\beta + (m-1)\delta. \end{aligned}$$

This can be written as a product of two sums which simplifies to zero. \square

Define a bilinear form $(\mid) : U^{\geq 0} \times U^{\leq 0} \rightarrow \mathbf{C}(q)$ by

$$(4.7) \quad (x|y) = \zeta(y)(x) \quad (x \in U^{\geq 0}, y \in U^{\leq 0}).$$

For $n \in \mathbf{Z}_{>0}$, we denote by $\Delta_n : U \rightarrow U^{\otimes(n+1)}$, the algebra homomorphism defined by $\Delta_1 = \Delta$, $\Delta_n = (\Delta \otimes 1) \circ \Delta_{n-1}$, and we write

$$(4.8) \quad \Delta_n(x) = \sum_{(x)_n} x_{(0)} \otimes x_{(1)} \otimes \cdots \otimes x_{(n)}.$$

For homogeneous elements $x_i \in U_{\beta_i}^{\geq 0}$, $y_i \in U_{-\gamma_i}^{\leq 0}$ ($i = 1, 2$), we define $(x_1 \otimes x_2 | y_1 \otimes y_2) = \theta(\beta_2, -\gamma_1)(x_1|y_1)(x_2|y_2)$ and extend it by linearity. For $x \in U_\beta$, $y \in U_\gamma$, we will write $\theta(x, y)$ to mean $\theta(\beta, \gamma)$ and define $P : U \otimes U \rightarrow U \otimes U$ by $P(x \otimes y) = \theta(x, y)y \otimes x$ on homogeneous elements and extend it by linearity.

Proposition 4.2. *The bilinear form (\mid) on $U^{\geq 0} \times U^{\leq 0}$ defined by (4.7) satisfies:*

$$(4.9) \quad (x|y_1 y_2) = (\Delta(x)|y_1 \otimes y_2) \quad (x \in U^{\geq 0}, y_1, y_2 \in U^{\leq 0}),$$

$$(4.10) \quad (x_1 x_2 | y) = (P(x_1 \otimes x_2) | \Delta(y)) \quad (x_1, x_2 \in U^{\geq 0}, y \in U^{\leq 0}),$$

$$(4.11) \quad (q^h | q^{h'}) = q^{-(h|h')} \quad (h, h' \in P^\vee),$$

$$(4.12) \quad (q^h | f_{i,k}) = 0,$$

$$(4.13) \quad (e_{i,k} | q^h) = 0,$$

$$(4.14) \quad (e_{i,k} | f_{j,l}) = -\frac{1}{\xi_i} \delta_{i,j} \delta_{k,l}$$

for $i, j \in I$, $k = 1, 2, \dots, m_j$.

Moreover, the bilinear form on $U^{\geq 0} \times U^{\leq 0}$ satisfying the above equations is unique.

Proof. Everything including uniqueness is straightforward except for (4.10). It is proved by induction. Here we just show the induction part. We suppress the summation signs for simplicity. Assume $(x^1 x^2 | y^i) = \theta(x^1, x^2)(x^2 \otimes x^1 | \Delta(y^i))$ for $i = 1, 2$. Then,

$$\begin{aligned} (x^1 x^2 | y^1 y^2) &= \zeta(y^1) \zeta(y^2) (x^1 x^2) \\ &= \zeta(y^1) \otimes \zeta(y^2) ((x_{(0)}^1 \otimes x_{(1)}^1) \cdot (x_{(0)}^2 \otimes x_{(1)}^2)) \end{aligned}$$

$$\begin{aligned}
&= \zeta(y^1) \otimes \zeta(y^2)(\theta(x_{(1)}^1, x_{(0)}^2)x_{(0)}^1x_{(0)}^2 \otimes x_{(1)}^1x_{(1)}^2) \\
&= \theta(x_{(1)}^1, x_{(0)}^2)\theta(y^2, x_{(0)}^1)\theta(y^2, x_{(0)}^2)(x_{(0)}^1x_{(0)}^2|y^1)(x_{(1)}^1x_{(1)}^2|y^2) \\
&= \theta(x_{(0)}^1, x_{(1)}^1)\theta(x_{(0)}^1, x_{(1)}^2)\theta(x_{(0)}^2, x_{(1)}^2)\theta(x_{(0)}^1, x_{(0)}^2)\theta(x_{(1)}^1, x_{(1)}^2) \\
&\quad \cdot (x_{(0)}^2 \otimes x_{(0)}^1|y_{(0)}^1 \otimes y_{(1)}^1)(x_{(1)}^2 \otimes x_{(1)}^1|y_{(0)}^2 \otimes y_{(1)}^2) \\
&= \theta(x_{(0)}^1, x_{(1)}^1)\theta(x_{(0)}^1, x_{(1)}^2)\theta(x_{(0)}^2, x_{(1)}^2) \\
&\quad \cdot (x_{(0)}^2|y_{(0)}^1)(x_{(0)}^1|y_{(1)}^1)(x_{(1)}^2|y_{(0)}^2)(x_{(1)}^1|y_{(1)}^2) \\
&= \theta(y_{(1)}^1, y_{(0)}^2)(\Delta(x^2)|y_{(0)}^1 \otimes y_{(0)}^2)(\Delta(x^1)|y_{(1)}^1 \otimes y_{(1)}^2) \\
&= \theta(y_{(1)}^1, y_{(0)}^2)\theta(y_{(0)}^1y_{(0)}^2, x^1)(x^2 \otimes x^1|y_{(0)}^1y_{(0)}^2 \otimes y_{(1)}^1y_{(1)}^2) \\
&= \theta(x^1, x^2)(x^2 \otimes x^1|(y_{(0)}^1 \otimes y_{(1)}^1) \cdot (y_{(0)}^2 \otimes y_{(1)}^2)) \\
&= \theta(x^1, x^2)(x^2 \otimes x^1|\Delta(y^1y^2)).
\end{aligned}$$

This completes the proof. \square

Lemma 4.3.

- (a) $(S(x)|S(y)) = (x|y)$
- (b) $(xq^h|yq^{h'}) = q^{-(h|h')}(x|y) \quad (h, h' \in P^\vee, x \in U^+, y \in U^-)$.
- (c) $(U_\beta^+|U_{-\gamma}^-) = 0 \quad \text{if } \gamma \neq \beta$.

Proof. To prove (a), we set $(|)' = (S(|)S(|))$ and show $(|)'$ satisfies conditions of Proposition 4.2. The remaining two are easy. \square

Lemma 4.4. For $x \in U^{\geq 0}$, $y \in U^{\leq 0}$, that are homogeneous, we have

$$(4.15) \quad \theta(x, y)yx = \sum_{(x)_2, (y)_2} \Theta_{xy}(x_{(0)}|S(y_{(0)}))(x_{(2)}|y_{(2)})x_{(1)}y_{(1)}$$

and

$$(4.16) \quad xy = \sum_{(x)_2, (y)_2} \theta(x_{(1)}, y_{(1)})\Theta_{xy}(x_{(0)}|y_{(0)})(x_{(2)}|S(y_{(2)}))y_{(1)}x_{(1)}$$

with $\Theta_{xy} = \theta(x_{(1)}, y_{(0)})\theta(x_{(2)}, y_{(0)})\theta(x_{(2)}, y_{(1)})$.

Proof. By substituting (4.15) into the right hand side of (4.16), we can show that (4.15) implies (4.16).

To prove (4.15), we use induction on y and reduce the problem to showing this true for $y = q^h$ and $y = f_{i,k}$. The case $y = q^h$ is easy. The case $y = f_{i,k}$ turns out to be equivalent to showing

$$\begin{aligned}
(4.17) \quad \theta(x, f_{i,k})f_{i,k}x &= \sum_{x_{(1)}} \left\{ (x_{(1)}|f_{i,k})x_{(0)} \right. \\
&\quad \left. + \theta(x_{(1)}, f_{i,k})(x_{(1)}|K_i^{-1})x_{(0)}f_{i,k} \right. \\
&\quad \left. - \theta(x_{(1)}, f_{i,k})(x_{(0)}|f_{i,k})K_i^{-1}x_{(1)} \right\},
\end{aligned}$$

which is proved by induction on the length of x . \square

Lemma 4.5. Let $\beta \in Q^+ \setminus \{0\}$ and $y \in U_{-\beta}^-$. If $e_{i,k}y = \theta(\alpha_i, -\beta)y e_{i,k}$ for all $i \in I$, $k = 1, 2, \dots, m_i$, then $y = 0$.

Proof. Choose $\lambda \in P^+$ satisfying the assumptions of Lemma 3.2. Since $e_{i,k}(y \cdot v_\lambda) = \theta(\alpha_i, -\beta)y(e_{i,k} \cdot v_\lambda) = 0$ for all $i \in I$, $k = 1, 2, \dots, m_i$, and $\text{wt}(y \cdot v_\lambda) = \lambda - \beta \preceq \lambda$, $y \cdot v_\lambda$ generates a proper submodule of $V^q(\lambda)$. Hence $y \cdot v_\lambda = 0$. Lemma 3.2 now says $y = 0$. \square

Theorem 4.6. *For $\beta \in Q^+$, the bilinear form $(\cdot | \cdot) : U^{\geq 0} \times U^{\leq 0} \rightarrow \mathbf{C}(q)$ defined by (4.7) is nondegenerate when restricted to $U_\beta^+ \times U_{-\beta}^-$.*

Proof. Since $\dim U_\beta^+ = \dim U_{-\beta}^-$, nondegeneracy on one side implies the nondegeneracy on the other side. So we will just prove the statement

$$(4.18) \quad \text{if } y \in U_{-\beta}^-, \text{ and } (U_\beta^+ | y) = 0 \text{ then } y = 0.$$

We use induction on β .

The case $\beta = 0$ or α_i is easy.

Assume (4.18) is true for all $\gamma < \beta$ with $\beta \in Q^+ \setminus (\{0\} \cup \{\alpha_i\}_{i \in I})$. Recall the notation $K_\gamma = q^{\sum_{i \in I} n_i s_i h_i}$ for $\gamma = \sum_{i \in I} n_i \alpha_i \in Q$. By definition of Δ , we see that

$$(4.19) \quad \Delta(y) = \sum_{0 \leq \gamma \leq \beta} y_\gamma (1 \otimes K_{-\gamma}), \quad y_\gamma \in U_{-\gamma}^- \otimes U_{-(\beta-\gamma)}^-$$

with $y_0 = 1 \otimes y$ and $y_\beta = y \otimes 1$. Fix $0 < \gamma < \beta$. For any $u \in U_{\beta-\gamma}^+$ and $v \in U_\gamma^+$, we have

$$(4.20) \quad (v \otimes u | y_\gamma) = (v \otimes u | y_\gamma (1 \otimes K_{-\gamma}))$$

$$(4.21) \quad = (v \otimes u | \Delta(y))$$

$$(4.22) \quad = \theta(\gamma, \beta - \gamma)(uv | y) \quad \text{by (4.10)}$$

$$(4.23) \quad = 0.$$

Hence $(U_\gamma^+ \otimes U_{\beta-\gamma}^+ | y_\gamma) = 0$. This implies $y_\gamma = 0$ by our induction hypothesis. Therefore $\Delta(y) = y \otimes K_{-\beta} + 1 \otimes y$. We apply Lemma 4.4 to

$$(4.24) \quad \Delta_2(e_{i,k}) = e_{i,k} \otimes 1 \otimes 1 + K_i \otimes e_{i,k} \otimes 1 + K_i \otimes K_i \otimes e_{i,k}$$

$$(4.25) \quad \Delta_2(y) = y \otimes K_{-\beta} \otimes K_{-\beta} + 1 \otimes y \otimes K_{-\beta} + 1 \otimes 1 \otimes y$$

and get

$$(4.26) \quad \theta(\alpha_i, -\beta)y e_{i,k} = e_{i,k}y \quad \text{for all } i \in I.$$

Hence $y = 0$ by Lemma 4.5. \square

4.2. The Killing form. Recall from Proposition 2.3 that $U \cong U^+ \otimes U^0 \otimes S(U^-) \cong U^- \otimes U^0 \otimes S(U^+)$. Using the bilinear form defined in the previous section, we define a new bilinear form

$$\langle \cdot | \cdot \rangle : U \times U \rightarrow \mathbf{C}(q^{\frac{1}{2}})$$

by setting,

$$(4.27) \quad \langle x_1 q^{h_1} S(y_1) | y_2 q^{h_2} S(x_2) \rangle = (x_1 | y_2)(x_2 | y_1) q^{-(h_1 | h_2)/2} \theta(y_1, y_2) \theta(y_1, x_2)$$

for homogeneous $x_i \in U^+$, $y_i \in U^-$, $h_i \in P^\vee$ and extending by linearity.

For homogeneous $u, v \in U$, we define

$$(4.28) \quad \text{ad}(u) \cdot v = \sum_{(u)_1} \theta(u_{(1)}, v) u_{(0)} v S(u_{(1)})$$

$$(4.29) \quad v \cdot \widetilde{\text{ad}}(u) = \sum_{(u)_1} \theta(v, u_{(0)}) S(u_0) v u_{(1)}.$$

It is easy to check that these define left and right actions of U on U .

The bilinear form on U defined above is *invariant* in that:

Proposition 4.7. For $u, v, v' \in U$,

$$(4.30) \quad \langle \text{ad}(u) \cdot v | v' \rangle = \langle v | v' \cdot \widetilde{\text{ad}}(u) \rangle \theta(u, v) \theta(u, v').$$

Proof. It suffices to check the formula for $u = q^{h''}$ ($h'' \in P^\vee$), $e_{i,k}$, $f_{i,k}$ ($i \in I, k = 1, 2, \dots, m_i$) and for $v = xq^h S(y)$ and $v' = y'q^{h'} S(x')$ with $x \in U_\beta^+$, $x' \in U_{\beta'}^+$, $y \in U_{-\gamma}^-$, $y' \in U_{-\gamma'}^-$ ($\beta, \beta', \gamma, \gamma' \in Q^+$). Since the case $u = f_{i,k}$ is similar to the case $u = e_{i,k}$, we will omit the case $u = f_{i,k}$.

(i) $u = q^{h''}$

The left hand side is

$$\langle \text{ad}(u) \cdot v | v' \rangle = \langle q^{h''} v q^{-h''} | v' \rangle = q^{(\beta-\gamma)(h'')} \langle v | v' \rangle,$$

and the right hand side is

$$\langle v | v' \cdot \widetilde{\text{ad}}(u) \rangle = \langle v | q^{-h''} v' q^{h''} \rangle = q^{(\gamma'-\beta')(h'')} \langle v | v' \rangle.$$

Since $\langle v | v' \rangle \neq 0$ only when $\beta = \gamma'$ and $\beta' = \gamma$, we are done.

(ii) $u = e_{i,k}$

Applying Lemma 4.4, we obtain

$$\begin{aligned} \text{ad}(u) \cdot v &= e_{i,k} x q^h S(y) + \theta(\alpha_i, \beta) q^{(\alpha_i|\beta)} x K_i q^h S(e_{i,k} y) \\ &= e_{i,k} x q^h S(y) + \theta(\alpha_i, \beta) q^{(\alpha_i|\beta)} \sum_{(y)_2} \{A - B + C\}, \end{aligned}$$

where

$$\begin{aligned} A &= (e_{i,k} | y_{(0)}) (1 | S(y_{(2)})) x K_i q^h S(y_{(1)}), \\ B &= q^{\alpha_i(h)} (K_i | y_{(0)}) (1 | S(y_{(2)})) x e_{i,k} q^h S(y_{(1)}), \\ C &= \theta(e_{i,k}, y_{(1)}) (K_i | y_{(0)}) (e_{i,k} | S(y_{(2)})) x q^h S(y_{(1)}). \end{aligned}$$

and

$$\begin{aligned} v' \cdot \widetilde{\text{ad}}(u) &= -\theta(\beta' - \gamma', \alpha_i) q^{(\gamma' - \alpha_i|\alpha_i)} e_{i,k} y' K_i^{-1} q^{h'} S(x') \\ &\quad - \theta(\beta', \alpha_i) q^{(\gamma' - \beta'|\alpha_i)} y' q^{h'} S(e_{i,k} x') \\ &= -\theta(\beta' - \gamma', \alpha_i) q^{(\gamma' - \alpha_i|\alpha_i)} \sum_{(y')_2} \{A' - B' + C'\} \\ &\quad - \theta(\beta', \alpha_i) q^{(\gamma' - \beta'|\alpha_i)} y' q^{h'} S(e_{i,k} x'), \end{aligned}$$

where

$$\begin{aligned} A' &= (e_{i,k}|y'_{(0)})(1|S(y'_{(2)}))y'_{(1)}K_i^{-1}q^{h'}S(x'), \\ B' &= \theta(\alpha_i, \beta' - \gamma')q^{(\alpha_i|\alpha_i)}q^{-\alpha_i(h')}(K_i|y'_{(0)})(1|S(y'_{(2)}))y'_{(1)}q^{h'}S(x'e_{i,k}), \\ C' &= \theta(e_{i,k}, y'_{(1)})(K_i|y'_{(0)})(e_{i,k}|S(y'_{(2)}))y'_{(1)}q^{h'}S(x'). \end{aligned}$$

There are only two cases to consider.

- $\gamma' = \beta + \alpha_i$ and $\gamma = \beta'$,
- $\gamma' = \beta$ and $\gamma = \beta' + \alpha_i$.

Since the latter case is similar to the former, we will only check the first case. Assume $\gamma' = \beta + \alpha_i$ and $\gamma = \beta'$. Then, in order to have $B \neq 0$, we must have $y_{(0)}, y_{(2)} \in U^0$ and $y_{(1)} \in U_{-\gamma'}^-$. Similarly, $A' \neq 0$ implies $y'_{(0)} \in U_{-\alpha_i}^{\leq 0}$, $y'_{(1)} \in U_{-\beta}^{\leq 0}$, and $y'_{(2)} \in U^0$. In this case, we get $y'_{(1)} = \tilde{y}'_1 K_i^{-1}$ for some $\tilde{y}'_1 \in U_{-\beta}^-$. Also $C' \neq 0$ implies $y'_{(0)} \in U^0$, $y'_{(1)} \in U_{-\beta}^{\leq 0}$, and $y'_{(2)} \in U_{-\alpha_i}^{\leq 0}$. In this case, we have $y'_{(2)} = \tilde{y}'_2 K_{\gamma' - \alpha_i}^{-1}$ for some $\tilde{y}'_2 \in U_{-\alpha_i}^-$. We need to prepare one more fact. Using Proposition 4.2, we obtain the following formula.

$$\begin{aligned} (x_1 x_2 x_3 | y) &= \sum_{(y)_2} \theta(x_1 x_2, x_3) \theta(x_1 x_2, y_{(0)}) \theta(x_1, x_2) \theta(x_1, y_{(1)}) \\ &\quad \times (x_3 | y_{(0)}) (x_2 | y_{(1)}) (x_1 | y_{(2)}) \end{aligned}$$

for any $x_i \in U^+$ ($i = 1, 2, 3$) and $y \in U^-$. From this formula, we get

$$\begin{aligned} (x' | y) &= (x' K_i | y) = \sum_{(y)_2} (K_i | y_{(0)}) (x' | y_{(1)}) (1 | y_{(2)}), \\ (x e_{i,k} | y') &= \sum_{(y')_2} \theta(x, e_{i,k}) \theta(x, y'_{(0)}) (e_{i,k} | y'_{(0)}) (x | y'_{(1)}) (1 | y'_{(2)}), \\ (e_{i,k} x | y') &= (e_{i,k} x K_i | y') \\ &= \sum_{(y')_2} \theta(e_{i,k}, x) \theta(e_{i,k}, y'_{(1)}) (K_i | y'_{(0)}) (x | y'_{(1)}) (e_{i,k} | y'_{(2)}). \end{aligned}$$

Now, we obtain

$$\begin{aligned} \langle \text{ad}(u) \cdot v | v' \rangle &= \langle e_{i,k} x q^h S(y) | y' q^{h'} S(x') \rangle \\ &\quad - \theta(\alpha_i, \beta) q^{(\alpha_i|\beta)} q^{\alpha_i(h)} \sum_{(y)_2} (K_i | y_{(0)}) \langle x e_{i,k} q^h S(y_{(1)}) | y' q^{h'} S(x') \rangle \\ &= \theta(\gamma, \gamma' - \beta') q^{-(h|h')/2} \\ &\quad \times \left\{ (e_{i,k} x | y') (x' | y) - \theta(\alpha_i, \beta) q^{(\alpha_i|\beta)} q^{\alpha_i(h)} \sum_{(y)_2} (K_i | y_{(0)}) (x e_{i,k} | y') (x' | y_{(1)}) \right\} \\ &= \theta(\gamma, \gamma' - \beta') q^{-(h|h')/2} \\ &\quad \times \sum_{(y)_2, (y')_2} (K_i | y_{(0)}) (x' | y_{(1)}) (x | y'_{(1)}) \\ &\quad \times \left\{ (K_i | y'_{(0)}) (e_{i,k} | y'_{(2)}) - \theta(x, y'_{(0)}) q^{(\alpha_i|\beta)} q^{\alpha_i(h)} (e_{i,k} | y'_{(0)}) \right\} \end{aligned}$$

and

$$\langle v | v' \cdot \widetilde{\text{ad}}(u) \rangle = -\theta(\beta' - \gamma', \alpha_i) q^{(\gamma' - \alpha_i|\alpha_i)}$$

$$\begin{aligned}
& \times \sum_{(y')_2} \left\{ (e_{i,k}|y'_{(0)}) \langle xq^h S(y)|y'_{(1)} K_i^{-1} q^{h'} S(x') \rangle \right. \\
& \quad \left. + \theta(e_{i,k}, y'_{(1)})(K_i|y'_{(0)})(e_{i,k}|S(y'_{(2)})) \langle xq^h S(y)|y'_{(1)} q^{h'} S(x') \rangle \right\} \\
& = \theta(\beta' - \gamma', \alpha_i) \theta(\gamma, \beta - \beta') q^{-(h|h')/2} q^{(\beta|\alpha_i)} \\
& \times \sum_{(y')_2} \left\{ \theta(e_{i,k}, y'_{(1)})(K_i|y'_{(0)})(e_{i,k} K_i^{-1} |\tilde{y}'_2 K_{\gamma' - \alpha_i}^{-1})(x|y'_{(1)})(x'|y) \right. \\
& \quad \left. - (e_{i,k}|y'_{(0)})(x|\tilde{y}'_1)(x'|y) q^{\alpha_i(h)} \right\} \\
& = \theta(\beta' - \gamma', \alpha_i) \theta(\gamma, \beta - \beta') q^{-(h|h')/2} \\
& \times \sum_{(y)_2, (y')_2} (K_i|y_{(0)})(x'|y_{(1)})(x|y'_{(1)}) \\
& \quad \times \left\{ \theta(e_{i,k}, y'_{(1)})(K_i|y'_{(0)})(e_{i,k}|y'_{(2)}) - q^{(\alpha_i|\beta)} q^{\alpha_i(h)} (e_{i,k}|y'_{(0)}) \right\}.
\end{aligned}$$

Comparing these two, we get the desired formula. \square

This proposition allows us to define a right U -module structure on some subalgebra of U^* . Define $\zeta : U \rightarrow U^*$ by setting

$$(4.31) \quad [\zeta(u)](v) = \langle v|u \rangle$$

for $u, v \in U$. Here, the dual space on the right should be viewed as the set of linear maps from U to $\mathbf{C}(q^{\frac{1}{2}})$. For $\zeta(u) \in \zeta(U)$, $x \in U$, define $\zeta(u) \cdot x$ by,

$$[\zeta(u) \cdot x](v) = \theta(u, x) \theta(v, x) [\zeta(u)](\text{ad}(x) \cdot v).$$

Proposition 4.7 allows us to check $\zeta(u) \cdot x = \zeta(u \cdot \widetilde{\text{ad}}(x))$. So this gives a right U -module structure on $\zeta(U)$ and $\zeta : U \rightarrow \zeta(U)$ becomes a U -module homomorphism.

Proposition 4.8. *The bilinear form $\langle | \rangle$ is nondegenerate. Hence, the map ζ is injective.*

Proof. Let $u \in U_{-\alpha}^- U^0 S(U_{\beta}^+)$ with $\langle v|u \rangle = 0$ for all $v \in U_{\alpha}^+ U^0 S(U_{-\beta}^-)$. It suffices to show $u = 0$. For each $\gamma \in Q^+ - \{0\}$, choose a basis $\{u_i^{\gamma}\}_i$ of U_{γ}^+ . And let $\{v_i^{\gamma}\}_i$ be a basis of $U_{-\gamma}^-$ dual to $\{u_i^{\gamma}\}_i$ with respect to the nondegenerate bilinear form $\langle | \rangle$. Notice that the elements $u_i^{\alpha} q^h S(v_j^{\beta})$ with $h \in P^{\vee}$ and i, j going over appropriate indices, form a basis for $U_{\alpha}^+ U^0 U_{-\beta}^-$. Similarly, the elements $v_k^{\alpha} q^{h'} S(u_l^{\beta})$ form a basis for $U_{-\alpha}^- U^0 U_{\beta}^+$. Writing $u = \sum_{k,h,l} a_{k,h,l} v_k^{\alpha} q^{h'} S(u_l^{\beta})$ with $a_{k,h,l} \in \mathbf{C}(q)$, and using

$$\langle u_i^{\alpha} q^h S(v_j^{\beta}) | v_k^{\alpha} q^{h'} S(u_l^{\beta}) \rangle = \delta_{i,k} \delta_{j,l} q^{-(h|h')/2} \theta(\beta, \alpha) \theta(\beta, \beta),$$

we arrive at,

$$\sum_{h' \in P^{\vee}} a_{k,h',l} q^{-(h|h')/2} = 0$$

for each k, l , and $h \in P^{\vee}$. Now, each map $h \mapsto q^{-(h|h')/2}$ is a group homomorphism from P^{\vee} to the multiplicative group $\mathbf{C}(q^{\frac{1}{2}})^{\times}$. Since $q^{\frac{1}{2}}$ is not a root of unity, distinct h' produces distinct homomorphisms. So, by Artin's Theorem on linear independence of characters, every $a_{k,h',l} = 0$. We have $u = 0$ as claimed. \square

5. HARISH-CHANDRA HOMOMORPHISM

We denote the center of U by \mathfrak{z} . For each $i \in I$ with $a_{i,i} \neq 0$, define the *simple reflection* $r_i \in \mathrm{GL}(\mathfrak{h})$ by,

$$r_i(h) = h - \frac{2}{a_{i,i}} \alpha_i(h) h_i,$$

and let $\widetilde{W} = \langle r_i \mid i \in I, a_{i,i} \neq 0 \rangle \subset \mathrm{GL}(\mathfrak{h})$. Let $(U^0)^{\widetilde{W}}$ be the subspace of U^0 consisting of the elements $\sum_{h \in P^\vee} c_h q^h$ ($c_h \in \mathbf{C}(q)$) such that $c_h \neq 0$ implies $w(h) \in P^\vee$ and $c_{w(h)} = c_h$ for any $w \in \widetilde{W}$.

We define an algebra automorphism $\phi : U^0 \rightarrow U^0$ by setting $\phi(q^h) = q^{-\rho(h)} q^h$ for $h \in P^\vee$. The *Harish-Chandra homomorphism* $\xi : \mathfrak{z} \rightarrow U^0$ is the restriction to \mathfrak{z} of the map

$$U \xrightarrow{\sim} U^- \otimes U^0 \otimes U^+ \xrightarrow{\varepsilon \otimes 1 \otimes \varepsilon} U^0 \xrightarrow{\phi} U^0.$$

For later use, we define the algebra homomorphism $\chi_\lambda : U^0 \rightarrow \mathbf{C}(q)$ for each $\lambda \in P^+$ by $\chi_\lambda(q^h) = q^{\lambda(h)}$.

Proposition 5.1.

- (a) ξ is an algebra homomorphism.
- (b) ξ is injective.

Proof. We will just prove (b). Let $z \in \mathfrak{z}$ be such that $\xi(z) = 0$. Writing $z = \sum_{\beta \in Q^+} z_\beta$ with $z_\beta \in U_{-\beta}^- U^0 U_\beta^+$, we see that $z_0 = 0$. Fix any $\beta \in Q^+$ minimal with the property that $z_\beta \neq 0$. Also choose basis $\{y_r\}_r$ and $\{x_s\}_s$ of $U_{-\beta}^-$ and U_β^+ , respectively. We may write $z_\beta = \sum_{r,s} y_r u_{r,s} x_s$ for some $u_{r,s} \in U^0$. Then,

$$\begin{aligned} 0 = e_{i,k} z - z e_{i,k} &= \sum_{\gamma \neq \beta} (e_{i,k} z_\gamma - z_\gamma e_{i,k}) \\ &+ \sum_{r,s} (e_{i,k} y_r - \theta(\alpha_i, -\beta) y_r e_{i,k}) u_{r,s} x_s \\ &+ \sum_{r,s} y_r (\theta(\alpha_i, -\beta) e_{i,k} u_{r,s} x_s - u_{r,s} x_s e_{i,k}) \end{aligned}$$

Recalling the minimality of β , we see that only the second term on the right belongs to $U_{-(\gamma-\alpha_i)}^- U^0 U_\gamma^+$. So we have $\sum_{r,s} (e_{i,k} y_r - \theta(\alpha_i, -\beta) y_r e_{i,k}) u_{r,s} x_s = 0$. $\{x_s\}_s$ was chosen to be a basis, so $e_{i,k} \sum_r y_r u_{r,s} = \theta(\alpha_i, -\beta) \sum_r y_r e_{i,k} u_{r,s}$ for all $i \in I$ and s .

Let $v_\lambda \in V^q(\lambda)$ denote the highest weight vector. Set $v = \sum_r \chi_\lambda(u_{r,s}) y_r v_\lambda$. Then $e_{i,k} v = \theta(\alpha_i, -\beta) \sum_r y_r e_{i,k} u_{r,s} v_\lambda = 0$ for all $i \in I$, so the irreducibility of $V^q(\lambda)$ says $v = 0$. Choosing an appropriate $\lambda \in P^+$, we may use Lemma 3.2 and say $\sum_r \chi_\lambda(u_{r,s}) y_r = 0$. Again, $\{y_r\}_r$ was a basis, so $\chi_\lambda(u_{r,s}) = 0$ for all r, s . By choosing a suitable set of λ , we may show $u_{r,s} = 0$ for all r, s and we have $z_\beta = 0$. This contradicts the choice of z_β . \square

We now try to close in on the image of ξ . For each $J \subset \{(i, k) \mid i \in I, k = 1, 2, \dots, m_i\}$, let $U_J = \langle e_{i,k}, f_{i,k}, U^0 \mid (i, k) \in J \rangle$. We denote by \mathfrak{z}_J the center of the algebra U_J and by $\xi_J : \mathfrak{z}_J \rightarrow U^0$ the Harish-Chandra homomorphism for U_J . Let U_J^+ (respectively, U_J^-) be the subalgebra of U_J generated by $e_{i,k}$ (respectively

$f_{i,k}$) with $(i, k) \in J$, and set

$$(5.1) \quad R_J^+ = \{x \in U^+ \mid (x|U_J^-) = 0\} = \{x \in U^+ \mid (x|U_J^- U^0) = 0\}$$

$$(5.2) \quad R_J^- = \{y \in U^- \mid (U_J^+|y) = 0\} = \{y \in U^- \mid (U^0 U_J^+|y) = 0\}$$

$$(5.3) \quad R_J = R_J^- U^0 U^+ + U^- U^0 R_J^+$$

The following may be proved as in [15].

Lemma 5.2.

- (a) $U = U_J \oplus R_J$,
- (b) $U_J R_J U_J \subset R_J$,
- (c) $(\varepsilon \otimes 1 \otimes \varepsilon)(R_J) = 0$.

Define $U_r^0 = \bigoplus_h \mathbf{C}(q)q^h$, where the direct sum is over all $h \in P^\vee$ satisfying,

- $\alpha_i(h) \in s_i a_{i,i} \mathbf{Z}$ if $i \in I^{\text{ev}}$,
- $\alpha_i(h) \in 2s_i a_{i,i} \mathbf{Z}$ if $i \in I^{\text{odd}}$ and $a_{i,i} \neq 0$.

Proposition 5.3.

- (a) $\text{Im}(\xi) \subset (U^0)^{\widetilde{W}}$.
- (b) $\text{Im}(\xi) \subset U_r^0$.
- (c) $\text{Im}(\xi) \subset \text{Im}(\xi_J)$.

Proof. (a) Let $z \in \mathfrak{z}$. Let $v_\lambda \in M^q(\lambda)$ be the highest weight vector. Then, $z v_\lambda = \chi_{\lambda+\rho}(\xi(z)) v_\lambda$. Since z commutes with every element of U , z acts as $\chi_{\lambda+\rho}(\xi(z))$ on every element of $M^q(\lambda)$. Now, fix $i \in I$ such that $a_{i,i} \neq 0$. We may calculate

$$e_{i,k} f_{i,k}^n = \theta_{i,i}^n f_{i,k}^n e_{i,k} + \theta_{i,i}^{n-1} f_{i,k}^{n-1} \frac{1}{\xi_i} \left(\frac{1 - \theta_{i,i}^{-n} q_i^{-n a_{i,i}}}{1 - \theta_{i,i}^{-1} q_i^{-a_{i,i}}} K_i - \frac{1 - \theta_{i,i}^n q_i^{n a_{i,i}}}{1 - \theta_{i,i} q_i^{a_{i,i}}} K_i^{-1} \right).$$

So that, for each $\lambda \in P$ satisfying $n(\lambda) := \frac{2}{a_{i,i}} \lambda(h_i) \in \mathbf{Z}_{\geq 0}$, we can check that $f_{i,k}^{n(\lambda)+1} v_\lambda$ is a highest weight vector. Its weight is

$$\begin{aligned} \lambda - \left(\frac{2}{a_{i,i}} \lambda(h_i) + 1 \right) \alpha_i &= \lambda - \frac{2}{a_{i,i}} (\lambda + \rho)(h_i) \alpha_i \\ &= r_i (\lambda + \rho) - \rho. \end{aligned}$$

The argument at the beginning of this proof applies to any highest weight vector and we have,

$$\chi_{\lambda+\rho}(\xi(z)) = \chi_{r_i(\lambda+\rho)}(\xi(z))$$

under the condition $\frac{2}{a_{i,i}} \lambda(h_i) \in \mathbf{Z}$. Checking $\chi_{r_i \mu}(q^h) = \chi_\mu(r_i q^h)$, for any $\mu \in \mathfrak{h}^*$, the above may now be written as

$$\chi_{\lambda+\rho}(\xi(z) - r_i \xi(z)) = 0$$

for every $\lambda \in P$ satisfying $\lambda(h_i) \in \frac{a_{i,i}}{2} \mathbf{Z}_{\geq 0}$. By choosing a suitable set of λ , we may show $\xi(z) = r_i \xi(z)$.

(b) Let $z = \sum_{\beta \in Q^+} z_\beta \in \mathfrak{z}$ with $z_\beta \in U_{-\beta}^- U^0 U_\beta^+$. Set $x = \sum_{n=0}^{\infty} z_{n\alpha_i}$ and $y = z - x$. Then $z = x + y$ with $x \in U_{\{(i,1)\}}$ and $y \in R_{\{(i,1)\}}$. Looking at

$$0 = e_{i,k} z - z e_{i,k} = (e_{i,k} x - x e_{i,k}) + (e_{i,k} y - y e_{i,k})$$

with Lemma 5.2 in mind, we see that $x \in \mathfrak{z}_{\{(i,1)\}}$. By the results of Section 6.1, all of which may be obtained by direct calculation, we have,

- $z_0 \in \langle K_i, q^h \mid \alpha_i(h) = 0 \rangle$ if $i \in I^{\text{ev}}$,
- $z_0 \in \langle K_i^2, q^h \mid \alpha_i(h) = 0 \rangle$ if $i \in I^{\text{odd}}$ and $a_{i,i} \neq 0$.

The result follows.

(c) For $z \in \mathfrak{z}$, write $z = x + y$ with $x \in U_J$ and $y \in R_J$. As in the proof for (b), we may show $x \in \mathfrak{z}_J$. So we have, $\xi(z) = \xi(x) + \xi(y) = \xi(x) = \xi_J(x) \in \text{Im}(\xi_J)$. \square

6. THE CENTER OF $U_q(\mathfrak{g})$

6.1. Rank 1. In this section, we list the center for the case when the index set is of size 1. All results may be obtained by direct calculation using induction after choosing a suitable basis of $U \cong U^- \otimes U^0 \otimes U^+$.

If $a_{i,i} \neq 0$, $\theta_{i,i} = 1$, define

$$C_i = f_{i,1}e_{i,1} + \frac{1}{\xi_i} \left(\frac{1}{1 - q^{-s_i a_{i,i}}} K_i - \frac{1}{1 - q^{s_i a_{i,i}}} K_i^{-1} \right).$$

If $a_{i,i} \neq 0$, $\theta_{i,i} = -1$, define

$$\begin{aligned} C_i &= f_{i,1}^2 e_{i,1}^2 + \frac{1}{\xi_i} f_{i,1} \left(\frac{1 - q^{s_i a_{i,i}}}{1 + q^{s_i a_{i,i}}} K_i - \frac{1 - q^{-s_i a_{i,i}}}{1 + q^{-s_i a_{i,i}}} K_i^{-1} \right) e_{i,1} \\ &\quad - \frac{1}{\xi_i^2} \left\{ \frac{1}{(1 + q^{-s_i a_{i,i}})^2} K_i^2 + \frac{1}{(1 + q^{s_i a_{i,i}})^2} K_i^{-2} \right\}. \end{aligned}$$

If $h \in P^\vee$ satisfy $\alpha_i(h) \neq 0$, define

$$C_{ih} = f_{i,1} q^h e_{i,1} + \frac{1}{\xi_i} \frac{1}{1 - q^{-\alpha_i(h)}} q^h (K_i - K_i^{-1}).$$

Proposition 6.1.

- If $J = \{(i, 1)\}$ and $a_{i,i} \neq 0$,
then $\mathfrak{z}_J = \langle C_i, q^h \mid h \in P^\vee, \alpha_i(h) = 0 \rangle$.
- If $J = \{(i, 1)\}$, $a_{i,i} = 0$, and $\theta_{i,i} = 1$,
then $\mathfrak{z}_J = \langle q^h \mid h \in P^\vee, \alpha_i(h) = 0 \rangle \subset U^0$.
- If $J = \{(i, 1)\}$, $a_{i,i} = 0$, and $\theta_{i,i} = -1$,
then $\mathfrak{z}_J = \langle C_{ih}, q^{h'} \mid h, h' \in P^\vee, \alpha_i(h) \neq 0, \alpha_i(h') = 0 \rangle$.

6.2. Finite type. In this section, we give a structure theorem for the center of $U_q(\mathfrak{g})$ when the Borchers-Cartan matrix is of finite type. We take the Borchers-Cartan matrix to be of finite type throughout this section. To simplify arguments, we redefine

$$\begin{aligned} P^\vee &= \bigoplus_{i \in I} \mathbf{Z}h_i, \\ \mathfrak{h} &= \bigoplus_{i \in I} \mathbf{C}h_i \end{aligned}$$

for this section. Notice that the bilinear form (\mid) is still nondegenerate on the redefined \mathfrak{h} .

The irreducible highest weight module has a natural grading.

$$V^q(\lambda) = \bigoplus_{\alpha \in Q^+} V^q(\lambda)_{\lambda - \alpha}.$$

Define a map $\eta \in \text{End}(V^q(\lambda))$ by setting $\eta(v) = \theta(\alpha, \alpha)v$ for $v \in V^q(\lambda)_{\lambda - \alpha}$. When the Borchers-Cartan matrix A is of finite type, it is known([12]) that the irreducible

highest weight module $V(\lambda)$ over $\mathfrak{g}(A)$ is finite dimensional for $\lambda \in P^+$. Since the classical limit([2]) of $V^q(\lambda)$ is $V(\lambda)$, $V^q(\lambda)$ is also of finite dimension when $\lambda \in P^+$. So we may define the *supertrace* for $x \in U_q(\mathfrak{g})$ acting on $V^q(\lambda)$ by

$$(6.1) \quad \text{str}(x; V^q(\lambda)) = \text{tr}(\eta \circ x; V^q(\lambda)).$$

For homogeneous elements $x, y \in U$, we can easily check

$$(6.2) \quad \text{str}(xy) = \theta(x, y) \text{str}(yx).$$

Lemma 6.2. $u \in \mathfrak{z}$ if and only if $u \cdot \widetilde{\text{ad}}(x) = \varepsilon(x)u$ for all $x \in U$.

Proof. Let $u \in \mathfrak{z}$. Then, $u \in U_0$ and

$$\begin{aligned} u \cdot \widetilde{\text{ad}}(x) &= \sum_{(x)_1} S(x_{(0)})ux_{(1)} \\ &= u \sum_{(x)_1} S(x_{(0)})x_{(1)} = \varepsilon(x)u. \end{aligned}$$

Conversely, if $u \cdot \widetilde{\text{ad}}(x) = \varepsilon(x)u$ for all $x \in U$,

$$q^{-h}uq^h = u \cdot \widetilde{\text{ad}}(x) = \varepsilon(q^h)u = u.$$

So $u \in U_0$ and we have,

$$0 = \varepsilon(e_{i,k})u = u \cdot \widetilde{\text{ad}}(e_{i,k}) = -K_i^{-1}e_{i,k}u + K_i^{-1}ue_{i,k}.$$

This shows $e_{i,k}u = ue_{i,k}$. We may similarly show $f_{i,k}u = uf_{i,k}$ and hence $u \in \mathfrak{z}$. \square

For each $\lambda \in P^+$, define $f_\lambda \in U^*$ by

$$(6.3) \quad f_\lambda(u) = \text{str}(uK_{2\rho}^{-1}; V^q(\lambda)).$$

Let $\nu : \mathfrak{h} \rightarrow \mathfrak{h}^*$ denote the isomorphism given by the nondegenerate symmetric bilinear form (\mid) . Define

$$\widehat{Q} = \nu(P^\vee) = \bigoplus_{i \in I} \mathbf{Z} \frac{1}{s_i} \alpha_i.$$

Recall the map $\zeta : U \rightarrow U^*$ defined in (4.31).

Lemma 6.3. For $\lambda \in P^+$, $f_\lambda \in \text{Im}(\zeta)$ if and only if $\lambda \in \frac{1}{2}\widehat{Q}$.

Proof. From Proposition 4.8, we see that the image of ζ is the restricted dual of $U_q(\mathfrak{g})$. So

$$\text{Im}(\zeta) = \left(\bigoplus_{\beta \in Q^+} (U_{-\beta}^-)^* \right) \otimes \left(\bigoplus_{\mu \in \frac{1}{2}\widehat{Q}} \mathbf{C}(q)\chi_\mu \right) \otimes \left(\bigoplus_{\beta \in Q^+} (U_\beta^+)^* \right)$$

under the identification $U \cong U^- \otimes U^0 \otimes U^+$. The finite dimensionality of $V^q(\lambda)$ allows us to show $f_\lambda \in \text{Im}(\zeta)$ if and only if $\lambda \in \frac{1}{2}\widehat{Q}$. \square

The next proposition gives elements of the center.

Proposition 6.4. For each $\lambda \in P^+ \cap \frac{1}{2}\widehat{Q}$, we have $z_\lambda := \zeta^{-1}(f_\lambda) \in \mathfrak{z}$.

Proof. Recall from the theory of finite dimensional simple Lie algebras, that ρ may be written as a half sum of positive roots. Since the simple roots for the super case is identical to the non-super case, we have $2\rho \in Q^+$ in either case. Hence, in the notation given on page 4, $K_{2\rho}$ is a well-defined element of U^0 . Using the fact that $K_{2\rho}^{-1}xK_{2\rho} = S^2(x)$ for any $x \in U$ and using the property of supertrace given by (6.2), we have for any $u \in U$,

$$\begin{aligned}
 (f_\lambda \cdot x)(u) &= f_\lambda(\text{ad}(x) \cdot u)\theta(u, x) \\
 &= \sum_{(x)_1} \text{str}(x_{(0)}uS(x_{(1)})K_{2\rho}^{-1}; V^q(\lambda))\theta(x_{(1)}, u)\theta(u, x) \\
 &= \sum_{(x)_1} \text{str}(uS(x_{(1)})K_{2\rho}^{-1}x_{(0)}; V^q(\lambda))\theta(x_{(0)}, x_{(1)}) \\
 &= \text{str}(uS(\sum_{(x)_1} S(x_{(0)})x_{(1)})K_{2\rho}^{-1}; V^q(\lambda)) \\
 &= \varepsilon(x) \text{str}(uK_{2\rho}^{-1}; V^q(\lambda)) \\
 &= \varepsilon(x)f_\lambda(u).
 \end{aligned}$$

Thus $f_\lambda \cdot x = \varepsilon(x)f_\lambda$. Recall from Proposition 4.8 that ζ is injective, and notice

$$f_\lambda \cdot x = \zeta(\zeta^{-1}(f_\lambda)) \cdot x = \zeta(\zeta^{-1}(f_\lambda) \cdot \widetilde{\text{ad}}(x)).$$

This shows $\zeta^{-1}(f_\lambda) \cdot \widetilde{\text{ad}}(x) = \varepsilon(x)\zeta^{-1}(f_\lambda)$. From Lemma 6.2, we get $\zeta^{-1}(f_\lambda) \in \mathfrak{z}$. \square

We finally show that the above elements generate the whole center.

Theorem 6.5. *Suppose that the Borcherds-Cartan matrix $A = (a_{i,j})_{i,j \in I}$ is indecomposable and of finite type. Then, $\xi : \mathfrak{z} \rightarrow (U_r^0)^{\widetilde{W}}$ is an isomorphism.*

Proof. Let us calculate $\xi(z_\lambda)$. We extend the notation K_β introduced in page 4 to $\beta \in \widehat{Q}$ by setting $K_{\frac{1}{s_i}\alpha_i} = q^{h_i}$. We have the commutative diagram

$$\begin{array}{ccc}
 U & \xrightarrow{\zeta} & U^* \\
 \varepsilon \otimes \text{id} \otimes \varepsilon \downarrow & & \downarrow \\
 U^0 & \longrightarrow & (U^0)^*
 \end{array}$$

where the right vertical arrow is the restriction map and the lower horizontal arrow is given by $K_\mu \mapsto \chi_{-\mu/2}$. Now, as maps on U^0 ,

$$f_\lambda = \sum_{\mu \leq \lambda} \theta(\lambda - \mu, \lambda - \mu) \dim(V(\lambda)_\mu) q^{-2(\rho|\mu)} \chi_\mu.$$

This shows,

$$(6.4) \quad \xi(z_\lambda) = \sum_{\mu \leq \lambda} \theta(\lambda - \mu, \lambda - \mu) \dim(V(\lambda)_\mu) K_{-2\mu}$$

for $\lambda \in P^+ \cap \frac{1}{2}\widehat{Q}$.

Define \widehat{P} to be the set of elements $\mu \in \mathfrak{h}^*$ such that $\mu(h_i) \in \mathbf{Z}$ if $i \in I$ is even and $\mu(h_i) \in 2\mathbf{Z}$ if $i \in I$ is odd. Notice $P^+ \subset \widehat{P}$. We can now write

$$U_r^0 = \bigoplus_{\mu \in 2\widehat{P} \cap \widehat{Q}} \mathbf{C}(q)K_\mu.$$

Action of the Weyl groups \widetilde{W} and W defined on \mathfrak{h} and \mathfrak{h}^* are compatible with the isomorphism ν . By Proposition 6.4, it suffices to show that the elements $\xi(z_\lambda)$ with $\lambda \in P^+ \cap \frac{1}{2}\widehat{Q}$ generate $(U_r^0)^W$.

Set $\bar{\mu} = \sum_{w \in W} K_{-w\mu}$ for any $\mu \in \widehat{Q}$. We know that the elements $\bar{\mu}$ with $\mu \in 2P^+ \cap \widehat{Q}$ generate $(U_r^0)^W$. Let us use induction to show that each of them belong to $\text{Im}(\xi)$. The element $\bar{0} \in U_r^0$ is given by $\xi(z_\lambda)$ with $\lambda = 0$. Choose any $\lambda \in 2P^+ \cap \widehat{Q}$. Then, $\frac{1}{2}\lambda \in P^+ \cap \frac{1}{2}\widehat{Q}$ so that $z_{\frac{1}{2}\lambda}$ is an element of the center. Recall that $\text{Im}(\xi)$ is invariant under the action of W (Proposition 5.3). Using $\dim V^q(\frac{1}{2}\lambda)_{\frac{1}{2}\lambda} = 1$, we may rewrite (6.4) as

$$\xi(z_{\frac{1}{2}\lambda}) = \bar{\lambda} + \sum_{\mu} n_{\mu} \bar{\mu},$$

with $n_{\mu} \in \mathbf{Z}$ and $\frac{1}{2}\mu$ running over some set of weights of $V^q(\frac{1}{2}\lambda)$. Since all $\mu \leq \lambda$, induction hypothesis show that each $\bar{\mu}$ belong to $\text{Im}(\xi)$. Hence $\bar{\lambda} \in \text{Im}(\xi)$ and the induction step is complete. \square

6.3. Other Cases. Let 2_i , 0_i , and \ominus_i denote the fact that $a_{i,i}$ is respectively, 2, 0, and negative. We will sometimes add a \pm to these to reflect the sign of $\theta_{i,i}$. So, for example, 2_i^- implies that i is an odd real index. For $i, j \in I$, let us say \ominus_i is *connected directly* to \ominus_j if $a_{i,j} \neq 0$, where \ominus can be any one of 2, 0, or \ominus . Here are some results for the case when $|J| = 2$.

Lemma 6.6. *Assume one of the following.*

- (a) $J = \{(i, 1), (i, 2)\}$ with \ominus_i
- (b) $J = \{(i, 1), (j, 1)\}$ with 0_i^+ connected directly to 0_j^-
- (c) $J = \{(i, 1), (j, 1)\}$ with 0_i^- connected directly to 0_j^-
- (d) $J = \{(i, 1), (j, 1)\}$ with \ominus_i connected directly to 0_j^-
- (e) $J = \{(i, 1), (j, 1)\}$ with 2_i connected directly to \ominus_j
- (f) $J = \{(i, 1), (j, 1)\}$ with \ominus_i connected directly to \ominus_j

Then, $\mathfrak{z}_J \subset U^0$.

Proof. (a) and (f) may be proved as in [15, Proposition 4.5]. And (e) may be proved as in [15, Proposition 4.6]. (c) is proved by explicit calculation.

Let us prove (b) and (d) simultaneously. Let $z \in \mathfrak{z}_J$. Since it commutes with q^h for all $h \in P^\vee$, $z = \sum z_\beta$ with $z_\beta \in U_\beta^- \otimes U^0 \otimes U_\beta^+$, where the sum is over all $\beta \in \mathbf{Z}_{\geq 0}\alpha_i \oplus \mathbf{Z}_{\geq 0}\alpha_j$. Let α be maximal among those $\beta \in \mathbf{Z}_{\geq 0}\alpha_i \oplus \mathbf{Z}_{\geq 0}\alpha_j$ for which z_β is nonzero and suppose $\alpha \neq 0$. Let $\{x_\mu\}$ and $\{y_\lambda\}$ be any bases of $(U_J^+)_\alpha$ and $(U_J^-)_{-\alpha}$ respectively. We can now write

$$z = \left(\sum_{\lambda, \mu, h} c_h^{\lambda, \mu} y_\lambda q^h x_\mu \right) + z'.$$

Recall Lemma 4.4 and notice

$$\begin{aligned} \Delta_2(e_{i,k}) &= e_{i,k} \otimes 1 \otimes 1 + K_i \otimes e_{i,k} \otimes 1 + K_i \otimes K_i \otimes e_{i,k}, \\ \Delta_2(y_\lambda) &= 1 \otimes y_\lambda \otimes K_{-\alpha} + \text{“other terms”}. \end{aligned}$$

This shows that the only part of $e_{i,k}z - ze_{i,k}$ belonging to the direct sum component $U_{-\alpha}^- \otimes U^0 \otimes U_{\alpha+\alpha_i}^+$ is

$$\begin{aligned} & \left(\sum_{\lambda, \mu, h} c_h^{\lambda, \mu} \theta(\alpha_i, -\alpha) y_\lambda e_{i,k} q^h x_\mu \right) - \left(\sum_{\lambda, \mu, h} c_h^{\lambda, \mu} y_\lambda q^h x_\mu e_{i,k} \right) \\ &= \sum_{\lambda, h} y_\lambda q^h \sum_{\mu} c_h^{\lambda, \mu} \left(q^{-\alpha_i(h)} \theta(\alpha, \alpha_i) e_{i,k} x_\mu - x_\mu e_{i,k} \right). \end{aligned}$$

Hence, for each $h \in P^\vee$ and λ ,

$$\sum_{\mu} c_h^{\lambda, \mu} \left(q^{-\alpha_i(h)} \theta(\alpha, \alpha_i) e_{i,k} x_\mu - x_\mu e_{i,k} \right) = 0,$$

and the same statement with i replaced by j also holds. Now, $e_{j,1}^2 = 0$ is the only relation in U_j^+ for the case we are considering, so we may take an explicit set of monomials in $e_{i,1}$ and $e_{j,1}$ for the basis of U_α^+ and using these, we can show that the two equations cannot be simultaneously true. \square

Proposition 6.7. *Assume that A is indecomposable. Suppose that every 0_j^- is connected directly to a 0_i or a \ominus_i . If there is a nonempty subset J of $\{(i, k) \mid i \in I, k = 1, \dots, m_i\}$ such that $\mathfrak{z}_J \subset U^0$, then \mathfrak{z} is contained in U^0 .*

Proof. Let $\bar{J} = \{i \in I \mid (i, k) \in J \text{ for some } k\}$. For $i \in I$, set

$$T_i = \bigoplus_{h \in P^\vee, \alpha_i(h)=0} \mathbf{C}(q)q^h.$$

We then have $\mathfrak{z} \cap U^0 = \bigcap_{i \in I} T_i$ and similarly, $\mathfrak{z}_J \cap U^0 = \bigcap_{i \in \bar{J}} T_i$. It suffices to show $\text{Im}(\xi) \subset \bigcap_{i \in I} T_i$.

We already have $\text{Im}(\xi) \subset \text{Im}(\xi_J) \subset T_i$ for every $i \in \bar{J}$. Also if 0_i^+ , we have, $\text{Im}(\xi) \subset \text{Im}(\xi_{\{(i,1)\}}) \subset T_i$ by Proposition 6.1 (b). If 0_i^- , the conditions on the matrix shows we may use Lemma 6.6 and say $\text{Im}(\xi) \subset T_i$.

We now show that if $a_{j,j} \neq 0$ and $a_{i,j} \neq 0$, then $T_i \cap (U^0)^{\bar{W}} \subset T_j$. Let $c = \sum c_h q^h \in T_i \cap (U^0)^{\bar{W}}$. We must have $r_j c = c \in T_i$, so if $c_h \neq 0$, then $\alpha_i(h) = 0$ and $\alpha_i(r_j h) = 0$. But $\alpha_i(r_j h) = -\frac{2}{a_{j,j}} a_{j,i} \alpha_j(h)$ so $\alpha_j(h) = 0$. We have $c \in T_j$ as wanted.

Fix any $j \in I - (\bar{J} \cup \{i \mid a_{i,i} = 0\})$. By the indecomposability of A , there exists a finite sequence $i = i_0, i_1, \dots, i_n = j$ such that $i \in \bar{J} \cup \{i \mid a_{i,i} = 0\}$, $i_k \notin \bar{J} \cup \{i \mid a_{i,i} = 0\}$ for $k \geq 1$, and $a_{i_k, i_{k+1}} \neq 0$ for all k . What we have found above allows us to recursively show $\text{Im}(\xi) \subset T_{i_k}$ and in particular, $\text{Im}(\xi) \subset T_j$. \square

Proposition 6.8. *Suppose there exists some finite $J \subset I$ such that for every $j \in J$, $a_{j,j} = 2$ and for which the corresponding submatrix $A_J = (a_{i,j})_{i,j \in J}$ is indecomposable and not of finite type. Then, $\mathfrak{z}_J \subset U^0$.*


Proof. By [13, Proposition 4.9], we have $|W_J| = \infty$. Let $J' \subset J$ be such that $|W_{J'}| = \infty$ and $|W_{J''}| \lesssim \infty$ for all $J'' \subsetneq J'$. We may use Proposition 6.7 if we can show $\mathfrak{z}_{J'} \subset U^0$. Hence it suffices to show that if $h \in P^\vee$, $|W_{J'}(h)| \lesssim \infty$, then $\alpha_i(h) = 0$, for all $i \in J'$. Give partial order to \mathfrak{h} by setting $h_1 \geq h_2$ if and only if $h_1 - h_2 \in (\sum_i \mathbf{Z}_{\geq 0} h_i) + (\sum_i \mathbf{Z}_{\geq 0} d_i)$. Let $h' \in W_{J'}(h)$ be maximal with respect to this order. Then for each $i \in J'$, if $\alpha_i(h') < 0$, then $h' < r_i h'$, so $\alpha_i(h') \geq 0$ for all $i \in J'$. Set $W_{h'} = \{w \in W \mid w(h') = h'\}$. By [13, Proposition 3.12(a)], $W_{h'} = W_{J''}$

with $J'' = \{i \in J \mid \alpha_i(h') = 0\}$. If $J'' \subsetneq J'$, then $|W_{J'}(h')| = |W_{J'}/W_{h'}| = \infty$. Hence we must have $J'' = J'$ and $\alpha_i(h') = 0$ for all $i \in J'$. $\{h'\} = W_{J'}(h') = W_{J'}(h)$. So $h = h'$ and $\alpha_i(h) = 0$ for all $i \in J'$. \square

Proposition 6.9. *Let A be indecomposable, not of finite type, and $a_{i,i} = 2$ for all $i \in I$. Then, $\mathfrak{z} \subset U^0$.*

Proof. Suppose there exists some finite indecomposable submatrix which is not of finite type. Then we may use Proposition 6.7 and Proposition 6.8 to obtain the result.

If, to the contrary, every finite submatrix of A is of finite type, it must be one of the following types :



In all cases, with I naturally ordered, the matrix satisfies the following condition.

For each $i \in I$, there exists some $j > i$ such that $a_{i,j} \neq 0$, and $a_{i,k} = 0$ for $k > j$.

Let $c = \sum_h c_h q^h \in \text{Im}(\xi) \subset (U^0)^{\widetilde{W}}$. Fix $h \in P^\vee$ for which $c_h \neq 0$. We aim to show $|\widetilde{W}(h)| = \infty$ if $\alpha_j(h) \neq 0$ for some $j \in I$. We may assume that only finitely many $j \in I$ satisfy $\alpha_j(h) \neq 0$. Let $k \in I$ be the maximal of those so that $\alpha_j(h) = 0$ for all $j > k$ and $\alpha_k(h) \neq 0$. Set $i_0 = k$, and using property (6.3), recursively choose i_n so that $i_{n+1} > i_n$ and $a_{i_n, i_{n+1}} \neq 0$. Put $h_0 = h$ and $h_{n+1} = r_{i_n} h_n$. Then, h_n cannot form a closed orbit and $|\widetilde{W}(h)| = \infty$. Hence $\text{Im}(\xi) \subset \bigoplus_{h \in P^\vee, \alpha_i(h)=0} \mathbf{C}q^h$ and $\mathfrak{z} \subset U^0$. \square

We can now collect all results and state :

Theorem 6.10. *Assume that the Borcherds-Cartan matrix $A = (a_{i,j})_{i,j \in I}$ is indecomposable and not of finite type. Suppose that every 0_j^- is connected directly to a 0_i or a \ominus_i . Except for the case when $|I| = 1$ with $m_i = 1$, the center \mathfrak{z} belongs to U^0 .*

Proof. We apply Proposition 6.7 to each possible case.

If $|I| = 1$, the conditions imply either a 0_i^+ or a \ominus_i with $m_i \geq 2$. These cases may be handled by Proposition 6.1 (b) and Lemma 6.6 (a), respectively.

Now suppose $|I| \geq 2$. Proposition 6.9 does away with the case when all $a_{i,i} = 2$. If it contains a 0_i^+ , we may again use Proposition 6.1 (b). If it contains a 0_i^- but no 0_i^+ , we use Lemma 6.6 (c),(d). The only other case is covered by Lemma 6.6 (e),(f). \square

7. THE UNIVERSAL R-MATRIX

In this section, we find the universal R-matrix for the quantum group $U_q(\mathfrak{g})$.

A Hopf superalgebra (a colored Hopf algebra) H together with an element $\mathbf{R} \in H \otimes H$ is called a *quasi-triangular Hopf superalgebra* if it satisfies:

- (a) \mathbf{R} is invertible,

- (b) $\mathbf{R} \cdot \Delta(a) = \Delta'(a) \cdot \mathbf{R}$ for all $a \in H$,
- (c) $(\Delta \otimes 1)(\mathbf{R}) = \mathbf{R}_{13}\mathbf{R}_{23}$,
- (d) $(1 \otimes \Delta)(\mathbf{R}) = \mathbf{R}_{13}\mathbf{R}_{12}$,

where $\Delta' = P \circ \Delta$ with P a colored twisting map, and where \mathbf{R}_{ij} is an element of $H \otimes H \otimes H$ such that the i 'th and j 'th components are given by \mathbf{R} and the remaining component is 1. The element \mathbf{R} is called the *universal R-matrix*. It satisfies the Yang-Baxter equation

$$(7.1) \quad \mathbf{R}_{12}\mathbf{R}_{13}\mathbf{R}_{23} = \mathbf{R}_{23}\mathbf{R}_{13}\mathbf{R}_{12}.$$

A Hopf superalgebra H together with an element $C \in H \otimes H$ and an algebra homomorphism $\Phi : H \otimes H \rightarrow H \otimes H$ is called a *pre-triangular Hopf superalgebra* if it satisfies:

- (P1) C is invertible,
- (P2) $C \cdot \Delta(a) = \Phi(\Delta'(a)) \cdot C$ for all $a \in H$,
- (P3) $\Phi_{23} \circ \Phi_{13}(C_{12}) = C_{12}$,
- (P4) $\Phi_{12} \circ \Phi_{13}(C_{23}) = C_{23}$,
- (P5) $\Phi_{23}(C_{13}) \cdot C_{23} = (\Delta \otimes 1)(C)$,
- (P6) $\Phi_{12}(C_{13}) \cdot C_{12} = (1 \otimes \Delta)(C)$.

Under some conditions, it is possible to show that a pre-triangular Hopf superalgebra becomes a quasi-triangular Hopf superalgebra.

We set $U^{+,\beta} = \bigoplus_{\gamma \in Q^+, \gamma \not\leq \beta} U_\gamma^+$ for each $\beta \in Q^+$ and define the completion \widehat{U} of U by:

$$(7.2) \quad \widehat{U} = \varprojlim_{\beta} U/UU^{+,\beta}.$$

There is a natural embedding of U in \widehat{U} and there is a natural algebra structure on \widehat{U} which extends that of U under this embedding.

The completion of $U^{\otimes n}$ is similarly defined. We will write $\widehat{U} \widehat{\otimes} \widehat{U}$ for the completion of $U \otimes U$.

Define an algebra automorphism $\Phi : U \otimes U \rightarrow U \otimes U$ by

$$(7.3) \quad \Phi(q^h \otimes q^{h'}) = q^h \otimes q^{h'},$$

$$(7.4) \quad \Phi(e_{i,k} \otimes 1) = e_{i,k} \otimes K_i, \quad \Phi(1 \otimes e_{i,k}) = K_i \otimes e_{i,k},$$

$$(7.5) \quad \Phi(f_{i,k} \otimes 1) = f_{i,k} \otimes K_i^{-1}, \quad \Phi(1 \otimes f_{i,k}) = K_i^{-1} \otimes f_{i,k}.$$

It can be shown that Φ naturally extends to an algebra automorphism of $\widehat{U} \widehat{\otimes} \widehat{U}$.

We denote by $C_\beta \in U_\beta^+ \otimes U_{-\beta}^-$ the canonical element of the bilinear form $(\mid) : U_\beta^+ \times U_{-\beta}^- \rightarrow \mathbf{C}$. Define

$$(7.6) \quad C = \sum_{\beta \in Q^+} \theta(\beta, \beta) q^{(h_\beta | h_\beta)} (K_\beta^{-1} \otimes K_\beta) C_\beta \in \widehat{U} \widehat{\otimes} \widehat{U}.$$

Lemma 7.1.

- (a) $C \cdot \Delta(q^h) = \Phi(\Delta'(q^h)) \cdot C \quad (h \in P^\vee)$,
- (b) $(\Phi_{23} \circ \Phi_{13})(C_{12}) = C_{12}$,
- (c) $(\Phi_{12} \circ \Phi_{13})(C_{23}) = C_{23}$.

Proof. This is just straightforward calculation. □

Lemma 7.2. *Let $\beta \in Q^+$.*

- (a) $\sum_{\substack{\gamma, \delta \in Q^+ \\ \gamma + \delta = \beta}} C_\gamma(K_\delta \otimes 1)(S \otimes 1)(C_\delta) = \delta_{\beta, 0}$.
- (b) $\sum_{\substack{\gamma, \delta \in Q^+ \\ \gamma + \delta = \beta}} (K_\gamma \otimes 1)(S \otimes 1)(C_\gamma)C_\delta = \delta_{\beta, 0}$.
- (c) $\theta_{i,i}[1 \otimes e_{i,k}, C_{\beta+\alpha_i}] = C_\beta(e_{i,k} \otimes K_i^{-1}) - (e_{i,k} \otimes K_i)C_\beta$.
- (d) $\theta_{i,i}[f_{i,k} \otimes 1, C_{\beta+\alpha_i}] = C_\beta(K_i \otimes f_{i,k}) - (K_i^{-1} \otimes f_{i,k})C_\beta$.
- (e) $(\Delta \otimes 1)(C_\beta) = \sum_{\substack{\gamma, \delta \in Q^+ \\ \gamma + \delta = \beta}} q^{-(h_\gamma|h_\delta)}(K_\delta \otimes 1 \otimes 1)(C_\gamma)_{13}(C_\delta)_{23}$.
- (f) $(1 \otimes \Delta)(C_\beta) = \sum_{\substack{\gamma, \delta \in Q^+ \\ \gamma + \delta = \beta}} q^{-(h_\gamma|h_\delta)}(1 \otimes 1 \otimes K_{-\delta})(C_\gamma)_{13}(C_\delta)_{12}$.

Proof. Here we show the proof for (a) only. Other cases may be proved in a similar spirit.

The case $\beta = 0$ is trivial. So assume $\beta \in Q^+ \setminus \{0\}$. The left hand side is contained in $U_\beta^+ \otimes U$, so by Theorem 4.6 it suffices to show that the application of $(\cdot|w) \otimes 1$ is zero for all $w \in U_{-\beta}^-$. We may write

$$\Delta(w) = \sum_{\substack{\gamma, \delta \in Q^+ \\ \gamma + \delta = \beta}} w^{\delta, \gamma}(1 \otimes K_\delta) \quad \text{with} \quad w^{\delta, \gamma} \in U_{-\delta}^- \otimes U_{-\gamma}^-$$

and

$$w^{\delta, \gamma} = \sum_m w_{\delta, m}^{\delta, \gamma} \otimes w_{\gamma, m}^{\delta, \gamma} \quad \text{with} \quad w_{\delta, m}^{\delta, \gamma} \in U_{-\delta}^-, \quad w_{\gamma, m}^{\delta, \gamma} \in U_{-\gamma}^-.$$

We may also fix basis $\{x_r^\gamma\}_r$ and $\{y_r^\gamma\}_r$ of U_γ^+ and $U_{-\gamma}^-$, respectively, which are dual with respect to the bilinear form. Now,

$$\begin{aligned} ((|w) \otimes 1)(\text{LHS}) &= ((|w) \otimes 1) \left(\sum_{\gamma, \delta, r, s} \theta(y_r^\gamma, x_s^\delta) x_r^\gamma K_\delta S(x_s^\delta) \otimes y_r^\gamma y_s^\delta \right) \\ &= \sum_{\gamma, \delta, r, s} \theta(x_r^\gamma, K_\delta S(x_s^\delta)) \theta(-\gamma, \delta) (K_\delta S(x_s^\delta) \otimes x_r^\gamma | \Delta(w)) y_r^\gamma y_s^\delta \\ &= \sum_{\gamma, \delta, r, s, m} \theta(\delta, \gamma) (k_\delta S(x_s^\delta) | w_{\delta, m}^{\delta, \gamma}) (x_r^\gamma | w_{\gamma, m}^{\delta, \gamma} K_{-\delta}) y_r^\gamma y_s^\delta \\ &= \sum_{\gamma, \delta, m} \theta(\delta, \gamma) \left(\sum_r (x_r^\gamma | w_{\gamma, m}^{\delta, \gamma} K_{-\delta}) y_r^\gamma \right) \left(\sum_s (K_\delta S(x_s^\delta) | w_{\delta, m}^{\delta, \gamma}) y_s^\delta \right) \\ &= \sum_{\gamma, \delta, m} \theta(\delta, \gamma) \left(\sum_r (x_r^\gamma | w_{\gamma, m}^{\delta, \gamma}) y_r^\gamma \right) \left(\sum_s ((x_s^\delta | K_\delta^{-1} S^{-1}(w_{\delta, m}^{\delta, \gamma})) y_s^\delta) \right) \\ &= \sum_{\gamma, \delta, m} \theta(\delta, \gamma) w_{\gamma, m}^{\delta, \gamma} K_\delta^{-1} S^{-1}(w_{\delta, m}^{\delta, \gamma}) \\ &= S^{-1} \left(\sum_{\gamma, \delta, m} w_{\delta, m}^{\delta, \gamma} S(w_{\gamma, m}^{\delta, \gamma} K_\delta^{-1}) \right) \\ &= (S^{-1} \circ \text{m} \circ (1 \otimes S) \circ \Delta)(w) \\ &= \varepsilon(w) \\ &= 0. \end{aligned}$$

Hence the left hand side is zero when $\beta \neq 0$. □

Proposition 7.3. *Let*

$$(7.7) \quad C' = \sum_{\beta \in Q^+} \theta(\beta, \beta) q^{(h_\beta | h_\beta)} (1 \otimes K_\beta) (S \otimes 1) (C_\beta) \in \widehat{U} \widehat{\otimes} \widehat{U}.$$

Then $CC' = C'C = 1$.

Proof.

$$\begin{aligned} CC' &= \left(\sum_{\gamma \in Q^+} \theta(\gamma, \gamma) q^{(h_\gamma | h_\gamma)} (K_\gamma^{-1} \otimes K_\gamma) C_\gamma \right) \\ &\quad \cdot \left(\sum_{\delta \in Q^+} \theta(\delta, \delta) q^{(h_\delta | h_\delta)} (1 \otimes K_\delta) (S \otimes 1) (C_\delta) \right) \\ &= \sum_{\beta \in Q^+} \sum_{\substack{\gamma + \delta = \beta \\ \gamma, \delta \in Q^+}} \theta(\beta, \beta) q^{(h_\gamma | h_\gamma) + (h_\delta | h_\delta)} (K_\gamma^{-1} \otimes K_\gamma) C_\gamma (1 \otimes K_\delta) (S \otimes 1) (C_\delta) \\ &= \sum_{\beta \in Q^+} \theta(\beta, \beta) q^{(h_\beta | h_\beta)} (K_\beta^{-1} \otimes K_\beta) \sum_{\substack{\gamma + \delta = \beta \\ \gamma, \delta \in Q^+}} C_\gamma (K_\delta \otimes 1) (S \otimes 1) (C_\delta) \end{aligned}$$

We may now apply Lemma 7.2. The other part is done similarly. \square

Proposition 7.4. *We have*

$$(7.8) \quad C \cdot \Delta(e_{i,k}) = \Phi(\Delta'(e_{i,k})) \cdot C,$$

$$(7.9) \quad C \cdot \Delta(f_{i,k}) = \Phi(\Delta'(f_{i,k})) \cdot C.$$

Proof.

$$\begin{aligned} C \cdot \Delta(e_{i,k}) &= \sum_{\beta \in Q^+} \theta(\beta, \beta) q^{(h_{\beta+\alpha_i} | h_\beta)} (K_\beta^{-1} \otimes K_{\beta+\alpha_i}) C_\beta (e_{i,k} \otimes K_i^{-1}) \\ &\quad + \sum_{\beta \in Q^+} \theta(\beta, \beta) q^{(h_{\beta-\alpha_i} | h_\beta)} (K_{\beta-\alpha_i}^{-1} \otimes K_\beta) C_\beta (1 \otimes e_{i,k}) \\ \Phi(\Delta'(e_{i,k})) \cdot C &= \sum_{\beta \in Q^+} \theta(\beta, \beta) q^{(h_\beta | h_{\beta-\alpha_i})} (K_{\beta-\alpha_i}^{-1} \otimes K_\beta) (1 \otimes e_{i,k}) C_\beta \\ &\quad + \sum_{\beta \in Q^+} \theta(\beta, \beta) q^{(h_{\beta+\alpha_i} | h_\beta)} (K_\beta^{-1} \otimes K_{\beta+\alpha_i}) (e_{i,k} \otimes K_i) C_\beta \end{aligned}$$

$$\begin{aligned} &C \cdot \Delta(e_{i,k}) - \Phi(\Delta'(e_{i,k})) \cdot C \\ &= \sum_{\beta \in Q^+} \theta(\beta, \beta) q^{(h_\beta | h_{\beta+\alpha_i})} (K_\beta^{-1} \otimes K_{\beta+\alpha_i}) \\ &\quad \cdot \left\{ C_\beta (e_{i,k} \otimes K_i^{-1}) - (e_{i,k} \otimes K_i) C_\beta - \theta(\alpha_i, \alpha_i) [1 \otimes e_{i,k}, C_{\beta+\alpha_i}] \right\} \end{aligned}$$

We apply Lemma 7.2 to obtain the result. The other case is similar. \square

Proposition 7.5. *We have*

$$(7.10) \quad \Phi_{23}(C_{13}) \cdot C_{23} = (\Delta \otimes 1)(C),$$

$$(7.11) \quad \Phi_{12}(C_{13}) \cdot C_{12} = (1 \otimes \Delta)(C).$$

Proof.

$$\begin{aligned}\Phi_{23}(C_{13}) &= \sum_{\gamma \in Q^+} \theta(\gamma, \gamma) q^{(h_\gamma|h_\gamma)} (K_\gamma^{-1} \otimes K_\gamma^{-1} \otimes K_\gamma) (C_\gamma)_{13} \\ (\Phi_{23}(C_{13}))C_{23} &= \sum_{\beta \in Q^+} \theta(\beta, \beta) q^{(h_\beta|h_\beta)} (K_\beta^{-1} \otimes K_\beta^{-1} \otimes K_\beta) \\ &\quad \cdot \sum_{\substack{\gamma, \delta \in Q^+ \\ \gamma + \delta = \beta}} q^{-(h_\delta|h_\gamma)} (K_\delta \otimes 1 \otimes 1) (C_\gamma)_{13} (C_\delta)_{23} \\ (\Delta \otimes 1)(C) &= \sum_{\beta \in Q^+} \theta(\beta, \beta) q^{(h_\beta|h_\beta)} (K_\beta^{-1} \otimes K_\beta^{-1} \otimes K_\beta) (\Delta \otimes 1)(C_\beta)\end{aligned}$$

The second case is done similarly. \square

The Propositions tell us that U is almost a pre-triangular Hopf superalgebra.

Theorem 7.6. *The statements (P1) and (P2) hold in $\widehat{U} \widehat{\otimes} \widehat{U}$ and the relations (P3)–(P6) hold in $\widehat{U} \widehat{\otimes} \widehat{U} \widehat{\otimes} \widehat{U}$.*

A weight module is P -weighted if all its weights belong to P . Notice $(P|P) \subset \mathbf{Z}$. This allows us to define $\mathfrak{Z} \in \text{End}(V \otimes W)$ for any P -weighted $U_q(\mathfrak{g})$ -modules V and W by setting,

$$(7.12) \quad \mathfrak{Z}(v \otimes w) = q^{(\text{wt}(v)|\text{wt}(w))} v \otimes w$$

on homogeneous elements and extending by linearity. The map \mathfrak{Z} is certainly invertible. There is a natural action of $U \otimes U$ on $V \otimes W$ and as endomorphisms on $V \otimes W$,

$$(7.13) \quad \Phi(a \otimes b) = \mathfrak{Z} \circ (a \otimes b) \circ \mathfrak{Z}^{-1}$$

for every $a \otimes b \in U \otimes U$.

Set $\mathbf{R} = \mathfrak{Z}^{-1}C$. Then we finally have:

Theorem 7.7. *Let V_i ($i = 1, 2, 3$) be P -weighted $U_q(\mathfrak{g})$ -modules. As endomorphisms on $V_1 \otimes V_2 \otimes V_3$, when it can be defined, \mathbf{R} satisfies the Yang-Baxter equation (7.1).*

Proof. From (P5) and equation (7.13), we have

$$(7.14) \quad \mathfrak{Z}_{23} C_{13} \mathfrak{Z}_{23}^{-1} C_{23} = (\Delta \otimes 1)(C)$$

$$(7.15) \quad \mathbf{R}_{13} \mathbf{R}_{23} = \mathfrak{Z}_{13}^{-1} \mathfrak{Z}_{23}^{-1} (\Delta \otimes 1)(C)$$

Applying $P \otimes 1$ to both sides of (P5) and working as above, we get

$$(7.16) \quad \mathbf{R}_{23} \mathbf{R}_{13} = \mathfrak{Z}_{23}^{-1} \mathfrak{Z}_{13}^{-1} (\Delta' \otimes 1)(C).$$

The use of (P2) shows,

$$(7.17) \quad \mathbf{R}_{23} \mathbf{R}_{13} \mathbf{R}_{12} = \mathfrak{Z}_{23}^{-1} \mathfrak{Z}_{13}^{-1} ((\Delta' \otimes 1)(C)) \mathbf{R}_{12}$$

$$(7.18) \quad = \mathfrak{Z}_{23}^{-1} \mathfrak{Z}_{13}^{-1} ((\Delta' \otimes 1)(C)) (\mathfrak{Z}^{-1}C \otimes 1)$$

$$(7.19) \quad = \mathfrak{Z}_{23}^{-1} \mathfrak{Z}_{13}^{-1} (\mathfrak{Z}^{-1}C \otimes 1) (\Delta \otimes 1)(C)$$

Now, the \mathfrak{Z}_{ij} commute with each other and (P3) with (7.13) says C_{12} commutes with $\mathfrak{Z}_{13}^{-1}\mathfrak{Z}_{23}^{-1}$, so we may use (7.15) to write

$$(7.20) \quad \mathfrak{Z}_{23}^{-1}\mathfrak{Z}_{13}^{-1}(\mathfrak{Z}^{-1}C \otimes 1)(\Delta \otimes 1)(C) = \mathbf{R}_{12}\mathbf{R}_{13}\mathbf{R}_{23}.$$

Putting things together, we have the result. □

Acknowledgments I would like to thank Professor Seok-Jin Kang and one of my colleagues, Kyu-Hwan Lee, for help throughout this work. Lee has contributed much in finding the structure theorem for the center of finite type cases.

REFERENCES

[1] P. Baumann, *On the center of quantized enveloping algebras*, J. Algebra **203** (1998), no. 1, 244–260.
 [2] G. Benkart, S.-J. Kang, and D. J. Melville, *Quantized enveloping algebras for Borcherds superalgebras*, Trans. Amer. Math. Soc. **350** (1998), no. 8, 3297–3319.
 [3] R. E. Borcherds, *Generalized Kac-Moody algebras*, J. Algebra **115** (1988), no. 2, 501–512.
 [4] ———, *Monstrous moonshine and monstrous Lie superalgebras*, Invent. Math. **109** (1992), no. 2, 405–444.
 [5] V. G. Drinfel'd, *Hopf algebras and the quantum Yang-Baxter equation*, Soviet Math. Dokl. **32** (1985), no. 1, 254–258.
 [6] ———, *Quantum groups*, Proceedings of the International Congress of Mathematicians, vol. 1,2 (Berkeley, Calif., 1986), Amer. Math. Soc., Providence, RI, 1987, pp. 798–820.
 [7] ———, *Almost cocommutative Hopf algebras*, Leningrad Math. J. **1** (1990), no. 2, 321–342.
 [8] P. I. Etingof, *Central elements for quantum affine algebras and affine Macdonald's operators*, Math. Res. Lett. **2** (1995), no. 5, 611–628.
 [9] M. D. Gould, R. B. Zhang, and A. J. Bracken, *Quantum double construction for graded Hopf algebras*, Bull. Austral. Math. Soc. **47** (1993), 353–375.
 [10] M. Jimbo, *A q-difference analogue of U(g) and the Yang-Baxter equation*, Lett. Math. Phys. **10** (1985), no. 1, 63–69.
 [11] A. Joseph and G. Letzter, *Separation of variables for quantized enveloping algebras*, Amer. J. Math. **116** (1994), no. 1, 127–177.
 [12] V. G. Kac, *Lie superalgebras*, Adv. Math. **26** (1977), 8–96.
 [13] ———, *Infinite dimensional Lie algebras*, third ed., Cambridge University Press, 1990.
 [14] S.-J. Kang, *Quantum deformations of generalized Kac-Moody algebras and their modules*, J. Algebra **175** (1995), no. 3, 1041–1066.
 [15] S.-J. Kang and T. Tanisaki, *Universal R-matrices and the center of the quantum generalized Kac-Moody algebras*, Hiroshima Math. J. **27** (1997), no. 2, 347–360.
 [16] M. Miyamoto, *A generalization of Borcherds algebra and denominator formula*, J. Algebra **180** (1996), 631–651.
 [17] U. Ray, *A character formula for generalized Kac-Moody superalgebras*, J. Algebra **177** (1995), 154–163.
 [18] N. Yu. Reshetikhin, *Quasitriangle Hopf algebras and invariants of links*, Leningrad Math. J. **1** (1990), no. 2, 291–513.
 [19] N. Yu. Reshetikhin, L. A. Takhtadzhyan, and L. D. Faddeev, *Quantization of Lie groups and Lie algebras*, Leningrad Math. J. **1** (1990), no. 1, 193–225.
 [20] M. Rosso, *Analogues de la forme Killing et du théorème d'Harish-Chandra pour les groupes quantiques*, Ann. Sci. École. Norm. Sup. (4) **23** (1990), no. 3, 445–467.
 [21] T. Tanisaki, *Killing forms, Harish-Chandra isomorphisms, and universal R-matrix for quantum algebras*, Inter. J. Mod. Phys. A **Vol. 7, Supp. 1B** (1992), 941–961.

DEPARTMENT OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 151-742, KOREA
 E-mail address: jhong@math.snu.ac.kr