ON THE QUANTUM COHOMOLOGY OF BLOW-UPS OF PROJECTIVE SPACES ALONG LINEAR SUBSPACES.

MARCO MAGGESI

ABSTRACT. We give an explicit presentation with generators and relations of the quantum cohomology ring of the blow-up of a projective space along a linear subspace.

1. INTRODUCTION.

Let \mathbb{P}^m be the complex projective space, $\Lambda \subset \mathbb{P}^m$ a linear subspace of dimension p and $\alpha : \mathbb{P}^m \to \mathbb{P}^m$ the blow-up of \mathbb{P}^m along Λ . Let k be the hyperplane class on \mathbb{P}^m and η the exceptional divisor on \mathbb{P}^m . The aim of the present paper is to show a way to compute the quantum cohomology ring $H_Q(\mathbb{P}^m)$ of \mathbb{P}^m . The (classical) cohomology ring of \mathbb{P}^m can be expressed as (compare with lemma 2.6)

(1)
$$\mathrm{H}^*(\widetilde{\mathbb{P}}^m) = \frac{\mathbb{Z}[k,\eta]}{(g_1,g_2)},$$

where the two relations g_1, g_2 are

(2) $g_1(k,\eta) = (k-\eta)^{m-p}, \qquad g_2(k,\eta) = k^{p+1}\eta.$

The main result of this work is the following

Theorem 1.1 (Main theorem). Suppose that 2p + 3 < m. Then the quantum cohomology ring of $\widetilde{\mathbb{P}}^m$ can be expressed as

(3)
$$\mathrm{H}_{Q}^{*}(\widetilde{\mathbb{P}}^{m}) = \frac{\mathbb{Z}[k,\eta]}{(\tilde{g}_{1},\tilde{g}_{2})}$$

where the two relations g_1, g_2 are

(4)
$$\tilde{g}_1(k,\eta) = (k-\eta)^{m-p} - \eta, \qquad \tilde{g}_2(k,\eta) = k^{p+1}\eta - 1.$$

The blow-up \mathbb{P}^m can be regarded as a projectivized vector bundle on a projective space (theorem 2.1). The quantum cohomology of projectivized bundles on projective spaces was studied by Qin and Ruan in [9]. The relevant material from their work will be enclosed here for completeness. Other studies on quantum cohomology of blow-ups of projective spaces can be found in [7], [5] and [6].

I wish to thank professor V. Ancona for many useful discussions and for his encouragement.

2. $\widetilde{\mathbb{P}}^m$ as projective bundle.

Let n := m - p - 1, r := p + 2 and V be the rank r vector bundle on \mathbb{P}^n given by

(5)
$$V := \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus r-1} \oplus \mathcal{O}_{\mathbb{P}^n}(2)$$

and consider the associate projectivized vector bundle $\pi : \mathbb{P}(V) \to \mathbb{P}^n$.

Theorem 2.1. The two varieties $\widetilde{\mathbb{P}}^m$ and $\mathbb{P}(V)$ are isomorphic.

Proof. Consider \mathbb{P}^m as the projective space $\mathbf{P}(U)$ of lines of a vector space U of dimension m+1 and $\Lambda = \mathbf{P}(U_0)$, where U_0 is a subspace of U of dimension p+1 (We will use " \mathbb{P} " for projective spaces of hyperplanes and " \mathbf{P} " for projective spaces of lines). Subspaces L, W of U of dimension 1 and p+2 determine points $[L] \in \mathbf{P}(U)$ and $[W/U_0] \in \mathbf{P}(U/U_0)$ respectively. The blow-up $\widetilde{\mathbb{P}}^m$ can be identified

with the closed subvariety of incidence $Q \subset \mathbf{P}(U) \times \mathbf{P}(U/U_0)$ defined by $Q := \{([L], [W/U_0]) : L \subset W\}$ and $\alpha : \widetilde{\mathbb{P}}^m \to \mathbb{P}^m$ is the projection on the first factor restricted to Q:

Note that $Q \to \mathbf{P}(U/U_0)$ is a projectivized vector bundles $\mathbf{P}(E) \to \mathbf{P}(U/U_0)$ where E is the vector bundle $E = (w, [W]) \in U \times \mathbf{P}(U/U_0)$. We want to show that $\mathbf{P}(E)$ and $\mathbb{P}(V)$ are isomorphic.

The exact sequence $0 \to U_0 \to W \to U/U_0 \to 0$ give rise to an exact sequence of vector bundles

(7)
$$0 \to U_0 \otimes \mathcal{O}_{\mathbb{P}^n} \to E \to \mathcal{O}_{\mathbb{P}^n}(-1) \to 0.$$

The above sequence splits $(\text{Ext}^1(\mathcal{O}_{\mathbb{P}^n}(-1), U_0 \otimes \mathcal{O}_{\mathbb{P}^n}) = 0)$ so

(8)
$$E = (U_0 \otimes \mathcal{O}_{\mathbb{P}^n}) \oplus \mathcal{O}_{\mathbb{P}^n}(-1)$$

Since $V = E^*(1)$, $\mathbb{P}(V) \simeq \mathbb{P}(E^*(1)) \simeq \mathbb{P}(E^*) \simeq \mathbf{P}(E)$.

Let h be the hyperplane bundle on \mathbb{P}^n and $\xi = \xi_V$ the tautological line bundle in $\mathbb{P}(V)$. We will make no distinction between h and their corresponding pull-back π^*h on $\mathbb{P}(V)$. Note that $\xi - h$ and h are nef on $\mathbb{P}(V)$. It is well known that the classical cohomology ring of $\mathbb{P}(V)$ is generated by h, ξ with the two relations

(9)
$$f_1(h,\xi) = h^{n+1},$$

(10)
$$f_2(h,\xi) = (\xi-h)^{r-1}(\xi-2h) = \sum_{k=0}^r (-1)^k c_k h^k \xi^{r-k}.$$

where $c_i = c_i(V)$ are the Chern classes of V.

The homology classes

(11)
$$A_1 := (h^n \xi^{r-2})_*, \qquad A_2 := (h^{n-1} \xi^{r-1} - rh^n \xi^{r-2})_*$$

form a system of generators for $H_2 \mathbb{P}(V)$ ((-)_{*} stays for the Poincare dual) and

(12)
$$(\xi - h)(A_1) = 1, \quad (\xi - h)(A_2) = 0, \quad h(A_1) = 0, \quad h(A_2) = 1$$

Proposition 2.2. $\mathbb{P}(V)$ is a Fano variety and the homology classes A_1 , A_2 are extremal rays for $NE(\widetilde{\mathbb{P}}^m)$. The canonical divisor is $-K = -K_{\mathbb{P}(V)} = r(\xi - h) + nh$.

Proof. In fact, $-K = r\xi - (n+1-c_1)h = r(\xi - h) + nh$. Let C be an effective curve and $[C] = a_1A_1 + a_2A_2$. From (12) we have $(\xi - h)[C] = a_1$ and $h[C] = a_2$. Since $\xi - h$ and h are nef, $a_1 \ge 0$ and $a_2 \ge 0$. Moreover, if C is non constant, $a_1 > 0$ or $a_2 > 0$ and $-K[C] = ra_1 + na_2 > 0$. Since h and $\xi - h$ are globally generated, $\mathbb{P}(V)$ is Fano.

Lemma 2.3. Let $\ell \subset \mathbb{P}^n$ be a line. To give a section $s : \ell \to \mathbb{P}(V|_\ell)$ of the projective vector bundle $\pi|_{\mathbb{P}(V|_\ell)} \to \ell$ is equivalent to give a surjective morphism of vector bundles $f : V|_\ell \to \mathcal{O}_\ell(t+1)$ and $t = c_1(s^*(\xi))$.

Proof. See [Ha] II Prop. 7.12

We will denote by $\mathbb{P}(f)$ the section s induced by f in the previous lemma.

Lemma 2.4. Let $\ell \subset \mathbb{P}^n$ be a line.

- 1. $[\mathbb{P}(f)(\ell)] = tA_1 + A_2$ for every surjective morphism $f: V|_{\ell} \to \mathcal{O}_{\ell}(t+1)$.
- 2. Vice versa, for any integer t, $tA_1 + A_2$ can be represented only by curves of the form $\mathbb{P}(f)(\ell)$ for some surjective morphism of vector bundles $f: V|_{\ell} \to \mathcal{O}_{\ell}(t+1)$ in particular $tA_1 + A_2$ is effective if and only if $t \ge 0$.

Proof. Let $A = tA_1 + A_2$.

- 1. Simply note that $h[\mathbb{P}(f)(\ell)] = 1 = h[tA_1 + A_2]$ and $\xi[\mathbb{P}(f)(\ell)] = t + 1 = \xi[tA_1 + A_2]$.
- 2. Let $\tilde{\ell}$ be a curve which represents $tA_1 + A_2$ and $\ell = \pi(\tilde{\ell})$. Since $h[\tilde{\ell}] = 1$, $\pi|_{\tilde{\ell}}\tilde{\ell} \to \ell$ is an isomorphism and ℓ is a line in \mathbb{P}^n . Then $\tilde{\ell}$ is a section of $\pi|_{\mathbb{P}(V|_{\ell})} : \mathbb{P}(V|_{\ell}) \to \ell$.

]

From (11) and the previous two lemmas we have:

Corollary 2.5. 1. A_1 , A_2 cannot be represented by reducible or non reduced curves.

- 2. A_1 is represented by a line in the fiber of π .
- 3. A_2 is represented by a rational smooth curve isomorphic to a line in \mathbb{P}^n via π .

Lemma 2.6. Under the isomorphism $\widetilde{\mathbb{P}}^m \simeq \mathbb{P}(V)$, the cohomology classes k, η correspond respectively to $\xi - h, \xi - 2h$. The classical cohomology ring of $\widetilde{\mathbb{P}}^m$ is given by

(13)
$$\mathrm{H}^{*}(\widetilde{\mathbb{P}}^{m},\mathbb{Z}) = \frac{\mathbb{Z}[k,\eta]}{(g_{1},g_{2})}$$

where $g_1(k,\eta) = (k-\eta)^{n+1} = (k-\eta)^{m-p}, \ g_2(k,\eta) = k^{r-1} = k^{p+1}\eta.$

Proof. Let $\tilde{\ell}_1 \subset \widetilde{\mathbb{P}}^m$ be the strict transform in the blow-up of a line $\ell_1 \subset \mathbb{P}^m$ not included in Λ . Note that the preimage $\alpha^{-1}(\Lambda)$ of Λ is isomorphic to $\Lambda \times \mathbb{P}^n \subset \widetilde{\mathbb{P}}^m$ and $\alpha|_{\Lambda \times \mathbb{P}^n} : \Lambda \times \mathbb{P}^n \to \Lambda$ is the canonical projection on the first factor. Let ℓ_2 be a line in the fiber of $\alpha|_{\Lambda \times \mathbb{P}^n}$ and $B_1 := [\tilde{\ell}_1]$, $B_2 := [\ell_2]$. After the characterization given in corollary 2.5, it easy to show that A_1, A_2 correspond respectively to B_1, B_2 under the isomorphism $\widetilde{\mathbb{P}}^m \simeq \mathbb{P}(V)$ described in theorem 2.1 and $k[B_1] = 1$, $k[B_2] = 0, \eta[B_1] = 1$ and $\eta[B_2] = -1$.

3. Study of $\mathfrak{M}(A, 0)$.

Let $\mathfrak{M}(A,0)$ be the moduli space of morphism $f: \mathbb{P}^1 \to \mathbb{P}(V)$ with $[\operatorname{Im} f] = A$ where A is a class in $\operatorname{H}_2 \mathbb{P}(V)$.

Recall that the virtual dimension of $\mathfrak{M}(A,0)$ is

(14)
$$\dim_{\operatorname{virt}}(\mathfrak{M}(A,0)) = -K[A] + n + r - 1.$$

Lemma 3.1. $\mathfrak{M}(A_1, 0)$ is smooth compact of expected dimension n + 2r - 1.

Proof. By lemma 2.5, A_1 is represented by a line $\ell \subset \mathbb{P}(V)$ in the fiber of $\pi : \mathbb{P}(V) \to \mathbb{P}^n$, then $\mathfrak{M}(A_1, 0)$ is compact of dimension $3 + \dim \mathbb{P}^n + \dim G(2, r) = n + 2r - 1$. It remains to show that $\mathfrak{M}(A_1, 0)$ is smooth.

From the inclusion $\ell \subset \pi^{-1}(p) \subset \mathbb{P}(V)$ we have the short exact sequence of normal bundles

(15)
$$0 \to N_{\ell|\pi^{-1}(p)} \to N_{\ell|\mathbb{P}(V)} \to N_{\pi^{-1}(p)|\mathbb{P}(V)} \to 0.$$

Since

(

16)
$$N_{\ell|\pi^{-1}(p)} = N_{\ell|\mathbb{P}^{r-1}} = \mathcal{O}_{\ell}(1)^{\oplus r-2},$$

(17)
$$N_{\pi^{-1}(p)|\mathbb{P}(V)} = N_{\pi^{-1}(p)|\mathbb{P}^n \times \mathbb{P}^{r-1}} = (\pi_{|\pi^{-1}(p)})^* (N_{\{p\}|\mathbb{P}^n}) \\ = (\pi_{|\pi^{-1}(p)})^* (T_p \mathbb{P}^n),$$

so $\mathrm{H}^1(N_{\ell|\mathbb{P}(V)}) = 0.$

Lemma 3.2. Let t be an integer and $A_{(t)} := tA_1 + A_2 \in H_2 \mathbb{P}(V)$. Then $\mathfrak{M}(A_{(t)}, 0)$ is empty for t < 0 and compact of expected dimension 2n + tr + r - 1 if $t \ge 0$.

Proof. By lemma 2.4, morphisms $f : \mathbb{P}^1 \to \mathbb{P}(V)$ with $[\operatorname{Im} f] = A_{(t)}$ exist only for $t \ge 0$ and in this case they are parametrized by an open subset of $G(2, n + 1) \times \mathbb{P}(\mathrm{Hom}(V|_{\ell}, \mathcal{O}_{\ell}(t + 1))))$ where G(2, n + 1) is the grassmannian of lines in \mathbb{P}^n . Then

(18)
$$\dim \mathfrak{M}(A_{(t)}, 0) = 3 + \dim G(2, n+1) + \dim \mathrm{H}^{0}(\mathrm{Hom}(V|_{\ell}, \mathcal{O}_{\ell}(t+1))) - 1.$$

Since $\operatorname{Hom}(V|_{\ell}, \mathcal{O}_{\ell}(t+1)) = \mathcal{O}_{\ell}(t)^{\oplus r-1} \oplus \mathcal{O}_{\ell}(t-1)$, so for any $t \ge 0$

(19)
$$\dim \mathrm{H}^{0}(\mathrm{Hom}(V|_{\ell}, \mathfrak{O}_{\ell}(t+1))) = (r-1)(t+1) + t$$
$$= tr + r - 1$$

and

(20)
$$\dim \mathfrak{M}(A_{(t)}, 0) = 3 + 2(n-1) + (tr + r - 2) \\ = 2n + tr + r - 1.$$

Proof. After lemma 3.2, we only need to prove smoothness. Let ℓ be a line in \mathbb{P}^n . The natural projection $V|_{\ell} = \mathcal{O}_{\ell}(1)^{\oplus (r-1)} \oplus \mathcal{O}_{\ell}(2) \to \mathcal{O}_{\ell}(1)^{\oplus (r-2)}$ induces an embedding $\ell \times \mathbb{P}^{r-2} = \mathbb{P}(\mathcal{O}_{\ell}(1)^{\oplus k}) \subset \mathbb{P}(V|_{\ell})$. If f is a morphism belonging to $\mathfrak{M}(A_2, 0)$, we know (lemma 2.4) that Im f is of the form $\ell \times \{p\}$ where p is a point in \mathbb{P}^{r-2} . The three inclusions

(21)
$$\ell \times \{p\} \subset \ell \times \mathbb{P}^{r-2} \subset \mathbb{P}(V|_{\ell}) \subset \mathbb{P}(V)$$

give rise to two exact sequences relating normal bundles:

(22)
$$0 \to N_{\ell \times \{p\} | \mathbb{P}(V|_{\ell})} \to N_{\ell \times \{p\} | \mathbb{P}(V)} \to N_{\mathbb{P}(V|_{\ell}) | \mathbb{P}(V)} \to 0,$$

(23)
$$0 \to N_{\ell \times \{p\} \mid \ell \times \mathbb{P}^{r-2}} \to N_{\ell \times \{p\} \mid \mathbb{P}(V \mid_{\ell})} \to N_{\ell \times \mathbb{P}^{r-2} \mid \mathbb{P}(V \mid_{\ell})} \to 0.$$

Notice that

(24)
$$N_{\mathbb{P}(V|_{\ell})|\mathbb{P}(V)} = (\pi|_{\mathbb{P}(V|_{\ell})})^* (N_{\ell}|_{\mathbb{P}^n}) = (\pi|_{\mathbb{P}(V|_{\ell})})^* (\mathcal{O}_{\ell}(1)^{\oplus (n-1)}),$$

(25)
$$N_{\ell \times \{p\} \mid \ell \times \mathbb{P}^{r-2}} = \mathcal{O}_{\ell} \otimes N_{\{p\} \mid \mathbb{P}^{r-2}} = \mathcal{O}_{\ell}^{\oplus (r-2)}.$$

Since $\xi|_{\ell \times \mathbb{P}^{r-2}} = \mathcal{O}_{\ell}(1) \otimes \mathcal{O}_{\mathbb{P}}^{r-2}(1),$

(26)
$$N_{\ell \times \mathbb{P}^{r-2}|\mathbb{P}(V|_{\ell})} = \mathcal{O}_{\ell}(-2) \otimes \xi|_{\ell \times \mathbb{P}^{r-2}} = \mathcal{O}_{\ell}(-1) \otimes \mathcal{O}_{\mathbb{P}^{r-2}}(1).$$

Then, from long cohomology sequences of (22) and (23) we get $\mathrm{H}^1(N_{\ell \times \{p\} \mid \mathbb{P}(V)}) = 0$.

4. Computation of Gromov-Witten invariants.

If A belongs to $H_2 \mathbb{P}(V)$ and $\alpha_1, \ldots, \alpha_s$ are classes in $H^*(\mathbb{P}(V))$, $I_A(\alpha_1, \alpha_2, \ldots, \alpha_s)$ denotes, as usual, the Gromov-Witten invariant.

Lemma 4.1. If $q_1 + q_2 < br$,

(27)
$$I_{bA_1}(h^{p_1}\xi^{p_1}, h^{p_2}\xi^{p_2}, \alpha) = 0.$$

Proof. We can choose α of the form $h^p h^q$ with $q \leq r-1$ and $p+p_1+p_2+q+q_1+q_2 = n+r-1+br$. If $I_{bA_1}(h^{p_1}\xi^{p_1}, h^{p_2}\xi^{p_2}, \alpha) \neq 0$, it exists an effective connected curve E of class A_1 meeting three varieties B, C, D dual to $h^{p_1}\xi^{q_1}, h^{p_2}\xi^{q_1}$, $h^p\xi^q$ respectively. Notice that $\pi(B), \pi(C), \pi(D)$ in \mathbb{P}^n are dual to h^{p_1}, h^{p_2}, h^p . Since A_1 is an extremal ray, every irreducible component of E is a multiple of A_1 , then contained in a fiber of π . Therefore the whole E is contained in a fiber of π as E is connected. We deduce that $\pi(B) \cap \pi(C) \cap \pi(D) \neq \emptyset$ thus $p_1 + p_2 + p \leq n$ and

(28)
$$q_1 + q_2 = n + r - 1 + br - p_1 - p_2 - p - q$$
$$\geq (n + r - 1) + br - (n + r - 1)$$
$$\geq br.$$

Lemma 4.2.

(29)
$$I_{A_1}(\xi,\xi^{r-1},h^n\xi^{r-1}) = 1.$$

Proof. By lemma 3.1, $\mathfrak{M}(A_1, 0)$ is smooth compact of expected dimension. The class $h^n \xi^{r-1}$ is the dual of a point q in $\mathbb{P}(V)$. Let $p := \pi(q)$. A parametrized curve in $\mathfrak{M}(A_1, 0)$ meeting q has support on a line $\ell \subset \pi^{-1}(q)$. Since $\xi|_{\pi^{-1}(q)}$ is the cohomology class of a hyperplane in $\pi^{-1}(q)$, then $I_{A_1}(\xi,\xi^{r-1},h^n\xi^{r-1}) = 1$.

Lemma 4.3. Let \tilde{n} be the largest integer less or equal to (n+1)/2. Then

(30)
$$I_{A_2}(h^{\tilde{n}}, h^{n+1-\tilde{n}}, h^n \xi^{r-2}) = 1,$$

(31)
$$I_{A_2}(h^{\tilde{n}}, h^{n+1-\tilde{n}}, h^{n-1}\xi^{r-1}) = r - 1.$$

Proof. Let S be the universal bundle, Q the quotient bundle on the grassmannian G(2, n + 1) and

(32)
$$0 \to S \to \left(\mathcal{O}_{G(2,n+1)}\right)^{\oplus (n+1)} \to Q \to 0$$

the tautological exact sequence. The bundles S and Q have ranks 2 and n-1 respectively. Let α , β be the virtual classes such that $\alpha + \beta = -c_1(S)$ and $\alpha\beta = c_2(S)$. Then

(33)
$$\operatorname{cl}(\{\ell \in G(2, n+1) | \ell \cap h^p \neq \emptyset\}) = \frac{\alpha^p - \beta^p}{\alpha - \beta},$$

where cl(-) denotes the fundamental class and h^p stands for a fixed linear subspace of \mathbb{P}^n of codimension p. As explained in [1], a polynomial of maximal degree in the Chern classes of the bundle S can be written as homogeneous polynomial $P(\alpha, \beta)$ of degree 2n - 2 and

(34)
$$\int_{G(2,n+1)} P(\alpha,\beta) = \text{``the coefficient of } \alpha^n \beta^n \text{ in } -\frac{1}{2}(\alpha-\beta)^2 P(\alpha,\beta).$$

If \tilde{h} is the hyperplane class on \mathbb{P}^{r-2} , the restriction of ξ to $\mathbb{P}^n \times \mathbb{P}^{r-2} = \mathbb{P}(\mathcal{O}_{\mathbb{P}^n}(1)^{r-1})$ is $(h + \tilde{h})$ and

(35)
$$(h^n \xi^{r-2})|_{\mathbb{P}^n \times \mathbb{P}^{r-2}} = h^n \tilde{h}^{r-2},$$

(36)
$$(h^{n-1}\xi^{r-1})|_{\mathbb{P}^n \times \mathbb{P}^{r-2}} = h^{n-1}\tilde{h}^{r-1} + (r-1)h^n\tilde{h}^{r-2}.$$

Thus

(37)
$$I_{A_2}(h^{\tilde{n}}, h^{n+1-\tilde{n}}, h^n \xi^{r-2}) = \int_{G(2,n+1) \times \mathbb{P}^{r-2}} P_1(\alpha, \beta),$$

(38)
$$I_{A_2}(h^{\tilde{n}}, h^{n+1-\tilde{n}}, h^{n-1}\xi^{r-1}) = \int_{G(2,n+1)\times\mathbb{P}^{r-2}} P_2(\alpha, \beta),$$

where

(39)
$$P_{1}(\alpha,\beta) = \frac{\alpha^{\tilde{n}} - \beta^{\tilde{n}}}{\alpha - \beta} \cdot \frac{\alpha^{n+1-\tilde{n}} - \beta^{n+1-\tilde{n}}}{\alpha - \beta} \cdot \frac{\alpha^{n} - \beta^{n}}{\alpha - \beta} \cdot \tilde{h}^{r-2}$$
$$= \frac{\alpha^{n+1} - \alpha^{n+1-\tilde{n}}\beta^{n} - \alpha^{n}\beta^{n+1-\tilde{n}} + \beta^{n+1}}{(\alpha - \beta)^{2}} \cdot \frac{\alpha^{n} - \beta^{n}}{\alpha - \beta} \cdot \tilde{h}^{r-2}$$

and, analogously,

(40)

$$P_{2}(\alpha,\beta) = \frac{\alpha^{n+1} - \alpha^{n+1-\tilde{n}}\beta^{n} - \alpha^{n}\beta^{n+1-\tilde{n}} + \beta^{n+1}}{(\alpha-\beta)^{2}} \times \left[\frac{\alpha^{n-1} - \beta^{n-1}}{\alpha-\beta} \cdot \tilde{h}^{r-1} + (r-1)\frac{\alpha^{n} - \beta^{n}}{\alpha-\beta} \cdot \tilde{h}^{r-2}\right]$$
Using (24) we obtain (20) and (21)

Using (34) we obtain (30) and (31).

5. The quantum cohomology ring of $\mathbb{P}(V)$: proof of the main theorem.

As we are going to work with two ring structure, classical and quantistic, on the cohomology group $\mathrm{H}^*(\mathbb{P}(V))$, we have to fix some notation. If α,β are cohomology classes in $\mathrm{H}^*\mathbb{P}(V)$ then $\alpha^{\bullet i} \bullet \beta^{\bullet j}$ is the quantum product of *i* copies of α and *j* copies of β . For the classical product we write $\alpha^i \beta^j$ as usual. For any homology class $A \in \mathrm{H}_2 \mathbb{P}(V)$ we denote by $(\alpha^{\bullet i} \bullet \beta^{\bullet j})_A$ the contribute of *A* to the product $\alpha^{\bullet i} \bullet \beta^{\bullet j}$ defined as follows. Let *d* be the integer $d := i \deg \alpha + j \deg \beta + K[A]$. Then $(\alpha^{\bullet i} \bullet \beta^{\bullet j})_A = 0$ for d < 0. For $d \ge 0$, $(\alpha^{\bullet i} \bullet \beta^{\bullet j})_A$ is the class of degree *d* in $\mathrm{H}^* \mathbb{P}(V)$ satisfying the property

(41)
$$(\alpha^{\bullet i} \bullet \beta^{\bullet j})_A[\gamma_*] = I_A(\alpha, \dots, \alpha, \beta, \dots, \beta, \gamma)$$

for every $\gamma \in H^*$ with deg $\gamma = \dim \mathbb{P}(V) - d$, being $I_A(\alpha, \ldots, \alpha, \beta, \ldots, \beta, \gamma)$ the Gromov-Witten invariant.

Then

(42)
$$\alpha^{\bullet i} \bullet \beta^{\bullet j} = \sum_{A \in \mathcal{H}_2 \mathbb{P}(V)} (\alpha^{\bullet i} \bullet \beta^{\bullet j})_A$$
$$= \alpha^i \beta^j + (\alpha^{\bullet i} \bullet \beta^{\bullet j})_{[E_1]} + (\alpha^{\bullet i} \bullet \beta^{\bullet j})_{[E_2]} + \dots$$

for some non constant effective curves $[E_1], [E_2], \ldots$

Theorem 5.1. When r < n the two relations

(43)
$$h^{\bullet(n+1)} = \xi - 2h$$
$$(\xi - h)^{\bullet(r-1)} \bullet (\xi - 2h) = 1$$

$$(\xi -$$

hold in $\operatorname{H}^*_O \mathbb{P}(V)$.

Proof. Since $r = \min(-K[A_1], -K[A_2])$, we have $h^{\bullet p}\xi^{\bullet q} = h^p\xi^q$ whenever p + q < r. Moreover for p + q < n, the contributions to the quantum product $h^{\bullet p}\xi^{\bullet q}$ can only arise from curves of type tA_1 . In particular, lemma 4.1 implies that

(44)
$$h^{\bullet p} \xi^{\bullet q} = h^p \xi^q \quad \text{for } q < r,$$

(45)
$$h^{\bullet p} = h^p \quad \text{for } p < n.$$

By lemma 4.2, $\xi^{\bullet r} = 1$, then

(46)
$$(\xi - h)^{\bullet (r-1)} \bullet (\xi - 2h) = (\xi - h)^{(r-1)} (\xi - 2h) + 1$$
$$= 1.$$

Analogously, by lemma 4.3, $(h^{\tilde{n}} \bullet h^{n+1-\tilde{n}})_{A_2} = \xi - 2h$, then

(47)
$$h^{n+1} = h^{n} \bullet h^{n+1-n} = h^n \bullet h^{n+1-1}$$
$$= h^{n+1} + (h^{\tilde{n}} \bullet h^{n+1-\tilde{n}})_{A_2}$$
$$= \xi - 2h.$$

We are finally ready to prove the main result of this work.

Proof of theorem 1.1. If 2p+3 < m then r < n and theorem 5.1 applies. Relations (43) can be easily translated to the ring $H_Q^* \widetilde{\mathbb{P}}^m$ with the aid of lemma 2.6 obtaining the two relations (4). Then \tilde{g}_1, \tilde{g}_2 are the classical relations g_1, g_2 in (2) evaluated in the quantum cohomology ring. By theorem 2.2 in [2], the quantum cohomology ring of $H_Q^* \widetilde{\mathbb{P}}^m$ is the ring generated by k, η with the two relations \tilde{g}_1, \tilde{g}_2 (compare also with proposition 10 of [4]).

References

- 1. A.Beauville, Quantum cohomology of complete intersection, alg-geom/9501008.
- 2. B.Siebert and G.Tian, On quantum cohomology rings of Fano manifolds and a formula of Vafa and Intriligator, alg-geom/9403010.
- C.Okonek, M.Schneider, and H.Spindler, Vector bundles on complex projective spaces, Progress in mathematics, Birkhüser, 1980.
- 4. W. Fulton and R. Pandharipande, Notes on stable maps and quantum cohomology, alg-geom/9608011.
- 5. Andreas Gathmann, Counting rational curves with multiple points and Gromov-Witten invariants of blow-ups, alggeom/9609010.
- 6. _____, Gromov-Witten invariants of blow-ups, math/9804043.
- 7. Lothar Göttsche and Rahul Pandharipande, The quantum cohomology of blow-ups of \mathbb{P}^2 and enumerative geometry, alg-geom/9611012.
- 8. Robin Hartsorn, Algebraic geometry, Graduate Text in Mathematics, vol. 52, Springer-Verlag, 1978.
- 9. Zhenbo Qin and Yongbin Ruan, Quantum cohomology of projectivized bundles over \mathbb{P}^n , alg-geom/9510010. E-mail address: maggesi@math.unifi.it

URL: http://www.math.unifi.it/~maggesi/