# THE OBSTACLE PROBLEM FOR FUNCTIONS OF LEAST GRADIENT

WILLIAM P. ZIEMER KEVIN ZUMBRUN Dedicated to Alois Kufner on the occasion of his sixty-fifth birthday, with the authors' best wishes.

ABSTRACT. For a given domain  $\Omega \subset \mathbb{R}^n$ , we consider the variational problem of minimizing the  $L^1$ -norm of the gradient on  $\Omega$  of a function u with prescribed continuous boundary values and satisfying a continuous lower obstacle condition  $u \geq \Psi$  inside  $\Omega$ . Under the assumption of strictly positive mean curvature of the boundary  $\partial\Omega$ , we show existence of a continuous solution, with Hölder exponent half of that of data and obstacle.

This generalizes previous results obtained for the unconstrained and doubleobstacle problems. The main new feature in the present analysis is the need to extend various maximum principles from the case of two area-minimizing sets to the case of one sub- and one superminimizing set. This we accomplish subject to a weak regularity assumption on one of the sets, sufficient to carry out the analysis. Interesting open questions include the uniqueness of solutions and a complete analysis of the regularity properties of area superminimizing sets. We provide some preliminary results in the latter direction, namely a new monotonicity principle for superminimizing sets, and the existence of "foamy" superminimizers in two dimensions.

### 1. INTRODUCTION

A rather complete and extensive literature is now in place concerning existence and regularity of solutions to a wide range of variational problems for which the following is prototypical:

(1.1) 
$$\inf\left\{\int_{\Omega} |\nabla u|^p : u \in W^{1,p}(\Omega), u - g \in W^{1,p}_0(\Omega)\right\}.$$

Here,  $\Omega \subset \mathbb{R}^n$  is a bounded, open set,  $1 and <math>g \in W^{1,p}(\Omega) \cap C^0(\overline{\Omega})$ . The Euler-Lagrange equation for (1.1) is the *p*-Laplacian div $(|\nabla u|^{p-2} \nabla u) = 0$ . The interested reader can consult recent books on this subject and the references therein, [AH], [HKM], and [MaZ]. The theory related to the case corresponding to p = 1 is far less complete. In spite of the fact that there is a vast literature relating to the least area functional,

$$\inf_{u}\left\{\int_{\Omega}\sqrt{1+\left|\nabla u\right|^{2}}\right\},$$

Typeset by  $\mathcal{AMS}$ -TEX

<sup>1991</sup> Mathematics Subject Classification. Primary 49Q05; Secondary 35J85.

Key words and phrases. least gradient, sets of finite perimeter, area-minimizing, obstacle.

Research of first author supported in part while visiting Charles University, June, 1997

Research of second author supported in part by a grant from the National Science Foundation

there are many open questions concerning other functionals with linear growth in  $|\nabla u|$ . Investigations concerning such questions were considered in [SWZ], [SZ], ???. In particular, the Dirichlet problem was investigated; that is, for a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^n$ , and for  $g : \partial\Omega \to \mathbb{R}^1$  continuous, the questions of existence and regularity of solutions to

(1.2) 
$$\inf \{ \|\nabla u\|(\Omega) : u \in BV(\Omega), \ u = g \text{ on } \partial\Omega \}$$

were examined. Here  $\|\nabla u\|(\Omega)$  denotes the total variation of the vector-valued measure  $\nabla u$  evaluated on  $\Omega$ . It was shown that a solution  $u \in BV(\Omega) \cap C^0(\Omega)$ exists provided that  $\partial\Omega$  satisfies two conditions, namely, that  $\partial\Omega$  has non-negative curvature (in a weak sense) and that  $\partial\Omega$  is not locally area-minimizing. See Section 2 below for notation and definitions.

In this paper we consider the obstacle problem

(1.3) 
$$\inf \left\{ \|\nabla u\| (\Omega) : u \in C^0(\overline{\Omega}), \ u \ge \psi \text{ on } \Omega, u = g \ge \psi \text{ on } \partial\Omega \right\}$$

where  $\Omega \subset \mathbb{R}^n$  is a bounded Lipschitz domian,  $q: \partial \Omega \to \mathbb{R}^1$  is continuous and  $\psi$  is a continuous function on  $\overline{\Omega}$ . The analogous obstacle problem for (1.1) was investigated by several authors and is now well understood, cf. [CL], [Li], [MiZ], [MuZ]. One of the difficulties encountered in the analysis of both (1.1) and (1.3) is the fact that the compactness in  $L^1(\Omega)$  of a sequence whose BV-norms are bounded does not ensure, a *priori*, continuity of the limiting function or that it will assume the boundary values q, thus making the question of existence problematic. In this paper as well as in [SWZ], we rely heavily on the discovery made in [BDG] that the superlevel sets of a function of least gradient are area-minimizing. This fact, along with the co-area formula (see (2.10) below), suggests that the existence of a function of least gradient subject to an obstacle constraint can be established by actually constructing each of its superlevel sets in such a way that it reflects both the appropriate boundary condition and the obstacle condition. The main thrust of this paper is to show that this is possible. Thus we show that there exists a continuous solution to (1.3) and we also show it inherits essentially the same regularity as the boundary data and obstacle.

As in [SWZ], both existence and regularity are developed by extensive use of BV theory and sets of finite perimeter as well as certain maximum principles. One of the main contributions of this paper is a new maximum principle that involves a super area-minimizing set and an area-minimizing set, Theorem 3.3. The similar result involving two area-minimizing sets, due independently to [Mo] and [S2], played a crucial role in [SWZ].

Our extended maximum principle requires a weak regularity property on one of the sets, that the set be contained in the (topological) closure of its interior. This is clearly satisfied in the contexts that we apply it, for which one of the sets is always area-minimizing. But, an interesting open question is whether or not this technical assumption can be dropped.

This issue leads us to consider a question of interest in its own right: "What is the regularity of a (sub)superminimizing set?" We conclude by presenting some separate, preliminary results on this subject, including a new monotonicity principle for (sub)superminimizing sets, and the existence of unusual, "foamy" (sub) superminimizers in two dimensions. It is our hope that these results will stimulate further investigation into the topic of regularity.

### 2. Preliminaries

The Lebesgue measure of a set  $E \subset \mathbb{R}^n$  will be denoted |E| and  $H^{\alpha}(E), \alpha > 0$ , will denote  $\alpha$ -dimensional Hausdorff measure of E. Throughout the paper, we almost exclusively employ  $H^{n-1}$ . The Euclidean distance between two points  $x, y \in \mathbb{R}^n$  will be denoted by |x - y|. The open ball of radius r centered at x is denoted by B(x, r) and  $\overline{B}(x, r)$  denotes its closure.

If  $\Omega \subset \mathbb{R}^n$  is an open set, the class of function  $u \in L^1(\Omega)$  whose partial derivatives in the sense of distribution are measures with finite total variation in  $\Omega$  is denoted by  $BV(\Omega)$  and is called the space of functions of bounded variation on  $\Omega$ . The space  $BV(\Omega)$  is endowed with the norm

(2.1) 
$$\|u\|_{BV(\Omega)} = \|u\|_{1;\Omega} + \|\nabla u\|(\Omega)$$

where  $||u||_{1;\Omega}$  denotes the  $L^1$ -norm of u on  $\Omega$  and where  $||\nabla u||$  is the total variation of the vector-valued measure  $\nabla u$ .

The following compactness result for  $BV(\Omega)$  will be needed later, cf. [Gi] or [Z].

**Theorem 2.1.** If  $\Omega \subset \mathbb{R}^n$  is a bounded Lipschitz domain, then

$$BF(\Omega) \cap \left\{ u : \|u\|_{BV(\Omega)} \le 1 \right\}$$

is compact in  $L^1(\Omega)$ . Moreover, if  $u_i \to u$  in  $L^1(\Omega)$  and  $U \subset \Omega$  is open, then

$$\liminf_{x \to \infty} \|\nabla u_i\|(U) \ge \|\nabla u\|(U)$$

A Borel set  $E \subset \mathbb{R}^n$  is said to have *finite perimeter in*  $\Omega$  provided the characteristic function of  $E, \chi_E$ , is a function of bounded variation in  $\Omega$ . Thus, the partial derivatives of  $\chi_E$  are Radon measures on  $\Omega$  and the perimeter of E in  $\Omega$  is defined as

(2.2) 
$$P(E,\Omega) = \|\nabla \chi_E\|(\Omega).$$

A set E is said to be of *locally finite perimeter* if  $P(E, \Omega) < \infty$  for every bounded open set  $\Omega \subset \mathbb{R}^n$ .

One of the fundamental results in the theory of sets of finite perimeter is that they possess a measure-theoretic exterior normal which is suitably general to ensure the validity of the Gauss-Green theorem. A unit vector  $\nu$  is defined as the measuretheoretic exterior normal to E at x provided

$$\lim_{r \to 0} r^{-n} |B(x,r) \cap \{y : (y-x) \cdot \nu < 0, \ y \notin E\}| = 0$$

and

(2.3) 
$$\lim_{r \to 0} r^{-n} |B(x,r) \cap \{y : (y-x) \cdot \nu > 0, \ y \notin E\}| = 0.$$

The measure-theoretic normal of E at x will be denoted by  $\nu(x, E)$  and we define

(2.4) 
$$\partial_* E = \{x : \nu(x, E) \ exists\}.$$

The Gauss-Green theorem in this context states that if E is a set of locally finite perimeter and  $V: \mathbb{R}^n \to \mathbb{R}^n$  is a Lipschitz vector field, then

(2.5) 
$$\int_{E} \operatorname{div} V(x) \, dx = \int_{\partial_{*}E} V(x) \cdot \nu(x, E) \, dH^{n-1}(x),$$

cf. [Fe2, §4.5.6]. Clearly,  $\partial_* E \subset \partial E$ , where  $\partial E$  denotes the topological boundary of E. Also, the topological interior of E is denoted by  $E^i = (\mathbb{R}^n \setminus \partial E) \cap E$ , the topological exterior by  $E^e = (\mathbb{R}^n \setminus \partial E) \cap (\mathbb{R}^n \setminus E)$  and  $E^c$  to denote the complement  $\mathbb{R}^n \setminus E$ . The notation  $E \subset \subset F$  means that the closure of E,  $\overline{E}$ , is a compact subset of  $F^i$ .

For measurable sets E, the measure-theoretic interior,  $E_m^i$ , is the set of all points at which the metric density of E is 1 and the measure-theoretic exterior,  $E_m^e$ , is all points at which the metric density is 0. The measure theoretic-boundary,  $\partial_m E := \mathbb{R}^n \setminus (E_m^i \cup E_m^e)$ . Clearly,  $\partial_* E \subset \partial_m E \subset \partial E$ . Moreover, it is well known that

(2.6) E is of finite perimeter if and only if 
$$H^{n-1}(\partial_m E) < \infty$$

and that

(2.7) 
$$P(E,\Omega) = H^{n-1}(\Omega \cap \partial_m E) = H^{n-1}(\Omega \cap \partial_* E) \text{ whenever } P(E,\Omega) < \infty$$

cf. [Fe2 §4.5]. From this it easily follows that

(2.8) 
$$P(E \cup F, \Omega) + P(E \cap F, \Omega) \le P(E, \Omega) + P(F, \Omega),$$

thus implying that sets of finite perimeter are closed under finite unions and intersections.

The definition implies that sets of finite perimeter are defined only up to sets of measure 0. In other words, each set determines an equivalence class of sets of finite perimeter. In order to avoid this ambiguity, we will employ

(2.9) 
$$\tilde{E} := E \cup E_1 \setminus E_0$$

as the distinguished representative for E. Here,

$$E_i := \{x : |E \cap B(x, r)| / |B(x, r)| = i \text{ for all small } r > 0\}, \ i = 0, 1$$

Thus, with this convention, it easy to see that

(2.10) 
$$\overline{\partial_* E} = \partial E.$$

This convention will apply, in particular, to all competitors of the variational problems (4.20) and (2.22) below as well as to the sets defined by (2.17).

Of particular importance to us are sets of finite perimeter whose boundaries are area-minimizing. If E is a set of locally finite perimeter and U a bounded, open set, then E is said to be *area-minimizing in* U if  $P(E,U) \leq P(F,U)$  whenever  $E\Delta F \subset C U$ . Also, E is said to be *super area-miniming in* U (sub area-minimizing in U) if  $P(E,U) \leq P(E \cup F,U)$  ( $P(E,U) \leq P(E \cap F,U)$ ) whenever  $E\Delta F \subset C U$ .

A tool that will play a significant role in this paper is the co-area formula. It states that if  $u \in BV(\Omega)$ , then

(2.11) 
$$\|\nabla u\|(\Omega) = \int_{-\infty}^{\infty} P(E_t, \Omega) dt$$

where  $E_t = \{u \ge t\}$ . In case u is Lipschitz, we have

$$\int_{\Omega} |\nabla u| dx = \int_{-\infty}^{\infty} H^{n-1} \left( u^{-1}(t) \cap \Omega \right) dt$$

Conversely, if u is integrable on  $\Omega$  then

(2.12) 
$$\int_{-\infty}^{\infty} P(E_t, \Omega) dt < \infty \text{ implies } u \in BV(\Omega),$$

cf. [Fe1], [FR].

Another fundamental result is the isoperimetric inequality for sets of finite perimeter. It states that there is a constant C = C(n) such that

(2.13) 
$$P(E)^{n/(n-1)} \le C|E|$$

whenever  $E \subset \mathbb{R}^n$  is a set of finite perimeter. Furthermore, equality holds if and only if E is a ball.

The regularity of  $\partial E$  plays a crucial role in our development. In particular, we will employ the notion of tangent cone. Suppose E is area-minimizing in U and for convenience of notation, suppose  $0 \in U \cap \partial E$ . For each r > 0, let,  $E_r = \mathbb{R}^n \cap \{x : rx \in E\}$ . It is known (cf. [S1, §35]) that for each sequence  $\{r_i\} \to 0$ , there exists a subsequence (denoted by the full sequence) such that  $\chi_{E_i}$  converges in  $L^1_{loc}(\mathbb{R}^n)$  to  $\chi_C$ , where C is a set of locally finite perimeter. In fact, C is area-minimizing and is called the tangent cone to E at 0. Although it is not immediate, C is a cone and therefore the union of half-lines issuing from 0. It follows from [S1, §37.6] that if  $\overline{C}$  is contained in  $\overline{H}$  where H is any half-space in  $\mathbb{R}^n$  with  $0 \in \partial H$ , then  $\partial H$  is regular at). That is there exists r > 0 such that

(2.14)  $B(0,r) \cap \partial E$  is a real analytic hypersurface.

Furthermore,  $\partial E$  is regular at all points of  $\partial_* E$  and

(2.15) 
$$H^{\alpha}\left(\left(\partial E \setminus \partial_* E\right) \cap U\right) = 0 \text{ for all } \alpha > n-8,$$

cf. [Gi, Theorem 11.8].

The boundary data g admits a continuous extension  $G \in BV(\mathbb{R}^n \setminus \overline{\Omega}) \cap C^0(\mathbb{R}^n \setminus \Omega)$ , [Gi, Theorem 2.16]. In fact,  $G \in C^{\infty}(\mathbb{R}^n \setminus \overline{\Omega})$ , but we only need that G is continuous on the complement of  $\Omega$ . Clearly, we can require that the support of G is contained in B(0, R) where R is chosen so that  $\Omega \subset C B(0, R)$ . We have

(2.16) 
$$G \in BV(\mathbb{R}^n \setminus \overline{\Omega}) \cap C^0(\mathbb{R}^n \setminus \Omega) \text{ with } G = g \text{ on } \partial\Omega.$$

We now introduce sets that will ensure that our constructed solution satisfies the required Dirichlet condition u = g on  $\partial \Omega$  and the obstacle condition  $u \ge \psi$  in  $\Omega$ . Thus, for each  $t \in [a, b]$ , let

(2.17) 
$$\mathcal{L}_t = (\mathbb{R}^n \setminus \Omega) \cap \{x : G(x) \ge t\},\$$

(2.18) 
$$L_t = \text{closure}(\{x : x \in \Omega, \psi(x) > t\})$$

Note that the co-area formula (2.11) and the fact that  $G \in BV(\mathbb{R}^n \setminus \overline{\Omega})$  imply that  $P(\mathcal{L}_t, \mathbb{R}^n \setminus \overline{\Omega}) < \infty$  for almost all t. For all such t, we remind the reader that we employ our convention (2.9) in defining  $\mathcal{L}_t$ .

We let [a, b] denote the smallest interval containing  $g(\partial \Omega) \cup \psi(\overline{\Omega})$  and define

(2.19) 
$$T := [a, b] \cap \left\{ t : P(\mathcal{L}_t, \mathbb{R}^n \setminus \overline{\Omega}) < \infty \right\}$$

Thus, by (2.7) and the fact that  $H^{n-1}(\partial\Omega) < \infty$ , we obtain

(2.20) 
$$H^{n-1}(\partial_m \mathcal{L}_t) = P(\mathcal{L}_t, \mathbb{R}^n \setminus \overline{\Omega}) + H^{n-1}\left[(\partial_m \mathcal{L}_t) \cap (\partial \Omega)\right] < \infty.$$

For each  $t \in T$ , the variational problems

(2.21) 
$$\min\left\{P(E,\mathbb{R}^n): E \setminus \overline{\Omega} = \mathcal{L}_t \setminus \overline{\Omega}, \ E \supset L_t\right\}$$

$$(2.22) \qquad \max\{|E|: E \text{ is a solution of } (2.21)\}$$

will play a central role in our development. In light of Theorem 2.1, a solution to both problems can be obtained from the direct method. (2.20) is also used to obtain existence for (2.21). We will denote by  $E_t$  the solution to (2.22). In this regard, note that our convention (2.9) ensures that  $E_t \setminus \overline{\Omega} = \mathcal{L}_t \setminus \overline{\Omega}$ ; furthermore, because of our convention,  $\mathcal{L}_t$  need not be a closed set. Also, observe that that  $E_t$ is super area-minimizing in  $\Omega$ .

# 3. A MAXIMUM PRINCIPLE

First, we begin with a result which is a direct consequence of a maximum principle for area-minimizing hypersurfaces established independently in [Mo] and [S2].

**3.1 Theorem.** Let  $E_1 \subset E_2$  and suppose both  $E_1$  and  $E_2$  are area-minimizing in an open set  $U \subset \mathbb{R}^n$ . Further, suppose  $x \in (\partial E_1) \cap (\partial E_2) \cap U$ . Then  $\partial E_1$  and  $\partial E_2$  agree in some neighborhood of x.

**3.2 Lemma.** For arbitrary measurable sets  $A, B \subset \mathbb{R}^n$ , it holds that

$$H^{n-1}(\partial_m(A \cup B)) \le H^{n-1}(\partial_m A \cap (B^i_m)^c) + H^{n-1}(\partial_m B \cap (\overline{A}_m)^c)$$
$$H^{n-1}(\partial_m (A \cap B)) \le H^{n-1}(\partial_m A \cap B^i_m) + H^{n-1}(\partial_m B \cap \overline{A}_m).$$

*Proof.* It follows immediately from definitions that

 $(\partial_m A \cap (B^i_m)^c) \cup (\partial_m B \cap (A^i_m)^c) = (\partial_m A \cap (B^i_m)^c) \cup (\partial_m B \cap (\overline{A}_m)^c),$ 

which yields the first inequality. The result for intersections then follows from  $\partial_m A = \partial_m A'$  and  $A \cap B = (A^c \cup B^c)^c$ .

**3.3 Theorem.** Let E be sub area-minimizing and F super area-minimizing relative to an open set U, with  $E \subset F$  and  $\partial E \cap \partial F \subset U$ . Further, suppose that  $\overline{E} \cap U = \overline{E^i} \cap U$ . Then, relative to U, either  $\partial E \cap \partial F = \emptyset$  or else  $\partial E = \partial F$  in a neighborhood of  $\partial E \cap \partial F$ .

*Proof.* Suppose  $\partial E \cap \partial F \neq \emptyset$ . The set  $\partial E \cap \partial F$  is contained in open neighborhood  $V \subset \subset U$  and thus, for sufficiently small  $|w|, w \in \mathbb{R}^n$ , we have

$$(3.1) (E+w) \setminus F \subset V + w \subset U.$$

Choose  $x_0 \in \partial E \cap \partial F$ . Since  $E = \overline{E^i}$ , there exists  $w \in \mathbb{R}^n$  with |w| arbitrarily small such that  $x_0 - w \in E^i$ , or equivalently

$$(3.2) x_0 \in (E+w)^i.$$

Denote the translated set E + w by  $E_w$ . By shrinking U if necessary, we can arrange that  $E_w$  is sub area-minimizing in U.

Now we will show that F is a rea-minimizing in the open set  $U \cap E_w^i$ . For, suppose to the contrary that there were a set G with

$$(3.3) G\Delta F \subset \subset U \cap E_w^i$$

and

$$(3.4) P(G, U \cap E_w^i) < P(F, U \cap E_w^i).$$

By (3.3),  $G \cap U = F \cap U$  near  $\partial E_w$ , while, by (3.4),

(3.5) 
$$H^{n-1}(\partial_m G \cap E^i_w \cap U) < H^{n-1}(\partial_m F \cap E^i_w \cap U)).$$

Since F and G agree on  $(E_w)^i_m \setminus E^i_w \subset \partial_m E_w$ , it follows that

(3.6) 
$$H^{n-1}(\partial_m G \cap (E_w)^i_m \cap U) < H^{n-1}(\partial_m F \cap (E_w)^i_m \cap U)).$$

On the other hand, super a rea-minimality of F in U implies that  $P(F \cup E_w, U) \ge P(F, U)$ . With Lemma 3.2, this gives

$$\begin{aligned} H^{n-1}(\partial_m F \cap ((E_w)^i_m)^c \cap U) + H^{n-1}(\partial_m E_w \cap (\overline{F}_m)^c \cap U) \\ &\geq H^{n-1}(\partial_m (F \cup E_w) \cap U) \\ &\geq H^{n-1}(\partial_m F \cap U) \\ &= H^{n-1}(\partial F \cap ((E_w)^i_m)^c \cap U) \\ &+ H^{n-1}(\partial F \cap (E_w)^i_m \cap U), \end{aligned}$$

and thus

(3.7) 
$$H^{n-1}(\partial E_w \cap (\overline{F}_m)^c \cap U) \ge H^{n-1}(\partial F \cap (E_w)^i_m \cap U).$$

Therefore,

$$(3.8) H^{n-1}(\partial (G \cap E_w) \cap U) \leq H^{n-1}(\partial E_w \cap \overline{G}_m \cap U) + H^{n-1}(\partial G \cap (E_w)^i_m \cap U) = H^{n-1}(\partial E_w \cap \overline{F}_m \cap U) + H^{n-1}(\partial G \cap (E_w)^i_m \cap U) < H^{n-1}(\partial E_w \cap \overline{F}_m \cap U) + H^{n-1}(\partial F \cap (E_w)^i_m \cap U) \leq H^{n-1}(\partial E_w \cap \overline{F}_m \cap U) + H^{n-1}(\partial E_w \cap (\overline{F}_m)^c \cap U) = H^{n-1}(\partial E_w \cap U),$$

where the first inequality follows by Lemma 3.2, the second by substituting F for G in the vicinity of  $\partial E_w$ , the third by (3.5), the fourth by (3.7), and the last by set decomposition. In other words,  $P(G \cap E_w, U) < P(E_w, U)$ . But, at the same time,

$$(G \cap E_w)\Delta E_w = E_w \setminus G \subset (E_w \setminus F) \cup (G\Delta F)$$

is compactly supported in U, by (3.1) and (3.3). contradicting the sub areaminimality of  $E_w$  in U. By contradiction, we have that F is area-minimizing in  $E_w^i \cap U$ , as claimed.

By basic regularity results, we thus have also that  $\overline{(F)} = \overline{F^i}$  in a neighborhood of  $x_0$ . By a symmetric argument, it follows that E is area-minimizing near  $x_0$  as well, and therefore we can appeal to Theorem 3.1 to obtain our conclusion.

We do not know whether the hypothesis  $E \cap U = \overline{E^i} \cap U$  in the previous result is necessary. However, in the case where E is area-minimizing in U, the regularity results (2.15) show that the hypothesis is satisfied and this is sufficient for the purposes of this paper. The following result is what we need and it now follows immediately from Theorem 3.3.

**3.4 Corollary.** Let E be area-minimizing and F super area-minimizing relative to an open set U, with  $E \subset F$  and  $\partial E \cap \partial F \subset U$ . Then, relative to U, either  $\partial E \cap \partial F = \emptyset$  or else  $\partial E = \partial F$  in a neighborhood of  $\partial E \cap \partial F$ .

# 4. Construction of the solution

In this section we will construct a solution u of (1.3) by using  $E_t \cap \overline{\Omega}$  to define the set  $\{u \ge t\}$  up to a set of measure zero for almost all t. This construction will be possible for bounded Lipschitz domains  $\Omega$  whose boundaries satisfy the following two conditions.

(i) For every  $x \in \partial \Omega$  there exists  $\varepsilon_0 > 0$  such that for every set of finite perimeter  $A \subset B(x, \varepsilon_0)$ 

$$(4.1) P(\Omega, \mathbb{R}^n) \le P(\Omega \cup A, \mathbb{R}^n)$$

(ii) For every  $x \in \partial \Omega$ , and every  $\varepsilon \geq 0$  there exists a set of finite perimeter  $A \subset \subset B(x, \varepsilon)$  such that

(4.2) 
$$P\left(\Omega, B(x,\varepsilon)\right) > P\left(\Omega \setminus A, B(x,\varepsilon)\right).$$

Clearly, we may assume that  $x \in A$ .

The first condition states that  $\partial\Omega$  has non-negative mean curvature (in the weak sense) while the second states that  $\Omega$  is not locally area-minimizing with respect to interior variations. Also, it can be easily verified that if  $\partial\Omega$  is smooth, then both conditions together are equivalent to the condition that the mean curvature of  $\partial\Omega$  is positive on a dense set of  $\partial\Omega$ .

Since  $\Omega$  is a Lipschitz domain, for each  $x_0 \in \partial\Omega$ ,  $\partial\Omega$  can be represented as the graph of a nonnegative Lipschitz function h defined on some ball  $B(x'_0, r) \subset \mathbb{R}^{n-1}$  where  $x'_0 \in \mathbb{R}^{n-1}$ . That is,  $\{(x', h(x')) : x' \in B'(x'_0, r)\} \subset \partial\Omega$ . Throughout we will use the notation  $B'(x'_0, r)$  and X' to denote elements in  $\mathbb{R}^{n-1}$  and thus they will be distinguished them from their n-dimensional counterparts  $B(x_0, r)$  and x.

We assume our configuration is oriented in such a way that

$$\{(x', x'') : 0 < x'' < h(x')\} \subset \Omega.$$

With  $S = \{(x', h(x')) : x' \in B'(x'_0, r)\}$  we have that

$$H^{n-1}(S) = \int_{B'(x'_0,r)} \sqrt{1 + |\nabla h|^2} \, dH^{n-1}(x')$$

These facts lead immediately to the following result.

**4.1 Lemma.** If  $\Omega$  is a Lipschitz domain with non-negative mean curvature in the sense of (4.1), then the function h, whose graph represents  $\partial\Omega$  locally, is a weak supersolution of the minimizing surface equation. That is, for r sufficiently small,

$$\int_{B'(x'_0,r)} \frac{\nabla h \cdot \nabla \phi}{\sqrt{1+|\nabla h|^2}} \, dx' \ge 0$$

whenever  $\phi \in C_0^{1,1}\left(B'(x'_0,r)\right), \phi > 0.$ 

We will also need the following result from [SWZ, Lemma 4.2] whose proof is an easy consequence of the weak Harnack inequality.

**4.2 Lemma.** Suppose W is an open subset of  $\mathbb{R}^{n-1}$ . If  $v_1, v_2 \in C^{0,1}(W)$  are respectively weak super and subsolutions of the minimal surface equation in W and if  $v_1(x'_0) = v_2(x'_0)$  for some  $x'_0 \in W$  while  $v_1(x') \ge v_2(x')$  for all  $x' \in W$ , then

$$v_1(x') = v_2(x')$$

for all x' in some closed ball contained in W centered at x'.

An important step in our development is the following lemma.

**4.3 Lemma.** For almost all  $t \in [a, b]$ ,  $\partial E_t \cap \partial \Omega \subset g^{-1}(t)$ .

*Proof.* First note that if  $t > \max_{x \in \partial \Omega} g(x)$ , then  $\partial E_t \cap \partial \Omega = \emptyset$ . So we may assume that  $t \in T$  and  $t \le \max_{x \in \partial \Omega} g(x)$ . The proof will proceed by contradiction and we first show that  $\partial E_t$  is locally area minimizing in a neighborhood of each point  $x_0 \in (\partial E_t \cap \partial \Omega \setminus g^{-1}(t))$ , i.e., we claim that there exists  $\varepsilon > 0$ , such that for every set F with the property that  $F\Delta E_t \subset C B(x_0, \varepsilon)$ , we have

(4.3) 
$$P(E_t, B(x_0, \varepsilon)) \le P(F, B(x_0, \varepsilon))$$

or equivalently,  $P(E_t, \mathbb{R}^n) \leq P(F, \mathbb{R}^n)$ .

By our assumption, either  $g(x_0) < t$  or  $g(x_0) > t$ . First consider the case  $g(x_0) < t$ . Since  $G(x_0) = g(x_0) < t$  and G is continuous on  $\mathbb{R}^n \setminus \Omega$ , there exists  $\varepsilon > 0$ , such that  $B(x_0, \varepsilon) \cap \mathcal{L}_t = \emptyset$ . Also,  $\psi$  is continuous on  $\overline{\Omega}$  and  $\psi(x_0) \leq g(x_0) < t$ , so we may take  $\varepsilon$  small enough such that  $L_t \cap B(x_0, \varepsilon) = \emptyset$ . We will assume that  $\varepsilon < \varepsilon_0$ , where  $\varepsilon_0$  appears in condition (4.1). We proceed by taking a variation F that satisfies  $F \Delta E_t \subset \subset B(x_0, \varepsilon)$ . Because of (4.1) and

(4.4, note that) 
$$P(E \cup F, \Omega) + P(E \cap F, \Omega) \le P(E, \Omega) + P(F, \Omega)$$

for every  $A \subset \subset B(x_0, \varepsilon)$ ,

$$P(A \cup \Omega, \mathbb{R}^n) + P(A \cap \Omega, \mathbb{R}^n) \le P(A, \mathbb{R}^n) + P(\Omega, \mathbb{R}^n)$$
$$\le P(A, \mathbb{R}^n) + P(A \cup \Omega, \mathbb{R}^n)$$

Hence

(4.5) 
$$P(A \cap \Omega, \mathbb{R}^n) \le P(A, \mathbb{R}^n).$$

Define  $F' = (F \setminus B(x_0, \varepsilon)) \cup (F \cap \overline{\Omega})$ , clearly

$$F' \setminus \overline{\Omega} = (F \setminus B(x_0, \varepsilon)) \setminus \overline{\Omega}$$
$$= (F \setminus \overline{\Omega}) \setminus B(x_0, \varepsilon)$$
$$= E_t \setminus \overline{\Omega} \setminus B(x_0, \varepsilon)$$
$$= \mathcal{L}_t \setminus \overline{\Omega} \setminus B(x_0, \varepsilon)$$
$$= \mathcal{L}_t \setminus \overline{\Omega}$$

and  $F' \supset L_t$ . Thus F' is admissible in (2.20) and therefore

$$P(E_t, \mathbb{R}^n) \le P(F', \mathbb{R}^n)$$

Now we will show that  $P(F', \mathbb{R}^n) \leq P(F, \mathbb{R}^n)$  which, with the previous inequality, will imply (4.3). First observe from  $E_t \Delta F \subset \subset B(x_0, \varepsilon)$  and  $(E_t \setminus \overline{\Omega}) \cap B(x_0, \varepsilon) = (\mathcal{L}_t \setminus \overline{\Omega}) \cap B(x_0, \varepsilon) = \emptyset$  that  $F' \cap B(x_0, \varepsilon) = F \cap B(x, \varepsilon) \cap \overline{\Omega}$  and  $F' \Delta F \subset \subset B(x_0, \varepsilon)$ . Hence we obtain by (4.5) (4.6)  $P(F, \mathbb{R}^n) - P(F', \mathbb{R}^n) = P(F, B(x_0, \varepsilon)) - P(F', B(x_0, \varepsilon))$ 

$$= P(F \cap B(x_0,\varepsilon), \mathbb{R}^n) - P(F \cap B(x_0,\varepsilon) \cap \Omega, B(x_0,\varepsilon))$$
$$= P(F \cap B(x_0,\varepsilon), \mathbb{R}^n) - P(F \cap B(x_0,\varepsilon) \cap \Omega, \mathbb{R}^n)$$
$$\ge 0$$

This establishes (4.3) when  $g(x_0) < t$ .

The argument to establish (4.3) when  $g(x_0) > t$  requires a slightly different treatment from the previous case. Since  $G(x_0) = g(x_0) > t$ , the continuity of Gin  $\Omega^c$  implies that  $\overline{B(x_0,\varepsilon)} \setminus \Omega \subset \mathcal{L}_t$ , provided  $\varepsilon$  is sufficiently small. Thus, we have  $\overline{B(x_0,\varepsilon)} \setminus \Omega \subset E_t$ . Clearly, we may assume  $\varepsilon$  chosen to be smaller than  $\varepsilon_0$  of (4.1). Observe that the assumption that  $\partial\Omega$  is locally Lipschitz implies that  $P(\Omega, B(x_0, \varepsilon)) = P(\mathbb{R}^n \setminus \Omega, B(x_0, \varepsilon))$ . Consequently, we can appeal to (4.1) to conclude that  $\mathbb{R}^n \setminus \Omega$  is sub area-minimizing in  $B(x_0, \varepsilon)$ . On the other hand,  $E_t$  is super area-minimizing. Since  $E_t \cap B(x_0, \varepsilon) \setminus \Omega \supset B(x_0, \varepsilon) \setminus \Omega$ , we may apply Theorem 3.3 to find that  $\partial E_t = \partial(\mathbb{R}^n \setminus \Omega)$  in some open neighborhood U of  $x_0$ . This implies that  $L_t \cap U = \emptyset$  since  $L_t \subset E_t$  and  $\partial E_t \cap U \cap \Omega = \emptyset$ . Consequently,  $E_t$  must be area-minimizing in U.

Thus far, we have shown that if either  $g(x_0) > t$  or  $g(x_0) < t$ , then  $\partial E_t$  is area minimizing in a neighborhood of  $x_0$ , say  $B(x_0, \varepsilon)$ . We will show this leads to a contradiction. Assume first that  $g(x_0) < t$  so that G < t on  $(\mathbb{R}^n \setminus \Omega) \cap B(x_0, \varepsilon)$ provided that  $\varepsilon$  has been chosen sufficiently small. Consequently

(4.7) 
$$E_t \cap B(x_0,\varepsilon) \subset \overline{\Omega} \cap B(x_0,\varepsilon).$$

We recall the notation concerning the representation of  $\partial\Omega$  as the graph of a Lipschitz function that preceded Lemma 3.1. Thus with  $x_0 \in \partial E_t \cap \partial\Omega \setminus g^{-1}(t)$ , we express  $\partial\Omega$  locally around  $x_0$  as  $\{(x'h(x')) : x' \in B(x'_0, \varepsilon)\}$ , where  $x_0 = (x'_0, x''_0)$  and  $x''_0 = h(x''_0) > 0$ . For simplicity of notation, we take  $x'_0 = 0$ . The number  $\varepsilon'$  is chosen so that  $\varepsilon' < \varepsilon$  and

(4.8) 
$$\{(x', h(x')) : |x'| < \varepsilon'\} \subset B(x_0, \varepsilon).$$

we define the half-infinite cylinder above  $B'(0,\varepsilon')$  as  $C = B'(0,\varepsilon') \times [0,\infty]$ . Because of the local nature of the argument we may assume that

$$\Omega \cap C = \{ (x', x'') : |x'|, 0 \le x'' < h(x') \}.$$

Now consider the solution to the minimal surface equation on  $B'(0, \varepsilon')$  relative to the boundary data  $f = h|_{\partial B'(0,\varepsilon')}$  [M.M. Chapter. 3]. Thus we let v be the unique solution of

$$div\left(\frac{\nabla v}{\sqrt{1+|\nabla v|^2}}\right) = 0 \quad \text{on } B'(0,\varepsilon').$$
$$v = f \quad \text{on } \partial B'(0,\varepsilon')$$

Since h is a weak supersolution of the minimal surface equation, by Lemma 3.1, we have that  $h \ge v$  on  $\overline{B'(0,\varepsilon)}$ , cf [GT, Theorem 10.7].; In fact, h > v on  $B'(0,\varepsilon')$  because the set  $\{h = v\}$  is obviously closed in  $B'(0,\varepsilon')$  and it is also open in  $B'(0,\varepsilon')$  because of Lemma 3.2. Hence, if this set is non-empty, h = v in  $B'(0,\varepsilon')$  which would contradict (4.2). Consequently, with  $\delta = h(0) - v(0)$ , we have  $\delta > 0$ . Now consider a 1-parameter family of graphs  $v_{\tau}(x') = v(x') + \tau$  and let

$$\tau_m = \max\left\{\tau : \text{there exists } x'' \in \overline{B'(0,\varepsilon')} \text{ such that } : (x',v_\tau(x')) \in \partial E_t \cap \overline{\Omega}\right\}$$

Note that  $\tau_m \geq \delta$  since  $v(0) + \delta = h(0)$  and  $(0, h(0)) \in \partial E_t \cap \overline{\Omega}$ . Let  $V_{\tau_m} = \{(x', x'') : |x'| < \varepsilon', x'' \leq v(x') + \tau_m\}$  and in view of our choice of  $\varepsilon'$ , observe that

$$E_t \cap \{x : |x'| < \varepsilon'\} \subset V_{\tau_m}$$

Observe also that if a point  $(x', v_{\tau_m}(x'))$  is an element of  $\partial E_t \cap \overline{\Omega}$ , then  $|x'| < \varepsilon'$ , for otherwise we would have  $v(x') + \tau_m \leq h(x')$  which would imply that  $\tau_m \leq 0$ . Thus the set  $\partial [E_t \cap \{x : |x'| < \varepsilon'\}] \cap \{(x', v_{\tau_m}(x')) : |x'| < \varepsilon'\}$  is non-empty and according to Theorem 2.2, it is open as well as closed in the connected set  $\{(x', v_{\tau_m}(x')) : |x'| < \varepsilon'\}$ . This implies that

(4.9) 
$$\partial \left[ E_t \cap \{ x : |x'| < \varepsilon' \} \right] \supset \left\{ (x', v_{\tau_m}(x')) : |x'| < \varepsilon' \right\}.$$

Since  $t_m > 0$ , it follows that  $v_{\tau_m}(x') > h(x'')$  whenever  $|x'| = \varepsilon'$ . consequently, using the continuity of  $v_{\tau_m}$ , the graph  $\{(x', v_{\tau_m}(x')), |x'| < \varepsilon'\}$  contains points in  $\mathbb{R}^n \setminus \overline{\Omega}$ , say  $(y', v_{\tau_m}(y')), |y'| < \varepsilon'$ , as well as points in  $\overline{\Omega} \cap B(x_0, \varepsilon)$ , say  $(z', v_{\tau_m}(z')), |z'| < \varepsilon'$ . The point  $(y', v_{\tau_m}(y')), |y'| < \varepsilon'$ , could possibly be an element of  $\mathbb{R}^n \setminus B(x_0, \varepsilon)$ . Consider the line segment,  $\lambda$ , in  $B'(x'_0, \varepsilon')$  that joins y' and z'. Let a' be the closet point to y' on  $\lambda$  with the property that  $(a', v_{\tau_m}(a')) \in \partial\Omega$ . Then all points a on  $\lambda$  that are closer to y' than a' and that are sufficiently close to a' have the property that  $(a, v_{\tau_m}(a)) \in (\mathbb{R}^n \setminus \overline{\Omega}) \cap B(x_0, \varepsilon)$ . Here we have used (4.8) and the continuity of  $v_{\tau_m}$ . In view of (4.9), this implies that  $E_t \cap B(x_0, \varepsilon) \cap (\mathbb{R}^n \setminus \partial\Omega) \neq \emptyset$ , contradicting (4.7). This contradiction was reached under the assumption that  $g(x_0) < t$  and the fact that  $E_t$  is area-minimizing in  $B(x_0, \varepsilon)$ . A similar argument is employed in case  $g(x_0) > t$ .

In order to ultimately identify  $E_t \cap \overline{\Omega}$  as the set  $\{u \ge t\}$  (up to a set of measure zero) for almost all t, we will need the following result.

**4.4 Lemma.** If  $s, t \in T$  with s < t, then  $E_t \subset \subset E_s$ .

*Proof.* We first show that  $E_t \subset E_s$ . Note that

$$(E_s \cap E_t) \setminus \overline{\Omega} = (E_s \setminus \overline{\Omega}) \cap (E_t \setminus \overline{\Omega})$$
$$= (\mathcal{L}_s \setminus \overline{\Omega}) \cap (\mathcal{L}_t \setminus \overline{\Omega})$$
$$= \mathcal{L}_t \setminus \overline{\Omega}$$

and

$$L_t \subset E_t, L_t \subset E_S \Longrightarrow L_t \subset E_s \cap E_t.$$

Thus,  $E_s \cap E_t$  is a competitor with  $E_t$ . Similarly,

Ì

$$(E_s \cup E_t) \setminus \overline{\Omega} = (E_s \setminus \overline{\Omega}) \cup (E_t \setminus \overline{\Omega})$$
$$= (\mathcal{L}_s \setminus \overline{\Omega}) \cup (\mathcal{L}_t \setminus \overline{\Omega})$$
$$= \mathcal{L}_s \setminus \overline{\Omega}$$

and

$$L_t \subset E_t, L_s \subset E_s \Rightarrow L_s \subset E_s \cup E_t.$$

So  $E_s \cup E_t$  is a competitor with  $E_s$ . Then employing (2.8), we have

$$P(E_s) + P(E_t) \le P(E_s \cup E_t, \mathbb{R}^n) + P(E_s \cap E_t, \mathbb{R}^n) \le P(E_s, \mathbb{R}^n) + P(E_t, \mathbb{R}^n),$$

and thus, since  $E_t$  and  $E_s$  are minimizers,

$$P(E_s \cup E_t, \mathbb{R}^n) = P(E_s, \mathbb{R}^n)$$

and

$$P(E_s \cap E_t, \mathbb{R}^n) = P(E_t, \mathbb{R}^n).$$

Reference to (2.21) yields  $|E_s \cup E_t| = |E_s|$ , which in turn implies  $|E_s \setminus E_t| = 0$ . In view of ( )

$$x \in E$$
 if and only if  $\limsup_{r \to 0} \frac{|E \cap B(x,r)|}{|B(x,r)|} > 0$ 

we conclude that  $E_t \subset E_s$ .

Now we come to the crucial part of the argument which is to show that this containment is in fact strict. For this purpose, first note that

(4.10) 
$$E_t \setminus \overline{\Omega} = \mathcal{L}_t \setminus \overline{\Omega} \subset \subset \mathcal{L}_s \setminus \overline{\Omega} = E_s \setminus \overline{\Omega}.$$

Now observe that implies

(4.11) 
$$\partial E_t \cap \partial E_s \cap \partial \Omega = \emptyset.$$

In review of (4.10) and (4.11), it remains to show that

(4.12) 
$$\partial E_t \cap \partial E_s \cap \Omega = \emptyset$$

in order to establish the lemma. For this purpose, let  $S \equiv \partial E_s \cap \partial E_t \cap \Omega$ . Then for  $x_0 \in S$ , there are three possible cases with case (ii) being the central issue of this paper.

- (i) For any  $\varepsilon > 0$ ,  $L_s \cap L_t \cap \Omega \cap B(x_0, \varepsilon)$  is non-empty.
- (ii)  $x_0 \in \overline{L_s}$  and  $B(x_0, \varepsilon) \cap L_t = \emptyset$  for some  $\varepsilon > 0$ .
- (iii)  $B(x_0,\varepsilon) \cap L_t = \emptyset = B(x_0,\varepsilon) \cap L_s$  for some  $\varepsilon > 0$ , thus implying that both  $\partial E_s$  and  $\partial E_t$  are area-minimizing in  $B(x_0,\varepsilon)$ .

Next, we will prove that above 3 cases are impossible, i.e.  $S = \emptyset$ , which implies that  $E_t \subset \subset E_s$ .

For case (i), we can choose some sequence  $\{y_n\} \subset L_s \cap L_t$ , such that  $\lim_{n\to\infty} y_n = x_0$ . Since  $\psi$  is continuous, we have  $\lim_{n\to\infty} \psi(y_n) = \psi(x_0) \ge t$ . Since t > s, there exists an  $\varepsilon > 0$ , such that  $B(x_0, \varepsilon) \subset E_s$  which conradicts the fact that  $x \in \partial E_s$ .

For case (ii), first observe that  $E_s$  is super area-minimizing and that  $E_t$  is areaminimizing near  $x_0$ . Since  $E_t \subset E_s$ , if follows from the maximum principle that  $\partial E_s$  and  $\partial E_t$  agree in a neighborhood of  $x_0$ .

For case (iii), since  $\partial E_s$  and  $\partial E_t$  are area minimizing in  $B(x_0, \varepsilon)$  and  $E_t \subset E_s$ , we apply the maximum principle again to conclude that  $\partial E_t$  and  $\partial E_s$  agree in a neighborhood of  $x_0$ .

Now combining above (i), (ii) and (iii), we conclude that S is area-minimizing and consists only of components of  $\partial E_s$  that do not intersect  $\partial \Omega$ .

Let C be a component of S and sing  $\partial E_s$  the set of singular points of  $\partial E_s$ . Intuitively, C is a bounding cycle that is area-minimizing, which is impossible. Our next step is to rigorously show that C is empty. We divide the proof of this into the following three parts. **Part 1.** There exists an open set  $V \subset \mathbb{R}^n$  such that  $\overline{V} \subset \Omega$  and  $\partial V \supset S$ . For this purpose, we first find an open set U such that

- (i)  $\partial U$  is an (n-1)-manifold with finitely many components,
- (ii)  $\partial U \cap \partial E_s = \emptyset$ ,
- (iii)  $U \cap E_s = C$ ,
- (iv)  $\overline{U} \subset \Omega$ ,
- (v) U is connected.

To find such a set, consider a smooth approximation,  $\rho$ , to the distance function d(x) = d(x, C), i.e., let  $\rho \in C^{\infty}(\mathbb{R}^n \setminus C)$  be such that

$$k^{-1}d(x) \le \rho(x) \le k \, d(x)$$

for all  $x \in \mathbb{R}^n$ , where k is a positive number, cf [Z, Lemma 3.6.1]. Since C is relatively open in  $\overline{\Omega \cap \partial E_s}$ , it follows that  $\partial \{x : \rho(x) < r\} \cap \partial E_s = \emptyset$  for all small values of r. Moreover, by Sard's Theorem and the Implicit Function Theorem,  $\rho^{-1}(r)$  is a smooth (n-1)-manifold for almost all values of r. For any such value of r, let U be the component of  $\{x : \rho(x) < r\}$  that contains C to produce a set satisfying all conditions (i)–(v) above except possibly (iv). By choosing r sufficiently small, this too will be satisfied because  $(\partial E_s \cap \partial E_t) \cap \partial \Omega = \emptyset$ .

Using only the fact that  $\partial U$  is a compact (n-1)-manifold, we invoke Alexander Duality of algebraic topology to conclude that  $\mathbb{R}^n - \partial U$  consists of finitely many components, one more than the number of components in  $\partial U$ , [GH, Theorem 27.10]. Moreover, each component of  $\partial U$  is the boundary of precisely one bounded open set. Note that  $\partial U_{\infty}$  is connected, where  $U_{\infty}$  denotes the unbounded component of  $\mathbb{R}^n - \partial U$ . Indeed, since  $U_{\infty}$  is connected, it is one of the components of  $\mathbb{R}^n - \partial U$ . thus there is a one-to-one correspondence between the bounded components of  $\mathbb{R}^n - \partial U$  and the components of  $\partial U$  which implies that  $\partial U_{\infty}$  is connected.

Since  $\partial U_{\infty}$  is connected, either  $\partial U_{\infty} \subset (E_s)^i$  or  $\partial U_{\infty} \subset (E_s)^e$ , because  $\partial U_{\infty} \cap \partial E_s = \emptyset$ . In case  $\partial U_{\infty} \cap (E_s)^i = \emptyset$ , define V by  $V = (U_{\infty})^e \cap (E_s)^i$ . Since  $(U_{\infty})^e \supset U$  and  $U \cap \partial E_s = C$ , it follows that

$$\partial V = \left(\partial U_{\infty} \cap (E_s)^i\right) \cup \left((U_{\infty})^e \cap \partial E_s\right) \supset C.$$

Similarly, if  $(E_s)^e \cap \partial U_\infty = \emptyset$ , define V by  $V = (U_\infty)^e \cap (E_s)^e$ , so that

$$\partial V = (\partial U_{\infty} \cap (E_s)^e) \cup ((U_{\infty})^e \cap \partial E_s) \supset C.$$

Thus we have established the existence of an open set V that is either a subset of  $(E_s)^i$  or a subset of  $(E_s)^e$  and satisfies

$$\overline{V} \subset \Omega, \partial V \supset C.$$

To finish the proof of the Lemma, we will now show that this leads to a contradiction.

**Part 2.** If  $V \subset (E_s)^i$ , then  $E_s$  is not a minimizer of (2.20).

There are two cases here.

Case 1.  $V \cap L_s = \emptyset$ . This implies that the closed set  $F_s$  defined by  $F_s = \overline{E_s - V}$  is admissible in the minimization problem (2.20).

15

If we can show that

(4.13) 
$$H^{n-1}(\Omega \cap \partial E_s) \ge H^{n-1}(\Omega \cap \partial F_s) + H^{n-1}(\partial V),$$

the desired conclusion is reached since then  $H^{n-1}(\Omega \cap \partial E_s) > H^{n-1}(\Omega \cap \partial F_s)$ , contradicting the minimality of  $H^{n-1}(\Omega \cap \partial E_s)$ .

To establish (4.13), it is sufficient to prove

(4.14) 
$$\partial F_s \cap \partial V = \emptyset$$

since  $\Omega \cap \partial E_s \supset [(\Omega \cap \partial F_s) - (\partial F_s) \cap (\partial V)] \cup \partial V$ . Because  $\partial V = (U_{\infty})^e \cap \partial E_s$ , it follows that for all sufficiently small r > 0,

$$B(x,r) \cap \partial V = B(x,r) \cap \partial E_s.$$

Furthermore, for all small r > 0

$$B(x,r) \cap \overline{V} = B(x,r) \cap E_s.$$

It follows immediately that  $x \notin \partial F_s$  and therefore (4.14) is established.

Case 2.  $V \cap L_s \neq \emptyset$ , but  $V \cap L_t = \emptyset$ . Then replace  $E_s$  and  $F_s$  by  $F_t = \overline{E_t \setminus \overline{V}}$  as in Case 1, follow the same line of proof as in Case 1, we will conclude that  $E_t$  is not a minimizer of (2.20).

Case 3.  $V \cap L_s \neq \emptyset$ ,  $V \cap L_t \neq \emptyset$ . Then there exist components  $L_{s_1}$  and  $L_{t_1}$  of  $L_s$  and  $L_t$  respectively such that  $L_{t_1} \subset L_{s_1} \subset V$ . Since  $\partial V \subset \partial E_s$ , it follows that V is a minimizer of the of the obstacle problem with  $L_{t_1}$  as the obstacle. But this is not possible since  $L_{t_1} \subset V$ .

**Part 3.** If  $V \subset (E_s)^e$ , then  $E_s$  is not a minimizer of (2.20). Let  $G_s = E_s \cup \overline{V}$ , then  $G_s$  is an admissible competitor in (2.20). Now repeat the argument of Part 2 case 1 with  $F_s$  replaced by  $G_s$  to contradict the minimality of  $\partial E_s$ .

We now are in a position to construct the solution u to problem (1.3). For this purpose, we first define for  $t \in T$ ,

$$A_t = \overline{E_t \cap \Omega}.$$

With the help of Lemma 4.4, observe that for  $t \in T$ ,

(4.15)

$$\{q > t\} \subset (E_t)^i \cap \partial\Omega \subset A_t \cap \partial\Omega$$

(4.16)

$$\overline{\{g > t\}} \subset A_t \cap \partial \Omega \subset \overline{E_t} \cap \partial \Omega = [(E_t)^i \cup \partial E_t] \cap \partial \Omega \subset \{g \ge t\}.$$

Finally, note that (4.16) and Lemma 4.3 imply

relative to the topology on  $\overline{\Omega}$  whenever  $s, t \in T$  with s < t. We now define our solution u by

(4.18) 
$$u(x) = \sup\{t : x \in A_t\}.$$

**4.5 Theorem.** The function u defined by (4.18) satisfies the following:

- (i)  $u = g \text{ on } \partial \Omega$
- (ii) u is continuous on  $\overline{\Omega}$ ,
- (iii)  $A_t \subset \{u \ge t\}$  for all  $t \in T$  and  $|\{u \ge t\} A_t| = 0$  for almost all  $t \in T$ . (iv)  $u \ge \psi$  on  $\overline{\Omega}$ .

Proof. To show that u = g on  $\partial\Omega$ , let  $x_0 \in \partial\Omega$  and suppose  $g(x_0) = t$ . If s < t, then G(x) > s for all  $x \in \Omega^c$  near  $x_0$ . Hence,  $x_0 \in (E_s)^i \cap \partial\Omega$  by (4.15) and consequently,  $x_0 \in A_s$  for all  $s \in T$  such that s < t. By (4.18), this implies  $u(x) \ge t$ . To show that u(x) = t suppose by contradiction that  $u(x) = \tau > t$ . Select  $r \in (t, \tau) \cap T$ . Then  $x \in A_r$ . But  $A_r \cap \partial\Omega \subset \{g \ge r\}$  by (4.16), a contradiction since g(x) = t < r.

For the proof of (ii), it is easy to verify that

$$\{u \ge t\} = \{\bigcap A_s : s \in T, s < t\} \text{ and } \{u > t\} = \{\bigcup A_s : s \in T, s > t\}.$$

The first set is obviously closed while the second is open relative to  $\overline{\Omega}$  by (4.17). Hence, u is continuous on  $\overline{\Omega}$ .

For (iii), it is clear that  $\{u \ge t\} \supset A_t$ . Now,  $\{u \ge t\} - A_t \subset u^{-1}(t)$ . But  $|u^{-1}(t)| = 0$  for almost all t because  $|\Omega| < \infty$ .

In (iv), it is sufficient to show  $u(x_0) \ge \psi(x_0)$  for  $x_0 \in \Omega$ . Let  $t = u(x_0)$  and  $r = \psi(x_0)$  and suppose t < r. Then  $x_0 \in L_{r'} \subset E_{r'}$  for t < r' < r. But then,  $x_0 \notin A_{r'}$  by the definition of u, a contradiction.

**4.6 Theorem.** If  $\Omega$  is a bounded Lipschitz domain that satisfies (4.1) and (4.2), then the function u defined by (4.18) is a solution to (1.3).

Proof. Let  $v \in BV(\Omega)$ , v = g on  $\partial\Omega$  be a competitor in problem (1.3). We recall the extension  $G \in BV(\mathbb{R}^n - \overline{\Omega})$  of g, (2.16). Now define an extension  $\overline{v} \in BV(\mathbb{R}^n)$ of v by  $\overline{v} = G$  in  $\mathbb{R}^n - \overline{\Omega}$ . Let  $F_t = \{\overline{v} \ge t\}$ . It is sufficient to show that

$$(4.19) P(E_t, \Omega) \le P(F_t, \Omega)$$

for almost every  $t \in T$  (see (2.19)), because then  $v \in BV(\Omega)$  and (2.11) would imply

$$\int_{a}^{b} P(E_{t}, \Omega) dt \leq \int_{-\infty}^{\infty} P(F_{t}, \Omega) dt = \|\nabla v\|(\Omega) < \infty.$$

Hence, by (2.12),  $u \in BV(\Omega)$ ; furthermore,  $\|\nabla u\|(\Omega) \le \|\nabla v\|(\Omega)$  by (2.11).

We know that  $E_t$  is a solution of

(4.20) 
$$\min\left\{P(E,\mathbb{R}^n): E\setminus\overline{\Omega}=\mathcal{L}_t\setminus\overline{\Omega}, \ E\supset L_t\right\},\$$

while  $F_t - \overline{\Omega} = \mathcal{L}_t - \overline{\Omega}$  and  $F_t \supset L_t$ . Hence,

$$(4.21) P(E_t, R^n) \le P(F_t, R^n).$$

Next, note that

(4.22) 
$$P(E_t, R^n) = H^{n-1}(\partial_* E_t - \overline{\Omega}) + H^{n-1}(\partial_* E_t \cap \partial \Omega) + H^{n-1}(\partial_* E_t \cap \Omega)$$
$$\geq H^{n-1}(\partial_* \mathcal{L}_t - \overline{\Omega}) + P(E_t, \Omega).$$

We will now show that

(4.23) 
$$P(F_t, R^n) = H^{n-1}(\partial_* \mathcal{L}_t - \overline{\Omega}) + H^{n-1}(\partial_* F_t \cap \Omega)$$
$$= H^{n-1}(\partial_* \mathcal{L}_t - \overline{\Omega}) + P(F_t, \Omega),$$

which will establish (4.19) in light of (4.21) and (4.22).

Observe

$$P(F_t, R^n) = H^{n-1}(\partial_* \mathcal{L}_t - \overline{\Omega}) + H^{n-1}(\partial_* F_t \cap \partial\Omega) + H^{n-1}(\partial_* F_t \cap \Omega).$$

We claim that  $H^{n-1}(\partial_* F_t \cap \partial \Omega) = 0$  for almost all t because  $\partial_* F_t \subset \partial F_t \subset \overline{v}^{-1}(t)$ since  $\overline{v} \in C^0(\mathbb{R}^n)$ . But  $H^{n-1}(\overline{v}^{-1}(t) \cap \partial \Omega) = 0$  for all but countably many t since  $H^{n-1}(\partial \Omega) < \infty$ .

#### 5. Modulus of continuity of the solution

**5.1 Lemma.** Suppose  $\Omega$  is a bounded, open subset of  $\mathbb{R}^n$  whose boundary is  $C^2$  with mean curvature bounded below by a > 0. Assume  $g \in C^{0,\alpha}(\partial\Omega)$ , and  $\psi \in C^{0,\alpha/2}(\Omega)$  for some  $0 < \alpha \leq 1$ . Let  $u \in C^0(\overline{\Omega}) \cap BV(\Omega)$  be a solution to (4.1). Then, there exist positive numbers  $\delta$  and C depending only on a,  $\|g\|_{C^{0,\alpha}(\partial\Omega)}$ ,  $\|g\|_{C^{0,\alpha}(\partial\Omega)}$ ,  $\|g\|_{C^{0,\alpha/2}(\Omega)}$  and  $\|u\|_{C^{0}(\Omega)}$  such that

$$|u(x) - u(x_0)| \le C |x - x_0|^{\alpha/2}$$

wherever  $x_0 \in \partial \Omega$  and  $x \in \overline{\Omega}$  with  $|x - x_0| < \delta$ .

*Proof.* For each  $x_0 \in \partial \Omega$  we will construct functions  $\omega^+, \omega^- \in C^0(\overline{U})$  where  $U(x_0, \delta) := B(x_0, \delta) \cap \Omega$  and  $\delta > 0$  is sufficiently small, such that

(i)  $\omega^+(x_0) = \omega^-(x_0) = g(x_0),$ (ii) For  $x \in U(x_0, \delta)$ 

$$|\omega^{+}(x) - g(x_{0})| \le C |x - x_{0}|^{\alpha/2}$$
$$|\omega^{-}(x) - g(x_{0})| \le C |x - x_{0}|^{\alpha/2},$$

(iii)  $\omega^{-} \leq u \leq \omega^{+}$  in  $U(x_0, \delta)$ 

We begin with the construction of  $\omega^-$ . To this end, let  $d(x) = \text{dist}(x, \partial\Omega)$ . Since  $\partial\Omega \in C^2$  recall that  $d \in C^2(\{x : 0 \le d(x) < \delta_0\})$  for some  $\delta_0 > 0$ . Furthermore, since  $\partial\Omega$  has positive mean curvature and  $|\nabla d| = 1$ , it follows that

(5.1) 
$$\operatorname{div}\left(\frac{\nabla d}{|\nabla d|}\right) = \Delta d \le -a,$$

in  $\{x: 0 \le d(x) < \delta_0\}$  for some a > 0. For each  $\varepsilon > 0$ , set

$$v(x) = |x - x_0|^2 + \lambda d(x)$$
  

$$\omega^{-}(x) = \max\{\psi, -Kv^{\alpha/2}(x) + g(x_0)\},\$$

where  $\lambda > 0$  is to be determined later. Clearly (i) is satisfied.

Next, in the open set  $\{\omega^- > \psi\}$ , observe that

$$\begin{aligned} \nabla \omega^{-} &| = K \frac{\alpha}{2} v^{\frac{\alpha}{2} - 1} |\nabla v|, \\ &| \nabla v | = |2(x - x_0) + \lambda \nabla d| \ge \lambda |\nabla d| - 2 |x - x_0| \\ &= \lambda - 2 |x - x_0| > 0, \end{aligned}$$

provided we choose  $\delta$  and  $\lambda$  such that  $\lambda > 2\delta$ . Further, we note that

$$\operatorname{div}\left(\frac{\nabla\omega^{-}}{|\nabla\omega^{-}|}\right) = -\operatorname{div}\left(\frac{\nabla v}{|\nabla v|}\right) = \frac{-1}{|\nabla v|^{3}}Av,$$

where  $Av = |\nabla v|^2 \Delta v - D_i v D_j v D_{ij} v$ . Finally, observe that Av < 0 for  $\lambda$  sufficiently large and  $\delta$  sufficiently small. Indeed, using  $D_i dD_{ij} d = 0$  for any j, one readily obtains

$$Av = |\nabla v|^2 (\lambda \Delta d + 2(n-1)) - 4\lambda (x - x_0)_i (x - x_0)_j D_{ij} d$$

and

$$|\nabla v|^2 = \lambda^2 + 4 |x - x_0|^2 + 4\lambda(x - x_0) \cdot \nabla d$$

so that

$$Av \leq -a\lambda^3 + O(\lambda^2)$$
, as  $\lambda \to \infty$ 

uniformly for  $\delta < \delta_0$ .

Clearly, we can choose K sufficiently large so that  $\omega^- = \psi$  on  $\partial U(x_0, \delta)$  and that (ii) is satisfied, where K depends only on  $\|g\|_{C^{0,\alpha}(\partial\Omega)}, \|g\|_{C^0(\partial\Omega)}, \|\psi\|_{C^{0,\alpha/2}(\Omega)}$  and  $\|u\|_{C^0(\Omega)}$ . Also, on  $\Delta := \{\omega^- > u\} \cap U(x_0, \delta)$ , we have  $\omega^- = -Kv^{\alpha/2} + g(x_0)$  and therefore

(5.2) 
$$|\nabla \omega^-| > 0 \text{ and } \operatorname{div}\left(\frac{\nabla \omega^-}{|\nabla \omega^-|}\right) > 0 \text{ on } \Delta.$$

We now proceed to show that  $\Delta = \emptyset$ , which will establish the first of the inequalities in (iii). For this purpose, note that  $\omega^- \in BV(\Delta)$ . Next, for t > 0, let  $\Delta_t := \{\omega^- - t > u\}$  and note that

(5.3) 
$$\Delta = \bigcup_{t>0} \Delta_t \qquad \Delta_t \subset \subset \Delta \subset \Omega.$$

Let  $\omega^* := \max(u, \omega^- - t)$  and note that  $\omega^* \in BV(\Omega) \cap C^0(\overline{\Omega})$  since  $\omega^- - t = \psi - t < u$ on  $\partial \Delta_t$ . For all but countably many t > 0, it follows from basic measure theory that

(5.4) 
$$\|\nabla \omega^*\| (\partial \Delta_t) = 0 = \|\nabla u\| (\partial \Delta_t).$$

For the remainder of this argument, we will consider only such t. Since  $\omega^* \ge u \ge \psi$ , it follows that

(5.5) 
$$\|\nabla u\|(\Omega) \le \|\nabla \omega^*\|(\Omega).$$

Now let  $\eta \in C_0^{\infty}(\Delta)$  satisfy  $\eta = 1$  on  $\Delta_t$  and  $0 \le \eta \le 1$  in  $\Delta$ . Set

$$h = \eta \frac{\nabla \omega^-}{|\nabla \omega^-|}$$

18

so that  $h \in [C_0^1(\Delta)]^n$ . Since  $\omega^* = u$  on  $\Delta - \Delta_t$ , it follows from

$$\int_{\Delta} u \operatorname{div} h \, dx = -\nabla u(h),$$

and

$$\int_{\Delta} \omega^* \operatorname{div} h \, dx = -\nabla \omega^*(h),$$

that

$$\int_{\Delta} u - \omega^* \, dx = \int_{\Delta_t} (u - \omega^- + t) \, \operatorname{div} h \, dx = [\nabla(\omega^* - u)](h).$$

It follows from (5.4) and the definition of the BV norm that

$$\|\nabla\omega^*\| \left(\partial\Delta_t\right) \le \|\nabla u\| \left(\partial\Delta_t\right) + \int_{\partial\Delta_t} |\nabla\omega^*| \ dx = 0,$$

so that

$$\int_{\Delta_t} (u - \omega^* + t) \operatorname{div} h \, dx = \nabla \omega^* (h \chi_{\Delta_t}) - \nabla u (h \chi_{\Delta_t})$$
$$\geq \int_{\Delta_t} |\nabla \omega^*| \, dx - \|\nabla u\| (\Delta_t).$$

Since  $u - \omega^- + t < 0$  and div h > 0 on  $\Delta_t$ , we have

$$\int_{\Delta_t} |\nabla \omega^*| \, dx < \|\nabla u\| \, (\Delta_t).$$

That is,

$$\left\|\nabla\omega^{*}\right\|\left(\Delta_{t}\right) < \left\|\nabla u\right\|\left(\Delta_{t}\right).$$

Since  $\omega^* = u$  on  $\mathbb{R}^n \setminus \Delta_t$ , we obtain from (5.4) that  $\|\nabla \omega^*\|(\Omega) < \|\nabla u\|(\Omega)$ , which contradicts (5.5). Thus we conclude that  $\omega^- \leq u$  on  $U(x_0, \delta)$ .

The proof of the second inequality in (iii) is obtained by a similar argument using  $\omega^+(x) := K v^{\alpha/2}(x) + g(x_0)$ .

**5.2 Theorem.** Suppose  $\Omega$  is a bounded, open subset of  $\mathbf{R}^n$  with  $C^2$  boundary having mean curvature bounded below by a > 0. Suppose  $g \in C^{0,\alpha}(\partial\Omega)$ , and  $\psi \in C^{0,\alpha/2}$  for some  $0 < \alpha \leq 1$ . If  $u \in C^0(\overline{\Omega}) \cap BV(\Omega)$  is a solution to (4.1), then  $u \in C^{0,\alpha/2}(\overline{\Omega})$ .

*Proof.* For s < t, consider the superlevel sets  $E_s$ ,  $E_t$  of u and assume that  $\operatorname{dist}(\partial E_s, \partial E_t) = |y - x|$  where  $x \in E_t$  and  $x \in E_s$ . Assume t - s is small enough to ensure that  $|y - x| < \delta$ , where  $\delta$  is given by Lemma 5.1. Observe that  $L_t \subset E_t \subset \subset E_s$ . Theorems 4.5 and 4.6 imply that u is continuous on  $\overline{\Omega}$  and therefore bounded. Hence it is sufficient to show that  $|u(y) - u(x)| = |t - x| \leq C |x - y|^{\alpha/2}$  whenever  $|y - x| < \delta$ . This will be accomplished by examining the following five cases.

(i) If either x or y belongs to  $\partial\Omega$ , then our result follows from Lemma 5.2. (ii)  $y \in \partial E_s \setminus \overline{L_s}$  and  $x \in \partial E_t \setminus \overline{L_t}$ : Let  $[\partial E_t]_v$  denote the translation of  $\partial E_t$  by the vector v := (y - x) |y - x|. Since both  $\partial E_s$  and  $\partial E_t$  are area-minimizing in some neighborhoods of y and x respectively, we can apply Theorem 3.3 to conclude that

19

 $\partial E_s$  and  $[\partial E_t]_v$  agree on some connected component of  $\partial E_s$ , say S, that contains y.

If  $S \cap \partial \Omega = \emptyset$ , it follows that S is area-minimizing in some open set  $U \supset S$ . Now let S' be a component of the set of regular points of S. We first show that S' is a cycle in the sense of currents; that if, we wish to show that

(5.6) 
$$\int_{S'} \mathrm{d}\varphi = 0$$

whenever  $\varphi$  is a smooth (n-2)-form supported in B(0,R) where B(0,R) is the ball having the property that  $\Omega \subset B(0,R)$ . Since S' is area-minimizing in  $\Omega$ , we appeal to the monotonicity formula [S1, §17.6] to conclude that only a finite number of components of  $(\partial E_s)$  can intersect any given compact subset of  $\Omega$ , in particular, spt  $\varphi \cap \overline{S'}$ . Thus, there exists a smooth function  $\zeta$  that is 1 on spt  $\varphi \cap \overline{S'}$ and 0 in a neighborhood of  $\partial E_s - \overline{S'}$ . Then, (5.6) is established by

$$\int_{S'} \mathrm{d} \varphi = \int_{S'} \mathrm{d} (\zeta \varphi) = \int_{\partial E_s} \mathrm{d} (\zeta \varphi) = 0$$

Thus, S' is an (n-1)-rectifiable cycle in the sense of currents; that is,  $\partial S' = 0$ . Now appeal to [S1, 27.6] to conclude that there is a measurable set  $F \subset B(0, R)$  such that  $\partial F = S'$ . It follows from elementary considerations that for a given vector  $\nu \in \mathbb{R}^n$ , there is a hyperplane, P, with normal  $\nu$  such that  $P \cap \overline{S'} \neq \emptyset$  and

$$F \subset \{x : (x - x_0) \cdot \nu \le 0\}$$

where  $x_0 \in P \cap \overline{S'}$ . Theorem 3.3 implies  $P \cap \overline{S'}$  is open as well as closed in P, thus leading to a contradiction in case  $S \cap \partial \Omega = \emptyset$ . If  $S \cap \partial \Omega \neq \emptyset$ , we then are led to a situation covered by (i).

(iii)  $y \in \partial E_s \setminus \overline{L_s}, x \in \partial E_t \cap \overline{L_t}$ : Since  $L_s \supset L_t$ , there exists  $y' \in \partial L_s$  such that y' - x = c(y - x), 0 < c < 1, and therefore

(5.7) 
$$|u(y) - u(x)| = |t - s| = |\psi(y') - \psi(x)| \le C |y' - x|^{\alpha} \le C |y - x|^{\alpha}.$$

(iv)  $y \in \partial E_s \cap \overline{L_s}, x \in \partial E_t \cap \overline{L_t}$ : This is treated as in (5.7).

(v)  $y \in \partial E_s \cap \overline{L_s}, x \in \partial E_t \setminus \overline{L_t}$ : In this case we can apply Corollary 3.4 to obtain an area minimizing connected component S of  $\partial E_s$  which can be treated as in (ii) above.

# 6. A MONOTONICITY PRINCIPLE FOR SUPERMINIMIZING SETS

An issue left open in our development is whether the regularity requirement  $\overline{E} \cap U = \overline{E^i} \cap U$  is necessary in Theorem 3.3, the extended maximum principle for sub and superminimizing sets.

This suggests the question, of interest in its own right, of what regularity, if any, is enjoyed by (sub)superminimizing sets. For example, do (sub)superminimizers have tangent cones? Are they  $C^1$  or analytic  $H^{n-1}$  almost-everywhere? And, the question begged by Theorem 3.3, is a subminimizer necessarily the closure of its interior? In the next section, we will give an explicit example showing that the last conjecture is false. In this section, we present some preliminary results in the direction of regularity, consisting of a new monotonicity principle and consequent one-sided mass bound for (sub)superminimizing sets.

Let  $B_r = B(0, r)$  denote the ball of radius r about the origin in  $\mathbb{R}^n$ . Let F be a superminimizing set in U, and without loss of generality, assume  $B_1 \subset U$ .

**6.1 Lemma.** Let  $A = \{x \in A^c : \text{the metric density of } A \text{ is one at } x\}$ . Then,  $H^{n-1}(\partial B_r \cap \tilde{A}) = 0$  for almost all r.

*Proof.* The Lebesgue measure of  $\tilde{A} \cap B_1$  is zero. But, by the co-area formula, (2.11), it is also equal to  $\int_0^1 H^{n-1}(\partial B_r \cap \tilde{A}) dr$ , whence the result follows.  $\square$ 

**6.2 Lemma.** Let E area subminimizing in U,  $B_1 \subset U$ , and r such that  $H^{n-1}(\partial B_r \cap$  $\tilde{E}$ ) = 0. Then,  $P(E, B_r) \leq H^{n-1}(E \cap \partial B_r)$ .

*Proof.* The set  $G = E \setminus B_r$  is a competitor to E. Exterior to  $B_r$ , G has the same reduced boundary as does E, but interior to  $B_r$ , it has no reduced boundary. On  $\partial B_r$ , G has reduced boundary contained in the set of points at which E has density one, which by assumption is contained in E except for a set of  $H^{n-1}$ -measure zero.

Therefore, by the subminimality of E, we have

$$0 \le P(E \setminus B_r, U) - P(E, U) \le H^{n-1}(E \cap \partial B_r) - P(E, B_r)$$

giving the result.

Define the dimension-dependent constant  $0 < \delta(n) < 1/2$  by

$$\delta(n) = |D_1|/|B_1|,$$

where  $D_1 \subset B_1$  is a set bounded by a hemispherical cap of radius one, orthogonal to  $\partial B_1$ .

**6.3 Lemma.** If 
$$|A \cap B_r| / |B_r| \le \delta(n)$$
, then  $P(A, B_r) / H^{n-1}(\partial B_r) \ge |A \cap B_r| / |B_r|$ .

**Remark.** Another way of stating this result is that  $P(A, B_r) > (n/r)|A \cap B_r|$ . It could also be rephrased as an isoperimetric inequality.

*Proof.* By rearrangement, we find that the set D of minimum perimeter  $P(D, B_r)$ subject to  $|D \cap B_r| = |A \cap B_r|$  is the set bounded by a hemispherical cap meeting  $\partial B$  orthogonally. Trivially, we have

$$(6.1) P(D,B) \le P(A,B).$$

Let  $D_r$  be the set bounded by a spherical cap of radius r, intersecting  $\partial B_r$ orthogonally, so that  $|D \cap B_r|/|B_r| = \delta(n)$ . Since  $|D \cap B_r|/|B_r| = |A \cap B_r|/|B_r| \leq 1$  $\delta(n)$ , we thus have that  $|D| \leq |D_r|$  and so the radius of the hemispherical cap bounding D is less than or equal to r. It follows by elementary geometry that

(6.2) 
$$H^{n-1}(\partial D \cap \partial B) \le P(D,B).$$

(To see this, e.g., one can reflect the hemispherical cap D about the plane of its intersection with  $B_r$ , to obtain a surface oriented in the same direction as the patch  $\overline{D} \cap \partial B_r$  and containing the patch in its interior. Since the patch has positive mean curvature, it follows that this outer surface has greater area than does  $\overline{D} \cap \partial B_r$ .)

But, D is entirely contained in the cone C from  $\partial D \cap \partial B_r$  to the center of  $B_r$ and tangent to D at  $\partial B_r$ . That is,  $|A \cap B_r| \leq |C|$ . On the other hand, the volume ratio  $|C|/|B_r|$  for a cone is exactly its surface ratio,  $H^{n-1}(\partial D \cap \partial B_r)/H^{n-1}(\partial B_r)$ . Combining these facts with (6.2) and (6.1), we have

$$\begin{aligned} |A|/|B_r| &\leq |C|/|B_r| = H^{n-1}(\partial D \cap \partial B_r)/H^{n-1}(\partial B_r) \\ &\leq P(D, B_r)/H^{n-1}(\partial B_r) \leq P(A, B_r)/H^{n-1}(\partial B_r), \end{aligned}$$
  
s to our desired conclusion.

which leads to our desired conclusion.

We now prove our main result, a *volume* monotonicity principle for superminimizing sets.

**6.4 Proposition.** Let E be subminimizing in U,  $B_1 \subset U$ . If  $|E \cap B_1|/|B_1| < \delta(n)$ (  $0 < \delta(n) < 1/2$  as defined above 6.2), then the ratio  $|E \cap B_r|/|B_r|$  is increasing in r for  $0 \le r \le 1$ .

Proof. From Lemmas 6.1 and 6.2, we have

 $H^{n-1}(E \cap \partial B_r))/H^{n-1}(\partial B_r) \ge P(E, B_r)/H^{n-1}(\partial B_r) \ge |E \cap B_r|/|B_r)|$ 

for almost all r, so long as  $|E \cap B_r|/|B_r| < \delta(n)$ .

By the co-area formula, (2.11),

$$(d/dr)|B_r| = H^{n-1}(\partial B)$$
 and  $(d/dr)|E| = H^{n-1}(E \cap \partial B_r)$ 

Thus,

$$d|E \cap B_r|/d|B_r| = H^{n-1}(E \cap \partial B_r)/H^{n-1}(\partial B) \ge |E \cap B_r|/|B_r|$$

giving monotonicity so long as  $|E \cap B_r|/|B_r| < \delta(n)$ . But, because of monotonicity, this property persists for all  $0 \le r \le 1$ .

This property has many implications. Among them is the following important one, a one-sided bound on the average density.

**Proposition 6.5.** Let *E* be subminimizing in *U*,  $B_1 \subset U$ . If  $0 \in \partial E$ , then  $|E \cap B_1|/|B_1| \ge \delta(n)$ .

*Proof.* Suppose to the contrary that  $|E \cap B_1|/|B_1| < \delta(n)$ . Then, for some R < 1,  $|E \cap B(x,R)| < \delta(n)$  for every  $x \in B_{1-R}$ . By the monotonicity property of Proposition 6.4, we thus have  $|E \cap B(x,r)|/|B(x,r)| < \delta(n)$  for  $r \leq R$ . Thus,

$$|E \cap \ddot{B}|/|\ddot{B}| < \delta(n) < 1/2$$

for any ball contained in  $B_{1-R}$ ; hence the density of E is strictly less than 1/2 at each point of  $B_{1-R}$ .

But, since the density of E must be zero or one at almost every point of  $B_{1-R}$ , the density of E must be zero at almost every point in  $B_{1-R}$ , and therefore  $|E \cap B_{1-R}| = 0$ . But, by our convention in choosing set representatives, this would imply that  $B_{1-R} \subset E^e$ , in particular  $0 \in E^e$ , a contradiction.

**6.6 Corollary.** If E is subminimizing, then  $\overline{E_m^i} = (E_m^e)^c = \overline{E}$ .

*Proof.* By Proposition 6.5, the density of E at any  $x \in \partial E$  is strictly greater than 0, hence  $\partial E \cap E_m^e = \emptyset$ . It follows that  $\partial E$ , and therefore  $\overline{E}$  as well, is contained in  $(E_m^e)^c \subset \overline{E_m^i}$ . Since  $\overline{E_m^i}$  is always contained in  $\overline{E}$ , we thus obtain

$$\overline{E_m^i} = (E_m^e)^c = \overline{E},$$

as claimed.

**6.7 Corollary.** Let E be minimizing in U and  $x \in \partial E$ . Then, in any ball  $B(x,r) \subset U$ , the relative volume fractions of E and  $E^c$  are bounded below by  $\delta(n) > 0$ .

*Proof.* By the previous Proposition applied to E and  $E^c$ , we find that violation of this bound would imply that x were in the interior of E or of  $E^c$ . But,  $x \in \partial E$  by assumption, a contradiction.

**6.8 Corollary.** Let E be minimizing in U and  $x \in \partial E$ . Then, in any ball  $B(x,r) \subset U$ ,  $P(E, B_r) \geq \delta r^{n-1}$ , where  $\delta > 0$  is an independent constant.

*Proof.* This follows from Corollary 6.7 plus the explicit form of the minimizer of  $P(A, B_r)$  among sets with |A| = |E|.

**Remark.** Propositions 6.4 and 6.5 give an alternative, and more elementary route to regularity of minimizing sets than the usual path via the Isoperimetric Theorem for minimal surfaces, cf., [Gi, Chapter 8]. Using Corollary 6.8, one can go on to show existence of tangent cones, etc. This standard result is usually proved by reference to the Isometric Theorem for minimal surfaces, cf. [Gi, Chapter 5].

#### 7. "Foamy" sets.

We conclude by demonstrating existence of sparse, "foamy" superminimizing sets having topological boundary with positive Lebesgue measure, thus indicating possible limitations of a regularity theory for (sub)superminimizing sets.

For  $\overline{B}(x_1, r), \overline{B}(x_0, R) \subset U \subset \mathbb{R}^2, \overline{B}(x_1, r) \cap \overline{B}(x_0, R) = \emptyset$ , consider the obstacle problem

(7.1) 
$$\inf\{P(F,U): B(x_1,r) \cup B(x_0,R) \subset F \subset U\}.$$

**7.1 Lemma.** For r sufficiently small, the solution of (7.1) is

$$E = B(x_1, r) \cup B(x_0, R).$$

Moreover, for any connected set  $\tilde{F}$  containing  $B(x_1, r) \cup B(x_0, R)$ , there holds

(7.2) 
$$P(\tilde{F}, U) > P(F, U) + \delta,$$

for some  $\delta > 0$ .

*Proof.* Without loss of generality, take U to be all of  $\mathbb{R}^2$ . Since we are in two dimensions, minimal surfaces for (7.1) are easily characterized as arcs of  $\partial B(x_0, R)$ ,  $\partial B(x_1, r)$  joined by straight lines. By explicit comparison, it is then found that the connected competitor  $\tilde{F}$  with least perimeter is the convex hull of  $\partial B(x_0, R)$ ,  $\partial B(x_1, r)$ , which for r sufficiently small satisfies (7.2). Among disconnected competitors, the best is  $F = B(x_1, r) \cup B(x_0, R)$ , by (2.13).

**7.2 Proposition.** For any open  $V \subset U \subset \mathbb{R}^2$ , and any  $\varepsilon > 0$ , there exists a superminimizing set F in U such that  $\overline{F} = \overline{V}$  and  $|F| \leq \pi \varepsilon^2$ .

*Proof.* Enumerate the rationals as  $\{x_j\}$ .

**Claim:** For suitably chosen  $r_j$ ,

$$F_J := \bigcup_{j < J} B(x_j, r_j)$$

has the properties:

(i) Any set  $F_J \subset G \subset U$  with a connected component containing two  $B(x_j, r_j)$  with  $j \leq J$ , satisfies

$$P(G,U) > P(F_J,U) + \delta_J, \quad \delta_J > 0,$$

(ii)

(7.3) 
$$\sum_{j=J+1}^{\infty} P(B(x_j, r_j) < \delta_J.$$

**Proof of claim:** The radii  $r_i$  may be chosen inductively, as follows:

Choose  $r_1 < \varepsilon/2$  sufficiently small that  $B(x_1, r_1) \subset V$ . If  $x_{j+1} \in F_j$ , then take  $r_{j+1} = 0$ . Otherwise, choose  $r_{j+1}$  so small that  $B(x_{j+1}, r_{j+1}) \subset V \setminus F_j$ ,

(7.4) 
$$P((B(x_{j+1}, r_{j+1}), U) < \delta_j/2,$$

and, by Lemma 7.1, any connected set G containing  $B(x, r_{i+1})$  and any  $B(x_k, r_k)$ ,  $k \leq j$  satisfies

(7.5) 
$$P(G,U) > P(B(x_{j+1},r_{j+1}),U) + P(B(x_k,r_k),U) + \delta_{j+1}$$

for some  $\delta_{j+1} > 0$ . By (7.4), (ii) is clearly satisfied. Further, (7.4) and (7.5) together give (i). For, if G has a component containing any  $B(x_k, r_k)$ ,  $B(x_l, r_l)$ ,  $k \neq l \leq j$ , then (7.3) holds by the induction hypothesis. Likewise, if no component of G contains  $B(x_{i+1}, r_{i+1})$  and any  $B(x_k, r_k), k \leq j$ . The remaining case is that precisely one  $B(x_k, r_k), k \leq j$ , lies in a component with  $B(x_{j+1}, r_{j+1})$ , and the rest lie each in distinct components. In this case, (7.3) follows by (7.5) and (2.13).

Defining  $F := \bigcup_i B(x_i, r_i)$ , we find that F is superminimizing in U. For, let G be any competitor. If G has any component containing  $B(x_i, r_i)$  and  $B(x_k, r_k)$ , j < k, then (i)-(ii) together give

$$P(G,U) > P(F_k,U) + \delta_k$$
  
>  $P(F_k,U) + \sum_{k+1}^{\infty} P(B(x_j,r_j),U)$   
 $\geq P(F,U).$ 

On the other hand, if each  $B(x_i, r_i)$  lies in a distinct component  $G_i$  of G, then either  $G_j \equiv B(x_j, r_j)$ , or, by the Isoperimetric Theorem,  $P(G_j, U) \geq P(B(x_j, r_j), U)$ , with strict inequality for some J. Noting that  $P(G, U) \geq \sum_{j=1}^k P(G_j, U)$  for any finite sum, and recalling (ii), we thus obtain P(G,U) > P(F,U) as claimed. By (ii), and the choice  $r_1 < \varepsilon$ , we have  $|F| \le \pi \varepsilon^2 \sum_{j=1}^{\infty} (1/2)^{2j} < \pi \varepsilon^2$ . But,

clearly, also, F is dense in V, giving  $\overline{F} = \overline{V}$  as claimed.  $\square$ 

**Remark.** It is not clear whether such a construction can be carried out in higher dimensions, since Lemma 7.1 no longer holds with positive  $\delta$ .

#### **Consequences:**

1. The construction of Proposition 7.2 shows that in general  $\overline{E} = \overline{E^i}$  is false for subminimizing sets E, in contrast to the result of Corollary 6.7. It would seem that some form of connectivity must be assumed on E, if this property is to hold.

2. A similar construction with U = B(0,1) yields a superminimizing set G contained in and dense in the lower hemisphere  $B^{-}(0,1) := \{x : s \in B(0,1), x_n \leq 0\}$ 0}. Taking  $E := G^c$ ,  $F = B^-(0, 1)$ , we find that the strong maximum principle as

24

25

stated in Theorem 2.2 is violated. However, we remark that in the original form as stated in [S2], the conclusion of the theorem was that  $\partial E$  and  $\partial F$  should agree on their components of  $x_0$ . This version of the theorem remains valid also for the above example, though the two statements are equivalent for minimizing sets. Here, again,  $\overline{E} \neq \overline{E^i} = \emptyset$ , violating the regularity assumption of 3.3.

Evidently, the issue of a maximum principle for sub- and superminimizing sets is a delicate one, requiring ideas beyond those in this paper. This would appear to be an interesting area for further study.

#### WILLIAM P. ZIEMER KEVIN ZUMBRUN

#### References

- [AH] Adams, D. R. and Hedberg, L. I., Function spaces and potential theory, Springer Verlag, 1996.
- [BMo] Biroli, M. and Mosco, U., Wiener criterion and potential estimates for obstacle problems relative to degenerate elliptic operators, Ann. Mat. Pura Appl. 159 (1991), 255–281.
- [BDG] Bombieri, E., E. De Giorgi, E. Giusti, Minimal cones and the Bernstein problem, Invent. Math. 7 (1969), 255-267..
- [CL] Choe, H. J. and Lewis, J. L., On the obstacle problem for quasilinear elliptic equations of p Laplacian type, SIAM J. Math. Anal. 22 (1991), 623–638.
- [Fe1] Federer, H., Curvature measures, Trans. Amer. Math. Soc. 93 (1959), 418-491.
- [Fe2] \_\_\_\_\_, Geometric measure theory, Springer Verlag, New York, Heidelberg, 1969.
- [FR] Fleming, W. H., and R. Rishel, An integral formula for total gradient variation, Arch. Math. 11 (1960), 218-222.
- [FM1] Frehse, J. and Mosco, U., Variational inequalities with one-sided irregular obstacles, Manuscripta Math 28 (1979), 219–233.
- [Gi] Giusti, E., Minimal surfaces and functions of bounded variation, Birkhäuser, 1985.
- [HKM] Heinonen, J., Kilpeläinen, T. and Martio, O., Nonlinear potential theory of degenerate elliptic equations., Oxford University Press, Oxford,, 1993.
- [Li] Lieberman, G., Regularity of solutions to some degenerate double obstacle problems, Indiana Univ. Math. J. 40 (1991), 1009–1028.
- [MaZ] Malý, Jan and Ziemer, W. P., Fine regularity of elliptic equations, Mathematical Surveys and Monographs, Vol. 51, American Mathematical Society., 1997.
- [Mo] Moschen, Maria Pia, Principio di massimo forte per le frontiere di misura minima, Ann. Univ. Ferrara, Sez. VII, Sc. Mat. 23 (1977), 165-168.
- [MuZ] Mu, Jun and Ziemer, W. P., Smooth regularity of solutions of double obstacle problems involving degenerate elliptic equations, Comm. Partial Diff. Equations 16 (1991), 821– 843.
- [MiZ] Michael, J. and Ziemer, W. P., Existence of solutions to nonlinear obstacle problems, Nonlinear Analysis, TMA 17 (1991), 45–73.
- [S1] Simon, L., Lectures on geometric measure theory, vol. 3, Proc. Centre Math. Analysis, ANU, 1983.
- [S2] \_\_\_\_\_, A strict maximum principle for area minimizing hypersurfaces, J. Diff. Geometry 26 (1987), 327-335.
- [SZ] Sternberg, P. and Ziemer, W. P., The Dirchlet problem for functions of least gradient, IMA Volumes in Math. and Appl. 47 (1993), 197-214.
- [SWZ] Sternberg, P., Williams, G. and Ziemer, W. P., Existence, uniqueness, and regularity for functions of least gradient, J. f
  ür die Reine und Angewandte Mathematik 430 (1992), 35-60.
- Ziemer, W. P., Weakly differentiable functions: Sobolev spaces and functions of bounded variation, Springer-Verlag, Graduate Texts in Math., New York, 1989.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405-5701