

# ON AN ANALOG OF SELBERG'S EIGENVALUE CONJECTURE FOR $SL_3(\mathbf{Z})$

Sultan Catto<sup>1</sup>

*The Graduate School and University Center, and Baruch College  
The City University Of New York  
17 Lexington Avenue  
New York, NY 10010  
(e-mail: catto@gursey.baruch.cuny.edu)*

and

*Physics Department, The Rockefeller University  
1230 York Avenue, New York, NY 10021-6399  
(e-mail: cattos@rockvax.rockefeller.edu)*

Jonathan Huntley<sup>2</sup>

*Department of Mathematics, Baruch College, CUNY  
17 Lexington Avenue, New York, NY 10010  
(e-mail: huntley@gursey.baruch.cuny.edu)*

Jay Jorgenson<sup>3</sup>

*School of Mathematics  
Institute for Advanced Study  
Princeton, NJ 08540*

David Tepper<sup>4</sup>

*Department of Mathematics, Baruch College, CUNY  
17 Lexington Avenue, New York, NY 10010  
(e-mail: tepper@gursey.baruch.cuny.edu)*

(1) Work supported in part by DOE contracts No. DE-AC-0276-ER 03074 and 03075; NSF Grant No. DMS-8917754; and PSC-CUNY Research Award Nos. 6-6(6407, 7418, 8445).

(2) Work supported by PSC-CUNY Research Award No. 9203393 and NSF Award No. DMS-9200317.

(3) Work supported from NSF grant DMS-93-07023 and from the Sloan Foundation.

(4) Work partially supported by PSC-CUNY Research Award No. 62335.

## Abstract

Let  $\mathcal{H}$  be the homogeneous space associated to the group  $PGL_3(\mathbf{R})$ . Let  $X = \Gamma \backslash \mathcal{H}$  where  $\Gamma = SL_3(\mathbf{Z})$  and consider the first non-trivial eigenvalue  $\lambda_1$  of the Laplacian on  $L^2(X)$ . Using geometric considerations, we prove the inequality  $\lambda_1 > 3\pi^2/10$ . Since the continuous spectrum is represented by the band  $[1, \infty)$ , our bound on  $\lambda_1$  can be viewed as an analogue of Selberg's eigenvalue conjecture for quotients of the hyperbolic half space.

## §1. Statement of the main theorem

A fundamental question in the spectral theory of automorphic forms is whether small eigenvalues exist. More specifically, let  $G$  be a noncompact reductive group with finite center,  $\Gamma$  a nonuniform lattice,  $K$  a maximal compact subgroup of  $G$ , and set  $X = \Gamma \backslash G/K$ . It is well known from the theory of Eisenstein series that  $L^2(X)$  has continuous spectrum for the ring of invariant differential operators, and in particular for the positive Laplacian,  $\Delta$ . The continuous spectrum will be, in cases of interest such as  $PGL_n(\mathbf{R})$ , an interval  $[a, \infty)$  with  $a > 0$ . The question we referred to above is: Do non-constant square integrable eigenforms exist with eigenvalue  $\lambda < a$ ? This problem is important for various considerations in number theory. In the case  $G = PGL_2(\mathbf{R})$  and  $\Gamma$  is a congruence subgroup, Selberg conjectured that no such nontrivial small eigenvalues exist.

In this paper, we consider the case when

$$G = PGL_3(\mathbf{R}) \quad \text{and} \quad \Gamma = SL_3(\mathbf{Z}).$$

Our main result is the following.

**Theorem** *Let  $\lambda_1$  be the eigenvalue for the first nontrivial eigenform on  $L^2(X)$ . Then*

$$\lambda_1 > 3\pi^2/10 > 2.96088.$$

## §2. Notation.

Let  $\mathcal{H} = \mathcal{G}/\mathcal{K}$  and set  $X = \Gamma \backslash \mathcal{H}$ . Explicit coordinates for  $\tau \in \mathcal{H}$  via the Iwasawa decomposition are given by

$$\tau = \begin{pmatrix} 1 & x_2 & x_3 \\ 0 & 1 & x_1 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 y_2 & 0 & 0 \\ 0 & y_1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

with  $y_1, y_2 > 0$ , from which one can compute that the (positive) Laplacian  $\Delta$  can be written as

$$-\Delta = y_1^2 \frac{\partial^2}{\partial y_1^2} - y_1 y_2 \frac{\partial^2}{\partial y_1 \partial y_2} + y_2^2 \frac{\partial^2}{\partial y_2^2} + y_1^2 (x_2^2 + y_2^2) \frac{\partial^2}{\partial x_3^2} + y_1^2 \frac{\partial^2}{\partial x_1^2} + y_2^2 \frac{\partial^2}{\partial x_2^2} + 2y_1^2 x_2 \frac{\partial^2}{\partial x_1 \partial x_3}$$

(see pages 17 and 33 of [1]). The ring of invariant differential operators is spanned by the identity, the Laplacian  $\Delta$ , and a third order operator  $\Delta_3$  (see [1]). The invariant volume element is given by

$$dV = \frac{dx_1 dx_2 dx_3 dy_1 dy_2}{y_1^3 y_2^3}.$$

We shall not use the explicit formula for the invariant volume element; however, the above expression for the Laplacian will be necessary in our proof of the main theorem.

For our purposes, it is more convenient to work with functions on  $\mathcal{H}$  that are  $SL_3(\mathbf{Z})$  invariant rather than considering functions on the quotient space  $X$ . To this end, we introduce a fundamental domain for  $\Gamma \backslash \mathcal{H}$ . Specifically, computations on page 56 of [2] show that a fundamental domain  $\mathcal{D}$  is described through the following set of inequalities:

$$\begin{aligned} v^{\frac{3}{2}} &< v^{\frac{3}{2}}(1 - x_2 + x_3)^2 + w(1 - x_1)^2 + w^{-1}; \\ v^{\frac{3}{2}} &< v^{\frac{3}{2}}(x_2 - x_3)^2 + w(1 - x_1)^2 + w^{-1}; \\ v^{\frac{3}{2}} &< v^{\frac{3}{2}}x_2^2 + w; \quad v^{\frac{3}{2}} < v^{\frac{3}{2}}x_3^2 + wx_1^2 + w^{-1}; \\ 1 &< w^{-2} + x_1^2; \quad 0 < x_1 < \frac{1}{2}; \quad 0 < x_2 < \frac{1}{2} \quad -\frac{1}{2} < x_3 < \frac{1}{2}, \end{aligned}$$

where we have used the notation  $w^{-1} = y_1$  and  $v^{-\frac{3}{2}} = y_2^2 y_1$ . Let  $S$  denote the Siegel set described via the inequalities

$$0 < x_1 < \frac{1}{2}, \quad 0 < x_2 < \frac{1}{2}, \quad -\frac{1}{2} < x_3 < \frac{1}{2}, \quad y_1 > \frac{\sqrt{3}}{2}, \quad y_2 > \frac{\sqrt{3}}{2}.$$

The set  $S$  contains the fundamental domain  $\mathcal{D}$ . Further, results from page 61 [2] show the existence of elements  $\gamma_1, \dots, \gamma_{10} \in SL_3(\mathbf{Z})$  such that  $S \subset \bigcup_{i=1}^{10} \mathcal{D}\gamma_i$  (we have used the notation  $\mathcal{D}\gamma$  to denote the image of the fundamental domain  $\mathcal{D}$  under left multiplication by  $\gamma$ ). The main aspects of the above points which we shall use are the assertions that for any  $\tau \in S$  we have  $y_1(\tau) > \sqrt{3}/2$  and that  $S$  is contained in ten translates of  $\mathcal{D}$ .

Recall that an automorphic form is a  $C^\infty$  function  $\phi$  on  $\mathcal{H}$  which satisfies the following properties:

- (i)  $\phi(\gamma \circ \tau) = \phi(\tau)$  for  $\gamma \in SL_3(\mathbf{Z})$ ;
- (ii)  $|\phi(\tau)| \ll y_1^{N_1} y_2^{N_2}$  for  $\tau \in \mathcal{D}$  and integers  $N_1, N_2$ ;

(iii)  $\phi$  is an eigenform for the ring of invariant differential operators.

An automorphic form is said to be a cusp form if it satisfies the additional property

$$(iv) \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \phi \left( \begin{bmatrix} 1 & 0 & \xi_3 \\ 0 & 1 & \xi_1 \\ 0 & 0 & 1 \end{bmatrix} \tau \right) d\xi_1 d\xi_3 = \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \phi \left( \begin{bmatrix} 1 & \xi_2 & \xi_3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \tau \right) d\xi_2 d\xi_3 = 0$$

Cusp forms are square integrable. Although we shall not need this fact, let us note that, from the theory of Eisenstein series, the only noncuspidal square integrable automorphic forms on  $X$  are constant.

### §3. Proof of the main theorem

Our method of proof is a modification of that used by Roelcke to show that the small eigenvalue  $\lambda_1$  for the quotient space  $SL_2(\mathbf{Z}) \backslash SL_2(\mathbf{R}) / SO_2(\mathbf{R})$  satisfies the bound  $\lambda_1 > 3\pi^2/2$  (see page 511 of [3]). We shall use the Fourier expansion of automorphic forms associated to  $SL_3(\mathbf{Z})$ , as developed in Chapter IV of [1].

Assume that  $\phi$  is a non-constant automorphic form, so then  $\Delta\phi = \lambda\phi$  and  $\phi\Delta\phi = \lambda\phi^2$ . Through integration by parts, using the automorphic boundary conditions, and the fact that the Siegel domain  $S$  is contained in ten translates of the fundamental domain  $\mathcal{D}$ , we obtain the inequality

$$\frac{\int_S |\nabla\phi|^2 dV}{\int_S |\phi|^2 dV} < 10\lambda.$$

As on page 67 of [1], let us expand  $\phi$  in a Fourier expansion with respect to the abelian group

$$\left\{ \begin{bmatrix} 1 & 0 & \xi_3 \\ 0 & 1 & \xi_1 \\ 0 & 0 & 1 \end{bmatrix} \mid \xi_1, \xi_3 \in \mathbf{R} \right\}.$$

Specifically, we have  $\phi(\tau) = \sum_{n_1, n_3} \phi_{n_1}^{n_3}(\tau)$  where

$$\phi_{n_1}^{n_3}(\tau) = \int_0^1 \int_0^1 \phi \left( \begin{bmatrix} 1 & 0 & \xi_3 \\ 0 & 1 & \xi_1 \\ 0 & 0 & 1 \end{bmatrix} \tau \right) e^{-2\pi i(n_1 \xi_1 + n_3 \xi_3)} d\xi_1 d\xi_3.$$

Observe that  $\phi_0^0 = 0$  since  $\phi$  is not constant and square integrable, hence cuspidal. Let

$$\Gamma_1^2 = \left\{ \begin{bmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & 1 \end{bmatrix} : \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2(\mathbf{Z}) \right\},$$

and set  $\Gamma_\infty^2$  to be the subgroup of  $\Gamma_1^2$  which stabilizes infinity. As on page 69 of [1], we then can write  $\phi_{n_1}^{n_3}$  as

$$\phi_{n_1}^{n_3}(\tau) = \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \sum_{n_1=1}^{\infty} \phi_{n_1}^0(\gamma \circ \tau).$$

By a standard application of elliptic regularity,  $\phi$  is necessarily  $C^\infty$ , hence we can interchange integration and summation and apply Parseval's theorem to obtain the inequality

$$\frac{\int_S \sum_{n_1=1}^{\infty} \left| \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \nabla_\tau \phi_{n_1}^0(\gamma \circ \tau) \right|^2 dV}{\int_S \sum_{n_1=1}^{\infty} \left| \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \phi_{n_1}^0(\gamma \circ \tau) \right|^2 dV} < 10\lambda.$$

Since  $\nabla$  is an invariant operator, we may differentiate the expressions in the numerator and then evaluate at  $\gamma \circ \tau$ , thus yielding

$$\frac{\int_S \sum_{n_1=1}^{\infty} \left| \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \nabla_\tau \phi_{n_1}^0(\tau)|_{\gamma \circ \tau} \right|^2 dV}{\int_S \sum_{n=1}^{\infty} \left| \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \phi_{n_1}^0(\tau)|_{\gamma \circ \tau} \right|^2 dV} < 10\lambda.$$

We now integrate by parts and consider the action of the Laplacian  $\Delta$  on functions of the form  $\phi_{n_1}^0$ . Since each function  $\phi_{n_1}^0$  is independent of  $x_3$ , these terms in  $\Delta$  annihilate  $\phi_{n_1}^0$ . Observe that all terms involving  $y_1, y_2$  and  $x_2$  are positive operators (compare with line (2.31) on page 32 of [1]), so we obtain the bound

$$\Delta \phi_{n_1}^0 \geq -y_1^2 \cdot \frac{\partial^2}{\partial x_1^2} \phi_{n_1}^0 = y_1^2 \cdot 4\pi^2 n_1^2 \phi_{n_1}^0.$$

Since  $y_1^2 > \frac{3}{4}$ , we have  $\Delta \phi_{n_1}^0 \geq \frac{3}{4} \cdot 4\pi^2 n_1^2 \phi_{n_1}^0 = 3\pi^2 n_1^2 \phi_{n_1}^0$ . Combining this inequality with the above calculations and the cuspidality condition  $\phi_0^0 = 0$ , we obtain

$$10\lambda > \frac{\int_S \sum_{n_1=1}^{\infty} \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \Delta_\tau \phi_{n_1}^0(\tau)|_{\gamma \circ \tau} \cdot \phi_{n_1}^0(\tau)|_{\gamma \circ \tau} dV}{\int_S \sum_{n=1}^{\infty} \left| \sum_{\gamma \in \Gamma_\infty^2 \setminus \Gamma_1^2} \phi_{n_1}^0(\tau)|_{\gamma \circ \tau} \right|^2 dV} \geq 3\pi^2,$$

hence  $\lambda \geq 3\pi^2/10$ . Since  $\phi$  was any cusp form, we obtain the bound as asserted in the theorem.

#### §4. Concluding remarks

As the continuous spectrum in this situation is  $[1, \infty)$ , our theorem implies an analogue of Selberg's eigenvalue conjecture. Note that our bound is stronger than result for  $SL_3(\mathbf{Z})$  from [4] who proved  $\lambda_1 \geq 1$ . In general, our method applies to  $G = SL_n(\mathbf{R})$  with  $\Gamma = SL_n(\mathbf{Z})$  to give the bound  $\lambda_1 > 3\pi^2/M$  where  $M$  is the number of fundamental domains which intersect a Siegel set containing the fundamental domain constructed in [2]; however, for  $n \geq 4$ , this bound is rather weak. Finally, let us remark that our theorem is indeed a consequence of the Ramanujan conjecture, which asserts that all nontrivial automorphic representations come from tempered representations.

## References

- [1] BUMP, D.: *Automorphic forms on  $GL(3, \mathbf{R})$* . Lecture Notes in Mathematics **1083** New York: Springer-Verlag (1984).
- [2] GREINER, D.: On the shape of fundamental domains in  $GL(n, R)$ . *Pacific Journal Math.* **160** (1993) 53-65.
- [3] HEJHAL, D. A.: *The Selberg Trace Formula for  $PSL(2, \mathbf{R})$* , vol. 2. Lecture Notes in Mathematics **1001** New York: Springer-Verlag (1983).
- [4] MILLER, S.: Spectral and cohomological applications of the Rankin-Selberg method. *Inter. Math. Res. Notices* **1** (1996) 15-26.