

ON THE DUFLO FORMULA FOR L_∞ -ALGEBRAS AND Q -MANIFOLDS

BORIS SHOIKHET

ABSTRACT. We prove a direct analogue of the classical Duflo formula in the case of L_∞ -algebras. We conjecture an analogous formula in the case of an arbitrary Q -manifold. When G is a compact connected Lie group, the Duflo theorem for the Q -manifold $(\Pi T G, d_{\text{DR}})$ is exactly the Duflo theorem for the Lie algebra $\mathfrak{g} = \text{Lie } G$. The corresponding theorem for the Q -manifold $(\Pi T M, d_{\text{DR}})$, where M is an arbitrary smooth manifold, is a generalization of the Duflo theorem for the case of smooth manifolds. On the other hand, the Duflo theorem for the Q -manifold $(\Pi \overline{T}_{\text{hol}} M, \overline{\partial})$, where M is a complex manifold, is a generalization of the M. Kontsevich's "theorem on complex manifold" [K1], Sect. 8.4.

CONTENTS

1. The classical Duflo formula
 2. Strong homotopy Lie algebras and Q -manifolds
 3. Relationship with the Formality conjecture
 4. Duflo formula for L_∞ -algebras
 5. Duflo formula for Q -manifolds
- References

1. THE CLASSICAL DUFLO FORMULA [D], [K1]

Let \mathfrak{g} be a finite-dimensional Lie algebra, $S^\bullet(\mathfrak{g})$ be the symmetric algebra of the vector space \mathfrak{g} , and $U(\mathfrak{g})$ be the universal enveloping algebra of the Lie algebra \mathfrak{g} . Both spaces $S^\bullet(\mathfrak{g})$ and $U(\mathfrak{g})$ are \mathfrak{g} -modules with respect to the adjoint action; it follows from the Poincaré–Birkhoff–Witt theorem that these modules are isomorphic. Therefore, the vector spaces of invariants $[S^\bullet(\mathfrak{g})]^\mathfrak{g}$ and $[U(\mathfrak{g})]^\mathfrak{g}$ are isomorphic. The Duflo theorem states that $[S^\bullet(\mathfrak{g})]^\mathfrak{g}$ and $[U(\mathfrak{g})]^\mathfrak{g}$ are canonically isomorphic as *algebras*. Moreover ([K1], Sect. 8.3) the algebras $H_{\text{Lie}}^\bullet(\mathfrak{g}; S^\bullet(\mathfrak{g}))$ and $H_{\text{Lie}}^\bullet(\mathfrak{g}; U(\mathfrak{g}))$ are canonically isomorphic.

Let us recall the construction of this isomorphism. First of all, the map $\varphi_{PBW}: S^\bullet(\mathfrak{g}) \rightarrow U(\mathfrak{g})$, defined as follows

$$(1) \quad \varphi_{PBW}(g_1 \cdot g_2 \cdot \dots \cdot g_k) = \frac{1}{k!} \sum_{\sigma \in \Sigma_k} g_{\sigma(1)} * g_{\sigma(2)} * \dots * g_{\sigma(k)}$$

is an isomorphism of the \mathfrak{g} -modules. The corresponding map $\varphi_{PBW}: [S^\bullet(\mathfrak{g})]^\mathfrak{g} \xrightarrow{\sim} [U(\mathfrak{g})]^\mathfrak{g}$ is *not* an isomorphism of the algebras.

There exists an isomorphism of \mathfrak{g} -modules $\varphi_{\text{strange}}: S^\bullet(\mathfrak{g}) \xrightarrow{\sim} S^\bullet(\mathfrak{g})$ such that the composition

$$[S^\bullet(\mathfrak{g})]^\mathfrak{g} \xrightarrow{\varphi_{\text{strange}}} [S^\bullet(\mathfrak{g})]^\mathfrak{g} \xrightarrow{\varphi_{PBW}} [U(\mathfrak{g})]^\mathfrak{g}$$

Date: 29.11.1998.

is an isomorphism of the algebras.

The map $\varphi_{\text{strange}}: S^\bullet(\mathfrak{g}) \xrightarrow{\sim} S^\bullet(\mathfrak{g})$ is defined as follows. Let us consider elements of \mathfrak{g}^* as derivations of the symmetric algebra $S^\bullet(\mathfrak{g})$, then elements of $S^k(\mathfrak{g}^*)$ are differential operators with constant coefficients acting on $S^\bullet(\mathfrak{g})$. If the Lie algebra \mathfrak{g} is finite-dimensional, there exists a canonical element $c_k \in S^k(\mathfrak{g}^*)$ for every $k \geq 1$, namely, it is the trace of the k -th power of the adjoint action, $c_k = \{g \mapsto \text{Tr ad}^k g\}$. The elements c_k are invariant elements in $S^k(\mathfrak{g}^*)$. We set:

$$(2) \quad \varphi_{\text{strange}} = \exp \left(\sum_{k \geq 1} \alpha_{2k} \cdot c_{2k} \right)$$

where the rational numbers α_{2k} are defined by the formula:

$$(3) \quad \sum_{k \geq 1} \alpha_{2k} \cdot x^{2k} = \log \sqrt{\frac{e^{\frac{x}{2}} - e^{-\frac{x}{2}}}{x}}.$$

It is clear, that $\varphi_{\text{strange}}: S^\bullet(\mathfrak{g}) \rightarrow S^\bullet(\mathfrak{g})$ is an isomorphism of the \mathfrak{g} -modules.

Theorem ([D]). *For any finite-dimensional Lie algebra \mathfrak{g} the composition*

$$\varphi_{PBW} \circ \varphi_{\text{strange}}: [S^\bullet(\mathfrak{g})]^\mathfrak{g} \rightarrow [U(\mathfrak{g})]^\mathfrak{g}$$

is an isomorphism of the algebras.

□

Theorem ([K1], Sect. 8.3). *For any finite-dimensional Lie super-algebra \mathfrak{g} the composition*

$$\varphi_{PBW} \circ \varphi_{\text{strange}}: H_{\text{Lie}}^\bullet(\mathfrak{g}; S^\bullet(\mathfrak{g})) \rightarrow H^\bullet(\mathfrak{g}; U(\mathfrak{g})).$$

is an isomorphism of the algebras.

□

In the present paper we prove the last Theorem also for differential graded Lie algebras. Moreover, after minor modifications the analogous statement is true also for strong homotopy Lie algebras (L_∞ -algebras). In fact, this result is a direct consequence of [KSh].

2. STRONG HOMOTOPY LIE ALGEBRAS AND Q -MANIFOLDS

2.1. A strong homotopy Lie algebra \mathfrak{g} (L_∞ -algebra) is, by definition, a \mathbb{Z} -graded vector space \mathfrak{g} and an odd vector field Q of degree +1 on the space $\mathfrak{g}[1]$ such that $[Q, Q] = 0$.

On the other hand, it is an odd derivation Q of degree +1 on the algebra $\bigwedge^\bullet(\mathfrak{g}^*)$, such that $Q^2 = 0$ (here \bigwedge^\bullet stands for the super-exterior algebra).

In the simplest case when \mathfrak{g} is a Lie algebra such a differential Q on $\bigwedge^\bullet(\mathfrak{g}^*)$ is the cochain differential, it contains only a quadratic part. When \mathfrak{g} is a DG Lie algebra, the cochain differential on $\bigwedge^\bullet(\mathfrak{g}^*)$ contain a linear and a quadratic parts, In other words, differential graded Lie algebras give us examples of L_∞ -algebras.

In the general case, the odd vector field Q contains also parts of 3-rd, 4-th, ... degree.

A Q -manifold is a smooth \mathbb{Z} -graded manifold X and an odd vector field Q on X of degree +1 such that $[Q, Q] = 0$. In other words, it is an odd derivation Q of the algebra of smooth functions $C^\infty(X)$ such that $Q^2 = 0$. It is clear, that the case when X is an \mathbb{Z} -graded vector space is exactly the case of L_∞ -algebras.

Examples. (1) Let Y be a smooth (purely even) manifold, TY be its tangent bundle, and $X = T[1]Y$ be the tangent bundle with fibers $T_x[1]$. The algebra of functions on $T[1]Y$ is the algebra of differential forms on Y , the de Rham differential d_{DR} acts on this algebra. Therefore $(T[1]Y, d_{\text{DR}})$ is an example of Q -manifold.

(2) Let Y be a complex manifold, $X = \overline{T}_{\text{hol}}[1]Y$, where $T_{\text{hol}}Y$ is the holomorphic tangent bundle. Then $(\overline{T}_{\text{hol}}[1]Y, \overline{\partial})$ is an example of Q -manifold. The corresponding complex is the Dolbeault complex of the manifold Y .

2.2.

Definition (algebra of polyvector fields on a Q -manifold). Let (A, Q) be a differential graded commutative algebra, $\text{Der } A$ be the Lie algebra of derivations of the algebra A . Then $\text{Der } A$ is a complex, the differential $D: \text{Der } A \rightarrow (\text{Der } A)[1]$ is defined as the bracket with the derivation Q :

$$D(\xi) = [Q, \xi], \quad \xi \in \text{Der } A.$$

It is clear that $D(f \cdot \xi) = Q(f) \cdot \xi + f \cdot D\xi$, $f \in A$, $\xi \in \text{Der } A$. The differential D acts by the Leibniz rule on the (super) exterior algebra over the algebra A , $\bigwedge_A^\bullet(\text{Der } A)$, we denote the last DG algebra by $T_{\text{poly}}^\bullet(A, Q)$, the algebra of polyvector fields. When X is a Q -manifold, the algebra $T_{\text{poly}}^\bullet(X, Q)$ is, by definition, the DG algebra $T_{\text{poly}}^\bullet(C^\infty(X), Q)$.

Basic example. Let \mathfrak{g} be a Lie algebra, $(\mathfrak{g}[1], d_{\text{Lie}})$ be the corresponding Q -manifold. The DG algebra of functions on this Q -manifold is $C_{\text{Lie}}^\bullet(\mathfrak{g}; \mathbb{C})$, the cochain complex of the Lie algebra \mathfrak{g} . The DG algebra of polyvector fields $T_{\text{poly}}^\bullet(\mathfrak{g}[1], d_{\text{Lie}})$ is the cochain complex $C_{\text{Lie}}^\bullet(\mathfrak{g}, S^\bullet(\mathfrak{g}))$ with the coefficient in the symmetric algebra of \mathfrak{g} .

2.3.

Conjecture. Let (A_1, Q_1) and (A_2, Q_2) be two commutative smooth DG algebras, which are quasi-isomorphic. Then

$$H^\bullet(T_{\text{poly}}^\bullet(A_1, Q_1)) \simeq H^\bullet(T_{\text{poly}}^\bullet(A_2, Q_2))$$

as algebras.

(A DG algebra (A, Q) is called smooth if it is the algebra of functions on a smooth Q -manifold.)

3. RELATIONSHIP WITH THE FORMALITY CONJECTURE [K1], [KSH]

3.1. Let $(\mathfrak{g}[1], Q)$ be an L_∞ -algebra, $(\bigwedge^\bullet \mathfrak{g}^*, Q)$ be the corresponding DG algebra of functions. In the case when \mathfrak{g} is a Lie algebra, $Q = d_{\text{Lie}}$, it is well-known result that

$$HH^\bullet(C_{\text{Lie}}^\bullet(\mathfrak{g}; \mathbb{C})) \simeq HH^\bullet(U(\mathfrak{g}))$$

as algebras, where HH^\bullet here stands for the Hochschild cohomology. On the other hand,

$$T_{\text{poly}}^\bullet(\mathfrak{g}[1], d_{\text{Lie}}) = C_{\text{Lie}}^\bullet(\mathfrak{g}, S^\bullet(\mathfrak{g})).$$

Therefore, the following theorem can be considered as a generalization of the Duflo formula for the case of L_∞ -algebras.

Theorem. The algebras $H^\bullet(T_{\text{poly}}^\bullet(\mathfrak{g}[1], Q))$ and $HH^\bullet(\bigwedge^\bullet \mathfrak{g}^*, Q)$ are canonically isomorphic.

We prove this theorem and construct an explicit isomorphism, analogous to the isomorphism $\varphi_{PBW} \circ \varphi_{\text{strange}}$ from Sect. 1, in the next Section. Here we explain, following [KSh], Sect. 4, how the Duflo theorem itself follows from the Formality conjecture of Maxim Kontsevich and his theorem on the cup-products on tangent cohomology.

Theorem (Formality conjecture; proved in [K1]). *Let $T_{\text{poly}}^{\bullet}(\mathbb{R}^{m|n})$ be the DG Lie algebra of polyvector field on $\mathbb{R}^{m|n}$ with zero-differential and the Schouten–Nijenhuis bracket and let $\mathcal{D}_{\text{poly}}^{\bullet}(\mathbb{R}^{m|n})$ be the DG Lie algebra of polydifferential operators on $\mathbb{R}^{m|n}$ with the Hochschild differential and the Gerstenhaber bracket. Then there exists an L_{∞} -quasi-isomorphism $\mathcal{U}: T_{\text{poly}}^{\bullet} \rightarrow \mathcal{D}_{\text{poly}}^{\bullet}$.* □

(Here $\mathbb{R}^{m|n}$ is a super-space $\mathbb{R}^m \oplus \mathbb{R}^n[1]$.)

More generally, one can consider an arbitrary finite-dimensional \mathbb{Z} -graded vector space.

3.2. An L_{∞} -morphism of two L_{∞} -algebras is a Q -equivariant (may be nonlinear) map $\varphi: (\mathfrak{g}_1[1], 0) \rightarrow (\mathfrak{g}_2[1], 0)$. In particular, if $\gamma \in (\mathfrak{g}_1)^1$ is such that $Q_1|_{\gamma} = 0$ than $Q_2|_{\varphi(\gamma)} = 0$. In the case of DG Lie algebra \mathfrak{g} the equation $Q|_{\gamma} = 0$ is exactly the Maurer–Cartan equation:

$$(4) \quad \gamma \in \mathfrak{g}^1, \quad d\gamma + \frac{1}{2}[\gamma, \gamma] = 0.$$

Also, $d + \text{ad } \gamma$ defines a new differential on \mathfrak{g} . Moreover, if $\varphi: (\mathfrak{g}_1[1], 0) \rightarrow (\mathfrak{g}_2[1], 0)$ is an L_{∞} -morphism of DG Lie algebras, it defines a map of the *complexes*

$$(\mathfrak{g}_1, d_1 + \text{ad } \gamma) \rightarrow (\mathfrak{g}_2, d_2 + \text{ad } \varphi(\gamma))$$

for each solution $\gamma \in \mathfrak{g}_1^1$ of the Maurer–Cartan equation. We denote the *tangent complex* $(\mathfrak{g}, d + \text{ad } \gamma)$ by $T_{\gamma}(\mathfrak{g})$ and we denote by $\varphi_{\gamma}: T_{\gamma}(\mathfrak{g}_1) \rightarrow T_{\varphi(\gamma)}(\mathfrak{g}_2)$ the corresponding tangent map. The map φ_{γ} is a map of the complexes.

For any solution $\gamma \in T_{\text{poly}}^1(\mathbb{R}^{m|n})$ of the Maurer–Cartan equation there exists a product on the tangent complex $T_{\gamma}(T_{\text{poly}}^{\bullet}(\mathbb{R}^{m|n}))$ which coincides with the usual cup-product of the polyvector fields. On the other hand, for any solution $\gamma \in \mathcal{D}_{\text{poly}}^1(\mathbb{R}^{m|n})$ of the Maurer–Cartan equation the usual product of Hochschild cochains:

$$(5) \quad (\varphi \circ \psi)(a_1 \otimes \dots \otimes a_{k_1+k_2}) = \varphi(a_1 \otimes \dots \otimes a_{k_1}) \cdot \psi(a_{k_1+1}, \dots, a_{k_1+k_2})$$

(here $\varphi: A^{\otimes k_1} \rightarrow A$, $\psi: A^{\otimes k_2} \rightarrow A$ are Hochschild cochains on an algebra A) defines a product on the tangent complex $T_{\gamma}(\mathcal{D}_{\text{poly}}^{\bullet}(\mathbb{R}^{m|n}))$.

Theorem (Theorem on cup-products on tangent cohomology, [K1], Sect. 8). *Let $\mathcal{U}: T_{\text{poly}}^{\bullet}(\mathbb{R}^{m|n}) \rightarrow \mathcal{D}_{\text{poly}}^{\bullet}(\mathbb{R}^{m|n})$ be the Formality L_{∞} -morphism, and let $\gamma \in T_{\text{poly}}^1(\mathbb{R}^{m|n})$ be a solution of the Maurer–Cartan equation. Then the map*

$$[\mathcal{U}_{\gamma}]: H^{\bullet}(T_{\gamma}(T_{\text{poly}}^{\bullet})) \rightarrow H^{\bullet}(T_{\mathcal{U}(\gamma)}(\mathcal{D}_{\text{poly}}^{\bullet})),$$

defined by the tangent map of the complexes:

$$\mathcal{U}_{\gamma}: T_{\gamma}(T_{\text{poly}}^{\bullet}) \rightarrow T_{\mathcal{U}(\gamma)}(\mathcal{D}_{\text{poly}}^{\bullet}),$$

is a morphism of the algebras. □

3.3. The differential graded Lie algebra $T_{\text{poly}}^\bullet(\mathbb{R}^m)$ is graded as follows:

$$T_{\text{poly}}^i(\mathbb{R}^m) = \{ (i+1)\text{-polyvector fields} \}.$$

In particular,

$$T_{\text{poly}}^0(\mathbb{R}^m) = \{ \text{vector fields on } \mathbb{R}^m \}.$$

However, every *odd* vector field Q of degree $+1$ on $\mathbb{R}^{m|n}$ lies in $T_{\text{poly}}^1(\mathbb{R}^{m|n})$, and the Maurer–Cartan equation is exactly the equation $[Q, Q] = 0$.

Let \mathfrak{g} be purely even finite-dimensional Lie algebra, and let $\gamma = \sum_{i,j,k} c_{ij}^k \xi_i \xi_j \frac{\partial}{\partial \xi_k}$ be the corresponding odd vector field on $\mathfrak{g}[1]$; the identity $[\gamma, \gamma] = 0$ is exactly the Jacobi identity.

Let us summarize some simplest facts on the tangent complex in this case.

- Lemma.** (i) $T_\gamma(T_{\text{poly}}^\bullet(\mathfrak{g}[1])) = C_{\text{Lie}}^\bullet(\mathfrak{g}; S^\bullet(\mathfrak{g}))$;
(ii) $\mathcal{U}(\gamma) = \left\{ f \mapsto \sum_{i,j,k} c_{ij}^k \xi_i \xi_j \frac{\partial f}{\partial \xi_k} \right\} \in \mathcal{D}_{\text{poly}}^1(\mathfrak{g}[1])$;
(iii) the tangent complex $T_{\mathcal{U}(\gamma)}(\mathcal{D}_{\text{poly}}^\bullet(\mathfrak{g}[1])) = CH^\bullet(C_{\text{Lie}}^\bullet(\mathfrak{g}; \mathbb{C}))$, the Hochschild cohomological complex of the cochain complex of the Lie algebra \mathfrak{g} . □

It follows from Theorem 3.2 that the Formality L_∞ -morphism produces a map of the algebras

$$[\mathcal{U}_\gamma]: H_{\text{Lie}}^\bullet(\mathfrak{g}; S^\bullet(\mathfrak{g})) \rightarrow HH^\bullet(C_{\text{Lie}}^\bullet(\mathfrak{g}; \mathbb{C})).$$

3.3.1.

Lemma. The map $[\mathcal{U}_\gamma]$ is an isomorphism (of the vector spaces).

Proof. The statement of the Lemma follows from the homotopy theory of L_∞ -algebras, see [K1], Sect. 4.5.1. If $\mathfrak{g}_1, \mathfrak{g}_2$ are two DG algebras, and

$$\varphi: (\mathfrak{g}_1[1], 0) \rightarrow (\mathfrak{g}_2[1], 0)$$

is an L_∞ -quasi-isomorphism between them, and a solution $\gamma \in \mathfrak{g}_1^1$ of the Maurer–Cartan equation is *sufficiently small*, i.e. lies in an open neighbourhood of 0 in $\mathfrak{g}_1[1]$, than the tangent map

$$[\varphi]: H^\bullet T_\gamma(\mathfrak{g}_1) \rightarrow H^\bullet T_{\varphi(\gamma)}(\mathfrak{g}_2)$$

is an *isomorphism* of the vector spaces. In our case, we can consider the vector field $\gamma_t = \sum_{i,j,k} t \cdot c_{ij}^k \xi_i \xi_j \frac{\partial f}{\partial \xi_k}$, $t \in \mathbb{C}$, instead of the vector field γ . For sufficiently small t the vector field γ_t lies in any open neighbourhood of 0 in $T_{\text{poly}}^\bullet(\mathfrak{g}[1])$; on the other hand, if $[\mathcal{U}_{\gamma_t}]$ is an isomorphism for some $t \neq 0$, than $[\mathcal{U}_\gamma]$ is also an isomorphism. □

4. DUFLO FORMULA FOR L_∞ -ALGEBRAS

We want to describe explicitly the tangent map

$$\mathcal{U}_\gamma: C_{\text{Lie}}^\bullet(\mathfrak{g}; S^\bullet(\mathfrak{g})) \rightarrow CH^\bullet(C_{\text{Lie}}^\bullet(\mathfrak{g}; \mathbb{C})).$$

Let $\gamma \in T_{\text{poly}}^1(\mathbb{R}^{m|n})$ be an arbitrary solution of the Maurer–Cartan equation, which is a *vector field*. The following description of the tangent map \mathcal{U}_γ was found in [KSh].

We fix a basis x_1, \dots, x_{m+n} on $\mathbb{R}^{m|n}$.

(i) The map $\varphi_{\text{HKR}}: T_{\text{poly}}^\bullet(\mathbb{R}^{m|n}) \rightarrow \mathcal{D}_{\text{poly}}^\bullet(\mathbb{R}^{m|n})$

This map is defined as follows:

if $\eta = \xi \wedge \dots \wedge \xi_k$, $\eta \in T_{\text{poly}}^\bullet(\mathbb{R}^{m|n})$, ξ_1, \dots, ξ_k are vector fields, then

$$(6) \quad \varphi_{\text{HKR}}(\eta)(f_1 \otimes \dots \otimes f_k) = \frac{1}{k!} \sum_{\sigma \in \Sigma_k} \text{supersign}(\sigma) \cdot \xi_{\sigma(1)}(f_1) \cdot \dots \cdot \xi_{\sigma(k)}(f_k).$$

The *supersign*(σ) is defined by the formula

$$\xi_{\sigma(1)} \wedge \dots \wedge \xi_{\sigma(k)} = \text{supersign}(\sigma) \cdot \xi_1 \wedge \dots \wedge \xi_k.$$

In particular, if ξ_1, \dots, ξ_k are usual (even) vector fields on \mathbb{R}^m , then *supersign*(σ) = *sgn*(σ); if all the vector fields ξ_1, \dots, ξ_k are odd, then *supersign*(σ) = 1 for any $\sigma \in \Sigma_k$.

(ii) the map c_T :

$$\text{Vect}(\mathbb{R}^{m|n}) \rightarrow \text{Hom}_{\text{Fun}(\mathbb{R}^{m|n})}(\text{Vect}(\mathbb{R}^{m|n}) \rightarrow \text{Vect}(\mathbb{R}^{m|n}))$$

(the ‘‘Atiyah class’’)

Let $\eta \in \text{Vect}(\mathbb{R}^{m|n})$. We set

$$(7) \quad c_T(\eta) = \left\{ \partial_{I(2)} \mapsto \sum_{I(1), I(3)} \eta(dx^{I(1)}) \cdot \partial_{I(1)} \partial_{I(2)} \langle \gamma, dx^{I(3)} \rangle \partial_{I(3)} \right\}.$$

Here $\partial_i = \frac{\partial}{\partial x_i}$ and $\langle dx^i, \partial_j \rangle = \delta_{ij}$, and I runs through all possible maps

$$I: \{1, 2, 3\} \mapsto \{1, 2, \dots, m+n\}.$$

The k -th power of the map c_T is the map

$$c_T^k: \text{Vect}^{\otimes k}(\mathbb{R}^{m|n}) \rightarrow \text{Hom}_{\text{Fun}(\mathbb{R}^{m|n})}^{\otimes k}(\text{Vect}, \text{Vect})$$

There exists the trace map

$$\text{Tr}: \text{Hom}_{\text{Fun}}^{\otimes k}(\text{Vect}, \text{Vect}) \rightarrow \text{Fun},$$

and the composition $\text{Tr} \circ c_T^k$ is a map

$$(8) \quad \text{Tr} \circ c_T^k = c_k: \text{Vect}^{\otimes k}(\mathbb{R}^{m|n}) \rightarrow \text{Fun}(\mathbb{R}^{m|n}).$$

After the (super-) symmetrization we consider c_k as an operator

$$c_k: T_{\text{poly}}^\bullet(\mathbb{R}^{m|n}) \rightarrow T_{\text{poly}}^\bullet(\mathbb{R}^{m|n}).$$

Example. In the case of the odd field $\gamma = \sum_{i,j,k} c_{ij}^k \xi_i \xi_j \frac{\partial}{\partial_k}$ on the space $\mathfrak{g}[1]$, where \mathfrak{g} is a finite-dimensional Lie algebra, we have

$$\langle \gamma, dx^{I(3)} \rangle = \sum_{i,j} c_{i,j}^{I(3)} \xi_I \xi_j.$$

Let $\eta = \partial_l$ for some $l = 1, \dots, \dim \mathfrak{g}$. Then, by formula (7), the map

$$c_T(\partial_l) = \left\{ \partial_{I(2)} \mapsto \sum_{I(3)} c_{l,I(2)}^{I(3)} \partial_{I(3)} \right\} = -\text{ad}(\partial_l)$$

coincides, up to a sign, with the adjoint action.

Theorem ([KSh]). *Let $\gamma \in T_{\text{poly}}^1(\mathbb{R}^{m|n})$ be an odd vector field such that $[\gamma, \gamma] = 0$. Then the tangent map*

$$\mathcal{U}_\gamma: T_\gamma(T_{\text{poly}}^\bullet(\mathbb{R}^{m|n})) \rightarrow T_{\mathcal{U}(\gamma)}(\mathcal{D}_{\text{poly}}^\bullet(\mathbb{R}^{m|n}))$$

coincides with the composition

$$\mathcal{U}_\gamma = \varphi_{\text{HKR}} \circ \varphi_{\text{strange}},$$

where

$$\varphi_{\text{strange}} = \exp \left(\sum_{k \geq 1} \alpha_{2k} c_{2k} \right).$$

(the rational numbers α_{2k} were defined in Sect. 1)

□

Theorem. *Let $(\mathfrak{g}[1], Q)$ be a finite-dimensional L_∞ -algebra, and let $Q = Q^{(1)} + Q^{(2)} + Q^{(3)} + \dots$, where $Q^{(i)}$ is the part of the vector field Q of i -th degree. Then the map*

$$\varphi_{\text{HKR}} \circ \varphi_{\text{strange}}: H^\bullet(T_{\text{poly}}^\bullet(\mathfrak{g}[1], Q)) \rightarrow HH^\bullet(\bigwedge^\bullet g^*, Q)$$

is an isomorphism of the algebras, where the operators

$$c_{2k}: T_{\text{poly}}^\bullet(\mathfrak{g}[1], Q) \rightarrow T_{\text{poly}}^\bullet(\mathfrak{g}[1], Q)$$

are defined by formulas (7) and (8) when $\gamma = Q$.

□

Corollary. *Let \mathfrak{g} be a finite-dimensional differential graded Lie algebra, i.e. $Q = Q^{(1)} + Q^{(2)}$. Then c_T, c_{2k} do not depend on $Q^{(1)}$, i.e. on the differential in the DG Lie algebra \mathfrak{g} . Therefore, the usual Duflo formula (Section 1) defines an isomorphism of the algebras*

$$\varphi_{\text{PBW}} \circ \varphi_{\text{strange}}: H_{\text{Lie}}^\bullet(\mathfrak{g}; S^\bullet(\mathfrak{g})) \rightarrow HH^\bullet(C_{\text{Lie}}^\bullet(\mathfrak{g}; \mathbb{C})).$$

Proof. The expression for the ‘‘Atiyah class’’ c_T depends only on the second derivatives of the vector field Q (see formula (7)), but $Q^{(1)}$ is the linear term.

□

5. DUFLO FORMULA FOR Q -MANIFOLDS

Here we propose a conjecture giving an explicit formula for the isomorphism of the algebras:

$$H^\bullet(T_{\text{poly}}^\bullet(C^\infty(X), Q)) \rightarrow HH^\bullet(C^\infty(X), Q)$$

for any smooth Q -manifold X .

5.1. Let us explain why the Duflo isomorphism for a smooth Q -manifold X should exist.

Theorem (Formality conjecture for smooth manifold; proved in [K1], Sect. 7). *Let X be a smooth super-manifold, $T_{\text{poly}}^\bullet(X)$ be the DG Lie algebra of smooth polyvector fields on X , and let $\mathcal{D}_{\text{poly}}^\bullet(X)$ be the DG Lie algebra of smooth polydifferential operators on X . Then there exists an L_∞ -quasi-isomorphism $\mathcal{U}: T_{\text{poly}}^\bullet(X) \rightarrow \mathcal{D}_{\text{poly}}^\bullet(X)$.* □

More generally, one can consider any smooth \mathbb{Z} -graded manifold X .

There does not exist a canonical choice of this L_∞ -quasi-isomorphism. It was constructed in [K1], Sect. 7 canonically “up to a homotopy”.

Conjecture ([K2], Sect. 6.4). *Let X be a smooth super-manifold, \mathcal{U} be an L_∞ -quasi-isomorphism $\mathcal{U}: T_{\text{poly}}^\bullet(X) \rightarrow \mathcal{D}_{\text{poly}}^\bullet(X)$ constructed in [K1], Sect. 7, and let $\gamma \in T_{\text{poly}}^1(X)$ be such that $[\gamma, \gamma] = 0$ (the Maurer–Cartan equation). Then the map*

$$[\mathcal{U}_\gamma]: H^\bullet T_\gamma(T_{\text{poly}}^\bullet(X)) \rightarrow H^\bullet T_{\mathcal{U}(\gamma)}(\mathcal{D}_{\text{poly}}^\bullet(X))$$

induced by the tangent map \mathcal{U}_γ , is a morphism of the algebras.

One can also suppose, that $\mathcal{U}_1 = \varphi_{\text{HKR}}$ and that $\mathcal{U}(\gamma) = \varphi_{\text{HKR}}(\gamma)$.

Let X be a smooth super-manifold, Q be an odd vector field of degree +1 on X such that $[Q, Q] = 0$ (i.e., X is a Q -manifold). It follows from the above Conjecture that the tangent map

$$[\mathcal{U}_Q]: H^\bullet T_{\text{poly}}^\bullet(C^\infty(X), Q) \rightarrow HH^\bullet(C^\infty(X), Q)$$

is a morphism of the algebras. Then the arguments analogous to Lemma 3.3.1 shows that $[\mathcal{U}_Q]$ is in fact an *isomorphism* of the algebras.

It would be very interesting to find a description of the tangent map

$$\mathcal{U}_Q: T_{\text{poly}}^\bullet(C^\infty(X), Q) \rightarrow CH^\bullet(C^\infty(X), Q),$$

analogous to the description given in Theorem 6 in the local case. The problem is that the L_∞ -quasi-isomorphism $\mathcal{U}: T_{\text{poly}}^\bullet(X) \rightarrow \mathcal{D}_{\text{poly}}^\bullet(X)$ is not defined canonically and therefore the tangent map \mathcal{U}_Q also is not defined canonically. On the other hand, the map

$$[\mathcal{U}_Q]: H^\bullet T_{\text{poly}}^\bullet(C^\infty(X), Q) \rightarrow HH^\bullet(C^\infty(X), Q)$$

is defined canonically. The Conjecture 5.2 below (Duflo formula in the case of Q -manifolds) describes explicitly the map $[\mathcal{U}_Q]$.

5.1.1. One can apply Conjecture 2.3 instead of Conjecture 5.1. Indeed, the DG algebra $(C^\infty(X), Q)$ is quasi-isomorphic to a finite-dimensional L_∞ -algebra $(\mathfrak{g}[1], \tilde{Q})$. It is easy to see that the Hochschild cohomology of quasi-isomorphic DG algebras coincides. On the other hand, Conjecture 2.3 states that

$$H^\bullet(T_{\text{poly}}^\bullet(C^\infty(X), Q)) \simeq H^\bullet(T_{\text{poly}}^\bullet(\mathfrak{g}[1], \tilde{Q}))$$

as algebras. However, this approach does not lead us to the explicit formula.

5.2. In this section we formulate a Conjecture about the tangent map $[\mathcal{U}_Q]$.

5.2.1. *The Atiyah class in Lie algebra cohomology.* Let X be a smooth super-manifold, $\text{Vect}(X)$ be the graded Lie algebra of the smooth vector fields on X . Let TX be the tangent bundle of the manifold X . We denote by $J^1(TX)$ the bundle of 1-jets of the tangent bundle. The space of global sections $\Gamma_X(J^1(TX))$ has a natural structure of $\text{Vect}(X)$ -module. There exists the canonical map of the $\text{Vect}(X)$ -modules

$$\tilde{p}: \Gamma_X(J^1(TX)) \rightarrow \text{Vect}(X).$$

It is clear that the kernel of the map \tilde{p} is the $\text{Vect}(X)$ -module $\Omega_X^1 \otimes_{C^\infty(X)} \text{Vect}(X)$. We obtain a short exact sequence

$$(9) \quad 0 \rightarrow \Omega_X^1 \otimes_{C^\infty(X)} \text{Vect}(X) \rightarrow \Gamma_X(J^1(TX)) \rightarrow \text{Vect}(X) \rightarrow 0.$$

Let us note that both maps in (9) are maps of $C^\infty(X)$ -modules. The short exact sequence (9) defines the ‘‘Atiyah class’’

$$(10) \quad c_T \in C_{\text{Lie}}^1(\text{Vect}(X); \Omega_X^1 \otimes_{C^\infty(X)} \text{Hom}_{C^\infty(X)}(\text{Vect}(X), \text{Vect}(X))).$$

5.2.2. Let X be a smooth Q -manifold. The value $c_T(Q)$ of the Atiyah class (10) on the odd vector field Q gives us an element

$$c_T(Q) \in \Omega_X^1 \otimes_{C^\infty(X)} \text{Hom}_{C^\infty(X)}(\text{Vect}(X), \text{Vect}(X)).$$

This is an explicit analogue of the Atiyah class in the case of L_∞ -algebras given by formula (7).

The k -th power of the element $c_T(Q)$ is a map

$$c_T^k(Q): \text{Vect}^{\otimes k}(X) \rightarrow \text{Hom}_{C^\infty(X)}^{\otimes k}(\text{Vect}(X), \text{Vect}(X)).$$

Futhermore, there exists the trace map

$$\text{Tr}: \text{Hom}_{C^\infty(X)}^{\otimes k}(\text{Vect}(X), \text{Vect}(X)) \rightarrow C^\infty(X),$$

and we obtain an element

$$(11) \quad c_k = \text{Tr} \circ c_T^k(Q): \text{Vect}^{\otimes k}(X) \rightarrow C^\infty(X)$$

After the (super-) symmetrization we can consider c_k as operators

$$c_k: T_{\text{poly}}^\bullet(X) \rightarrow T_{\text{poly}}^\bullet(X).$$

Lemma. *The map c_k is a map of the complexes*

$$c_k: T_{\text{poly}}^\bullet(X, Q) \rightarrow T_{\text{poly}}^\bullet(X, Q).$$

Proof. It is sufficient to prove that $[Q, c_T(Q)] = 0$. But c_T is a 1-cocycle on $\text{Vect}(X)$, and therefore $(d_{\text{Lie}} c_T)(Q, Q) = 0$. By the definition of the cochain differential d_{Lie} , we have

$$(d_{\text{Lie}} c_T)(Q, Q) = c_T([Q, Q]) - 2[Q, c_T(Q)].$$

The desired result follows now from the identity $[Q, Q] = 0$. □

We set:

$$\varphi_{\text{strange}} = \exp \left(\sum_{k \geq 1} \alpha_{2k} c_{2k} \right) : T_{\text{poly}}^\bullet(X, Q) \rightarrow T_{\text{poly}}^\bullet(X, Q),$$

where the numbers α_{2k} are defined by formula (3).

Conjecture (Duflo formula for Q -manifolds). *Let X be a smooth Q -manifold.*

(i) *the map*

$$[\varphi_{\text{HKR}} \circ \varphi_{\text{strange}}]: H^\bullet(T_{\text{poly}}^\bullet(X, Q)) \rightarrow HH^\bullet(C^\infty(X), Q)$$

is an isomorphism of the algebras;

(ii) *the map $[\mathcal{U}_Q]$, induced by the tangent map \mathcal{U}_Q , coincides with the map $[\varphi_{\text{HKR}} \circ \varphi_{\text{strange}}]$.*

5.3. Examples.

5.3.1. (*the de Rham complex*). Let Y be a (purely even) smooth manifold, and let $X = (T[1]Y, d_{\text{DR}})$. We consider X as a Q -manifold, the DG algebra of functions $C^\infty(X, Q)$ coincides with the de Rham complex of the manifold Y .

In the case when $Y = G$ be a (connected compact) Lie group, the corresponding Duflo formula for Q -manifold $(T[1]G, d_{\text{DR}})$ can be considered as the classical Duflo formula for the Lie algebra \mathfrak{g} . We consider the Duflo formula for the Q -manifold $X = (T[1]Y, d_{\text{DR}})$ as an analogue of the classical Duflo formula for smooth manifolds.

5.3.2. (*the Dolbeault complex*). Let Y be a complex manifold, $T_{\text{hol}}Y$ be its holomorphic tangent bundle, $\overline{T}_{\text{hol}}Y$ be its anti-holomorphic tangent bundle. We consider $X = (\overline{T}_{\text{hol}}[1]Y, \overline{\partial})$ as a Q -manifold, the DG algebra of functions $C^\infty(X, Q)$ coincides with the Dolbeault complex of the manifold Y .

There exist at least two different ways to define the notion of the ‘‘Hochschild cohomology of the structural sheaf \mathcal{O}_Y .’’

5.3.2.1 (M. Kontsevich). One can define $HH^\bullet(\mathcal{O}_Y)$ as the algebra of Ext -s $Ext_{\text{Coh}(Y \times Y)}^\bullet(\mathcal{O}_{\text{diag}}, \mathcal{O}_{\text{diag}})$. The direct analogue of the Hochschild–Kostant–Rosenberg theorem states that

$$(12) \quad Ext_{\text{Coh}(Y \times Y)}^k(\mathcal{O}_{\text{diag}}, \mathcal{O}_{\text{diag}}) = \bigoplus_{i+j=k} H_{\text{sheaf}}^i(X, T^j)$$

(here T^j be the sheaf of holomorphic j -polyvector fields on Y). There exist canonical products on both sides of (12): the Yoneda product on the Hochschild cohomology (see formula (5)) and the product induced by the usual cup-product of polyvector fields on the right-hand side.

The ‘‘theorem on complex manifold’’ of M. Kontsevich states that both algebras are canonically isomorphic. Let us recall the construction of this isomorphism.

Let $\alpha_T \in H_{\text{sheaf}}^1(Y, \Omega_{\text{hol}}^1 \otimes_{\mathcal{O}} \text{End } T_{\text{hol}})$ be the Atiyah class of the holomorphic tangent bundle (in the classical sense), it define the elements $c_k = \text{Tr} \circ \alpha_T^k \in H_{\text{sheaf}}^k(Y, \Omega_{\text{hol}}^k)$, which can be considered as the Chern classes of the tangent bundle. One can consider c_k as an operator

$$c_k: H_{\text{sheaf}}^\bullet(Y, T_{\text{poly}}^\bullet) \rightarrow H_{\text{sheaf}}^\bullet(Y, T_{\text{poly}}^\bullet).$$

Theorem (M. Kontsevich). *The map*

$$\varphi_{\text{HKR}} \circ \varphi_{\text{strange}}: H_{\text{sheaf}}^\bullet(Y, T_{\text{poly}}^\bullet) \rightarrow Ext_{\text{Coh}(Y \times Y)}^\bullet(\mathcal{O}_{\text{diag}}, \mathcal{O}_{\text{diag}})$$

is an isomorphism of the algebras.

□

5.3.2.2. We define the Hochschild cohomology $HH^\bullet(\mathcal{O}_Y)$ of the structural sheaf \mathcal{O}_Y as the Hochschild cohomology of the corresponding Dolbeault complex, $HH^\bullet(C^\infty(X), \bar{\partial})$. We claim, that this definition does *not* coincide with the definition given in Sect. 5.3.2.1. Conjecture 5.2 states, that the algebras $HH^\bullet(C^\infty(X), \bar{\partial})$ and $H^\bullet T_{\text{poly}}^\bullet(C^\infty(X), \bar{\partial})$ are isomorphic, and gives an explicit formula for the isomorphism.

We claim, that $\bigoplus_{i,j} H^i(Y, T_{\text{poly}}^j)$ is a *proper* subalgebra of the algebra $H^\bullet T_{\text{poly}}^\bullet(C^\infty(X), \bar{\partial})$, and $Ext_{\text{Coh}(Y,Y)}^\bullet(\mathcal{O}_{\text{diag}}, \mathcal{O}_{\text{diag}})$ is a *proper* subalgebra of the algebra $HH^\bullet(C^\infty(X), \bar{\partial})$. Indeed, let us consider the derivations of the Dolbeault complex of the manifold Y of the form $C^\infty(X) \otimes_{\mathcal{O}} w$, where w is a *holomorphic* vector field on Y ; the differential $\text{ad}(\bar{\partial})$ on these derivations is the same as the differential in the Dolbeault complex $D^\bullet(\text{Hol})$ of the sheaf of holomorphic vector fields on Y . It is clear that $\bigwedge_{(C^\infty(X), \bar{\partial})}^\bullet D^\bullet(\text{Hol})$ is a proper DG subalgebra in $T_{\text{poly}}^\bullet(C^\infty(X), \bar{\partial})$; on the other hand,

$$H^\bullet \left(\bigwedge_{(C^\infty(X), \bar{\partial})}^\bullet D^\bullet(\text{Hol}) \right) \simeq \bigoplus H_{\text{sheaf}}^\bullet(Y, T_{\text{poly}}^\bullet).$$

Therefore, Conjecture 5.2 in the case of the Q -manifold $X = (\bar{T}_{\text{hol}}[1]Y, \bar{\partial})$ gives us a generalization of the M. Kontsevich's theorem on complex manifold.

ACKNOWLEDGEMENTS

I am grateful to Boris Feigin and to Maxim Kontsevich for many useful discussions.

REFERENCES

- [D] M. Duflo, Caractères des algèbres de Lie résolubles, C. R. Acad. Sci, 269(1969), série a, p. 437-438
- [K1] M. Kontsevich, *Deformation quantization of Poisson manifolds I*, preprint q-alg/9709040.
- [K2] M. Kontsevich, *Formality conjecture*, preprint.
- [KSh] M. Kontsevich, B. Shoikhet, *Formality conjecture, geometry of complex manifolds, and combinatorics of the graph-complex*, in preparation.
- [Sh] B. Shoikhet, *On the A_∞ -Formality conjecture*, preprint math. QA/9809117

IUM, 11 BOL'SHOJ VLAS'EVSKIJ PER., MOSCOW 121002, RUSSIA
E-mail address: borya@mccme.ru