

# The Representation Theory of Co-triangular Semisimple Hopf Algebras

Pavel Etingof

Department of Mathematics  
Harvard University  
Cambridge, MA 02138

Shlomo Gelaki

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90089

December 18, 1998

## 1 Introduction

In [EG1, Theorem 2.1] we prove that *any* semisimple triangular Hopf algebra  $A$  over an algebraically closed field of characteristic 0 (say the field of complex numbers  $\mathbf{C}$ ) is obtained from a finite group after twisting the ordinary comultiplication of its group algebra in the sense of Drinfeld [D]; that is  $A = \mathbf{C}[G]^J$  for some finite group  $G$  and a twist  $J \in \mathbf{C}[G] \otimes \mathbf{C}[G]$ . In [EG2] we show how to construct twists for certain solvable non-abelian groups by iterating twists of their abelian subgroups, and thus obtain new non-trivial semisimple triangular Hopf algebras. We also show how any non-abelian finite group which admits a bijective 1-cocycle with coefficients in an abelian group, gives rise to a non-trivial semisimple minimal triangular Hopf algebra. Such non-abelian groups (which are necessarily solvable [ESS]) exist in abundance and were constructed in [ESS] in connection with set-theoretical solutions to the quantum Yang-Baxter equation.

If  $A$  is minimal triangular then  $A$  and  $A^{*op}$  are isomorphic as Hopf algebras. But any non-trivial semisimple triangular  $A$  which is not minimal, gives rise to a new Hopf algebra  $A^*$ , which is also semisimple by [LR]. These are very interesting semisimple Hopf algebras which arise from finite groups, and they are abundant by the constructions given in [EG2]. Generally, the dual Hopf algebra of a triangular Hopf algebra is called *co-triangular* in the literature.

In this paper we explicitly describe the representation theory of co-triangular semisimple Hopf algebras  $A^* = (\mathbf{C}[G]^J)^*$  in terms of representations of some associated groups. As a corollary we prove that Kaplansky's 6th conjecture [K] holds for  $A^*$ ; that is that the dimension of any irreducible representation of  $A^*$  divides the dimension of  $A$ .

We note that we have used in an essential way the results of the paper [Mo], from which we learned a great deal.

## 2 Preliminaries

### 2.1 Projective Representations and Central Extensions

Here we recall some basic facts about projective representations and central extensions. They can be found in textbooks, e.g. [CR, Section 11E].

A projective representation of a group  $\Gamma$  is a vector space  $V$  together with a homomorphism of groups  $\pi_V : \Gamma \rightarrow PGL(V)$ , where  $PGL(V) \cong GL(V)/\mathbf{C}$  is the projective linear group.

A linearization of a projective representation  $V$  of  $\Gamma$  is a central extension  $\hat{\Gamma}$  of  $\Gamma$  by a central subgroup  $\zeta$  together with a linear representation  $\tilde{\pi}_V : \hat{\Gamma} \rightarrow GL(V)$  which descends to  $\pi_V$ . If  $V$  is a finite-dimensional projective representation of  $\Gamma$  then there exists a linearization of  $V$  such that  $\zeta$  is finite (in fact, one can make  $\zeta = \mathbf{Z}/(\dim V)\mathbf{Z}$ ).

Any projective representation  $V$  of  $\Gamma$  canonically defines a cohomology class  $[V] \in H^2(\Gamma, \mathbf{C}^*)$ . The representation  $V$  can be lifted to a linear representation of  $\Gamma$  if and only if  $[V] = 0$ .

### 2.2 The Algebras Associated With a Twist

Let  $H$  be a finite group, and let  $J \in \mathbf{C}[H] \otimes \mathbf{C}[H]$  be a *minimal twist* (see [EG1]). That is, the right (and left) components of the R-matrix  $R = J_{21}^{-1}J$  span  $\mathbf{C}[H]$ . Define two coalgebras  $(A_1, \Delta_1, \varepsilon), (A_2, \Delta_2, \varepsilon)$  as follows:  $A_1 = A_2 = \mathbf{C}[H]$  as vector spaces, the coproducts are determined by

$$\Delta_1(x) = (x \otimes x)J, \quad \Delta_2(x) = J^{-1}(x \otimes x)$$

for all  $x \in H$ , and  $\varepsilon$  is the ordinary counit of  $\mathbf{C}[H]$ . Note that since  $J$  is a twist,  $\Delta_1$  and  $\Delta_2$  are indeed coassociative. Clearly the dual algebras  $A_1^*$  and  $A_2^*$  are spanned by  $\{\delta_h | h \in H\}$ , where  $\delta_h(h') = \delta_{hh'}$ .

**Theorem 2.2.1** *Let  $A_1^*$  and  $A_2^*$  be as above. The following hold:*

1.  $A_1^*$  and  $A_2^*$  are  $H$ -algebras via

$$\rho_1(h)\delta_y = \delta_{hy}, \quad \rho_2(h)\delta_y = \delta_{yh^{-1}}$$

*respectively.*

2.  $A_1^* \cong A_2^{*op}$  as  $H$ -algebras (where  $H$  acts on  $A_2^{*op}$  as it does on  $A_2^*$ ).
3. The algebras  $A_1^*$  and  $A_2^*$  are simple, and are isomorphic as  $H$ -modules to the regular representation  $R_H$  of  $H$ .

**Proof:** The proof of part 1 is straightforward.

The proof of part 3 follows from the results in [Mo]. Namely, it follows from [Mo, Proposition 14] that in the case of a minimal twist the group  $St$  defined in [Mo] (which is, by definition, a subgroup of  $H$ ) coincides with  $H$ . Therefore, by [Mo, Propositions 6,7] the algebras  $A_1^*$  and  $A_2^*$  are simple. Furthermore, by [Mo, Propositions 11,12], the actions of  $H$  on  $A_1^*$  and  $A_2^*$  are isomorphic to the regular representation of  $H$ .

Let us prove part 2. Let  $S_0, \Delta_0, m_0$  denote the standard antipode, coproduct and multiplication of  $\mathbf{C}[H]$ , and define  $Q = m_0(S_0 \otimes I)(J)$ . Then it is straightforward to verify that  $Q$  is invertible, and  $(S_0 \otimes S_0)(J) = (Q \otimes Q)J_{21}^{-1}\Delta_0(Q)^{-1}$  (see e.g. (2.17) in [Ma, Section 2.3]). Hence the map  $A_2^* \rightarrow A_1^{*op}$ ,  $\delta_x \mapsto \delta_{S_0(x)Q^{-1}}$  determines an  $H$ -algebra isomorphism. ■

**Corollary 2.2.2** *Let  $A$  be a semisimple minimal triangular Hopf algebra over  $\mathbf{C}$  with Drinfeld element  $u$ . If  $u = 1$  then  $\dim A$  is a square, and if  $u \neq 1$  then  $2\dim A$  or  $\dim A$  is a square.*

**Proof:** The first statement follows from [EG1, Theorem 2.1] and part 3 of Theorem 2.2.1. To prove the second statement, let  $(A, R, u)$  be a semisimple minimal triangular Hopf algebra with  $u \neq 1$ , and  $(A, R', u')$  be obtained from  $(A, R, u)$  by changing  $R$  so that the new Drinfeld element  $u' = 1$  (as in [EG1]). Then  $(A, R') = (\mathbf{C}[H]^J, J_{21}^{-1}J)$  for some finite group  $H$ . Let  $A_{min} = \mathbf{C}[H']^J$  be the minimal triangular Hopf subalgebra of  $(A, R')$  where  $H' \subset H$  is a subgroup, and  $J$  is a minimal twist for  $H'$ . It is clear that  $H$  is generated by  $H'$  and  $u$ . Since  $u$  is a central grouplike element of order 2, we get that the index of  $H'$  is at most 2. This implies our statement. ■

Let  $A_1^*, A_2^*$  be the  $H$ -algebras as in Theorem 2.2.1. Since the algebras  $A_1^*, A_2^*$  are simple, the actions of  $H$  on  $A_1^*, A_2^*$  give rise to projective representations  $H \rightarrow PGL(|H|^{1/2}, \mathbf{C})$ . We will denote these projective representations by  $V_1, V_2$  (they can be thought of as the simple modules over  $A_1^*, A_2^*$ , with the induced projective action of  $H$ ). Note that part 2 of Theorem 2.2.1 implies that  $V_1, V_2$  are dual to each other, hence that  $[V_1] = -[V_2]$ .

### 3 The Main Result

Let  $(A, R)$  be a semisimple triangular Hopf algebra over  $\mathbf{C}$ , and assume that the Drinfeld element  $u$  is 1 (this can be always achieved by a simple modification of  $R$ , without changing the Hopf algebra structure [EG1]). Then by [EG1, Theorem 2.1], there exist finite groups  $H \subset G$  and a minimal twist  $J \in \mathbf{C}[H] \otimes \mathbf{C}[H]$  such that  $(A, R) \cong (\mathbf{C}[G]^J, J_{21}^{-1}J)$  as triangular Hopf algebras. So from now on we will assume that  $A$  is of this form.

Consider the dual Hopf algebra  $A^*$ . It has a basis of  $\delta$ -functions  $\delta_g$ . The first simple but important fact about the structure of  $A^*$  as an algebra is:

**Proposition 3.1** *Let  $Z$  be a double coset of  $H$  in  $G$ , and  $A_Z^* = \bigoplus_{g \in Z} \mathbf{C}\delta_g \subset A^*$ . Then  $A_Z^*$  is a subalgebra of  $A^*$ , and  $A^* = \bigoplus_Z A_Z^*$  as algebras.*

**Proof:** Straightforward. ■

Thus, to study the representation theory of  $A^*$ , it is sufficient to describe the representations of  $A_Z^*$  for any  $Z$ .

Let  $Z$  be a double coset of  $H$  in  $G$ , and let  $g \in Z$ . Let  $K_g = H \cap gHg^{-1}$ , and define the embeddings  $\theta_1, \theta_2 : K_g \rightarrow H$  given by  $\theta_1(a) = g^{-1}ag$ ,  $\theta_2(a) = a$ . Denote by  $W_i$  the pullback of the projective  $H$ -representation  $V_i$  to  $K_g$  by means of  $\theta_i$ ,  $i = 1, 2$ .

Our main result is the following theorem, which is proved in the next section.

**Theorem 3.2** *Let  $W_1, W_2$  be as above, and let  $(\hat{K}_g, \tilde{\pi}_W)$  be any linearization of the projective representation  $W = W_1 \otimes W_2$  of  $K_g$ . Let  $\zeta$  be the kernel of the projection  $\hat{K}_g \rightarrow K_g$ , and  $\chi : \zeta \rightarrow \mathbf{C}^*$  be the character by which  $\zeta$  acts in  $W$ . Then there exists a 1-1 correspondence between isomorphism classes of irreducible representations of  $A_Z^*$  and isomorphism classes of irreducible representations of  $\hat{K}_g$  with  $\zeta$  acting by  $\chi$ . If a representation  $Y$  of  $A_Z^*$  corresponds to a representation  $X$  of  $\hat{K}_g$  then  $\dim Y = \frac{|H|}{|K_g|} \dim X$ .*

As a corollary we get Kaplansky's 6th conjecture [K] for semisimple co-triangular Hopf algebras.

**Corollary 3.3** *The dimension of any irreducible representation of a semisimple co-triangular Hopf algebra divides the dimension of the Hopf algebra.*

**Proof:** Since  $\dim X$  divides  $|K_g|$  (see e.g. [CR, Proposition 11.44]), we have that  $\frac{|G|}{\frac{|H|}{|K_g|} \dim X} = \frac{|G|}{|H|} \frac{|K_g|}{\dim X}$  and the result follows. ■

In some cases the classification of representations of  $A_Z^*$  is even simpler. Namely, let  $\bar{g} \in \text{Aut}(K_g)$  be given by  $a \mapsto g^{-1}ag$ . Then we have:

**Corollary 3.4** *If the cohomology class  $[W_1]$  is  $\bar{g}$ -invariant then irreducible representations of  $A_Z^*$  correspond in a 1-1 manner to irreducible representations of  $K_g$ , and if  $Y$  corresponds to  $X$  then  $\dim Y = \frac{|H|}{|K_g|} \dim X$ .*

**Proof:** For any  $\alpha \in \text{Aut}(K_g)$  and  $f \in \text{Hom}((K_g)^n, \mathbf{C}^*)$ , let  $\alpha \circ f \in \text{Hom}((K_g)^n, \mathbf{C}^*)$  be given by  $(\alpha \circ f)(h_1, \dots, h_n) = f(\alpha(h_1), \dots, \alpha(h_n))$  (which determines the action of  $\alpha$  on  $H^i(K_g, \mathbf{C}^*)$ ). Then it follows from the identity  $[V_1] = -[V_2]$ , given at the end of Section 2, that  $[W_1] = -\bar{g} \circ [W_2]$ . Thus, in our situation  $[W] = 0$ , hence  $W$  comes from a linear representation of  $K_g$ . Thus, we can set  $\hat{K}_g = K_g$  in the theorem, and the result follows. ■

**Example 3.5** Let  $p > 2$  be a prime number, and  $H = (\mathbb{Z}/p\mathbb{Z})^2$  with the standard symplectic form  $(, ) : H \times H \rightarrow \mathbf{C}^*$  given by  $((x, y), (x', y')) = e^{2\pi i(xy' - yx')/p}$ . Then the element

$J = p^{-2} \sum_{a,b \in H} (a,b)a \otimes b$  is a minimal twist for  $\mathbf{C}[H]$ . Let  $g \in GL_2(\mathbb{Z}/p\mathbb{Z})$  be an automorphism of  $H$ , and  $G_0$  be the cyclic group generated by  $g$ . Let  $G$  be the semidirect product of  $G_0$  and  $H$ . It is easy to see that in this case, the double cosets are ordinary cosets  $g^k H$ , and  $K_{g^k} = H$ . Moreover, one can show either explicitly or using [Mo, Proposition 9], that  $[W_1]$  is a generator of  $H^2(H, \mathbf{C}^*)$  which is isomorphic to  $\mathbb{Z}/p\mathbb{Z}$ . The element  $g^k$  acts on  $[W_1]$  by multiplication by  $\det(g^k)$ . Therefore, by Corollary 3.4, the algebra  $A_{g^k H}^*$  has  $p^2$  1-dimensional representations (corresponding to linear representations of  $H$ ) if  $\det(g^k) = 1$ .

However, if  $\det(g^k) \neq 1$ , then  $[W]$  generates  $H^2(H, \mathbf{C}^*)$ . Thus,  $W$  comes from a linear representation of the Heisenberg group  $\hat{H}$  (a central extension of  $H$  by  $\mathbb{Z}/p\mathbb{Z}$ ) with some central character  $\chi$ . Thus,  $A_{g^k H}^*$  has one  $p$ -dimensional irreducible representation, corresponding to the unique irreducible representation of  $\hat{H}$  with central character  $\chi$  (which is  $W$ ).

## 4 Proof of Theorem 3.2

Let  $Z \subset G$  be a double coset of  $H$  in  $G$ , and let  $A_1, A_2$  be as in Subsection 2.2. For any  $g \in Z$  define the linear map

$$F_g : A_Z^* \rightarrow A_2^* \otimes A_1^*, \quad \delta_y \mapsto \sum_{h,h' \in H: y=hgh'} \delta_h \otimes \delta_{h'}.$$

**Proposition 4.1** *Let  $\rho_1, \rho_2$  be as in Theorem 2.2.1. Then:*

1. *The map  $F_g$  is an injective homomorphism of algebras.*
2.  *$F_{aga'}(\varphi) = (\rho_2(a) \otimes \rho_1(a')^{-1})F_g(\varphi)$  for any  $a, a' \in H$ ,  $\varphi \in A_Z^*$ .*

**Proof:** 1. It is straightforward to verify that the map  $F_g^* : A_2 \otimes A_1 \rightarrow A_Z$  is determined by  $h \otimes h' \mapsto hgh'$ , and that it is a surjective homomorphism of coalgebras. Hence the result follows.

2. Straightforward. ■

For any  $a \in K_g$  define  $\rho(a) \in \text{Aut}(A_2^* \otimes A_1^*)$  by  $\rho(a) = \rho_2(a) \otimes \rho_1(a^g)$ , where  $a^g = g^{-1}ag$  and  $\rho_1, \rho_2$  are as in Theorem 2.2.1. Then  $\rho$  is an action of  $K_g$  on  $A_2^* \otimes A_1^*$ .

**Proposition 4.2** *Let  $U_g = (A_2^* \otimes A_1^*)^{\rho(K_g)}$  be the algebra of invariants. Then  $\text{Im}(F_g) = U_g$ , so  $A_Z^* \cong U_g$  as algebras.*

**Proof:** It follows from Proposition 4.1 that  $\text{Im}(F_g) \subseteq U_g$ , and  $\text{rk}(F_g) = \dim A_Z^* = \frac{|H|^2}{|K_g|}$ . On the other hand, by Theorem 2.2.1,  $A_1^*, A_2^*$  are isomorphic to the regular representation  $R_H$

of  $H$ . Thus,  $A_1^*, A_2^*$  are isomorphic to  $\frac{|H|}{|K_g|}R_{K_g}$  as representations of  $K_g$ , via  $\rho_1(a), \rho_2(a^g)$ . Thus,  $A_2^* \otimes A_1^* \cong \frac{|H|^2}{|K_g|^2}(R_{K_g} \otimes R_{K_g}) \cong \frac{|H|^2}{|K_g|}R_{K_g}$ . So  $U_g$  has dimension  $|H|^2/|K_g|$ , and the result follows. ■

Now we are in a position to prove Theorem 3.2. Since  $W_i \otimes W_i^* \cong A_i^*$  for  $i = 1, 2$ , it follows from Theorem 2.2.1 that  $W_1 \otimes W_2 \otimes W_1^* \otimes W_2^* \cong \frac{|H|^2}{|K_g|}R_{K_g}$  as  $\hat{K}_g$  modules. Thus, if  $\chi_w$  is the character of  $W = W_1 \otimes W_2$  as a  $\hat{K}_g$  module then

$$|\chi_w(x)|^2 = 0, x \notin \zeta \text{ and } |\chi_w(x)|^2 = |H|^2, x \in \zeta.$$

Therefore,

$$\chi_w(x) = 0, x \notin \zeta \text{ and } \chi_w(x) = |H| \cdot x_w, x \in \zeta,$$

where  $x_w$  is the root of unity by which  $x$  acts in  $W$ . Now, it is clear from the definition of  $U_g$  (see Proposition 4.2) that  $U_g = \text{End}_{\hat{K}_g}(W)$ . Thus if  $W = \bigoplus_{M \in \text{Irr}(\hat{K}_g)} W(M) \otimes M$ , where  $W(M) = \text{Hom}_{\hat{K}_g}(M, W)$  is the multiplicity space, then  $U_g = \bigoplus_{M: W(M) \neq 0} \text{End}_{\mathbf{C}}(W(M))$ . So  $\{W(M) | W(M) \neq 0\}$  are the irreducible representations of  $U_g$ . Thus the following implies the theorem:

**Lemma.**

1.  $W(M) \neq 0$  if and only if for all  $x \in \zeta$ ,  $x_{|M} = x_{|W}$ .
2. If  $W(M) \neq 0$  then  $\dim W(M) = \frac{|H|}{|K_g|} \dim M$ .

**Proof of the Lemma.** The "only if" part of 1 is clear. For the "if" part compute  $\dim W(M)$  as the inner product  $(\chi_w, \chi_M)$ . We have

$$(\chi_w, \chi_M) = \sum_{x \in \zeta} \frac{|H|}{|\hat{K}_g|} x_{|W} \cdot \dim M \cdot \bar{x}_{|M}.$$

If  $x_{|M} = x_{|W}$  then

$$(\chi_w, \chi_M) = \sum_{x \in \zeta} \frac{|H|}{|\hat{K}_g|} \dim M = \frac{|H||\zeta|}{|\hat{K}_g|} \dim M = \frac{|H|}{|K_g|} \dim M.$$

This proves part 2 as well, and hence concludes the proof of the theorem. ■

## References

- [CR] C. Curtis and I. Reiner, *Methods of Representation Theory* **1**, John Wiley & Sons, Inc. (1981).
- [D] V. Drinfeld, On Almost Cocommutative Hopf Algebras, *Leningrad Math. J* **1** (1990), 321-342.
- [EG1] P. Etingof and S. Gelaki, Some Properties of Finite-Dimensional Semisimple Hopf Algebras, *Mathematical Research Letters* **5** (1998), 191-197.
- [EG2] P. Etingof and S. Gelaki, A Method of Construction of Finite-Dimensional Triangular Semisimple Hopf Algebras, *Mathematical Research Letters* **5** (1998), 551-561.
- [ESS] P. Etingof, T. Schedler and A. Soloviev, Set-Theoretic Solutions to the Quantum Yang-Baxter Equation, *Duke Math. J.*, to appear, math.QA/9801047.
- [K] I. Kaplansky, *Bialgebras*, University of Chicago, 1975.
- [LR] R. G. Larson and D. E. Radford, Finite-Dimensional Cosemisimple Hopf Algebras in Characteristic 0 are Semisimple, *J. Algebra* **117** (1988), 267-289.
- [Ma] S. Majid, *Foundations of quantum group theory*, Cambridge University Press, 1995.
- [Mo] M. Movshev, Twisting in group algebras of finite groups, *Func. Anal. Appl.* **27** (1994), 240-244.