

Quantum Analogue of the Neumann Function of Integer Order

H. Ahmedov¹ and I. H. Duru^{2,1}

1. Feza Gürsey Institute, P.O. Box 6, 81220, Çengelköy, Istanbul, Turkey ¹.
2. Trakya University, Mathematics Department, P.O. Box 126, Edirne, Turkey.

Abstract: q -Neumann function of integer order $N_n(x; q)$ is obtained and some of its properties are given. q -Psi function which is used in deriving $N_n(x; q)$ is also introduced and some of its properties are presented.

March 1999

1. Introduction

The Hahn-Exton q -Bessel functions $J_\nu(x; q)$ which are closely connected to the quantum group of the plane motions are well studied [1, 2, 3, 4]. Note that for $\nu = n$ is integer the functions $J_n(x; q)$ and $J_{-n}(x; q)$ are not independent of each other. To our knowledge the quantum analogues of the Bessel functions of integer order which are independent of $J_n(x; q)$ have not been addressed.

The main purpose of this note is to introduce the second independent solution $N_n(x; q)$ of the q -Bessel difference equation. This function possess the same recurrence relations as the Hahn-Exton q -Bessel function and in $q \rightarrow 1^-$ becomes Neumann function of the order n . We call it the q -Neumann function of order n . This solution is non regular at $x = 0$. It is well known that non-regular solutions are important, since the Green functions are given

¹E-mail : hagi@gursey.gov.tr and duru@gursey.gov.tr

in terms of them. For example the q -Legendre function of the second kind shows up as the Green function on the quantum sphere [5]. The recently obtained Green function on the quantum plane is in fact the superposition of the Hahn-Exton q -Bessel and q -Neumann functions of the order 0 [6].

Since the classical Neumann function of integer order $N_n(x)$ is obtained by taking the derivatives of the Bessel functions $J_\nu(x)$ and $J_{-\nu}(x)$ with respect to the order ν , it involves the psi functions. Therefore to derive the q -Neumann function of integer order we first have to have q -psi function in hand.

The Section 2 is devoted to the introduction of the q -psi function and some of its properties which are employed to derive the q -Neumann function of integer order.

In Section 3 we obtained the q -Neumann function of integer order and presented some relations involving it.

2. q -Psi Function

We define the q -psi function as

$$\psi_q(\nu) = \frac{d}{d\nu} \log \Gamma_q(\nu), \quad (1)$$

where the q -gamma function $\Gamma_q(\nu)$ is defined by ($0 < q < 1$)

$$\Gamma_q(\nu) = (1 - q)^{1-\nu} \prod_{l=1}^{\infty} \frac{1 - q^l}{1 - q^{\nu+l}}. \quad (2)$$

Many properties of the gamma function were derived by Askey [7]. It is obvious from (2) that $\Gamma_q(\nu)$ has poles at $\nu = 0, 1, 2, \dots$. The residue at $\nu = -n$ is

$$\lim_{\nu \rightarrow -n} (\nu + n) \Gamma_q(\nu) = (-1)^n \frac{(q - 1) q^{-n(n+1)/2}}{\log q \Gamma_q(n + 1)}. \quad (3)$$

The explicit form of $\psi_q(\nu)$ is

$$\psi_q(\nu) = -\log(1 - q) + \log q \sum_{l=0}^{\infty} \frac{q^{\nu+l}}{1 - q^{\nu+l}} \quad (4)$$

The recurrence relations and asymptotic conditions satisfied by this function are

$$\psi_q(\nu + n) = \psi_q(\nu) - \log q \sum_{l=0}^{n-1} \frac{q^{\nu+l}}{1 - q^{\nu+l}}, \quad (5)$$

$$\psi_q(\nu - n) = \psi_q(\nu) + \log q \sum_{l=0}^n \frac{q^{\nu-l}}{1 - q^{\nu-l}} \quad (6)$$

and

$$\lim_{\nu \rightarrow \infty} \psi_q(\nu) = -\log(1 - q); \quad \lim_{\nu \rightarrow -\infty} \psi_q(\nu) = \infty. \quad (7)$$

$\psi_q(\nu)$ has poles at $\nu = 0, 1, 2, \dots$ with the residue

$$\begin{aligned} \lim_{\nu \rightarrow -n} (\nu + n) \psi_q(\nu) &= \log q \lim_{\nu \rightarrow -n} (\nu + n) \sum_{l=0}^{\infty} \frac{q^{\nu+l}}{1 - q^{\nu+l}} \\ &= \log q \lim_{\nu \rightarrow +n} \frac{(\nu + n)}{1 - q^{\nu+n}} = -1 \end{aligned} \quad (8)$$

Equations (3) and (8) imply that

$$\lim_{\nu \rightarrow -n} \frac{\psi_q(\nu)}{\Gamma_q(\nu)} = (-1)^n q^{-n(n+1)/2} \frac{\log q}{1 - q} \Gamma_q(n + 1). \quad (9)$$

Before closing this section we like to present the $q \rightarrow 1^-$ limit of $\psi_q(\nu)$. We first rewrite it as

$$\psi_q(\nu) = \lim_{n \rightarrow \infty} \sum_{l=1}^n \left(\log \frac{1 - q^{l+1}}{1 - q^l} + \frac{q^{\nu+l-1} \log q}{1 - q^{\nu+l-1}} \right) \quad (10)$$

Taking the $q \rightarrow 1^-$ limit in the finite sum in the above formula we have

$$\begin{aligned} \lim_{q \rightarrow 1^-} \psi_q(\nu) &= \lim_{n \rightarrow \infty} \left(\log(n + 1) - \sum_{l=1}^n \frac{1}{\nu + l - 1} \right) \\ &= \lim_{n \rightarrow \infty} \left(\log(n + 1) - \sum_{l=1}^{n+1} \frac{1}{l} + \sum_{l=1}^n \left(\frac{1}{l} - \frac{1}{\nu + l - 1} \right) + \frac{1}{n + 1} \right) \\ &= -C + \sum_{l=0}^{\infty} \left(\frac{1}{l + 1} - \frac{1}{\nu + l} \right) = \psi(\nu), \end{aligned} \quad (11)$$

where

$$C = \lim_{n \rightarrow \infty} \left(\sum_{l=1}^n \frac{1}{l} - \log n \right) = -\psi(1) \quad (12)$$

is the Euler number [8]. It is then natural to define the q-Euler number

$$C_q \equiv -\psi_q(1). \quad (13)$$

3. q-Neumann Function of Integer Order

The Hahn-Exton q-Bessel function of order ν is defined as

$$J_\nu(x; q) = \sum_{k=0}^{\infty} \frac{(-1)^k q^{\frac{k(k+1)}{2}}}{\Gamma_q(k+1)\Gamma_q(k+\nu+1)} x^{2k+\nu} \quad (14)$$

satisfies the q-difference equation

$$J_\nu(q^{1/2}x; q) + J_\nu(q^{-1/2}x; q) + q^{-\nu/2}((1-q)^2x^2 - q^\nu - 1)J_\nu(x; q) = 0. \quad (15)$$

For non-integer ν let us define the function

$$N_\nu(x; q) = \frac{\cos(\pi\nu)J_\nu(x; q) - q^{-\nu/2}J_{-\nu}(q^{-\nu/2}x; q)}{\sin(\pi\nu)} \quad (16)$$

which is the second independent solution of the q-difference equation (15). it satisfies the same recurrence relations as the Hahn-Exton q-Bessel functions. For example we have

$$q^{(\nu+1)/2}N_{\nu+1}(q^{1/2}x; q) - N_{\nu+1}(x; q) = (q-1)xN_\nu(x; q), \quad (17)$$

$$q^{\nu/2}N_\nu(q^{-1/2}x; q) - N_\nu(x; q) = (q-1)xN_{\nu+1}(x; q), \quad (18)$$

Note that for integer $\nu = n$ one has the property

$$J_{-n}(x; q) = (-1)^n q^{n/2} J_n(q^{n/2}x; q). \quad (19)$$

Therefore to derive the form of (16) for integer order we use the L'Hospital rule

$$\pi N_n(x; q) = \frac{d}{d\nu} J_\nu(x; q) \Big|_{\nu=n} - (-1)^n \frac{d}{d\nu} (q^{-\nu/2} J_{-\nu}(q^{-\nu/2}x; q)) \Big|_{\nu=n}. \quad (20)$$

Making use of (9) and (19) for $n \in Z_+$ we get

$$\begin{aligned} \pi N_n(x; q) &= 2J_n(x; q) \log(q^{1/4}x) + \frac{\log q}{1-q} \sum_{k=0}^{n-1} \frac{\Gamma_q(n-k)}{\Gamma_q(k+1)} x^{2k-n} \\ &- \sum_{k=0}^{\infty} \frac{(-1)^k q^{\frac{k(k+1)}{2}} x^{2k+n}}{\Gamma_q(k+1)\Gamma_q(k+n+1)} (\psi_q(k+n+1) + \psi_q(k+1) + k \log q). \end{aligned} \quad (21)$$

Using the recurrence relation (5) and the definition (13) we arrive at

$$\begin{aligned} \pi N_n(x; q) &= 2J_n(x; q) (\log(q^{1/4}x) + C_q) + \frac{\log q}{1-q} \sum_{k=0}^{n-1} \frac{\Gamma_q(n-k)}{\Gamma_q(k+1)} x^{2k-n} \\ &+ \log q \sum_{k=1}^{\infty} \frac{(-1)^k q^{\frac{k(k+1)}{2}} x^{2k+n}}{\Gamma_q(k+1)\Gamma_q(k+n+1)} \left(\sum_{l=1}^{k+n} \frac{q^l}{1-q^l} + \sum_{l=1}^k \frac{1}{1-q^l} \right) \\ &+ \log q \frac{x^n}{\Gamma_q(n+1)} \sum_{k=1}^n \frac{q^k}{1-q^k}. \end{aligned} \quad (22)$$

For $n = 0$ we follow the similar steps and obtain

$$\begin{aligned} \pi N_0(x; q) &= 2J_0(x; q) (\log(q^{1/4}x) + C_q) \\ &+ \log q \sum_{k=1}^{\infty} \frac{(-1)^k q^{\frac{k(k+1)}{2}} x^{2k}}{\Gamma_q(k+1)\Gamma_q(k+1)} \sum_{l=1}^k \frac{1+q^l}{1-q^l}. \end{aligned} \quad (23)$$

It is obvious that (22) and (23) become the usual Neumann functions in $q \rightarrow 1^-$ limit. From the construction it is clear that the q -Neumann functions of integer order satisfy the difference equation (15) and possess all recurrence relations satisfied by the Harh-Exton q -Bessel functions of integer order. We also have

$$N_{-n}(x; q) = (-1)^n q^{n/2} N_n(q^{n/2}x; q). \quad (24)$$

Before closing the section we note that several relations involving the q -Neumann functions can be obtained from those of the Hahn-Exton q -Bessel functions. For example using the product formula [9]

$$\begin{aligned} \sum_{s=-\infty}^{\infty} q^z J_x(q^{z/2}; q) J_{x-\nu}(q^{z/2}; q) J_{\nu}(rq^{(y+\nu+z)/2}; q) &= \\ &= J_0(rq^{(x+y)/2}; q) J_{\nu}(rq^{(\nu+y)/2}; q) \end{aligned} \quad (25)$$

which is valid for $r, x, y, \nu \in C$; $Re(x) > -1$, $|r|^2 q^{1+Re(x)+Re(y)} < 1$ and $r \neq 0$ we obtain the product formula for $x = -y = \nu/2$

$$\sum_{s=-\infty}^{\infty} q^z J_{\nu/2}(q^{z/2}; q) J_{-\nu/2}(q^{z/2}; q) N_{\nu}(rq^{\nu/4+z/2}; q) = J_0(r; q) N_{\nu}(rq^{\nu/4}; q). \quad (26)$$

References

- [1] Koelink, H. T., Duke Math. J., **76**, 483 (1994).
- [2] Vaksman, L. L., Korogodski, L. I., Soviet Math. Dokl., **39**, 173 (1989).
- [3] Woronowicz, S. L., Lett. Math. Phys., **23**, 251 (1991); Commun. Math. Phys., **144**, 417 (1992); Commun. Math. Phys., **136**, 399 (1991).
- [4] Koornwinder, T. H. and Swarttouw, R. F., Trans. Amer. Math. Soc., **333**, 445 (1992).
- [5] Ahmedov, H. and Duru, I. H., J. Phys. A: Math. Gen, **31**, 5741 (1998).
- [6] Ahmedov, H. and Duru, I. H.; Green function on the quantum plane, math.QA/9812108 (1998).
- [7] Askey, R., Applicable Analysis **8**, 125, (1978)
- [8] Gradshteyn, I. S. and Ryzhik, I. M.; Tables of Integrals, Series and Products. Academic Press, New York (1980).
- [9] Koelink, H. T. and Swarttouw, R. F., J. Approx. Theory **81**, 260 (1995).