

EQUIVARIANT AND BOTT-TYPE SEIBERG-WITTEN FLOER HOMOLOGY: PART I

GUOFANG WANG AND RUGANG YE

ABSTRACT. We construct Bott-type and equivariant Seiberg-Witten Floer homology and cohomology for 3-manifolds, in particular rational homology spheres, and prove their diffeomorphism invariance. This paper is a revised version of [26].

TABLE OF CONTENTS

- 1. Introduction
- 2. Preliminaries
- 3. Seiberg-Witten moduli space over Y
- 4. Seiberg-Witten trajectories: transversality
- 5. Index and orientation
- 6. The temporal model and compactification
- 7. Bott-type homology and cohomology
- 8. Invariance
- Appendix A. The ordinary Seiberg-Witten Floer homology
- Appendix B. Local gauge fixing
- Appendix C. A transversality

1. INTRODUCTION

At the very beginning of the development of the Seiberg-Witten gauge theory it was clear that, at least formally, the celebrated instanton homology theory of A. Floer for 3-manifolds (homology spheres) [10] could be adapted to the Seiberg-Witten set-up. Indeed, the original 4-dimensional Seiberg-Witten equation leads naturally to a 3-dimensional Seiberg-Witten equation via a limit process, as first observed by Kronheimer and Mrowka [15]. To establish a Seiberg-Witten Floer homology theory for a 3-manifold Y , the obvious idea is to replace flat connections in Floer's set-up by solutions of the 3-dimensional Seiberg-Witten equation on Y (henceforth called *Seiberg-Witten points*), and instanton trajectories by Seiberg-Witten trajectories, which are solutions of the 4-dimensional Seiberg-Witten equation on the infinite cylinder $Y \times \mathbb{R}$. Note that the Seiberg-Witten points are precisely the critical points of the Seiberg-Witten type Chern-Simons functional, and that the Seiberg-Witten trajectories are precisely the trajectories (negative gradient flow lines) of this functional. Hence the said idea amounts to establishing a Morse-Floer theory for the Seiberg-Witten type Chern-Simons functional. However, one encounters various difficulties when trying to implement this idea. The most serious problem is that the Seiberg-Witten Floer homology for a homology sphere (or rational homology sphere) may depend on the underlying Riemannian metric, and

hence is generally not a diffeomorphism invariant. Indeed, the Euler number of the Seiberg-Witten Floer homology already exhibits dependence on Riemannian metrics, see e.g. [8] [5] and [17]. The purpose of this series of papers, which consists of the present paper (Part I), [28] (Part II) and [29] (Part III), is to resolve this problem.

The trouble of non-invariance is caused by the reducible Seiberg-Witten point, which is the trivial Seiberg-Witten point in the case of a homology sphere: the trivial connection coupled with the zero spinor field. It is a fixed point of the action of the group S^1 of constant gauges. Under reasonable perturbations of the Seiberg-Witten equation, this reducible Seiberg-Witten point always survives. To deal with it, one can use suitable perturbations to make it a transversal point for the Seiberg-Witten equation. Then one can construct a Seiberg-Witten Floer homology, see Appendix A. However, one encounters a serious obstruction when trying to compare the homologies for two different perturbation parameters (e.g. metrics). A canonical way of such comparison is to construct chain maps in terms of parameter-dependent Seiberg-Witten trajectories which connect the 3-dimensional Seiberg-Witten equation of one parameter to that of another. We shall call them *transition trajectories*. The said obstruction is the presence of reducible transition trajectories with negative spectral flow of the linearized Seiberg-Witten operator. Such trajectories are not in transversal position and may appear in the compactification of the moduli spaces of transition trajectories between irreducible Seiberg-Witten points. Consequently, the compactified moduli spaces of transition trajectories may be very pathological and cannot be used to define the desired chain maps.

The appearance of such trajectories can be explained in the following way. The spectral flow along a reducible Seiberg-Witten trajectory for a fixed parameter is 1. That along a reducible transition trajectory from a given generic parameter to a nearby one is also 1. When passing from one generic parameter to another through certain degenerate parameters, the spectral flow jumps and becomes negative. Here, typically, the Seiberg-Witten Floer homology also jumps.

In many ways, the above situation is comparable with the situation of 4-manifolds with $b_2^+ = 1$ where the Seiberg-Witten invariants exhibit dependence on chambers of Riemannian metrics, and the wall crossing phenomena appear. The root of the whole trouble lies in the singularity of the gauge quotient of the configuration space $\mathcal{A}(Y) \times \Gamma(S)$ (see the sequel for its definition). If we factorize the gauge group \mathcal{G} by the subgroup \mathcal{G}^0 of based gauges, then the said singularity is seen to be the same as the singularity of the S^1 quotient of the space $\mathcal{B}^0 = (\mathcal{A}(Y) \times \Gamma(S))/\mathcal{G}^0$, where S^1 is the subgroup of constant gauges.

New Constructions

Since the conventional Seiberg-Witten Floer homology may not be diffeomorphism invariant for rational homology spheres, we seek alternative constructions. Our constructions are based on two ideas: one is that we work on the level of the based gauge quotient, the other is that we multiply the based gauge quotient by a suitable space (e.g. the circle), and then pass to the S^1 quotient. The first idea leads to the Bott-type theory, while the second leads to the equivariant theory. In the present Part I, we present the Bott-type theory. The equivariant theory will be presented in Part II. We would like to emphasize that both the Bott-type and equivariant constructions are natural from the point of view of comparison with

classical Morse-Bott theory and equivariant homology theory. On the other hand, we would like to point out that as shown in [5] and [17] the Euler number of the ordinary Seiberg-Witten Floer homology can be corrected into an invariant (indeed the Euler number of the instanton Floer homology) by adding in an explicit way a certain (metric dependent) term to it. It is therefore natural to seek similar corrections of the entire homology. We believe however that the information provided by such possible corrections is already contained in our constructions. Indeed, detailed comparison of our constructions with the conventional construction should reveal the possible precise forms of desired corrections. We shall discuss this point in more details in [29].

The Morse-Floer-Bott Flow Complex

Now, in the set-up of based gauge quotient, the irreducible part of the moduli space of gauge classes of Seiberg-Witten points consists of finitely many circles, while its reducible part consists of a single point, provided that we choose a generic parameter. These circles and the reducible point are precisely the critical submanifolds of the (Seiberg-Witten type) Chern-Simons functional. Our goal here amounts to establishing a Bott-type Morse-Floer theory for the Chern-Simons functional on the based quotient configuration space \mathcal{B}^0 . The basic strategy is to use the moduli spaces of trajectories between critical submanifolds to send (co)homological chains from one critical submanifold to others. Combining this map with the ordinary boundary operator in (co)homology theory then yields the desired boundary operator for the (co)chain complex. This is a natural extension of Floer's construction and was first used by Austin-Braam [3] and Fukaya [12] in Floer's set-up. The former authors use equivariant differential forms as chains and cochains, while the latter uses "geometric chains". Our core constructions use *generalized cubical singular chains and cochains*. For delicate technical reasons cubical singular chains are more suitable than ordinary singular chains, see [28] for details of this point. For convenience, we shall sometimes use "singular chains (cochains)" to refer to generalized cubical singular chains (cochains).

A major point of the construction using generalized cubical singular chains (and cochains) is that we can restrict to subcomplexes generated by generalized singular cubes which are \mathbf{F} -*transversal* (see Section 7) with respect to collections \mathbf{F} of suitable maps. This is crucial for the invariance proof, see Section 8.

The compactified, suitably defined moduli space of trajectories (flow lines) will be called the *Morse-Floer-Bott flow complex*. The main technical point here is to construct this flow complex along with its projection to critical submanifolds, and to establish its compactness and smooth structure. In Floer's work, the Morse-Floer-Bott flow complex does not appear. Instead, he uses the much simpler Morse-Floer flow complex (indeed only the lower dimensional part of it). In Fukaya's work [12], a Morse-Floer-Bott flow complex appears, but his set-up is the full gauge quotient. Our situation is very different, and the construction of the Morse-Floer-Bott flow complex along with its projection to critical submanifolds is considerably more delicate. We cannot find adequate treatments of this problem in the Seiberg-Witten set-up or instanton set-up in the literature.

A crucial point here is how to define the endpoint projections of trajectories to critical submanifolds. Obviously, one should take the limits of a trajectory at time infinities. The subtlety lies in the choice of gauges. We need to send based gauge classes of trajectories to based gauge classes of critical points. The

trouble is that based gauges on $Y \times \mathbb{R}$ may not give rise to based gauges on Y at time infinities. More precisely, if two trajectories u_1 and u_2 are equivalent under based gauges, their limits at time infinities may not be equivalent under based gauges. To resolve this issue, we first transform a given trajectory into temporal form, and then take endpoint limits. This way we arrive at the essential concept of “temporal projections”. Two other resulting important concepts are “consistent multiple temporal trajectory class” and “consistent piecewise trajectory”, which are used for compactifying our moduli spaces. The consistent condition means e.g. that the temporal endpoint projections of the trajectories in a given piecewise trajectory match each other. We also introduce the concept “twisted time translation”.

We have two different, but equivalent models for our moduli spaces of trajectories. One is the *temporal model* $\mathcal{M}_T^0(S_\alpha, S_\beta)$ (see Section 6), the other is the *fixed-end model* $\mathcal{M}^0(p, q)$ (see Section 4). The Fredholm theory for the Seiberg-Witten trajectory equation is worked out for the second model, and the acquired information is then carried over to the first one. On the other hand, the temporal projections are based on the first model. The ordinary time translation is used for the temporal model, while it is necessary to use the twisted time translation for the fixed-end model.

A number of further delicate issues have to be taken care of in order to compactify the moduli spaces and establish the smooth structure of the compactified moduli spaces. For example, in compactifying the moduli spaces, we have to establish that after suitable gauge adjustments, sequences of temporal Seiberg-Witten trajectories of finite energy converge to consistent piecewise trajectories. Further discussions on the analysis in this regard, in particular on convergence analysis and gluing analysis, can be found in Part II [28].

Spinor Perturbation

To prove the diffeomorphism invariance of the Bott-type Seiberg-Witten Floer (co)homology, we employ the *transition Morse-Floer-Bott flow complex* (along with its projection to critical submanifolds) which is the flow complex built out of the transition trajectories. Here, we have to overcome the obstruction of reducible transition trajectories with negative spectral flow described before. Our strategy for this is to perturb the spinor equation in the transition trajectory equation in order to eliminate these transition trajectories. We utilize the vanishing of the rational homology group to construct suitable vector fields which are equivariant under based gauges. Note that *they are not equivariant under constant gauges*. A desired perturbation is then gotten by adding one of these vector fields to the spinor equation. This is our key technique. Thus the source of our trouble, namely the vanishing of the rational homology group, also works to our benefit - a rather amusing phenomenon.

The same kind of amusing phenomenon occurs in the proof of the transversality along reducible Seiberg-Witten trajectories, see Appendix C. (Note that here we are not talking about transition trajectories.) This transversality plays a role for establishing the smooth structure of certain expanded moduli spaces of transition trajectories with additional parameters, which is used in one stage of the invariance proof, see [28].

Using the transition flow complex with the spinor perturbation we construct the desired chain map from one parameter to another, which induces an isomorphism of the homologies. In this paper, we present the construction of this chain map, while

leaving the proof the isomorphism property to Part II and Part III. Indeed, in Part II the isomorphism property in the equivariant case will be shown. In Part III, we shall present an additional perturbation technique- the cokernel perturbation, and use it to prove the isomorphism property in the Bott-type case of the present paper.

We would like to mention that the spinor perturbations cause additional analytical difficulties which demand special treatments. For example, one has to establish a uniform L^∞ estimate for the spinor part of the transition trajectories. With the presence of the spinor perturbations, the ordinary pointwise maximum principle argument no longer works. Instead, we apply the 3-dimensional Weitzenböck formula (rather than the 4-dimensional one) to obtain an initial local integral estimate in terms of the Seiberg-Witten energy. Then we apply the 4-dimensional Weitzenböck formula and the technique of Moser iteration to derive the desired L^∞ estimate.

Singular Chains/Cochains vs. Differential Forms

Instead of generalized cubical singular chains and cochains on critical submanifolds, we can also use differential forms to build chain and cochain complexes, and therewith a Bott-type Seiberg-Witten Floer homology and cohomology. The boundary operator in this set-up is again constructed by using the Morse-Floer-Bott flow complex along with its projection to critical submanifolds. Here, the fibration property of the projection is essential for the construction. On the other hand, the projection of the transition flow complex to critical submanifolds is not a fibration in general, and hence is not good enough for producing chain maps in the set-up of differential forms. In contrast, in the set-up of generalized cubical singular chains, one has the freedom of using \mathbf{F} -transversal chains for a suitable \mathbf{F} and thereby bypasses this problem, see Section 8.

Nevertheless, we can prove that the differential form or de Rham version of Bott theory is diffeomorphism invariant. Indeed, it is isomorphic to the singular version with real coefficients. The construction of the de Rham version and the proof of its equivalence to the singular version will be given in Part II.

Uniform Formulation for All 3-Manifolds

The Bott-type and equivariant constructions can easily be extended to 3-manifolds with nonzero first Betti number. They can be shown to be isomorphic to the ordinary Seiberg-Witten Floer homology (under the assumption that the first Betti number is nonzero). (For accounts of the ordinary Seiberg-Witten Floer homology, see [18], [25] or Appendix A.) Hence we obtain a uniform formulation of Seiberg-Witten Floer homology for all 3-manifolds. Details will be given in Part III. In the present Part I and Part II, we focus on rational homology spheres. Note that the ordinary Seiberg-Witten Floer homology for manifolds with $b_1 = 1$ is not a diffeomorphism invariant, but its metric dependence is reduced to dependence on polarizations of the first homology.

A major part of the results in this paper were obtained in Spring 1996 while both authors were at Bochum University.

2. PRELIMINARIES

To fix notations, we first recall the definitions of the Seiberg-Witten equations on 3- and 4-dimensional manifolds.

Let (X_0, g) be an oriented Riemannian manifold of dimension n and $Spin^c(X_0)$ the set of isomorphism classes of $spin^c$ structures on X_0 . Consider a $spin^c$ structure $c \in Spin^c(X_0)$ and its associated spinor bundle W and line bundle L . (More precisely, c is a representative of an element in $Spin^c(X_0)$. The homology invariants we are going to construct are independent of the choice of the representative.) We have the associated configuration space $\mathcal{A} \times \Gamma(W)$, where \mathcal{A} denotes the space of smooth unitary connections on L and Γ the space of smooth sections of a vector bundle. (We suppress the dependence on c in the notations.) The gauge transformation group (the group of gauges) is $\mathcal{G} = C^\infty(X_0, S^1)$, where $S^1 \equiv U(1)$ denotes the unit circle in \mathbb{C} . (\mathcal{G} depends only on X_0 .) The action of \mathcal{G} on $\mathcal{A} \times \Gamma(W)$ is defined by $((A, \Phi), g) \rightarrow (A + g^{-1}dg, g^{-1}\Phi)$. This formula also defines the (separate) actions of \mathcal{G} on \mathcal{A} and $\Gamma(W)$. \mathcal{G} acts freely on the subspace of pairs (A, Φ) with $\Phi \neq 0$. Such pairs are called irreducible. The isotropy subgroup at any reducible pair $(A, 0)$ is the subgroup of gauge transformations which are constants on each component of X_0 . If X_0 is connected, we identify it with S^1 . In this case, we fix a reference point $x_0 \in X_0$ and set $\mathcal{G}^0 = \{g \in \mathcal{G} : g(x_0) = 1\}$, which is called the group of based gauges. Then the quotient \mathcal{G}/S^1 is represented by \mathcal{G}^o .

The action of a gauge g will be denoted by g^* . We set $\mathcal{B} = (\mathcal{A} \times \Gamma)/\mathcal{G}$ and $\mathcal{B}^* = (\mathcal{A} \times (\Gamma - \{0\}))/\mathcal{G}$. Let $\Omega^k(X_0)$ denote the space of smooth imaginary valued k -forms, and $\Omega^+(X_0)$ the space of smooth imaginary valued self-dual 2-forms (in the case that $\dim X_0 = 4$).

We define the gauge actions on forms to be the trivial action, i.e. $g^*\alpha = \alpha$. Note that the tangent space of the configuration space $\mathcal{A} \times \Gamma(W)$ at any (A, Φ) can be identified with the space $\Omega^1(X_0) \times \Gamma(W)$. The induced gauge action on this space is then the product action, more precisely, $g^*(\alpha, \Phi) = (g^*\alpha, g^*\Phi) = (\alpha, g^{-1}\Phi)$.

We shall need the following

Lemma 2.1. *Assume that X_0 is closed. Then the map from \mathcal{G} to $H^1(X_0; \mathbb{Z})/\{\text{torsions}\}$ given by $g \rightarrow$ the deRham class of $g^{-1}dg$ is surjective and induces an isomorphism from the component group of \mathcal{G} to $H^1(X_0; \mathbb{Z})/\{\text{torsions}\}$. Moreover, there is a unique harmonic map g with $g(x_0) = 1$ in each component of \mathcal{G} , provided that X_0 is connected and $x_0 \in X_0$ is a fixed point. In particular, \mathcal{G} is connected if X_0 is connected and $H^1(X_0; \mathbb{Z})$ is torsion.*

Proof. For simplicity, assume that X_0 is connected. The surjectivity of the said map follows from integration along paths. If $g^{-1}dg$ and $g_1^{-1}dg_1$ represent the same cohomology class, then $g_1 = ge^f$ for some $f \in \Omega^0(X_0)$ as one easily sees. Hence g_1 and g lie in the same component group.

The statement about harmonic representative follows from the standard theory of harmonic maps. It can also be derived quickly in an elementary way. For example, if $g_1 = ge^f$ and g are two harmonic maps, then f is a harmonic function, hence constant. \square

We continue with the above $spin^c$ structure c on X_0 . A connection $A \in \mathcal{A}$ induces along with the Levi-Civita connection a connection ∇^A on the spinor bundle W and the associated Dirac operator $D_A : \Gamma(W) \rightarrow \Gamma(W)$,

$$D_A = \sum_{e_i}^n e_i \cdot \nabla_{e_i}^A,$$

where $\{e_i\}$ denotes a local orthonormal tangent frame and the dot denotes the Clifford multiplication. The Dirac operator is gauge equivariant, i.e. $D_A(g^{-1}\Phi) = g^{-1}D_A\Phi$, and satisfies the following fundamental Weitzenböck formula for the Dirac operator

$$(2.1) \quad D_A^*D_A\Phi = -\Delta_A\Phi + \frac{s}{4}\Phi - \frac{1}{2}F_A \cdot \Phi,$$

where s denotes the scalar curvature of (X, g) and F_A the curvature of A .

Now we specify to the dimension $n = 4$. There is a canonical decomposition $W = W^+ \oplus W^-$ of the spinor bundle W . The Dirac operator splits: $D_A : \Gamma(W^+) \rightarrow \Gamma(W^-)$, $D_A : \Gamma(W^-) \rightarrow \Gamma(W^+)$. For a positive spinor field $\Phi \in \Gamma(W^+)$, the curvature F_A in the above Weitzenböck formula reduces to its self-dual part F_A^+ .

Definition 2.2. The Seiberg-Witten equation with the given $spin^c$ structure c is

$$(2.2) \quad \begin{cases} F_A^+ &= \frac{1}{4}\langle e_i e_j \Phi, \Phi \rangle e^i \wedge e^j, \\ D_A \Phi &= 0, \end{cases}$$

for $(A, \Phi) \in \mathcal{A} \times \Gamma(W^+)$, where $\{e^i\}$ denotes the dual of $\{e_i\}$ (a local orthonormal tangent frame). The Seiberg-Witten operator is

$$\mathbf{SW}(A, \Phi) = (F_A^+ - \frac{1}{4}\langle e_i e_j \Phi, \Phi \rangle e^i \wedge e^j, D_A \Phi).$$

It is easy to see that the Seiberg-Witten operator is gauge equivariant, i.e. $\mathbf{SW}(g^*A, g^*\Phi) = g^*\mathbf{SW}(A, \Phi) = (F_A^+ - \frac{1}{4}\langle e_i e_j \Phi, \Phi \rangle e^i \wedge e^j, g^{-1}D_A\Phi)$. Consequently, the Seiberg-Witten equation is gauge invariant.

Next let (Y, h) be an oriented, closed Riemannian 3-manifold with metric h , and c a $spin^c$ structure on Y . We have the associated spinor bundle $S = S_c(Y)$, line bundle $L_Y = L_Y(c)$ and the other associated spaces: $\mathcal{G}(Y)$, $\mathcal{A}(Y)$, $\mathcal{B}(Y)$, $\mathcal{B}^*(Y)$ etc. We shall often use $\Gamma(Y)$ to denote $\Gamma(S)$. We set $X = Y \times \mathbb{R}$, which will be equipped with the product metric and given the orientation $(e_1, e_2, e_3, \frac{\partial}{\partial t})$, where (e_1, e_2, e_3) denotes a positive local orthonormal frame on Y . Let $\pi : X \rightarrow Y$ denote the projection. The $spin^c$ structure c induces a $spin^c$ structure π^*c on X with the associated line bundle $L_X = \pi^*L_Y$ and associated spinor bundles $W^+ = \pi^*S$, W^- . We have the following relation between the Clifford multiplications on S and on W^+ :

$$v \cdot \phi(y) = -(\frac{\partial}{\partial t} \cdot v \cdot \pi^*\phi)(y, 0).$$

The associated spaces for X will be indicated by the letter X . We also use $\Gamma^+(O)$ and $\Gamma^-(O)$ to denote $\Gamma(W^+|_O)$ and $\Gamma(W^-|_O)$ respectively for $O \subset X$. Let $i_t : Y \rightarrow X$ denote the inclusion map which sends $y \in Y$ to $(y, t) \in X$. A connection $A \in \mathcal{A}(X)$ can be written as

$$A = a(t) + f(\cdot, t)dt,$$

where $a(t) = i_t^*(A) \in \mathcal{A}(Y)$ and $f \in C^\infty(X, i\mathbb{R})$. We set $\phi(t) = \Phi(\cdot, t) = i_t^*(\Phi)$ for $\Phi \in \Gamma(W^+)$. With these notations, we have

$$(2.3) \quad F_A = F_c + dt \wedge \frac{\partial a}{\partial t} + d_Y f \wedge dt.$$

$$(2.4) \quad 2F_A^+ = dt \wedge \left(\frac{\partial a}{\partial t} - *_Y F_a - d_Y f \right) - *_Y \left(\frac{\partial a}{\partial t} - *_Y F_a - df \right),$$

and

$$(2.5) \quad D_A \Phi = \frac{\partial}{\partial t} \cdot \left(\frac{\partial \phi}{\partial t} + \not\partial_a p + f \phi \right).$$

where F_a denotes the curvature of a , $*_Y$ the Hodge star operator w.r.t. h , d_Y the differential on Y , and $\not\partial_a$ the Dirac operator associated with the connection a . Hence we can rewrite (2.2) as follows

$$(2.6) \quad \begin{cases} \frac{\partial a}{\partial t} &= *_Y F_a + d_Y f + \langle e_i \cdot \phi, \phi \rangle e^i, \\ \frac{\partial \phi}{\partial t} &= -\not\partial_a \phi - f \phi. \end{cases}$$

Here F_a denotes the curvature of a , $*_Y$ the Hodge star operator w.r.t. h , d_Y the exterior differential on Y , and $\not\partial_a$ the Dirac operator associated with the connection a .

Definition 2.3. The Seiberg-Witten energy of (A, Φ) is

$$E(A, \Phi) = \int_X \left(\left| \frac{\partial \phi}{\partial t} + f \phi \right|^2 + |\not\partial \phi|^2 + \left| \frac{\partial a}{\partial t} - d_Y f \right|^2 + \left| *_Y F_a + \langle e_i \cdot \phi, \phi \rangle e^i \right|^2 \right).$$

(The volume form is omitted.) One readily shows that it is gauge invariant. Since $F_A = F_a + dt \wedge \left(\frac{\partial a}{\partial t} - d_Y f \right)$, we have

$$E(A, \Phi) = \int_X \left(\left| \frac{\partial \phi}{\partial t} + f \phi \right|^2 + |\not\partial \phi|^2 + |F_A + \langle e_i \cdot \phi, \phi \rangle *_Y e^i|^2 \right).$$

Using the finite energy condition one easily derives from (2.6) the following limiting equation for a connection $a \in \mathcal{A}(Y)$ and a spinor field $\phi \in \Gamma(S)$

$$(2.7) \quad \begin{cases} *_Y F_a + \langle e_i \cdot \phi, \phi \rangle e^i &= 0, \\ \not\partial_a \phi &= 0. \end{cases}$$

Definition 2.4. The Seiberg-Witten equation on Y with the $spin^c$ structure c is defined to be (2.7). The Seiberg-Witten operator on Y is

$$\mathbf{sw}(a, \phi) = (*_Y F_a + \langle e_i \cdot \phi, \phi \rangle e^i, -\not\partial_a \phi).$$

As in dimension 4, the Seiberg-Witten operator \mathbf{sw} is gauge equivariant.

The following lemma is obvious.

Lemma 2.5. *Let $u = (A, \Phi) = (a + f dt, \phi)$ be a solution of (2.2). Denote the energy of u on a domain Ω by $E(u, \Omega)$. Then we have*

$$(2.8) \quad E(u, \Omega) = 2 \int_{\Omega} |\mathbf{sw}(a, \phi)|^2.$$

We shall need the following perturbed Seiberg-Witten equation

$$(2.9) \quad \begin{cases} *_Y F_a + \langle e_i \cdot \phi, \phi \rangle e^i & = \nabla H(a), \\ \not\partial_a \phi + \lambda \phi & = 0, \end{cases}$$

where λ denotes a real number and ∇H the L^2 -gradient of a $\mathcal{G}(Y)$ -invariant real valued function H on $\mathcal{A}(Y)$. (The L^2 -product is given in (2.11) below.) We have the associated perturbed Seiberg-Witten operator $\mathbf{sw}_{\lambda, H}$. Note that ∇H is gauge equivariant and belongs to $\ker d^*$, which are consequences of the gauge invariance of H .

The classical Chern-Simons functional plays a fundamental role in Floer's instanton homology theory. Similarly, a Chern-Simons functional associated with the 3-dimensional Seiberg-Witten equation will be important in our situation. This functional was first used by Kronheimer and Mrowka in their proof of the Thom conjecture [15].

Definition 2.6. The Chern-Simons functional with respect to a reference connection a_0 is

$$\mathbf{cs}(a, \phi) = \frac{1}{2} \int_Y (a - a_0) \wedge (F_a + F_{a_0}) + \int_Y \langle \phi, \not\partial_a \phi \rangle.$$

Let λ and H be as above. The perturbed Chern-Simons functional with perturbation (λ, H) is

$$\begin{aligned} \mathbf{cs}_{\lambda, H}(a, \phi) &= \frac{1}{2} \int_Y (a - a_0) \wedge (F_a + F_{a_0}) + \int_Y \langle \phi, \not\partial_a \phi \rangle \\ &\quad - \lambda \int_Y \langle \phi, \phi \rangle + H(a, \phi). \end{aligned}$$

Under a gauge g the perturbed Chern-Simons functional changes as follows:

$$(2.10) \quad \mathbf{cs}_{\lambda, H}(g^*(a, \phi)) = \mathbf{cs}_{(\lambda, H)}(a, \phi) + 2\pi i \int_Y c_1(L(Y)) \wedge g^{-1} dg.$$

This formula implies that $\mathbf{cs}_{\lambda, H}$ is invariant under the identity component of $\mathcal{G}(Y)$. Hence it descends to the quotient $\mathcal{B}(Y)$, provided that Y is a rational homology sphere.

We introduce an L^2 -product on $\Omega^1(Y) \oplus \Gamma(S)$:

$$(2.11) \quad \langle (\phi_1, a_1), (\phi_2, a_2) \rangle_{L^2} = \int_Y (Re \langle \phi_1, \phi_2 \rangle + \langle a_1, a_2 \rangle)$$

Here, $\langle a_1, a_2 \rangle$ denotes the (pointwise) Hermitian product. Easy computations lead to

Lemma 2.7. *The L^2 -gradient of the perturbed Chern-Simons functional is given by*

$$\nabla \mathbf{cs}_{\lambda, H}(a, \phi) = -\mathbf{sw}_{\lambda, H}.$$

It follows that the critical points of the perturbed Chern-Simons functional are precisely the solutions of the perturbed Seiberg-Witten equation.

Consider a solution (A, Φ) of the Seiberg-Witten equation on the product X . Using a suitable gauge we can transform it into temporal form. Let's assume that

it is already in temporal form, i.e. $f \equiv 0$ in the formula $A = a + f dt$. Then Lemma 2.7 and the equation (2.6) imply $(\phi(t) = \Phi(\cdot, t))$

$$(2.12) \quad \frac{\partial}{\partial t}(a, \phi) = -\nabla \mathbf{cs}(a, \phi).$$

Hence solutions of the Seiberg-Witten equation on the product X can be interpreted as trajectories (negative gradient flow lines) of the Chern-Simons functional. A similar formula and statement hold for solutions of the perturbed Seiberg-Witten equation on X , which is

$$(2.13) \quad \begin{cases} F_A^+ = \frac{1}{4} \langle e_i e_j \Phi, \Phi \rangle e^i \wedge e^j \\ \quad + \nabla H(a) \wedge dt + *(\nabla H(a) \wedge dt), \\ D_A \Phi = -\lambda \frac{\partial}{\partial t} \cdot \Phi, \end{cases}$$

where $\frac{d}{dt} \cdot \Phi$ denotes the Clifford multiplication on X , $A = a + f dt$, $\phi(t) = \Phi(\cdot, t)$ as before, and $*$ means the Hodge star operator on X . The operator $\mathbf{SW}_{\lambda, H}$ is defined in an obvious way. Obviously, we can rewrite (2.13) as follows

$$(2.14) \quad \begin{cases} \frac{\partial a}{\partial t} = *_Y F_a + d_Y f + \langle e_i \cdot \phi, \phi \rangle e^i - \nabla H(a), \\ \frac{\partial \phi}{\partial t} = -\not\partial_a \phi - f \phi - \lambda \phi. \end{cases}$$

Next we introduce the perturbed Seiberg-Witten energy :

$$(2.15) \quad \begin{aligned} E_{\lambda, H}(A, \Phi) &= \int_X (|\frac{\partial \phi}{\partial t} + f \phi|^2 + |\not\partial \phi + \lambda \phi|^2 + |\frac{\partial a}{\partial t} - d_Y f|^2 \\ &\quad + |*_Y F_a + \langle e_i \cdot \phi, \phi \rangle e^i - \nabla H(a, \phi)|^2) \\ &= \int_X (|\frac{\partial \phi}{\partial t} + f \phi|^2 + |\not\partial \phi + \lambda \phi|^2 \\ &\quad + |F_A + \langle e_i \cdot \phi, \phi \rangle *_Y e^i - *_Y \nabla H(a, \phi)|^2). \end{aligned}$$

Note that it is invariant under the action of $\mathcal{G}(X)$. We have an analogue of Lemma 2.5.

Lemma 2.8. *Let $u = (A, \Phi) = (a + f dt, \phi)$ be a solution of (2.13). Then we have*

$$(2.16) \quad E_{\lambda, H}(u, \Omega) = 2 \int_{\Omega} |\mathbf{sw}_{\lambda, H}|^2.$$

Lemma 2.9. *Assume that Y is a rational homology sphere. Let $A = a + f dt$ and assume that the gauge equivalence class of (a, ϕ) converges to $\alpha, \beta \in \mathcal{B}(Y)$ as $t \rightarrow -\infty, \infty$ respectively, then we have*

$$(2.17) \quad \begin{aligned} E_{\lambda, H}(A, \Phi) &= 2\mathbf{cs}_{\lambda, H}(\alpha) - 2\mathbf{cs}_{\lambda, H}(\beta) + \int_X |D_A \Phi|^2 \\ &\quad + 2 \int |F_A^+ - \frac{1}{4} \langle e_i e_j \Phi, \Phi \rangle e^i \wedge e^j - dt \wedge \nabla H(a) - *(dt \wedge \nabla H(a))|^2. \end{aligned}$$

In particular, there holds for a solution (A, Φ) of (2.13)

$$(2.18) \quad E_{\lambda, H}(A, \Phi) = 2\mathbf{cs}_{\lambda, H}(\alpha) - 2\mathbf{cs}_{\lambda, H}(\beta).$$

Proof. We have

$$\begin{aligned} & \int_X |D_A \Phi|^2 + 2 \int_X |F_A^+ - \frac{1}{4} \langle e_i e_j \Phi, \Phi \rangle e^i \wedge e^j - dt \wedge \nabla H(a) - *(dt \wedge \nabla H(a))|^2 \\ &= \int_X (|\frac{\partial \phi}{\partial t} + f\phi + \not\partial \phi + \lambda\phi|^2 + |-\frac{\partial a}{\partial t} + d_Y f + *_Y F_a + \langle e_i \cdot \phi, \phi \rangle e^i - \nabla H(a, \phi)|^2) \\ &= E(\Phi, A) + 2 \int \langle \frac{\partial \phi}{\partial t} + f\phi, \not\partial_a \phi + \lambda\phi \rangle + 2 \langle -\frac{\partial a}{\partial t} + d_Y f, *_Y F_a + \langle e_i \cdot \phi, \phi \rangle e^i - \nabla H(a, \phi) \rangle \\ &= 2 \frac{d}{dt} \int (|\phi|^2 + \langle a - a_o, F_a \rangle) + 2H(a) + 2 \int (\langle \frac{\partial \phi}{\partial t}, \not\partial_a \phi \rangle + \langle \frac{\partial a}{\partial t} \phi, \phi \rangle). \end{aligned}$$

Since the metric on $X = Y \times \mathbb{R}$ is the product metric, it follows that

$$\frac{d}{dt} \not\partial_a \phi - \not\partial_a \frac{d}{dt} \phi = \frac{\partial a}{\partial t} \phi.$$

The desired conclusion follows. \square

3. SEIBERG-WITTEN MODULI SPACES OVER Y

We continue with the (Y, h) and c of the last section. While our theory applies to arbitrary closed Y , we assume for convenience that Y is connected. Fix a reference connection a_0 . If $L(Y)$ is a trivial bundle, we choose a_0 to be the trivial connection. We have $\mathcal{A}(Y) = a_0 + \Omega^1(Y)$. We shall use the (l, p) -Sobolev norms (the $L^{l,p}$ -norms) for $l \geq 0$ and $p > 0$:

$$\|u\|_{l,p} = (\sum_{0 \leq k \leq l} \int_Y |\nabla^k u|^p)^{1/p}.$$

Consider the Sobolev spaces $\mathcal{A}_{l,p}(Y)$ and $\Gamma_{l,p}(S)$, which are the completions of $\mathcal{A}(Y)$ and $\Gamma(S)$ with respect to the (l, p) -Sobolev norm respectively. Similarly, we have the Sobolev spaces $\Omega_{l,p}^k(Y)$. The corresponding group of gauges is $\mathcal{G}_{l+1,p}(Y)$, which is the completion of $\mathcal{G}(Y)$ with respect to the $(l+1, p)$ -Sobolev norm.

We need to make a choice of the configuration spaces $\mathcal{A}_{l,p}(Y) \times \Gamma_{l,p}(S)$. We require $3p/(3-lp) > 3$ or $lp > 3$, for then all elements in $\mathcal{A}_{l,p}(Y) \times \Gamma_{l,p}(S)$ are continuous, and hence the holonomy perturbations in the sequel can be performed. Moreover, the corresponding gauges on the product space $X = Y \times \mathbb{R}$ are continuous. In particular, if we choose $l = 2$, then we require $p > 3/2$.

Definition 3.1. We have the following spaces of Seiberg-Witten points

$$\mathcal{SW}_{l,p} = \mathcal{SW}_{h,\lambda,H,l,p} = \{(a, \phi) \in \mathcal{A}_{l,p}(Y) \times \Gamma_{l,p}(S) : \mathbf{sw}_{\lambda,H}((a, \phi)) = 0\}$$

and the following moduli spaces of Seiberg-Witten points

$$\mathcal{R}_{l,p} = \mathcal{R}_{h,\lambda,H,l,p} = \mathcal{SW}_{l,p} / \mathcal{G}_{l+1,p}(Y),$$

The irreducible part of e.g. $\mathcal{R}_{l,p}$ will be denoted by $\mathcal{R}_{l,p}^*$.

We choose to work with the configuration space $\mathcal{A}_{2,2}(Y) \times \Gamma_{2,2}(Y)$. The reason for this choice is that the L^2 spaces are more convenient to work with from the viewpoint of characterizing the cokernel of the relevant operators. But this is concerned with the situation on the product X rather than on Y . Henceforth, the subscript l stands for $(l, 2)$, e.g. $\|u\|_l = \|u\|_{l,2}$, $\mathcal{A}_l = \mathcal{A}_{l,2}$. We have the quotients $\mathcal{B}_2(Y)$ and $\mathcal{B}_2^0(Y)$ of the chosen configuration space $\mathcal{A}_2(Y) \times \Gamma_2(Y)$ under the group of gauges $\mathcal{G}_3(Y)$ and group of based gauges \mathcal{G}_3^0 . The gauge class of $(a, \phi) \in \mathcal{A}_2(Y) \times \Gamma_2(S)$ with respect to the full gauge group $\mathcal{G}_3(Y)$ will be denoted by $[a, \phi]$. Its gauge class with respect to based gauges will be denoted by $[a, \phi]_0$.

Local slices of the gauge actions are provided by the following lemma, which give rise to the Banach manifold structure for \mathcal{B}_2^* and \mathcal{B}_2^0 .

Lemma 3.2. (1) Consider an irreducible pair $(\bar{a}, \bar{\phi}) \in \mathcal{A}_l(Y) \times \Gamma_l(Y)$. If $u = (a, \phi) \in \mathcal{A}_l(Y) \times \Gamma_l(Y)$ is sufficiently close to $(\bar{a}, \bar{\phi})$ in $L^{l,2}$ -norm, then there is a unique gauge $g = e^f \in \mathcal{G}_{l+1}$ such that $\tilde{u} = (\tilde{a}, \tilde{\phi}) = g^*u$ satisfies the gauge fixing equation

$$(3.1) \quad d_Y^*(\tilde{a} - \bar{a}) + \text{Im} \langle \bar{\phi}, \tilde{\phi} \rangle = 0.$$

Moreover, the $L^{l+1,2}$ -norm of f can be estimated in terms of the $L^{l,2}$ -norm of $u - \bar{u}$.

(2) Given pairs $(\bar{a}, \bar{\phi}), (a, \phi) \in \mathcal{A}_l(Y) \times \Gamma_l(Y)$, there is a unique based gauge $g = e^f \in \mathcal{G}_{l+1}$ such that $\int_Y f = 0$ and $\tilde{u} = (\tilde{a}, \tilde{\phi}) = g^*u$ satisfies the gauge fixing equation

$$(3.2) \quad d_Y^*(\tilde{a} - \bar{a}) = 0.$$

Moreover, the $L^{l+1,2}$ -norm of f can be estimated in terms of the $L^{l,2}$ -norm of $a - \bar{a}$.

Proof. (1) The said gauge fixing equation amounts to

$$d_Y^* d_Y f + d_Y^*(a - \bar{a}) + \text{Im} \langle \bar{\phi}, e^{-f} \phi \rangle = 0.$$

The linearization of this equation with respect to f at $(\bar{a}, \bar{\phi})$ is the following equation

$$d_Y^* d_Y f + f \langle \bar{\phi}, \bar{\phi} \rangle = 0.$$

Since $\bar{\phi}$ is nonzero, integration shows that this equation has only the trivial solution. The desired results follow from elementary elliptic theory and the implicit function theorem.

(2) This part is a consequence of elementary theory of harmonic functions. \square

Lemma 3.3. (1) Each element in \mathcal{R}_2 or \mathcal{R}_2^0 can be represented by a smooth pair (a, ϕ) .

(2) If $\|\nabla H\|_{L^\infty} < C$ and $\|\nabla^2 H\|_{L^\infty} < C$ for a constant C , then \mathcal{R}_2 and \mathcal{R}_2^0 are compact.

Proof. We present the proof for (2), which contains the argument for (1). Let $(a, \phi) \in \mathcal{A}_2(Y) \times \Gamma_2(S)$ be a solution of (2.5). Applying the 3-dimensional Weitzenböck formula, the bound $\|\nabla H\|_{L^\infty} < C$ and Moser's weak maximum principle (cf. the proof of Lemma 8.6), we obtain $\|a\|_{L^\infty} \leq C$ for a constant C . (Here and in the

sequel, we use the same letter C to denote all constants which appear in a priori estimation). Since $H^1(Y, \mathbb{R}) = 0$, by Hodge decomposition, $a - a_0 = d_Y \gamma + d_Y^* \delta$ for some $\gamma \in \Omega_0^0(Y)$ and $\delta \in \Omega_0^2(Y)$. By gauge fixing, we can assume $d_Y^*(a - a_0) = 0$. Note that we can achieve this gauge fixing by a based gauge. Hence $a - a_0 = d_Y^* \delta$. Furthermore, we can assume that $d_Y \delta = 0$. Hence we have $\Delta \delta = F_a - F_{a_0}$. Since $\|F_a - F_{a_0}\| \leq C \|\phi\|^2 + C \leq C$, we have $\|\delta\|_2 \leq C$ by elliptic estimates. This implies $\|a\|_1 \leq C$. Applying this, the second equation of (2.5) and elliptic estimates, we deduce $\|\phi\|_1 \leq C$. Higher regularity and estimates follow from elliptic estimates and imply the desired compactness. \square

Convention 3.4. *Henceforth we drop the subscript 2 in SW_2 , \mathcal{R}_2 and \mathcal{R}_2^0 .*

Obviously, $\mathcal{R} = \mathcal{R}^0/S^1$, where S^1 is the group of constant gauges. We deal with \mathcal{R}^* and \mathcal{R}^0 separately.

The moduli space \mathcal{R}^*

For a given $(a, \phi) \in \mathcal{A}_2(Y) \times \Gamma_2(Y)$, let $G_Y = G_{Y,(a,\phi)} : \Omega_1^0(Y) \rightarrow \Omega_0^1(Y) \oplus \Gamma_0(S)$ be the infinitesimal gauge action operator at (a, ϕ) , i.e. $G_Y(f) = (d_Y f, -f\phi)$. Let $G_Y^* = G_{Y,(a,\phi)}^* : \Omega_2^1(Y) \oplus \Gamma_2(S) \rightarrow \Omega_1^0(Y)$ be the formal adjoint operator of G_Y w.r.t. the inner product (2.7). We have

$$G_Y^*(b, \psi) = d_Y^* b + \text{Im} \langle \phi, \psi \rangle.$$

There is a decomposition $\Omega_2^1(Y) \oplus \Gamma_2(S) = \ker G_Y^* \oplus \text{im} G_Y$. To be more precise, we write $\Omega_2^1(Y) \oplus \Gamma_2(S) = \ker_2 G_Y^* \oplus \text{im}_2 G_Y$. It follows that the tangent space $T_{[a,\phi]} \mathcal{B}_2^*(Y)$ of $\mathcal{B}_2^*(Y)$ at $[a, \phi]$ is represented by $\ker_2 G_{Y,(a,\phi)}^*$ (for any representative (a, ϕ) in $[a, \phi]$). Indeed, the latter gives rise to a vector bundle $\text{Ker}_2 G_Y^* \rightarrow \mathcal{A}_2 \times (\Gamma_2(S) - \{0\})$, whose quotient bundle $\text{Ker}_2 G_Y^*$ under the action of $\mathcal{G}_3(Y)$ can be identified with $T\mathcal{B}_2^*$. Next note that by the gauge invariance of the Chern-Simons functional and Lemma 2.7, we have

$$G_Y^* \mathbf{sw}_{\lambda,H}(a, \phi) = 0$$

for any $(a, \phi) \in \mathcal{A}_2(Y) \times \Gamma_2(Y)$. For this reason, and by the gauge equivariance, the operator $\mathbf{sw}_{\lambda,H}$ defines a section $[\mathbf{sw}_{\lambda,H}]$ of the quotient bundle $\ker_1 G_Y^*$, whose fiber at $[a, \phi]$ is represented by the kernel $\ker_1 G_Y^*$ of G_Y^* in $\Omega_1(Y) \oplus \Gamma_1(S)$. The moduli space \mathcal{R}^* is precisely the zero locus of this section.

In the sequel we omit the subscripts λ, H in the notation $\mathbf{sw}_{\lambda,H}$. Consider the operator $d\mathbf{sw}|_{(a,\phi)} : \Omega_2^1(Y) \oplus \Gamma_2(S) \rightarrow \Omega_1^1(Y) \oplus \Gamma_1(S)$, where $d\mathbf{sw}$ means the derivative, i.e. the tangent map of the operator \mathbf{sw} . By Lemma 2.6, it is formally self-adjoint. Assume $[a, \phi] \in \mathcal{R}_2$. Then the gauge invariance of the equation $\mathbf{sw} = 0$ implies $d\mathbf{sw} \circ G_Y = 0$. It follows that $G_Y^* \circ d\mathbf{sw} = 0$. Hence we obtain an operator $d\mathbf{sw}|_{\ker_2 G_{Y,(a,\phi)}^*} : \ker_2 G_{Y,(a,\phi)}^* \rightarrow \ker_1 G_{Y,(a,\phi)}^*$ (for any representative (a, ϕ) in $[a, \phi]$). It is easy to see that this operator is Fredholm of index zero and represents the linearization of the section $[\mathbf{sw}]$. Let it be denoted by $\mathcal{D} = \mathcal{D}_{a,\phi}$.

Lemma 2.7 implies that the operator \mathcal{D} coincides with the Hessian operator of the Chern-Simons functional with respect to the product (2.11). Note that it extends straightforwardly to reducible Seiberg-Witten points. On the other hand, \mathcal{D} is gauge equivariant, i.e.

This is a simple consequence of the gauge equivariance of the operator \mathbf{sw} . It follows that \mathcal{D} induces a section of Fredholm operators over Y , where the corresponding domain and target vector bundles are the quotient bundles induced from $\Omega_2^1(Y) \oplus \Gamma_2(S)$ and $\Omega_1^1(Y) \oplus \Gamma_1(S)$ respectively. This point of view provides a conceptual set-up. It will appear again in the 4-dimensional discussions in the sequel and be used more explicitly there.

Definition 3.5. *Let (a, ϕ) be a Seiberg-Witten point, i.e. a solution of the (perturbed) Seiberg-Witten equation (2.5). It is called non-degenerate if $\mathcal{D}_{a,\phi}$ is onto. The classes $[a, \phi]$ or $[a, \phi]_0$ are called nondegenerate if a representative is nondegenerate. (This is independent of the choice of the representative.)*

Lemma 3.6. *If all elements in \mathcal{R}^* are nondegenerate, then it is a naturally oriented smooth manifold of dimension zero. (The orientation means that every point in \mathcal{R}^* is assigned a sign.)*

Proof. By the above discussions, we only need to produce the natural orientation. We can use either the degree of the operator \mathbf{sw} or the spectral flow of the operator Q below as in [22]. (They give the same orientation.)

To analyse the operator \mathcal{D} , we introduce another closely related formally self-adjoint Fredholm operator Q . (The Fredholm property of \mathcal{D} is also a consequence of the Fredholm property of Q .) First notice the following deformation complex

$$(3.3) \quad 0 \longrightarrow \Omega^0 \xrightarrow{G_{a,\phi}} \Omega^1 \oplus \Gamma \xrightarrow{d\mathbf{sw}} \Omega^1 \oplus \Gamma \xrightarrow{G_{a,\phi}^*} \Omega^0 \longrightarrow 0$$

where the letters Y and S and the Sobolev subscripts are omitted in the notations. We define $Q = Q_{(a,\phi)} : (\Omega_2^1(Y) \oplus \Gamma_2(S)) \oplus \Omega_2^0(Y) \rightarrow (\Omega_1^1(Y) \oplus \Gamma_1(S)) \oplus \Omega_1^0(Y)$ by the following formula:

$$Q = \begin{pmatrix} d\mathbf{sw} & G \\ G^* & 0 \end{pmatrix}$$

Lemma 3.7. *Let $(a, \phi) \in \mathcal{SW}$. Then we have*

$$\ker Q \cong \begin{cases} \ker \mathcal{D} \oplus \mathbb{R}, & \text{if } (a, \phi) \text{ is reducible;} \\ \ker \mathcal{D}, & \text{if } (a, \phi) \text{ is irreducible.} \end{cases}$$

and

$$\text{coker } Q \cong \begin{cases} \text{coker } \mathcal{D} \oplus \mathbb{R}, & \text{if } (\phi, a) \text{ is reducible;} \\ \text{coker } \mathcal{D}, & \text{if } (\phi, a) \text{ is irreducible.} \end{cases}$$

We omit the simple proof.

The moduli space \mathcal{R}^0 .

The above treatment does not apply to the reducible elements of \mathcal{R} , because the tangent bundle of $\mathcal{B}_2^*(Y)$ does not extend smoothly across the reducibles. To analyse the structure of \mathcal{R}^0 around reducibles, one can use a quotient bundle formulation on the level of the based gauge quotient. But we choose a different approach which gives somewhat stronger results. Henceforth we make

Assumption 3.8. *Y is a rational homology sphere, i.e. its first Betti number is zero.*

Lemma 3.9. *There is a canonical diffeomorphism from $\Sigma \equiv (a_0 + \ker_2 d^*) \times \Gamma_2(S)$ to $\mathcal{B}_2^0(Y)$. In other words, the former space is a global slice of the action of the group $\mathcal{G}_3^0(Y)$ on the space $\mathcal{A}_2(Y) \times \Gamma_2(S)$.*

Proof. To show that the natural map from the former space to the latter is one to one, consider $b_1, b_2 \in \ker_2 d_Y^*$ and $\phi_1, \phi_2 \in \Gamma_2(S)$ such that $(a_0 + b_2, \phi_2) = g^*(a_0 + b_1, \phi_1)$ for some gauge $g \in \mathcal{G}_3^0$. Then $d_Y^*(g^{-1}d_Y g) = 0$ and $g(y_0) = 1$. Since Y is a rational homology sphere, we have $dg \equiv 0$, and hence $g \equiv 1$. The remaining part of the proof is obvious. \square

This lemma enables us to reduce the Seiberg-Witten operator to the said global slice. But the operator Q is no longer suitable for analysing the linearization of the Seiberg-Witten operator. Instead, we consider the following augmented Seiberg-Witten equation

$$(3.4) \quad \begin{cases} *_Y F_a + d_Y f + \langle e_i \cdot \phi, \phi \rangle e^i & = \nabla H(a), \\ \partial_a \phi + \lambda \phi + f \phi & = 0, \end{cases}$$

where $a \in \mathcal{A}_2(Y)$, $\phi \in \Gamma_2(S)$ and $f \in \Omega_2^0(Y)$.

Lemma 3.10. *Let (a, ϕ, f) be a solution of (3.2). Then (a, ϕ) satisfies the Seiberg-Witten equation (2.5) and f is a constant. Moreover, if (a, ϕ) is irreducible, then f must be zero.*

Proof. Applying d_Y^* to the first equation of (3.2), we deduce

$$d_Y^* d_Y f + f |\phi|^2 = 0.$$

The desired conclusion follows. \square

We denote the left hand side of (3.4) by $\mathbf{swa}(a, \phi, f)$. The linearization of the restriction of \mathbf{swa} to Σ will be denoted by \mathcal{D}_1 . One readily checks that it is a Fredholm operator of index 1.

Definition 3.11. Let (a, ϕ) be a Seiberg-Witten point. It is called *based-nondegenerate*, if \mathcal{D}_1 is onto at (a, ϕ, f) , where f is an arbitrary constant if $\phi = 0$ and zero if $\phi \neq 0$. It is easy to see that the based-nondegenerate property is invariant under gauge transformations. In particular, this definition makes sense for based gauge classes $[a, \phi]_0$.

Let the gauges act on f trivially. The moduli space of based gauge classes of solutions of the equation (3.2) will be denoted by \mathcal{R}_a^0 . An element in it is called based-nondegenerate, if its corresponding element in \mathcal{R}^0 is so. As an immediate consequence of the above discussions we obtain

Lemma 3.12. *If all elements of \mathcal{R}_a^0 are based-nondegenerate, then it is a smooth oriented manifold of dimension one. If moreover \mathcal{R}^0 is compact, then the irreducible part of \mathcal{R}_a^0 consists of finitely many disjoint circles and its reducible part consists of finitely many disjoint lines. Consequently, the irreducible part of \mathcal{R}^0 consists of finitely many disjoint circles and its reducible part consists of finitely many points.*

Next we give the definition of holonomy perturbations. We follow [10]. Let D denote the unit disk in \mathbb{C} . Consider a triple (ω, α, D) , where $\omega \in Y$, $\alpha \in T^*Y$ and

$I : D^2 \rightarrow Y$ is a smooth embedding such that $I(0) = y_o$ and $dI(T_0D)$ is transversal to v_0 . Fix a point $s_0 \in S^1$. Let $P(y_0, v_0, I)$ be the set of all smooth embeddings $\gamma : S^1 \times D \rightarrow Y$ such that $\gamma(s_0, \theta) = I(\theta)$ for $\theta \in D$ and $\frac{\partial \gamma}{\partial s}(s_0, 0) = v_0$. Here s_0 is a fixed point in S^1 . We set

$$P^{(m)} = \cup_{(y_o, v_0, I)} (P(y_0, v_0, I))^m,$$

for $m \in \mathbf{N}$. Now we define a map $\gamma^h : \mathcal{A}(Y) \times P^{(m)} \rightarrow C^\infty(D^2, U(1)^m)$ by

$$\gamma^h(a, (\gamma^1, \gamma^2, \dots, \gamma^m))(\theta) = (\gamma_\theta^1(a), \gamma_\theta^2(a), \dots, \gamma_\theta^m(a)),$$

where $\gamma_\theta^i : \mathcal{A}(Y) \rightarrow U(1)$ denotes the holonomy map along the loop $\gamma^i(\cdot, \theta)$ (at the base point y_0). It is easy to see that γ^h is gauge invariant. Next we choose a sequence $\{\epsilon_i\}$ of positive numbers as in [10] such that

$$C^\epsilon(U(1)^m, \mathbb{R}) = \{u \in C^\infty(U(1)^m, \mathbb{R}) : \|u\|_\epsilon < \infty\}$$

is complete. Here

$$\|u\|_\epsilon = \sum_{i=0}^{\infty} \epsilon_i \max_{U(1)^m} |\nabla^i u|.$$

Now we set

$$\Pi = \cup_{m \in \mathbf{N}} (P_m \times C^\epsilon(U(1)^m, \mathbb{R})).$$

This is the parameter space of holonomy perturbations. Choose a smooth function ξ with support in the interior of D . For each $\pi = (\gamma, u) \in \Pi$, we define the holonomy perturbation $H_\pi : \mathcal{A}(Y) \rightarrow \mathbb{R}$ by

$$H_\pi(a) = \int_{D^2} u(\gamma^h(a)) \xi(\theta) d^2\theta.$$

It is clear that H_π extends to $\mathcal{A}_1(Y)$.

Lemma 3.13. *For any $\pi = (\gamma, u) \in \Pi$, H_π is a smooth $\mathcal{G}_3(Y)$ -invariant function. Moreover, the L^2 -gradient ∇H_π satisfies*

$$\|\nabla H_\pi(a)\|_{L^\infty} \leq C,$$

with $C > 0$ independent of $a \in \mathcal{A}$. Similar bounds hold for the higher derivatives of H_π . The bounds can be made arbitrarily small by choosing u small.

Proof. For simplicity, we only consider $m = 1$. Set $H = H_\pi$. We can write $H(a) = \int_{D^2} u(\gamma_\theta(a)) \xi d^2\theta$. It follows that

$$dH(a)(b) = \int_{D^2} du|_{\gamma_\theta(a)} (\gamma'_\theta(a)b) \xi d^2\theta.$$

Elementary computations lead to

$$\gamma'_\theta(a)b = -\gamma_\theta(a) \int b.$$

We deduce

$$\begin{aligned}
 dH(a)(b) &= - \int_{D^2} \xi \langle \nabla u, \gamma_\theta(a) \rangle d^2\theta \int_{\gamma_\theta} b \\
 &= \int_{\gamma(S^1 \times D^2)} (\xi \circ \gamma^{-1}) \langle b, \overline{\langle \nabla u, \gamma_\theta(a) \rangle} (\gamma^{-1})^*(dt) \rangle \left| \frac{\partial \gamma}{\partial t} \right|^{-2} (\gamma^{-1})^*(dtd^2\theta), \\
 &= \int_Y f(\xi \circ \gamma^{-1}) \langle b, \overline{\langle \nabla u, \gamma_\theta(a) \rangle} (\gamma^{-1})^*(dt) \rangle
 \end{aligned}$$

where

$$f = \left| \frac{\partial \gamma}{\partial t} \right|^{-2} |(\gamma^{-1})^*(dtd^2\theta)/dvol|.$$

Consequently, $\nabla H(a) = f(\xi \circ \gamma^{-1}) \overline{\langle \nabla u, \gamma_\theta(a) \rangle} (\gamma^{-1})^*(dt)$. The desired estimate for ∇H follows. The higher order derivatives can easily be computed by using the above formula. \square

We make

Assumption 3.14. *Henceforth we choose H in (2.5) and (3.4) to be H_π .*

We remark in passing that for the purpose of achieving transversality for the moduli spaces \mathcal{R}^* and \mathcal{R}^0 it is not necessary to introduce the holonomy perturbations. However, they are important for achieving transversality for Seiberg-Witten trajectories as will be seen in the next section.

Lemma 3.15. *For perturbation $\pi \in \Pi$ such that $\nabla^2 H_\pi$ and $\nabla^3 H_\pi$ are small enough in L^∞ -norm (the set of such π is a nonempty open set), there exists a unique reducible element $[(a, 0)] \in \mathcal{R}$. Equivalently, there is a unique $a \in \mathcal{A}_2(Y)$ such that*

$$\begin{aligned}
 (3.5) \quad *_{\mathcal{Y}} F_a - \nabla H_\pi(a) &= 0, \\
 d_{\mathcal{Y}}^* a &= 0.
 \end{aligned}$$

Proof. Since Y is a rational homology sphere, the operator $*_{\mathcal{Y}} d_{\mathcal{Y}} : \ker d_{\mathcal{Y}}^* \rightarrow \ker d_{\mathcal{Y}}^*$ is a bounded isomorphism. Hence the existence follows from the implicit function theorem. To prove the uniqueness, consider connections a and a_1 satisfying (3.5). We set $b = a - a_1$ and deduce

$$*_{\mathcal{Y}} d_{\mathcal{Y}} b = \nabla H_\pi(a_1) - \nabla H_\pi(a) \text{ and } d_{\mathcal{Y}}^* b = 0.$$

By the implicit function theorem, for π with the property stated in the lemma, $b = 0$. \square

The unique solution of (3.5) will be denoted by $a_{(h, \pi)}$.

Lemma 3.16. *For $\pi \in \Pi$ satisfying the condition of Lemma 3.15, let $\sigma(\mathcal{J}_{a_{(h, \pi)}})$ be the set of eigenvalues of $\mathcal{J}_{a_{(h, \pi)}}$. Assume that $\nabla^2 H_\pi$ and $\nabla^3 H_\pi$ are small enough in L^∞ -norm (the set of such π is a nonempty open set). Then for $\lambda \in (\mathbb{R} - \sigma(\mathcal{J}_{a_{(h, \pi)}}))$, all elements of $\mathcal{R}(Y)$ are nondegenerate and all elements in $\mathcal{R}^0(Y)$ are based-nondegenerate.*

Proof. We only present the proof for the statement concerning the non-degeneracy. The based nondegeneracy can be treated in a similar way. Consider $(\sigma, \tau) \in [\sigma, \tau] \subset$

$\mathcal{R}(Y)$, we are going to show that \mathcal{D} at (a, ϕ) is onto. By gauge equivariance, we can choose $a = a_{(h, \pi)}$ for the reducible element. By Lemma 3.7, it suffices to analyse the operator Q . We have

$$Q(b, \psi, f) = \begin{pmatrix} *_Y db + 2\langle e_i \cdot \phi, \psi \rangle e^i + d_Y f - \nabla^2 H(a)b \\ \mathfrak{D}_a \psi + \lambda \psi + f\phi + b\phi \\ d_Y^* b + \text{Im} \langle \phi, \psi \rangle \end{pmatrix}.$$

Consider an element $(b_1, \psi_1, f_1) \in \Omega_1^1(Y) \oplus \Gamma_1(S) \oplus \Omega_1^0(Y)$ satisfying

$$(3.4) \quad \langle Q(b, \psi, f), (b_1, \psi_1, f_1) \rangle_{L^2} = 0$$

for all $(b, \psi, f) \in \Omega_2^1(Y) \oplus \Gamma_2(S) \oplus \Omega_2^0(Y)$. We first derive that (b_1, ψ_1, f_1) satisfies the adjoint equation $Q^* = 0$ (hence it satisfies $Q = 0$ because $Q^* = Q$) and is smooth.

Case 1 $\phi = 0$ and $a = a_{(h, \pi)}$.

We have $\mathfrak{D}_a \psi_1 + \lambda \psi_1 = 0$. By the choice of λ , we conclude that $\psi_1 \equiv 0$. Now (b_1, f_1) satisfies the following equation

$$(3.7) \quad \begin{aligned} *_Y d_Y b_1 + d_Y f_1 - \nabla^2 H(a)b_1 &= 0, \\ d_Y^* b_1 &= 0. \end{aligned}$$

Since Y is a rational homology sphere, the operator $(b_1, f_1) \rightarrow (*_Y d_Y b_1 + d_Y f_1, d_Y^* b_1)$ is an isomorphism from $\Omega_2^1(Y) \oplus (\Omega_2^0(Y))^0$ onto $\Omega_1^1(Y) \oplus (\Omega_1^0(Y))^0$, where the superscript 0 means the condition that the average be zero. As in the proof of Lemma 3.15, we deduce that if $\nabla^2 H$ and $\nabla^3 H$ are small enough, f_1 must be a constant and $b_1 = 0$. We conclude that $\text{coker } Q \cong \mathbb{R}$. By Lemma 3.7, this implies that \mathcal{D} is onto.

Case 2 $\phi \neq 0$.

By the unique continuation, the set $U = \{\phi \neq 0\}$ is an open dense set. For $y \in U$, $e_1 \cdot \phi(y), e_2 \cdot \phi(y), e_3 \cdot \phi(y)$ and $\phi(y)$ span S_y , where S_y denotes the fiber of S at $y \in Y$. We deduce that $\psi_1(y) = 0$ for $y \in U$, whence $\psi_1 \equiv 0$.

Now we easily see that (b_1, f_1) satisfies the equation (3.7). Hence $b_1 \equiv 0$ and the equation $Q(b_1, \psi_1, f_1) = 0$ reduces to $f_1 \phi = 0$. It follows that $f_1 = 0$ in U and consequently $f_1 \equiv 0$. We conclude that Q is onto. By Lemma 3.7, \mathcal{D} is onto. \square

As a consequence of the previous lemmas, we deduce

Proposition 3.17. *Let π and λ satisfy the same conditions as in Lemma 3.15 and Lemma 3.16. Then \mathcal{R} consists of finitely many signed points, the irreducible part of \mathcal{R}^0 consists of finitely many disjoint oriented circles, and its reducible part is a signed point.*

Definition 3.18. We call π and λ “ Y -generic”, provided that they satisfy the conditions of Lemma 3.15 and Lemma 3.16. The set of Y -generic parameters (π, λ) is a dense open set

4. SEIBERG-WITTEN TRAJECTORIES: TRANSVERSALITY

By Seiberg-Witten trajectories we mean solutions of the (perturbed) Seiberg-Witten trajectory equation (2.13) (we choose $H = H_\pi$). As stated in the introduction, our goal is to establish a Morse-Floer theory for the Chern-Simons functional on the quotient space $\mathcal{B}_2^0(Y) = (\mathcal{A}_2(Y) \times \Gamma_2(Y))/\mathcal{G}_3^0(Y)$. The union of the critical submanifolds of the Chern-Simons functional is precisely the moduli space \mathcal{R}^0 . The negative gradient flow lines of the Chern-Simons functional are given by the temporal form of the Seiberg-Witten trajectories, cf. Section 2. Setting $A = a(t) + f(\cdot, t)dt$ and $\phi(t) = \Phi(\cdot, t)$ in (2.10) we can rewrite it as follows

$$(4.1) \quad \begin{cases} \frac{\partial a}{\partial t} - *_Y F_a - d_Y f - \langle e_i \cdot \phi, \phi \rangle e^i & = \nabla H(a), \\ \frac{\partial \phi}{\partial t} + \not\partial_a \phi + \lambda \phi + f \phi & = 0. \end{cases}$$

(We omit the subscript π in H_π .)

We shall use various spaces of local $(l, 2)$ Sobolev class ($L_{loc}^{l,2}$ class), e.g. $\mathcal{A}_{l,loc} = \mathcal{A}_{l,2,loc}(X)$, $\Gamma_{l,2,loc}^\pm = \Gamma_{l,2,loc}(W^\pm)$, $\Omega_{l,loc}^k = \Omega_{l,2,loc}^k(X)$ and $\mathcal{G}_{l,loc}(X) = L_{loc}^{l,2}(X, S^1)$.

Definition 4.1. *We have the following spaces of Seiberg-Witten trajectories:*

$\mathcal{N} = \{(A, \Phi) \in \mathcal{A}_{2,loc} \times \Gamma_{2,loc}^+ : (A, \Phi) \text{ solves (4.1) and has finite (perturbed) Seiberg-Witten energy}\}$.

The corresponding moduli spaces are:

$\mathcal{M} = \mathcal{N}/\mathcal{G}_{3,loc}(X)$ and $\mathcal{M}^0 = \mathcal{N}/\mathcal{G}_{3,loc}^0(X)$, where $\mathcal{G}_{3,loc}^0(X) = \{g \in \mathcal{G}_{3,loc}(X) : g(y_0, 0) = 1\}$. (Recall that y_0 is a fixed reference point in Y .)

For sets B_1 and B_2 of Seiberg-Witten points we set

$$\mathcal{N}(B_1, B_2) = \{u \in \mathcal{N} : u(\cdot, t) \text{ converges pointwise smoothly to some } p \in B_1 \text{ as } t \rightarrow -\infty \text{ and to some } q \in B_2 \text{ as } t \rightarrow +\infty\}$$

and

$$\begin{aligned} \mathcal{N}^G(B_1, B_2) &= \{u \in \mathcal{N} : \text{there is a } g \in \mathcal{G}_{3,loc}(X) \text{ such that} \\ &g^*u(\cdot, t) \text{ converges pointwise smoothly to some } p \in B_1 \text{ as } t \rightarrow -\infty \\ &\text{and to some } q \in B_2 \text{ as } t \rightarrow +\infty\}. \end{aligned}$$

For a positive function ξ on X we consider the following ξ -weighted $(l, 2)$ -Sobolev norms

$$(4.3) \quad \|u\|_{l|\xi} = \left(\sum_{k \leq l} \int_X \xi^2 |\nabla^k u|^2 dy dt \right)^{\frac{1}{2}}.$$

For each pair of nonnegative numbers $\delta = (\delta_-, \delta_+)$ we choose a positive smooth function δ_F on \mathbb{R} such that $\delta_F(t) = \delta_\pm |t|$ near $\pm\infty$. We have the following δ -weighted $(l, 2)$ -Sobolev norms ($L_\delta^{l,2}$ -norms)

$$\|u\|_{l,\delta} = \|u\|_{l|e^{\delta_F}}.$$

If Q is a domain in $Y \times \mathbb{R}$, then $\|u\|_{l,\delta}$ means the $L_\delta^{l,2}$ norm of u on Q .

Here and in the sequel we adopt the following convention for Sobolev indices: in the context for X , the pair of indices l, δ refers to the above Sobolev space $L_\delta^{l,2}$, while l, loc refer to the local Sobolev space $L_{loc}^{l,2}$. On the other hand, in the context of Y , the pair of indices l, p refers to the Sobolev space $L^{l,p}$, with the index l alone standing for the pair $l, 2$.

Let $\Omega_{l,\delta}^k = \Omega_{l,\delta}^k(X)$, $\mathcal{A}_{l,\delta} = \mathcal{A}_{l,\delta}(X)$ and $\Gamma_{l,\delta}^\pm = \Gamma_{l,\delta}(W^\pm)$ denote the completion of the obvious spaces w.r.t. the $L_\delta^{l,2}$ -norm.

Proposition 4.2. *Assume that λ and π are Y -generic. Then there are positive constants δ_0 and E_0 depending only on h, λ and π with the following properties. For any temporal Seiberg-Witten trajectory $u = (A, \Phi) = (\phi, a)$ of local $(2, 2)$ -Sobolev class and finite energy, there exist a gauge $g \in \mathcal{G}_3(Y)$ and two smooth solutions u_-, u_+ of (2.6) such that g^*u is smooth and the following holds. For all l ,*

$$\|g^*u - u_+\|_{l,\delta_0;Y \times [T_1, \infty)} \leq C(l),$$

$$\|g^*u - u_-\|_{l,\delta_0;Y \times [-\infty, T_2]},$$

where $C(l)$ depends only on h, λ, π, l and an upper bound of the energy of u , and $T_1 > 0, T_2 < 0$ satisfy $E(u, Y \times [T_1, \infty)) \leq E_0, E(u, Y \times (-\infty, T_2]) \leq E_0$.

The proof will be presented in [28].

Corollary 4.3. *We have*

$$\mathcal{M} = \cup_{\alpha, \beta \in \mathcal{R}} \mathcal{M}(\alpha, \beta), \mathcal{M}^0 = \cup_{\alpha, \beta \in \mathcal{R}} \mathcal{M}^0(\alpha, \beta),$$

where $\mathcal{M}(\alpha, \beta) = \mathcal{N}^G(\alpha, \beta) / \mathcal{G}_{3,loc}(X)$ and $\mathcal{M}^0(\alpha, \beta) = \mathcal{N}^G(\alpha, \beta) / \mathcal{G}_{3,loc}^0(X)$.

We shall use the moduli spaces $\mathcal{M}^0(\alpha, \beta)$ (or rather suitable equivalent models for them) to construct the boundary operator in our Bott-type chain (cochain) complex.

Proposition 4.2 suggests that we can work in the set-up of trajectories with exponential asymptotics.

We introduce the relevant spaces. For $u \in \mathcal{A}_2(Y) \times \Gamma_2(S)$, let $G_u \subset \mathcal{G}(Y)$ denote its isotropy group of gauge actions. It is trivial if u is irreducible and S^1 if u is reducible. The isotropy groups are identical for gauge equivalent elements, hence $G_{[u]}$ is well-defined. Choose a reference connection $a_0 \in \mathcal{A}(Y)$ and set $u_0 = (a_0, 0) \in \mathcal{A}(Y) \times \Gamma(Y)$. For $p, q \in \mathcal{A}_2(Y) \times \Gamma_2(Y)$ we define the extension of p, q

$$(4.4) \quad Ext(p, q) = u_0 + \chi(p - u_0) + (1 - \chi)(q - u_0),$$

where χ is a cut-off function satisfying

$$(4.5) \quad \chi(t) = \begin{cases} 0, & \text{if } t \geq 0, \\ 1, & \text{if } t \leq -1. \end{cases}$$

Definition 4.4. For $p, q \in \mathcal{A}_l(Y) \times \Gamma_l(S)$ we introduce

$$L_{l,\delta}(p, q) = \{u \in \mathcal{A}_{l,loc} \times \Gamma_{l,loc}^+ : u - Ext(p, q) \in \Omega_{l,\delta}^1 \times \Gamma_{l,\delta}^+\}$$

$\mathcal{G}_{l+1,\delta}(p, q) = \{g \in \mathcal{G}_{l+1,loc}(X) : g - g_0 \in L_\delta^{l+1,2}(X, \mathbb{C}) \text{ for some } g_0 \in \mathcal{G}_{l+1,loc} \text{ which is t-independent and belongs to } G_p(G_q) \text{ near } \infty(-\infty)\}$,

$$\mathcal{G}_{l+1,\delta}^0(p, q) = \{g \in \mathcal{G}_{l+1,\delta}(p, q) : g(y_0, 0) = 1\},$$

$$\mathcal{G}_{l+1,\delta}^I = \{g \in \mathcal{G}_{l+1,loc}(X) : g - 1 \in L_\delta^{l+1,2}(X, \mathbb{C})\},$$

$$\mathcal{G}_{l+1,\delta}^{I,0} = \{g \in \mathcal{G}_{l+1,\delta}^I : g(y_0, 0) = 1\}.$$

Note that the second and third groups act freely. The first acts freely on the irreducible part of $L_{l,\delta}(p, q)$. We abbreviate e.g. $L_{2,\delta}(p, q)$ to $L_\delta(p, q)$, and e.g. $\mathcal{G}_{2+1,\delta}(p, q)$ to $\mathcal{G}_\delta(p, q)$. We have the quotients:

$$\mathcal{B}_\delta = \mathcal{B}_\delta(p, q) = L_\delta(p, q) / \mathcal{G}_\delta,$$

$$\mathcal{B}_\delta^0 = L_\delta / \mathcal{G}_\delta^0,$$

$$\mathcal{B}_\delta^I = L_\delta / \mathcal{G}_\delta^I, \text{ and}$$

$$\mathcal{B}_\delta^{I,0} = L_\delta / \mathcal{G}_\delta^{I,0}.$$

A suitable pair (A, Φ) induces the following elements (gauge classes) in these spaces respectively: $[(A, \Phi)]$, $[(A, \Phi)]_0$, $[(A, \Phi)]_I$ and $[(A, \Phi)]_{I,0}$.

Because $\mathcal{G}_3(Y)$ is connected, we obtain equivalent (in terms of suitable gauges) spaces for Seiberg-Witten points p', q' which are gauge equivalent to p, q respectively.

Definition 4.5. For $B_1, B_2 \subset \mathcal{A}_l(Y) \times \Gamma_l(Y)$, we set

$$L_{l,\delta}(B_1, B_2) = \cup_{p \in B_1, q \in B_2} L_{l,\delta}(p, q).$$

We abbreviate $L_{2,\delta}(B_1, B_2)$ to $L_\delta(B_1, B_2)$,

For example, we have $L_\delta(\alpha, \beta)$ for $\alpha, \beta \in \mathcal{B}_2(Y)$. This is a Hilbert manifold, indeed an affine Hilbert space bundle. We can represent it by the product $\alpha \times \beta \times L_\delta(p_0, q_0)$ for any chosen $p_0 \in \alpha, q_0 \in \beta$. Indeed, we can send $u \in L_\delta(p, q)$ to (p, q, \tilde{u}) with $\tilde{u} = u - \chi(p - p_0) - (1 - \chi)(q - q_0)$. This map defines the said Hilbert manifold structure.

The natural topology on $L_{l,\delta}(\mathcal{A}_l(Y) \times \Gamma_l(Y), \mathcal{A}_l(Y) \times \Gamma_l(Y))$ can be given in terms of the following distance.

Definition 4.6. For $u \in L_{l,\delta}(p_1, q_1), v \in L_{l,\delta}(p_2, q_2)$ with $p_1, p_2, q_1, q_2 \in \mathcal{A}_l(Y) \times \Gamma_l(Y)$, we set

$$d_{l,\delta}(u, v)^2 = \|p_1 - p_2\|_{l,2}^2 + \|q_1 - q_2\|_{l,2}^2 + \|(u - Ext(p_1, q_1)) - (v - Ext(p_2, q_2))\|_{l,\delta}^2.$$

Convergence with respect to this distance will be called ‘‘convergence in exponential Sobolev $(l, 2)$ -norm, or ‘‘exponential convergence in Sobolev $(l, 2)$ -norm. Smooth exponential convergence means exponential convergence in Sobolev $(l, 2)$ -norm for all l .

These distance and convergence concepts naturally descend to various quotient spaces. Indeed, we define for equivalence classes ω_1, ω_2

$$d_{l,\delta}(\omega_1, \omega_2) = \inf\{d_{l,\delta}(u, v) : u \in \omega_1, v \in \omega_2\}.$$

Definition 4.7. We set

$$\mathcal{G}_\delta^\infty = \{g \in \mathcal{G}_{3,loc}(X) : g - g_0 \in L_\delta^{3,2}(X, \mathbb{C}) \text{ for some } g_0 \in L_{loc}^{3,2} \text{ which is } t\text{-independent near } \pm\infty\},$$

$$\mathcal{G}_\delta^{\infty,0} = \{g \in \mathcal{G}_\delta^\infty : g(y_0, 0) = 1\}.$$

Definition 4.8. For $p, q \in \mathcal{SW}$, we set $\mathcal{N}_\delta(p, q) = L_\delta(p, q) \cap \mathcal{N}$. We have the moduli spaces of trajectories $\mathcal{M}_\delta(p, q) = \mathcal{N}_\delta(p, q) / \mathcal{G}_\delta(p, q)$, $\mathcal{M}_\delta^0(p, q) = \mathcal{N}_\delta(p, q) / \mathcal{G}_\delta^0(p, q)$ and $\mathcal{M}_\delta^I(p, q) = \mathcal{N}_\delta(p, q) / \mathcal{G}_\delta^I$.

Definition 4.9. For $B_1, B_2 \subset \mathcal{SW}$, we set

$$\mathcal{N}_\delta(B_1, B_2) = \cup_{p \in B_1, q \in B_2} \mathcal{N}_\delta(p, q).$$

For $\alpha, \beta \in \mathcal{R}$, We have the moduli spaces:

$$\mathcal{M}_\delta(\alpha, \beta) = \mathcal{N}_\delta(\alpha, \beta) / \mathcal{G}_\delta^\infty, \quad \mathcal{M}_\delta^0(\alpha, \beta) = \mathcal{N}_\delta(\alpha, \beta) / \mathcal{G}_\delta^{\infty,0}.$$

The irreducible part of e.g. \mathcal{M}_δ will be denoted by \mathcal{M}_δ^* . The spaces $\mathcal{M}_\delta(p, q)$ and $\mathcal{M}_\delta^0(p, q)$ are canonically isomorphic to $\mathcal{M}_\delta([p], [q])$ and $\mathcal{M}_\delta^0([p], [q])$ respectively. Moreover, we have the following easy lemma.

Lemma 4.10. *If (π, λ) is Y -generic, p, q are smooth representatives of α and β respectively, and δ_-, δ_+ are positive and do not exceed the exponent δ_0 in Proposition 4.3, then there are canonical isomorphisms*

$$\mathcal{M}_\delta(\alpha, \beta) \cong \mathcal{M}(\alpha, \beta), \quad \mathcal{M}_\delta^0(\alpha, \beta) \cong \mathcal{M}^0(\alpha, \beta).$$

We use these isomorphisms to topologize $\mathcal{M}(\alpha, \beta)$ and $\mathcal{M}^0(\alpha, \beta)$.

Thus, $\mathcal{M}(\alpha, \beta)$, $\mathcal{M}_\delta(\alpha, \beta)$ and $\mathcal{M}_\delta(p, q)$ can be viewed as three different models of the same space. The same holds for the spaces $\mathcal{M}^0(\alpha, \beta)$ etc.. To analyse the structures of these moduli spaces, we focus on the set-ups $\mathcal{B}_\delta^I(p, q)$ and $\mathcal{M}_\delta^I(p, q)$.

For $p, q \in \mathcal{A}_2(Y) \times \Gamma_2(S)$, consider the infinitesimal gauge action operator $G_X = G_{X,(A,\Phi)} : \Omega_{2,\delta}^0 \rightarrow \Omega_{1,\delta}^1 \oplus \Gamma_{1,\delta}^+$ at a given $(A, \Phi) \in L_\delta(p, q)$, $G_X(f) = (df, -f\Phi)$. Let G_X^* be the formal adjoint operator of G_X w.r.t. the following inner product

$$(4.6) \quad \langle (\Phi_1, A_1), (\Phi_2, A_2) \rangle_\delta = \int_X (Re\langle \Phi_1, \Phi_2 \rangle + \langle A_1, A_2 \rangle) e^{2\delta_F} dydt.$$

There holds

$$G_X^*(A', \Phi') = d_\delta^* A' + Im\langle \Phi, \Phi' \rangle,$$

where $d_\delta^* = e^{-2\delta_F} d^* e^{2\delta_F}$ is the formal adjoint of d with respect to (4.6). We have the following elementary lemma, which is analogous to Proposition 2a.1 in [10].

Lemma 4.11. *If δ_- and δ_+ are positive and small enough, then $\Omega_{2,\delta}^1 \oplus \Gamma_{2,\delta}^+ = im_2 G_X \oplus ker_2 G_X^*$. Consequently, the tangent space $T_{[A,\Phi]} \mathcal{B}_\delta(p, q)$ is represented by $ker G_X^*$.*

Proof. Choosing δ_- and δ_+ positive and small enough, we can make sure that $G_X^* G_X : \Omega_{2,\delta}^0 \rightarrow \Omega_{0,\delta}^0$ is Fredholm. Then the desired conclusions follow easily. \square

Now we assume that δ satisfies the condition of Lemma 4.10. Let $\mathcal{U}_\delta \rightarrow \mathcal{B}_\delta^I$ denote the quotient bundle of the trivial bundle $L(p, q) \times (\Omega_{2,\delta}^+ \oplus \Gamma_{2,\delta}^-) \rightarrow L(p, q)$

($\mathcal{G}_{3,\delta}$ acts on Ω^+ trivially and on Γ^- by $g^*\Psi = g^{-1}\Psi$). The (perturbed) Seiberg-Witten operator $\mathbf{SW} = \mathbf{SW}_{\lambda,H}$ (cf. Section 2) induces a section $[\mathbf{SW}]$ of this bundle. If $p, q \in \mathcal{SW}$, then its zero locus is precisely the moduli space $\mathcal{M}_\delta(p, q)$.

The linearization of the section $[\mathbf{SW}]$ is given by the restriction of the operator $d\mathbf{SW}|_{(A,\Phi)}$ to $\ker_2 G_X^*$, which will be denoted by $\mathcal{D}_X = \mathcal{D}_{X,(A,\Phi)}$. We introduce another closely related operator $\mathcal{F}_{p,q} : \Omega_{2,\delta}^1 \oplus \Gamma_{2,\delta}^+ \rightarrow \Omega_{1,\delta}^+ \oplus \Gamma_{1,\delta}^- \oplus \Omega_{1,\delta}^0$

$$\mathcal{F}_{p,q} = \begin{pmatrix} d\mathbf{SW} \\ -G_X^* \end{pmatrix}.$$

Lemma 4.12. *If $p, q \in \mathcal{SW}, (A, \Phi) \in \mathcal{N}_\delta(p, q)$, then $\ker \mathcal{D}_X = \ker \mathcal{F}_{p,q}$ and $\text{coker } \mathcal{D}_X = \text{coker } \mathcal{F}_{p,q}$. Consequently, \mathcal{D}_X is Fredholm iff $\mathcal{F}_{p,q}$ is Fredholm. If they are Fredholm, they have the same index.*

Proof. The kernel equality is clear. By the gauge invariance of the Seiberg-Witten equation, we have

$$(4.7) \quad d\mathbf{SW} \circ G_X = 0.$$

Applying this and Lemma 4.11, we derive $\text{im } \mathcal{F}_{p,q} = \text{im } \mathcal{D}_X \oplus G_X^*(\text{im}_2 G_X)$. But the second summand equals $\Omega_{1,\delta}^0$. Indeed, the cokernel of the operator $G_X^* G_X$ is precisely $\ker G_X$, which is trivial because the exponential weight δ implies vanishing at infinity. \square

Lemma 4.13. *Assume that δ_- and d_+ are small enough. If p (q) is reducible, we assume in addition that δ_- (δ_+) is positive. Then $\mathcal{F}_{p,q}$ is Fredholm for all $p, q \in \mathcal{SW}$.*

Proof. We first convert $\mathcal{F}_{p,q}$ into an equivalent form $\tilde{\mathcal{F}}_{p,q}$. Let $\Omega^{k,Y}$ denote the subspace of Ω^k consisting of forms which do not contain dt . Then $\Omega_{2,\delta}^+$ can be identified with $\Omega_{2,\delta}^{1,Y}$. On the other hand, we can identify $\Gamma_{2,\delta}^-$ with $\Gamma_{2,\delta}^+$ by the multiplication with $-\partial/\partial t$. Using these identifications we obtain for a given $(A, \Phi) = (a + fdt, \phi)$

$$\tilde{\mathcal{F}}_{p,q}(b + \tilde{f}dt, \psi) = \frac{\partial}{\partial t} \begin{pmatrix} b \\ \psi \\ \tilde{f} \end{pmatrix} - \begin{pmatrix} *_Y d_Y b + d_Y \tilde{f} + 2\text{Im}\langle e_i \phi, \psi \rangle e^i - \nabla^2 H(a) \cdot b \\ -\partial_a \psi - \lambda \psi - b\phi - f\psi - \tilde{f}\phi \\ d_Y^* b + 2\delta'_F \tilde{f} + \text{Im}\langle \phi, \psi \rangle \end{pmatrix}.$$

Hence $\tilde{\mathcal{F}}_{p,q} - (\frac{\partial}{\partial t} - Q - (0, 0, 2\delta'_F)) = (0, f, 0)$, where Q was defined in Section 3. Since f decays exponentially, its multiplication is a compact operator. Now the limits of the operator $Q + (0, 0, 2\delta'_F)$ at $\pm\infty$ are formally self-adjoint. Hence we can follow [10] or [21] to show that $\frac{d}{dt} - Q - 2\delta'_F$ is Fredholm. Consequently, $\mathcal{F}_{p,q}$ is Fredholm. \square

Consider a Y -generic pair (π_0, λ) . Choose a neighborhood Π_0 of π_0 such that $\Pi_0 \times \{\lambda\}$ consists of Y -generic pairs and the smallness conditions in Lemma 4.13 is uniform for all $\pi \in \Pi_0$ (with λ fixed).

Theorem 4.14. *Assume that δ_- and δ_+ are positive, satisfy the above smallness condition for all $\pi \in \Pi_0$ and do not exceed the constant δ_0 in Proposition 4.2. Then for generic $\pi \in \Pi_0$ the following holds. For all $p, q \in \mathcal{SW}$, $[\mathbf{SW}]$ is transverse*

to the zero section, and hence \mathcal{M}_δ^I is a smooth manifold of dimension $\text{ind } \mathcal{F}_{p,q}$. Consequently, \mathcal{M}_δ is a smooth manifold of dimension

$$\text{ind } \mathcal{F}_{p,q} - \max\{\dim G_p, \dim G_q\}$$

and \mathcal{M}_δ^0 is a smooth manifold of dimension

$$\text{ind } \mathcal{F}_{p,q} - \max\{\dim G_p, \dim G_q\} + 1.$$

Proof. First assume that at least one of p, q is irreducible. Then all elements of $L_\delta(p, q)$ are irreducible. We extend the bundle $\mathcal{U}_\delta \rightarrow \mathcal{B}_\delta^I \times \Pi_0$ in the trivial way. Then [SW] gives rise to a section of the extended bundle. By the Sard-Smale theorem, it suffices to show that this section is transversal to the zero section, which amounts to the surjectivity of the operator $\mathcal{D}_X \oplus d_\pi \mathbf{SW}$ at all (A, Φ) which solve the Seiberg-Witten equation with parameter $\pi \in \Pi_0$ (and λ). By Lemma 4.12, the latter is equivalent to the surjectivity of the operator $\mathcal{F}_{p,q} \oplus d_\pi \mathbf{SW}$, which in turn follows from the spinor part of the transversality argument in [15] (which is similar to the argument in the proof of Lemma 3.16) and Floer's transversality argument in [10] based on holonomy perturbations. This establishes the statement about the space \mathcal{M}_δ^I . The statements about \mathcal{M}_δ^0 and \mathcal{M}_δ follow via the involved group actions. For example, assume that p is reducible. We can choose a smooth imaginary valued function f_0 on \mathbb{R} which is supported in the positive half \mathbb{R} and equals $2\pi\sqrt{-1}$ near $+\infty$. Then the family of gauges $\exp(tf_0), 0 \leq t < 1$ represent the quotient $\mathcal{G}_\delta(p, q)/\mathcal{G}_\delta^I$. Dividing by these gauges reduces the dimension by one.

If both p and q are reducible, then they represent the same (unique) reducible element in \mathcal{R} . By Lemma 2.9, the energy of every Seiberg-Witten trajectory equals zero. By the gauge equivariance of the operator $\mathcal{F}_{p,q}$, we can use temporal gauges and assume that $p = q = (a_{(h,\pi)}, 0)$ and $(A, \Phi) \equiv p$. The asserted transversality follows from Lemma C.1 in Appendix C. Both \mathcal{M}_δ^I and \mathcal{M}_δ^0 are circles in this case. Note that although there is a family of gauges $\exp(tf_0)$ as above corresponding to each end (p or q), the dimension of \mathcal{M}_δ is one less than that of the dimension of \mathcal{M}_δ^I , rather than two less. This can be understood in the following way: suitable combinations of the two families are equivalent to constant gauges, which do not change the constant solution $(A, \Phi) \equiv p$.

□

Note that by gauge equivariance, the transversality property is independent of the choice of the representatives p, q in their gauge classes. Since \mathcal{R} consists of finitely many points, for generic π , the transversality property is shared by all p, q with $[p], [q] \in \mathcal{R}$.

Definition 4.15. We shall say that those π and the corresponding λ as described above are *generic*.

By gauge equivariance, $d\mathbf{SW}, G_X$ and G_X^* are equivariant as can easily be verified. More precisely, we have e.g.

$$(4.8) \quad d\mathbf{SW}_{g^*u} = (g^{-1})^* d\mathbf{SW} g^*.$$

By this and the homotopy invariance of Fredholm index, we also have

Lemma 4.16. *For given $\alpha, \beta \in \mathcal{R}$, $\text{ind } \mathcal{F}_{p,q}$ is independent of the choice of $p \in \alpha, q \in \beta$ and $(A, \Phi) \in L_\delta(p, q)$. It is also independent of the choice of (δ_-, δ_+) (satisfying the smallness condition in Theorem 4.14).*

Finally, we note an important consequence of transversality.

Lemma 4.17. *Let (π, λ) be generic, $p, q \in \mathcal{SW}$ and $u \in \mathcal{N}_\delta(p, q)$. Choose a reference $u_0 \in L_\delta(p, q)$. Then $d\mathbf{SW}_u$ has a right inverse $Q_u : \Omega_{1,\delta}^+ \oplus \Gamma_{1,\delta}^- \rightarrow \Omega_{2,\delta}^1 \oplus \Gamma_{2,\delta}^+$ with*

$$\|Q_u\| \leq C,$$

where C depends only on $\|u - u_0\|_{2,\delta}$ and (π, λ) . Q_u is equivariant under gauge actions. In particular, $\|Q_{g^*u}\| \leq \|g\|_{C^1} \|Q_u\|$ for $g \in \mathcal{G}_\delta^\infty \cap C^1(X, S^1)$.

Proof. Let \mathcal{O}_u denote the L_δ^2 -orthogonal complement of $\ker \mathcal{F}_{p,q}$, where $\mathcal{F}_{p,q} = \mathcal{F}_{p,q}|_{\mathcal{O}_u}$. Then $\mathcal{F}_{p,q}|_{\mathcal{O}_u}$ is a bounded isomorphism onto $\Omega_{1,\delta}^+ \oplus \Gamma_{1,\delta}^- \oplus \Omega_{1,\delta}^0$. Let \tilde{Q}_u denote its inverse. By elliptic and Fredholm estimates, we derive $\|\tilde{Q}_u\| \leq C$, where C depends only on $\|u - u_0\|_{2,\delta}$ and (π, λ) . We set $Q_u = \tilde{Q}_u|_{\Omega_{1,\delta}^+ \oplus \Gamma_{1,\delta}^-}$.

The gauge equivariance of Q_u follows from that of $d\mathbf{SW}$ and G_X . \square

5. INDEX AND ORIENTATION

Consider a Y -generic pair (π, λ) . Let $O = O_{\pi,\lambda}$ be the unique reducible element in \mathcal{R} . For $\alpha \in \mathcal{R}$ we define

$$\mu(\alpha) = \text{ind } \mathcal{F}_{p,q} - 1,$$

where $p \in \alpha, q \in O$.

Note that μ can easily be extended to all elements of $\mathcal{B}_2(Y)$. It depends on h, π and λ . Elementary computation shows $\mu(O) = 0$.

Lemma 5.1. *For $p, q, r \in \mathcal{SW}$ there holds*

$$\text{ind } \mathcal{F}_{p,r} = \text{ind } \mathcal{F}_{p,q} + \text{ind } \mathcal{F}_{q,r} - \dim G_q.$$

Proof. This is similar to the corresponding index addition formula in Floer's theory [10]. Floer's argument can be applied directly. Another argument is as follows. Composing with exponential weight multiplication operators, we can transform the operators to Sobolev spaces without weight. Then the addition formula is the consequence of a linear gluing argument. The term $\dim G_q$ arises because of the "jumping" across the kernel of the operator $d_Y^* + \text{Im}\langle \phi, \cdot \rangle$ which is caused by the operator $(0, 0, \delta'_F)$.

Corollary 5.2. *There holds*

$$\text{ind } \mathcal{F}_{p,q} = \mu([p]) - \mu([q]) + \dim G_q.$$

Consequently, if (π, λ) is a generic pair, then we have $\dim \mathcal{M}_\delta(p, q) = \mu([p]) - \mu([q]) - \dim G_p$, $\dim \mathcal{M}_\delta^0(p, q) = \mu([p]) - \mu([q]) + 1 - \dim G_p$.

Next we study the orientation of the moduli spaces of Seiberg-Witten trajectories

Proposition 5.3. *Assume that (π, λ) is generic. Then $\mathcal{M}_\delta^I(p, q)$, $\mathcal{M}_\delta(p, q)$ and $\mathcal{M}_\delta^0(p, q)$ are orientable. Indeed, their orientations are canonically determined after some choices are made, which will be given in the proof below. (Consequently, $\mathcal{M}_\delta([p], [q])$, $\mathcal{M}_\delta^0([p], [q])$, $\mathcal{M}([p], [q])$ and $\mathcal{M}^0([p], [q])$ are orientable.) Moreover, the orientations are consistent with the gluing construction used in the proof below, namely the orientation of $\mathcal{M}_\delta^I(p, q)$ is the same as the product orientation induced from gluing $\mathcal{M}_\delta^I(p, r)$ to $\mathcal{M}_\delta^I(r, q)$.*

Proof. Without loss of generality, assume that δ_+ and δ_- are equal and positive. The operator $\mathcal{F}_{p,q}$ defines a section of Fredholm operators over $L_\delta(p, q)$. Since it is gauge equivariant, i.e. $\mathcal{F}_{p,q}^{g^*u} = g^* \mathcal{F}_{p,q}^g (g^{-1})^*$, it induces a section of Fredholm operators over \mathcal{B}_δ^I , for which we use the same notation $\mathcal{F}_{p,q}$. Let $\det(p, q) = \det \mathcal{F}_{p,q}$ be its determinant line bundle.

Next we fix a smooth irreducible pair $p_0 = (a_0, \phi_0)$ and construct a new section of Fredholm operators $\tilde{\mathcal{F}}_{p_0, p_0}$ over $L_\delta(p_0, p_0)$ by interpolating \mathcal{F}_{p_0, p_0} along the time direction with the operator $\mathcal{F}_0 = \mathcal{F}_0^{(A, \Phi)} = (d^+ d_\delta^*, D_A)$, which is defined near time infinities. More precisely, for every $u = (A, \Phi)$, we set

$$\tilde{\mathcal{F}}_{p_0, p_0} u = \eta \mathcal{F}_{p,q}^u + (1 - \eta) \mathcal{F}_0^u,$$

where $\eta = \eta(t)$ is a nonnegative smooth function taking the value zero near infinities. By the arguments in the proof of Lemma 4.11 one readily deduces that $\tilde{\mathcal{F}}_{p_0, p_0}$ is Fredholm. On the other hand, it is gauge equivariant. It follows that $\tilde{\mathcal{F}}_{p_0, p_0}$ is Fredholm and gauge equivariant, and hence induces a section of Fredholm operators over $\mathcal{B}^I(p_0, p_0)$. Let its determinant line bundle be denoted by $\tilde{\det}(p_0, p_0)$.

By the uniform structure of $\tilde{\mathcal{F}}_{p_0, p_0}$ near time infinities, we can deform it through a continuous family of Fredholm operator sections to the standard section \mathcal{F}_0 , whose determinant line bundle is trivial. Hence $\tilde{\det}_0(p_0, p_0)$ is trivial. An orientation of the vector space $H_\delta^0 \oplus H_\delta^1 \oplus H_\delta^+$ (the homology of the complex associated with the operator $d^+ + d_\delta^*$) then determines an orientation of $\tilde{\det}_0(p_0, p_0)$, cf. [27]. We fix an orientation, i.e. a trivialization of $\tilde{\det}_0(p_0, p_0)$.

For each $p \in \mathcal{SW}$, we interpolate \mathcal{F}_{p, p_0} with $(d^+ + d_\delta^*, D_A)$ near plus and minus infinity to get $\tilde{\mathcal{F}}_{p, p_0}$ and $\tilde{\mathcal{F}}_{p_0, p}$ respectively.

For $p, q \in \mathcal{SW}$ we construct an embedding by a simple gluing process

$$L_\delta(p_0, p) \times L_\delta(p, q) \times L_\delta(q, p_0) \rightarrow L_\delta(p_0, p_0).$$

(Compare [13].) On the other hand, we choose reference elements $u_0 \in L_\delta(p_0, p)$, $u_1 \in L_\delta(p, q)$ and $u_2 \in L_\delta(q, p_0)$. Then it is easy to show that $u_0 + (\ker d_\delta^* \oplus \Gamma_{2, \delta}^+)$, $u_1 + (\ker d_\delta^* \oplus \Gamma_{2, \delta}^+)$ and $u_2 + (\ker d_\delta^* \times \Gamma_{2, \delta}^+)$ are global slices in $L_\delta(p_0, p)$, $L_\delta(p, q)$ and $L_\delta(q, p_0)$ for the action of the groups \mathcal{G}^I respectively. Using them and the above embedding we obtain an embedding Θ :

$$\mathcal{B}_\delta^I(p_0, p) \times \mathcal{B}_\delta^I(p, q) \times \mathcal{B}_\delta^I(q, p_0) \rightarrow \mathcal{B}_1(p_0, p_0).$$

We have the projections π_0 , π_1 and π_2 of the above product to its factors. In addition let π_p and π_q be its projections to p and q respectively. Now the index addition formula (Lemma 5.1) leads to an addition formula for the index bundle

which in turn implies a product formula for the determinant line bundle. We apply the last formula to the present situation to deduce

$$\pi_0^* \tilde{det}(p_0, p) \otimes \pi_1^* \det(p, q) \otimes \pi_2^* \tilde{det}(q, p_0) \otimes \pi_p^* l_p \otimes \pi_q^* l_q \cong \tilde{det}(p_0, p_0) |_{\text{im } \Theta},$$

where l_p (l_q) is the dual of the kernel of the operator $d^* + \langle \phi, \cdot \rangle$ at p (q).

The above isomorphism implies that $\tilde{det}(p_0, p)$, $\det(p, q)$ and $\tilde{det}(q, p_0)$ are trivial. Indeed, the above tensor product contains e.g. for $u_1 \in \mathcal{B}_\delta^I(p, q)$ and $u_2 \in \mathcal{B}_\delta^I(q, p_0)$

$$\pi_0^* \tilde{det}(p_0, p) |_{\mathcal{B}_\delta^I(p_0, p) \times \{u_1\} \times \{u_2\}} \otimes \det(p, q) |_{u_1} \otimes \tilde{det}(q, p_0) |_{u_2} \otimes l_p \otimes l_q,$$

whose triviality clearly implies the triviality of $\tilde{det}(p_0, p)$.

We choose an orientation for l_O (note that the l_p 's are canonically equivalent to each other for $p \in O$), and an orientation for each $\tilde{det}(p_0, p)$ and $\tilde{det}(q, p_0)$. Then the above isomorphism determines an orientation of $\det(p, q)$, which gives rise to an orientation of $\mathcal{M}_\delta^I(p, q)$. The desired consistency follows from the construction.

If both p and q are reducible, then \mathcal{M}_δ^0 is a circle generated by gauge actions, and hence inherits a canonical orientation from the actions. Otherwise, the moduli space \mathcal{M}_δ is the quotient of \mathcal{M}_δ^I by a free S^1 action if one of p, q is reducible, and equals \mathcal{M}_δ^I if neither is reducible. Hence the orientation of \mathcal{M}_δ^I induces an orientation of \mathcal{M}_δ . On the other hand, \mathcal{M}_δ is the quotient of \mathcal{M}_δ^0 by another free S^1 action, hence we arrive at an orientation of \mathcal{M}_δ^0 . \square

6. THE TEMPORAL MODEL AND COMPACTIFICATION

First we introduce

$$L_{T,\delta}(B_1, B_2) = \{u \in L_\delta(B_1, B_2) : u \text{ is temporal, i.e.}$$

$$u = (A, \Phi) = (a + f dt, \Phi) \text{ with } f \equiv 0\}.$$

$L_{T,\delta}(\alpha, \beta)$ is a Hilbert submanifold of $L_\delta(\alpha, \beta)$. The group $\mathcal{G}_3^0(Y)$ acts on it freely. We obtain the quotient topology on $L_{T,\delta}(\alpha, \beta)/\mathcal{G}_3^0(Y)$. Although there is a natural smooth structure on this quotient space, we shall not use it. Our moduli space $\mathcal{M}_T^0(S_\alpha, S_\alpha)$ below is a subspace of $L_{T,\delta}(\alpha, \beta)/\mathcal{G}_3^0(Y)$, but its smooth structure will be established via its equivalent model $\mathcal{M}^0(p, q)$, $p \in \alpha, q \in b$.

Next we consider temporal Seiberg-Witten trajectories. For $B_1, B_2 \subset \mathcal{SW}$ and positive number δ we set

$$\mathcal{N}_{T,\delta}(B_1, B_2) = \{u \in \mathcal{N}_\delta(B_1, B_2) : u \text{ is temporal}\}.$$

For $\alpha \in \mathcal{R}$ let S_α denote its lift to \mathcal{R}^0 . For $\alpha, \beta \in \mathcal{R}$ we set

$$\mathcal{M}_{T,\delta}(S_\alpha, S_\beta) = \mathcal{N}_{T,\delta}(S_\alpha, S_\beta)/\mathcal{G}_3(Y),$$

$$\mathcal{M}_{T,\delta}^0(S_\alpha, S_\beta) = \mathcal{N}_{T,\delta}(S_\alpha, S_\beta)/\mathcal{G}_3^0(Y).$$

The second space will play the major role in our constructions.

We shall only consider Y -generic parameters and fix a sufficiently small δ . Henceforth we shall omit the subscript δ in the various notations such as $\mathcal{M}^0(S_\alpha, S_\alpha)$.

Lemma 6.1. *Assume that (π, λ) is generic. Then $\mathcal{M}_T^0(S_\alpha, S_\beta)$ (the “temporal model”) are smooth manifolds which are canonically diffeomorphic to $\mathcal{M}^0(p, q)$ (the “fixed-end model”) for $p \in \alpha, q \in \beta$.*

Proof. The space $\mathcal{N}(p, q)$ is a smooth Hilbert fibration over $\mathcal{M}^0(p, q)$, and hence a Hilbert manifold. Consider the temporal transformation $T_G : L(\alpha, \beta) \rightarrow L_T(\alpha, \beta)$: for $u = (a + fdt, \Phi)$, set

$$g_T(u) = e^{-\int_0^t f}$$

and $T_G(u) = g_T(u)^*u$. The restriction of T_G to $\mathcal{N}(p, q)$ has image $\mathcal{N}_T(\alpha, \beta) \subset L_T(\alpha, \beta)$. The induced temporal transformation $T_G : \mathcal{M}^0(p, q) \rightarrow \mathcal{M}_T^0(S_\alpha, S_\beta) \subset L_T(\alpha, \beta)/\mathcal{G}_3^0(Y)$ is a homeomorphism. We use this natural homeomorphism to define the canonical smooth structure for $\mathcal{M}_T^0(S_\alpha, S_\beta)$. \square

Definition 6.2. For $u \in \mathcal{N}(p, q)$, let p', q' be the endpoints of $T_G(u)$, i.e. its limits at $-\infty$ and $+\infty$ respectively. We set $\pi_-(u) = [p']_0, \pi_+(u) = [q']_0$. π_+ and π_- are called the *temporal projections*. For $[u]_0^T \in \mathcal{M}_T^0(S_\alpha, S_\beta)$ we define the temporal projections as follows: $\pi_\mp([u]_0^T) = \pi_\mp(u)$. This is independent of the choice of u . We have a similar definition for $\mathcal{M}_\delta^0(p, q)$.

Lemma 6.3. *The temporal projections are invariant under the action of $\mathcal{G}_\delta^{\infty, 0}$.*

Proof. Consider $u_1, u_2 \in \mathcal{N}(p, q)$ such that $u_2 = g^*u_1$ with $g \in \mathcal{G}_\delta^{\infty, 0}$. Then $T_G(u_1) = \tilde{g}^*T_G(u_2)$ for some $\tilde{g} \in \mathcal{G}_\delta^{\infty, 0}$. Since both $T_G(u_1)$ and $T_G(u_2)$ are temporal, it follows that $\tilde{g} \in \mathcal{G}_3^0(Y)$. Hence we conclude that the temporal projections of u_1 are the same as those of u_2 .

We need to compactify our moduli spaces of trajectories. For this purpose, we introduce the following definitions.

Definition 6.4. Let $\mathcal{M}_T^0(S_{\alpha_0}, \dots, S_{\alpha_k})$ denote the fibered product

$$\mathcal{M}_T^0(S_{\alpha_0}, S_{\alpha_1}) \times_{S_{\alpha_1}} \mathcal{M}_T^0(S_{\alpha_1}, S_{\alpha_2}) \dots \times_{S_{\alpha_{k-1}}} \mathcal{M}_T^0(S_{\alpha_{k-1}}, S_{\alpha_k}),$$

i.e.

$$\begin{aligned} \mathcal{M}_T^0(S_{\alpha_0}, \dots, S_{\alpha_k}) &= \{([u_1]_0^T, \dots, [u_k]_0^T) \in \mathcal{M}_T^0(S_{\alpha_0}, S_{\alpha_1}) \times \dots \times \mathcal{M}_T^0(S_{\alpha_{k-1}}, S_{\alpha_k}) : \\ &\quad \pi_+([u_i]_0^T) = \pi_-([u_{i+1}]_0^T), i = 1, \dots, k-1\}. \end{aligned}$$

We call elements in $\mathcal{M}_T^0(S_{\alpha_i}, S_{\alpha_{i+1}})$ *temporal trajectory classes*, elements in

$$\mathcal{M}_T^0(S_{\alpha_0}, S_{\alpha_1}) \times \dots \times \mathcal{M}_T^0(S_{\alpha_{k-1}}, S_{\alpha_k})$$

multiple temporal trajectory classes, and elements in

$$\mathcal{M}_T^0(S_{\alpha_0}, \dots, S_{\alpha_k})$$

consistent multiple temporal trajectory classes. We call S_{α_i} the *i-th juncture manifold* of these consistent multiple temporal trajectory classes. We call $[u_i]_0^T$ the *i-th piece* of $([u_1]_0^T, \dots, [u_k]_0^T)$.

The temporal projections π_\pm are naturally extended to consistent multiple temporal trajectory classes. Indeed, we define π_- in terms of the first piece, and π_+ in terms of the last piece.

For distinct α and β , we define $\mathcal{M}_T^0(S_\alpha, S_\beta)_k$ to be the union of all $\mathcal{M}_T^0(S_{\alpha_0}, \dots, S_{\alpha_k})$ with distinct $\alpha_0, \dots, \alpha_k$ and $\alpha_0 = \alpha, \alpha_k = \beta$. Finally, we set for distinct α, β

$$\mathbf{M}_T^0(S_\alpha, S_\beta) = \cup_k \mathcal{M}_T^0(S_\alpha, S_\beta)_k.$$

By the definition, $\mathbf{M}_T^0(S_\alpha, S_\beta)$ has a natural stratified structure.

For $\alpha = \beta$, we set $\mathbf{M}_T^0(S_\alpha, S_\alpha) = \mathcal{M}_T^0(S_\alpha, S_\alpha)$, which consist of time-independent trajectories.

Definition 6.5. Let $p, q \in \mathcal{SW}$. A *piecewise trajectory* $u = (u_1, \dots, u_m)$ of length k from p to q with consecutive *junctures* $p_0 = p, \dots, p_k = q \in \mathcal{SW}$ is an element in $\mathcal{N}(p_0, p_1) \times \dots \times \mathcal{N}(p_{k-1}, p_k)$ with $u_i \in \mathcal{N}(p_{i-1}, p_i)$. u_i is called the *i-th piece* of u . p, q are called the *endpoints* of u at $+\infty$ and $-\infty$ respectively. u is called *proper*, if $E(u_i) > 0$ for every i . We also call a piecewise trajectory of length k a *k-trajectory*.

Definition 6.6. A piecewise trajectory $u = (u_1, \dots, u_k)$ is called *consistent*, provided that $\pi_+(u_i) = \pi_-(u_{i+1}), i = 1, \dots, k-1$.

For $p_0, \dots, p_k \in \mathcal{SW}$, let $\mathcal{N}(p_0, \dots, p_k)$ denote the space of consistent k -trajectories with junctures p_0, \dots, p_k . For $p, q \in \mathcal{SW}$ with $[p] \neq [q]$, let $\mathcal{N}(p, q)_k$ denote the space of proper and consistent k -trajectories from p to q . We set

$$\mathbf{N}(p, q) = \cup_k \mathcal{N}(p, q)_k.$$

The temporal projections π_+ and π_- naturally extend to consistent piecewise trajectories, namely π_- is defined in terms of the first portion and π_+ in terms of the last portion.

Definition 6.7. For $p_0, \dots, p_k \in \mathcal{SW}$, we set

$$\mathcal{M}^0(p_0, \dots, p_k) = \mathcal{N}(p_0, \dots, p_k) / (\mathcal{G}_\delta^0(p_0, p_1) \times \dots \times \mathcal{G}_\delta^0(p_{k-1}, p_k)).$$

For each $\alpha \in \mathcal{R}$, we choose an element $p_\alpha \in \alpha$. We fix this choice henceforth and denote the set of these elements by \mathcal{SW}_0 . For distinct $p, q \in \mathcal{SW}_0$, let $\mathcal{M}^0(p, q; \mathcal{SW}_0)_k$ denote the union of all $\mathcal{M}^0(p_0, \dots, p_k)$ with distinct $p_0 = p, \dots, p_k = q \in \mathcal{SW}_0$, and let $\mathbf{M}^0(p, q; \mathcal{SW}_0)$ denote the union of $\mathcal{M}^0(p, q; \mathcal{SW}_0)_k$ over all possible k . By the definition, $\mathbf{M}^0(p, q; \mathcal{SW}_0)$ has a natural stratification.

Proposition 6.8. *Assume that (π, λ) is generic. Consider $\alpha_i \in \mathcal{R}$ with $\mu(\alpha_i) \geq \mu(\alpha_{i+1}), i = 0, \dots, k-1$. Also consider $p_i \in \alpha_i, i = 0, \dots, k$. The moduli spaces $\mathcal{M}^0(p_0, \dots, p_k)$ and $\mathcal{M}_T^0(S_{\alpha_0}, \dots, S_{\alpha_k})$ are smooth manifolds of dimension*

$$\begin{aligned} & \sum_{0 \leq i \leq k-1} \dim \mathcal{M}^0(p_i, p_{i+1}) - \sum_{1 \leq i \leq k-1} \dim S_{\alpha_i} = \\ & \sum_{0 \leq i \leq k-1} \dim \mathcal{M}_T^0(S_{\alpha_i}, S_{\alpha_{i+1}}) - \sum_{1 \leq i \leq k-1} \dim S_{\alpha_i}. \end{aligned}$$

Moreover, $\mathcal{M}^0(p_0, \dots, p_k)$ is canonically diffeomorphic to $\mathcal{M}_T^0(S_{\alpha_0}, \dots, S_{\alpha_k})$.

Proof. Consider e.g. the case $\mathcal{M}_T^0(S_{\alpha_0}, S_{\alpha_1}, S_{\alpha_2})$. The general case is similar. We have the temporal projections $\pi_+ : \mathcal{M}_T^0(S_{\alpha_0}, S_{\alpha_1}) \rightarrow S_{\alpha_1}$ and $\pi_- : \mathcal{M}_T^0(S_{\alpha_1}, S_{\alpha_2}) \rightarrow S_{\alpha_1}$. They are smooth maps and transversal to each other because of the action of

the constant gauges. The moduli space $\mathcal{M}_T^0(S_{\alpha_0}, S_{\alpha_1}, S_{\alpha_2})$ is precisely the preimage of the digonal in $S_{\alpha_1} \times S_{\alpha_1}$ under the map $\pi_+ \times \pi_-$. Hence we infer that it is a smooth manifold. The dimension formula follows easily.

The proof for $\mathcal{M}^0(p_0, \dots, p_k)$ is similar. Here constant gauges are replaced by the family of gauges e^{tf_0} , $0 \leq t < 1$, where f_0 is a compactly supported imaginary valued function on \mathbb{R} with $f_0(y_0, 0) = 2\pi\sqrt{-1}$.

The claimed diffeomorphism is given by the temporal transformation. \square

Next we introduce suitable convergence concepts. First, we set for $p_i \in \mathcal{A}_l(Y) \times \Gamma_l(Y)$,

$$L_{l,\delta}(p_0, \dots, p_k) = L_{l,\delta}(p_0, p_1) \times L_{l,\delta}(p_1, p_2) \times \dots \times L_{l,\delta}(p_{k-1}, p_k).$$

Definition 6.9. Let τ_{t_0} denote the time translation operator with translation amount $-t_0$, i.e. $\tau_{t_0}(u)(y, t) = u(y, t - t_0)$.

Definition 6.10. We define the ‘‘suspension’’ or ‘‘pre-gluing’’ operator

$$\# : L_{l,\delta}(p_0, \dots, p_k) \times \mathbb{R}_+^{k-1} \rightarrow L_{l,\delta}(p_0, p_k),$$

$\mathbb{R}_+ = \{t \in \mathbb{R} : t > 0\}$, as follows. Let χ be the cut-off function introduced before. For $(u, \mathbf{r}) = ((u_1, \dots, u_k), (r_1, \dots, r_{k-1})) \in L(p_0, \dots, p_k) \times \mathbb{R}_+^{k-1}$, we set

$$\#(u, \mathbf{r}) = \begin{cases} u_1, & \text{if } t \leq r_1 - 1; \\ u_k, & \text{if } t \geq 2 \sum_{i \leq k-2} r_i + r_{k-1} + 1; \\ p_1 + \tau_{r_1}(\chi)(u_1 - p_1) + \tau_{r_1}(1 - \chi)\tau_{2r_1}(u_2 - p_1) + \\ & + 1; \\ \dots \\ p_{k-1} + \tau_{2r_1 + \dots + 2r_{k-2} + r_{k-1}}(\chi)\tau_{2r_1 + \dots + 2r_{k-2}}(u_{k-1} - p_{k-1}) + \\ & \tau_{2r_1 + \dots + r_{k-2} + r_{k-1}}(1 - \chi)\tau_{2r_1 + \dots + 2r_{k-1}}(u_k - p_{k-1}), & \text{if } 2 \sum_{j \leq k-2} r_j + \\ & r_{k-1} - 1 \leq t \leq 2 \sum_{j \leq k-2} r_j + r_{k-1} + 1. \end{cases}$$

We also denote $\#(u, \mathbf{r})$ by $u\#\mathbf{r}$.

Definition 6.10. For $\mathbf{r} = (r_1, \dots, r_{k-1}) \in \mathbb{R}_+^{k-1}$ we introduce the weight $w_{\mathbf{r}} \equiv \exp(w_{\mathbf{r}}^*)$,

$$w_{\mathbf{r}}^* = \tau_{r_1}(\chi)\delta_F + \tau_{r_1}(1 - \chi)\tau_{2r_1+r_2}(\chi)\tau_{2r_1}(\delta_F) + \dots + \\ \tau_{2r_1 + \dots + 2r_{k-2} + r_{k-1}}(1 - \chi)\tau_{2r_1 + \dots + 2r_{k-1}}(\delta_F).$$

The weighted Sobolev norms $\|\cdot\|_{l, w_{\mathbf{r}}}$ are obtained by replacing δ_F in $\|\cdot\|_{l,\delta}$ with $w_{\mathbf{r}}$. For $\bar{u} = (u, \mathbf{r})$, we set

$$w_{\bar{u}} = w_{\mathbf{r}}.$$

Definition 6.11. For $u \in L_{l,\delta}(p, q) \cup L_{loc}^{l+1,2}$ with $p, q \in \mathcal{A}_l(Y) \times \Gamma_l(Y)$ and $\mathbf{r} = (r_1, \dots, r_{k-1}) \in \mathbb{R}_+^{k-1}$ we define the \mathbf{r} -interpolation of u as follows

$$\text{Int}_{\mathbf{r}}(u) = u_0 + \chi(p - u_0) + \tau_{r_1}(\eta)(u(\cdot, r_1) - u_0) + \dots + \\ \tau_{2r_1 + \dots + 2r_{k-2} + r_{k-1}}(\eta)(u(\cdot, 2r_1 + \dots + 2r_{k-2} + r_{k-1}) - u_0) + \\ \tau_{2r_1 + \dots + 2r_{k-1}}(1 - \chi)(q - u_0),$$

where u_0 is the reference element introduced in Section 4 and η is a cut-off function satisfying

$$\eta(t) = \begin{cases} 1, & \text{if } -1 \leq t \leq 1; \\ 0, & \text{if } t \leq -2 \text{ or } t \geq 2. \end{cases}$$

Definition 6.12. For $\mathbf{r} = (r_1, \dots, r_{k-1}) \in \mathbb{R}_+^{k-1}$, $l \geq 0$ and $u \in L_{l,\delta}(p_1, q_2) \cap L_{loc}^{l+1,2}$, $v \in L_{l,\delta}(p_2, q_2) \cap L_{loc}^{l+1,2}$ with $p_1, p_2, q_1, q_2 \in \mathcal{A}_l(Y) \times \Gamma_l(Y)$, we introduce the (l, \mathbf{r}) -distance

$$d_{l,\mathbf{r}}(u, v)^2 = \|p_1 - p_2\|_{l,2}^2 + \|q_1 - q_2\|_{l,2}^2 + \sum_{1 \leq i \leq k-1} \|u(\cdot, r_i) - v(\cdot, r_i)\|_{l,2}^2 + \|(u - \text{Int}_{\mathbf{r}}(u)) - (v - \text{Int}_{\mathbf{r}}(v))\|_{l, w_{\mathbf{r}}}^2.$$

For gauge equivalence classes ω_1, ω_2 we set

$$d_{l,\mathbf{r}}(\omega_1, \omega_2) = \inf\{d_{l,\mathbf{r}}(u, v) : u \in \omega_1, v \in \omega_2\}.$$

Definition 6.13. We say that a sequence $u_j \in L_{l,\delta}(\mathcal{A}_l(Y) \times \Gamma_l(S), \mathcal{A}_l(Y) \times \Gamma_l(S)) \cap L_{loc}^{l+1,2}$ converges to $u = (u^1, \dots, u^k) \in L_{l,\delta}(p_0, \dots, p_k)$ in “piecewise exponential Sobolev $(l, 2)$ -norm”, or “piecewise exponentially in Sobolev $(l, 2)$ -norm”, provided that there is a sequence $\mathbf{r}_j = (r_{j,1}, \dots, r_{j,k-1}) \in \mathbb{R}_+^{k-1}$ such that $r_{j,i} \rightarrow \infty$ for each i and $d_{l,\mathbf{r}_j}(u_j, \#(u, \mathbf{r}_j)) \rightarrow 0$ as $j \rightarrow \infty$.

The essence of this convergence concept is this: roughly speaking, we can divide each u_j into k portions to produce k new sequences, such that each of them converges in exponential Sobolev $(l, 2)$ -norm. One important feature is the “equal distance property” of the exponential weight $w_{\mathbf{r}}$. Namely it consists of $k-1$ weights, such that the distance from the i -th “lowest position” r_i to the “center position” of the i -th weight is equal to its distance to the center position of the $(i+1)$ -th weight. (The picture is clear when one draws the graph of the weight $w_{\mathbf{r}}$.)

Smooth piecewise exponential convergence is defined to be piecewise exponential convergence with respect to all Sobolev norms.

Note that the one piece case $k = 1$ of piecewise exponential convergence is identical to the concept of exponential convergence given in Definition 4.6.

A consequence of the piecewise exponential convergence is the energy convergence, namely

$$E(u_j) \rightarrow E(u) \equiv \sum_i E(u^i).$$

Definition 6.14. The convergence concepts in Definition 6.13 are extended in an obvious way to convergence of elements in $L_{l,\delta}(p_0, \dots, p_k)$ (with a fixed k but varying p_i in general). Namely the convergence is defined to be that of each piece.

The piecewise exponential convergence of equivalence classes $[u^j]$ (in various contexts and set-ups) is defined in terms of the piecewise exponential convergence of \tilde{u}^j for some representatives $\tilde{u}^j \in [u^j]$. Equivalently, we can use $d_{l,\mathbf{r}}$ for equivalence classes.

Then the concept of piecewise exponential convergence is also naturally extended to multiple temporal trajectories classes. It is defined to be the piecewise exponential convergence of each piece.

The real line \mathbb{R} acts on trajectories in terms of the time translation. We define the time translation action on multiple temporal trajectories classes or piecewise trajectories to be the separate time translation action on individual pieces of the classes. It gives rise to an action of \mathbb{R}^k . Let the underline denote quotient under the time translation action in the *context of temporal trajectories*, e.g. $\underline{\mathcal{M}}_T^0(S_\alpha, S_\beta) = \mathcal{M}_T^0(S_\alpha, S_\beta) / \mathbb{R}^k$.

Theorem 6.15. *Assme that (π, λ) is Y -generic. Then the moduli spaces $\underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)$ are compact Hausdorff spaces, where the topology is given by smooth piecewise exponential convergence. (This topology is equivalent to that given by piecewise exponential convergence in the Sobolev $(2, 2)$ -norm.) Similarly, for $p, q \in \mathcal{SW}_0$, the quotient of $\mathbf{M}^0(p, q)$ under the time translation action are compact Hausdorff spaces.*

The proof of this theorem will be presented in Part II.

Consider $p \in \alpha, q \in \beta$ such that $p, q \in \mathcal{SW}_0$. The spaces $\mathbf{M}^0(p, q)$ (or $\mathcal{M}^0(p, q)$) and $\mathbf{M}_T^0(S_\alpha, S_\beta)$ (or $\mathcal{M}_T^0(S_\alpha, S_\beta)$) are isomorphic. But their quotients under the time translation action are not isomorphic. For our purpose, the time translation action on the temporal model is more suitable. For this reason, we consider the action of \mathbb{R} on $\mathcal{N}(p, q)$ induced from the time translation action on $\mathcal{N}_T(S_\alpha, S_\beta)$, which we call the “twisted time translation”.

Definition 6.16. The twisted time translation T_R by R is defined as follows. Let $u \in \mathcal{N}(p, q)$. Then $T_R u = (g_T(u)^{-1})^*(\tau_R(T_G(u)))$. (See the proof of Lemma 6.1 for g_T and T_G .) The twisted time translation acts on each piece of k -trajectories separately, giving rise to a \mathbb{R}^k action. We use the underline to denote the quotient under the twisted time translation action.

Lemma 6.17. *Assume that (π, λ) is Y -generic. Then the moduli spaces $\underline{\mathcal{M}}^0(p_0, \dots, p_k)$ and $\underline{\mathcal{M}}_T^0(S_{\alpha_0}, \dots, S_{\alpha_k})$ are canonically diffeomorphic smooth manifolds, where p_i, α_i are the same as in Lemma 6.8.*

Proposition 6.18. *Assume that (π, λ) is Y -generic. Then the moduli spaces $\underline{\mathbf{M}}^0(p, q; \mathcal{SW}_0)$ are compact Hausdorff spaces, where $p, q \in \mathcal{SW}_0, p \neq q$.*

The proof of this proposition will be presented in Part II.

Notice that the compact space $\underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)$ contains $\underline{\mathcal{M}}_T^0(S_\alpha, S_\beta)$ as a subspace. The compactification of $\underline{\mathcal{M}}_T^0(S_\alpha, S_\beta)$ is given by its closure $\bar{\underline{\mathcal{M}}}_T^0(S_\alpha, S_\beta)$.

Theorem 6.19. *For generic (π, λ) , we have $\bar{\underline{\mathcal{M}}}_T^0(S_\alpha, S_\beta) = \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)$. Moreover, the following hold:*

(1) $\underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)$ has the structure of d -dimensional smooth orientable manifolds with corners (i.e. modelled on the first quadrant of \mathbb{R}^d), where $d = \mu(\alpha) - \mu(\beta) - \dim G_p + 1$.

(2) This structure is compatible with the stratification $\underline{\mathbf{M}}_T^0(S_\alpha, S_\beta) = \cup_k \underline{\mathcal{M}}_T^0(S_\alpha, S_\beta)_k$, i.e. the interior of the k -dimensional face of $\underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)$ is precisely $\underline{\mathcal{M}}_T^0(S_\alpha, S_\beta)_k$.

(3) The temporal projections $\pi_- : \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta) \rightarrow S_\alpha$ and $\pi_+ : \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta) \rightarrow S_\beta$ are smooth fibrations.

Consequently, we have

$$(6.1) \quad \begin{aligned} \partial \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta) &= \cup_{\mu(\alpha) > \mu(\gamma) > \mu(\beta)} \underline{\mathbf{M}}_T^0(S_\alpha, S_\gamma, S_\beta) \\ &\equiv \cup_{\mu(\alpha) > \mu(\gamma) > \mu(\beta)} \underline{\mathbf{M}}_T^0(S_\alpha, S_\gamma) \times_{S_\gamma} \underline{\mathbf{M}}_T^0(S_\gamma, S_\beta), \end{aligned}$$

where the fiber product space $\underline{\mathbf{M}}_T^0(\alpha, S_\gamma) \times_{S_\gamma} \underline{\mathbf{M}}_T^0(S_\gamma, S_\beta)$ is defined to be

$$\{(\gamma, \alpha) \in \mathbf{M}^0(S_\alpha, S_\gamma) \times \mathbf{M}^0(S_\gamma, S_\beta) : \tau_-(\gamma) = \tau_+(\alpha)\}$$

(Of course, only the nonempty moduli spaces appear in (6.1).)

The situation with the isomorphic model $\mathcal{M}^0(p, q; \mathcal{SW}_0)$ is similar. Namely, the compact space $\underline{\mathbf{M}}^0(p, q; \mathcal{SW}_0)$ contains the subspace $\mathcal{M}^0(p, q; \mathcal{SW}_0)$, whose closure we denote by $\overline{\underline{\mathbf{M}}}^0(p, q; \mathcal{SW}_0)$. Then we have the following equivalent formulation for Theorem 6.19. (We only give part of the statements.)

Theorem 6.20. *In a generic situation, we have $\overline{\underline{\mathbf{M}}}^0(p, q; \mathcal{SW}_0) = \underline{\mathbf{M}}^0(p, q; \mathcal{SW}_0)$. Moreover, $\underline{\mathbf{M}}^0(p, q; \mathcal{SW}_0)$ has the structure of d -dimensional manifolds with corners which is compatible with its natural stratification.*

The said equivalence means the following:

Proposition 6.21. *In a generic situation, $\underline{\mathbf{M}}^0(p, q)$ is canonically diffeomorphic to $\underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)$.*

This proposition is an easy consequence of the temporal transformation. Clearly, Theorem 6.20 is a consequence of Theorem 6.19. The proof of Theorem 6.19 will be presented in Part II.

The two different models- the *temporal model* $\underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)$ and the *fixed-end model* $\underline{\mathbf{M}}^0(p, q; \mathcal{SW}_0)$ - provide different aspects of our basic set-up. It is good to have both of them for the purpose of conceptual understanding. Since the former is canonical (while the latter involves the choice of \mathcal{SW}_0), we shall use it in the formulation of our main constructions in the sequel. Of course, we can use equally well the latter.

7. BOTT-TYPE HOMOLOGY

We first introduce a few orientation conventions. We follow those used in [12]. For an oriented smooth manifold with corners \mathcal{X} , its boundary is oriented in such a way that $\text{span}\{n_{\partial\mathcal{X}}\} \oplus T\partial\mathcal{X} = TX|_{\partial\mathcal{X}}$ as oriented vector bundles (away from the corners of $\partial\mathcal{X}$), where $n_{\partial\mathcal{X}}$ is an inward normal field of the boundary. Given transversal smooth maps $F_1 : \mathcal{X}_1 \rightarrow S$ and $F_2 : \mathcal{X}_2 \rightarrow S$ from two oriented smooth manifolds with corners into an oriented smooth manifold S , the fibered product $\mathcal{X}_1 \times_S \mathcal{X}_2 = (F_1 \times F_2)^{-1}(\text{Diag}(S \times S))$ (*Diag* means *diagonal*) has a canonical orientation such that $T(\mathcal{X}_1 \times_S \mathcal{X}_2) \oplus N = (-1)^{\dim S \cdot \dim \mathcal{X}_2} TX_1 \oplus TX_2$ as oriented bundles, where N denotes the oriented bundle $(d(F_1 \times F_2))^{-1}((TS \oplus \{0\})|_{\text{Diag}(S \times S)})$. The following lemma can be found in [12] and is easy to verify.

Lemma 7.1. *There holds for oriented boundaries*

$$(7.1) \quad \partial(\mathcal{X}_1 \times_S \mathcal{X}_2) = \partial\mathcal{X}_1 \times_S \mathcal{X}_2 + (-1)^{\dim \mathcal{X}_1 + \dim S} \mathcal{X}_1 \times_S \partial\mathcal{X}_2,$$

where the summation sign means taking union. We also have

$$(7.2) \quad (X_1 \times_S \mathcal{X}_2) \times_{S'} \mathcal{X}_3 = X_1 \times_S (\mathcal{X}_2 \times_{S'} \mathcal{X}_3).$$

We shall use the natural orientation of S_α induced from the S^1 action. (We can also use any other orientations.) By Theorem 6.13, the boundary of $\underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)$ is a union of fibered products. We need to arrange the orientation of these spaces so that a suitable consistency holds in regard of the natural orientation of fibered products as defined above and boundary orientations. Indeed, we have

Lemma 7.2. *We can choose the orientation of $\underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)$ such that*

(7.3)

$$\partial \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta) = (-1)^{\mu(\alpha) + \dim S_\alpha} \Sigma_{\mu(\alpha) > \mu(\gamma) > \mu(\beta)} \underline{\mathbf{M}}_T^0(S_\alpha, S_\gamma) \times_{S_\gamma} \underline{\mathbf{M}}_T^0(S_\gamma, S_\beta)$$

as oriented manifolds with corners.

This lemma is analogous to Sublemma 1.20 in [12] and can be proven by the same arguments as there.

For a topological space \mathcal{X} and an abelian group \mathbf{G} , let $Q_j(\mathcal{X}; \mathbf{G}) = Q_j(\mathcal{X}; \mathbb{Z}) \otimes \mathbf{G}$ denote the abelian group generated by generalized singular j -cubes in \mathcal{X} with coefficients in \mathbf{G} . Here we define a generalized singular j -cube to be a continuous map $f : \Delta \rightarrow \mathcal{X}$, where Δ is an oriented j -cube, namely an oriented smooth manifold with corners which is diffeomorphic to the euclidean j -cube $[0, 1]^j$. (A continuous map $f : [0, 1]^j \rightarrow \mathcal{X}$ is called a singular j -cube, see [19].) Let $D_j(\mathcal{X}; \mathbf{G})$ be the subgroup of $Q_j(\mathcal{X}; \mathbf{G})$ generated by degenerate generalized singular j -cubes. Here, a generalized singular j -cube $f : \Delta \rightarrow \mathcal{X}$ is called degenerate, if $f \circ F$ is independent of some variable t_i , where F is a diffeomorphism from $[0, 1]^j$ to Δ .

Next consider a j -dimensional compact oriented manifold with corners Σ and a continuous map $f : \Sigma \rightarrow \mathcal{X}$. We divide Σ into oriented cubes and produce this way a chain $\sigma \in Q_j(\mathcal{X}; \mathbb{Z})$. Choosing a different way of dividing, we obtain another chain σ' . Let $Div_j(\mathcal{X}; \mathbf{G})$ be the subgroup of $Q_j(\mathcal{X}; \mathbf{G})$ generated by elements of the form $(\sigma - \sigma') \otimes g, g \in \mathbf{G}$.

Let $C_j(\mathcal{X}; \mathbf{G})$ denote the quotient group $Q_j(\mathcal{X}; \mathbf{G}) / (D_j(\mathcal{X}; \mathbf{G}) + Div_j(\mathcal{X}; \mathbf{G}))$. This is the group of generalized cubical singular j -chains in \mathcal{X} (with coefficients in \mathbf{G}). The corresponding cochain group is $C^j(\mathcal{X}; \mathbf{G}) = Hom(C_j(\mathcal{X}; \mathbb{Z}), \mathbf{G})$.

The ordinary boundary operator ∂_O for singular j -cubes extend to generalized singular j -cubes canonically via orientation preserving diffeomorphisms between j -cubes and $[0, 1]^j$. It then extends to $Q_j(\mathcal{X}; \mathbf{G})$ by linearity, and to $C_j(\mathcal{X}; \mathbf{G})$ by descending. We have the corresponding coboundary operator ∂_O^* . It is easy to show that the homology $H_*(C_*(\mathcal{X}; \mathbf{G}), \partial_O)$ and cohomology $H^*(C^*(\mathcal{X}; \mathbf{G}), \partial_O^*)$ are canonically isomorphic to the singular homology and cohomology of \mathcal{X} with coefficient group \mathbf{G} .

Remark 7.3. For $\sigma = ([0, 1]^j, f) \in C_j(\mathcal{X}; \mathbb{Z})$, the map $(\partial[0, 1]^j, f|_{\partial[0, 1]^j})$ from the oriented boundary induces a chain in a natural way. By our convention for boundary orientation, this chain equals $-\partial_O \sigma$.

Now we fix a coefficient group \mathbf{G} and abbreviate e.g. $C_j(\mathcal{X}; \mathbf{G})$ to $C_j(\mathcal{X})$. We set

$$(7.4) \quad S_i = \cup \{S_\alpha : \alpha \in \mathcal{R}, \mu(\alpha) = i\}$$

and introduce our Bott complexes C_*^{Bott} and C_{Bott}^* ,

$$C_k^{Bott} = \oplus_{i+j=k} C_j(S_i), C_{Bott}^k = \oplus_{i+j=k} C^j(S_i).$$

Note $C_*^{Bott} = C_*(\mathcal{R}^0)$, $C_{Bott}^* = C^*(\mathcal{R}^0)$.

Next we define a boundary operator $\partial_{Bott} : C_k^{Bott} \rightarrow C_{k-1}^{Bott}$ along with a coboundary operator $\partial_{Bott}^* : C_{Bott}^{k-1} \rightarrow C_{Bott}^k$ for each k . First, for each pair $\alpha, \beta \in \mathcal{R}$ with

$\mu(\alpha) > \mu(\beta)$, we define a boundary operator $\partial_{\alpha,\beta} : C_k^{Bott} \rightarrow C_{k-1}^{Bott}$. If the moduli space $\underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)$ is empty, we define $\partial_{\alpha,\beta}$ to be the zero operator. If it is nonempty, we define $\partial_{\alpha,\beta}$ as follows. For $\sigma \notin C_*(S_\alpha)$, we set $\partial_{\alpha,\beta}\sigma = 0$. For $\sigma = [(\Delta, f)] \in C_j(S_\alpha)$ with $\mu(\alpha) + j = k$, consider the fibered product

$$\tilde{\Delta} = \Delta \times_{S_\alpha} \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta) = \{(z, u) \in \Delta \times \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta) : f(z) = \pi_-(u)\}.$$

We have a natural projection map $\pi_+ : \tilde{\Delta} \rightarrow S_\beta$, $\pi_+((z, u)) = \pi_+(u)$. Since the projection π_- is a submersion, $\tilde{\Delta}$ is a compact oriented manifold with corners of dimension $j + \mu(\alpha) - \mu(\beta) - 1$. We divide $\tilde{\Delta}$ into oriented cubes. Using the projection map π_+ we then obtain from the divided $\tilde{\Delta}$ an element $\tilde{\Delta}_f$ in $Q_{j+\mu(\alpha)-\mu(\beta)-1}(S_\beta)$. We define

$$\partial_{\alpha,\beta}(\sigma) = [\tilde{\Delta}_f].$$

It is easy to see that this definition is independent of the choice of (Δ, f) and the way of dividing $\tilde{\Delta}$.

Clearly, we indeed have

$$\partial_{\alpha,\beta} : C_k^{Bott} \rightarrow C_{k-1}^{Bott}$$

for all k .

Definition 7.4. We define $\partial_{Bott} : C_k^{Bott} \rightarrow C_{k-1}^{Bott}$ as follows:

$$\partial_{Bott} = \partial_0 + \sum_{\mu(\alpha) > \mu(\beta)} \partial_{\alpha,\beta},$$

where $\partial_0 = (-1)^k \partial_O$. The boundary operator $\partial_{Bott} : C_*^{Bott} \rightarrow C_*^{Bott}$ is defined to be the direct sum of these boundary operators.

The boundary operator ∂_{Bott} induces a coboundary operator $\partial_{Bott}^* : C_{Bott}^k \rightarrow C_{Bott}^{k+1}$. Indeed, we have $\partial_{Bott}^* = (-1)^k \partial_O^* + \sum \partial_{\alpha,\beta}^*$.

Lemma 7.5. *We have $\partial_{Bott}^2 = 0$, $(\partial_{Bott}^*)^2 = 0$. Hence $(C_*^{Bott}, \partial_{Bott})$ is a chain complex and $(C_{Bott}^*, \partial_{Bott}^*)$ is a cochain complex.*

Proof. We handle the case $\mathbf{G} = \mathbb{Z}$ and ∂_{Bott} . The arguments easily extend to ∂_{Bott}^* and general coefficient groups. Consider $\sigma = [(\Delta, f)] \in C_j(S_\alpha)$ with $j + \mu(\alpha) = k$. We have $\partial_{Bott}^2 \sigma = \sum_{\mu(\alpha) < \mu(\beta)} I_\beta$, where

$$I_\beta = \partial_0 \partial_{\alpha,\beta} \sigma + \partial_{\alpha,\beta} \partial_0 \sigma + \sum_{\mu(\alpha) > \mu(\gamma) > \mu(\beta)} \partial_{\gamma,\beta} \partial_{\alpha,\gamma} \sigma.$$

On the other hand, by (7.1) we have

$$\begin{aligned} (7.5) \quad \partial(\Delta \times_{S_\alpha} \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta)) &= \partial \Delta \times_{S_\alpha} \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta) + (-1)^{j+\dim S_\alpha} \Delta_j \times_{S_\alpha} \partial \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta) \\ &= \partial \Delta \times_{S_\alpha} \underline{\mathbf{M}}_T^0(S_\alpha, S_\beta) + \\ &\quad (-1)^{j+\mu(\alpha)} \sum_{\mu(\alpha) > \mu(\gamma) > \mu(\beta)} \Delta \times_{S_\alpha} \underline{\mathbf{M}}_T^0(S_\alpha, S_\gamma) \times_{S_\gamma} \underline{\mathbf{M}}_T^0(S_\gamma, S_\beta). \end{aligned}$$

Multiplying this equation by $(-1)^{j+\mu(\alpha)-1} = (-1)^{k-1}$ and applying Definition 7.4 we then deduce

$$(7.6) \quad \partial_0 \partial_{\alpha,\beta} \sigma = -\partial_{\alpha,\beta} \partial_0 \sigma - \sum_{\mu(\alpha) > \mu(\gamma) > \mu(\beta)} \partial_{\alpha,\gamma} \partial_{\gamma,\beta} \sigma.$$

This shows that $I_\beta = 0$. \square

Definition 7.6. We define the Bott-type Seiberg-Witten Floer homology

$$FH_{Bott; \mathbf{G}}^{SW}(c)_*$$

to be the homology $H_*(C_*^{Bott}, \partial_{Bott})$ of the chain complex $(C_*^{Bott}, \partial_{Bott})$. (Recall that c is the given $spin^c$ structure.) We define the Bott-type Seiberg-Witten Floer cohomology $FH_{Bott; \mathbf{G}}^{SW}(c)^*$ to be the cohomology $H^*(C_{Bott}^*, \partial_{Bott}^*)$ of the cochain complex $(C_{Bott}^*, \partial_{Bott}^*)$.

Next we introduce a natural filtration -the index filtration

$$\mathcal{F}_*^{Bott} = \dots \subset \mathcal{F}_k^{Bott} \subset \mathcal{F}_{k+1}^{Bott} \subset \dots$$

on our chain complex $(C_*^{Bott}, \partial_{Bott})$. We put

$$\mathcal{F}_k^{Bott} = \bigoplus_{m \leq k} C_*(S_m).$$

It is clear that this defines a filtration by subcomplexes. We also have a dual filtration $\mathcal{F}_{Bott}^* = \dots \mathcal{F}_{Bott}^k \subset \mathcal{F}_{Bott}^{k-1} \subset \dots$ for the cochain complex,

$$\mathcal{F}_{Bott}^k = \bigoplus_{m \geq k} C_m^*.$$

Theorem 7.7. *The index filtration induces a spectral sequence $E(Bott)_{**}^*$ which converges to FH_{Bott}^{SW} and satisfies*

$$(7.7) \quad E(Bott)_{ij}^1 \cong H_j(S_i, \mathbf{G}).$$

*The dual filtration induces a spectral sequence $E(Bott, dual)_{**}^*$ which converges to FH_{Bott}^{SW*} and satisfies*

$$(7.8) \quad E(Bott, dual)_{ij}^1 \cong H^j(S_i; \mathbf{G}).$$

Proof. The construction of the spectral sequence out of the filtration is standard. The associated graded complex GC_*^{Bott} for our filtered complex $(C_*^{Bott}, \mathcal{F}_*^{Bott})$ is given by

$$GC_*^{Bott} = \bigoplus_k C_*(S_k),$$

which leads to the formula (7.8). The convergence of the spectral sequence to the Bott-type Seiberg-Witten Floer homology is a consequence of the basic theory of spectral sequences. The case of the dual filtration is similar. \square

Finally, we introduce a tool which will be used for proving the invariance of our Seiberg-Witten Floer homology theories.

Definition 7.8. Consider a finite collection $\mathbf{F} = \{f_1, \dots, f_m\}$, where each f_i is a smooth map from a compact smooth manifold with corners to some S_α . A generalized singular cube $f : \Delta \rightarrow S_\alpha$ for some α is called \mathbf{F} -transversal, provided that the following holds:

- 1) f is smooth,
- 2) f and $f|_{\partial \Delta}$ are transversal to every f_i and

3) for each sequence β_1, \dots, β_j with $\mu(\alpha) > \mu(\beta_1) > \dots > \mu(\beta_j)$, the induced maps $\pi_+ : \Delta \times_{S_\alpha} (\mathbf{M}_T^0(S_\alpha, S_{\beta_1}) \times_{S_{\beta_1}} \dots \times_{S_{\beta_{j-1}}} \mathbf{M}_T^0(S_{\beta_{j-1}}, S_{\beta_j})) \rightarrow S_{\beta_j}$ and $\pi_+ : \partial\Delta \times_{S_\alpha} (\mathbf{M}_T^0(S_\alpha, S_{\beta_1}) \times_{S_{\beta_1}} \dots \times_{S_{\beta_{j-1}}} \mathbf{M}_T^0(S_{\beta_{j-1}}, S_{\beta_j})) \rightarrow S_{\beta_j}$ are transversal to every f_i .

We have the subgroup $Q_j^{\mathbf{F}}(\mathcal{R}^0)$ of $Q_j(\mathcal{R}^0)$ generated by \mathbf{F} -transversal generalized singular cubes, which yields a subgroup $C_j^{\mathbf{F}}(\mathcal{R}^0)$ of $C_j(\mathcal{R}^0)$, whose elements will be called \mathbf{F} -transversal chains. We define

$$C_k^{Bott, \mathbf{F}} = \bigoplus_{i+j=k} C_j^{\mathbf{F}}(S_i)$$

and obtain therewith a subcomplex $C_*^{Bott, \mathbf{F}}$ of C_*^{Bott} . It is easy to see that this is indeed a subcomplex. We have the corresponding cochain complex $C_{Bott, \mathbf{F}}^*$ dual to $C_*^{Bott, \mathbf{F}}$.

For each given \mathbf{F} , we obtain a homology $H_*(C_*^{Bott, \mathbf{F}}, \partial_{Bott})$ and a cohomology $H^*(C_{Bott, \mathbf{F}}^*, \partial_{Bott}^*)$.

Lemma 7.8. *The homology $H_*(C_*^{Bott, \mathbf{F}}, \partial_{Bott})$ is canonically isomorphic to the homology $H_*(C_*^{Bott}, \partial_{Bott})$. Similarly, the cohomology $H^*(C_{Bott, \mathbf{F}}^*, \partial_{Bott}^*)$ is canonically isomorphic to the cohomology $H^*(C_{Bott}^*, \partial_{Bott}^*)$.*

Proof. We have the inclusion homomorphism $I : C_*^{Bott, \mathbf{F}} \rightarrow C_*^{Bott}$ and the induced homomorphism $I_* : H_*(C_*^{Bott, \mathbf{F}}, \partial_{Bott}) \rightarrow H_*(C_*^{Bott}, \partial_{Bott})$. We claim that I_* is an isomorphism. First, consider a closed chain σ in $C_*^{Bott, \mathbf{F}}$ such that $F(\sigma) = \partial_{Bott}\sigma_1$ with $\sigma_1 \in C_*^{Bott}$. By smooth approximation and transversal perturbations we deform σ_1 to obtain $\sigma_2 \in C_*^{Bott, \mathbf{F}}$ such that $\sigma = \partial_{Bott}\sigma_2$. This shows that F_* is a monomorphism. On the other hand, we can use smooth approximation and transversal perturbations to deform an arbitrary closed chain $\sigma \in C_*^{Bott}$ to obtain a closed chain $\sigma' \in C_*^{Bott, \mathbf{F}}$ such that $F(\sigma') - \sigma$ is homologous to zero. This shows that F_* is an epimorphism.

The arguments for the cohomologies are similar. \square

8. INVARIANCE

Consider two chain complexes (C_*, ∂) and $(\bar{C}_*, \bar{\partial})$. A chain map of degree $m \in \mathbb{Z}$ from the former to the latter consists of homomorphisms $F : C_k \rightarrow \bar{C}_{k+m}$ such that $F \cdot \partial = \bar{\partial} \cdot F$. A shifting homomorphism $F : H_*(C_*, \partial) \rightarrow H_*(\bar{C}_*, \bar{\partial})$ of degree $m \in \mathbb{Z}$ consists of homomorphisms $F : H_k(C_*, \partial) \rightarrow H_{k+m}(\bar{C}_*, \bar{\partial})$. These concepts are also defined for cochain complexes and cohomologies in a similar way, with the degree shifting in the opposite way, i.e. $k \rightarrow k - m$.

We shall establish the following invariance result.

Theorem 8.1. *The Bott-type Seiberg-Witten Floer homology and cohomology are diffeomorphism invariants modulo shifting isomorphisms.*

In other words, these homology and cohomology are independent of the metric h and generic parameter (π, λ) modulo shifting isomorphisms.

In this section, we construct a shifting homomorphism, which will be shown in [29] to be a desired shifting isomorphism. We would like to point out that the construction already contains some basic arguments for establishing the isomorphism property.

Consider two metrics h_+ and h_- on Y and Y -generic parameters (π_+, λ_+) for h_+ and (π_-, λ_-) for h_- respectively. We would like to construct a shifting isomorphism between our homologies (cohomologies) constructed with respect to (h_+, π_+, λ_+) and (h_-, π_-, λ_-) respectively.

Choose a smooth path of metrics $h(t)$ on Y such that

$$h(t) = \begin{cases} h_-, & \text{if } t < -1, \\ h_+, & \text{if } t > 1, \end{cases}$$

a smooth path of $\pi(t) \in \Pi$

$$\pi(t) = \begin{cases} \pi_-, & \text{if } t < -1, \\ \pi_+, & \text{if } t > 1, \end{cases}$$

and a smooth function $\lambda(t) \in \mathbb{R}$ such that

$$\lambda = \begin{cases} \lambda_-, & \text{if } t < -1, \\ \lambda_+, & \text{if } t > 1. \end{cases}$$

The following lemma is an immediate consequence of Lemma B.1 in Appendix B.

Lemma 8.2. *Fix an $(A_0, \Phi_0) \in \mathcal{A}(X) \times \Gamma^+(X)$. Indeed, we use the reference element u_0 introduced before as (A_0, Φ_0) . For any $R > 0$, set $X_R = Y \times [-R, R]$ and*

$$\begin{aligned} \mathcal{S}_R = \{u = (A_0, \Phi_0) + (A, \Phi) : (A, \Phi) \in \Omega_2^1(X_R) \times \Gamma_2^+(X_R) \\ \text{with } d^*A = 0 \text{ and } A|_{\partial X_R}(\frac{\partial}{\partial t}) = 0.\} \end{aligned}$$

Then \mathcal{S}_R is a global slice for the action of $\mathcal{G}_3^0(X_R)$ on $\mathcal{A}_2(X_R) \times \Gamma_2^+(X_R)$. In other words, $\mathcal{A}_2(X_R) \times \Gamma_2^+(X_R) = \mathcal{G}_3^0(X_R) \cdot \mathcal{S}_R$.

The following definition is a crucial construction.

Definition 8.3. Choose a nonzero $\Psi_0 \in \Gamma^-(X)$ with support contained in the interior of X_1 . We define a smooth vector field Z on $\mathcal{A}_2(X_1) \times \Gamma_2^+(X_1)$ by

$$Z(g^*u) = g^{-1}\Psi_0,$$

for $g \in \mathcal{G}_3^0(X_1), u \in \mathcal{S}_1$. Furthermore, we extend Z to $\mathcal{A}_{2,loc}(X) \times \Gamma_{2,loc}^+(X)$ by setting

$$Z(u) = Z(u|_{X_1}).$$

The following lemma is readily proved.

Lemma 8.4. *Z is equivariant with respect to the action of $\mathcal{G}_{3,loc}^0$.*

Let X be endowed with the warped product metric determined by the family of metrics $h(t)$ and the standard metric on \mathbb{R} . We introduce the (perturbed) transition trajectory equation

$$(8.1) \quad \begin{cases} F_A^+ = & \frac{1}{4} \langle e_i e_j \Phi, \Phi \rangle e^i \wedge e^j \\ & + (\nabla H_{\pi(t)}(a) + b_0) \wedge dt + *((\nabla H_{\pi(t)}(a) + b_0) \wedge dt), \\ & \partial_t \lambda = \lambda \left(\frac{\partial}{\partial t} \log \left(\frac{h_+}{h_-} \right) \right) \end{cases}$$

where b_0 denotes a smooth 1-form of compact support on X without dt -part. Setting $A = a + fdt$, $\Phi = \phi$ as usual we can rewrite (8.1) as follows

$$(8.2) \quad \begin{cases} \frac{\partial a}{\partial t} &= *_Y F_a + d_Y f + \langle e_i \cdot \phi, \phi \rangle e^i + \nabla H_{\pi(t)}(a) + b_0, \\ \frac{\partial \phi}{\partial t} &= -\not\partial_a \phi - \lambda(t)\phi + Z. \end{cases}$$

Of course, in those places in (8.2) where the metric on Y is involved, we use the metric $h(t)$. For example, the Hodge $*_Y$ at time t in the equation is that of the metric $h(t)$. The perturbation term Z (or $\partial/\partial t \cdot Z$) is called a *spinor perturbation*. We have the following obvious, but crucial lemma.

Lemma 8.5. *The equation (8.1) is invariant with respect to the action of $\mathcal{G}_{3,loc}^0$. Moreover, it has no reducible solution.*

A fundamental property of the Seiberg-Witten equation is a pointwise maximum principle for the spinor field, which is a consequence of the Weitzenböck formula (2.1). With the presence of Z , this principle no longer holds. Instead, we have the following result.

Lemma 8.6. *Let $(A, \Phi) = (a + fdt, \phi)$ be a solution of (8.1). Then there holds*

$$\|\Phi\|_{L^\infty} \leq C(1 + E(A, \Phi))^{1/4}$$

for a constant C depending only on the families $h(t), \pi(t), \lambda(t)$ and the geometry of Y .

Proof. Before proceeding with the proof, we first observe that by (2.17) and (8.1) the energy can be estimated in the following way

$$(8.3) \quad \begin{aligned} E((A, \Phi)) &= 2 \lim_{t \rightarrow -\infty} \mathbf{cs}_{(\lambda, H)}(a(\cdot, t), \phi(\cdot, t)) \\ &\quad - 2 \lim_{t \rightarrow +\infty} \mathbf{cs}_{(\lambda, H)}(a(\cdot, t), \phi(\cdot, t)) + \int_X |Z|^2, \end{aligned}$$

$$(8.4) \quad \int_X |Z|^2 < C.$$

Using local Columb gauges provided by Lemma B.1 in Appendix B and a patching argument, we can perform a gauge transformation to convert (A, Φ) into a smooth solution. Since the L^∞ norm of Φ is invariant, we can assume that (A, Φ) is already smooth. Furthermore, we can assume that (A, Φ) is in temporal form.

For simplicity, we assume $\lambda(t) \equiv 0$ in the following argument. It is easy to modify the argument to handle $\lambda(t)$. Put

$$I_1 = \not\partial_a \phi, I_2 = *_Y F_a + \langle e_i \cdot \phi, \phi \rangle e^i - \nabla H(a, \phi), I_3 = I_2 + \nabla H(a, \phi).$$

Then

$$\int (|I_1|^2 + |I_2|^2) = \frac{1}{2} E((A, \Phi), \Omega),$$

for any domain $\Omega \subset X$. For each $t \in \mathbb{R}$, we use the 3-dimensional Weitzenböck formula (2.1) on $Y \times \{t\}$ to derive

$$(8.5) \quad \partial_a I_1 = -\Delta_a \phi + \frac{\bar{s}}{4} \phi + \frac{1}{2} |\phi|^2 \phi + I_3 \phi,$$

where \bar{s} denotes the scalar curvature function of $(Y, h(t))$. Multiplying (8.5) by ϕ and integrating by parts, we infer

$$\int_{Y \times \{t\}} (|\nabla_a \phi|^2 + \frac{s}{4} |\phi|^2 + \frac{1}{2} |\phi|^4) \leq \int_{Y \times \{t\}} |I_1| \cdot |\partial_a \phi| + |I_3| |\phi|^2.$$

Using the Hölder inequality we then deduce that

$$\int_{Y \times \{t\}} (|\nabla_a \phi|^2 + \frac{1}{4} |\phi|^4) \leq C(1 + \int_{Y \times \{t\}} (|I_1|^2 + |I_2|^2)),$$

where C depends only on $\|\nabla H\|_{L^\infty}$, which can be estimated by appealing to Lemma 3.13. This last estimate implies

$$(8.6) \quad \int_{X_{R-2, R+2}} (|\nabla_a \phi|^2 + |\phi|^4) \leq C(1 + E(A, \Phi, X_{R-2, R+2}))$$

for any $R > 0$, where $X_{r, R} = Y \times [r, R]$.

Next we apply the Moser iteration to deduce the desired L^∞ estimate. Let $\xi : X \rightarrow [0, 1]$ be a cut-off function such that $\text{supp } \xi \subset X_{R-2, R+2}$ and $\xi(t) = 1$ for $t \in X_{R-1, R+1}$. By the 4-dimensional Weitzenböck formula (2.1) on X (recall that X is endowed with warped product metric), we have

$$(8.7) \quad D_A Z = -\Delta_A \Phi + \frac{s}{4} \Phi - \frac{1}{4} |\Phi|^2 \Phi.$$

Choosing $\xi^2 |\Phi|^p \Phi$ as a test function, where $p > 0$ will be determined later, we obtain

$$(8.8) \quad \begin{aligned} & \int_X \xi^2 (\nabla_A \Phi \cdot \nabla_A (|\Phi|^p \Phi) + \frac{s}{4} |\Phi|^{p+2} + \frac{1}{4} |\Phi|^{p+4}) \\ &= - \int_X (\xi^2 Z D_A (|\Phi|^p \Phi) + 2\xi \nabla \xi Z |\Phi|^p \Phi + 2\xi \nabla \xi \cdot \nabla \Phi |\Phi|^p \Phi). \end{aligned}$$

We have

$$\begin{aligned} \nabla_A (|\Phi|^p \Phi) &= |\Phi|^p \nabla_A \Phi + d |\Phi|^p \Phi \\ &= |\Phi|^p \nabla_A \Phi + p |\Phi|^{p-1} \frac{\langle \nabla_A \Phi, \Phi \rangle}{|\Phi|} \Phi, \nabla_A \Phi \cdot \nabla_A (|\Phi|^p \Phi) \\ &= |\Phi|^p |\nabla_A \Phi|^2 + p |\Phi|^{p-2} \langle \nabla_A \Phi, \Phi \rangle^2. \end{aligned}$$

On the other hand,

$$\eta |\Phi|^{p-2} \langle \nabla_A \Phi, \Phi \rangle^2 \leq C_\varepsilon |\Phi|^p |\nabla_A \Phi|^2 + \frac{C p^2}{\varepsilon} |\Phi|^p.$$

where $\varepsilon > 0$ is arbitrary. Similarly,

$$\begin{aligned} |D_A(|\Phi|^p\Phi)| &\leq |\Phi|^p|D_A\Phi| + |d|\Phi|^p \cdot \Phi| \\ &\leq C|\Phi|^p|\nabla_A\Phi| + p|\Phi|^{p-1}|\nabla_A\Phi| \\ &\leq C\varepsilon|\Phi|^p|\nabla_A\Phi|^2 + \frac{C}{\varepsilon}(p^2 + 1)|\Phi|^p, \end{aligned}$$

$$|\nabla\xi||\xi||Z||\Phi|^{p+1} \leq C\xi^2|\Phi|^p + C|\nabla\xi|^2|\Phi|^{p+2},$$

and

$$|\nabla\xi||\xi||\nabla\Phi||\Phi|^{p+1} \leq \varepsilon\xi^2|\Phi|^p|\nabla\Phi|^2 + \frac{C}{\varepsilon}|\nabla\xi|^2|\Phi|^{p+2}.$$

Choosing ε suitably, we deduce

$$\int \xi^2|\Phi|^p|\nabla_A\Phi|^2 \leq C \int ((p^2 + 1)\xi^2|\Phi|^p + |\nabla\xi|^2|\Phi|^{p+2}).$$

Consequently,

$$\int_X \xi^2|\Phi|^p|\nabla|\Phi||^2 \leq C \int_X ((p^2 + 1)\xi^2|\Phi|^p + |\nabla\xi|^2|\Phi|^{p+2}),$$

or

$$\int_X \xi^2|\nabla|\Phi|^{\frac{p+2}{2}}|^2 \leq C(p+1)^2 \int_X (\xi^2|\Phi|^p + |\nabla\xi|^2|\Phi|^{p+2}).$$

Now we set $w = |\Phi|^{(p+2)/2}$. By the Sobolev inequality and Hölder inequality, we arrive at

$$\begin{aligned} \|\xi w\|_{L^2}^2 &\leq C(p+1)^2 \int (|\xi\nabla w|^2 + |w\nabla\xi|^2) \\ &\leq C(p+1)^2 (\|\xi w\|_{L^2}^{\frac{p}{p+2}} + \|\nabla\xi|w\|_{L^2}). \end{aligned}$$

Then we use the iteration process as presented in [14] to infer

$$\sup_{X_{R-1, R+1}} |\Phi| \leq C\|\Phi\|_{L^4(X_{R-2, R+2})}.$$

Combining it with (8.6) we are done. \square

Since (8.1) is identical to (2.13) near time infinities, we have

Proposition 8.7. *An analogue of Proposition 4.2 for (8.1) holds.*

We have various configuration spaces and moduli spaces associated with (8.1) which are analogous to the spaces introduced in Sections 4 and 6. All the analysis in Sections 4, 5 and 6 carries over. We shall be brief in formulating the relevant results.

Let e.g. \mathcal{R}_\pm denote the \mathcal{R} for $(h_\pm, \pi_\pm, \lambda_\pm)$. Consider $\alpha_- \in \mathcal{R}_-$, $\alpha_+ \in \mathcal{R}_+$ and $p_\pm \in \alpha_\pm$. We have the spaces of transition trajectories $\mathcal{N}(p_-, p_+)$ and the moduli spaces $\mathcal{M}^0(p_-, p_+)$, $\mathcal{M}_T^0(S_{\alpha_-}, S_{\alpha_+})$ etc.. We also have the various spaces of consistent multiple temporal transition trajectory classes, and those of consistent piecewise transition trajectories. In particular, we have $\mathbf{M}_T^0(S_{\alpha_-}, S_{\alpha_+})$. For example a multiple temporal transition trajectory class is a tuple $([y_+], [T_+], [y_-], [T_-]) \in$

$\mathcal{M}_T^0(S_{\alpha_0^-}, S_{\alpha_1^-}) \times_{S_{\alpha_1^-}} \dots \mathcal{M}_T^0(S_{\alpha_{m-1}^-}, S_{\alpha_m^+}) \dots \mathcal{M}_T^0(S_{\alpha_{k-1}^+}, S_{\alpha_k^+})$ with a distinguished piece $[u_m]_0^T, 1 \leq m \leq k$. The time translation acts on all pieces except the distinguished one. We define the \mathbb{R} -action on the distinguished one as the trivial action. In particular, the \mathbb{R} -action on $\mathcal{M}_T^0(S_{\alpha_-}, S_{\alpha_+})$ is defined to be the trivial action. As before, we use the underline to denote quotient by the \mathbb{R}^k -action in the case of temporal transition trajectories and multiple temporal transition trajectory classes. In the case of general transition trajectories and piecewise transition trajectories, we have the \mathbb{R}^k -action induced from the twisted time translation, where the \mathbb{R} -action on the distinguished portion is again trivial.

Let O_{\pm} be the unique reducible elements in \mathcal{R}_{\pm} respectively. We set

$$(8.9) \quad m_0 = \text{ind } \mathcal{F}_{o_-, o_+} - 1,$$

where $o_- \in O_-, o_+ \in O_+$.

The following lemma is analogous to Corollary 5.2.

Lemma 8.8. *We have*

$$(8.10) \quad \text{ind } \mathcal{F}_{p_-, p_+} = \mu_-([p_-]) - \mu_+([p_+]) + m_0 + \dim G_{p_+}.$$

We have the following analogues of Theorem 4.14, Theorem 6.15 and Theorem 6.19.

Theorem 8.9. *The moduli spaces $\underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+})$ are compact Hausdorff spaces with respect to the topology induced by the piecewise exponential convergence. Moreover, for generic b_0 , we have for all $\alpha_- \in \mathcal{R}_-, \alpha_+ \in \mathcal{R}_+$*

(1) $\underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+})$ has the structure of d -dimensional smooth oriented manifolds with corners, where

$$\begin{aligned} d &= \text{ind } \mathcal{F}_{p_-, p_+} - \max\{\dim G_{p_-}, \dim G_{p_+}\} + 1 \\ &= \mu(\alpha_-) - \mu(\alpha_+) + m_0 + \dim G_{p_+} - \max\{\dim G_{p_-}, \dim G_{p_+}\} + 1 \end{aligned}$$

for $p_- \in \alpha_-, p_+ \in \alpha_+$.

(2) *This structure is compatible with the natural stratification of $\underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+})$.*

(3) *The temporal projections $\pi_- : \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+}) \rightarrow S_{\alpha_-}$ and $\pi_+ : \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+}) \rightarrow S_{\alpha_+}$ are smooth maps. (But they may not be fibrations in general.)*

Consequently, there holds

$$(8.11) \quad \begin{aligned} \partial \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+}) &= (\cup_{\mu(\alpha_-) > \mu(\alpha'_-) \geq \mu(\alpha_+) - m_0} \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha'_-}) \times_{S_{\alpha'_-}} \underline{\mathbf{M}}_T^0(S_{\alpha'_-}, S_{\alpha_+})) \cup \\ &(\cup_{\mu(\alpha_-) \geq \mu(\alpha'_+) - m_0 > \mu(\alpha_+) - m_0} \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha'_+}) \times_{S_{\alpha'_+}} \underline{\mathbf{M}}_T^0(S_{\alpha'_+}, S_{\alpha_+})). \end{aligned}$$

(Of course, only the nonempty moduli spaces appear in this equation.)

The key point here is that Lemma 8.5 rules out reducible transition trajectories. Note that instead of using holonomy perturbations we now use the perturbation b_0 as in [15]. (This perturbation is not time translation equivariant, and hence can't be applied in the construction of our homologies.)

Now we proceed to construct the desired shifting isomorphism

Consider the Bott complexes C_*^{Bott-} and C_*^{Bott+} associated with the parameters (h_-, π_-, λ_-) and (h_+, π_+, λ_+) respectively. Let \mathbf{F} denote the collection of all the projections $\pi_- : \mathbf{M}_T^0(S_{\alpha_-}, S_{\alpha_+}) \rightarrow S_{\alpha_-}$ and $\pi_+ : \mathbf{M}_T^0(S_{\alpha_-}, S_{\alpha_+}) \rightarrow S_{\alpha_+}$. We construct our chain homomorphism $F : C_k^{Bott-, \mathbf{F}} \rightarrow C_{k+m_0}^{Bott+}$. For simplicity, we assume that the coefficient group is \mathbb{Z} . The general case is similar. Consider $\sigma \in C_j^{\mathbf{F}}(S_{\alpha_-})$ with $\sigma = [(\Delta, f)]$ and $j + \mu(\alpha_-) = k$. For each α_+ with the corresponding moduli space nonempty, we follow the construction of the boundary operator $\partial_{\alpha, \beta}$ in Section 7 to obtain a generalized cubical singular chain $\sigma' \in C_{j'}(S_{\alpha_+})$ with $j' + \mu_+(\alpha_+) = j + \mu_-(\alpha_-) + m_0$. We define $F_{\alpha_-, \alpha_+}(\sigma)$ to be σ' . We define it to be zero if the moduli space is empty. Then we set

$$F(\sigma) = \sum_{\mu_+(\alpha_+) \leq \mu_-(\alpha_-) + m_0} F_{\alpha_-, \alpha_+}(\sigma).$$

It is easy to see that we indeed have $F : C_k^{Bott-, \mathbf{F}} \rightarrow C_{k+m_0}^{Bott+}$.

Theorem 8.10. *We have $\partial_{Bott} \cdot F = F \cdot \partial_{Bott}$, hence F is a chain map of degree m_0 from $(C_*^{Bott-, \mathbf{F}}, \partial_{Bott})$ to $(C_*^{Bott+}, \partial_{Bott})$. The induced shifting homomorphism between the homologies is denoted F_* . From the construction of F we easily obtain a cochain map of degree m_0 and the induced shifting homomorphism F^* between cohomologies.*

Proof. The proof goes along the same lines as the proof of Lemma 7.5. First, we arrange the orientations such that

$$(8.12) \quad \partial \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+}) = (-1)^{\mu(\alpha_-) + \dim S_{\alpha_-} + m_0 + 1} (\Sigma_{\alpha'_-} \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha'_-}) \times_{S_{\alpha'_-}} \underline{\mathbf{M}}_T^0(S_{\alpha'_-}, S_{\alpha_-})) + (\Sigma_{\alpha'_+} \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha'_+}) \times_{S_{\alpha'_+}} \underline{\mathbf{M}}_T^0(S_{\alpha'_+}, S_{\alpha_+})).$$

In comparison with (7.3), we see an additional 1 appearing in the sign exponent (note that m_0 reduces to zero in the situation of (7.3)). This is because of the difference in the time translation action. Consider $\sigma = [(\Delta, f)] \in C_j(S_{\alpha_-})$. Analogous to (7.5) we have for α_+ with $\mu(\alpha_+) \leq \mu_-(\alpha_-) + m_0$

$$\begin{aligned} \partial(\Delta \times_{S_{\alpha_-}} \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+})) &= \partial \Delta \times_{S_{\alpha_-}} \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+}) + \\ &(-1)^{j + \dim S_{\alpha_-}} \Delta \times_{S_{\alpha_-}} \partial \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+}) = \\ &\partial \Delta \times_{S_{\alpha_-}} \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha_+}) + \\ &(-1)^{j + \mu(\alpha_-) + m_0 + 1} \sum_{\alpha'_-} \Delta \times_{S_{\alpha'_-}} \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha'_-}) \times_{S_{\alpha'_-}} \underline{\mathbf{M}}_T(S_{\alpha'_-}, S_{\alpha_+}) + \\ &(-1)^{j + \mu(\alpha_-) + m_0 + 1} \sum_{\alpha'_+} \Delta \times_{S_{\alpha'_+}} \underline{\mathbf{M}}_T^0(S_{\alpha_-}, S_{\alpha'_+}) \times_{S_{\alpha'_+}} \underline{\mathbf{M}}_T(S_{\alpha'_+}, S_{\alpha_+}). \end{aligned}$$

This implies

$$\begin{aligned} \partial_0 F_{\alpha_-, \alpha_+} \sigma &= F_{\alpha_-, \alpha_+} \partial_0 \sigma + \sum_{\mu(\alpha_-) > \mu(\alpha'_-) \geq \mu(\alpha_+) - m_0} F_{\alpha'_-, \alpha_+} \partial_{\alpha_-, \alpha'_-} \sigma - \\ &\partial_{\alpha'_+, \alpha_+} F_{\alpha_-, \alpha'_+} \sigma. \end{aligned}$$

Summing over all α_+ we arrive at the desired chain map property. \square

APPENDIX A. THE ORDINARY SEIBERG-WITTEN FLOER HOMOLOGY

Here, we only consider rational homology spheres. It is not hard to extend the construction to general manifolds.

For simplicity, we formulate the theory with integer coefficients. Let Y be a rational homology sphere with a given metric h and c a $spin^c$ structure on Y as before. Set $C_i = \mathbb{Z}\{\alpha \in \mathcal{R}^* | \mu(\alpha) = i\}$. We define a boundary operator $\partial : C_i \rightarrow C_{i-1}$ in terms of the moduli spaces $\widetilde{\mathcal{M}}(\alpha, \beta)$, or equivalently $\widetilde{\mathcal{M}}(p, q), p \in \alpha, q \in \beta$, where the tilde means quotient by the time translation action. For a generic pair (π, λ) , $\widetilde{\mathcal{M}}(\alpha, \beta)$ is a compact oriented manifold of zero dimension, provided that $\mu(\alpha) - \mu(\beta) = 1$. The orientation (sign) is given by pairing the orientation of $\mathcal{M}(\alpha, \beta)$ provided by Proposition 5.3 with its orientation induced by the time translation action. For $\alpha \in C_i$ we then set

$$(A.1) \quad \partial\alpha = \sum_{\mu(\alpha) - \mu(\beta) = 1} \# \widetilde{\mathcal{M}}(\alpha, \beta) \beta,$$

where $\# \widetilde{\mathcal{M}}(p, q)$ is the algebraic sum of $\widetilde{\mathcal{M}}(\alpha, \beta)$.

The compactification of the moduli spaces $\widetilde{\mathcal{M}}(\alpha, \gamma)$ with $\mu(\alpha) - \mu(\gamma) = 2$ is similar to the results in Section 6. For dimensional reasons, no trajectory connecting to the reducible point appears in the compactification. Using these compactified moduli spaces and the consistency of orientation (Proposition 5.3) we obtain

Lemma A.1. $\partial^2 = 0$.

Definition A.2. The ordinary Seiberg-Witten Floer homology $FH_*^{SW}(c, h, \pi, \lambda)$ and cohomology $FH^{SW*}(c, h, \pi, \lambda)$ for the $spin^c$ structure c and the parameters h, π, λ are defined to be the homology and cohomology of the chain complex (C_*, ∂) .

APPENDIX B. LOCAL GAUGE FIXING

In this appendix we prove a result on local Columb gauge fixing. We assume that Y is a rational homology sphere. Let c be a fixed $spin^c$ structure on Y and a_0 a smooth reference connection as in Section 3. It induces a time-independent connection A_0 over X . For positive numbers $r < R$, let $x_0 \in X_{r,R}$ be a reference point.

Lemma B.1. *For any $A \in \mathcal{A}_l(X_{r,R})$ with $l \geq 1$, there exists a unique gauge $g \in \mathcal{G}_{l+1}(X_{r,R})$ with $g(x_0) = 1$ such that $\widetilde{A} = g^*(A)$ satisfies*

$$(B.1) \quad \begin{cases} d^*(\widetilde{A} - A_0) = 0, \\ (\widetilde{A} - A_0)(\nu) = 0 \text{ on } \partial X_{r,R}, \end{cases}$$

where ν denotes the unit outer normal of $\partial X_{r,R}$. Moreover, we have

$$\|\widetilde{A} - A_0\|_{1,2} \leq C \|F_{\widetilde{A}} - F_{A_0}\|_{0,2}$$

for a positive constant C depending only on Y, r and R .

Proof. The associated gauge fixing equation is

$$(B.2) \quad \begin{cases} d^*(g^{-1}dg) + d^*(A - A_0) = 0, \\ \dots \end{cases}$$

If we choose $g = e^f$, then the equation reduces to

$$(B.3) \quad \begin{cases} d^*df + d^*(A - A_0) &= 0, \\ \frac{\partial f}{\partial \nu} + (A - A_0)(\nu) &= 0 \text{ on } \partial X_{r,R}. \end{cases}$$

It is clear that a solution $f \in \Omega_{l+1}^0(X_{r,R})$ with $f(x_0) = 0$ exists.

Now assume that there are g_1 and g_2 satisfying (B.2) with $g_1(x_0) = g_2(x_0) = 1$. We set $g = g_1^{-1}g_2$. Then g is a solution of the problem

$$\begin{cases} d^*(g^{-1}dg) &= 0, \\ g^{-1}dg(\nu) &= 0 \text{ on } \partial X_{r,R}. \end{cases}$$

Taking two copies of $X_{r,R}$ and gluing them along their common boundary, we obtain the Riemannian manifold $Y \times S^1$. The two copies of g then yield a solution $g_0 : Y \times S^1 \rightarrow S^1$ of the harmonic map equation $d^*(g^{-1}dg) = 0$. This means that the 1-form $g^{-1}dg$ is harmonic.

We claim that every harmonic 1-form on $Y \times S^1$ is a constant multiple of the base harmonic form ds , where s is the arclength parameter on S^1 . Indeed, let $\alpha = \alpha_Y + f ds$ be an arbitrary harmonic 1-form, where α_Y has no ds component. Then we deduce from the equation $d\alpha = 0$

$$\begin{cases} d_Y \alpha_Y &= 0, \\ \frac{\partial \alpha_Y}{\partial s} &= d_Y f. \end{cases}$$

Since Y is a rational homology sphere, the first equation above implies that $\alpha_Y = d_Y h$ for a function h . Then the second equation implies that $f = \frac{\partial h}{\partial s} + f_0$ for a function f_0 depending only on s . It follows that $\alpha = dh + f_0 ds$. Now we use the equation $d^* \alpha = 0$ to deduce

$$d^*dh + \frac{df_0}{ds} = 0.$$

Integrating f_0 along S^1 we obtain another solution h_0 of this equation, which depends only on s . Clearly h differs from h_0 by a constant. We deduce that $\frac{dh}{ds} + f_0$ is a constant, whence α is a constant multiple of ds .

Since $g^{-1}dg$ is harmonic, it is a constant multiple of ds . On the other hand, for each $y \in Y$, $g_0(y, \cdot)$ is a map from S^1 to S^1 with degree zero, hence *i.e.* $\int_{\{y\} \times S^1} g_0^{-1}dg_0 = 0$. It follows that $g^{-1}dg = 0$. We deduce that g is constant. Consequently, $g_1 \equiv g_2$.

Next consider an A satisfying (B.1). By the above gluing argument, the form $A - A_0$ leads to a form A_1 on $Y \times S^1$ satisfying $d^*A_1 = 0$. The Hodge decomposition of A_1 takes the following form

$$A_1 = cds + d^*\beta,$$

where c denotes a constant and $\beta \in \Omega_2^2(Y \times S^1)$. The constant c is given by the L^2 product between A_1 and ds . By the construction of A_1 , it is easy to see that this product is zero. Now we deduce

$$\|A_1\|_{1,2} = \|d^*\beta\|_{1,2} \leq C\|dd^*\beta\|_{0,2} = C\|dA_1\|_{0,2}.$$

But $\|dA_1\|_{0,2} = \sqrt{2}\|F_A - F_{A_0}\|_{0,2}$, hence the desired estimate follows. \square

Remark B.2. If Y is not a homology sphere, then we can still achieve the gauge fixing (B.1) provided that the L^2 norm of the curvature of A is sufficiently small. This is similar to Uhlenbeck's gauge fixing lemma in the Yang-Mills theory, cf. [9] [24]

APPENDIX C. A TRANSVERSALITY

In this appendix we prove transversality at reducible trajectories.

Lemma C.1. *Let (π, λ) be a pair of Y -generic parameters such that $\nabla^2 H$ is sufficiently small. Choose δ_-, δ_+ small enough (but positive) in the set-up of Definition 4.4. Then the operator $\mathcal{F}_{p,q}$ for $p, q \in O = O_{\pi, \lambda}$ (the unique reducible element in \mathcal{R}) at a Seiberg-Witten trajectory is onto.*

Proof. Consider $p, q \in O$ and $\mathcal{F} = \mathcal{F}_{p,q}$ at a trajectory (A_0, Φ_0) . By the arguments in the proof of Theorem 4.13 we can assume that $p = q = (a_0, 0)$ and $(A_0, \Phi_0) \equiv (a_0, 0)$. The formal adjoint \mathcal{F}^* of $\mathcal{F} = \mathcal{F}_{p,q}$ with respect to the product (4.4) is given by

$$(C.1) \quad \mathcal{F}^*(v) = -\frac{\partial v}{\partial t} - \begin{pmatrix} *_Y db + d_Y f - \nabla^2 H(a_0) \cdot b \\ -\not{\partial}_{a_0} \psi - \lambda \psi \\ d_Y^* b + 2\delta'_F f \end{pmatrix} - 2\delta'_F v$$

for $v = (\psi, b, f)$. The surjectivity of \mathcal{F} is equivalent to the vanishing of the kernel of \mathcal{F}^* . Let v satisfy $\mathcal{F}^*v = 0$. Then we have

$$(C.2) \quad \frac{\partial}{\partial t} \begin{pmatrix} b \\ f \end{pmatrix} + \begin{pmatrix} *_Y d_Y b + d_Y f \\ d_Y^* b \end{pmatrix} + \begin{pmatrix} \delta'_F b + \nabla^2 H(a_0) \cdot b \\ 0 \end{pmatrix} = 0.$$

We define the operator L by

$$L \begin{pmatrix} b \\ f \end{pmatrix} = - \begin{pmatrix} *_Y d_Y b + d_Y f \\ d_Y^* b \end{pmatrix}.$$

L is formally self-adjoint and satisfies $L^2 = \Delta$, where Δ denotes the Hodge Laplacian. Let $\{\xi_i = \begin{pmatrix} b_i \\ f_i \end{pmatrix}\}$ be a complete L^2 orthonormal system of eigenvectors of L with $L\xi_i = \lambda_i \xi_i$. From the above discussion we deduce

$$\Delta b_i = \lambda_i^2 b_i, \Delta f_i = \lambda_i^2 f_i.$$

Now we write

$$\begin{pmatrix} b \\ f \end{pmatrix} = \sum_{-\infty}^{+\infty} l_i(t) \xi_i.$$

Then it follows from C.2 that

$$(C.3) \quad \sum l'_i(t) \xi_i + \lambda_i l_i(t) \xi_i + \begin{pmatrix} \delta'_F b + \nabla^2 H(a_0) \cdot b \\ 0 \end{pmatrix} = 0.$$

Consequently, f_j is a nonzero constant. Then we deduce $l'_j(t) \equiv 0$, hence l_j is a constant. But (b, f) is L^2 integrable, which forces l_j to be zero. We conclude that the above expansion of (b, f) does not contain terms with zero eigenvalue. Using the elementary arguments in e.g. [21] it is then easy to show that (b, f) must vanish, provided that $\nabla^2 H$, δ_+ and δ_- have been chosen small enough. Using the same arguments one also infers that ψ vanishes. Thus $v = 0$. \square

REFERENCES

- [1] M.F. Atiyah, *New invariants of 3- and 4-dimensional manifolds*, Symp. Pure Math. **48** (1988), 285-299.
- [2] M.F. Atiyah, V.K. Patodi and I.M. Singer, *Spectral asymmetry and Riemannian geometry I, II, III*, Math. Proc. Cambridge Soc. **77**.
- [3] D. M. Austin and P. J. Braam, *Equivariant Floer theory and gluing Donaldson polynomials*, Topology **35** (1996), 167-200.
- [4] ———, *Morse-Bott theory and equivariant cohomology*, Floer Memorial Volume, Birkhäuser, 1996, pp. 123-164.
- [5] W. Chen, *Casson's invariant and Seiberg-Witten gauge theory*, preprint.
- [6] R. Cohen, J. Jones and G. Segal, *Floer's infinite dimensional Morse theory and homotopy theory*, Floer Memorial Volume, Birkhäuser, 1996, pp. 297-326.
- [7] S.K. Donaldson, *The orientation of Yang-Mills moduli spaces and 4-manifold topology*, J. Diff. Geo. **26** (1987), 397-428.
- [8] ———, *The Seiberg-Witten equations and the 4-manifold topology*, Bull. Amer. Math. Soc. (1996).
- [9] S.K. Donaldson and P. B. Kronheimer, *The geometry of four-manifolds*, Oxford Science Publications, 1990.
- [10] A. Floer, *An instanton invariant for 3-manifolds*, Comm. Math. Phys. **118** (1988), 215-240.
- [11] A. Floer, *Holomorphic spheres and symplectic fixed points*, Comm. Math. Phys..
- [12] Fukaya, *Floer homology of connected sum of homology 3-sphere*, Topology **36** (1996).
- [13] ———, *Aspects of low dimensional manifolds*, Advanced studies in pure mathematics, 1992, pp. 1-99.
- [14] Gilbarg and Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, 1981.
- [15] P. B. Kronheimer and T. S. Mrowka, *The genus of embedded surfaces in the projective space*, Math. Res. Letters **1** (1994), 797-808.
- [16] H. B. Lawson and M.-L. Michelsohn, *Spin geometry*, Princeton, New Jersey, 1989.
- [17] Y. Lim, *The equivalence of Seiberg-Witten and Casson invariants for homology 3-spheres*, preprint.
- [18] M. Marcolli, *Seiberg-Witten-Floer homology and Heegard splittings*, Inter. J. Math. **7** (1996), 671-696.
- [19] W. S. Massey, *Singular homology theory*, Graduate Texts in Mathematics, Springer-Verlag, 1980.
- [20] D. Salamon, *Spin geometry and the Seiberg-Witten invariants*, Lectures notes, Warwick, 1995.
- [21] D. Salamon and E. Zehnder, *Morse theory for periodic solutions of Hamiltonian system and the Maslov index*, Comm. Pure Appl. Math. **45** (1992), 1303-1360.
- [22] C. Taubes, *Casson's invariant and gauge theory*, J. Diff. Geom. **31** (1990), 547-599.
- [23] ———, *The Seiberg-Witten invariants and the Gromov invariants*, preprint.
- [24] K. K. Uhlenbeck, *Connections with L^p bounds on curvature*, Comm. Math. Phys. **83** (1982), 31-42.
- [25] R. Wang, *On Seiberg-Witten Floer invariants and the generalized Thom problem*, preprint.
- [26] G. Wang and Rugang Ye, *Bott-type and equivariant Seiberg-Witten Floer homology I*, DG9701010 (1997).
- [27] E. Witten, *Monopoles and 4-manifolds*, Math. Res. Letters **1** (1994), 769-796.
- [28] Rugang Ye, *Equivariant and Bott-type Seiberg-Witten Floer homology: Part II*, preprint.
- [29] Rugang Ye, *Equivariant and Bott-type Seiberg-Witten Floer homology: Part III*, in preparation.

Acknowledgement: The first named author was supported by SFB 237 and the Leibniz program of DFG. He is grateful to Prof. J. Jost for providing these supports.

E-mail address: `gwang@mis.mpg.de`

INSTITUTE OF SYSTEM SCIENCE, ACADEMIA SINICA, 100080 BEIJING, CHINA

E-mail address: `gfwang@iss06.iss.ac.cn`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106,
USA

E-mail address: `yer@math.ucsb.edu`

RUHR-UNIVERSITÄT BOCHUM, FAKULTÄT FÜR MATHEMATIK, 44780 BOCHUM, GERMANY

E-mail address: `ye@dgeo.ruhr-uni-bochum.de`