

A NOTE ON A CONJECTURE OF XIAO

Miguel A. BARJA¹.

Departament de Matemàtica Aplicada I. Universitat Politècnica de Catalunya. Barcelona. Spain

Francesco ZUCCONI².

Dipartimento di Matematica e Informatica. Università degli Studi di Udine. Udine. Italy

When $f : S \rightarrow B$ is a surjective morphism of a complex, smooth surface S onto a complex, smooth, genus b curve B , such that the fibre F of f has genus g , it is well known that $f_*\omega_{S/B} = \mathcal{E}$ is a locally free sheaf of rank g and degree $d = \mathcal{X}\mathcal{O}_S - (b-1)(g-1)$ and that f is not an holomorphic fibre bundle if and only if $d > 0$. In this case the *slope*, $\lambda(f) = \frac{K_S^2 - 8(b-1)(g-1)}{d}$, is a natural invariant associated by Xiao to f (cf. [7]). In [7][Conjecture 2] he conjectured that $\mathcal{E}$ has no locally free quotient of degree zero (i.e., $\mathcal{E}$ is ample) if $\lambda(f) < 4$. We give a partial affirmative answer to this conjecture:

Theorem 1. *Let $f : S \rightarrow B$ be a relatively minimal fibration with general fibre F . Let $b = g(B)$ and assume that $g = g(F) \geq 2$ and that f is not locally trivial.*

If $\lambda(f) < 4$ then $\mathcal{E} = f_\omega_{S/B}$ is ample provided one of the following conditions hold*

- (i) F is non hyperelliptic.
- (ii) $b \leq 1$.
- (iii) $g(F) \leq 3$.

Proof. (i) If $q(S) > b$ the result follows from [7] Corollary 2.1. Now assume $q(S) = b$. By Fujita's decomposition theorem (see [3], [4] and also [5] for a proof)

$$\mathcal{E} = \mathcal{A} \oplus \mathcal{F}_1 \oplus \dots \oplus \mathcal{F}_r$$

where $h^0(B, (\mathcal{A} \oplus \mathcal{F}_1 \oplus \dots \oplus \mathcal{F}_r)^*) = 0$, $\mathcal{A}$ is an ample sheaf and $\mathcal{F}_i$ are non trivial stable degree zero sheaves. Then we only must prove that $\mathcal{F}_i = 0$. If F is not hyperelliptic and $\text{rank}(\mathcal{F}_i) \geq 2$ the claim is the content of [7] Proposition 3.1. If $\text{rank}(\mathcal{F}_i) = 1$ we can use

¹Partially supported by CICYT PS93-0790 and HCM project n.ERBCHRXCT-940557.

²Partially supported by HCM project n.ERBCHRXCT-940557.

[2] §4.2 or [1] Theorem 3.4 to conclude that $\mathcal{F}_i$ is torsion in $\text{Pic}^0(B)$. Hence it induces an étale base change

$$\begin{array}{ccc} \tilde{S} & \longrightarrow & S \\ \downarrow \tilde{f} & & \downarrow f \\ \tilde{B} & \xrightarrow{\sigma} & B \end{array}$$

By flatness $\tilde{f}_*\omega_{\tilde{S}/\tilde{B}} = \sigma^*(f_*\omega_{S/B})$. Since σ is étale $\lambda(f) = \lambda(\tilde{f})$ and $\sigma^*(\mathcal{F}_i) = \mathcal{O}_{\tilde{B}}$ is a direct summand of $\tilde{f}_*\omega_{\tilde{S}/\tilde{B}}$. In particular by [3] $q(\tilde{S}) > \tilde{b} = g(\tilde{B})$ hence $\lambda(\tilde{f}) \geq 4$ by [7] Theorem 3.3: a contradiction.

(ii) If $b = 0$ the claim is trivial. If $b = 1$, any stable degree zero sheaf has rank one, then as in (i) we conclude.

(iii) If $g = 2$ and $\mathcal{E} \neq \mathcal{A}$, then $\mathcal{E} = \mathcal{A} \oplus \mathcal{L}$ where $\mathcal{L}$ torsion and we are done. The only non trivial case if $g = 3$ is $\mathcal{E} = \mathcal{A} \oplus \mathcal{F}$ where $\mathcal{A}$ an ample line bundle and $\mathcal{F}$ a stable, degree zero, rank two vector bundle. Then $K_{S/B}^2 \geq (2g - 2)\deg \mathcal{A} = 4d$ and we are done by [7][Theorem 2] $\square$

Theorem 3.3 of [7] Xiao says that if $q(S) > b$ and $\lambda(f) = 4$ then $\mathcal{E} = \mathcal{F} \oplus \mathcal{O}_B$, where $\mathcal{F}$ is a semistable sheaf. We have the following improvement:

Theorem 2. *Let $f : S \longrightarrow B$ be a relatively minimal non locally trivial fibration. If $\lambda(f) = 4$ then $\mathcal{E} = f_*\omega_{S/B}$ has at most one degree zero, rank one quotient $\mathcal{L}$. Moreover, in this case $\mathcal{E} = \mathcal{A} \oplus \mathcal{L}$ with $\mathcal{A}$ semistable and $\mathcal{L}$ torsion.*

Proof. As in the previous theorem the torsion subsheaf $\mathcal{L}$ becomes the trivial one after an étale base change; thus

$$\tilde{f}_*\omega_{\tilde{S}/\tilde{B}} = \tilde{\mathcal{A}} \oplus \mathcal{O}_{\tilde{B}}, \quad \tilde{\mathcal{A}} = \sigma^*\mathcal{A}.$$

By [7][Theorem 3.3], $\tilde{\mathcal{A}}$ is semistable. Then $\mathcal{A}$ is also semistable by [6][Proposition 3.2]. $\square$

References

- [1] Barja, M.A. *On the slope and geography of fibred surfaces and threefolds*. Ph. D. Thesis. 1998.
- [2] P. Deligne. *Théorie de Hodge II*. Publ. Math. I.H.E.S., **40** (1971), 5-58.
- [3] Fujita, T. *On Kahler fibre spaces over curves*. J. Math. Soc. Japan, **30** (1978), 779–794.
- [4] Fujita, T. *The sheaf of relative canonical forms of a Kahler fiber space over a curve*. Proc. Japan. Acad., **54**, Ser.A (1978), 183–184.
- [5] J. Kollár. *Subadditivity of the Kodaira dimension: fibers of general type*. Adv. studies in Pure Math. 10(1987), Algebraic Geometry, Sendai 1985. 361-398.
- [6] Y. Miyaoka. *The Chern classes and Kodaira dimension of a minimal variety*. Algebraic Geometry, Sendai (1985); Adv. Study in Pure Math. **10** (1987), 449-476.
- [7] Xiao, G. *Fibered algebraic surfaces with low slope*. Math. Ann. **276** (1987), 449–466.