

CHARACTERIZATION OF ABELIAN VARIETIES

JUNGKAI A. CHEN, CHRISTOPHER D. HACON

March 15, 1999

ABSTRACT. We prove that any smooth complex projective variety X with plurigenera $P_1(X) = P_2(X) = 1$ and irregularity $q(X) = \dim(X)$ is birational to an abelian variety.

INTRODUCTION

Given a smooth projective variety X , one would like to characterize it by its birational invariants. Classical results include Castelnuovo's criterion for rational surfaces, which states that a complex surface with plurigenera $P_2(X) = 0$ and irregularity $q(X) = 0$ must be rational. Another well known result due to Enriques, characterizes abelian surfaces as the only surfaces with $P_1(X) = P_2(X) = 1$ and $q(X) = 2$.

In an attempt to generalize Enriques' result to higher dimensional varieties, Kawamata proved that a variety X with Kodaira dimension $\kappa(X) = 0$ and irregularity $q(X) = \dim(X)$ admits a birational map to an abelian variety [Ka]. In [Ko1], Kollár gives an effective characterization by proving that if $P_1(X) = P_4(X) = 1$ and X has maximal Albanese dimension, then X is birational to an abelian variety. Ein and Lazarsfeld prove a similar result by using generic vanishing theorems [EL1]. Kollár subsequently improved his result by weakening the hypothesis to $P_3(X) = 1$ and $q(X) = \dim(X)$. He conjectured that the result still holds if one assumes that $P_2(X) = 1$ and X is generically finite over an abelian variety [Ko2].

In this paper, we verify Kollár's conjecture. Our method is based on the approaches of Kollár and of Ein and Lazarsfeld.

Acknowledgment. We are in debt to R. Lazarsfeld, L. Ein and J. Kollár for valuable conversations during the preparation of this paper.

CONVENTIONS AND NOTATIONS

(0.1) Throughout this paper, we work over the field of complex numbers \mathbb{C} .

(0.2) For D_1, D_2 \mathbb{Q} -divisors on a variety X , we write $D_1 \prec D_2$ if $D_2 - D_1$ is effective, and $D_1 \equiv D_2$ if D_1 and D_2 are numerically equivalent.

(0.3) $|D|$ will denote the linear series associated to the divisor D , and $Bs|D|$ denotes the base locus of $|D|$.

The first author was partially supported by National Science Council (NSC-87-2119-M-194-007)

1991 *Mathematics Subject Classification.* Primary 14H45, 14H99; Secondary 14H50.

(0.4) For a real number a , let $\lfloor a \rfloor$ be the largest integer $\leq a$ and $\lceil a \rceil$ be the smallest integer $\geq a$. For a \mathbb{Q} -divisor $D = \sum a_i D_i$, let $\lfloor D \rfloor = \sum \lfloor a_i \rfloor D_i$ and $\lceil D \rceil = \sum \lceil a_i \rceil D_i$.

(0.5) Let \mathcal{F} be a coherent sheaf on X , then $h^i(X, \mathcal{F})$ denotes the complex dimension of $H^i(X, \mathcal{F})$. In particular, the plurigenera $h^0(X, \omega_X^{\otimes m})$ are denoted by $P_m(X)$ and the irregularity $h^0(X, \Omega_X^1)$ is denoted by $q(X)$.

(0.6) Let A be an abelian variety, for all positive integers n , we will denote by A_n the set of n -division points of A . For all subgroups T of A , T_0 will be the connected component of T containing the origin. For any line bundle L there is a canonical homomorphism $\phi_L : A \rightarrow \text{Pic}^0(A)$ defined by $x \rightarrow t_x^* L \otimes L^{-1}$. Denote by $K(L)$ the kernel of ϕ_L , and by $p_L : A \rightarrow A(L) := A/K(L)_0$ the induced quotient map.

1. COHOMOLOGICAL SUPPORT LOCI

We start by recalling some results about cohomological support loci that will be needed in this paper. Most of these results may be found in [EL1] and [EV].

Let $f : X \rightarrow A$ be a morphism from a smooth projective variety X to an abelian variety A . If \mathcal{F} is a coherent sheaf on X , then one can define the cohomological support loci by

Definition 1.1.

$$V_i(X, A, \mathcal{F}) := \{P \in \text{Pic}^0(A) \mid h^i(X, \mathcal{F} \otimes f^* P) \neq 0\}.$$

In particular, if $f = \text{alb}_X : X \rightarrow \text{Alb}(X)$, then we simply write

$$V_i(X, \mathcal{F}) := \{P \in \text{Pic}^0(X) \mid h^i(X, \mathcal{F} \otimes P) \neq 0\}.$$

We say that X has maximal Albanese dimension if $\dim(\text{alb}_X(X)) = \dim(X)$. The loci $V_i(X, \omega_X)$ defined above are very useful in the study of irregular varieties and in particular of varieties of maximal Albanese dimension. The geometry of these loci is governed by the following:

Theorem 1.2 ([GL1],[GL2]).

a. Any irreducible component of $V_i(X, \omega_X)$ is a translate of a sub-torus of $\text{Pic}^0(X)$ and is of codimension at least $i - (\dim(X) - \dim(\text{alb}_X(X)))$.

b. Let $P \in S$ be a general point of an irreducible component T of $V_i(X, \omega_X)$. Suppose that $v \in H^1(X, \mathcal{O}_X) \cong T_P \text{Pic}^0(X)$ is not tangent to T . Then the sequence

$$H^{n-i-1}(X, \omega_X \otimes P) \xrightarrow{\cup v} H^{n-i}(X, \omega_X \otimes P) \xrightarrow{\cup v} H^{n-i+1}(X, \omega_X \otimes P)$$

is exact. If v is tangent to T , then the maps in the above sequence vanish.

c. If X is a variety of maximal Albanese dimension, then

$$\text{Pic}^0(X) \supset V_0(X, \omega_X) \supset V_1(X, \omega_X) \supset \dots \supset V_n(X, \omega_X) = \{\mathcal{O}_X\}.$$

Theorem 1.3 ([S]). *Under the above hypothesis, every irreducible component of $V_i(X, \omega_X)$ is a translate of a sub-torus of $\text{Pic}^0(X)$ by a torsion point.*

Remark 1. The results listed above also hold when we consider an abelian variety A and a morphism $f : X \rightarrow A$ with $\dim f(X) = \dim(X)$ and we replace $\text{Pic}^0(X)$ by $\text{Pic}^0(A)$.

Proposition 1.4 [EL1, Proposition 2.1]. *If $P_1(X) = P_2(X) = 1$, then the origin is an isolated point of $V_0(X, \omega_X)$.*

Proposition 1.5 [EL1, Proposition 2.2]. *If the origin is an isolated point of $V_0(X, \omega_X)$, then the Albanese mapping $\text{alb}_X : X \rightarrow \text{Alb}(X)$ is surjective.*

If X has maximal Albanese dimension, then X admits a generically finite map to an abelian variety A . It is easy to see that $P_1(X) \geq 1$ by pulling back sections of Ω_A^n , where $n = \dim(X)$. Assume furthermore that $P_2(X) = 1$, then necessarily $P_1(X) = 1$, and so the Albanese map is surjective. In particular $q(X) = \dim(X)$. Therefore one has:

Corollary 1.6. *The following are equivalent.*

- a. $P_1(X) = P_2(X) = 1$, and $q(X) = \dim(X)$.
- b. $P_2(X) = 1$ and X has maximal Albanese dimension.

Next we illustrate some examples in which the geometry of X can be recovered from information on the loci $V_i(X, \omega_X)$.

Theorem 1.7 ([EL2]). *If X is a variety with maximal Albanese dimension and $\dim V_0(X, \omega_X) = 0$. Then X is birational to an abelian variety.*

The idea of the proof is as follows. There exists a map $f : X \rightarrow A$ to an abelian variety A , such that X is generically finite over its image. We may assume that the map $f : X \rightarrow A$ is surjective, as otherwise by Proposition 1.5, there exists a positive dimensional component of $V_0(X, \omega_X)$. Consider the sheaf $f_*\omega_X$. By Theorem 1.2, the groups $H^i(A, f_*\omega_X \otimes P)$ do not vanish only for finitely many $P \in \text{Pic}^0(A)$. (In fact it is not hard to see [EL2] that $h^i(f_*\omega_X \otimes P) = 0$ for all $P \neq \mathcal{O}_X$.) By [M, Example 2.9 and §3], $c_1(f_*\omega_X) = 0$, and so the map $X \rightarrow A$ is étale in codimension one and hence étale.

Corollary 1.8 ([EL2]). *Let X be a variety of maximal Albanese dimension, and of positive Kodaira dimension ($\kappa(X) > 0$). Then there exists a torsion line bundle $Q \in \text{Pic}^0(X)$ and a positive dimensional subgroup T of $\text{Pic}^0(X)$, such that $h^0(2K_X \otimes 2Q \otimes P) \geq 2$ for all $P \in T$. In particular, if $V_0(X, \omega_X)$ contains a positive dimensional component through the origin \mathcal{O}_X (or through any 2-torsion point $P \in \text{Pic}^0(A)_2$), then $P_2(X) \geq 2$.*

Proof. Let S_Q be any positive dimensional component of $V^0(X, \omega_X)$ containing $Q \in \text{Pic}^0(X)$ and let $S = S_Q - Q$. Since S is a subgroup of $\text{Pic}^0(X)$, one sees that $-S = S$. Consider the natural map

$$H^0(X, \omega_X \otimes Q \otimes P_y) \otimes H^0(X, \omega_X \otimes Q \otimes P_y^*) \rightarrow H^0(X, \omega_X^{\otimes 2} \otimes Q^{\otimes 2})$$

and let P_y vary in S . Both $H^0(X, \omega_X \otimes Q \otimes P_y)$ and $H^0(X, \omega_X \otimes Q \otimes P_y^*)$ are non-zero. The natural map above gives rise to a map of linear series

$$|K_X + Q + P_y| \times |K_X + Q - P_y| \rightarrow |2K_X + 2Q|.$$

Since there are infinitely many $P_y \subset S$, $|2K_X + 2Q|$ cannot consist of only one element. Therefore $h^0(X, \omega_X^{\otimes 2} \otimes Q^{\otimes 2}) \geq 2$. \square

Corollary 1.9 [EL1]. *Let D be a reduced irreducible divisor of an abelian variety A . Then there exists a positive dimensional sub-group $T_D \subset \text{Pic}^0(X)$ such that for any desingularization $\nu : \tilde{D} \rightarrow D$, $h^0(\omega_{\tilde{D}} \otimes \nu^*P) > 0$ for all $P \in T_D$.*

We end this section by recalling a few results that will frequently be used in what follows.

Theorem 1.10 [Ko1, EV2]. *Let $f : X \rightarrow Y$ be a surjective morphism from a smooth projective variety X to a normal variety Y . Let L be a line bundle on X such that $L \equiv f^*M + \Delta$, where M is a \mathbb{Q} -divisor on Y and (X, Δ) is klt. Then*

- a. $R^j f_*(K_X \otimes L)$ is torsion free for $j \geq 0$.
- b. Assume furthermore that M is nef and big. Then $H^i(Y, R^j f_*(K_X \otimes L)) = 0$ for $i > 0$, $j \geq 0$.

Theorem 1.11 [Ko2]. *Let $f : X \rightarrow Y$ be a surjective morphism from a smooth projective variety X to a smooth variety Y , birational to an abelian variety. Let L be a nef and big \mathbb{Q} -divisor on Y , Δ be a \mathbb{Q} -divisor with normal crossings such that $[\Delta] = 0$. Let $U \subset Y$ be a dense open set, and N a Cartier divisor on X , $N \equiv K_X + \Delta + f^*L$. Assume that $N|_{g^{-1}U}$ is linearly equivalent to an effective divisor. Then $h^0(X, N) \neq 0$.*

Theorem 1.12 [Ko2]. *Let $f : X \rightarrow Y$ be a surjective morphism from a smooth projective variety X to a normal variety Y . Let L be a line bundle on X such that $L \equiv f^*M + \Delta$, where M is a \mathbb{Q} -divisor on Y and (X, Δ) is klt. Let D be a effective divisor on X such that $f(D) \neq Y$. Then*

$$H^j(X, K_X \otimes L) \rightarrow H^j(X, K_X \otimes L(D)) \text{ is injective.}$$

Remark. Suppose X admits a generically finite map $a : X \rightarrow A$ to an abelian variety A . And let $f : X \rightarrow Y$ be its Iitaka fibration. Let D be an non-exceptional irreducible component of the ramification divisor with respect to the map $a : X \rightarrow A$. Then $P_1(D) \geq 1$, and $f(D) \neq Y$ [Ko2, 17.6.1].

2. BIRATIONAL INVARIANTS

In this section we study birational invariants of varieties with maximal Albanese dimension. Let $f : X \rightarrow Y$ be a birational model of the Iitaka fibration of X . We may assume that X and Y are smooth projective varieties. Let K be the image of the generic fiber of f via the Albanese map $\text{alb}_X : X \rightarrow \text{Alb}(X)$. K is a sub-group of $\text{Alb}(X)$, and there is a generically finite map $Y \rightarrow \text{Alb}(X)/K$. Thus Y is also of maximal Albanese dimension [Ko2, 17.5.1].

Lemma 2.1. *Suppose that X has maximal Albanese dimension, and positive Kodaira dimension $\kappa(X) > 0$. Fix Q a torsion element of $\text{Pic}^0(X)$. Then $h^0(X, 2K_X \otimes Q \otimes f^*P)$ is constant for all torsion $P \in \text{Pic}^0(Y)$.*

Proof. Fix a nef and big divisor A on Y . The linear series $|mK_X - f^*A|$ is nonempty for all m sufficiently large [Ko2, Lemma 17.6.1]. Fix an appropriate divisor $B_2 \in |m_2K_X - f^*A|$. For an arbitrary integer m_1 , one may consider the map

induced by multiplication by $B_2 \in |m_2 K_X - f^* A|$. Replacing X by an appropriate birational model, we may assume that for all sufficiently big and divisible integers m_1 , $|m_1 K_X| = |M_{m_1}| + F_{m_1}$ and $|m_2 K_X - f^* A| = |M_{m_1, A}| + F_{m_1, A}$, where M_{m_1} and $M_{m_1, A}$ are free and $F_{m_1} + F_{m_1, A}$ is a divisor with normal crossings. In particular we may assume that the divisors $F_{m_1} = Bs|m_1 K_X|$ and $F_{m_1+m_2, A} = Bs|(m_1 + m_2) K_X - f^* A|$ have normal crossing singularities. It follows that

$$\lfloor \frac{Bs|(m_1 + m_2) K_X - f^* A|}{m_1 + m_2} \rfloor \prec \lfloor \frac{Bs|m_1 K_X| + B_2}{m_1 + m_2} \rfloor.$$

We will show next that for all sufficiently big and divisible integers m_1 ,

$$\lfloor \frac{Bs|m_1 K_X| + B_2}{m_1 + m_2} \rfloor \prec \frac{2Bs|m_1 K_X|}{m_1}.$$

We proceed by comparing the multiplicities of each component. One may assume that for all sufficiently big and divisible integers m_1 , the support of $Bs|m_1 K_X|$ and of B_2 , are contained in a fixed divisor D with irreducible components D_i , $i = 1 \dots r$. (Fix for example $m_1 = 2^j m_0$, $j \geq 0$.) We may therefore write $Bs|m_1 K_X| = \sum_{i=1}^r b_{m_1, i} D_i$, and $B_2 = \sum_{i=1}^r \delta_i D_i$. If $b_{m_1, i}$ is bounded, then $\lfloor \frac{b_{m_1, i} + \delta_i}{m_1 + m_2} \rfloor = 0$, and hence is bounded by the positive rational numbers $\frac{2b_{m_1, i}}{m_1}$ for all m_1 sufficiently big and divisible. Conversely, assume that $b_{m_1, i}$ is not bounded. Then for all m_1 sufficiently big and divisible

$$\frac{b_{m_1, i} + \delta_i}{m_1 + m_2} < \frac{2b_{m_1, i}}{m_1 + m_2} < \frac{2b_{m_1, i}}{m_1}.$$

Therefore the desired inequality holds.

Since Q and P are torsion, we may assume that $m_1 = 2m'_1$ is chosen so that $m'_1(f^*P) = m'_1 Q = \mathcal{O}_X$, and therefore

$$\frac{2Bs|m_1 K_X|}{m_1} = \frac{Bs|m'_1(2K_X \otimes Q \otimes f^*P)|}{m'_1} \prec Bs|2K_X \otimes Q \otimes f^*P|.$$

Therefore one can choose a divisor $B \in |(m_1 + m_2) K_X - f^* A|$ such that

$$\lfloor \frac{B}{m_1 + m_2} \rfloor = \lfloor \frac{Bs|(m_1 + m_2) K_X - f^* A|}{m_1 + m_2} \rfloor \prec Bs|2K_X \otimes Q \otimes f^*P|,$$

and such that

$$\lfloor \frac{B}{m_1 + m_2} \rfloor \prec Bs|2K_X \otimes Q|.$$

Now let $m = m_1 + m_2$, and define

$$M := K_X(-\lfloor \frac{B}{m} \rfloor) \equiv f^*\left(\frac{A}{m}\right) + \left\{\frac{B}{m}\right\}.$$

One sees that $h^0(X, K_X + M + Q + f^*P) = h^0(X, 2K_X + Q + f^*P)$, and $h^0(X, K_X + M + Q) = h^0(X, 2K_X + Q)$. Since the support of B has only normal crossing singularities, $\lfloor \frac{B}{m} \rfloor$ is Kawamata log terminal. By Theorem 1.10, it follows that

$H^i(Y, R^j f_*(K_X \otimes M \otimes T) \otimes P) = 0$ for all $i > 0$, $j \geq 0$ and all $T \in \text{Pic}^0(X)$. In particular

$$h^0(Y, f_*(K_X \otimes M \otimes Q) \otimes P) = \chi(Y, f_*(K_X \otimes M \otimes Q) \otimes P),$$

which is deformation invariant. So, the quantity $h^0(X, K_X \otimes M \otimes Q \otimes f^*P)$ is constant for all $P \in \text{Pic}^0(Y)$. Therefore, for any torsion element P of $\text{Pic}^0(Y)$,

$$h^0(X, 2K_X \otimes Q \otimes f^*P) = h^0(X, K_X \otimes M \otimes Q \otimes f^*P) =$$

$$h^0(X, K_X \otimes M \otimes Q) = h^0(X, 2K_X \otimes Q).$$

□

An immediate consequence of Lemma 2.1 is the following

Theorem 2.2. *Let X be a smooth projective variety with maximal Albanese dimension. If X is of general type, then $P_2(X) = h^0(X, \omega_X^{\otimes 2}) \geq 2$.*

Proof. Assume that $P_2(X) = 1$, then by Lemma 1.6, the map $X \rightarrow \text{Alb}(X)$ is surjective and generically finite. Since X is not birational to an abelian variety, by Corollary 1.9, there exists a torsion line bundle $Q \in \text{Pic}^0(X)$, such that $h^0(X, 2K_X + 2Q) \geq 2$. By Lemma 2.1, $h^0(X, 2K_X) \geq 2$, and this is the required contradiction. □

Lemma 2.3. *Keeping the notation as in Lemma 2.1. Assume furthermore that the Albanese map $g : X \rightarrow \text{Alb}(X)$ is surjective and generically finite with ramification divisor R . Let R_0 be any reduced irreducible smooth divisor contained in R with multiplicity $r_0 \geq 2$, then $h^0(X, 2K_X - (r_0 - 1)R_0 \otimes f^*P) \geq 1$ for all $P \in \text{Pic}^0(Y)$. Moreover, if the divisor R_0 is not exceptional with respect to the map $g : X \rightarrow \text{Alb}(X)$, then $P_2(X) \geq 2$.*

Proof. The proof is a technical improvement on a result of Kollár. We closely follow the methods and the notation of [Ko2, §17]. There is no harm in assuming that R_0 be a smooth divisor as this can always be arranged replacing X by an appropriate birational model.

Consider the divisor

$$K' = K_X - (r_0 - 1)R_0,$$

we have the following inclusion of effective divisors

$$K_X \prec r_0 K' \prec r_0 K_X.$$

It follows by [Ko2, 17.6.1] that for any fixed ample divisor A on Y , the linear series $|mK' - f^*A|$ is not empty, for all sufficiently big and divisible m . Let B' be a general member of the above linear series, then define

$$M' := K'(-\lfloor (1/m)B' \rfloor) \equiv f^*\left(\frac{A}{m}\right) + \left\{\frac{B'}{m}\right\}.$$

We may further assume that R_0 is not a component of B' as by [Ko2], one can arrange that R_0 is not contained in the base locus of $|mK_X - f^*A|$. However,

and $r_0K' - K_X$ is an effective divisor not containing R_0 .

By Theorem 1.10, $H^i(Y, f_*(K_X \otimes M') \otimes P) = 0$ for all $i > 0$ and all $P \in \text{Pic}^0(Y)$. Hence the quantity $h^0(f_*(K_X \otimes M') \otimes P)$ is positive and independent of $P \in \text{Pic}^0(Y)$. Therefore $h^0(X, 2K_X - (r_0 - 1)R_0 \otimes f^*P) \geq 1$ for all $P \in \text{Pic}^0(Y)$.

Consider the exact sequence [Ko2, 17.7],

$$0 \longrightarrow K_X \otimes M' \longrightarrow K_X \otimes M'(R_0) \longrightarrow K_{R_0} \otimes M' \longrightarrow 0.$$

By Theorem 1.12, the map $H^1(X, K_X \otimes M') \longrightarrow H^1(X, K_X \otimes M'(R_0))$ is injective. Since, M' and $M'|_{R_0} \equiv f^*A|_{R_0} + \{B/m|_{R_0}\}$ satisfy the assumptions of Theorem 1.11, it follows that the quantities $h^0(K_X \otimes M')$, $h^0(K_{R_0} \otimes M')$ are both positive, and hence $h^0(K_X + M'(R_0)) \geq 2$. The assertion now follows from the map of linear series $|K_X \otimes M'(R_0)| \longrightarrow |2K_X|$ induced by multiplication by the effective divisor $(r_0 - 2)R_0 + \lfloor (1/m)B' \rfloor$. \square

Theorem 2.4. *Let X be a smooth projective variety with maximal Albanese dimension and positive Kodaira dimension $\kappa(X) > 0$. Let $f : X \rightarrow Y$ be the Iitaka fibration. If Y is birational to an abelian variety, then $P_2(X) \geq 2$.*

Proof. Let $q : Y \rightarrow S$ be a birational morphism to an abelian variety S . In what follows, we assume that $h^0(X, 2K_X) = 1$ and derive a contradiction.

Step 1. Since X is not birational to an abelian variety, by Corollary 1.8, we may choose a torsion line bundle Q such that $h^0(X, 2K_X + 2Q) \geq 2$. By Lemma 2.1, $h^0(X, 2K_X + 2Q + P) \geq 2$ for all torsion $P \in \text{Pic}^0(Y)$. It follows by the upper semicontinuity of the dimension of cohomology groups, that $h^0(X, 2K_X + 2Q + P) \geq 2$ for all $P \in \text{Pic}^0(Y)$.

Step 2. Again by Lemma 2.1, since the dimension of cohomology groups is upper semicontinuous, $h^0(X, 2K_X + P) \geq 1$ for all $P \in \text{Pic}^0(Y)$. As $h^0(2K_X) = 1$, equality holds for all P in an appropriate neighborhood $U \subset \text{Pic}^0(Y)$ of the origin \mathcal{O}_Y .

Step 3. There exists an effective divisor G which is contained in the base locus of $|2K_X + f^*P|$ for all $P \in U$.

To this end, let M be a divisor chosen as in the proof of Lemma 2.1 (in particular $\lfloor \frac{B}{m} \rfloor \prec Bs|2K_X|$). We wish to compute the cohomology groups of the direct image sheaf $q_*f_*(K_X + M)$. From the Leray spectral sequence associated to the morphism $q : Y \rightarrow S$, one sees that for all $P \in \text{Pic}^0(Y)$,

$$h^0(X, K_X \otimes M \otimes f^*P) = h^0(S, q_*f_*(K_X + M) \otimes P) = 1,$$

and there is an injection

$$H^i(S, q_*f_*(K_X \otimes M) \otimes P) \hookrightarrow H^i(Y, f_*(K_X \otimes M) \otimes P).$$

It follows that for all $i > 0$, both of the above cohomology groups vanish. So the sheaf $q_*f_*(K_X \otimes M)$ is coherent, generically of rank one which is a "cohomological principal polarization". In other words, for all $P \in \text{Pic}^0(S)$, the cohomology groups of $q_*f_*(K_X \otimes M) \otimes P$ are of the same dimension as the cohomology groups of a principal polarization of S . By [H, Proposition 2.2], there exists an appropriate theta-divisor Θ on S and an isomorphism of sheaves $q_*f_*(K_X \otimes M) \cong \mathcal{O}_S(\Theta)$.

Pulling back with respect to q and f , for all $P \in \text{Pic}^0(Y)$, we get a map of linear series

$$f^*q^*|\mathcal{O}_S(\Theta) \otimes P| \xrightarrow{\times G} |2K_X \otimes f^*P|$$

which factors through $|K_X \otimes M \otimes f^*P|$. Since for $P \in U$, the groups $H^0(X, 2K_X \otimes f^*P)$ are one dimensional, the above map of linear series is an isomorphism, and so G is contained in $Bs|2K_X \otimes f^*P|$ for all $P \in U \subset \text{Pic}^0(Y)$.

Step 4. We wish to show that the divisor G is non-zero and not exceptional with respect to the map $X \rightarrow S$.

Let F be a maximal effective divisor such that for all $P \in \text{Pic}^0(Y)$, $2K_X \otimes 2Q \otimes f^*P$ is linearly equivalent to $F + V_P$, and $h^0(V_P) \geq 2$. We may assume that D_P, D'_P are two distinct general members of $|V_P|$, which do not contain any components of the ramification divisor of $X \rightarrow Y$. Therefore there exist divisors Δ_P, Δ'_P on Y such that $D_P = f^*(\Delta_P)$ and $D'_P = f^*(\Delta'_P)$. The map $Y \rightarrow S$ is birational, so there exist effective divisors $\bar{\Delta}_P$ and $\bar{\Delta}'_P$ on S and arbitrary q -exceptional divisors E_P, E'_P such that

$$\Delta_P = q^*(\bar{\Delta}_P) + E_P, \quad \Delta'_P = q^*(\bar{\Delta}'_P) + E'_P.$$

If $h^0(S, \mathcal{O}_S(\bar{\Delta}_P)) = 1$, then $\bar{\Delta}_P = \bar{\Delta}'_P$ and similarly since $Y \rightarrow S$ is birational, $\Delta_P = \Delta'_P$, which is a contradiction. So $\bar{\Delta}_P$ is a divisor on S with at least two sections. We may also assume that $\bar{\Delta}_P$ is an ample divisor on S , since otherwise there would exist a non trivial map of abelian varieties $S \rightarrow \Sigma$ such that $\bar{\Delta}_P$ is the pull back of a divisor on Σ . However then for general $P_1, P_2 \in \text{Pic}^0(S)$, $V_{P_1} - V_{P_2}$ is rationally equivalent to $P_1 - P_2$ and is an element of $\text{Pic}^0(\Sigma)$. The only way this can occur is if $S = \Sigma$ as required.

Hence, for H' the pull back of an ample divisor on $A = \text{Alb}(X)$, and H the pull back of an ample divisor on S , we have the numerical inequality

$$2K_X \cdot H^{s-1} \cdot (H')^{n-s} > (f^*q^*\Theta) \cdot H^{s-1} \cdot (H')^{n-s}.$$

From the injection of linear series

$$f^*q^*|\mathcal{O}_S(\Theta) \otimes P| \xrightarrow{\times G} |2K_X \otimes f^*P|,$$

one sees that the effective divisor G is not exceptional with respect to the map $X \rightarrow S$.

Step 5. Consider now G_0 an irreducible reduced component of G which is not exceptional with respect to the map $X \rightarrow S$. Since we assumed that $h^0(X, 2K_X) = 1$, by Proposition 1.4, $h^0(X, K_X \otimes P) = 0$ for $P \in (\text{Pic}^0(Y) - \mathcal{O}_Y)$ in an appropriate neighborhood U' of the origin \mathcal{O}_S . We may assume that $U = U'$. From the exact sequence of sheaves

$$0 \rightarrow K_X \otimes P \rightarrow K_X + G_0 \otimes P \rightarrow K_{G_0} \otimes P \rightarrow 0,$$

we see that for $P \in U^0 = U - \{\mathcal{O}_Y\}$, there is an isomorphism

$$H^0(X, K_X + G_0 \otimes P) \cong H^0(G_0, K_{G_0} \otimes P).$$

By Corollary 1.9, there exists a positive dimensional subgroup $T_{G_0} \subset \text{Pic}^0(X)$ such that for all $P \in T_{G_0}$, $h^0(X, K_X \otimes P) > 0$. Since $G_0 \subset 2K_X$, G_0 is vertical with

respect to the Iitaka fibration $X \rightarrow Y$. We may therefore assume that T_{G_0} is in fact contained in $\text{Pic}^0(Y)$. It follows that for $P \in T_{G_0} \cap U^0$, G_0 is not contained in the base locus of the linear series $|K_X + G_0 \otimes P|$. Let

$$|K_X + G_0 \otimes P| \xrightarrow{K_X - G_0} |2K_X \otimes P|$$

be the map of linear series induced by multiplication by the effective divisor $K_X - G_0$. Since G_0 is in the base locus of $|2K_X \otimes P|$, G_0 must be an irreducible component of $K_X - G_0$. In other words the multiplicity g_0 of G_0 in $R = K_X$ is at least 2. By Lemma 2.3, this is the required contradiction. \square

3. PROOF OF THE MAIN THEOREM

Lemma 3.1. *Let X and Y be smooth varieties and let A and C be abelian varieties. Suppose that there is a commutative diagram*

$$\begin{array}{ccc} X & \xrightarrow{g} & A \\ f \downarrow & & \downarrow \pi \\ Y & \xrightarrow{q} & C, \end{array}$$

such that both g and q are generically finite and surjective. Then $P_m(Y) \leq P_m(X)$, for all $m \geq 1$.

Proof. Let $\{\bar{z}_1, \dots, \bar{z}_k\}$ be coordinates on C such that $\{z_i = \bar{z}_i \circ \pi \mid 1 \leq i \leq k\}$ may be extended to coordinates $\{z_1, \dots, z_n\}$ on A , where n and k are the dimensions of A and of C respectively. Let $\Lambda = g^* dz_{k+1} \wedge \dots \wedge g^* dz_n \in H^0(X, \Omega_X^{n-k})$, we claim that

$$(f^* \omega_Y)^{\otimes m} \xrightarrow{\wedge \Lambda^{\otimes m}} \omega_X^{\otimes m}$$

induces the desired injective map

$$\Lambda_m : H^0(Y, \omega_Y^{\otimes m}) \rightarrow H^0(X, \omega_X^{\otimes m}).$$

Since $q : Y \rightarrow C$ is generically finite and surjective, for a general point $y \in Y$, there are isomorphic Euclidean open neighborhoods \mathcal{U}_y and $\mathcal{V}_{q(y)}$ of y and $q(y)$ respectively. For a non-zero section $\eta \in H^0(Y, \omega_Y^{\otimes m})$, we may assume that $\eta(y) \neq 0$. On \mathcal{U}_y , η is represented by $\zeta \cdot (q^* d\bar{z}_1 \wedge \dots \wedge q^* d\bar{z}_k)^m$, with $\zeta(y) \neq 0$. On $f^{-1}(\mathcal{U}_y)$

$$\Lambda_m(\eta)|_{f^{-1}(\mathcal{U}_y)} = f^* \zeta \cdot (g^* dz_1 \wedge \dots \wedge g^* dz_n)^m \neq 0.$$

Therefore (by choosing harmonic representatives, see [GL2, Lemma 3.1]), $\Lambda_m : H^0(Y, \omega_Y^{\otimes m}) \rightarrow H^0(X, \omega_X^{\otimes m})$ is injective and so $P_m(Y) \leq P_m(X)$. \square

Theorem 3.2. *Let X be a smooth projective variety with $P_1(X) = P_2(X) = 1$ and $q(X) = \dim(X)$. Then X is birational to an abelian variety.*

Proof. Since $P_1(X) = P_2(X)$ and $q(X) = \dim(X)$, by Proposition 1.5, it follows that the Albanese morphism is surjective, and hence a generically finite map.

If the Kodaira dimension of X is zero, then by Kawamata's theorem [Ka], X is birational to an abelian variety.

Suppose that $\kappa(X) > 0$. By Theorem 2.2, the variety X cannot be of general type. Therefore X admits a nontrivial Iitaka fibration. Let $f : X \rightarrow Y$ be a birational model of the Iitaka fibration. The generic fiber F has $\kappa(F) = 0$ and $\dim(\text{alb}_X(F)) = \dim(F)$. Thus $\text{alb}_X(F)$ is an abelian sub-variety which we denote by K . We obtain a diagram

$$\begin{array}{ccc} X & \xrightarrow{\text{alb}_X} & \text{Alb}(X) \\ f \downarrow & & \downarrow \pi \\ Y & \xrightarrow{g} & \text{Alb}(X)/K \end{array}$$

where the morphism $g : Y \rightarrow \text{Alb}(X)/K$ is generically finite (cf [Ko2, 17.5.1]). By Lemma 3.1, we have $P_1(Y) = P_2(Y) = 1$. By iterating this procedure, we eventually produce a variety Y' either of maximal Kodaira dimension $\kappa(Y') = \dim(Y')$, or with Kodaira dimension 0 (and hence birational to an abelian variety). By applying Theorem 2.2 or Theorem 2.4, respectively, in conjunction with Lemma 3.1, we conclude that $P_2(X) \geq 2$. \square

REFERENCES

- [EL1] L. Ein, R. Lazarsfeld, *Singularities of theta divisors, and birational geometry of irregular varieties*, Jour. AMS **10** (1997), no. 1, 243–258.
- [EL2] L. Ein, R. Lazarsfeld, *Unpublished personal communication*.
- [EV] H. Esnault, E. Viehweg, *Lectures on Vanishing Theorems DMV 20*, Birkhauser, 1992.
- [EV2] H. Esnault, E. Viehweg, *Revêtements cyclique II*, Géométrie Algébrique et Applications II, La Rábida. Herman, Paris (1987), 81–94.
- [H] C. D. Hacon, *Fourier transforms, generic vanishing theorems and polarizations of abelian varieties*, math.AG/9902078 (1999).
- [GL1] M. Green, R. Lazarsfeld, *Deformation theory, generic vanishing theorems, and some conjectures of Enriques, Catanese and Beauville*, Invent. Math. **90** (1987), 389–407.
- [GL2] ———, *Higher obstruction to deforming cohomology groups of line bundles*, Jour. Amer. Math. Soc. **4** (1991), 87–103.
- [Ka] Y. Kawamata, *Characterization of abelian varieties*, Comp. Math. **43** (1981), 253–276.
- [Ko1] J. Kollár, *Higher direct images of dualizing sheaves I*, Ann. Math. **123** (1986), 11–42.
- [Ko2] ———, *Shafarevich Maps and Automorphic Forms*, Princeton University Press, 1995.
- [M] S. Mukai, *Duality between $D(X)$ and $D(\hat{X})$, with application to Picard sheaves*, Nagoya math. J. **81** (1981), 153–175.
- [S] C. Simpson, *Subspaces of moduli spaces of rank one local systems*, Ann. Sci. École Norm. Sup. (4) **26** (1993), no. 3, 361–401.

JUNGKAI ALFRED CHEN, DEPARTMENT OF MATHEMATICS, NATIONAL CHUNG CHENG UNIVERSITY, MING HSIUNG, CHIA YI, 621, TAIWAN
E-mail address: jkchen@math.ccu.edu.tw

CHRISTOPHER DEREK HACON, UTAH, DEPARTMENT OF MATHEMATICS, SALT LAKE CITY, UT 84112, USA
E-mail address: chhacon@math.utah.edu