

Cross Product Bialgebras

Part II

Yuri Bespalov

Bernhard Drabant

July 1998/March 1999

Abstract

This is the central article of a series of three papers on cross product bialgebras. We present a universal theory of bialgebra factorizations (or cross product bialgebras) with cocycles and dual cocycles. We also provide an equivalent (co-)modular co-cyclic formulation. All known examples as for instance bi- or smash, doublecross and bicross product bialgebras as well as double biproduct bialgebras and bicrossed or cocycle bicross product bialgebras are now united within a single theory. Furthermore our construction yields various novel types of cross product bialgebras.

1991 Mathematics Subject Classification: 16W30, 16S40, 18D10, 16B50, 81R50

1 Introduction

By definition cross product bialgebras are isomorphic to factorizations $B = B_1 \otimes B_2$, in a braided category say, such that the given isomorphisms can be characterized universally in terms of certain projections and injections from the bialgebra into the particular tensor factors. In this context tensor product bialgebras, bi- or bismash product bialgebras [24, 30], doublecross product bialgebras, bicross product bialgebras and double biproduct bialgebras [18, 21] are cross product bialgebras. But also the bicrossed or cocycle bicross product bialgebras [22, 20] are cross product bialgebras in our terminology; their universal characterization is given in terms of cleft extensions and coextensions (see [7, 12, 23, 27] for a comprehensive study of cleft or normal basis Hopf-Galois extensions on the algebra level). The notion of cross product bialgebras therefore as well includes constructions involving cocycles and dual cocycles (or simply “cycles”). Well known examples of cross product bialgebras are Drinfel’d’s quantum double, the quantum Poincaré group, Radford’s 4-parameter Hopf algebra, Lusztig’s construction of the quantum enveloping algebra, the quantum Weyl group, the affine quantum groups $U_q(\hat{\mathfrak{g}})$, the Connes-Moscovici Hopf algebra in transverse differential geometry [10], etc. The universal constructions of cross product bialgebras in [24, 18, 20, 22, 30] additionally admit an equivalent characterization in terms of mutual weak (co-)modular, co-cyclic relations of the tensor factors B_1 and B_2 . This means that there exist certain weak (co-)multiplications, (co-)cycles and (co-)actions which are morphisms with tensor rank $(2, 1)$ or $(1, 2)$ defined on B_1 and B_2 and their two-fold tensor products, such that the multiplication m_B and the comultiplication Δ_B of the bialgebra B can be composed monoidally by these structure morphisms. For example we

encounter a weak left coaction $\nu_l : B_1 \rightarrow B_2 \otimes B_1$ which has tensor rank $(1, 2)$, a weak cocycle $\sigma : B_2 \otimes B_2 \rightarrow B_1$ with rank $(2, 1)$, etc. All these structure morphisms are subject to certain relations with rank $(2, 2)$, $(3, 1)$ and $(1, 3)$ which on the one hand follow from the bialgebra structure of $B = B_1 \otimes B_2$ and on the other hand determine the bialgebra structure of $B_1 \otimes B_2$ uniquely.

Despite those common properties of the cross product bialgebra constructions in [24, 18, 20, 22, 30] no theory exists which characterizes all of them as special versions of a single universal construction which in turn is uniquely determined by its (co-)modular co-cyclic structure. A first step towards this objective has been achieved in [5] where we discovered a method to describe cross product bialgebras without co-cycles, generalizing and uniting [18, 21, 24].

The present article considerably extends the results of [5]. The main outcome is a universal and (co-)modular, co-cyclic theory of cross product bialgebras with cocycles and cycles which unites all known constructions within a single setting and provides several new families of cross product bialgebras. This is the most comprising framework for cross product bialgebras so far, which equivalently takes into account both universal and (co-)modular co-cyclic aspects. Furthermore our construction is designed to work in arbitrary braided categories¹.

To derive these results we will proceed in two steps. We begin with the general definition of cross product bialgebras and find a universal characterization for them. We show that so-called cocycle cross product bialgebras possess a certain (co-)modular co-cyclic structure. Then we restrict our consideration to strong cross product bialgebras. They also admit a universal characterization and will turn out to be the central objects in our theory of cross product bialgebras.

In a second step we present a (co-)modular co-cyclic construction method by so-called Hopf data². A Hopf datum consists of a pair of objects (B_1, B_2) and weak (co-)actions, (co-)multiplications and co-cycles defined on B_1 and B_2 which obey certain interrelated identities with rank $(2, 2)$, $(3, 1)$ and $(1, 3)$. Then we introduce strong Hopf data for which additional “strong” relations hold. We show that strong Hopf data induce the structure of a strong cross product bialgebra on the tensor product $B_1 \otimes B_2$. Strong Hopf data and strong cross product bialgebras are different depictions of the same object.

Then we combine the universal characterization and the (co-)modular co-cyclic description in terms of strong Hopf data to obtain the theory of strong cross product bialgebras. This is the central result of the article. Eventually we apply our construction and investigate strong cross product bialgebras according to their (co-)modular co-cyclic structure. In particular we recover all known constructions [24, 18, 22, 20, 5] and find various new types of cross product bialgebras. All of them are special versions of the most general strong cross product bialgebra construction.

After introducing preliminary notations and definitions we study cross product algebras

¹Therefore we have been able to apply the results of [5] to categories of Hopf bimodules and Yetter-Drinfel'd modules [6]; we characterized two-sided bi- or bismash product bialgebras as certain cross product Hopf bimodule bialgebras. In particular we showed that the double biproduct is a twisted Hopf bimodule tensor product bialgebra.

²Observe that Hopf data without co-cycles have been introduced in [5] already. There should be no confusion with the more general notation of the same name which will be defined in the present article. In a very similar context the term “Hopf datum” has been used in [1] for special cases of bicrossed product bialgebras [20].

and cross product coalgebras from a universal point of view in Section 2. These results will be used in Section 3 to define cross product bialgebras. From this general definition we extract cocycle cross product bialgebras and strong cross product bialgebras. The structure of cocycle cross product bialgebras gives useful hints for the design of Hopf data. This will be done in Section 4 where we investigate the connection of Hopf data and cocycle cross product bialgebras. A Hopf datum is canonically assigned to every cocycle cross product bialgebra. The converse is not true in general. But Theorem 4.4 states that strong Hopf data yield strong cross product bialgebras. More precisely strong Hopf data and strong cross product bialgebras are equivalent constructions which are in one-to-one correspondence. The universal (co-)modular co-cyclic theory of strong cross product bialgebras culminates in the central Theorems 4.4 and 4.5. Finally we present a table of special strong cross product bialgebras which contains all known and several new types of cross product bialgebras embedded in our most general framework.

The rather intricate proof of Theorem 4.4 has been postponed to Section 5.

In the recent articles [9, 11, 26] cross product bialgebras have been studied as well. In [9] the universal properties of co-cycle free cross product bialgebras have been considered. The article [11] investigates certain cross product bialgebras without co-cycles in universal and (co-)modular co-cyclic terms. In [26] a special type of our cocycle cross product bialgebras with one cocycle has been studied in the special case where all objects are vector spaces over a field.

Preliminaries

We are working throughout in (strict) braided monoidal categories [15]. In our article we denote categories by calligraphic letters \mathcal{C} , \mathcal{D} , etc. For a braided monoidal category \mathcal{C} the tensor product is denoted by $\otimes_{\mathcal{C}}$, the unit object by $\mathbb{1}_{\mathcal{C}}$, and the braiding by ${}^{\mathcal{C}}\Psi$. If it is clear from the context we omit the index ‘ \mathcal{C} ’ at the various symbols. We confine ourselves to braided categories which admit split idempotents [3, 17]; for each idempotent $\Pi = \Pi^2 : M \rightarrow M$ of any object M in \mathcal{C} there exists an object M_{Π} and a pair of morphisms (i_{Π}, p_{Π}) such that $p_{\Pi} \circ i_{\Pi} = \text{id}_{M_{\Pi}}$ and $i_{\Pi} \circ p_{\Pi} = \Pi$. This is not a severe restriction of the categories since every braided category can be canonically embedded into a braided category which admits split idempotents [3, 17].

We use generalized algebraic structures like algebra, bialgebra, module, comodule, etc. in such categories. We assume the reader is familiar with these generalizations. A thorough introduction to braided algebraic structures and the graphical calculus coming along with it can be found in [13, 15, 16, 17, 19, 25, 31]. We denote by $m : A \otimes A \rightarrow A$ the multiplication and by $\eta : \mathbb{1} \rightarrow A$ the unit of an algebra A in \mathcal{C} . $\Delta : C \rightarrow C \otimes C$ is meant for the comultiplication and $\varepsilon : C \rightarrow \mathbb{1}$ for the counit of a coalgebra C in \mathcal{C} , $\mu_l : A \otimes M \rightarrow M$ is the left action of an algebra A on a module M , and $\nu_l : N \rightarrow C \otimes N$ denotes the left coaction of a coalgebra C on a comodule N . Right actions are denoted by μ_r and right coactions by ν_r . In the course of our work we often encounter weak versions of (co-)multiplication and (co-)action which do not necessarily obey the relations of ‘true’ (co-)algebras and (co-)modules. We will nevertheless use the same symbols as for proper (co-)multiplications and (co-)actions – it should become clear from the context whether these structure morphisms are weak or not.

In our article we make use of graphical calculus for (strict) braided monoidal categories which simplifies intricate categorical equations of morphisms and helps to uncover their

$$\begin{aligned}
\mathfrak{m} &= \begin{array}{c} \cup \\ | \\ \cup \end{array} , & \eta &= \begin{array}{c} \circ \\ | \\ \circ \end{array} , & \Delta &= \begin{array}{c} | \\ \cup \\ | \end{array} , & \varepsilon &= \begin{array}{c} | \\ \circ \\ | \end{array} , \\
\mu_l &= \begin{array}{c} \cup \\ | \\ \cup \end{array} , & \mu_r &= \begin{array}{c} \cup \\ | \\ \cup \end{array} , & \nu_l &= \begin{array}{c} | \\ \cup \\ | \end{array} , & \nu_r &= \begin{array}{c} | \\ \cup \\ | \end{array} , \\
\Psi &= \begin{array}{c} \diagdown \\ \diagup \end{array} , & \Psi^{-1} &= \begin{array}{c} \diagup \\ \diagdown \end{array} .
\end{aligned}$$

Figure 1: Graphical presentation of (weak) multiplication \mathfrak{m} , unit η , (weak) comultiplication Δ , counit ε , (weak) left action μ_l , (weak) right action μ_r , (weak) left coaction ν_l , (weak) right coaction ν_r , braiding Ψ , and inverse braiding Ψ^{-1} .

intrinsic structure. Morphisms will be composed from up to down, i. e. the domains of the morphisms are at the top and the codomains are at the bottom of the graphics. Tensor products are represented by horizontal concatenation in the corresponding order. We present our own conventions [3, 4, 5] in Figure 1. If there is no fear of confusion we omit the assignment of a specific object to the ends of the particular strings in the graphics. Below we elucidate the graphical calculus to readers unfamiliar with this technique. The first example is the associativity of the action of an algebra A on a right A -module M . The second is the naturality of the braiding in a braided category.

$$\begin{aligned}
\mu_r \circ (\mu_r \otimes \text{id}_A) &= \mu_r \circ (\text{id}_M \otimes \mathfrak{m}_A) && \text{corresponds to} && \begin{array}{c} \cup \\ | \\ \cup \end{array} = \begin{array}{c} \cup \\ | \\ \cup \end{array} , \\
\Psi \circ (f \otimes g) &= (g \otimes f) \circ \Psi && \text{corresponds to} && \begin{array}{c} \circ \quad \circ \\ \diagdown \quad \diagup \\ \circ \quad \circ \end{array} = \begin{array}{c} \circ \quad \circ \\ \diagup \quad \diagdown \\ \circ \quad \circ \end{array} .
\end{aligned}$$

Note that throughout the article there is (almost) no need to require invertibility of the braiding. Therefore most of the results can be derived if we assume that the underlying category \mathcal{C} is pre-braided. We do not discuss these generalizations further and confine to braided categories in what follows.

2 Cross Product Algebras and Coalgebras

In the first part of Section 2 we study cross product algebras. They have been considered also in [8]. It turns out that cross product algebras are universal constructions. They generalize crossed product algebras [12, 23, 27]. Cross product coalgebras will be studied in the second part of Section 2. Since the results for cross product coalgebras can be obtained easily by certain categorical dualization, we omit all the proofs in this case and refer to the analogous proofs for cross product algebras. Both structures, cross product algebras and cross product coalgebras, will be needed later in the definition of cross product bialgebras.

Cross Product Algebras

DEFINITION 2.1 *Let $(B_1, \mathfrak{m}_1, \eta_1)$ be an algebra and B_2 be an object in \mathcal{C} . Suppose there are morphisms $\eta_2 : \mathbb{1} \rightarrow B_2$, $\varphi_{2,1} : B_2 \otimes B_1 \rightarrow B_1 \otimes B_2$, and $\hat{\sigma} : B_2 \otimes B_2 \rightarrow B_1 \otimes B_2$ such that the relation*

$$\varphi_{2,1} = (\mathfrak{m}_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \hat{\sigma}) \circ (\varphi_{2,1} \otimes \eta_2) \tag{2.1}$$

holds. Suppose further that $B = B_1 \otimes B_2$ is an algebra through

$$m_B = (m_1 \otimes \text{id}_{B_2}) \circ (m_1 \otimes \hat{\sigma}) \circ (\text{id}_{B_1} \otimes \varphi_{2,1} \otimes \text{id}_{B_2}), \quad \eta_B = \eta_1 \otimes \eta_2 \quad (2.2)$$

then B is called cross product algebra and will be denoted by $B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$.

In the subsequent proposition we find equivalent constructive conditions for cross product algebras. Similar results have been obtained in [8].

PROPOSITION 2.2 *The following statements are equivalent.*

1. $B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ is a cross product algebra.
2. (B_1, m_1, η_1) is an algebra and B_2 is an object in \mathcal{C} . There exist morphisms $\eta_2 : \mathbb{1} \rightarrow B_2$, $\varphi_{2,1} : B_2 \otimes B_1 \rightarrow B_1 \otimes B_2$ and $\hat{\sigma} : B_2 \otimes B_2 \rightarrow B_1 \otimes B_2$ for which the subsequent identities hold.

$$\begin{aligned} \varphi_{2,1} \circ (\eta_2 \otimes \text{id}_{B_1}) &= \text{id}_{B_1} \otimes \eta_2, & \hat{\sigma} \circ (\eta_2 \otimes \text{id}_{B_2}) &= \eta_1 \otimes \text{id}_{B_2}, \\ \varphi_{2,1} \circ (\text{id}_{B_2} \otimes \eta_1) &= \eta_1 \otimes \text{id}_{B_2}, & \hat{\sigma} \circ (\text{id}_{B_2} \otimes \eta_2) &= \eta_1 \otimes \text{id}_{B_2}, \\ \varphi_{2,1} \circ (\text{id}_{B_2} \otimes m_1) &= (m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \varphi_{2,1}) \circ (\varphi_{2,1} \otimes \text{id}_{B_1}), \\ (m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \hat{\sigma}) \circ (\varphi_{2,1} \otimes \text{id}_{B_2}) \circ (\text{id}_{B_2} \otimes \hat{\sigma}) & & (2.3) \\ &= (m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \hat{\sigma}) \circ (\hat{\sigma} \otimes \text{id}_{B_2}), \\ (m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \hat{\sigma}) \circ (\varphi_{2,1} \otimes \text{id}_{B_2}) \circ (\text{id}_{B_2} \otimes \varphi_{2,1}) & & \\ &= (m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \varphi_{2,1}) \circ (\hat{\sigma} \otimes \text{id}_{B_1}). \end{aligned}$$

PROOF. Suppose that $B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ is a cross product algebra. Since by assumption (B_1, m_1, η_1) is itself an algebra one obtains the first and second identity of (2.3) by unitality of m_B and m_1 . Using these relations yields the fourth identity of (2.3) by application of $\text{id}_{B_2} \otimes \eta_1$ to (2.1). Again using the left unitality of m_B one concludes that $(m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \hat{\sigma} \circ (\eta_2 \otimes \text{id}_{B_2})) = \text{id}_{B_1} \otimes \text{id}_{B_2}$ from which one immediately derives the third identity of (2.3). Now the associativity $m_B \circ (m_B \otimes \text{id}_B) = m_B \circ (\text{id}_B \otimes m_B)$ yields the fifth identity of (2.3) by application of $\eta_1 \otimes \text{id} \otimes \text{id} \otimes \eta_1 \otimes \text{id} \otimes \eta_2$. The sixth relation is obtained by applying $\eta_1 \otimes \text{id} \otimes \eta_1 \otimes \text{id} \otimes \eta_1 \otimes \text{id}$. To derive the seventh identity one has to apply $\eta_1 \otimes \text{id} \otimes \eta_1 \otimes \text{id} \otimes \text{id} \otimes \eta_2$. Conversely suppose the conditions of the second item of the proposition are fulfilled. Define m_B according to (2.2). Then $m_B \circ (m_B \otimes \text{id}_B) = (m_1^{(5)} \otimes \text{id}) \circ (\text{id}^{(4)} \otimes \hat{\sigma}) \circ (\text{id}^{(3)} \otimes \varphi_{2,1} \otimes \text{id}) \circ (\text{id}^{(2)} \otimes \hat{\sigma} \otimes \text{id}^{(2)}) \circ (\text{id} \otimes \varphi_{2,1} \otimes \text{id}^{(3)})$ where $m_1^{(n)} : B_1^{\otimes n} \rightarrow B_1$ is the canonical n -fold multiplication, and $\text{id}^{(n)}$ is the abbreviation of the identity of an n -fold tensor product of (combined) B_1 's and B_2 's. On the other hand

$$\begin{aligned} m_B \circ (\text{id}_B \otimes m_B) &= (m_1^{(4)} \otimes \text{id}) \circ (\text{id}^{(4)} \otimes \hat{\sigma}) \circ (\text{id}^{(3)} \otimes \varphi_{2,1} \otimes \text{id}) \circ \\ &\quad \circ (\text{id}^{(2)} \otimes \varphi_{2,1} \otimes \hat{\sigma}) \circ (\text{id} \otimes \varphi_{2,1} \otimes \varphi_{2,1} \otimes \text{id}) \\ &= (m_1^{(5)} \otimes \text{id}) \circ (\text{id}^{(4)} \otimes \hat{\sigma}) \circ (\text{id}^{(3)} \otimes \hat{\sigma} \otimes \text{id}) \circ \\ &\quad \circ (\text{id}^{(2)} \otimes \varphi_{2,1} \otimes \text{id}^{(2)}) \circ (\text{id} \otimes \varphi_{2,1} \otimes \varphi_{2,1} \otimes \text{id}) \\ &= (m_1^{(5)} \otimes \text{id}) \circ (\text{id}^{(4)} \otimes \hat{\sigma}) \circ (\text{id}^{(3)} \otimes \varphi_{2,1} \otimes \text{id}) \circ \\ &\quad \circ (\text{id}^{(2)} \otimes \hat{\sigma} \otimes \text{id}^{(2)}) \circ (\text{id} \otimes \varphi_{2,1} \otimes \text{id}^{(3)}) \end{aligned}$$

where the fourth condition of (2.3) has been used twice to obtain the first equation, the sixth relation of (2.3) has been applied in the second equation, and with the help of the seventh relation of (2.3) we obtained the third equation. Hence associativity of m_B has been proven. Using the first four identities of (2.3) one easily proves (2.1) and unitality $m_B \circ (\eta_B \otimes \text{id}_B) = \text{id}_B = m_B \circ (\text{id}_B \otimes \eta_B)$. \blacksquare

REMARK 1 Condition (2.1) in Definition 2.1 can be replaced equivalently by the identity $(\text{id}_{B_1} \otimes \hat{\sigma}) = (m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \hat{\sigma}) \circ (\varphi_{2,1} \circ (\text{id}_{B_2} \otimes \eta_1) \otimes \text{id}_{B_2})$.

REMARK 2 There is another equivalent definition of cross product algebras which has been kindly reported to us by Gigel Militaru. Observe that the morphism Γ below is our morphism $m_{0,20}$ in (5.22). Then the following statements are equivalent.

1. $B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ is a cross product algebra.
2. (B_1, m_1, η_1) is an algebra and B_2 is an object in \mathcal{C} . There are morphisms $\eta_2 : \mathbb{1} \rightarrow B_2$ and $\Gamma : B_2 \otimes B_1 \otimes B_2 \rightarrow B_1 \otimes B_2$ such that $B = B_1 \otimes B_2$ is an algebra through

$$m_B = (m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \Gamma) \quad \text{and} \quad \eta_B = \eta_1 \otimes \eta_2. \quad (2.4)$$

3. (B_1, m_1, η_1) is an algebra and B_2 is an object in \mathcal{C} . There exist morphisms $\eta_2 : \mathbb{1} \rightarrow B_2$ and $\Gamma : B_2 \otimes B_1 \otimes B_2 \rightarrow B_1 \otimes B_2$ such that

$$\begin{aligned} \Gamma \circ (\eta_2 \otimes \text{id}_{B_1 \otimes B_2}) &= \text{id}_{B_1 \otimes B_2}, \quad \Gamma \circ (\text{id}_{B_2} \otimes \eta_1 \otimes \eta_2) = \eta_1 \otimes \text{id}_{B_2}, \\ (m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \Gamma) \circ (\Gamma \otimes \text{id}_{B_1 \otimes B_2}) & \\ = \Gamma \circ (\text{id}_{B_2} \otimes (m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \Gamma)) & \end{aligned} \quad (2.5)$$

The one-to-one correspondence is given by $\Gamma = (m_1 \otimes \text{id}_{B_2}) \circ (\text{id}_{B_1} \otimes \hat{\sigma}) \circ (\varphi_{2,1} \otimes \text{id}_{B_2})$ and its inverse $\varphi_{2,1} = \Gamma \circ (\text{id}_{B_2 \otimes B_1} \otimes \eta_2)$ and $\hat{\sigma} = \Gamma \circ (\text{id}_{B_2} \otimes \eta_1 \otimes \text{id}_{B_2})$.

We will show that cross product algebras are universal constructions. The first proposition describes equivalent projection and injection conditions for algebras isomorphic to cross product algebras (see also [5] for the cocycle-free case). The second proposition is closely related to the first one and characterizes the universal construction explicitly.

PROPOSITION 2.3 *Let A be an algebra in \mathcal{C} . Then it holds equivalently*

1. A is algebra isomorphic to a cross product algebra $B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$.
2. There is an algebra (B_1, m_1, η_1) , an object B_2 , morphisms $B_1 \xrightarrow{i_1} A \xleftarrow{i_2} B_2$ and $\eta_2 : \mathbb{1} \rightarrow B_2$ where i_1 is an algebra morphism and $i_2 \circ \eta_2 = \eta_A$ such that $m_A \circ (i_1 \otimes i_2) : B_1 \otimes B_2 \rightarrow A$ is an isomorphism in \mathcal{C} .

PROOF. If $\Lambda : B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2 \rightarrow A$ is an algebra isomorphism then define $i_1 := \Lambda \circ (\text{id}_{B_1} \otimes \eta_2)$ and $i_2 := \Lambda \circ (\eta_1 \otimes \text{id}_{B_2})$. Using the particular definition of $m_{B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2}$ in (2.2) and the identities (2.1), it is verified immediately that i_1 is an algebra morphism since Λ is an algebra morphism. Similarly one proves that $i_2 \circ \eta_2 = \eta_A$ and $m_A \circ (i_1 \otimes i_2) = \Lambda$ which is therefore an isomorphism. If on the other hand the conditions of the second statement of Proposition 2.3 is fulfilled, then $\Lambda := m_A \otimes (i_1 \otimes i_2)$ is an isomorphism by assumption. Therefore $B = B_1 \otimes B_2$ is canonically an algebra through $m_B := \Lambda^{-1} \circ m_A \circ (\Lambda \otimes \Lambda)$ and $\eta_B := \Lambda^{-1} \circ \eta_A$. We will show that this defines a cross product algebra structure on B by $\varphi_{2,1} := \Lambda^{-1} \circ m_A \circ (i_2 \otimes i_1)$ and $\hat{\sigma} := \Lambda^{-1} \circ m_A \circ (i_2 \otimes i_2)$. Using the explicit expression for Λ , inserting several times $\Lambda \circ \Lambda^{-1}$ and using the above definitions, as well as the fact that i_1 is algebra morphism, one obtains the following identities.

$$\begin{aligned} m_B &= \Lambda^{-1} \circ m_A^{(3)} \circ (\text{id} \otimes \Lambda \circ \Lambda^{-1} \otimes \text{id}) \circ (i_1 \otimes m_A \circ (i_2 \otimes i_1) \otimes i_2) \\ &= \Lambda^{-1} \circ m_A^{(4)} \circ (i_1 \otimes (i_1 \otimes i_2) \circ \varphi_{2,1} \otimes i_2) \\ &= \Lambda^{-1} \circ m_A^{(3)} \circ (i_1 \otimes i_1 \otimes i_2) \circ (m_1 \otimes \hat{\sigma}) \circ (\text{id} \otimes \varphi_{2,1} \otimes \text{id}) \\ &= (m_1 \otimes \text{id}) \circ (m_1 \otimes \hat{\sigma}) \circ (\text{id} \otimes \varphi_{2,1} \otimes \text{id}). \end{aligned}$$

Similarly one derives $\Lambda \circ (\eta_1 \otimes \eta_2) = \eta_A$, hence $\eta_B = \eta_1 \otimes \eta_2$. Therefore $(B, m_B, \eta_B) = B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$. \blacksquare

PROPOSITION 2.4 *Let $B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ be a cross product algebra, and A be an algebra. Suppose there are morphisms $\alpha : B_1 \rightarrow A$ and $\beta : B_2 \rightarrow A$ such that*

1. α is an algebra morphism.
2. $\beta \circ \eta_2 = \eta_A$.
3. $m_A \circ (\alpha \otimes \beta) \circ \varphi_{2,1} = m_A \circ (\beta \otimes \alpha)$.
4. $m_A \circ (\alpha \otimes \beta) \circ \hat{\sigma} = m_A \circ (\beta \otimes \beta)$.

Then there exists a unique algebra morphism $\gamma : B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2 \rightarrow A$ obeying the identities $\gamma \circ (\text{id}_{B_1} \otimes \eta_2) = \alpha$ and $\gamma \circ (\eta_1 \otimes \text{id}_{B_2}) = \beta$.

PROOF. Define $\gamma := m_A \circ (\alpha \otimes \beta)$. Then by assumption 1 and 2 of Proposition 2.4 it follows $\gamma \circ (\text{id}_{B_1} \otimes \eta_2) = \alpha$, $\gamma \circ (\eta_1 \otimes \text{id}_{B_2}) = \beta$, and $\gamma \circ (\eta_1 \otimes \eta_2) = \eta_A$. Using consecutively that α is an algebra morphism, assumption 4, and assumption 3 of Proposition 2.4 then yields $\gamma \circ m_{B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2} = m_A^{(4)} \circ (\alpha \otimes \alpha \otimes (\alpha \otimes \beta) \circ \hat{\sigma}) \circ (\text{id} \otimes \varphi_{2,1} \otimes \text{id}) = m_A^{(4)} \circ (\alpha \otimes (\alpha \otimes \beta) \circ \varphi_{2,1} \otimes \beta) = m_A \circ (m_A \circ (\alpha \otimes \beta) \otimes m_A \circ (\alpha \otimes \beta)) = m_A \circ (\gamma \otimes \gamma)$. Therefore γ is an algebra morphism obeying the conditions of Proposition 2.4. Suppose that there is another γ' meeting the same stipulations as γ . Then $\gamma = m_A \circ (\alpha \otimes \beta) = m_A \circ (\gamma' \circ (\text{id} \otimes \eta_2) \otimes \gamma' \circ (\eta_1 \otimes \text{id})) = \gamma' \circ m_{B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2} \circ (\text{id} \otimes \eta_2 \otimes \eta_1 \otimes \text{id}) = \gamma'$. In the last equation unital properties of cross product algebra have been used. This proves uniqueness of γ . \blacksquare

REMARK 3 The conditions of Proposition 2.4 can be realized on $A = B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ with $\alpha = \text{id}_{B_1} \otimes \eta_2$ and $\beta = \eta_1 \otimes \text{id}_{B_2}$. Then of course $\gamma = \text{id}$.

In the following we consider generalized smash product algebras [5] and demonstrate that they are special cases of cross product algebras³. A (generalized) *smash product algebra* $B_1 \times_{\phi_{2,1}} B_2$ consists of algebras B_1 and B_2 , and a morphism $\phi_{2,1} : B_2 \otimes B_1 \rightarrow B_1 \otimes B_2$, such that $B = B_1 \otimes B_2$ is an algebra through $m_B = (m_1 \otimes m_2) \circ (\text{id}_{B_1} \otimes \phi_{2,1} \otimes \text{id}_{B_2})$ and $\eta_B = \eta_1 \otimes \eta_2$. The next proposition shows that smash product algebras are indeed special cases of cross product algebras under certain natural conditions.

PROPOSITION 2.5 *Suppose there is a morphism $\varepsilon_1 : B_1 \rightarrow \mathbb{1}$ with $\varepsilon_1 \circ \eta_1 = \text{id}_{\mathbb{1}}$. Then*

$B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ is a cross product algebra with $\hat{\sigma} = \eta_1 \otimes m_2$ for some morphism $m_2 : B_2 \otimes B_2 \rightarrow B_2$.

$\Leftrightarrow B_1 \times_{\varphi_{2,1}} B_2$ is a smash product algebra.

PROOF. The proposition can be proven easily by (2.3) and triviality of $\hat{\sigma}$. ■

REMARK 4 Crossed product algebras $A \#_{\sigma} H$ [12, 23] are special examples of cross product algebras through $\varphi_{2,1} := (\mu_l \otimes \text{id}_H) \circ (\Delta_H \otimes \text{id}_A)$ and $\hat{\sigma} := (\sigma \otimes m_H) \circ (\text{id}_H \otimes \Psi_{H,H} \otimes \text{id}_H) \circ (\Delta_H \otimes \Delta_H)$ where A is an algebra, $\mu_l : H \otimes A \rightarrow A$ is a left H -measure on A , and $\sigma : H \otimes H \rightarrow A$ is a (convolution invertible) cocycle.

The next proposition yields criteria under which a morphism $f : B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_1 \rightarrow A$ is multiplicative.

PROPOSITION 2.6 *Let A be an algebra and $B = B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ be a cross product algebra. Then the subsequent statements are equivalent.*

1. *The morphism $f : B \rightarrow A$ is multiplicative, i. e. $f \circ m_B = m_A \circ (f \otimes f)$.*

2. *The identities*

$$\begin{aligned} m_A \circ (f \otimes f) \circ (\text{id}_{B_1} \otimes \eta_2 \otimes \text{id}_{B_1} \otimes \text{id}_{B_2}) &= f \circ m_B \circ (\text{id}_{B_1} \otimes \eta_2 \otimes \text{id}_{B_1} \otimes \text{id}_{B_2}), \\ m_A \circ (f \otimes f) \circ (\eta_1 \otimes \text{id}_{B_2} \otimes \text{id}_{B_1} \otimes \text{id}_{B_2}) &= f \circ m_B \circ (\eta_1 \otimes \text{id}_{B_2} \otimes \text{id}_{B_1} \otimes \text{id}_{B_2}) \end{aligned}$$

are satisfied.

3. *The identities*

$$\begin{aligned} m_A \circ (f \otimes f) \circ (\text{id}_{B_1} \otimes \text{id}_{B_2} \otimes \text{id}_{B_1} \otimes \eta_2) &= f \circ m_B \circ (\text{id}_{B_1} \otimes \text{id}_{B_2} \otimes \text{id}_{B_1} \otimes \eta_2), \\ m_A \circ (f \otimes f) \circ (\text{id}_{B_1} \otimes \text{id}_{B_2} \otimes \eta_1 \otimes \text{id}_{B_2}) &= f \circ m_B \circ (\text{id}_{B_1} \otimes \text{id}_{B_2} \otimes \eta_1 \otimes \text{id}_{B_2}) \end{aligned}$$

are satisfied.

4. *The identities*

$$\begin{aligned} m_A \circ (f \otimes f) \circ (\text{id}_{B_1} \otimes \eta_2 \otimes \text{id}_{B_1} \otimes \text{id}_{B_2}) &= f \circ m_B \circ (\text{id}_{B_1} \otimes \eta_2 \otimes \text{id}_{B_1} \otimes \text{id}_{B_2}), \\ m_A \circ (f \otimes f) \circ (\text{id}_{B_1} \otimes \text{id}_{B_2} \otimes \eta_1 \otimes \text{id}_{B_2}) &= f \circ m_B \circ (\text{id}_{B_1} \otimes \text{id}_{B_2} \otimes \eta_1 \otimes \text{id}_{B_2}), \\ m_A \circ (f \otimes f) \circ (\eta_1 \otimes \text{id}_{B_2} \otimes \text{id}_{B_1} \otimes \eta_2) &= f \circ m_B \circ (\eta_1 \otimes \text{id}_{B_2} \otimes \text{id}_{B_1} \otimes \eta_2) \end{aligned}$$

are satisfied.

³We adopt the naming "smash product algebras" from [9].

PROOF. The non-trivial part of the proposition can be verified easily with the help of the identity $m_B \circ (\text{id} \otimes \eta_2 \otimes \eta_1 \otimes \text{id}) = \text{id} \otimes \text{id}$, the associativity of A and B , and the assumptions 2.6.2, 2.6.3 or 2.6.4 respectively. \blacksquare

Cross Product Coalgebras

Cross product coalgebras are somehow dual constructions to cross product algebras. We will omit proofs since they can be obtained from the corresponding proofs for cross product algebras by an obvious kind of dualization.

DEFINITION 2.7 *Let $(C_2, \Delta_2, \varepsilon_2)$ be a coalgebra and C_1 be an object in \mathcal{C} . Suppose there exist morphisms $\varepsilon_1 : B_1 \rightarrow \mathbb{1}$, $\varphi_{1,2} : C_1 \otimes C_2 \rightarrow C_2 \otimes C_1$, $\hat{\rho} : C_1 \otimes C_2 \rightarrow C_1 \otimes C_1$ such that the relation $\varphi_{1,2} = (\varepsilon_1 \otimes \varphi_{1,2}) \circ (\hat{\rho} \otimes \text{id}_{C_2}) \circ (\text{id}_{C_1} \otimes \Delta_2)$ holds. Then $C = C_1 \otimes C_2$ is called cross product coalgebra if it is a coalgebra through $\Delta_C = (\text{id}_{B_1} \otimes \varphi_{1,2} \otimes \text{id}_{B_2}) \circ (\hat{\rho} \otimes \Delta_2) \circ (\text{id}_{B_1} \otimes \Delta_2)$ and $\varepsilon_C = \varepsilon_1 \otimes \varepsilon_2$. We denote C by $C_1 \underset{\varphi_{1,2}}{\hat{\rho}} \bowtie C_2$.*

PROPOSITION 2.8 *The following statements are equivalent.*

1. $C_1 \underset{\varphi_{1,2}}{\hat{\rho}} \bowtie C_2$ is a cross product coalgebra.
2. $(C_2, \Delta_2, \varepsilon_2)$ is a coalgebra and C_1 is an object in \mathcal{C} . There are morphisms $\varepsilon_1 : B_1 \rightarrow \mathbb{1}$ and $\varphi_{1,2} : C_1 \otimes C_2 \rightarrow C_2 \otimes C_1$ such that

$$\begin{aligned} (\text{id}_{C_2} \otimes \varepsilon_1) \circ \varphi_{1,2} &= \varepsilon_1 \otimes \text{id}_{C_2}, & (\text{id}_{C_1} \otimes \varepsilon_1) \circ \hat{\rho} &= \text{id}_{C_1} \otimes \varepsilon_2, \\ (\varepsilon_1 \otimes \text{id}_{C_2}) \circ \varphi_{1,2} &= \text{id}_{C_2} \otimes \varepsilon_1, & (\varepsilon_1 \otimes \text{id}_{C_1}) \circ \hat{\rho} &= \text{id}_{C_1} \otimes \varepsilon_2, \\ (\Delta_2 \otimes \text{id}_{C_1}) \circ \varphi_{1,2} &= (\text{id}_{C_2} \otimes \varphi_{1,2}) \circ (\varphi_{1,2} \otimes \text{id}_{C_2}) \circ (\text{id}_{C_1} \otimes \Delta_2), \\ (\hat{\rho} \otimes \text{id}_{C_1}) \circ (\text{id}_{C_1} \otimes \varphi_{1,2}) \circ (\hat{\rho} \otimes \text{id}_{C_2}) \circ (\text{id}_{C_1} \otimes \Delta_2) \\ &= (\text{id}_{C_1} \otimes \hat{\rho}) \circ (\hat{\rho} \otimes \text{id}_{C_2}) \circ (\text{id}_{C_1} \otimes \Delta_2), \\ (\varphi_{1,2} \otimes \text{id}_{C_1}) \circ (\text{id}_{C_1} \otimes \varphi_{1,2}) \circ (\hat{\rho} \otimes \text{id}_{C_2}) \circ (\text{id}_{C_1} \otimes \Delta_2) \\ &= (\text{id}_{C_2} \otimes \hat{\rho}) \circ (\varphi_{1,2} \otimes \text{id}_{C_2}) \circ (\text{id}_{C_1} \otimes \Delta_2). \end{aligned}$$

\blacksquare

Like cross product algebras the cross product coalgebras are universal constructions, too.

PROPOSITION 2.9 *Let C be a coalgebra. Then*

C is coalgebra isomorphic to a cross product coalgebra $C_1 \underset{\varphi_{1,2}}{\hat{\rho}} \bowtie C_2$.

\Leftrightarrow *There is a coalgebra $(C_2, \Delta_2, \varepsilon_2)$ and an object C_1 in \mathcal{C} , morphisms $C_1 \xleftarrow{p_1} C \xrightarrow{p_2} C_2$ and $\varepsilon_1 : C_1 \rightarrow \mathbb{1}$ where p_2 is a coalgebra morphism and $\varepsilon_1 \circ p_1 = \varepsilon_C$ such that $(p_1 \otimes p_2) \circ \Delta_C : C \rightarrow C_1 \otimes C_2$ is an isomorphism in \mathcal{C} .* \blacksquare

PROPOSITION 2.10 *Let $C_1 \hat{\varphi}_{1,2} \bowtie C_2$ be a cross product coalgebra and C be a coalgebra. Suppose that there exist morphisms a and b such that*

1. $a : C \rightarrow C_1$ and $\varepsilon_1 \circ a = \varepsilon_C$.
2. $b : C \rightarrow C_2$ is a coalgebra morphism.
3. $\varphi_{1,2} \circ (a \otimes b) \circ \Delta_C = (b \otimes a) \circ \Delta_C$.
4. $\hat{\rho} \circ (a \otimes b) \circ \Delta_C = (a \otimes a) \circ \Delta_C$.

Then there exists a unique coalgebra morphism $c : C \rightarrow C_1 \hat{\varphi}_{1,2} \bowtie C_2$ obeying the identities $a = (\text{id}_{C_1} \otimes \varepsilon_2) \circ c$ and $b = (\varepsilon_1 \otimes \text{id}_{C_2}) \circ c$. ■

Co-smash product coalgebras are special cases of cross product coalgebras. More precisely, a *co-smash product coalgebra* $C = C_1 \times^{\phi_{1,2}} C_2$ is given by coalgebras C_1 and C_2 , and a morphism $\phi_{1,2} : C_1 \otimes C_2 \rightarrow C_2 \otimes C_1$, such that $C = C_1 \otimes C_2$ is a coalgebra through $\Delta_C = (\text{id}_{C_1} \otimes \phi_{1,2} \otimes \text{id}_{C_2}) \circ (\Delta_1 \otimes \Delta_2)$ and $\varepsilon_C = \varepsilon_{C_1} \otimes \varepsilon_{C_2}$. Then the corresponding result of Proposition 2.5 holds for co-smash product coalgebras.

PROPOSITION 2.11 *Suppose C_2 is a coalgebra and there is a morphism $\eta_2 : \mathbb{1} \rightarrow C_2$ with $\varepsilon_2 \circ \eta_2 = \text{id}_{\mathbb{1}}$. Then*

$(C_1, C_2, \varphi_{1,2}, \hat{\rho})$ is a cross product coalgebra with $\hat{\rho} = \Delta_1 \otimes \varepsilon_2$ for some morphism $\Delta_2 : C_2 \rightarrow C_2 \otimes C_2$.

$\Leftrightarrow C_1 \times^{\varphi_{1,2}} C_2$ is a co-smash product coalgebra. ■

REMARK 5 (π -SYMMETRY) We would like to point out the following important observation to the reader. Definition 2.1 is not dual to Definition 2.7. Rather, both definitions can be obtained from each other by a combination of duality (followed by the usual exchanges of multiplication and comultiplication, unit and counit, cocycle and cycle, etc.) and use of the opposite tensor product (followed by exchange of indices "1 \leftrightarrow 2", and later also by exchange of "left/right (coaction) \leftrightarrow right/left (action)"). This is not an accidental fact but has to do with the subsequent definition of cross product bialgebras where we use precisely these algebra and coalgebra structures. Only then are we able to get all the known cross product bialgebras [24, 18, 22, 20, 5]. This kind of symmetry between cross product algebra and cross product coalgebra will be called **π -symmetry** henceforth. In terms of graphical calculus π -symmetry can be interpreted easily as rotation of the graphic by the angle π along an axis normal to the planar graphic followed by the above mentioned exchanges of indices and morphism types. We will often apply this principle to obtain π -symmetric results simply by π -rotation. Hence cross product bialgebras - as defined below - are therefore π -symmetric invariant rather than dual symmetric invariant. Dualization of our definition would lead to another kind of cross product bialgebras. The corresponding results for these dual versions follow from our results straightforwardly by dualization.

3 Cross Product Bialgebras

In Section 3 we investigate cross product bialgebras from a universal point of view. They are simultaneously cross product algebras and cross product coalgebras with compatible bialgebra structure. We find a universal description for the isomorphism classes of cross product bialgebras. Then we will consider special cases called cocycle and strong cross product bialgebras which are universal constructions as well. But in addition the given isomorphism class of a cocycle/strong cross product bialgebra can be characterized by certain interrelated (co-)modular co-cyclic structures on the tensor factors. Strong cross product bialgebras are the pivotal objects for the studies in Section 4.

Cross Product Bialgebras – General Definition

DEFINITION 3.1 *A bialgebra B is called cross product bialgebra if its underlying algebra is a cross product algebra $B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$, and its underlying coalgebra is a cross product coalgebra $B_1 \hat{\bowtie}_{\varphi_{1,2}} B_2$ on the same objects. The cross product bialgebra B will be denoted by $B_1 \hat{\bowtie}_{\varphi_{1,2}}^{\hat{\sigma}} \bowtie_{\varphi_{2,1}} B_2$. A cross product bialgebra is called normalized if $\varepsilon_1 \circ \eta_1 = \text{id}_{\mathbb{1}}$ (and then equivalently $\varepsilon_2 \circ \eta_2 = \text{id}_{\mathbb{1}}$).*

Cross product bialgebras are universal in the following sense.

THEOREM 3.2 *Let B be a bialgebra in \mathcal{C} . Then the subsequent equivalent conditions hold.*

1. *B is bialgebra isomorphic to a normalized cross product bialgebra $B_1 \hat{\bowtie}_{\varphi_{1,2}}^{\hat{\sigma}} \bowtie_{\varphi_{2,1}} B_2$.*
2. *There are idempotents $\Pi_1, \Pi_2 \in \text{End}(B)$ such that*

$$\begin{aligned} m_B \circ (\Pi_1 \otimes \Pi_1) &= \Pi_1 \circ m_B \circ (\Pi_1 \otimes \Pi_1), & \Pi_1 \circ \eta_B &= \eta_B, \\ (\Pi_2 \otimes \Pi_2) \circ \Delta_B &= (\Pi_2 \otimes \Pi_2) \circ \Delta_B \circ \Pi_2, & \varepsilon_B \circ \Pi_2 &= \varepsilon_B, \end{aligned} \quad (3.1)$$

and the sequence $B \otimes B \xrightarrow{m_B \circ (\Pi_1 \otimes \Pi_2)} B \xrightarrow{(\Pi_1 \otimes \Pi_2) \circ \Delta_B} B \otimes B$ is a splitting of the idempotent $\Pi_1 \otimes \Pi_2$ of $B \otimes B$.

3. *There are objects B_1 and B_2 and morphisms $B_1 \xrightarrow{i_1} B \xrightarrow{p_1} B_1$ and $B_2 \xrightarrow{i_2} B \xrightarrow{p_2} B_2$ where i_j is algebra morphism, p_j is coalgebra morphism and $p_j \circ i_j = \text{id}_{B_j}$ for $j \in \{1, 2\}$ such that $m_B \circ (i_1 \otimes i_2) : B_1 \otimes B_2 \rightarrow B$ and $(p_1 \otimes p_2) \circ \Delta_B : B \rightarrow B_1 \otimes B_2$ are mutually inverse isomorphisms.*

PROOF. “2. \Rightarrow 3.”: For $j \in \{1, 2\}$ we define i_j, p_j to be the morphisms splitting the idempotent Π_j as $\Pi_j = i_j \circ p_j$ and $p_j \circ i_j = \text{id}_{B_j}$ for some objects B_j . Then with the help of Theorem 3.2.2 it follows $m_B \circ (i_1 \otimes i_2) \circ (p_1 \otimes p_2) \circ \Delta_B = m_B \circ (\Pi_1 \otimes \Pi_2) \circ \Delta_B = \text{id}_B$ and $(p_1 \otimes p_2) \circ \Delta_B \circ m_B \circ (i_1 \otimes i_2) = (p_1 \otimes p_2) \circ (\Pi_1 \otimes \Pi_2) \circ \Delta_B \circ m_B \circ (\Pi_1 \otimes \Pi_2) \circ (i_1 \otimes i_2) = (p_1 \circ i_1 \otimes p_2 \circ i_2) = \text{id}_{B_1 \otimes B_2}$. Hence $(m_B \circ (i_1 \otimes i_2))^{-1} = (p_1 \otimes p_2) \circ \Delta_B$. Now we define $m_1 := p_1 \circ m_B \circ (i_1 \otimes i_1)$, $\eta_1 := p_1 \circ \eta_B$, $\Delta_2 := (p_2 \otimes p_2) \circ \Delta_B \circ i_2$, and $\varepsilon_2 := \varepsilon_B \circ i_2$. Then using (3.1) and the (co-)algebra property of B one verifies easily that (B_1, m_1, η_1) is an algebra and $(B_2, \Delta_2, \varepsilon_2)$ is a coalgebra. Furthermore i_1 is an algebra morphism because $i_1 \circ m_1 = \Pi_1 \circ m_B \circ (i_1 \otimes i_1) = m_B \circ (i_1 \otimes i_1)$, and $i_1 \circ \eta_1 = \Pi_1 \circ \eta_B = \eta_B$. In a π -symmetric manner one proves that p_2 is a coalgebra morphism.

“3. \Rightarrow 2.”: Conversely we define $\Pi_j := i_j \circ p_j$ for $j \in \{1, 2\}$. Since by assumption i_1 is an algebra morphism it follows $\Pi_1 \circ \eta_B = \Pi_1 \circ i_1 \circ \eta_1 = i_1 \circ \eta_1 = \eta_B$ and using $p_1 \circ i_1 = \text{id}_{B_1}$ it holds $\Pi_1 \circ m_B \circ (\Pi_1 \otimes \Pi_1) = i_1 \circ p_1 \circ i_1 \circ m_1 \circ (p_1 \otimes p_1) = m_B \circ (\Pi_1 \otimes \Pi_1)$. In π -symmetrical analogy it will be proven that $(\Pi_2 \otimes \Pi_2) \circ \Delta_B = (\Pi_2 \otimes \Pi_2) \circ \Delta_B \circ \Pi_2$ and $\varepsilon_B \circ \Pi_2 = \varepsilon_B$. From Theorem 3.2.3 we conclude $(\Pi_1 \otimes \Pi_2) \circ \Delta_B \circ m_B \circ (\Pi_1 \otimes \Pi_2) = (i_1 \otimes i_2) \circ \text{id}_{B_1 \otimes B_2} \circ (p_1 \otimes p_2) = \Pi_1 \otimes \Pi_2$ and $\text{id}_B = m_B \circ (\Pi_1 \otimes \Pi_2) \circ \Delta_B = m_B \circ (\Pi_1 \otimes \Pi_2) \circ (\Pi_1 \otimes \Pi_2) \circ \Delta_B$. Thus $m_B \circ (\Pi_1 \otimes \Pi_2)$ and $(\Pi_1 \otimes \Pi_2) \circ \Delta_B$ split the idempotent $(\Pi_1 \otimes \Pi_2)$.

“3. \Rightarrow 1.”: By assumption (B_1, m_1, η_1) is an algebra and i_1 is an algebra morphism, i_2 is a morphism in \mathcal{C} , and $m_B \circ (i_1 \otimes i_2)$ is an isomorphism. Define $\eta_2 := p_2 \circ \eta_B : \mathbb{1} \rightarrow B_2$. Then $i_2 \circ \eta_2 = \Pi_2 \circ \eta_B = m_B \circ (\Pi_1 \circ \eta_B \otimes \Pi_2 \circ \eta_B) = m_B \circ (i_1 \otimes i_2) \circ (p_1 \otimes p_2) \circ \Delta_B \circ \eta_B = \eta_B$. Therefore all conditions of Proposition 2.3.2 are fulfilled implying that B is algebra isomorphic to a cross product algebra $B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$. From the proof of Proposition 2.3 one reads off that the isomorphism is given by $\Lambda = m_B \circ (i_1 \otimes i_2)$. In a π -symmetric way one shows that B is coalgebra isomorphic to a cross product coalgebra $B_1 \bowtie_{\varphi_{1,2}}^{\hat{\rho}} B_2$ on the same tensor product with isomorphism $\tilde{\Lambda} = (p_1 \otimes p_2) \circ \Delta_B$. Thus by assumption $\tilde{\Lambda} = \Lambda^{-1}$. Since B is a bialgebra it follows then from Definition 3.1 that $B_1 \otimes B_2$ is cross product bialgebra $B_1 \bowtie_{\varphi_{1,2}}^{\hat{\rho}} \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ and $\Lambda : B_1 \bowtie_{\varphi_{1,2}}^{\hat{\rho}} \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2 \rightarrow B$ is bialgebra isomorphism. Furthermore the identities $\text{id}_{\mathbb{1}} = \varepsilon_B \circ \eta_B = \varepsilon_B \circ i_2 \circ \eta_2 = \varepsilon_2 \circ p_2 \circ i_2 \circ \eta_2 = \varepsilon_2 \circ \eta_2$ show that $B_1 \bowtie_{\varphi_{1,2}}^{\hat{\rho}} \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ is normalized.

“1. \Rightarrow 3.”: Given a bialgebra isomorphism $\Lambda : B_1 \bowtie_{\varphi_{1,2}}^{\hat{\rho}} \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2 \rightarrow B$. Then in particular $\Lambda : B_1 \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2 \rightarrow B$ is an algebra isomorphism and $\Lambda : B_1 \bowtie_{\varphi_{1,2}}^{\hat{\rho}} B_2 \rightarrow B$ is a coalgebra isomorphism. We define $i_1 := \Lambda \circ (\text{id}_{B_1} \otimes \eta_2)$, $i_2 := \Lambda \circ (\eta_1 \otimes \text{id}_{B_2})$, $p_1 := (\text{id}_{B_1} \otimes \varepsilon_2) \circ \Lambda^{-1}$, and $p_2 := (\varepsilon_1 \otimes \text{id}_{B_2}) \circ \Lambda^{-1}$. From Propositions 2.3 and 2.9 one derives that i_1 is an algebra morphism, p_2 is a coalgebra morphism, and $\Lambda = m_B \circ (i_1 \otimes i_2) = ((p_1 \otimes p_2) \circ \Delta_B)^{-1}$. By assumption $B_1 \bowtie_{\varphi_{1,2}}^{\hat{\rho}} \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ is normalized from which follows $p_1 \circ i_1 = (\text{id}_{B_1} \otimes \varepsilon_2) \circ \Lambda^{-1} \circ \Lambda \circ (\text{id}_{B_1} \otimes \eta_2) = \text{id}_{B_1}$ and similarly $p_2 \circ i_2 = \text{id}_{B_2}$. \blacksquare

The next corollary of Theorem 3.2 will be used frequently in the following.

COROLLARY 3.3 *Under the conditions of Theorem 3.2 it holds*

$$\begin{aligned} \Pi_2 \circ \eta_B &= \eta_B, & \Pi_1 \circ \Pi_2 &= \eta_B \circ \varepsilon_B, & \Pi_2 \circ m_B &= \Pi_2 \circ m_B \circ (\Pi_2 \otimes \text{id}_B), \\ \varepsilon_B \circ \Pi_1 &= \varepsilon_B, & \Pi_2 \circ \Pi_1 &= \eta_B \circ \varepsilon_B, & \Delta_B \circ \Pi_1 &= (\text{id}_B \otimes \Pi_1) \circ \Delta_B \circ \Pi_1, \\ & & \Pi_1 \circ m_B \circ (\Pi_1 \otimes \text{id}_B) &= \Pi_1 \circ m_B \circ (\Pi_1 \otimes \Pi_1), \\ & & (\text{id}_B \otimes \Pi_2) \circ \Delta_B \circ \Pi_2 &= (\Pi_2 \otimes \Pi_2) \circ \Delta_B \circ \Pi_2. \end{aligned} \tag{3.2}$$

The structure morphisms $\varphi_{1,2}$, $\varphi_{2,1}$, $\hat{\rho}$, and $\hat{\sigma}$ of the cross product bialgebra $B_1 \bowtie_{\varphi_{1,2}}^{\hat{\rho}} \bowtie_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ can be expressed through the projections and injections on B as

$$\begin{aligned} i_1 \circ \eta_1 &= \eta_B, & \varepsilon_1 \circ p_1 &= \varepsilon_B, & i_2 \circ \eta_2 &= \eta_B, & \varepsilon_2 \circ p_2 &= \varepsilon_B, \\ m_1 &= p_1 \circ m_B \circ (i_1 \otimes i_1), & \Delta_2 &= (p_2 \otimes p_2) \circ \Delta_B \circ i_2, \\ \varphi_{1,2} &= (p_2 \otimes p_1) \circ \Delta_B \circ m_B \circ (i_1 \otimes i_2), & \hat{\rho} &= (p_1 \otimes p_1) \circ \Delta_B \circ m_B \circ (i_1 \otimes i_2), \\ \varphi_{2,1} &= (p_1 \otimes p_2) \circ \Delta_B \circ m_B \circ (i_2 \otimes i_1), & \hat{\sigma} &= (p_1 \otimes p_2) \circ \Delta_B \circ m_B \circ (i_2 \otimes i_2). \end{aligned} \tag{3.3}$$

PROOF. The identities $i_1 \circ \eta_1 = \eta_B$ and $\varepsilon_2 \circ p_2 = \varepsilon_B$ hold since i_1 is an algebra morphism and p_2 is a coalgebra morphism, whereas the identities $i_2 \circ \eta_2 = \eta_B$ and $\varepsilon_1 \circ p_1 = \varepsilon_B$, as

well as the equations $\Pi_2 \circ \eta_B = \eta_B$ and $\varepsilon_B \circ \Pi_1 = \varepsilon_B$ have been shown in the proof of Theorem 3.2. From the proofs of Propositions 2.3 and 2.9 one can directly read off the structure of the morphisms $\varphi_{1,2}$, $\varphi_{2,1}$, $\hat{\rho}$, and $\hat{\sigma}$ given in (3.3) using $\Lambda = m_B \circ (i_1 \otimes i_2)$ and $\Lambda^{-1} = (p_1 \otimes p_2) \circ \Delta_B$. Since i_1 is algebra morphism, p_2 is coalgebra morphism and $p_j \circ i_j = \text{id}_{B_j}$ for $j \in \{1, 2\}$ it follows $m_1 = p_1 \circ m_B \circ (i_1 \otimes i_1)$ and $\Delta_2 = (p_2 \otimes p_2) \circ \Delta_B \circ i_2$. Because of (3.1) and Theorem 3.2.2 it follows $\Pi_1 \circ \Pi_2 = (\text{id}_B \otimes \varepsilon_B) \circ (\Pi_1 \otimes \Pi_2) \circ \Delta_B \circ m_B \circ (\Pi_1 \otimes \Pi_2) \circ (\eta_B \otimes \text{id}_B) = (\Pi_1 \circ \eta_B \otimes \varepsilon_B \circ \Pi_2) = \eta_B \circ \varepsilon_B$. Using $\Pi_2 \circ \eta_B = \eta_B$ and $\varepsilon_B \circ \Pi_1 = \varepsilon_B$ from above we obtain in a similar way $\Pi_2 \circ \Pi_1 = \eta_B \circ \varepsilon_B$. Using consecutively Theorem 3.2.2 two times, (3.1), the identity $\Pi_2 = (\varepsilon_B \circ \Pi_1 \otimes \Pi_2) \circ \Delta_B$, again Theorem 3.2.2, and the identity $\varepsilon_B \circ \Pi_1 = \varepsilon_B$ one obtains the following series of equations.

$$\begin{aligned}
\Pi_2 \circ m_B &= \Pi_2 \circ m_B \circ (m_B \circ (\Pi_1 \otimes \Pi_2) \circ \Delta_B \otimes \text{id}_B) \\
&= \Pi_2 \circ m_B^{(3)} \circ (\Pi_1 \otimes \Pi_1 \otimes \Pi_2) \circ (\text{id}_B \otimes \Delta_B \circ m_B) \circ \\
&\quad \circ (\Pi_1 \otimes \Pi_2 \otimes \text{id}_B) \circ (\Delta_B \otimes \text{id}_B) \\
&= \Pi_2 \circ m_B \circ (\Pi_1 \otimes \Pi_2) \circ (m_B \circ (\Pi_1 \otimes \Pi_1) \otimes \text{id}_B) \circ \\
&\quad \circ (\text{id}_B \otimes \Delta_B \circ m_B) \circ (\text{id}_B \otimes \Pi_2 \otimes \text{id}_B) \circ (\Delta_B \otimes \text{id}_B) \\
&= \Pi_2 \circ m_B \circ (\Pi_2 \otimes \text{id}_B)
\end{aligned}$$

and π -symmetrically $\Delta_B \circ \Pi_1 = (\text{id}_B \otimes \Pi_1) \circ \Delta_B \circ \Pi_1$. Finally one obtains in a similar manner

$$\begin{aligned}
\Pi_1 \circ m_B \circ (\Pi_1 \otimes \text{id}_B) &= \Pi_1 \circ m_B \circ (\Pi_1 \otimes m_B \circ (\Pi_1 \otimes \Pi_2) \circ \Delta_B) \\
&= \Pi_1 \circ m_B^{(3)} \circ (\Pi_1 \otimes \Pi_1 \otimes \Pi_2) \circ (\text{id}_B \otimes \Delta_B) \\
&= \Pi_1 \circ m_B \circ (\Pi_1 \otimes \Pi_2) \circ (m_B \circ (\Pi_1 \otimes \Pi_1) \otimes \Pi_2) \circ (\text{id}_B \otimes \Delta_B) \\
&= (\Pi_1 \otimes \varepsilon_B \circ \Pi_2) \circ \Delta_B \circ m_B \circ (\Pi_1 \otimes \Pi_2) \circ \\
&\quad \circ (m_B \circ (\Pi_1 \otimes \Pi_1) \otimes \Pi_2) \circ (\text{id}_B \otimes \Delta_B) \\
&= \Pi_1 \circ m_B \circ (\Pi_1 \otimes \Pi_1)
\end{aligned}$$

and $(\text{id}_B \otimes \Pi_2) \circ \Delta_B \circ \Pi_2 = (\Pi_2 \otimes \Pi_2) \circ \Delta_B \circ \Pi_2$ by π -symmetry. \blacksquare

REMARK 6 The statement of Theorem 3.2 can be refined in the following sense. For a given bialgebra B the tuples $(B, \Lambda, B_1 \xrightarrow{\hat{\rho}} \xrightarrow{\hat{\sigma}} B_2)$ and (B, i_1, i_2, p_1, p_2) obeying the conditions of Theorem 3.2.1 and 3.2.3 respectively, are in one-to-one correspondence. This correspondence has been constructed in the proof of Theorem 3.2.

Theorem 3.2 and the previous remark imply a useful corollary. Given a normalized cross product bialgebra $B_1 \xrightarrow{\hat{\rho}} \xrightarrow{\hat{\sigma}} B_2$. We set $i_{\alpha 1} := \text{id}_{B_1} \otimes \eta_2$, $i_{\alpha 2} := \eta_1 \otimes \text{id}_{B_2}$, $p_{\alpha 1} := \text{id}_{B_1} \otimes \varepsilon_2$ and $p_{\alpha 2} := \varepsilon_1 \otimes \text{id}_{B_2}$. Then we define

$$\begin{aligned}
m_{\triangleright i,j,k} &:= p_{\alpha i} \circ m_{B_1 \xrightarrow{\hat{\rho}} \xrightarrow{\hat{\sigma}} B_2} \circ (i_{\alpha j} \otimes i_{\alpha k}), \\
\Delta_{\triangleright i,j,k} &:= (p_{\alpha i} \otimes p_{\alpha j}) \circ \Delta_{B_1 \xrightarrow{\hat{\rho}} \xrightarrow{\hat{\sigma}} B_2} \circ i_{\alpha k}
\end{aligned} \tag{3.4}$$

for $i, j, k \in \{1, 2\}$. On the other hand suppose that B is a bialgebra which obeys the conditions of Theorem 3.2.3. Then we define

$$m_{B i,j,k} := p_i \circ m_B \circ (i_j \otimes i_k), \quad \Delta_{B i,j,k} := (p_i \otimes p_j) \circ \Delta_B \circ i_k \tag{3.5}$$

for $i, j, k \in \{1, 2\}$.

COROLLARY 3.4 *Let B be bialgebra and suppose that the tuples $(B, \Lambda, B_1 \varphi_{1,2}^{\hat{\rho}} \bowtie_{\varphi_{2,1}^{\hat{\sigma}}} B_2)$ and (B, i_1, i_2, p_1, p_2) are related according to Theorem 3.2. Then $m_{\bowtie i,j,k} = m_B i,j,k$ and $\Delta_{\bowtie ij,k} = \Delta_B ij,k$.*

PROOF. If the conditions of Theorem 3.6 hold, then by construction (see the proof of the theorem) the injections and projections of B and the morphisms $i_{\alpha 1}$, $i_{\alpha 2}$, $p_{\alpha 1}$ and $p_{\alpha 2}$ are related by $i_1 := \Lambda \circ i_{\alpha 1}$, $i_2 := \Lambda \circ i_{\alpha 2}$, $p_1 := p_{\alpha 1} \circ \Lambda^{-1}$, and $p_2 := p_{\alpha 2} \circ \Lambda^{-1}$, where $\Lambda : B_1 \otimes B_2 \rightarrow B$ is the given bialgebra isomorphism. Thus $m_{\bowtie i,j,k} = m_B i,j,k$ and $\Delta_{\bowtie ij,k} = \Delta_B ij,k$ follow directly. \blacksquare

Hence we will use the notation $m_{i,j,k}$ and $\Delta_{ij,k}$ for the corresponding morphisms henceforth.

Cocycle Cross Product Bialgebras

Up to now a (co-)modular co-cyclic structure of cross product bialgebras and their tensor factors B_1 and B_2 did not emerge. In Definition 3.5 below we will restrict our considerations to cocycle cross product bialgebras for which (co-)modular co-cyclic structures appear in a very natural way. The universal property of cocycle cross product bialgebras will be discussed subsequently. We define strong cross product bialgebras in Definition 3.8. They are the basic objects which eventually constitute the universal, (co-)modular co-cyclic theory of cross product bialgebras.

In Definition 3.1 the morphisms m_1 , η_1 , ε_1 , Δ_2 , η_2 , ε_2 , etc. occur. Definition 3.5 below requires additional morphisms Δ_1 , m_2 , μ_l , μ_r , ν_l , ν_r , σ , and ρ . This is the stage where we find it convenient to start working with graphical calculus. All m , Δ , η , ε , μ , and ν will be presented by the graphics displayed in Figure 1 respectively⁴. The cocycle and cycle morphisms σ and ρ will be presented henceforth graphically by $\sigma = \text{⌋} : B_2 \otimes B_2 \rightarrow B_1$ and $\rho = \text{⌋} : B_2 \rightarrow B_1 \otimes B_1$.

DEFINITION 3.5 *A normalized cross product bialgebra $B_1 \varphi_{1,2}^{\hat{\rho}} \bowtie_{\varphi_{2,1}^{\hat{\sigma}}} B_2$ is called cocycle cross product bialgebra if there exist additional morphisms*

$$\begin{aligned} \Delta_1 : B_1 &\rightarrow B_1 \otimes B_1, & m_2 : B_2 \otimes B_2 &\rightarrow B_2, \\ \mu_l : B_2 \otimes B_1 &\rightarrow B_1, & \mu_r : B_2 \otimes B_1 &\rightarrow B_2, \\ \nu_l : B_1 &\rightarrow B_2 \otimes B_1, & \nu_r : B_2 &\rightarrow B_2 \otimes B_1, \\ \sigma : B_2 \otimes B_2 &\rightarrow B_1, & \rho : B_2 &\rightarrow B_1 \otimes B_1 \end{aligned}$$

such that $\varphi_{1,2}$, $\varphi_{2,1}$, $\hat{\sigma}$ and $\hat{\rho}$ are of the form

$$\begin{aligned} \varphi_{1,2} &= (m_2 \otimes m_1) \circ (\text{id}_{B_1} \otimes \Psi_{B_1, B_2} \otimes \text{id}_{B_2}) \circ (\nu_l \otimes \nu_r), \\ \varphi_{2,1} &= (\mu_l \otimes \mu_r) \circ (\text{id}_{B_2} \otimes \Psi_{B_2, B_1} \otimes \text{id}_{B_1}) \circ (\Delta_2 \otimes \Delta_1), \\ \hat{\sigma} &= (\sigma \otimes m_2 \circ (\mu_r \otimes \text{id}_{B_2})) \circ (\text{id}_{B_2} \otimes \Psi_{B_2, B_2} \otimes \text{id}_{B_1 \otimes B_2}) \circ (\Delta_2 \otimes (\nu_r \otimes \text{id}_{B_2}) \circ \Delta_2), \\ \hat{\rho} &= (m_1 \circ (\text{id}_{B_1} \otimes \mu_l) \otimes m_1) \circ (\text{id}_{B_1 \otimes B_2} \otimes \Psi_{B_1, B_1} \otimes \text{id}_{B_1}) \circ ((\text{id}_{B_1} \otimes \nu_l) \circ \Delta_1 \otimes \rho). \end{aligned}$$

⁴Note that some of these morphisms might be weak. That is, weak (co-)multiplications, weak (co-)actions, etc.

Furthermore we require the following (co-)unital identities

$$\begin{aligned}
\Delta_1 \circ \eta_1 &= \eta_1 \otimes \eta_2, & (\text{id}_{B_1} \otimes \varepsilon_1) \circ \Delta_1 &= (\varepsilon_1 \otimes \text{id}_{B_1}) \circ \Delta_1 = \text{id}_{B_1}, \\
\varepsilon_2 \circ \mathfrak{m}_2 &= \varepsilon_2 \otimes \varepsilon_2, & \mathfrak{m}_2 \circ (\text{id}_{B_2} \otimes \eta_2) &= \mathfrak{m}_2 \circ (\eta_2 \otimes \text{id}_{B_2}) = \text{id}_{B_2}, \\
\mu_l \circ (\eta_2 \otimes \text{id}_{B_1}) &= \text{id}_{B_1}, & \mu_l \circ (\text{id}_{B_2} \otimes \eta_1) &= \eta_1 \circ \varepsilon_2, & \varepsilon_1 \circ \mu_l &= \varepsilon_2 \otimes \varepsilon_1, \\
\mu_r \circ (\text{id}_{B_2} \otimes \eta_1) &= \text{id}_{B_2}, & \mu_r \circ (\eta_2 \otimes \text{id}_{B_1}) &= \eta_2 \circ \varepsilon_1, & \varepsilon_2 \circ \mu_r &= \varepsilon_2 \otimes \varepsilon_1, \\
(\varepsilon_2 \otimes \text{id}_{B_1}) \circ \nu_l &= \text{id}_{B_1}, & (\text{id}_{B_2} \otimes \varepsilon_1) \circ \nu_l &= \eta_2 \circ \varepsilon_1, & \nu_l \circ \eta_1 &= \eta_2 \otimes \eta_1, \\
(\text{id}_{B_2} \otimes \varepsilon_1) \circ \nu_r &= \text{id}_{B_2}, & (\varepsilon_2 \otimes \text{id}_{B_1}) \circ \nu_r &= \eta_1 \circ \varepsilon_2, & \nu_r \circ \eta_2 &= \eta_2 \otimes \eta_1
\end{aligned} \tag{3.6}$$

and the “projection” relations

$$\begin{aligned}
(\mathfrak{m}_2 \otimes \text{id}_{B_1}) \circ (\text{id}_{B_2} \otimes \nu_r) &= (\mathfrak{m}_2 \otimes \text{id}_{B_1}) \circ (\mu_r \otimes \varphi_{1,2}) \circ (\text{id}_{B_2} \otimes (\rho \otimes \text{id}_{B_2}) \circ \Delta_2), \\
(\mu_l \otimes \text{id}_{B_1}) \circ (\text{id}_{B_2} \otimes \Delta_1) &= (\mathfrak{m}_1 \circ (\text{id}_{B_1} \otimes \sigma) \otimes \text{id}_{B_1}) \circ (\varphi_{2,1} \otimes \nu_l) \circ (\text{id}_{B_2} \otimes \Delta_1).
\end{aligned} \tag{3.7}$$

Cocycle cross product bialgebras will be denoted subsequently by $B_1 \overset{\hat{\rho}}{\underset{\hat{\sigma}}{\bowtie}}_{\varphi_{1,2}, \varphi_{2,1}} B_2$.

REMARK 7 For a cocycle cross product bialgebra the following (co-)unital identities can be derived easily.

$$\begin{aligned}
\Delta_2 \circ \eta_2 &= \eta_2 \otimes \eta_2, & \varepsilon_1 \circ \mathfrak{m}_1 &= \varepsilon_1 \otimes \varepsilon_1, \\
\varepsilon_1 \circ \sigma &= \varepsilon_2 \otimes \varepsilon_2, & \rho \circ \eta_2 &= \eta_1 \otimes \eta_1, \\
\sigma \circ (\eta_2 \otimes \text{id}_{B_2}) &= \sigma \circ (\text{id}_{B_2} \otimes \eta_2) = \eta_1 \circ \varepsilon_2, \\
(\text{id}_{B_1} \otimes \varepsilon_1) \circ \rho &= (\varepsilon_1 \otimes \text{id}_{B_1}) \circ \rho = \eta_1 \circ \varepsilon_2.
\end{aligned}$$

REMARK 8 By definition, cocycle cross product bialgebras always come with certain structure morphisms $\mathfrak{m}_1, \mathfrak{m}_2, \Delta_1, \Delta_2$, etc. There might be two different cocycle cross product bialgebras with the same underlying bialgebra structure. It is understood henceforth, that the notation $B_1 \overset{\hat{\rho}}{\underset{\hat{\sigma}}{\bowtie}}_{\varphi_{1,2}, \varphi_{2,1}} B_2$ always refers to the complete structure of cocycle cross product bialgebras.

The graphics of the morphisms $\varphi_{1,2}, \varphi_{2,1}, \hat{\rho}$ and $\hat{\sigma}$ of a cocycle cross product bialgebra $B_1 \overset{\hat{\rho}}{\underset{\hat{\sigma}}{\bowtie}}_{\varphi_{1,2}, \varphi_{2,1}} B_1$ are given by

$$\varphi_{1,2} := \begin{array}{c} \text{---} \\ | \quad | \\ \diagdown \quad \diagup \\ | \quad | \\ \text{---} \end{array}, \quad \varphi_{2,1} := \begin{array}{c} \text{---} \\ | \quad | \\ \diagup \quad \diagdown \\ | \quad | \\ \text{---} \end{array}, \quad \hat{\rho} := \begin{array}{c} \text{---} \\ | \quad | \\ \diagdown \quad \diagup \\ | \quad | \\ \text{---} \end{array}, \quad \hat{\sigma} := \begin{array}{c} \text{---} \\ | \quad | \\ \diagup \quad \diagdown \\ | \quad | \\ \text{---} \end{array} \tag{3.8}$$

Then the graphical shapes of the multiplication and the comultiplication look like

$$\mathfrak{m}_{B_1 \overset{\hat{\rho}}{\underset{\hat{\sigma}}{\bowtie}}_{\varphi_{1,2}, \varphi_{2,1}} B_1} = \begin{array}{c} \text{---} \\ | \quad | \\ \diagdown \quad \diagup \\ | \quad | \\ \text{---} \end{array} \quad \text{and} \quad \Delta_{B_1 \overset{\hat{\rho}}{\underset{\hat{\sigma}}{\bowtie}}_{\varphi_{1,2}, \varphi_{2,1}} B_1} = \begin{array}{c} \text{---} \\ | \quad | \\ \diagup \quad \diagdown \\ | \quad | \\ \text{---} \end{array} \tag{3.9}$$

It is an easy exercise to prove that Definition 3.5 is compatible with trivial (co-)actions $\mu_l, \mu_r, \nu_l, \nu_r$ and trivial (co-)cycles σ and ρ respectively, in the sense that no additional identities involving (co-)units have to be required for the remaining morphisms which define this special cocycle cross product bialgebra.

The following theorem is the central basic statement in Section 3. It describes universality of cocycle cross product bialgebras.

THEOREM 3.6 *Let B be a bialgebra in \mathcal{C} . Then the following statements are equivalent.*

1. B is bialgebra isomorphic to a cocycle cross product bialgebra $B_1 \mathop{\varphi_{1,2}}^{\hat{\rho}} \mathop{\varphi_{2,1}}^{\hat{\sigma}} B_2$.
2. There are idempotents $\Pi_1, \Pi_2 \in \text{End}(B)$ such that the conditions of Theorem 3.2.2 and the following “projection” relations hold.

$$\begin{aligned}
(\gamma_1) : & \quad (\text{id} \otimes \Pi_1) \circ \Delta_B = \delta_{(1,2,1)} \circ \gamma_1 \circ (\Pi_1 \otimes \Pi_2) \circ \Delta_B, \\
(\gamma_2) : & \quad m_B \circ (\Pi_2 \otimes \text{id}) = m_B \circ (\Pi_1 \otimes \Pi_2) \circ \gamma_2 \circ \delta_{(2,1,2)}, \\
(\delta_j) : & \quad (\Pi_j \otimes \text{id}) \circ \delta_{(0,0,0)} \circ (\text{id} \otimes \Pi_j) = (\Pi_j \otimes \text{id}) \circ \delta_{(0,j,0)} \circ (\text{id} \otimes \Pi_j)
\end{aligned} \tag{3.10}$$

for $j \in \{1, 2\}$. In (3.10) we used the abbreviations

$$\begin{aligned}
\gamma_j & := (m_B \otimes m_B) \circ (\text{id} \otimes \Pi_j \otimes \text{id} \otimes \text{id}) \circ (\text{id} \otimes \Psi_{B,B} \otimes \text{id}) \circ (\Delta_B \otimes \Delta_B), \\
\delta_{(i,j,k)} & := (m_B \otimes \text{id}) \circ (\Pi_i \otimes \Pi_j \otimes \Pi_k) \circ (\text{id} \otimes \Delta_B).
\end{aligned}$$

for $i, j, k \in \{0, 1, 2\}$, and we set $\Pi_0 := \text{id}_A$.

3. There are objects B_1 and B_2 and morphisms $B_1 \xrightarrow{i_1} B \xrightarrow{p_1} B_1$ and $B_2 \xrightarrow{i_2} B \xrightarrow{p_2} B_2$ for which the conditions of Theorem 3.2.3 and the “projection” relations (3.10) hold, with $\Pi_j := i_j \circ p_j$, $j \in \{1, 2\}$.

PROOF. Obviously statement 2. and 3. are equivalent due to Theorem 3.2. Hence suppose statements 2. and 3. hold. Then by Theorem 3.2 there exists a normalized cross product bialgebra $B_1 \mathop{\varphi_{1,2}}^{\hat{\rho}} \mathop{\varphi_{2,1}}^{\hat{\sigma}} B_2$ which is isomorphic to B via the isomorphism $\Lambda = m_B \circ (i_1 \otimes i_2)$. We define

$$\begin{aligned}
m_2 & := p_2 \circ m_B \circ (i_2 \otimes i_2), & \Delta_1 & := (p_1 \otimes p_1) \circ \Delta_B \circ i_1, \\
\sigma & := p_1 \circ m_B \circ (i_2 \otimes i_2), & \rho & := (p_1 \otimes p_1) \circ \Delta_B \circ i_2, \\
\mu_l & := p_1 \circ m_B \circ (i_2 \otimes i_1), & \nu_l & := (p_2 \otimes p_1) \circ \Delta_B \circ i_1, \\
\mu_r & := p_2 \circ m_B \circ (i_2 \otimes i_1), & \nu_r & := (p_2 \otimes p_1) \circ \Delta_B \circ i_2.
\end{aligned} \tag{3.11}$$

The morphisms $m_1, \Delta_2, \eta_1, \eta_2, \varepsilon_1$, and ε_2 are given in terms of the projections and injections by the corresponding relations in (3.3) in Corollary 3.3. With the help of these data we define structure morphisms $\tilde{\varphi}_{1,2}, \tilde{\varphi}_{2,1}, \hat{\sigma}$, and $\hat{\rho}$ analogous to (3.8). We will show that the morphisms $m_{B_1 \otimes B_2}$ and $\Delta_{B_1 \otimes B_2}$ defined with these structure morphisms according to (3.9) precisely coincide with $m_{B_1 \mathop{\varphi_{1,2}}^{\hat{\rho}} \mathop{\varphi_{2,1}}^{\hat{\sigma}} B_2}$ and $\Delta_{B_1 \mathop{\varphi_{1,2}}^{\hat{\rho}} \mathop{\varphi_{2,1}}^{\hat{\sigma}} B_2}$ respectively, if the assumptions of statement 2. (or 3.) are satisfied. Before we prove this we will provide

several auxiliary identities. From (3.1) and Corollary 3.3 we obtain

$$\begin{aligned} (\Pi_2 \otimes \Pi_2) \circ \Delta_B \circ m_B \circ (\Pi_2 \otimes \text{id}_B) &= (\Pi_2 \otimes \Pi_2) \circ \Delta_B \circ \Pi_2 \circ m_B \circ (\Pi_2 \otimes \text{id}_B) \\ &= (\Pi_2 \otimes \Pi_2) \circ \Delta_B \circ m_B, \end{aligned} \quad (3.12)$$

$$\begin{aligned} \Pi_1 \circ m_B^{(3)} \circ (\Pi_1 \otimes \Pi_1 \otimes \Pi_1) &= \Pi_1 \circ m_B \circ (\text{id}_B \otimes \Pi_1 \circ m_B) \circ (\Pi_1 \otimes \Pi_1 \otimes \Pi_1) \\ &= \Pi_1 \circ m_B \circ (\text{id}_B \otimes \Pi_1 \circ m_B) \circ (\Pi_1 \otimes \Pi_1 \otimes \text{id}_B) \\ &= \Pi_1 \circ m_B^{(3)} \circ (\Pi_1 \otimes \Pi_1 \otimes \text{id}_B). \end{aligned} \quad (3.13)$$

Furthermore it holds

$$(\Pi_1 \otimes \text{id}_B) \circ \delta_{(1,0,2)} = (\Pi_1 \otimes \Pi_2) \circ \Delta_B \circ m_B \circ (\Pi_1 \otimes \text{id}_B) \quad (3.14)$$

since the following equations are satisfied.

$$\begin{aligned} m_B \circ (\Pi_1 \otimes \Pi_2) \circ (m_B \otimes \text{id}_B) \circ (\Pi_1 \otimes \Delta_B) &= m_B \circ (\Pi_1 \otimes \text{id}_B) \circ (m_B \circ (\Pi_1 \otimes \Pi_1) \otimes \Pi_2) \circ \\ &\quad \circ (\text{id}_B \otimes \Delta_B) \\ &= m_B \circ (\Pi_1 \otimes m_B \circ (\Pi_1 \otimes \Pi_2) \circ \Delta_B) \\ &= m_B \circ (\Pi_1 \otimes \text{id}_B). \end{aligned}$$

Then (3.14) follows from Theorem 3.2.3. Observe that Corollary 3.3 implies the additional conditions (3.10) have not been used to derive (3.14). Subsequently we will prove $m_{B_1 \otimes B_2} = m_{B_1 \varphi_{1,2} \hat{\rho} \hat{\sigma} \tilde{\varphi}_{2,1} B_2}$ graphically. Henceforth we use the notation $\Pi_j = \overset{j}{\uparrow}$ for $j \in \{1, 2\}$.

$$\begin{aligned} m_{B_1 \otimes B_2} &= \text{Diagram 1} = \text{Diagram 2} = \text{Diagram 3} \\ &= \text{Diagram 4} = \text{Diagram 5} = (\text{p}_1 \otimes \text{p}_2) \circ \Delta_B \circ m_B^{(4)} \circ (\text{i}_1 \otimes \text{i}_2 \otimes \text{i}_1 \otimes \text{i}_2) \\ &= \Lambda^{-1} \circ m_B \circ (\Lambda \otimes \Lambda) = m_{B_1 \varphi_{1,2} \hat{\rho} \hat{\sigma} \tilde{\varphi}_{2,1} B_2}. \end{aligned} \quad (3.15)$$

To derive the first identity of (3.15) we used the specific form of the structure morphisms $\tilde{\varphi}_{1,2}$, $\tilde{\varphi}_{2,1}$, $\hat{\sigma}$, and $\hat{\rho}$. Then (3.1), (3.2), and the third identity of (3.10) for $j = 1$ yield

the second equality of (3.15). With Theorem 3.2.2, (3.2) and (3.12) we derive the third equation, whereas (3.13) and again use of Theorem 3.2.2 yield the fourth identity of (3.15). The fifth equality comes from application of the second “projection” relation of (3.10), and for the derivation of the sixth identity we used (3.14). Finally the definition of Λ , given in the proof of Theorem 3.2, yields the result. In a π -symmetric way the identity $\Delta_{B_1 \otimes B_2} = \Delta_{B_1 \varphi_{1,2} \hat{\rho} \bowtie \hat{\sigma} \varphi_{2,1} B_2}$ will be shown. The (co-)unital identities (3.6) can be verified straightforwardly from the definitions, the assumptions and Corollary 3.3. It remains to prove the “projection” relations (3.7). Observe that

$$\Pi_1 = \Lambda \circ (\text{id}_{B_1} \otimes \eta_2 \circ \varepsilon_2) \circ \Lambda^{-1}, \quad \Pi_2 = \Lambda \circ (\eta_1 \circ \varepsilon_1 \otimes \text{id}_{B_2}) \circ \Lambda^{-1}. \quad (3.16)$$

Taking into account the relation $\Delta_B = (\Lambda \otimes \Lambda) \circ \Delta_{B_1 \varphi_{1,2} \hat{\rho} \bowtie \hat{\sigma} \varphi_{2,1} B_2} \circ \Lambda^{-1}$ and the relation $m_B = \Lambda \circ m_{B_1 \varphi_{1,2} \hat{\rho} \bowtie \hat{\sigma} \varphi_{2,1} B_2} \circ (\Lambda^{-1} \otimes \Lambda^{-1})$ one obtains with the help of (3.16) and (3.6)

$$\begin{aligned} \Delta_B \circ \Pi_1 &= (\Lambda \otimes \Lambda \circ (\text{id}_{B_1} \otimes \eta_2)) \circ (\text{id}_{B_1} \otimes \nu_l) \circ \Delta_1 \circ (\text{id}_{B_1} \otimes \varepsilon_2) \circ \Lambda, \\ \Pi_1 \circ m_B &= \Lambda \circ (\text{id} \otimes \eta_2) \circ m_1 \circ (m_1 \otimes \sigma) \circ (\text{id} \otimes \varphi_{2,1} \otimes \text{id}) \circ (\Lambda^{-1} \otimes \Lambda^{-1}). \end{aligned} \quad (3.17)$$

Gluing the identities of (3.17) and Π_1 according to the left and right hand side of the third equation of (3.10) (for $j = 1$), using (3.16) again, and eventually multiplying both resulting sides with $|\circ (\Lambda \otimes \Lambda) \circ (\eta_1 \otimes \text{id} \otimes \text{id} \otimes \eta_2)$ and $(\text{id} \otimes \varepsilon_2 \otimes \text{id} \otimes \varepsilon_2) \circ (\Lambda^{-1} \otimes \Lambda^{-1}) \circ |$ yields $(\mu_l \otimes \text{id}_{B_1}) \circ (\text{id}_{B_2} \otimes \Delta_1) = (m_1 \circ (\text{id}_{B_1} \otimes \sigma) \otimes \text{id}_{B_1}) \circ (\varphi_{2,1} \otimes \nu_l) \circ (\text{id}_{B_2} \otimes \Delta_1)$ which is the second identity of (3.7). Analogously the first equation of (3.7) can be derived. This proves that $B_1 \varphi_{1,2} \hat{\rho} \bowtie \hat{\sigma} \varphi_{2,1} B_2$ is cocycle cross product bialgebra.

Conversely suppose that $\Lambda : B_1 \varphi_{1,2} \hat{\rho} \bowtie \hat{\sigma} \varphi_{2,1} B_2 \rightarrow B$ is an isomorphism of bialgebras. To prove statement 2. it suffices to verify relations (3.10). Like in the proof “1. \Rightarrow 3.” of Theorem 3.2 we define $i_1 := \Lambda \circ (\text{id}_{B_1} \otimes \eta_2)$, $i_2 := \Lambda \circ (\eta_1 \otimes \text{id}_{B_2})$, $p_1 := (\text{id}_{B_1} \otimes \varepsilon_2) \circ \Lambda^{-1}$, and $p_2 := (\varepsilon_1 \otimes \text{id}_{B_2}) \circ \Lambda^{-1}$. Then the identities (3.17) will be proven for $\Pi_j := i_j \circ p_j$ in the way described above. And the previously performed gluing of the identities (3.17) yields

$$\begin{aligned} (\Pi_1 \otimes \text{id}) \circ \delta_{(0,0,0)} \circ (\text{id} \otimes \Pi_1) &= (\Lambda \circ (\text{id} \otimes \eta_2) \otimes \Lambda \circ (\text{id} \otimes \eta_2)) \circ (m_1 \circ (m_1 \otimes \sigma) \otimes \text{id}) \circ \\ &\quad \circ (\text{id} \otimes \varphi_{2,1} \otimes \nu_l) \circ (\Lambda^{-1} \otimes \Delta_1 \circ (\text{id} \otimes \varepsilon_2) \circ \Lambda^{-1}), \\ (\Pi_1 \otimes \text{id}) \circ \delta_{(0,1,0)} \circ (\text{id} \otimes \Pi_1) &= (\Lambda \circ (\text{id} \otimes \eta_2) \otimes \Lambda \circ (\text{id} \otimes \eta_2)) \circ (m_1 \circ (\text{id} \otimes \mu_l) \otimes \text{id}) \circ \\ &\quad \circ (\Lambda^{-1} \otimes \Delta_1 \circ (\text{id} \otimes \varepsilon_2) \circ \Lambda^{-1}). \end{aligned}$$

Therefore by assumption (3.7) the third identity of (3.10) for $j = 1$ follows. Applying π -symmetry the third equation of (3.10) for $j = 2$ will be derived. Then all conditions are satisfied which have been needed to derive the first four identities in (3.15). Hence

$$\begin{aligned} m_{B_1 \varphi_{1,2} \hat{\rho} \bowtie \hat{\sigma} \varphi_{2,1} B_2} &= (p_1 \circ m_B^{(3)} \otimes p_1 \circ m_B) \circ (\text{id}^{(2)} \otimes \Psi_{B,B} \circ (\text{id} \otimes \Pi_2) \otimes \text{id}) \circ \\ &\quad \circ (i_1 \otimes \Delta_B \circ m_B \circ (i_2 \otimes i_1) \otimes \Delta_B \circ i_2) \end{aligned}$$

On the other hand it holds by assumption $m_{B_1 \varphi_{1,2} \hat{\rho} \bowtie \hat{\sigma} \varphi_{2,1} B_2} = \Lambda^{-1} \circ m_B \circ (\Lambda \otimes \Lambda) = (p_1 \otimes p_2) \circ \Delta_B \circ m_B^{(4)} \circ (i_1 \otimes i_2 \otimes i_1 \otimes i_2) = (p_1 \otimes p_2) \circ \delta_{(0,0,0)} \circ (i_1 \otimes m_B^{(3)} \circ (i_2 \otimes i_1 \otimes i_2))$, where we used (3.14) which has been derived under the more general assumption of Theorem 3.2.

Multiplying both expressions of $m_{B_1 \varphi_{1,2} \hat{\rho} \otimes_{\varphi_{2,1}} B_2}$ with $|\circ(\eta_1 \otimes p_2 \otimes p_1 \otimes p_2)$ and $(i_1 \otimes i_2) \circ |$, and using that i_1 is algebra morphism yields

$$\begin{aligned} & (\Pi_1 \circ m_B \otimes \Pi_1 \circ m_B) \circ (\text{id} \otimes \Psi_{B,B} \circ (\text{id} \otimes \Pi_2) \otimes \text{id}) \circ (\Delta_B \circ m_B \circ (\Pi_2 \otimes \Pi_1) \otimes \Delta_B \circ \Pi_2) \\ &= (\Pi_1 \otimes \Pi_1) \circ \Delta_B \circ m_B^{(3)} \circ (\Pi_2 \otimes \Pi_1 \otimes \Pi_2) \end{aligned}$$

from which the second identity of (3.10) can be derived easily with the help of Theorem 3.2.2. Similarly by π -symmetry the first equation of (3.10) will be proven. \blacksquare

REMARK 9 The “projection” relations (γ_1) and (γ_2) in (3.10) can be derived from the identities

$$(\beta_1) : \beta_1 = \beta_1 \circ (\text{id} \otimes \Pi_1) \quad \text{and} \quad (\beta_2) : \beta_2 = (\Pi_2 \otimes \text{id}) \circ \beta_2 \quad (3.18)$$

where $\beta_1 := (\Pi_1 \circ m_B \otimes \text{id}) \circ (\text{id} \otimes \Psi_{B,B}) \circ (\Delta_B \circ \Pi_1 \otimes \text{id})$ and $\beta_2 := (\text{id} \otimes \Pi_2 \circ m_B) \circ (\Psi_{B,B} \otimes \text{id}) \circ (\text{id} \otimes \Delta_B \circ \Pi_2)$. However, in general the conditions (β_1) and (β_2) are not equivalent to the conditions (γ_1) and (γ_2) in Theorem 3.6.

PROPOSITION 3.7 *Under the equivalent conditions of Theorem 3.6 the following statements hold.*

1. $\mu_l = \varepsilon_2 \otimes \text{id}_{B_1}$ is trivial $\Leftrightarrow \Pi_1 \circ m_B \circ (\text{id}_B \otimes \Pi_1) = \Pi_1 \circ m_B \circ (\Pi_1 \otimes \Pi_1)$.
2. $\mu_r = \text{id}_{B_2} \otimes \varepsilon_1$ is trivial $\Leftrightarrow \Pi_2 \circ m_B = \Pi_2 \circ m_B \circ (\Pi_2 \otimes \Pi_2) \Leftrightarrow p_2$ is algebra morphism.
3. $\nu_l = \eta_2 \otimes \text{id}_{B_1}$ is trivial $\Leftrightarrow \Delta_B \circ \Pi_1 = (\Pi_1 \otimes \Pi_1) \circ \Delta_B \circ \Pi_1 \Leftrightarrow i_1$ is coalgebra morphism.
4. $\nu_r = \text{id}_{B_2} \otimes \eta_1$ is trivial $\Leftrightarrow (\Pi_2 \otimes \text{id}_B) \circ \Delta_B \circ \Pi_2 = (\Pi_2 \otimes \Pi_2) \circ \Delta_B \circ \Pi_2$.
5. $\sigma = \eta_1 \circ (\varepsilon_2 \otimes \varepsilon_2)$ is trivial $\Leftrightarrow \hat{\sigma} = \eta_1 \otimes m_2 \Leftrightarrow m_B \circ (\Pi_2 \otimes \Pi_2) = \Pi_2 \circ m_B \circ (\Pi_2 \otimes \Pi_2) \Leftrightarrow i_2$ is algebra morphism.
6. $\rho = (\eta_1 \otimes \eta_1) \circ \varepsilon_2$ is trivial $\Leftrightarrow \hat{\rho} = \Delta_1 \otimes \varepsilon_2 \Leftrightarrow (\Pi_1 \otimes \Pi_1) \circ \Delta_B = (\Pi_1 \otimes \Pi_1) \circ \Delta_B \circ \Pi_1 \Leftrightarrow p_1$ is coalgebra morphism.
7. μ_l and σ are trivial $\Leftrightarrow \Pi_1 \circ m_B = \Pi_1 \circ m_B \circ (\Pi_1 \otimes \Pi_1) \Leftrightarrow p_1$ is algebra morphism.
8. ν_r and ρ are trivial $\Leftrightarrow \Delta_B \circ \Pi_2 = (\Pi_2 \otimes \Pi_2) \circ \Delta_B \circ \Pi_2 \Leftrightarrow i_2$ is coalgebra morphism.

PROOF. We prove Proposition 3.7.1 and 3.7.5. The remaining statements can be derived in a similar manner or follow directly by π -symmetric reasoning. Without loss of generality we may assume $\Pi_1 = \text{id}_{B_1} \otimes \eta_2 \circ \varepsilon_2$ and $\Pi_2 = \eta_1 \circ \varepsilon_1 \otimes \text{id}_{B_2}$.

Ad 1.: Suppose that μ_l is trivial. This means that $(\text{id}_{B_1} \otimes \varepsilon_2) \circ \varphi_{2,1} = \varepsilon_2 \otimes \text{id}_{B_1}$. Using (2.2) and the unital identities of (2.3) then yields $\Pi_1 \circ m_B \circ (\text{id}_B \otimes \Pi_1) = (m_{B_1} \otimes \eta_2 \circ \varepsilon_2) \circ (\text{id}_{B_1} \otimes \varphi_{2,1} \otimes \varepsilon_2) = (m_{B_1} \otimes \eta_2) \circ (\text{id}_{B_1} \otimes \mu_l \otimes \varepsilon_2) = (m_{B_1} \otimes \eta_2) \circ (\text{id}_{B_1} \otimes \varepsilon_2 \otimes \text{id}_{B_1} \otimes \varepsilon_2)$. On the other hand from the unital identities of (2.3) immediately follows $\Pi_1 \circ m_B \circ (\Pi_1 \otimes \Pi_1) = (m_{B_1} \otimes \eta_2) \circ (\text{id}_{B_1} \otimes \varepsilon_2 \otimes \text{id}_{B_1} \otimes \varepsilon_2)$. Conversely if $\Pi_1 \circ m_B \circ (\text{id}_B \otimes \Pi_1) = \Pi_1 \circ m_B \circ (\Pi_1 \otimes \Pi_1)$ holds then analogous calculations as before yield $(m_{B_1} \otimes \eta_2 \circ \varepsilon_2) \circ (\text{id}_{B_1} \otimes \varphi_{2,1} \otimes \varepsilon_2) =$

$(m_{B_1} \otimes \eta_2) \circ (\text{id}_{B_1} \otimes \varepsilon_2 \otimes \text{id}_{B_1} \otimes \varepsilon_2)$. And then the triviality of $\mu_l = (\text{id}_{B_1} \otimes \varepsilon_2) \circ \varphi_{2,1}$ follows straightforwardly.

Ad 5.: Suppose that $m_B \circ (\Pi_2 \otimes \Pi_2) = \Pi_2 \circ m_B \circ (\Pi_2 \otimes \Pi_2)$. Then $m_2 = p_2 \circ m_B \circ (i_2 \otimes i_2)$ is associative for as $m_2 \circ (\text{id} \otimes m_2) = p_2 \circ m_B \circ (i_2 \otimes \Pi_2 \circ m_B \circ (i_2 \otimes i_2)) = p_2 \circ m_B \circ (\text{id}_B \otimes m_B) \circ (i_2 \otimes i_2 \otimes i_2) = p_2 \circ m_B \circ (m_B \otimes \text{id}_B) \circ (i_2 \otimes i_2 \otimes i_2) = m_2 \circ (m_2 \otimes \text{id}_{B_2})$. Hence B_2 is an algebra. And $i_2 \circ m_2 = \Pi_2 \circ m_B \circ (i_2 \otimes i_2) = m_B \circ (i_2 \otimes i_2)$. Therefore i_2 is algebra morphism. Conversely, if i_2 is algebra morphism then $m_B \circ (\Pi_2 \otimes \Pi_2) = i_2 \circ m_2 \circ (p_2 \otimes p_2) = i_2 \circ p_2 \circ i_2 \circ m_2 \circ (p_2 \otimes p_2) = \Pi_2 \circ m_B \circ (\Pi_2 \otimes \Pi_2)$.

Let i_2 be algebra morphism. Then from the last identity of (3.3) and the second identity of (3.2) we derive $\hat{\sigma} = (p_1 \otimes p_2) \circ \Delta_B \circ m_B \circ (i_2 \otimes i_2) = (p_1 \otimes p_2) \circ \Delta_B \circ i_2 \circ m_2 = \eta_1 \otimes m_2$. If on the other hand $\hat{\sigma} = \eta_1 \otimes m_2$ then we use (2.2) and (2.3) to obtain $m_B \circ (i_2 \otimes i_2) = m_B \circ (\eta_1 \otimes \text{id}_{B_2} \otimes \eta_1 \otimes \text{id}_{B_2}) = (\eta_1 \otimes \text{id}_{B_2}) \circ m_2 = i_2 \circ m_2$ which shows that i_2 is algebra morphism.

It is an easy exercise to prove that triviality of $\hat{\sigma}$ and triviality of σ are equivalent. \blacksquare

REMARK 10 Under the equivalent conditions of Theorem 3.2 similar results like in Proposition 3.7 can be shown for general cross product bialgebras if m_2 , Δ_1 , μ_l , μ_r , ν_l and ν_r will be defined formally as in (3.11).

Strong Cross Product Bialgebras

In the following we define strong cross product bialgebras. They will be studied in more detail in Theorems 4.4 and 4.5 in connection with so-called strong Hopf data. It turns out that strong cross product bialgebras are the central objects in our universal, (co-)modular co-cyclic theory of cross product bialgebras ⁵.

DEFINITION 3.8 A cocycle cross product bialgebra $B_1 \overset{\hat{\rho}}{\underset{\hat{\sigma}}{\bowtie}}_{\varphi_{2,1}} B_2$ is called strong cross product bialgebra if in addition the identities

$$\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} = \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array}, \quad \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} = \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} \quad (3.19)$$

$$\begin{array}{c} \text{Diagram 9} \\ \text{Diagram 10} \end{array} = \begin{array}{c} \text{Diagram 11} \\ \text{Diagram 12} \end{array}, \quad \begin{array}{c} \text{Diagram 13} \\ \text{Diagram 14} \end{array} = \begin{array}{c} \text{Diagram 15} \\ \text{Diagram 16} \end{array}, \quad \begin{array}{c} \text{Diagram 17} \\ \text{Diagram 18} \end{array} = \begin{array}{c} \text{Diagram 19} \\ \text{Diagram 20} \end{array}, \quad \begin{array}{c} \text{Diagram 21} \\ \text{Diagram 22} \end{array} = \begin{array}{c} \text{Diagram 23} \\ \text{Diagram 24} \end{array} \quad (3.20)$$

$$\begin{array}{c} \text{Diagram 25} \\ \text{Diagram 26} \end{array} = \begin{array}{c} \text{Diagram 27} \\ \text{Diagram 28} \end{array}, \quad \begin{array}{c} \text{Diagram 29} \\ \text{Diagram 30} \end{array} = \begin{array}{c} \text{Diagram 31} \\ \text{Diagram 32} \end{array} \quad (3.21)$$

are fulfilled. We denote strong cross product bialgebras by ⁶ $B_1 \overset{\hat{h}}{\bowtie} B_2$.

⁵strong cross product bialgebras and strong Hopf data provide a unifying universal and (co-)modular co-cyclic theory of cross product bialgebras. But they are probably not the most general setting to meet the same demands. Therefore our notation “strong” should be specified further. However, in order to avoid terminological blow-up we will henceforth use our notation, always having in mind that there might be a weaker definition of “strong”. This is certainly an interesting direction for further study.

⁶The deeper meaning of this notation will become clear in Section 4.

REMARK 11 (!) A tedious calculation shows that the “projection” relations (γ_1) , (γ_2) , (δ_1) , and (δ_2) in (3.10) as well as (3.7) are redundant for strong cross product bialgebras. Therefore, if we use the morphisms $m_{i,jk}$ and $\Delta_{ij,k}$ from Corollary 3.4 an alternative definition of strong cross product bialgebras can be given as follows.

A strong cross product bialgebra is a cross product bialgebra for which the identities (3.19), (3.20) and (3.21) formally hold for the corresponding morphisms $m_{i,jk}$ and $\Delta_{ij,k}$.

We call a strong cross product bialgebra *regular* if all defining identities have rank $(2, 2)$, $(1, 3)$ or $(3, 1)$. We call it *pure* if (3.19), (3.20) and (3.21) are redundant.

Any bialgebra B which is isomorphic to a certain cocycle cross product bialgebra has injections and projections i_1, i_2, p_1 , and p_2 which are uniquely determined by the cocycle cross product bialgebra and the given isomorphism (see Theorem 3.6 and Remark 6). In the special case of strong cross product bialgebras the structure morphisms obey the additional identities (3.19) – (3.21). Using (3.3) and (3.11) these relations can be translated easily into equations of the idempotents Π_1 and Π_2 of the bialgebra B as follows.

$$\begin{aligned}
& (\Pi_2 \otimes \text{id}_B) \circ \Phi_{\text{id}_B} \circ (\Pi_1 \otimes \Pi_1) = (\Pi_2 \otimes \text{id}_B) \circ \Delta_B \circ \Pi_1 \otimes \varepsilon_B, \\
& ((\Pi_1 \otimes \Pi_1) \circ \Phi_{\Pi_2} \otimes \text{id}_B) \circ (\Pi_2 \otimes \Psi_{B,B}) \circ (\Delta_B \circ \Pi_1 \otimes \Pi_2) = \eta_B \otimes \eta_B \otimes \Pi_1 \otimes \varepsilon_B, \\
& (\Pi_1 \otimes \text{id}_B) \circ \delta_{(2,2,0)} \circ (\text{id}_B \otimes \Pi_1) = \eta_B \circ \varepsilon_B \otimes \Pi_1, \\
& (\Pi_1 \otimes \Pi_2 \circ m_B \circ (\text{id}_B \otimes \Pi_2) \otimes \text{id}_B \otimes \Pi_1) \circ (\Psi_{B,B} \otimes \Phi_{\Pi_2} \otimes \text{id}_B) \circ \quad (3.22) \\
& \quad \circ (\text{id}_B \otimes (\Psi_{B,B} \otimes \text{id}_B \otimes \text{id}_B) \circ (\Pi_1 \otimes (\Delta_B \circ \Pi_1 \otimes \text{id}_B) \circ \Delta_B \circ \Pi_2)) \\
& = ((\Pi_1 \otimes \Pi_2 \circ m_B) \circ (\Psi_{B,B} \otimes \text{id}_B) \circ (\text{id}_B \otimes \Delta_B) \otimes \Pi_1 \otimes \Pi_1) \circ \\
& \quad \circ (\text{id}_B \otimes (\Psi_{B,B} \otimes \text{id}_B) \circ (\text{id}_B \otimes (\Pi_1 \otimes \text{id}_B) \circ \Delta_B \circ \Pi_2))
\end{aligned}$$

and the corresponding π -symmetric counterparts.

We used $\Phi_f := (m_B \otimes \text{id}_B) \circ (f \otimes \Psi_{B,B}) \circ (\Delta_B \otimes \text{id}_B)$ for $f : B \rightarrow B$, and $\delta_{(i,j,k)}$ of Theorem 3.6.

Recall Remark 5 and observe that our construction of (strong/cocycle) cross product bialgebras is invariant under π -symmetry. However, cross product bialgebras in general fail to be invariant if duality or rotational symmetry along a vertical axis (in the plane of the paper) will be transformed separately. Without problems the dual versions of cross product bialgebras can be defined using our original definition. The corresponding results follow immediately. If the category \mathcal{C} has (right) duality then the following proposition shows how dual cross product bialgebras can be constructed explicitly.

PROPOSITION 3.9 *Suppose that \mathcal{C} is a braided category with duality functor $(-)^{\vee} : \mathcal{C} \rightarrow \mathcal{C}_{\text{op}}^{\text{op}}$ where $\mathcal{C}_{\text{op}}^{\text{op}}$ is the opposite category with opposite tensor product. If $(B; B_1, B_2, i_1, i_2, p_1, p_2)$ is a (strong/cocycle) cross product bialgebra in \mathcal{C} then the dual tuple $(B^{\vee}; B_1^{\vee}, B_2^{\vee}, i_1^{\vee}, i_2^{\vee}, p_1^{\vee}, p_2^{\vee})$ is a dual (strong/cocycle) cross product bialgebra in $\mathcal{C}_{\text{op}}^{\text{op}}$. \blacksquare*

4 Hopf Data

In Section 3 we studied cross product bialgebras from a universal point of view. We answered the question under which conditions a bialgebra is isomorphic to a cross product bialgebra. Now we present an explicit (co-)modular co-cyclic construction method in terms of Hopf data. A Hopf datum consists of two objects with certain interrelated (co-)modular co-cyclic identities. In Theorem 4.4 we will show that so-called strong Hopf data and strong cross product bialgebras are different descriptions of the same objects. Universality of our “strong” construction will be demonstrated in Theorem 4.5. We postpone the lengthy proof of Theorem 4.4 to Section 5. In the subsequent definition of Hopf data occur two objects B_1 and B_2 , and morphisms

$$\begin{aligned}
m_1 : B_1 \otimes B_1 &\rightarrow B_1, & m_2 : B_2 \otimes B_2 &\rightarrow B_2, \\
\Delta_1 : B_1 &\rightarrow B_1 \otimes B_1, & \Delta_2 : B_2 &\rightarrow B_2 \otimes B_2, \\
\eta_1 : \mathbb{1} &\rightarrow B_1, & \eta_2 : \mathbb{1} &\rightarrow B_2, \\
\varepsilon_1 : B_1 &\rightarrow \mathbb{1}, & \varepsilon_2 : B_2 &\rightarrow \mathbb{1}, \\
\mu_l : B_2 \otimes B_1 &\rightarrow B_1, & \nu_l : B_1 &\rightarrow B_2 \otimes B_1, \\
\mu_r : B_2 \otimes B_1 &\rightarrow B_2, & \nu_r : B_2 &\rightarrow B_2 \otimes B_1, \\
\sigma : B_2 \otimes B_2 &\rightarrow B_1, & \rho : B_2 &\rightarrow B_1 \otimes B_1.
\end{aligned} \tag{4.1}$$

Again we use the graphical presentation of Figure 1 for these morphisms, and we represent σ and ρ by $\sigma = \begin{array}{c} \bullet \\ \downarrow \end{array} : B_2 \otimes B_2 \rightarrow B_1$ and $\rho = \begin{array}{c} \bullet \\ \uparrow \end{array} : B_2 \rightarrow B_1 \otimes B_1$. Similarly as in (3.8) we define morphisms $\varphi_{1,2}$, $\varphi_{2,1}$, $\hat{\sigma}$, and $\hat{\rho}$ by

$$\begin{aligned}
\varphi_{1,2} &:= (m_2 \otimes m_1) \circ (\text{id}_{B_1} \otimes \Psi_{B_1, B_2} \otimes \text{id}_{B_2}) \circ (\nu_l \otimes \nu_r), \\
\varphi_{2,1} &:= (\mu_l \otimes \mu_r) \circ (\text{id}_{B_2} \otimes \Psi_{B_2, B_1} \otimes \text{id}_{B_1}) \circ (\Delta_2 \otimes \Delta_1), \\
\hat{\sigma} &:= (\sigma \otimes m_2 \circ (\mu_r \otimes \text{id}_{B_2})) \circ (\text{id}_{B_2} \otimes \Psi_{B_2, B_2} \otimes \text{id}_{B_1 \otimes B_2}) \circ (\Delta_2 \otimes (\nu_r \otimes \text{id}_{B_2}) \circ \Delta_2), \\
\hat{\rho} &:= (m_1 \circ (\text{id}_{B_1} \otimes \mu_l) \otimes m_1) \circ (\text{id}_{B_1 \otimes B_2} \otimes \Psi_{B_1, B_1} \otimes \text{id}_{B_1}) \circ ((\text{id}_{B_1} \otimes \nu_l) \circ \Delta_1 \otimes \rho)
\end{aligned}$$

or graphically

$$\varphi_{1,2} := \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \quad , \quad \varphi_{2,1} := \begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} \quad , \quad \hat{\rho} := \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \quad , \quad \hat{\sigma} := \begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} \tag{4.2}$$

Occasionally we also use the graphical abbreviations

$$\varphi_{1,2} = \begin{array}{c} \text{---} \\ \text{---} \end{array} \quad , \quad \varphi_{2,1} = \begin{array}{c} \text{---} \\ \text{---} \end{array} \quad , \quad \hat{\rho} = \begin{array}{c} \text{---} \\ \text{---} \end{array} \quad , \quad \hat{\sigma} = \begin{array}{c} \text{---} \\ \text{---} \end{array} \tag{4.3}$$

DEFINITION 4.1 *The tuple $\mathfrak{h} = ((B_1, m_1, \eta_1, \Delta_1, \varepsilon_1), (B_2, m_2, \eta_2, \Delta_2, \varepsilon_2); \mu_l, \mu_r, \nu_l, \nu_r, \rho, \sigma)$ is called Hopf datum if*

1. (B_1, m_1, η_1) is an algebra and $\varepsilon_1 : B_1 \rightarrow \mathbb{1}$ is an algebra morphism.
2. $(B_2, \Delta_2, \varepsilon_2)$ is a coalgebra and $\eta_2 : \mathbb{1} \rightarrow B_2$ is a coalgebra morphism.
3. (B_1, ν_l) is left B_2 -comodule.

4. (B_2, μ_r) is right B_1 -module.

5. The identities

$$\begin{aligned}
\Delta_1 \circ \eta_1 &= \eta_1 \otimes \eta_2, & (\text{id}_{B_1} \otimes \varepsilon_1) \circ \Delta_1 &= (\varepsilon_1 \otimes \text{id}_{B_1}) \circ \Delta_1 = \text{id}_{B_1}, \\
\varepsilon_2 \circ \mathfrak{m}_2 &= \varepsilon_2 \otimes \varepsilon_2, & \mathfrak{m}_2 \circ (\text{id}_{B_2} \otimes \eta_2) &= \mathfrak{m}_2 \circ (\eta_2 \otimes \text{id}_{B_2}) = \text{id}_{B_2}, \\
\mu_r \circ (\eta_2 \otimes \text{id}_{B_1}) &= \eta_2 \circ \varepsilon_1, & \varepsilon_2 \circ \mu_r &= \varepsilon_2 \otimes \varepsilon_1, \\
(\text{id}_{B_2} \otimes \varepsilon_1) \circ \nu_l &= \eta_2 \circ \varepsilon_1, & \nu_l \circ \eta_1 &= \eta_2 \otimes \eta_1, \\
\mu_l \circ (\eta_2 \otimes \text{id}_{B_1}) &= \text{id}_{B_1}, & \mu_l \circ (\text{id}_{B_2} \otimes \eta_1) &= \eta_1 \circ \varepsilon_2, & \varepsilon_1 \circ \mu_l &= \varepsilon_2 \otimes \varepsilon_1, \\
(\text{id}_{B_2} \otimes \varepsilon_1) \circ \nu_r &= \text{id}_{B_2}, & (\varepsilon_2 \otimes \text{id}_{B_1}) \circ \nu_r &= \eta_1 \circ \varepsilon_2, & \nu_r \circ \eta_2 &= \eta_2 \otimes \eta_1, \\
\sigma \circ (\eta_2 \otimes \text{id}_{B_2}) &= \sigma \circ (\text{id}_{B_2} \otimes \eta_2) = \eta_1 \circ \varepsilon_2, & \varepsilon_1 \circ \sigma &= \varepsilon_2 \otimes \varepsilon_2, \\
(\text{id}_{B_1} \otimes \varepsilon_1) \circ \rho &= (\varepsilon_1 \otimes \text{id}_{B_1}) \circ \rho = \eta_1 \circ \varepsilon_2, & \rho \circ \eta_2 &= \eta_1 \otimes \eta_1
\end{aligned} \tag{4.4}$$

are satisfied.

6. The subsequent compatibility relations hold.



Weak associativity of \mathfrak{m}_2 ,

Weak coassociativity of Δ_1 ,




Weak associativity of μ_l ,

Weak coassociativity of ν_r ,



Module-algebra compatibility,

Comodule-coalgebra compatibility,



Cocycle and cycle compatibilities,

$$B_1 B_1 = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}, \quad B_2 B_2 = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$$

Algebra-coalgebra compatibility,

$$\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}, \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}, \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$$

Module-coalgebra compatibility,

Comodule-algebra compatibility,

$$\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$$

Module-comodule compatibility,

$$\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$$

Cycle-cocycle compatibility.

(End of Definition 4.1)

Observe that Definition 4.1 is π -symmetric invariant. It will be verified in the next proposition that every cocycle cross product bialgebra canonically induces a Hopf datum.

PROPOSITION 4.2 *Let $B_1 \underset{\varphi_{1,2}}{\overset{\hat{\rho}}{\bowtie}} \underset{\varphi_{2,1}}{\overset{\hat{\sigma}}{\bowtie}} B_2$ be a cocycle cross product bialgebra with corresponding structure morphisms $m_1, \eta_1, \Delta_1, \varepsilon_1, m_2, \eta_2, \Delta_2, \varepsilon_2, \mu_l, \mu_r, \nu_l, \nu_r, \rho,$ and σ . Then $((B_1, m_1, \eta_1, \Delta_1, \varepsilon_1), (B_2, m_2, \eta_2, \Delta_2, \varepsilon_2); \mu_l, \mu_r, \nu_l, \nu_r, \rho, \sigma)$ is a Hopf datum.*

PROOF. By definition cocycle cross product bialgebras are cocycle cross product algebras and cycle cross product coalgebras in particular. Hence $(B_1, m_1 \eta_1)$ is an algebra and $(B_2, \Delta_2, \varepsilon_2)$ is a coalgebra. Since cocycle cross product bialgebras are normalized one deduces with (3.6) that $\eta_2 : \mathbb{1} \rightarrow B_2$ is an algebra morphism and $\varepsilon_1 : B_1 \rightarrow \mathbb{1}$ is a coalgebra morphism. Then the conditions of Definition 4.1.1 and 4.1.2 hold. The first sixteen (co-)unital identities of (4.4) hold by assumption (see (3.6)). Then from (3.8) one easily derives

$$\begin{aligned} \varphi_{1,2} \circ (\eta_1 \otimes \text{id}_{B_2}) &= \nu_r, & (\varepsilon_1 \otimes \text{id}_{B_2}) \circ \varphi_{2,1} &= \mu_r, & \hat{\rho} \circ (\eta_1 \otimes \text{id}_{B_1}) &= \rho \\ \varphi_{1,2} \circ (\text{id}_{B_1} \otimes \eta_2) &= \nu_l, & (\text{id}_{B_1} \otimes \varepsilon_2) \circ \varphi_{2,1} &= \mu_l, & (\text{id}_{B_2} \otimes \varepsilon_2) \circ \hat{\sigma} &= \sigma. \end{aligned} \quad (4.5)$$

Since $B_1 \hat{\varphi}_{1,2}^{\hat{\sigma}} \hat{\bowtie}_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ is especially cocycle cross product algebra, the identities (2.3) are satisfied. Composing the fifth equation of (2.3) with $(\varepsilon_1 \otimes \text{id}_{B_2}) \circ |$ and using (4.5) proves that (B_2, μ_r) is right B_1 -module. Similarly the weak associativity of μ_l is shown by composing with $(\text{id}_{B_1} \otimes \varepsilon_2) \circ |$. From (4.5) and (2.3) it will be concluded straightforwardly that $\sigma \circ (\eta_2 \otimes \text{id}_{B_2}) = (\text{id}_{B_1} \otimes \varepsilon_2) \circ \hat{\sigma} \circ (\eta_2 \otimes \text{id}_{B_2}) = (\text{id}_{B_1} \otimes \varepsilon_2) \circ (\eta_1 \otimes \text{id}_{B_2}) = \eta_1 \circ \varepsilon_2$ and similarly $\sigma \circ (\eta_2 \otimes \text{id}_{B_2}) = \eta_1 \circ \varepsilon_2$. Since $B_1 \hat{\varphi}_{1,2}^{\hat{\sigma}} \hat{\bowtie}_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ is cross product bialgebra, $\varepsilon_1 \otimes \varepsilon_2$ is algebra morphism, and therefore one easily shows that $\varepsilon_1 \circ \sigma = \varepsilon_2 \otimes \varepsilon_2$. Eventually all other (co-)unital identities of (4.4) follow then by π -symmetry. The weak associativity for m_2 follows now from the sixth equation of (2.3) by adjoining $(\varepsilon_1 \otimes \text{id}_{B_2}) \circ |$ on both sides. The module-algebra compatibility will be derived from the fifth equation of (2.3) through composition with $(\text{id}_{B_1} \otimes \varepsilon_2) \circ |$. Making use of the bialgebra identity $\Delta_B \circ m_B = (m_B \otimes m_B) \circ (\text{id}_B \otimes \Psi_{B,B} \otimes \text{id}_B) \circ (\Delta_B \otimes \Delta_B)$ for $B = B_1 \hat{\varphi}_{1,2}^{\hat{\sigma}} \hat{\bowtie}_{\varphi_{2,1}}^{\hat{\sigma}} B_2$ we prove the algebra-coalgebra compatibility by application of $| \circ (\text{id}_{B_1} \otimes \eta_2 \otimes \text{id}_{B_1} \otimes \eta_2)$ and $(\text{id}_{B_1} \otimes \text{id}_{\varepsilon_2} \otimes \text{id}_{B_1} \otimes \varepsilon_2) \circ |$ on both sides of the bialgebra identity. The first equation of the module-coalgebra compatibility will be shown similarly by composing with $| \circ (\eta_1 \otimes \text{id}_{B_2} \otimes \text{id}_{B_1} \otimes \eta_2)$ and $(\varepsilon_1 \otimes \text{id}_{B_2} \otimes \varepsilon_1 \otimes \text{id}_{B_2}) \circ |$. Application of $| \circ (\eta_1 \otimes \text{id}_{B_2} \otimes \text{id}_{B_1} \otimes \eta_2)$ and $(\text{id}_{B_1} \otimes \varepsilon_2 \otimes \text{id}_{B_1} \otimes \varepsilon_2) \circ |$ yields the second equation of the module-coalgebra compatibility. The module-comodule compatibility is derived from the bialgebra identity by composing with $| \circ (\eta_1 \otimes \text{id}_{B_2} \otimes \text{id}_{B_1} \otimes \eta_2)$ and $(\varepsilon_1 \otimes \text{id}_{B_2} \otimes \text{id}_{B_1} \otimes \varepsilon_2) \circ |$, whereas the cycle-cocycle compatibility comes from composition with $| \circ (\eta_1 \otimes \text{id}_{B_2} \otimes \eta_1 \otimes \text{id}_{B_2})$ and $(\text{id}_{B_1} \otimes \varepsilon_2 \otimes \text{id}_{B_1} \otimes \varepsilon_2) \circ |$. Application of $(\text{id}_{B_1} \otimes \varepsilon_2) \circ |$ to the sixth equation of (2.3) and use of the sixth identity of (4.5) yield the cocycle compatibility. All remaining compatibility relations of Definition 4.1 will be proven by application of π -symmetry to the former results. \blacksquare

Strong Hopf Data

The converse of Proposition 4.2 is not true in general. However, in the following we show that so-called strong Hopf data yield cross product bialgebras. The definition of strong Hopf data is closely related to the definition of strong cross product bialgebras.

DEFINITION 4.3 *A Hopf datum \mathfrak{h} is called strong if in addition the following identities hold.*

$$\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} = \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array}, \quad \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} = \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} \quad (4.6)$$

$$\begin{array}{c} \text{Diagram 9} \\ \text{Diagram 10} \end{array} = \begin{array}{c} \text{Diagram 11} \\ \text{Diagram 12} \end{array}, \quad \begin{array}{c} \text{Diagram 13} \\ \text{Diagram 14} \end{array} = \begin{array}{c} \text{Diagram 15} \\ \text{Diagram 16} \end{array}, \quad \begin{array}{c} \text{Diagram 17} \\ \text{Diagram 18} \end{array} = \begin{array}{c} \text{Diagram 19} \\ \text{Diagram 20} \end{array}, \quad \begin{array}{c} \text{Diagram 21} \\ \text{Diagram 22} \end{array} = \begin{array}{c} \text{Diagram 23} \\ \text{Diagram 24} \end{array} \quad (4.7)$$

$$\begin{array}{c} \text{Diagram 25} \\ \text{Diagram 26} \end{array} = \begin{array}{c} \text{Diagram 27} \\ \text{Diagram 28} \end{array}, \quad \begin{array}{c} \text{Diagram 29} \\ \text{Diagram 30} \end{array} = \begin{array}{c} \text{Diagram 31} \\ \text{Diagram 32} \end{array} \quad (4.8)$$

Strong Hopf data are invariant under π -symmetry. This fact will be used extensively in Section 5 where we prove Theorem 4.4.

REMARK 12 Observe that except of (4.8) and the first and the second identity of (4.7) the defining identities of a strong Hopf datum are either of rank (2, 2), (1, 3) or (3, 1).

Therefore in the same way as in Remark 11 we call a strong Hopf datum *regular* if the defining identities are of either rank (2, 2), (1, 3) or (3, 1). We call the strong Hopf datum *pure* if (4.6), (4.7) and (4.8) are redundant.

Main Results

This is the central part of the article. In the subsequent Theorems 4.4 and 4.5 we present the universal (co-)modular co-cyclic theory of (strong) cross product bialgebras. In Theorem 4.4 we describe the (co-)modular co-cyclic construction of strong cross product bialgebras. Theorem 4.5 exhibits its universality.

THEOREM 4.4 *Let $\mathfrak{h} = ((B_1, m_1, \eta_1, \Delta_1, \varepsilon_1), (B_2, m_2, \eta_2, \Delta_2, \varepsilon_2); \mu_l, \mu_r, \nu_l, \nu_r, \rho, \sigma)$ be a strong Hopf datum. Then $B_1 \otimes B_2$ with $\varphi_{1,2}$, $\varphi_{2,1}$, $\hat{\sigma}$, and $\hat{\rho}$ defined according to (4.2) is a strong cross product bialgebra $B_1 \overset{\mathfrak{h}}{\bowtie} B_2$. Strong Hopf data and strong cross product bialgebras are in one-to-one correspondence. ■*

Every strong Hopf datum \mathfrak{h} therefore induces a strong cross product bialgebra $B_1 \overset{\mathfrak{h}}{\bowtie} B_2$. For any bialgebra B which is isomorphic to $B_1 \overset{\mathfrak{h}}{\bowtie} B_2$ the additional relations (4.6) – (4.8) (or by one-to-one correspondence the equations (3.19) – (3.21)) imply the additional relations (3.22) for the idempotents Π_1 and Π_2 of B .

THEOREM 4.5 *Let B be a bialgebra in \mathcal{C} . Then the following statements are equivalent.*

1. *There is a strong Hopf datum \mathfrak{h} such that the corresponding strong cross product bialgebra $B_1 \overset{\mathfrak{h}}{\bowtie} B_2$ is bialgebra isomorphic to B .*
2. *There are idempotents $\Pi_1, \Pi_2 \in \text{End}(B)$ such that the conditions of Theorem 3.2.2 and the “strong projection” relations (3.22) hold.*
3. *There are objects B_1 and B_2 and morphisms $B_1 \xrightarrow{i_1} B \xrightarrow{p_1} B_1$ and $B_2 \xrightarrow{i_2} B \xrightarrow{p_2} B_2$ such that the conditions of Theorem 3.2.3 and the “strong projection” relations (3.22) hold with $\Pi_j := i_j \circ p_j$.*

PROOF. Theorem 4.5 can be derived straightforwardly from Theorem 4.4, Theorem 3.2, (Theorem 3.6), Remark 6, Remark 11 and the relations (3.22). ■

In the sequel we will consider special versions of strong cross product bialgebras. They admit equivalent universal and (co-)modular co-cyclic descriptions and can be derived from the most general construction given in Theorem 4.4. In particular all known constructions of cocycle cross product bialgebras will be recovered, and additionally we find several new types of cocycle cross product bialgebras. In Proposition 4.6 we describe comprehensively the universal and (co-)modular co-cyclic properties of one of these special types.

Theorem 4.4 and Proposition 3.7 provide the necessary tools to describe the various subclasses. We distinguish the different special versions of strong cross product bialgebras

with the help of the boxes $\begin{array}{|c|c|} \hline \mu_l & \mu_r \\ \hline \sigma & \rho \\ \hline \nu_l & \nu_r \\ \hline \end{array}$ where the particular entries will be left blank “ ” or take the values \bullet for the (weak) (co-)actions, \blacktriangledown for σ and \blacktriangle for ρ dependent on the respective










Figure 2: Graph representing the various special versions of strong cross product bialgebras.

morphisms are trivial or not. For example a strong cross product bialgebra with trivial cocycle $\sigma = \eta_1 \circ (\varepsilon_2 \otimes \varepsilon_2)$ and trivial right coaction $\nu_r = \text{id}_{B_2} \otimes \eta_1$ will be represented by the classification box $\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \square & \blacktriangle \\ \hline \bullet & \square \\ \hline \end{array}$. Thereby $2^6 = 64$ types of special strong cross products can be obtained; $2^4 = 16$ of them are co-cycle free. In a similar (dual) way the dual strong cross product bialgebras will be denoted by a classification box $\begin{array}{|c|c|} \hline \mu_l & \nu_r \\ \hline \sigma & \rho \\ \hline \nu_l & \nu_r \\ \hline \end{array}$ with entries \bullet for the (co)actions, \blacktriangle for the cocycle σ , and \blacktriangledown for the cycle ρ . Then analogously 64 special types of dual strong cross product bialgebras can be obtained from which $2^4 = 16$ (co-cycle free) types coincide with the corresponding 16 types of strong cross product bialgebras. Therefore the total number of different types of strong and dual strong cross product bialgebras which is classified by our scheme is $2^6 + (2^6 - 2^4) = 7 \cdot 2^4 = 112$. However, henceforth we do not tell between a certain type and its dual or π -symmetric counterparts

- it is an easy exercise to obtain one from the other⁷. Up to this $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetry we obtain 33 different classification boxes. In Figure 2 we present a graph where the boxes (modulo $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetry) are the vertices, and each box is linked with its descending “next neighbour” special versions. We stratify the whole graph into three layers. The layers consist of boxes with 2, 1 or 0 co-cycles respectively. The most general type of strong cross product bialgebra (Theorem 4.4) is at the top of the graph in the first row. In the second row all descendants with one trivial morphism $\mu_l, \mu_r, \sigma, \rho, \nu_l$, or ν_r are listed, in the third row all special types with two trivial morphisms are listed, etc. The cross product bialgebra constructions from [24, 18, 22, 20, 5] are descendants of the special cases  and  which are the co-cycle free cross product bialgebras of [5] and the braided versions of bicrossed product bialgebras [20, 22] respectively⁸. In the following tables we describe the different types of cocycle cross product bialgebras in more detail; we omit the description of the various cross product bialgebras studied in [24, 18, 22, 20, 5]. In the second column of the tables the structure of the multiplication m_B and the comultiplication Δ_B is given. In the third column we recall the corresponding equivalent properties of Proposition 3.7, and in the last column the status of the strong cross product bialgebra is listed.

⁷Observe that a π -rotation of the classification box in the plane of the graphic yields the corresponding π -symmetric counterpart of a certain type of cross product bialgebra. And analogously a reflection of the classification box along a horizontal line yields the dual counterpart.

⁸A special symmetric version of  has been studied recently in [26].

Type	Structure morphisms m_B, Δ_B	Proposition 3.7	Status
			
		3.7.4	
		3.7.3	
		3.7.6	
		3.7.1, 3.7.4	regular
		3.7.3, 3.7.4	

Type	Structure morphisms m_B, Δ_B	Proposition 3.7	Status
		3.7.2, 3.7.4	regular
		3.7.2, 3.7.3	pure
		3.7.2, 3.7.6	
		3.7.3, 3.7.6	pure
		3.7.8	regular
		3.7.1, 3.7.5	
		3.7.5, 3.7.6	regular
		3.7.6, 3.7.7	pure

Type	Structure morphisms m_B, Δ_B	Proposition 3.7	Status
		3.7.1, 3.7.3, 3.7.4	regular
		3.7.2, 3.7.8	regular
		3.7.3, 3.7.8	pure
		3.7.1, 3.7.2, 3.7.6	
		3.7.1, 3.7.3, 3.7.6	pure
		3.7.1, 3.7.8	regular
		3.7.1, 3.7.2, 3.7.8	regular
		3.7.1, 3.7.3, 3.7.8	pure

There are various redundant and special relations among the defining identities of the particular cases listed above. They can be derived from Theorem 4.4.

Exemplarily we will describe the explicit structure of the cross product bialgebras of type

1. $(B_1, m_1, \eta_1, \Delta_1, \varepsilon_1)$ is a bialgebra.
2. $(B_2, \Delta_2, \varepsilon_2, \mu_r, \nu_r)$ is B_1 -module coalgebra and B_1 -comodule coalgebra and $\eta_2 : \mathbb{1} \rightarrow B_2$ is coalgebra morphism.
- 3.

$$\begin{aligned}
\varepsilon_2 \circ m_2 &= \varepsilon_2 \otimes \varepsilon_2, & m_2 \circ (\text{id}_{B_2} \otimes \eta_2) &= m_2 \circ (\eta_2 \otimes \text{id}_{B_2}) = \text{id}_{B_2}, \\
\mu_r \circ (\eta_2 \otimes \text{id}_{B_1}) &= (\text{id}_{B_2} \otimes \varepsilon_1) \circ \nu_l = \eta_2 \circ \varepsilon_1, & \nu_l \circ \eta_1 &= \eta_2 \otimes \eta_1, \\
\mu_l \circ (\eta_2 \otimes \text{id}_{B_1}) &= \text{id}_{B_1}, & \mu_l \circ (\text{id}_{B_2} \otimes \eta_1) &= \eta_1 \circ \varepsilon_2, \\
\varepsilon_1 \circ \mu_l &= \varepsilon_2 \otimes \varepsilon_1, & \nu_r \circ \eta_2 &= \eta_2 \otimes \eta_1, \\
\sigma \circ (\eta_2 \otimes \text{id}_{B_2}) &= \sigma \circ (\text{id}_{B_2} \otimes \eta_2) = \eta_1 \circ \varepsilon_2, & \varepsilon_1 \circ \sigma &= \varepsilon_2 \otimes \varepsilon_2, \\
(\text{id}_{B_1} \otimes \varepsilon_1) \circ \rho &= (\varepsilon_1 \otimes \text{id}_{B_1}) \circ \rho = \eta_1 \circ \varepsilon_2, & \rho \circ \eta_2 &= \eta_1 \otimes \eta_1.
\end{aligned} \tag{4.9}$$

4. The weak associativity of m_2 and of μ_l in Definition 4.1 hold.
5. The module-algebra compatibility of Definition 4.1 is satisfied.
6. The cocycle compatibility of $\hat{\sigma}$ holds.
7. The algebra-coalgebra compatibility of B_2 holds.
8. The module-coalgebra compatibility of μ_l , the comodule-algebra compatibility of ν_r , the module-comodule compatibility and the cycle-cocycle compatibility are respectively given by

$$\begin{aligned}
\Delta_1 \circ \mu_l &= (\mu_l \otimes m_1 \circ (\text{id}_{B_1} \otimes \mu_l)) \circ (\text{id}_{B_2} \otimes \Psi_{B_1 \otimes B_2, B_1} \otimes \text{id}_{B_1}) \circ \\
&\quad \circ ((\nu_r \otimes \text{id}_{B_2}) \circ \Delta_2 \otimes \Delta_1), \\
(\text{id}_{B_2} \otimes m_1) \circ (\Psi_{B_1, B_2} \otimes \text{id}_{B_1}) \circ (\text{id}_{B_1} \otimes \nu_r) \circ \hat{\sigma} \\
&= (m_2 \otimes m_1 \circ (m_1 \otimes \sigma) \circ (\text{id}_{B_1} \otimes \varphi_{2,1} \otimes \text{id}_{B_2})) \circ (\text{id}_{B_2} \otimes \Psi_{B_1 \otimes B_2, B_2} \otimes \text{id}_{B_1 \otimes B_2}) \circ \\
&\quad \circ ((\nu_r \otimes \text{id}_{B_2}) \circ \Delta_2 \otimes (\nu_r \otimes \text{id}_{B_2}) \circ \Delta_2), \\
(\text{id}_{B_2} \otimes m_1) \circ (\Psi_{B_1, B_2} \otimes \text{id}_{B_1}) \circ (\text{id}_{B_1} \otimes \nu_r) \circ \varphi_{2,1} \\
&= (\mu_r \otimes m_1 \circ (\text{id}_{B_1} \otimes \mu_l)) \circ (\text{id}_{B_2} \otimes \Psi_{B_1 \otimes B_2, B_1} \otimes \text{id}_{B_1}) \circ ((\nu_r \otimes \text{id}_{B_2}) \circ \Delta_2 \otimes \Delta_1), \\
\Delta_1 \circ \sigma &= (\sigma \otimes m_1 \circ (m_1 \otimes \sigma) \circ (\text{id}_{B_1} \otimes \varphi_{2,1} \otimes \text{id}_{B_2})) \circ (\text{id}_{B_2} \otimes \Psi_{B_1 \otimes B_2, B_2} \otimes \text{id}_{B_1 \otimes B_2}) \circ \\
&\quad \circ ((\nu_r \otimes \text{id}_{B_2}) \circ \Delta_2 \otimes (\nu_r \otimes \text{id}_{B_2}) \circ \Delta_2).
\end{aligned}$$

The set of defining identities of the idempotents Π_1 and Π_2 and of the projections and injections p_1, p_2, i_1, i_2 can be determined similarly. Eventually the universal properties of the cross product bialgebras of type $\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \nabla & \square \\ \hline \bullet & \bullet \\ \hline \end{array}$ will be described by the following proposition. We will use the notations of Corollary 3.4 and Remark 11.

PROPOSITION 4.6 *Let B be a bialgebra in \mathcal{C} . Then the following equivalent conditions are satisfied.*

1. B is isomorphic to a cross product bialgebra $B_1 \bowtie B_2$ where $\Delta_{21,1} = \eta_2 \otimes \text{id}_{B_1}$ and $\Delta_{11,2} = (\eta_1 \otimes \eta_1) \circ \varepsilon_2$ are trivial.

2. There are idempotents $\Pi_1, \Pi_2 \in \text{End}(B)$ such that
- (a) $m_B \circ (\Pi_1 \otimes \Pi_1) = \Pi_1 \circ m_B \circ (\Pi_1 \otimes \Pi_1)$,
 - (b) Π_1 is coalgebra morphism,
 - (c) $(\Pi_2 \otimes \Pi_2) \circ \Delta_B = (\Pi_2 \otimes \Pi_2) \circ \Delta_B \circ \Pi_2$,
 - (d) $\Pi_1 \circ \eta_B = \eta_B$ and $\varepsilon_B \circ \Pi_2 = \varepsilon_B$,
 - (e) $m_B \circ (\Pi_1 \otimes \Pi_2)$ and $(\Pi_1 \otimes \Pi_2) \circ \Delta_B$ split the idempotent $\Pi_1 \otimes \Pi_2$.
3. There are objects B_1 and B_2 and morphisms $B_1 \xrightarrow{i_1} A \xrightarrow{p_1} B_1$ and $B_2 \xrightarrow{i_2} A \xrightarrow{p_2} B_2$ such that
- (a) i_1 is algebra and coalgebra morphism,
 - (b) p_1 is coalgebra morphism,
 - (c) p_2 is coalgebra morphism,
 - (d) $p_j \circ i_j = \text{id}_{B_j}$ for $j \in \{1, 2\}$.
 - (e) $m_A \circ (i_1 \otimes i_2) : B_1 \otimes B_2 \rightarrow A$ is isomorphism with inverse $(p_1 \otimes p_2) \circ \Delta_A$. \blacksquare

Concluding Remarks

We found a universal theory of strong cross product bialgebras with an equivalent (co-)modular co-cyclic characterization in terms of strong Hopf data. The theory unites all known cross product bialgebras [24, 18, 22, 20, 21, 5] in a single construction. Furthermore various new types of cross product bialgebras arise out of the most general construction. The (co-)modular co-cyclic structure of strong cross product bialgebras corresponds canonically to a strong Hopf datum which in turn completely determines the bialgebra structure.

Thus Hopf data basically provide a pattern for the realization of cross product bialgebras in terms of explicit examples - a task to be done in future investigations.

There is no conceptual explanation yet for the understanding of the strong conditions in Definition 4.3. A canonical origin of these conditions may be found in higher dimensional categorical constructions of cross product bialgebras where the two tensor factors of the strong cross product bialgebras will be considered as object of two different monoidal categories.

Since we are working throughout in braided categories, the results of the article may now be used to investigate cross product bialgebras in various types of braided categories (see [5, 6] for applications in Hopf bimodule categories).

Our directions of study of cross product bialgebras are particularly concerned with these questions as well as with extension theory and cohomological considerations (see [14, 28, 29]).

5 Proof of Theorem 4.4

This section is exclusively devoted to the proof of Theorem 4.4. In what follows we denote by $\mathfrak{h} = ((B_1, m_1, \eta_1, \Delta_1, \varepsilon_1), (B_2, m_2, \eta_2, \Delta_2, \varepsilon_2); \mu_l, \mu_r, \nu_l, \nu_r, \rho, \sigma)$ a strong Hopf datum. Although many of the subsequent results hold for more general Hopf data, we do not

The relative associativity of m_2 and μ_l , and by π -symmetry the relative coassociativity of Δ_2 and ν_r will be shown in the following lemma.

LEMMA 5.2 *For a strong Hopf datum \mathfrak{h} the identities*

$$\begin{array}{c} \begin{array}{ccc} \begin{array}{c} \text{---} \\ \diagup \\ \text{---} \\ \diagdown \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ \diagdown \\ \text{---} \\ \diagup \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ \diagdown \\ \text{---} \\ \diagup \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \\ \begin{array}{c} \text{---} \\ \diagdown \\ \text{---} \\ \diagup \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ \diagup \\ \text{---} \\ \diagdown \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \end{array}$$

and the corresponding π -symmetric versions of the relative coassociativity of Δ_1 and ν_r hold.

PROOF. The relative associativity of m_2 and μ_l will be derived from the respective weak associativity of Definition 4.1 taking into account (4.7) and (5.4) of Lemma 5.1. \blacksquare

REMARK 13 For a Hopf data satisfying $|\mathfrak{r}' = |\mathfrak{b}$ and $|\mathfrak{r} = |\mathfrak{q}$ the multiplication m_2 and the comultiplication Δ_1 are (co-)associative.

LEMMA 5.3 *The relative versions of the morphisms $\varphi_{1,2}$ and $\varphi_{2,1}$ are given by*

$$\begin{array}{c} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \diagdown \\ \text{---} \\ \diagup \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \diagup \\ \text{---} \\ \diagdown \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \quad \text{and} \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \diagup \\ \text{---} \\ \diagdown \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \diagdown \\ \text{---} \\ \diagup \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \end{array} \quad (5.5)$$

PROOF. Using that (B_2, μ_r) is a right module and applying Lemma 5.1 yields the first identity of (5.5). \blacksquare

Then the next lemma follows from the module-algebra and comodule-coalgebra compatibilities of Definition 4.1.

LEMMA 5.4 *The identities*

$$\begin{array}{c} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \end{array} \quad (5.6)$$

$$\begin{array}{c} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \end{array} \quad (5.7)$$

are satisfied for a strong Hopf datum \mathfrak{h} .

LEMMA 5.5 *Let \mathfrak{h} be strong Hopf datum. Then $(B_1, m_1, \eta_1, \nu_l)$ is a left B_2 -comodule algebra and $(B_2, \Delta_2, \varepsilon_2, \mu_r)$ is a right B_1 -module coalgebra.*

PROOF. We use Lemma 5.1 and the second identity of the comodule-algebra compatibility of Definition 4.1 to show that $(B_1, m_1, \eta_1, \nu_l)$ is a left B_2 -comodule algebra. In π -symmetric manner it will be proven that $(B_2, \Delta_2, \varepsilon_2, \mu_r)$ is a right B_1 -module coalgebra. \blacksquare

LEMMA 5.6 For a strong Hopf datum \mathfrak{h} the relativizations of the algebra-coalgebra compatibilities, the (left) module-coalgebra compatibility, and the (right) comodule-algebra compatibility are respectively given by

$$\begin{array}{c} \text{Diagram 1} = \text{Diagram 2} \quad , \quad \text{Diagram 3} = \text{Diagram 4} \end{array} \quad (5.8)$$

$$\begin{array}{c} \text{Diagram 5} = \text{Diagram 6} \quad , \quad \text{Diagram 7} = \text{Diagram 8} \end{array} \quad (5.9)$$

$$\begin{array}{c} \text{Diagram 9} = \text{Diagram 10} \quad , \quad \text{Diagram 11} = \text{Diagram 12} \end{array} \quad (5.10)$$

PROOF. Since $(B_1, m_1, \eta_1, \nu_l)$ is a left comodule algebra by Lemma 5.5 the first identity in (5.12) follows from the relative associativity of m_2 according to Lemma 5.2.

The verification of (5.9) needs a little more calculation. We prove the first identity of (5.9). The second one can be derived with similar techniques. We start with the second module-coalgebra compatibility in Definition 4.1. We apply the relativization (with μ_r) to the second tensor factor on both sides of the graphic. The left hand side of this relative module-coalgebra compatibility yields the left hand side of the first identity in (5.9) if we consecutively apply modularity of (B_2, μ_r) , the second relation of (4.7) and the second relation of (4.6). To obtain the right hand side of the first identity of (5.9) we transform the right hand side of the relative module-coalgebra compatibility using successively the third relation of (4.7), modularity of (B_2, μ_r) , the first equation of (5.5), the modularity of B_2 , the second identity of (4.7), and eventually again modularity of B_2 .

All other identities can be derived similarly, in particular because of π -symmetric reasons. ■

In the subsequent Lemmas 5.7 and 5.8 the entwining properties of the morphisms $\varphi_{1,1}$, $\varphi_{2,2}$, $\varphi_{1,2}$ and $\varphi_{2,1}$ will be investigated.

LEMMA 5.7 The morphism $\varphi_{1,1}$ entwines with the multiplication m_1 , and $\varphi_{2,2}$ entwines with the comultiplication Δ_2 according to

$$\begin{array}{c} \text{Diagram 13} = \text{Diagram 14} \quad , \quad \text{Diagram 15} = \text{Diagram 16} \\ \text{Diagram 17} = \text{Diagram 18} \quad , \quad \text{Diagram 19} = \text{Diagram 20} \\ \text{Diagram 21} = \text{Diagram 22} = \text{Diagram 23} \quad , \quad \text{Diagram 24} = \text{Diagram 25} = \text{Diagram 26} \end{array} \quad (5.11)$$

PROOF. For the proof of the first identity of Lemma 5.7 we use Lemma 5.5 and the fourth equation of Lemma 5.2. To verify the second identity of (5.11), the module-algebra compatibility for Hopf data, Lemma 5.3 and the comodularity of (B_1, ν_l) have to be applied successively. Using the module-algebra compatibility and the π -symmetric version of the fourth identity of Lemma 5.2 yields the fifth identity of (5.11). Simple calculations yield the sixth identity of (5.11). The remaining relations of the lemma follow by π -symmetric reasoning. \blacksquare

LEMMA 5.8 *The (relative) entwining identities for $\varphi_{1,2}$ and $\varphi_{2,1}$ are given by*

$$\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} = \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array}, \quad \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} = \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} \quad (5.12)$$

$$\begin{array}{c} \text{Diagram 9} \\ \text{Diagram 10} \end{array} = \begin{array}{c} \text{Diagram 11} \\ \text{Diagram 12} \end{array}, \quad \begin{array}{c} \text{Diagram 13} \\ \text{Diagram 14} \end{array} = \begin{array}{c} \text{Diagram 15} \\ \text{Diagram 16} \end{array} \quad (5.13)$$

$$\begin{array}{c} \text{Diagram 17} \\ \text{Diagram 18} \end{array} = \begin{array}{c} \text{Diagram 19} \\ \text{Diagram 20} \end{array}, \quad \begin{array}{c} \text{Diagram 21} \\ \text{Diagram 22} \end{array} = \begin{array}{c} \text{Diagram 23} \\ \text{Diagram 24} \end{array} \quad (5.14)$$

$$\begin{array}{c} \text{Diagram 25} \\ \text{Diagram 26} \end{array} = \begin{array}{c} \text{Diagram 27} \\ \text{Diagram 28} \end{array}, \quad \begin{array}{c} \text{Diagram 29} \\ \text{Diagram 30} \end{array} = \begin{array}{c} \text{Diagram 31} \\ \text{Diagram 32} \end{array} \quad (5.15)$$

PROOF. We use Lemma 5.5 for B_1 and the relative associativity of m_2 (Lemma 5.2) to obtain the first identity of (5.12).

The first identity of (5.13) will be used explicitly in the proof of Proposition 5.19. Below we will give its detailed derivation.

$$\varphi_{2,1} \circ (B_2 \otimes m_1) = \begin{array}{c} \text{Diagram 33} \\ \text{Diagram 34} \\ \text{Diagram 35} \end{array} = \begin{array}{c} \text{Diagram 36} \\ \text{Diagram 37} \\ \text{Diagram 38} \end{array} = \begin{array}{c} \text{Diagram 39} \\ \text{Diagram 40} \\ \text{Diagram 41} \end{array} = (m_1 \otimes B_2) \circ \varphi_{2,11}$$

where the first equation comes from the relative bialgebra property of B_1 according to (5.8). The second equation can be verified with the help of the module-algebra compatibility of Definition 4.1, and using that (B_2, μ_r) is a right module. In the third identity we use the relative associativity of Δ_1 proven in Lemma 5.2. Finally the result follows because $(B_2, \Delta_2, \varepsilon_2, \mu_r)$ is a module coalgebra by Lemma 5.5.

The left hand side of the first identity of (5.14) will be transformed to the right hand side of the identity by using successively the algebra-coalgebra compatibility for Δ_2 and m_2 , the comodularity of (B_1, ν_l) , the first relation of (4.6), the first identity of (5.6), the relative associativity of μ_l according to Lemma 5.2 and finally the second relation of (5.5).

The first identity of (5.15) immediately follows from the relative module coalgebra properties (5.9).

All other relations of the lemma can be derived easily by applying π -symmetry. \blacksquare

LEMMA 5.9 For the strong Hopf datum \mathfrak{h} the following identities are satisfied.

$$\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} = \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array}, \quad \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} = \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} \quad (5.16)$$

$$\begin{array}{c} \text{Diagram 9} \\ \text{Diagram 10} \end{array} = \begin{array}{c} \text{Diagram 11} \\ \text{Diagram 12} \end{array}, \quad \begin{array}{c} \text{Diagram 13} \\ \text{Diagram 14} \end{array} = \begin{array}{c} \text{Diagram 15} \\ \text{Diagram 16} \end{array} \quad (5.17)$$

$$\begin{array}{c} \text{Diagram 17} \\ \text{Diagram 18} \end{array} = \begin{array}{c} \text{Diagram 19} \\ \text{Diagram 20} \end{array}, \quad \begin{array}{c} \text{Diagram 21} \\ \text{Diagram 22} \end{array} = \begin{array}{c} \text{Diagram 23} \\ \text{Diagram 24} \end{array} \quad (5.18)$$

PROOF. The proof follows straightforwardly from the cocycle and cycle compatibilities of Definition 4.1 and the identities (4.6) and (4.7). \blacksquare

LEMMA 5.10 The subsequent relations involving $\hat{\rho}$ and $\hat{\sigma}$ hold in \mathfrak{h} .

$$\begin{array}{c} \text{Diagram 25} \\ \text{Diagram 26} \end{array} = \begin{array}{c} \text{Diagram 27} \\ \text{Diagram 28} \end{array}, \quad \begin{array}{c} \text{Diagram 29} \\ \text{Diagram 30} \end{array} = \begin{array}{c} \text{Diagram 31} \\ \text{Diagram 32} \end{array} \quad (5.19)$$

$$\begin{array}{c} \text{Diagram 33} \\ \text{Diagram 34} \end{array} = \begin{array}{c} \text{Diagram 35} \\ \text{Diagram 36} \end{array}, \quad \begin{array}{c} \text{Diagram 37} \\ \text{Diagram 38} \end{array} = \begin{array}{c} \text{Diagram 39} \\ \text{Diagram 40} \end{array} \quad (5.20)$$

PROOF. Because of π -symmetry we will only demonstrate the first identities of (5.19) and (5.20). Applying to the left hand side of (5.19) the algebra-coalgebra compatibility (for Δ_2 and m_2) and the entwining property of $\varphi_{2,2}$ (Lemma 5.7) yields the result.

To obtain the first identity of (5.20) we transform its left hand side consecutively using the relative bialgebra property (5.8) of Δ_2 and m_2 , the comodularity of (B_1, ν_l) , the first relation of (5.6), the first identity of (5.16), the relative associativity of m_2 , and again the fact that (B_1, ν_l) is left comodule. \blacksquare

REMARK 14 Note that (5.4) and (5.16) are special cases of (5.20). Furthermore (5.18) implies (4.7).

Special Properties of Strong Hopf Data

Before we will prove Theorem 4.4 we have to provide several auxiliary definitions and specific properties of strong Hopf Data. Similarly as in Remark 6 we define $p_1 = \text{id}_{B_1} \otimes \varepsilon_2$, $p_2 = \varepsilon_1 \otimes \text{id}_{B_2}$, $p_0 = \text{id}_B$, $i_1 = \text{id}_{B_1} \otimes \eta_2$, $i_2 = \eta_1 \otimes \text{id}_{B_2}$, and $i_0 = \text{id}_B$. In this context we occasionally use the notation $B_0 := B := B_0 \otimes B_1$. Given a strong Hopf datum \mathfrak{h} we can build the morphisms $m_B : B \otimes B \rightarrow B$ and $\Delta_B : B \rightarrow B \otimes B$ like in (3.9) as

$$m_B := \begin{array}{c} \text{Diagram 41} \\ \text{Diagram 42} \end{array} \quad \text{and} \quad \Delta_B := \begin{array}{c} \text{Diagram 43} \\ \text{Diagram 44} \end{array} \quad (5.21)$$

Similarly as in (3.4) and (3.5) we define

$$\begin{aligned} m_{i,j,k} &:= p_i \circ m_B \circ (i_j \otimes i_k) = \begin{array}{c} \text{Diagram 45} \\ \text{Diagram 46} \end{array} \quad \text{for all } i, j, k \in \{0, 1, 2\}, \\ \Delta_{i,j,k} &:= (p_i \otimes p_j) \circ \Delta_B \circ i_k = \begin{array}{c} \text{Diagram 47} \\ \text{Diagram 48} \end{array} \quad \text{for all } i, j, k \in \{0, 1, 2\}. \end{aligned} \quad (5.22)$$

and

$$\begin{aligned} m_{0,20}^* &:= \hat{\sigma} \circ (\mu_r \otimes \text{id}_{B_2}), & m_{l,2m}^* &:= \text{pl} \circ m_{0,20}^* \circ (\text{id}_{B_2} \otimes i_m) \\ \Delta_{01,0}^* &:= (\text{id}_{B_1} \otimes \nu_l) \circ \hat{\rho}, & \Delta_{l1,m}^* &:= (\text{pl} \otimes \text{id}_{B_1}) \circ \Delta_{01,0}^* \circ i_m \end{aligned} \quad (5.23)$$

for $l, m \in \{0, 1, 2\}$. In particular (5.22) implies

$$\begin{aligned} m_1 &= m_{1,11}, & \Delta_1 &= \Delta_{11,1}, & \mu_l &= m_{1,21}, & \nu_l &= \Delta_{21,1}, \\ m_2 &= m_{2,22}, & \Delta_2 &= \Delta_{22,2}, & \mu_r &= m_{2,21}, & \nu_r &= \Delta_{21,2}, \\ m_B &= m_{0,00}, & \Delta_B &= \Delta_{00,0}, & \sigma &= m_{1,22}, & \rho &= \Delta_{11,2}. \end{aligned}$$

The following morphisms $\langle \begin{smallmatrix} kl \\ ijmn \\ rs \end{smallmatrix} \rangle, \langle i_{rs}^{kl} |, |_{rs}^{klj} \rangle, \langle 2_{rs}^{kl} |^*, |_{rs}^{kl1} \rangle^* : B_k \otimes B_l \rightarrow B_r \otimes B_s$ turn out to be useful in the sequel. Let $i, j, m, n, r, s, k, l \in \{0, 1, 2\}$, then

$$\begin{aligned} \langle \begin{smallmatrix} kl \\ ijmn \\ rs \end{smallmatrix} \rangle &:= (m_{r,im} \otimes m_{s,jn}) \circ (\text{id}_{B_i} \otimes \Psi_{B_j, B_m} \otimes \text{id}_{B_n}) \circ (\Delta_{ij,k} \otimes \Delta_{mn,l}), \\ \langle i_{rs}^{kl} | &:= (m_{r,il} \otimes \text{id}_{B_s}) \circ (\text{id}_{B_i} \otimes \Psi_{B_s, B_l}) \circ (\Delta_{is,k} \otimes \text{id}_{B_l}), \\ |_{rs}^{klj} \rangle &:= (\text{id}_{B_r} \otimes m_{s,kj}) \circ (\Psi_{B_k, B_r} \otimes \text{id}_{B_j}) \circ (\text{id}_{B_k} \otimes \Delta_{rj,l}), \\ \langle 2_{rs}^{kl} |^* &:= (m_{r,2l}^* \otimes \text{id}_{B_s}) \circ (\text{id}_{B_2} \otimes \Psi_{B_s, B_l}) \circ (\Delta_{2s,k} \otimes \text{id}_{B_l}), \\ |_{rs}^{kl1} \rangle^* &:= (\text{id}_{B_r} \otimes m_{s,k1}) \circ (\Psi_{B_k, B_r} \otimes \text{id}_{B_1}) \circ (\text{id}_{B_k} \otimes \Delta_{r1,l}^*). \end{aligned} \quad (5.24)$$

For a strong Hopf datum \mathfrak{h} we obtain the following bra-ket decomposition of morphisms (5.24).

LEMMA 5.11

$$\langle \begin{smallmatrix} kl \\ i0mn \\ rs \end{smallmatrix} \rangle = (\text{id}_{B_r} \otimes m_{s,1s}) \circ (\langle i_{r1}^{km} | \otimes \text{id}_{B_s}) \circ (\text{id}_{B_k} \otimes |_{ms}^{2l n}) \circ (\Delta_{k2,k} \otimes \text{id}_{B_l}) \quad (5.25)$$

for any $i, k, l, m, n, r, s \in \{0, 1, 2\}$.

PROOF. The lemma follows straightforwardly from the (co-)associativity of m_B and Δ_B and the identities $m_{0,12} = \text{id}_{B_0} = \Delta_{12,0}$. \blacksquare

Below we define another set of auxiliary morphisms which will be used in the course of the proof of Theorem 4.4.

$$\tau_t^0 := \text{diagram}, \quad \tau_t^1 := \text{diagram}, \quad \tau_t^2 := \text{diagram}, \quad \tau_t^3 := \text{diagram} \quad (5.26)$$

$$\tau_b^0 := \text{diagram}, \quad \tau_b^1 := \text{diagram}, \quad \tau_b^2 := \text{diagram}, \quad \tau_b^3 := \text{diagram} \quad (5.27)$$

Besides we use the following definitions.

$$\begin{aligned} \sigma^{\text{tr}} &:= \text{diagram}, & \rho^{\text{tr}} &:= \text{diagram}, & (\tau_t^3)^{\text{red}_1} &:= \text{diagram}, & (\tau_b^3)^{\text{red}_1} &:= \text{diagram} \\ (\tau_t^3)^{\text{red}_2} &:= \text{diagram}, & (\tau_b^3)^{\text{red}_2} &:= \text{diagram}, & (\tau_t^3)^{\text{red}_3} &:= \text{diagram}, & (\tau_b^3)^{\text{red}_3} &:= \text{diagram} \end{aligned} \quad (5.28)$$

The previous lemma leads to

LEMMA 5.14

$$\begin{array}{c}
 \begin{array}{ccc}
 \begin{array}{c} \text{---} \\ | \langle 22_{01}^1 \rangle | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \boxed{(\tau_t^3) \text{red}_2} \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ | \langle 2_{11}^{20} \rangle | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \boxed{(\tau_b^3) \text{red}_2} \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ \langle \begin{array}{c} 22 \\ 2001 \\ 11 \end{array} \rangle \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \boxed{(\tau_t^3) \text{red}_2} \\ \text{---} \end{array}
 \end{array}
 \end{array} \quad (5.35)$$

$$\begin{array}{c}
 \begin{array}{ccc}
 \begin{array}{c} \text{---} \\ | \langle 2_{21}^{20} \rangle | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \boxed{(\tau_t^3) \text{red}_3} \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ | \langle 2_{10}^{21} \rangle | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \boxed{(\tau_b^3) \text{red}_3} \\ \text{---} \end{array}
 \end{array}
 \end{array} \quad (5.36)$$

$$\begin{array}{c}
 \begin{array}{ccc}
 \begin{array}{c} \text{---} \\ | \langle 2_{01}^{20} \rangle | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \boxed{(\tau_t^3) \text{red}_1} \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ | \langle 2_{01}^{20} \rangle^* | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \boxed{(\tau_b^3) \text{red}_1} \\ \text{---} \end{array}
 \end{array}
 \end{array} \quad (5.37)$$

PROOF. The first identity of (5.35) has been derived by successive application of the left module-algebra compatibility of Definition 4.1 and the relations (5.6) and (5.16). The second identity is its π -symmetric counterpart. To get the third identity we use (5.25), the first and the second equation of (5.35), the relations (5.17) and the module-comodule compatibility.

The first identity in (5.36) is obtained from the left module-algebra compatibility and (5.6). The identities in (5.37) are derived from (5.36) with the help of (5.20). \blacksquare

The next lemma is a straightforward implication of the previous result.

LEMMA 5.15

$$\begin{array}{ccc}
 \begin{array}{c} \text{---} \\ | \langle \begin{array}{c} 22 \\ 2001 \\ 11 \end{array} \rangle | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \langle \begin{array}{c} 22 \\ 2221 \\ 11 \end{array} \rangle | \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ | \langle \begin{array}{c} 22 \\ 2001 \\ 11 \end{array} \rangle | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \langle \begin{array}{c} 22 \\ 2111 \\ 11 \end{array} \rangle | \\ \text{---} \end{array}
 \end{array} \quad (5.38)$$

PROOF. Both identities follow from the third identity of (5.35) using (4.6), (5.16) and (4.7). \blacksquare

LEMMA 5.16 *Given a strong Hopf datum \mathfrak{h} , then it holds*

$$\begin{array}{ccc}
 \begin{array}{c} \text{---} \\ | \hat{\sigma} | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \langle \begin{array}{c} 22 \\ 2020 \\ 11 \end{array} \rangle | \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ | \hat{\rho} | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \langle \begin{array}{c} 22 \\ 0011 \\ 11 \end{array} \rangle | \\ \text{---} \end{array}
 \end{array} \quad (5.39)$$

$$\begin{array}{ccc}
 \begin{array}{c} \text{---} \\ | \hat{\sigma} | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \langle \begin{array}{c} 22 \\ 2020 \\ 10 \end{array} \rangle | \\ \text{---} \end{array} & , & \begin{array}{c} \text{---} \\ | \hat{\rho} | \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ | \langle \begin{array}{c} 22 \\ 0011 \\ 01 \end{array} \rangle | \\ \text{---} \end{array}
 \end{array} \quad (5.40)$$

PROOF. We prove the first identity of (5.39) graphically.

$$\begin{array}{c} \text{hook} \end{array} = \begin{array}{c} \langle \begin{array}{c} 22 \\ 2000 \\ 11 \end{array} \rangle = \begin{array}{c} \sigma^{\text{tr}} \\ \langle \begin{array}{c} 22 \\ 2001 \\ 11 \end{array} \rangle = \begin{array}{c} \sigma^{\text{tr}} \\ \langle \begin{array}{c} 22 \\ 2221 \\ 11 \end{array} \rangle = \begin{array}{c} \langle \begin{array}{c} 22 \\ 2020 \\ 11 \end{array} \rangle \end{array} \quad (5.41)$$

The first equation in the graphic has been obtained with the help of the cycle-cocycle compatibility (Definition 4.1), (4.6) and (4.7). In the second equation we use (5.25) and (5.32). The third equation is (5.38). Application of (5.32) to the fourth diagram yields the fifth relation. The identities (5.40) follow from (5.39) with the help of the entwining properties of $\varphi_{2,2}$ and $\varphi_{1,1}$ respectively. ■

LEMMA 5.17 *Let \mathfrak{h} be a strong Hopf datum. Then*

$$\begin{array}{c} \hat{\sigma} \\ \langle \begin{array}{c} 22 \\ 210 \end{array} \rangle \end{array} = \begin{array}{c} \langle \begin{array}{c} 22 \\ 210 \end{array} \rangle \end{array} \quad (5.42)$$

and the corresponding π -symmetric version for $\hat{\rho}$ holds.

PROOF. The identity follows straightforwardly from (5.34). ■

LEMMA 5.18

$$\begin{array}{c} \langle \begin{array}{c} 221 \\ 001 \end{array} \rangle^* = \begin{array}{c} \langle \begin{array}{c} 22 \\ 011 \end{array} \rangle^* \end{array}, \quad \langle \begin{array}{c} 200 \\ 11 \end{array} \rangle^* = \begin{array}{c} \langle \begin{array}{c} 220 \\ 11 \end{array} \rangle^* \end{array} \quad (5.43)$$

$$\begin{array}{c} \langle \begin{array}{c} 221 \\ 001 \end{array} \rangle = \begin{array}{c} \langle \begin{array}{c} 221 \\ 011 \end{array} \rangle \end{array}, \quad \langle \begin{array}{c} 200 \\ 11 \end{array} \rangle = \begin{array}{c} \langle \begin{array}{c} 220 \\ 11 \end{array} \rangle \end{array} \quad (5.44)$$

PROOF. The identities (5.43) follow from (5.16), (5.6), whereas (5.44) will be derived from (5.35) and the entwining properties (5.11), (5.13) of $\varphi_{2,2}$, $\varphi_{2,1}$. ■


Proof of Theorem 4.4

Given a strong Hopf datum \mathfrak{h} we have to prove that the object $B = B_1 \otimes B_2$ provided with the structure given in Theorem 4.4, is a strong cross product bialgebra. The (co-)unital identities (3.6) can be verified easily. The “strong conditions” of Definition 3.8 hold by construction. Using the first and the second identity of Lemma 5.4 it follows straightforwardly that the “projection relations” (3.7) are fulfilled. It remains to show that B is a bialgebra.

PROPOSITION 5.19 *Let \mathfrak{h} be a strong Hopf datum and $\varphi_{1,2}$, $\varphi_{2,1}$, $\hat{\rho}$, and $\hat{\sigma}$ be the morphisms defined in (4.2). Then $B = B_1 \otimes B_2$ is a cocycle cross product algebra $B_1 \bowtie_{\hat{\sigma}} B_2$ and a cycle cross product coalgebra $B_1 \hat{\bowtie}_{\hat{\rho}} B_2$.*

PROOF. According to Proposition 2.2 the identities (2.3) have to be verified in order to prove that B is a cross product algebra. Similar π -symmetric procedures are needed to demonstrate that B is a cross product coalgebra.

Without difficulties the unital identities of (2.3) can be verified. The fifth relation of (2.3) has been proven in Lemma 5.8 in the first identity of (5.13). The seventh identity of (2.3) will be proven subsequently.



where the first identity has been obtained from (5.19). We use the weak associativity of μ_l and the module-algebra compatibility of μ_r according to Definition 4.1 to get the second equation in the graphic. Then the entwining property of $\varphi_{2,2}$ (see Lemma 5.7) yields the third relation. In the fourth identity we use (5.18) and the relative coassociativity of Δ_1 corresponding to Lemma 5.2. To derive the fifth identity the module-comodule compatibility of Definition 4.1 has been used. In the sixth equation we apply (5.15), and in the seventh equation we use (5.12). Hence the seventh identity of (2.3) has been verified. Finally we will prove the sixth identity of (2.3). Its left hand side will be transformed

according to

(5.45)

where we used (5.12) and (5.19) in the first equation, and (5.40) and the right modularity of (B_2, μ_r) in the second equation. To get the third identity in the graphic one has to apply (5.42) and again the fact that (B_2, μ_r) is right module. In the fourth relation we use (5.12) and the relative comodule property of ν_r according to Lemma 5.2. The fifth identity can be verified with the help of (5.5). The module-comodule compatibility, the associativity of Δ_2 and the entwining property (5.11) of $\varphi_{2,2}$ yield the final equation in the graphic.

For the right hand side of the sixth identity of (2.3) we get

(5.46)

In the first equation of (5.46) we used (5.19). Then we applied (5.11) to get the second equation. To derive the third identity we make use of the cocycle compatibility and the

weak associativity of m_2 according to Definition 4.1. In the fourth equation we apply the right comodule-coalgebra compatibility, and the fifth identity follows from (5.33). The sixth equation in the graphical calculation (5.46) has been obtained from the entwining property of $\varphi_{2,2}$ and the left module-algebra compatibility.

Eventually, the seventh graphic in (5.45) and in (5.46) coincide. This can be verified by applying Lemma 5.2 for m_2 and ν_r , coassociativity of Δ_2 , the right modularity of (B_2, μ_r) and (5.6) of Lemma 5.4. \blacksquare

REMARK 15 To finish the proof of Theorem 4.4 we have to show that Δ_B is an algebra morphism. For this purpose we will use a sort of *dressing transformation* technique. This means we define two sequences of morphisms $(\beta_\ell^i)_{i=0}^4$ and $(\beta_r^i)_{i=0}^4$ and a sequence of “dressing transformations” $(T^{(i)})_{i=1}^4$ where $\beta_r^0 = (m_B \otimes m_B) \circ (\text{id} \otimes \Psi_{B,B} \otimes \text{id}) \circ (m_B \otimes m_B)$, $\beta_\ell^0 = \Delta_B \circ m_B$, $\beta_\ell^4 = \beta_r^4$, and $T^{(i)}(\beta_\ell^i) = \beta_\ell^{i-1}$, $T^{(i)}(\beta_r^i) = \beta_r^{i-1}$ for all $i \in \{1, 2, 3, 4\}$. This implies $\beta_\ell^0 = \beta_r^0$ and therefore the statement.

REMARK 16 Once it is proven that B is an algebra, Proposition 2.6 provides an alternative method to show that Δ_B is an algebra morphism. The present proof using dressing transformations is invariant under π -symmetry, however.

The “dressing transformations” will be defined as

$$T^{(1)}(f) := \begin{array}{c} \tau_\ell^1 \\ \boxed{f} \\ \tau_b^1 \end{array}, \quad T^{(2)}(f) := \begin{array}{c} \tau_\ell^2 \\ \boxed{f} \\ \tau_b^2 \end{array}, \quad T^{(3)}(g) := \begin{array}{c} \tau_\ell^3 \\ \boxed{g} \\ \tau_b^3 \end{array}, \quad T^{(4)}(h) := \begin{array}{c} \tau_\ell^4 \\ \boxed{h} \\ \tau_b^4 \end{array} \quad (5.47)$$

for any $f \in \text{Hom}(B_{212}, B_{121})$, $g \in \text{Hom}(B_{2122}, B_{1121})$ and $h \in \text{Hom}(B_{22122}, B_{11211})$.

The remainder of this section is devoted to the proof of the following proposition.

PROPOSITION 5.20 *Given a strong Hopf datum \mathfrak{h} , then there exist two sequence of morphisms $(\beta_\ell^i)_{i=0}^4$, $(\beta_r^i)_{i=0}^4$ with $\beta_\ell^0 := \Delta_B \circ m_B$ and $\beta_r^0 := (m_B \otimes m_B) \circ (\text{id}_B \otimes \Psi_{B,B} \otimes \text{id}_B) \circ (\Delta_B \otimes \Delta_B)$ such that*

$$\begin{array}{ccccccccc} \beta_\ell^0 & \xleftarrow{T^{(1)}} & \beta_\ell^1 & \xleftarrow{T^{(2)}} & \beta_\ell^2 & \xleftarrow{T^{(3)}} & \beta_\ell^3 & \xleftarrow{T^{(4)}} & \beta_\ell^4 \\ & & & & & & & & \parallel \\ \beta_r^0 & \xleftarrow{T^{(1)}} & \beta_r^1 & \xleftarrow{T^{(2)}} & \beta_r^2 & \xleftarrow{T^{(3)}} & \beta_r^3 & \xleftarrow{T^{(4)}} & \beta_r^4 \end{array} \quad (5.48)$$

Therefore $\beta_\ell^0 = \beta_r^0$ and $B = B_1 \otimes B_2$ is a bialgebra which proves Theorem 4.4.

PROOF. The proof will be splitted into several parts. At first we verify the following diagram.

$$\begin{array}{cccc}
 \beta_r^0 & \beta_r^1 & \beta_r^2 & \beta_r^3 \\
 \parallel & \parallel & \parallel & \parallel \\
 \langle 00 \rangle & \langle 20 \rangle & \langle 20 \rangle & \\
 \langle 0000 \rangle & \langle 0000 \rangle & \langle 2001 \rangle & \\
 \langle 00 \rangle & & \langle 01 \rangle & \\
 \parallel & \parallel & \parallel & \parallel \\
 \begin{array}{c} \boxed{\langle 20_0 \rangle} \\ \parallel \\ \langle 00_01 \rangle \end{array} & \xleftarrow{T(1)} & \begin{array}{c} \boxed{\langle 20_0 \rangle} \\ \parallel \\ \langle 02_01 \rangle \end{array} & \xleftarrow{T(2)} & \begin{array}{c} \boxed{\langle 20_1 \rangle} \\ \parallel \\ \langle 22_01 \rangle \end{array} & \xleftarrow{T(3)} & \text{Diagram}
 \end{array} \tag{5.49}$$

where the identities (\equiv) in the first row are definitions and the equalities ($=$) in the second row are special cases of (5.25). The morphisms in the third row will be obtained by applying $T^{(1)}$, $T^{(2)}$ and $T^{(3)}$ using the identities (5.30), (5.31) and (5.37) respectively.

A similar diagram can be set up for the (β_ℓ^i) . For that we use the following auxiliary definitions

$$\gamma_t^0 := (\text{id}_{B_1} \otimes \Delta_1) \circ m_{0,20} \quad , \quad \gamma_b^0 := \Delta_{01,0} \circ (m_1 \otimes \text{id}_{B_2}) \tag{5.50}$$

$$\begin{array}{ccc}
 \gamma_t^1 := \text{Diagram} & , & \gamma_t^2 := \text{Diagram} & , & \gamma_t^3 := \text{Diagram} \\
 \gamma_b^1 := \text{Diagram} & , & \gamma_b^2 := \text{Diagram} & , & \gamma_b^3 := \text{Diagram}
 \end{array} \tag{5.51}$$

Then

$$\begin{array}{cccc}
 \beta_\ell^0 & \beta_\ell^1 & \beta_\ell^2 & \beta_\ell^3 \\
 \parallel & \parallel & \parallel & \parallel \\
 \Delta_0 \circ m_0 & \Delta_{01,0} \circ m_{0,20} & & \\
 \parallel & \parallel & & \\
 \begin{array}{c} \boxed{\gamma_t^0} \\ \parallel \\ \gamma_b^0 \end{array} & \xleftarrow{T(1)} & \begin{array}{c} \boxed{\gamma_t^1} \\ \parallel \\ \gamma_b^1 \end{array} & \xleftarrow{T(2)} & \begin{array}{c} \boxed{\gamma_t^2} \\ \parallel \\ \gamma_b^2 \end{array} & \xleftarrow{T(3)} & \begin{array}{c} \boxed{\gamma_t^3} \\ \parallel \\ \gamma_b^3 \end{array}
 \end{array} \tag{5.52}$$

Again the identities (\equiv) in the first row are definitions. The remaining identities will be proven in the subsequent lemmas.

LEMMA 5.21

$$\gamma_t^0 = (m_{0,20} \otimes \text{id}_{B_2}) \circ \tau_t^1, \quad \gamma_b^0 = \tau_b^1 \circ (\text{id}_{B_1} \otimes \Delta_{01,0}) \quad (5.53)$$

and hence $\beta_\ell^0 = T^{(1)}(\beta_\ell^1)$.

PROOF. The identities are obtained using (5.19) and (5.12). \blacksquare


LEMMA 5.22

$$\beta_\ell^1 = (\text{id} \otimes m_1) \circ (\gamma_b^1 \otimes \text{id}_{B_1}) \circ (\text{id}_{B_2} \otimes \gamma_t^1) \circ (\Delta_2 \otimes \text{id}) \quad (5.54)$$

$$\gamma_t^1 = (\text{id} \otimes m_1) \circ (\gamma_t^2 \otimes \text{id}_{B_1}) \circ \tau_t^2, \quad \gamma_b^1 = \tau_b^2 \circ (\text{id}_{B_2} \otimes \gamma_b^2) \circ (\Delta_2 \otimes \text{id}) \quad (5.55)$$


and therefore $\beta_\ell^1 = T^{(2)}(\beta_\ell^2)$.

PROOF. The proof of the identity (5.54) will be given in the following graphical calculation.



where the first equation holds by definition and the second identity is an application of (5.19). In the third equation we use the cycle-cocycle compatibility, (5.25), the (co-)associativity of m_1 and Δ_2 , and (5.12).

To prove the first identity of (5.55) we transform in the next calculation the morphism γ_t^1 with the help of (5.32), the entwining properties (5.11) of $\varphi_{2,2}$, and (5.12) of $\varphi_{1,2}$.



From the right hand side of this equation we get the right hand side of (5.55) by applying (5.34) and again the entwining property (5.11) of $\varphi_{2,2}$. \blacksquare

LEMMA 5.23

$$\gamma_t^2 = \begin{array}{c} \tau_t^3 \\ \gamma_t^3 \end{array}, \quad \gamma_b^2 = \begin{array}{c} \gamma_b^3 \\ \tau_t^3 \end{array} \quad (5.56)$$

and therefore $\beta_\ell^2 = T^{(3)}(\beta_\ell^3)$.

PROOF. The following identities hold.

$$\begin{array}{c}
 \begin{array}{|c|} \hline (22) \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (22|01) \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (22|21) \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (21|2001) \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (21|21) \\ \hline \end{array}
 \end{array} = \begin{array}{c}
 \begin{array}{|c|} \hline (02|2) \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (21|2001) \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (21|21) \\ \hline \end{array}
 \end{array}, \quad \gamma_t^2 = \begin{array}{c}
 \begin{array}{|c|} \hline (\tau_t^3)^{\text{red}_1} \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (02|2) \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (21|2001) \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (21|21) \\ \hline \end{array}
 \end{array} \quad (5.57)$$

The first identity of (5.57) is a consequence of (5.44) and the module-comodule compatibility. Using this result we obtain the second relation in (5.57) with the help of (5.25) and the entwining property (5.11) of $\varphi_{2,2}$.

We denote

$$\tau_t^4 := \begin{array}{|c|} \hline (22) \\ \hline \end{array} \quad (5.58)$$

Then we obtain

$$\begin{array}{c}
 \begin{array}{|c|} \hline (02|2) \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (21|2001) \\ \hline \end{array} \\
 \begin{array}{|c|} \hline (21|21) \\ \hline \end{array}
 \end{array} = \begin{array}{c}
 \begin{array}{|c|} \hline \tau_t^4 \\ \hline \end{array} \\
 \begin{array}{|c|} \hline \gamma_t^3 \\ \hline \end{array}
 \end{array}, \quad \tau_t^3 = \begin{array}{c}
 \begin{array}{|c|} \hline (\tau_t^3)^{\text{red}_1} \\ \hline \end{array} \\
 \begin{array}{|c|} \hline \tau_t^4 \\ \hline \end{array}
 \end{array} \quad (5.59)$$

The first equation in (5.59) results from the algebra-coalgebra compatibility, the right comodule-algebra compatibility, the left module-algebra compatibility, (5.6), (5.16) and the right module algebra property of B_2 . The second identity can be obtained by subsequent application of the right comodule-coalgebra compatibility, the left module-algebra compatibility, (5.6), the right module-coalgebra property of B_2 , and the entwining property of $\varphi_{2,2}$.

The first relation of (5.56) can now be derived by recursive substitution of the identities (5.59) into the second identity of (5.57). \blacksquare

Eventually we define

$$\beta_\ell^4 := \text{[Diagram 1]} \quad \text{and} \quad \beta_r^4 := \text{[Diagram 2]} \quad (5.60)$$

In what follows we will tacitly use the various forms of (weak) (co-)associativity of m_1 , m_2 , Δ_1 , Δ_2 , μ_l , μ_r , ν_l and ν_r which are given by Lemma 5.2 in particular. The subsequent graphical calculation yields $\beta_\ell^3 = T^{(4)}(\beta_\ell^4)$.

$$\beta_\ell^3 = \text{[Diagram 3]} = \text{[Diagram 4]} = \text{[Diagram 5]} = T^{(4)}(\beta_\ell^4)$$

where we used the module-comodule compatibility and (5.6) in the first equation. In the second identity we use again (5.6). To get the third equation in the graphic we applied the right module-algebra and the left comodule-coalgebra compatibilities of Definition 4.1 as well as (5.17). With the help of the module-comodule compatibility the fourth identity can be derived from the definition of β_ℓ^4 in (5.60) and the definition of $T^{(4)}$ in (5.47).

Next we will prove the identity $\beta_r^3 = T^{(4)}(\beta_r^4)$. And therefore all relations in both rows of the diagram in Proposition 5.20 are satisfied.

$$\beta_r^3 = \quad = \quad = \quad = T^{(4)}(\beta_r^4)$$

The first identity in this graphical calculation has been obtained from (5.9) and (5.10) of Lemma 5.6. In the second equation we use (5.7), Lemma 5.5 and (5.8). With the help of the left module-algebra and right comodule-coalgebra compatibilities of Definition 4.1 and the relativizations (5.9) and (5.10) of Lemma 5.6 we derive the third identity. The fourth equation holds by definition.

In order to complete the proof of the proposition the equation $\beta_\ell^4 = \beta_r^4$ needs to be shown. This will be done in the following calculation.

$$\beta_\ell^4 = \quad = \quad = \quad = \beta_r^4$$

where the first equation has been derived with the help of (4.6). In the second and the third equation we use (5.6) and (5.7). Finally we apply (4.8) to obtain the fourth identity.

Hence Proposition 5.20 and therefore Theorem 4.4 have been proven. \blacksquare

References

- [1] N. Andruskiewitsch and J. Devoto, *Extensions of Hopf Algebras*, St. Petersburg Math. J., **7**, No. 1, 17-52 (1996).

- [2] Yu. N. Bespalov, *Crossed modules and quantum groups in braided categories*, Applied Categorical Structures **5**, no. 2, 155-204 (1997).
 ———, *Crossed modules, quantum braided groups and ribbon structures*, Teoreticheskaya i Matematicheskaya Fizika **103**, no. 3, 368-387 (1995).
- [3] Yu. N. Bespalov and B. Drabant, *Hopf (Bi-)Modules and Crossed Modules in Braided Categories*, J. Pure Appl. Algebra **123**, 105-129 (1998).
- [4] Yu. N. Bespalov and B. Drabant, *Differential Calculus in Braided Tensor Categories*, preprint q-alg/9703036 (1997).
 ———, *Bicovariant Differential Calculi and Cross Products on Braided Hopf Algebras*, in Proceedings “Quantum Groups and Quantum Spaces, 1995”, Banach Center Publications **40**, Institute of Math., Polish Acad. Sci., eds. R. Budzynski et al (Warsaw 1997).
- [5] Yu. N. Bespalov and B. Drabant, *Cross Product Bialgebras – Part I*, J. Algebra **219**, 466-505 (1999).
- [6] Yu. N. Bespalov and B. Drabant, *Cross Product Bialgebras – Part III*, Preprint DAMTP-98-134 (1998).
- [7] R. J. Blattner, M. Cohen and S. Montgomery, *Crossed products and inner actions of Hopf algebras*, Trans. AMS **298**, 671-711 (1986).
- [8] T. Brzezinski, *Crossed products by a coalgebra*, Commun. Algebra **25** (11), 3551-3575 (1997).
- [9] S. Caenepeel, B. Ion, G. Militaru and S. Zhu, *Factorisation Structures of Algebras and Coalgebras*, Preprint (1998). To appear in Algebras and Representation Theory.
- [10] A. Connes and H. Moscovici, *Hopf Algebras, Cyclic Cohomology and the Transverse Index Theorem*, Commun. Math. Phys. **198**, 199-246 (1998).
- [11] H.-X. Chen and S. Zhang, *The Double Biproduct in Braided Tensor Categories*, Preprint (1998).
- [12] Y. Doi and M. Takeuchi, *Cleft comodule algebras for a bialgebra*, Commun. Alg. **14**, 801-818 (1986).
- [13] P. Freyd and D. Yetter, *Braided compact closed categories with applications to low dimensional topology*, Adv. Math. **77**, 156-182 (1989).
 ———, *Coherence theorems via knot theory*, J. Pure Appl. Algebra **78**, 49-76 (1992).
- [14] I. Hofstetter, *Erweiterungen von Hopf-Algebren und ihre kohomologische Beschreibung*, Ph.D. thesis, Universität München (1990).
 ———, *Extensions of Hopf Algebras and Their Cohomological Description*, J. Algebra **164**, 264-298 (1994).
- [15] A. Joyal and R. Street, *Braided tensor categories*, Adv. Math. **102**, 20-78 (1993).
 ———, *Braided tensor categories*, Macquarie Math. Reports 860081 (1986).

- [16] C. Kassel, *Quantum Groups*, volume **155** of GTM, Springer (1995).
- [17] V. V. Lyubashenko, *Tangles and Hopf algebras in braided categories* J. Pure Appl. Algebra, **98**, 245 (1995).
 ———, *Modular transformations for tensor categories* J. Pure Appl. Algebra, **98**, 279 (1995).
- [18] S. Majid, *Physics for algebraists: Non-commutative and non-cocommutative Hopf algebras by a bicrossproduct construction*, J. Algebra **130**, no. 1, 17-64 (1990).
- [19] S. Majid, *Braided Groups*, J. Pure Appl. Algebra **86**, 187 (1993).
- [20] S. Majid, *More examples of bicrossproduct and double cross product Hopf algebras*, Isr. J. Math. **72**, 133-148 (1990).
- [21] S. Majid, *Double-bosonisation of braided groups and the construction of $u_q(g)$* , to appear in Math. Proc. Camb. Phil. Soc., preprint RIMS-1047, q-alg/9511001 (1995).
- [22] S. Majid and Ya. S. Soibelman, *Bicrossproduct structure of the quantum Weyl group*, J. Algebra **163**, 68-87 (1994).
- [23] S. Montgomery, *Hopf Algebras and Their Actions on Rings*, CBMS **82**, AMS (1992).
- [24] D. E. Radford, *The Structure of Hopf Algebras with a Projection*, J. Algebra **92**, 322 (1985).
- [25] N. Yu. Reshetikhin and V. G. Turaev, *Ribbon Graphs and Their Invariants Derived from Quantum Groups*, Commun. Math. Phys. **127**, 1 (1990).
- [26] P. Schauenburg. *The Structure of Hopf Algebras with a Weak Projection*, Preprint (1998).
- [27] H.-J. Schneider, *Hopf Galois Extensions, Crossed Products, and Clifford Theory*, Advances in Hopf Algebras. Lecture Notes in Pure Appl. Math. **158**, 267-297, (J. Bergen and S. Montgomery, ed.), Dekker (1994).
- [28] W. M. Singer, *Extension Theory for Connected Hopf Algebras*, J. Algebra **21**, 1-16 (1972).
- [29] M. E. Sweedler, *Cohomology of algebras over Hopf algebras*, Trans. Amer. Math. Soc. **133**, 205-239 (1968).
- [30] M. Takeuchi, *Matched pairs of groups and bismash products of Hopf algebras*, Comm. Alg. **9**, 841-882 (1981).
- [31] V. G. Turaev, *Quantum Invariants of Knots and 3-Manifolds*, volume **18** of Studies in Mathematics, Walter de Gruyter (Berlin, New York 1994).