### MANIN TRIPLES OF REAL SIMPLE LIE ALGEBRAS. PART 2

## A.PANOV

Samara State University ul.Akad.Panlova 1, Samara, 443011,Russia panov@info.ssu.samara.ru

**Abstract.** We complete the study of Manin triples of real simple Lie algebras. In the Part 2 of the article we classify the Manin triples  $(\mathbf{g}(\mathbf{R}), W, \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R}))$  (case 2 of the doubles) up to weak and gauge equivalence.

## 0.Introduction

First we recall the main definitions of the Part 1.

**Definition 1.** Let  $\mathbf{g}_1, \mathbf{g}_2, \mathbf{d}$  be Lie algebras over a field K and let Q be a symmetric nondegenerate bilinear form on  $\mathbf{d}$ . A triple  $(\mathbf{g}_1, \mathbf{g}_2, \mathbf{d})$  is called a Manin triple if Q(x, y) is ad-invariant and  $\mathbf{d}$  is a direct sum of maximum isotropic subspaces  $\mathbf{g}_1, \mathbf{g}_2$ .

**Definition 2.** We say that two Manin triples  $(\mathbf{g}, W, \mathbf{d})$  and  $(\mathbf{g}, W', \mathbf{d})$  are weak equivalent if there exists an element a in the adjoint group D of the double  $\mathbf{d}$  such that  $W' = Ad_a(W)$ .

**Definition 3.** We say that two Manin triples  $(\mathbf{g}, W, \mathbf{d})$  and  $(\mathbf{g}, W', \mathbf{d})$  are gauge equivalent if there exists an element a in the adjoint group D of double  $\mathbf{d}$  such that  $W' = Ad_a(W)$  and  $Ad_a(\mathbf{g}) = \mathbf{g}$ .

We study Manin triples as follows. First for given  $\mathbf{g}$  we find all the doubles  $\mathbf{d}$  and describe all forms  $Q(\cdot, \cdot)$  on  $\mathbf{d}$  such that  $\mathbf{g}$  is an isotropic subspace with respect to this form. Then fixing  $\mathbf{d}$  and Q we study complementary subalgebras W such that  $(\mathbf{g}, W, \mathbf{d})$  forms a Manin triple.

We are going the classify Manin triples of real simple Lie algebras up to weak and gauge equivalence. It is known that the double  $\mathbf{d}$  of complex simple Lie algebra  $\mathbf{g}$  is isomorphic to one of following Lie algebras  $\mathbf{g} \otimes A_i(\mathbf{C})$ , where  $A_1(\mathbf{C}) = \mathbf{C}[t]/t^2$ ,  $A_2(\mathbf{C}) = \mathbf{C}[t]/(t^2 - 1/4)$ . In Part 1 of this paper we study the action of conjugation  $\sigma$  of  $\mathbf{C}/\mathbf{R}$  on complex doubles and obtain that the double  $\mathbf{d}(\mathbf{R})$  of real simple Lie algebra  $\mathbf{g}(\mathbf{R})$  is isomorphic to one of following algebras  $\mathbf{g} \otimes A_i(\mathbf{R})$ , where  $A_1(\mathbf{R}) = \mathbf{R}[t]/t^2$ ,  $A_2(\mathbf{R}) = \mathbf{R}[t]/(t^2 - 1/4)$ ,  $A_3(\mathbf{R}) = \mathbf{C}$ .

So, according to the double, there exist 3 types of Manin triples. In Part 1 we consider Manin triples  $(\mathbf{g}(\mathbf{R}), W, \mathbf{g})$  (i.e. Case 3). In this paper we study Manin triples  $(\mathbf{g}, W, \mathbf{g} \oplus \mathbf{g})$  (i.e. Case 2.)

## On Manin triples $(g, W, g \oplus g)$

Let **g** be a semisimple Lie algebra over the field **C** of complex numbers. Let G be the adjoint group of **g**. Denote by  $\Delta$  (resp.  $\Delta^+$ ) the system of roots (resp.of positive

roots) and by  $\Pi = \{\alpha_1, \dots, \alpha_n\}$  the sistem of simple roots. Let  $K(\cdot, \cdot)$  be a Killing form. Consider the Weyl basis:

$$H_k = H_{\alpha_k}, \alpha_k \in \Pi; E_{\alpha}, E_{-a}, \alpha \in \Delta^+$$
$$[H, E_{\alpha}] = \alpha(H)E_{\alpha}, \quad [E_{\alpha}, E_{-\alpha}] = H_{\alpha},$$
$$\alpha(H) = K(H_{\alpha}, H), \quad K(E_{\alpha}, E_{-\alpha}) = 1$$

As usial **h** is a Cartan subalgebra and  $\mathbf{n}^{\pm}$  is upper and lower nilpotent subalgebras.

Let  $(\mathbf{g}, W, \mathbf{d})$  be a Manin triple with  $\mathbf{d} = \mathbf{g} \oplus \mathbf{g}$ . We recall the classification of Manin triples in this case. Denote by j the image of t in  $A_2(\mathbf{C})$ . and  $e = \frac{1}{2} + j$ ,  $f = \frac{1}{2} - j$ . The elements e, f are ortogonal idempotents, i.e. ef = fe = 0 and  $e^2 = f^2 = 1$ . The double  $\mathbf{d} = \mathbf{g}e + \mathbf{g}f$ . The simple calculations yield that

$$Q(ae + bf, a'e + b'f) = K(a, a') - K(b, b')$$

where  $K(\cdot, \cdot)$  is the Killing form of **g**.

Let  $\Pi_1$ ,  $\Pi_2$  be the subsets of  $\Pi$  and  $\phi:\Pi_1\mapsto\Pi_2$  is the map such that:

- i)  $\phi$  is a bijection,
- ii)  $(\phi(\alpha), \phi(\beta)) = (\alpha, \beta),$
- iii) for every  $\alpha \in \Pi_1$  there exists k with the property  $\phi(\alpha), \ldots, \phi^{k-1}(\alpha) \in \Pi_1$  and  $\phi^k(\alpha) \notin \Pi_1$ .

Notations:

 $\mathbf{g}_i$  the Lie semisimple subalgebra of  $\mathbf{g}$  with the system of simple roots  $\Pi_i$ , system of positive roots  $\Delta_i^+$  and Cartan subalgebra  $\mathbf{h}_i$ ;  $G_i$  is the subgroup with Lie algebra  $\mathbf{g}_i$ .  $\phi: \mathbf{g}_1 \mapsto \mathbf{g}_2$  isomorphism of semisimple Lie subalgebras defined via  $\phi: \Pi_1 \mapsto \Pi_2$ ,  $\mathbf{p}_i^{\pm}$  is upper and lower parabolic subalgebras,  $\mathbf{n}_i^{\pm}$  is the nilpotent radical of  $\mathbf{p}_i^{\pm}$ ,  $\mathbf{r}_i = \mathbf{g}_i + \mathbf{h}$  is the reductive part of  $\mathbf{p}_i^{\pm}$ ,  $R_i$  is Lie subgroup with Lie algebra  $\mathbf{r}_i$  and  $R' = R_1 \cap R_2$ .

Let  $\overline{\mathbf{h}}_1$ ,  $\overline{\mathbf{h}}_2$  be the subalgebras of  $\mathbf{h}$  such that  $\overline{\mathbf{h}}_i^{\perp} \subset \overline{\mathbf{h}}_i$  and  $\mathbf{h}_i \subset \overline{\mathbf{h}}_i$ .

Denote  $\mathbf{l}_i = \overline{\mathbf{h}}_i^{\perp} = KerK(\cdot, \cdot)|_{\overline{\mathbf{h}}_i}, \overline{\mathbf{r}}_i = \mathbf{g}_i + \overline{\mathbf{h}}_i$ 

Suppose that  $\phi$  is extended to the isomorphism

$$\phi: \overline{\mathbf{r}}_1/\mathbf{l}_1 \mapsto \overline{\mathbf{r}}_2/\mathbf{l}_2 \tag{1.1}$$

**Definition 1.1**. The following condition we shall call the condition iv):

iv)  $\phi$  preserve  $K(\cdot,\cdot)$  and has no fixed points in **h**.

(the last means that there is no  $x \in \overline{\mathbf{r}}_1 \cap \overline{\mathbf{r}}_2$ ,  $x \neq 0$  such that  $\phi(x) = x(mod(\mathbf{l}_2))$ . Consider

$$\mathbf{W}(\phi) = \mathbf{l}_1 e + \mathbf{n}_1^+ e + \sum_{x \in \overline{\mathbf{r}}_1} \mathbf{C}(xe + \phi(x)f) + \mathbf{n}_2^- f + \mathbf{l}_2 f$$
(1.2)

**Theorem 1.2**( [BD],[ST],[CGR],[S]). Let **g** be a semisimple Lie algebra. Let  $\phi$  satisfy i), ii), iii), iv). Then

- 1)  $(\mathbf{g}, \mathbf{W}(\phi), \mathbf{g} \oplus \mathbf{g}))$  is a Manin triple;
- 2) Every Manin triple  $(\mathbf{g}, W', \mathbf{g} \oplus \mathbf{g})$  is gauge equivalent to  $(\mathbf{g}, \mathbf{W}(\phi), \mathbf{g} \oplus \mathbf{g})$ .

On Manin triples 
$$(g(R), W, g(R) \oplus g(R))$$

Let  $\mathbf{g}(\mathbf{R})$  be a real simple Lie algebra. It is well known that  $\mathbf{g}(\mathbf{R})$  is a real form of a complex simple Lie algebra  $\mathbf{g}$  or coinsides with the realification of  $\mathbf{g}$  (see [VO],[GG]). It was proved in [CGR] and [CH] that Manin triples  $(\mathbf{g}(\mathbf{R}), W, \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R}))$  correspond to the solutions  $\tilde{Q} : \mathbf{g}(\mathbf{R}) \mapsto \mathbf{g}(\mathbf{R})$  of Modified Yang-Baxter equation

$$[\tilde{Q}x, \tilde{Q}y] - \tilde{Q}[\tilde{Q}x, y] - \tilde{Q}[x, \tilde{Q}y] = \lambda[x, y]$$

with  $\lambda < 0$ . A solution exists whenever the simple real Lie algebra  $\mathbf{g}(\mathbf{R})$  has no black roots in Satake-diagram [CGR]. It takes place for algebras:

- L1)  $g(\mathbf{R})$  is a split real form complex simple Lie algebra;
- L2)  $\mathbf{g}(\mathbf{R}) = \mathbf{g}_{\mathbf{R}}$  is the realification of  $\mathbf{g}$ ;
- L3)  $g(\mathbf{R}) = su(p, p), su(p, p + 1);$
- L4)  $g(\mathbf{R}) = so(p, p + 2);$
- L5)  $\mathbf{g}(\mathbf{R}) = EII$ ;

Let  $\mathbf{g}(\mathbf{R})$  be a real simple Lie algebra of L1-L5.

Denote by  $\Gamma = \{1, \sigma : \sigma^2 = 1\}$  the Galois group  $\mathbf{C}/\mathbf{R}$ . There exists a semilinear action of  $\Gamma$  on  $\mathbf{g}(\mathbf{C}) = \mathbf{g} \otimes \mathbf{C}$  such that  $\mathbf{g}(\mathbf{R}) = \mathbf{g}(\mathbf{C})^{\Gamma}$ . There exist the subalgebras  $\mathbf{h}$ ,  $\mathbf{n}^{\pm}$  in  $\mathbf{g}\mathbf{C}$  such that  ${}^{\sigma}\mathbf{n}^{+} = \mathbf{n}^{+}$ ,  ${}^{\sigma}\mathbf{n}^{-} = \mathbf{n}^{-}$ ,  ${}^{\sigma}\mathbf{h} = \mathbf{h}$  (see [CGR]). We are fixing further this subalgebras.

We have the action of  $\sigma$  on the roots  ${}^{\sigma}\alpha(h) = \overline{\alpha({}^{\sigma}h)}$ ,  $h \in \mathbf{h}$ ,  $\alpha \in \Delta$ . In all cases L1-L5  $\sigma(\Pi) = \Pi$ . For the action of  $\sigma$  on  $\Pi$  see Satake-diagrams [VO].

Let  $(\mathbf{g}(\mathbf{R}), W, \mathbf{d}(\mathbf{R}))$  be a Manin triple with  $\mathbf{d}(\mathbf{R}) = \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R})$  (i.e. Case 2). Then  $(\mathbf{g}(\mathbf{C}), W \otimes \mathbf{C}, \mathbf{d}(\mathbf{R}) \otimes \mathbf{C})$  is Manin triple over  $\mathbf{C}$ . Clearly,  $\mathbf{d}(\mathbf{C}) = \mathbf{d}(\mathbf{R}) \otimes \mathbf{C} = \mathbf{g}(\mathbf{C}) \oplus \mathbf{g}(\mathbf{C})$ . The action of  $\sigma$  on  $\mathbf{d}(\mathbf{C})$  is as follows  ${}^{\sigma}e = e$  and  ${}^{\sigma}f = f$ .

Recall that  $\mathbf{g}(\mathbf{C}) = \mathbf{g}$  in the case  $\mathbf{g}(\mathbf{R})$  is a real form of a complex simple Lie algebra  $\mathbf{g}$  and  $\mathbf{g}(\mathbf{C}) = \mathbf{g} + \mathbf{g}$  in the case  $\mathbf{g}(\mathbf{R}) = \mathbf{g}_{\mathbf{R}}$  is the realification of  $\mathbf{g}$ . To simplify the notations, we shall refer  $\mathbf{g}(\mathbf{C})$  as  $\mathbf{g}$  till the end of this section.

According to Theorem 1.2,  $W \otimes \mathbf{C}$  coinsides with some  $\sigma$ -invariant subalgebra  $Ad_g\mathbf{W}(\phi)$ , where  $g \in G$ . In this section we are going to study  $\sigma$ -invariant subalgebras in the orbit  $Ad_g\mathbf{W}(\phi)$  of the adjoint group G. The list of real forms of this subalgebras  $W = (Ad_g\mathbf{W}(\phi))^{\Gamma}$  provides the classification of Manin triples  $(\mathbf{g}(\mathbf{R}), W, \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R}))$ . Consider

$$^{\sigma}(Ad_g\mathbf{W}(\phi)) = Ad_g\mathbf{W}(\phi) \tag{2.1}$$

It follows

$$^{\sigma}(\mathbf{W}(\phi)) = Ad_{p}\mathbf{W}(\phi), \tag{2.2}$$

where  $p = {}^{\sigma}g^{-1}g$ . Note that  ${}^{\sigma}p = p^{-1}$ . Replacing  $\mathbf{W}(\phi)$  by (1.2) in (2.1), we get the equality of two subalgebras

$${}^{\sigma}\mathbf{l}_{1}e + {}^{\sigma}\mathbf{n}_{1}^{+}e + \sum_{x \in \overline{\mathbf{r}}_{1}} \mathbf{C}({}^{\sigma}xe + {}^{\sigma}(\phi(x))f) + {}^{\sigma}\mathbf{n}_{2}^{-}f + {}^{\sigma}\mathbf{l}_{2}f = Ad_{p}(\mathbf{l}_{1})e + Ad_{p}(\mathbf{n}_{1}^{+})e + \sum_{x \in \overline{\mathbf{r}}_{1}} \mathbf{C}(Ad_{p}(x)e + Ad_{p}(\phi(x))f) + Ad_{p}(\mathbf{n}_{2}^{-})f + Ad_{p}(\mathbf{l}_{2})f$$

$$(2.3)$$

Denote  $\mathbf{m}_1 = \mathbf{g}_1 \cap \mathbf{n}_2^-, \ \mathbf{m}_2 = \mathbf{g}_2 \cap \mathbf{n}_1^+.$ 

**Lemma 2.1**. We save the notation of section 1. Let  $\phi$  satisfy i), ii), iii), iv) and (2.3)

hold. Then the following statements are true.

- 1) The conjugation  $\sigma$  saves  $\mathbf{l}_1$ ,  $\mathbf{r}_1$ ,  $\overline{\mathbf{r}}_1$ ,  $\mathbf{g}_1$ ,  $\mathbf{m}_1$  and  $\mathbf{l}_2$ ,  $\mathbf{r}_2$ ,  $\overline{\mathbf{r}}_2$ ,  $\mathbf{g}_2$ ,  $\mathbf{m}_2$ . This defines the maps  $\sigma : \overline{\mathbf{r}}_1/\mathbf{l}_1 \mapsto \overline{\mathbf{r}}_1/\mathbf{l}_1$ ,  $\sigma : \overline{\mathbf{r}}_2/\mathbf{l}_2 \mapsto \overline{\mathbf{r}}_2/\mathbf{l}_2$ ;
- 2)  $p \in P_1^+ \cap P_2^-$ ; there exists the decomposition  $p = m_1 q m_2$  with  $m_1 \in \exp(\mathbf{m}_1)$ ,  $m_2 \in \exp(\mathbf{m}_2)$ ,  $q \in R'$ . The elements  $m_1$ ,  $m_2$ , q obey the following relations  $m_1 m_2 = m_2 m_1$ ,  ${}^{\sigma} m_1 = q^{-1} m_1^{-1} q$ ,  ${}^{\sigma} m_2 = q m_2^{-1} q^{-1}$ ,  ${}^{\sigma} q = q^{-1}$ ;
- 3) Denote  $p_1 = m_1 q$  and  $p_2 = q m_2$ . Then  ${}^{\sigma} p_1 = p_1^{-1}$  and  ${}^{\sigma} p_2 = p_2^{-1}$ ;
- 4)  $Ad_{p_1}(\overline{\mathbf{r}}_1) = \overline{\mathbf{r}}_1$ ,  $Ad_{p_1}(\mathbf{l}_1) = \mathbf{l}_1$ ,  $Ad_{p_2}(\overline{\mathbf{r}}_2) = \overline{\mathbf{r}}_2$ ,  $Ad_{p_2}(\mathbf{l}_2) = \mathbf{l}_2$ . This defines  $Ad_{p_1}: \overline{\mathbf{r}}_1/\mathbf{l}_1 \mapsto \overline{\mathbf{r}}_1/\mathbf{l}_1$ ,  $Ad_{p_2}: \overline{\mathbf{r}}_2/\mathbf{l}_2 \mapsto \overline{\mathbf{r}}_2/\mathbf{l}_2$  and

$$\phi(Ad_{p_1}^{-1\sigma}(x)) = Ad_{p_2}^{-1\sigma}\phi(x) \tag{2.4}$$

holds for  $x \in \overline{\mathbf{r}}_1/\mathbf{l}_1$ .

**Proof**. Consider the intesection of left and right sets of (2.3) with  $\mathbf{g}e$  and  $\mathbf{g}f$ .

We have

$$\begin{cases}
\sigma(\mathbf{l}_1 + \mathbf{n}_1^+) &= Ad_p(\mathbf{l}_1 + \mathbf{n}_1^+) \\
\sigma(\mathbf{l}_2 + \mathbf{n}_2^-) &= Ad_p(\mathbf{l}_2 + \mathbf{n}_2^-)
\end{cases}$$
(2.5)

Compairing the radicals of this subalgebras, we have

$$\begin{cases}
{}^{\sigma}\mathbf{n}_{1}^{+} = Ad_{p}\mathbf{n}_{1}^{+} \\
{}^{\sigma}\mathbf{n}_{2}^{-} = Ad_{p}\mathbf{n}_{2}^{-}
\end{cases} (2.6)$$

Since  ${}^{\sigma}\mathbf{n}_1^+ = \mathbf{n}_1^+$  and  ${}^{\sigma}\mathbf{n}_2^- = \mathbf{n}_2^-$ , then

$$\begin{cases}
\mathbf{n}_1^+ = Ad_p\mathbf{n}_1^+ \\
\mathbf{n}_2^- = Ad_p\mathbf{n}_2^-
\end{cases}$$
(2.7)

This implies

$$p \in P_1^+ \cap P_2^-$$
.

The element p can be uniquely decomposed

$$p = m_1 q m_2, \tag{2.8}$$

where  $m_1 \in exp(\mathbf{m}_1), m_2 \in exp(\mathbf{m}_2), q \in R'$ .

First we are going to prove statement 2) of the Lemma. Choose the elements  $\mu_1 \in \mathbf{m}_1$  and  $\mu_2 \in \mathbf{m}_2$  such that  $m_1 = exp(\mu_1)$ ,  $m_2 = exp(\mu_2)$ . Up to definition,

$$\mu_1 = \sum_{\alpha \in \Delta_1 - \Delta_2} \xi_{\alpha} E_{-\alpha}, \qquad \mu_2 = \sum_{\beta \in \Delta_2 - \Delta_1} \eta_{\beta} E_{\beta}$$

Suppose that  $[E_{-\alpha}, E_{\beta}] = cE_{\gamma}$  and  $c \neq 0$ . Then  $-\alpha + \beta = \gamma$  is a root. If  $\gamma > 0$ , then  $\beta = \alpha + \gamma$  and there exists the simple roots  $\Pi_1 - \Pi_2$  in decomposition of  $\beta$  in the sum of simple roots. This contradicts to  $\beta \in \Delta_2$ . If  $\gamma < 0$ , then  $\alpha = \beta - \gamma$  and similarly we get a contradiction. Hence  $[E_{-\alpha}, E_{\beta}] = 0$ . Therefore,  $[\mu_1, \mu_2] = 0$  and  $m_1 m_2 = m_2 m_1$ .

Since  ${}^{\sigma}p = p^{-1}$ , then

$${}^{\sigma}m_{1}{}^{\sigma}q^{\sigma}m_{2}=m_{2}^{-1}q^{-1}m_{1}^{-1}=m_{2}^{-1}(q^{-1}m_{1}^{-1}q)q^{-1}=$$

$$(q^{-1}m_1^{-1}q)m_2^{-1}q^{-1} = (q^{-1}m_1^{-1}q)q^{-1}(qm_2^{-1}q^{-1}).$$

Thus,  ${}^{\sigma}m_1=q^{-1}m_1^{-1}q$ ,  ${}^{\sigma}m_2=qm_2^{-1}q^{-1}$ ,  ${}^{\sigma}q=q^{-1}$ . This proves 2). One can deduce 3) from 2) by direct calculations.

Since  $\mathbf{g}_2 \cap \mathbf{l}_2 = 0$ , then  $Ad_{p_2}\mathbf{l}_2 = \mathbf{l}_2$ . Therefore,

$$Ad_p(\mathbf{l}_2 + \mathbf{n}_2^-) = Ad_p\mathbf{l}_2 + Ad_p\mathbf{n}_2^- = Ad_{m_1}\mathbf{l}_2 + \mathbf{n}_2^- = \mathbf{l}_2 + \mathbf{n}_2^-$$

Similarly,  $Ad_{p_1}\mathbf{l}_1 = \mathbf{l}_1$  and

$$Ad_p(\mathbf{l}_1 + \mathbf{n}_1^+) = Ad_p\mathbf{l}_1 + Ad_p\mathbf{n}_1^+ = Ad_{m_2}\mathbf{l}_1 + \mathbf{n}_1^+ = \mathbf{l}_1 + \mathbf{n}_1^+$$

Substituting this to (2.5), we have

$$\begin{cases}
\sigma(\mathbf{l}_1 + \mathbf{n}_1^+) = \mathbf{l}_1 + \mathbf{n}_1^+ \\
\sigma(\mathbf{l}_2 + \mathbf{n}_2^-) = \mathbf{l}_2 + \mathbf{n}_2^+
\end{cases}$$
(2.9)

Since  ${}^{\sigma}\mathbf{h} = \mathbf{h}$ , then

$$\begin{cases}
 \sigma(\mathbf{l}_1) &= \mathbf{l}_1 \\
 \sigma(\mathbf{l}_2) &= \mathbf{l}_2
\end{cases}$$
(2.10)

The (2.3) can be rewritten as

$$\mathbf{l}_{1}e + \mathbf{n}_{1}^{+}e + \sum_{x \in \overline{\mathbf{r}}_{1}} \mathbf{C}({}^{\sigma}xe + {}^{\sigma}(\phi(x))f) + \mathbf{n}_{2}^{-}f + \mathbf{l}_{2}f =$$

$$\mathbf{l}_{1}e + \mathbf{n}_{1}^{+}e + \sum_{x \in \overline{\mathbf{r}}_{1}} \mathbf{C}(Ad_{p}(x)e + Ad_{p}(\phi(x))f) + \mathbf{n}_{2}^{-}f + \mathbf{l}_{2}f$$

$$(2.11)$$

We stress that

 $Ad_p x = Ad_{p_1} x(mod(\mathbf{n}_1^+)), \ Ad_{p_1} x \in \overline{\mathbf{r}}_1 \subset \mathbf{r}_1 \text{ and } Ad_p \phi(x) = Ad_{p_2} \phi(x)(mod(\mathbf{n}_2^+)), Ad_{p_2} \phi(x) \in \overline{\mathbf{r}}_2 \subset \mathbf{r}_2.$  We obtain

$$\sum_{x \in \overline{\mathbf{r}}_1} \mathbf{C}({}^{\sigma}xe + {}^{\sigma}(\phi(x))f) \subset \mathbf{l}_1e + \mathbf{n}_1^+e + \sum_{x \in \overline{\mathbf{r}}_1} \mathbf{C}(Ad_{p_1}(x)e + Ad_{p_2}(\phi(x))f) + \mathbf{n}_2^-f + \mathbf{l}_2f \quad (2.12)$$

Since  $p_1 = h_1 g_1$  with  $h_1 \in H = exp(\mathbf{h}), g_1 \in G_1 = exp(\mathbf{g}_1)$ , then

$$\begin{cases}
Ad_{p_1}\overline{\mathbf{r}}_1 = \overline{\mathbf{r}}_1 \\
Ad_{p_2}\overline{\mathbf{r}}_2 = \overline{\mathbf{r}}_2
\end{cases}$$
(2.13)

$$\begin{cases}
Ad_{p_1}\mathbf{l}_1 &= \mathbf{l}_1 \\
Ad_{p_2}\mathbf{l}_2 &= \mathbf{l}_2
\end{cases}$$
(2.14)

The (2.12) implies that for every  $x \in \overline{\mathbf{r}}_1$  there exists  $y \in \overline{\mathbf{r}}_1$  such that

$$\begin{cases}
\sigma(x) = Ad_{p_1}y(mod(\mathbf{l}_1)) \\
\sigma(\phi(x)) = Ad_{p_2}\phi(y)(mod(\mathbf{l}_2))
\end{cases}$$
(2.15)

It follows

$$\begin{cases}
 \sigma(\overline{\mathbf{r}}_1) = \overline{\mathbf{r}}_1 \\
 \sigma(\overline{\mathbf{r}}_2) = \overline{\mathbf{r}}_2
\end{cases}$$
(2.16)

The (2.10) and (2.16) prove the statement 1).

We have seen above that  $Ad_{p_1}$  saves  $\overline{\mathbf{r}}_1$ ,  $\mathbf{l}_1$  and  $Ad_{p_2}$  saves  $\overline{\mathbf{r}}_2$ ,  $\mathbf{l}_2$ . Finally the (2.15) yields  $\phi(Ad_{p_1}^{-1\sigma}(x)) = Ad_{p_2}^{-1\sigma}\phi(x)$  for  $x \in \overline{\mathbf{r}}_1/\mathbf{l}_1$ .  $\square$ 

**Definition 2.2**. The following condition we shall call the condition  $v^*$ ) for  $\phi : \overline{\mathbf{r}}_1/\mathbf{l}_1 \mapsto \overline{\mathbf{r}}_2/\mathbf{l}_2$ :

 $\mathbf{v}^*$ )  ${}^{\sigma}\overline{\mathbf{r}}_i = \overline{\mathbf{r}}_i$ ,  ${}^{\sigma}\mathbf{l}_i = \mathbf{l}_i$  for i = 1, 2 and

$$^{\sigma}(\phi(x)) = \phi(^{\sigma}x) \tag{2.17}$$

for  $x \in \overline{\mathbf{r}}_1/\mathbf{l}_1$ .

Since  $\sigma: \Pi \to \Pi$ , then  ${}^{\sigma}\Pi_1 = \Pi_1$ ,  ${}^{\sigma}\Pi_2 = \Pi_2$  and  ${}^{\sigma}\phi(\alpha) = \phi({}^{\sigma}\alpha)$  for  $\alpha \in \Pi_1$ .

Notations:

 $\Pi_0 = \Pi_1 \bigcup \Pi_2;$ 

 $\mathbf{h}^{\phi} = \{ v \in \mathbf{h} | \alpha(v) = \phi(\alpha)v, \alpha \in \Pi_1 \} = \{ v \in \mathbf{h} | [v, \phi(x)] = \phi([v, x]), x \in \overline{\mathbf{r}}_1 \};$ 

The last formula can be rewritten as  $ad_v\phi(x) = \phi ad_v(x), x \in \overline{\mathbf{r}}_1$  We shall call  $\mathbf{h}^{\phi}$  the stabilizer subalgebra of  $\phi$ .

 $H^{\phi} = \{ p \in H | Ad_p \phi(x) = \phi Ad_p(x) \}$ . We shall call  $H^{\phi}$  the stabilizer of  $\phi$ . The Lie algebra of  $H^{\phi}$  is  $\mathbf{h}^{\phi}$ .

**Definition 2.3**[BD].Let  $\phi: \Pi_1 \mapsto \Pi_2$  satisfy i),ii),iii). Let  $\alpha, \beta \in \Pi_1$ . We shall say that  $\alpha < \beta$  if there exists integer k such that  $\phi^k(\alpha) = \beta$ .

Note that  $\alpha < \phi(\alpha) < \dots < \phi^k(\alpha) = \beta$ .

**Definition 2.4**. The set  $\alpha_1 < \alpha_2 < \cdots < \alpha_k$  is called a chain. We shall denote by  $C(\alpha)$  the maximal chain with  $\alpha$ .

**Lemma 2.5**. 1) The maximal chains either coinside or don't intersect.; 2) If  $\phi$  and  $\sigma$  obey v\*) (see Definition 2.2), then the image  $\sigma(C)$  of a maximal chain is also a maximal chain.

#### Proof.

- 1) Let  $C_1 = \{\alpha_1 < \alpha_2 < \dots < \alpha_k\}$  and  $C_2 = \{\beta_1 < \beta_2 < \dots < \beta_k\}$  are maximal chains. If  $\phi^n(\alpha) = \phi^m(\beta)$ ,  $n \ge m$  then  $\phi^{n-m}(\alpha) = \beta$  and, therefore,  $\beta \in C_1$ . Hence,  $C_2 \subset C_1$ . Since  $C_2$  is maximal,  $C_1 = C_2$ .
- 2) Let  $\{\alpha_1 < \alpha_2 < \dots < \alpha_k\}$  be a maximal chain with  $\alpha_{i+1} = \phi(\alpha_i)$ . Denote  $\beta_i = \sigma(\alpha_i)$ . We see  $\phi(\beta_i) = \phi(\sigma(\alpha_i)) = \sigma(\phi(\alpha_i)) = \sigma(\alpha_{i+1}) = \beta_{i+1}$ . Hence,  $\beta_1 < \beta_2 < \dots < \beta_k$ .  $\square$

Corollary 2.6 The  $\Pi_0$  is decomposed in maximal chains. The automorphism  $\sigma$  acts on the set of maximal chains.

**Theorem 2.7**. Let  $\mathbf{g}(\mathbf{R})$  be a simple Lie algebra of L1-L5 types. Suppose that the map  $\phi$  satisfies i)-iv) and the algebra  $Ad_g\mathbf{W}(\phi)$  is  $\sigma$ -invariant. Denote as above  $p = {}^{\sigma}g^{-1}g$ . Then  $\phi$  satisfies v\*) and  $p \in H^{\phi}$ .

**Proof**. The proof is divided into 2 steps.

1) We recall that (2.1) implies (2.2) with  $p = {}^{\sigma}g^{-1}g$ . We are going to show in this step that if  $Ad_q \mathbf{W}(\phi)$  is  $\sigma$ -invariant then  $\phi$  satisfies  $\mathbf{v}^*$ ).

It was proved that (2.1) implies (2.4). The equality (2.4) can be rewritten in the form

$$Ad_{p_2}^{-1\sigma}\phi^{\sigma}(Ad_{p_1}x) = \phi(x)$$
 (2.18)

for  $x \in \overline{\mathbf{r}}_1/\mathbf{l}_1$ .

Decompose  $\Pi_1$  and  $\Pi_2$  in connected components:

$$\Pi_1 = \bigcup_{i=1}^k \Pi_{1i}$$

$$\Pi_2 = \bigcup_{i=1}^k \Pi_{2i}$$

Then  $\mathbf{g}_1$  and  $\mathbf{g}_2$  are decomposed in direct sums:

$$\mathbf{g}_1 = \sum_{i=1}^k \mathbf{g}_{1i}$$

$$\mathbf{g}_2 = \sum_{i=1}^k \mathbf{g}_{2i}$$

We may assume that  $\phi(\mathbf{g}_{1i}) = \mathbf{g}_{2i}$ .

As we saw in Lemma 2.1  ${}^{\sigma}\mathbf{g}_1 = \mathbf{g}_1$ ,  ${}^{\sigma}\mathbf{g}_2 = \mathbf{g}_2$ . Since  $p_1 \in R_1$ , then  $Ad_{p_1}(\mathbf{g}_1) = \mathbf{g}_1$  and  $Ad_{p_1}$  save the decomposition of  $\mathbf{g}_1$  in the some of simple algebras. Simillarly,  $p_2 \in R_2$ , then  $Ad_{p_2}(\mathbf{g}_2) = \mathbf{g}_2$  and  $Ad_{p_2}$  save the decomposition of  $\mathbf{g}_2$  in the some of simple algebras. Denote  $\psi = {}^{\sigma}\phi^{\sigma}$ . According to (2.18), the maps  $\psi$  and  $\phi$  gave the same permutation of simple components. Thus  $\psi(\mathbf{g}_{1i}) = \mathbf{g}_{2i}$ . Therefore,  $\phi^{-1}\psi(\mathbf{g}_{1i}) = \mathbf{g}_{1i}$  and  $\phi^{-1}\psi(\Pi_{1i}) = \Pi_{1i}$ .

Denote  $\phi^{-1}\psi = s_{1i}$ . Then  $\psi = \phi s_{1i}$ . Substituting  $\phi s_{1i}$  for  $\psi$  in (2.18), we get

$$Ad_{p_2}^{-1}\phi s_{1i}(Ad_{p_1}x) = \phi(x)$$
 (2.19),

$$\phi^{-1}Ad_{p_2}^{-1}\phi s_{1i}(Ad_{p_1}x) = x \tag{2.19}$$

for  $x \in g_{1i}$ . Denote  $v_2 \in \mathbf{r}_2$  such that  $p_2 = exp(v_2)$ . Then

$$\phi^{-1}Ad_{p_2}^{-1}\phi(x) = \phi^{-1}exp(-ad_{v_2})\phi(x) = exp(-ad_{\phi^{-1}v_2})x$$

Therefore,  $\phi^{-1}Ad_{p_2}^{-1}\phi$  is an inner automorphism of  $\mathbf{g}_{1i}$ . We denote it by  $Ad_q$ . The equality  $Ad_qs_{1i}Ad_{p_1}=1$  implies that  $s_{1i}$  is inner automorphism of  $\mathbf{g}_{1i}$ . Since  $s_{1i}=\phi^{-1}\psi$  save  $\Pi_{1i}$ , then  $s_{1i}=1$ . Finally, we get  $\phi^{-1}\psi=1$  and  $\psi=\phi$ . This proves (2.17) for  $x \in \mathbf{g}_1$ . If x belongs to orthogonale complement of  $\mathbf{g}_1$  in  $\overline{\mathbf{r}}_1$ , then (2.4) is equivalent to (2.17). Consequently, (2.17) holds for all  $x \in \overline{\mathbf{r}}_1/\mathbf{l}_1$ .

2) The goal of this step is to prove that  $p_1 = p_2 = p \in H^{\phi}$ . As we prove in step 1), (2.17) and (2.4) hold. Then

$$\phi(Ad_{p_1}^{-1\sigma}(x)) = Ad_{p_2}^{-1\sigma}\phi(x) = Ad_{p_2}^{-1}\phi(\sigma x)$$

We get

$$\phi(Ad_{p_1}x) = Ad_{p_2}\phi(x), \tag{2.19}$$

for all  $x \in \overline{\mathbf{r}}_1/\mathbf{l}_1$ .

There exists  $v_i \in \mathbf{r}_i$  such that  $p_i = exp(v_i)$ . According Lemma 2.1, we have  $v_1 = \mu_1 + \theta + t$ ,  $v_2 = \mu_2 + \theta + t$  where  $\mu_i \in \mathbf{m}_i$ ,  $\theta \in (\mathbf{g}_1 \cap \mathbf{g}_2) \cap (\mathbf{n}^+ \oplus \mathbf{n}^-)$ ,  $t \in \mathbf{h}$ . We shall prove further that  $\mu_1 = \mu_2 = \theta = 0$ .

We can decompose  $t = t_i + t'_i$  where  $t_i \in \mathbf{h}_i = \mathbf{g}_i \cap \mathbf{h}$ ,  $t'_i \in \mathbf{h}$  and  $[t'_i, \mathbf{g}_i] = 0$ . Denoting  $g_i = exp(\mu + \theta + t_i)$ ,  $h_i = exp(t'_i)$ , we get  $p_i = g_i h_i = h_i g_i$ .

The (2.19) implies  $\phi(Ad_{g_1}x) = Ad_{g_2}\phi(x)$ . Let  $G_i$  be the subgroup of G with Lie algebra  $\mathbf{g}_i$ . We can extend  $\phi$  to the isomorphism of the groups  $\phi: G_1 \mapsto G_2$ . Then  $Ad_{\phi(g_1)}\phi(x) = Ad_{g_2}\phi(x)$ . Hence,  $\phi(g_1) = g_2$ . Therefore,  $\phi(\mu_1 + \theta + t_1) = \mu_2 + \theta + t_2$ . The element  $\theta$  is a sum  $\theta = \theta^+ + \theta^-$  where  $\theta^{\pm} \in \mathbf{g}_1 \cap \mathbf{g}_2 \cap \mathbf{n}^{\pm}$ . We obtain

$$\phi(\mu_1) + \phi(\theta^+) + \phi(\theta^-) + \phi(t_1) = \mu_2 + \theta^+ + \theta^- + t_2 \tag{2.20}$$

Recall that the map  $\phi$  is defined via the map  $\phi : \Pi_1 \mapsto \Pi_2$ . Thus,  $\phi(\mathbf{n}^+) \cap \mathbf{g}_1 \subset \mathbf{n}^+ \cap \mathbf{g}_2$ ,  $\phi(\mathbf{n}^-) \cap \mathbf{g}_1 \subset \mathbf{n}^- \cap \mathbf{g}_2$ ,  $\phi(\mathbf{h}_1) \subset \mathbf{h}_2$ . The equality (2.20) holds if and only if

$$\begin{cases}
\phi(\theta^{+}) = \mu_{2} + \theta^{+} \\
\phi(\mu_{1}) + \phi(\theta^{-}) = \theta^{-} \\
\phi(t_{1}) = t_{2}
\end{cases} (2.21)$$

Suppose that  $\theta^+ \neq 0$ . Denote  $\Delta_i^+$  the system if positive roots generated by  $\Pi_i$ . Then

$$\theta^+ = \sum_{\alpha \in \Delta_1^+} \xi_{\alpha} E_{\alpha}, \qquad \mu_2 = \sum_{\alpha \in \Delta_2^+ - \Delta_1^+} \xi_{\beta} E_{\beta}$$

The equality  $\phi(\theta^+) = \mu_2 + \theta^+$  implies

$$\sum_{\alpha \in \Delta_1^+} \xi_{\alpha} E_{\phi(\alpha)} = \sum_{\alpha \in \Delta_2^+ - \Delta_1^+} \xi_{\beta} E_{\beta} + \sum_{\alpha \in \Delta_1^+} \xi_{\alpha} E_{\alpha}$$
 (2.22)

For every simple root  $\alpha \in \Pi_1$  we denote  $l(\alpha)$  the length of the maximal chain  $C(\alpha)$  (see definition 2.4) from the beginning to  $\alpha$ . Denote

$$l_0 = \min\{l(\alpha) | \alpha \in \Delta_1^+, \xi_\alpha \neq 0\}$$

and  $l_0 = l(\alpha_0)$  for some positive root  $\alpha_0 \in \Delta_1^+$  The term  $\xi_{\alpha_0} E_{\alpha_0}$  belongs to the right side of (2.22) and dose not belong to to left side. A contradiction.

Therefore  $\theta^+ = 0$  and  $\mu_2 = 0$ . Similarly we get  $\theta^- = 0$  and  $\mu_1 = 0$ . This proves  $v_1 = v_2 = t$ ,  $p_1 = p_2 = exp(t) = p \in H$  and

$$Ad_p \phi = \phi Ad_p \tag{2.23}$$

This proves the Theorem.  $\Box$ 

Corollary 2.8 Let  $\phi: \Pi_1 \mapsto \Pi_2$  satisfy i)-iv). Consider the following conditions: 1)  $Ad_g \mathbf{W}(\phi)$  is  $\sigma$ -invariant, 2)  $\mathbf{W}(\phi)$  is q-invariant. Then 1) implies 2) and 2) is equivalent to  $\mathbf{v}^*$ ).

**Proof.** If  $Ad_g \mathbf{W}(\phi)$  is  $\sigma$ -invariant for some  $g \in G$ , then  $\phi$  obelies  $\mathbf{v}^*$ ). This proves  $1) \mapsto \mathbf{v}^*$  and  $2) \mapsto \mathbf{v}^*$ .

According to Lemma 2.1,if  $Ad_g \mathbf{W}(\phi)$  is  $\sigma$ -invariant, then  ${}^{\sigma} \mathbf{l}_i = \mathbf{l}_i$ ,  ${}^{\sigma} \mathbf{n}_1^+ = \mathbf{n}_1^+$ ,  ${}^{\sigma} \mathbf{n}_2^- = \mathbf{n}_2^-$ ,  ${}^{\sigma} \overline{\mathbf{r}}_i = \overline{\mathbf{r}}_i$ . The condition  $\mathbf{v}^*$ ) is true and yields

$$\sigma(xe + \phi(x)f) = \sigma xe + \sigma \phi(x)f = \sigma xe + \phi(\sigma x)f$$

for  $x \in \overline{\mathbf{r}}_1/\mathbf{l}_1$ . He have

$${}^{\sigma}\mathbf{W}(\phi) = {}^{\sigma}(\mathbf{l}_1e + \mathbf{n}_1^+e + \sum_{x \in \overline{\mathbf{r}}_1} \mathbf{C}(xe + \phi(x)f) + \mathbf{n}_2^-f + \mathbf{l}_2f) =$$

$$\mathbf{l}_1 e + \mathbf{n}_1^+ e + \sum_{x \in \overline{\mathbf{r}}_1} \mathbf{C}(xe + \phi(x)f) + \mathbf{n}_2^- f + \mathbf{l}_2 f = \mathbf{W}(\phi).$$

This proves  $1)\mapsto 2$ ) and  $v^*)\mapsto 2$ ).  $\square$ 

## Completion of classification

In this section we are going to classify all Manin triples  $\mathbf{g}(\mathbf{R}), W, \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R})$  up to the Ad-action of the group  $G(\mathbf{R}) = G(\mathbf{C})^{\Gamma}$ . As above  $\mathbf{g}(\mathbf{C}) = \mathbf{g}(\mathbf{R}) \otimes \mathbf{C}$ . Recall that  $\mathbf{g}(\mathbf{C}) = \mathbf{g}$  in the case  $\mathbf{g}(\mathbf{R})$  is a real form of a complex simple Lie algebra  $\mathbf{g}$  and  $\mathbf{g}(\mathbf{C}) = \mathbf{g} + \mathbf{g}$  in the case  $\mathbf{g}(\mathbf{R}) = \mathbf{g}_{\mathbf{R}}$  is the realification of  $\mathbf{g}$ . Denote  $\Pi^c$  the set of simple roots of  $\mathbf{g}(\mathbf{C})$ . So  $\Pi^c = \Pi$  in the case of real forms and  $\Pi^c = \Pi \cup \Pi'$  in the case  $\mathbf{g}(\mathbf{R})$  is the realification of  $\mathbf{g}$  (here  $\Pi,\Pi'$  are sets of simple roots of  $\mathbf{g}$ ). Let we have two subsets  $\Pi_1^c, \Pi_2^c$  in  $\Pi^c$ . Let  $\phi: \Pi_1^c \mapsto \Pi_2^c$  satisfy i)-iiv). The  $\mathbf{W}(\phi)$  is the subalgebra (1.2), constructed by the map  $\phi$  for  $\mathbf{g}(\mathbf{C})$ .

**Proposition 3.1.** Let  $\mathbf{g}(\mathbf{R})$  be a real simple Lie algebra if the type L1-L5. and let the subalgebra  $Ad_g\mathbf{W}(\phi)$  be  $\sigma$ -invariant. We assert  $(Ad_g\mathbf{W}(\phi))^{\Gamma} = Ad_r(\mathbf{W}(\phi))^{\Gamma}$  for some  $r \in G(\mathbf{R})$ .

**Proof.** Recall that we choose  $\sigma$ -invariant subalgebras  $\mathbf{n}^{\pm}$ ,  $\mathbf{h}$  in  $\mathbf{g}(\mathbf{C})$ . The conjugation  $\sigma$  saves the set of simple roots  $\Pi$ . According to Theorem 2.7., the element  $p = {}^{\sigma}g^{-1}g$  lies in  $H^{\phi}$ ,  $\mathbf{W}(\phi)$  is also  $\sigma$ -invariant and  $\phi$  satisfies conditions  $\mathbf{v}^*$ ). Then  $\phi({}^{\sigma}\alpha) = {}^{\sigma}\phi(\alpha)$  for  $\alpha \in \Pi_1$ .

Denote by v the element in  $\mathbf{h}$  such that p = exp(v). The element p uniquely determined by the system of numbers  $p_{\alpha} = exp(\alpha(v)), \ \alpha \in \Pi$ .

As p = exp(v), then  ${}^{\sigma}p = exp({}^{\sigma}v)$  and

$$({}^{\sigma}p)_{\alpha} = exp\alpha({}^{\sigma}v) = exp\overline{{}^{\sigma}\alpha(v)} = \overline{p_{\sigma(\alpha)}}$$

Easily

$$(p^{-1})_{\alpha} = \exp(\alpha(-v)) = p_{\alpha}^{-1}$$

The equality  ${}^{\sigma}p = p^{-1}$  implies

$$\overline{p_{\sigma(\alpha)}} = p_{\alpha}^{-1} \tag{3.1}$$

for all  $\alpha \in \Pi$ . Since  $p \in H^{\phi}$ , then  $Ad_p\phi(x) = \phi(Ad_p(x))$ ,  $x \in \overline{\mathbf{r}}_1/\mathbf{l}$ . This is equivalent to  $Ad_p\phi(E_{\alpha}) = \phi(Ad_p(E_{\alpha}))$ ,  $\alpha \in \Pi_1$ . We obtain

$$p_{\alpha} = p_{\phi(\alpha)} \tag{3.2}$$

for  $\alpha \in \Pi_1$ .

By Lemmas 2.5-2.6,  $\Pi_0$  is decomposes in maximal chains and  $\sigma$  acts on the chains by permutations. It follows from (3.2) that the system of elements  $p_{\alpha}$  is stable on maximal chains in  $\Pi_0$ . There exists a system  $h_{\alpha} \in \mathbb{C}^*$  which is stable on maximal chains and  $p_{\alpha} = \overline{h_{\sigma(\alpha)}}^{-1} h_{\alpha}$ ,  $\alpha \in \Pi$ . Therefore, there exists the element  $h \in H^{\phi}$  such that  $p = {}^{\sigma}h^{-1}h$ . By definition,  ${}^{\sigma}g^{-1}g = p = {}^{\sigma}h^{-1}h$ . Hence  ${}^{\sigma}(gh^{-1}) = gh^{-1}$  and  $gh^{-1} \in G(\mathbf{R})$ . Finally g = rh,  $r \in G(\mathbf{R})$ ,  $h \in H^{\phi}$ . We have

$$Ad_h \mathbf{W}(\phi) = Ad_h(\mathbf{l}_1 e + \mathbf{n}_1^+ e) + \sum_{x \in \overline{\mathbf{r}}_1} \mathbf{C}(Ad_h x e + Ad_h \phi(x) f) + Ad_h(\mathbf{n}_2^- f + \mathbf{l}_2 f) =$$

$$\mathbf{l}_1 e + \mathbf{n}_1^+ e + \sum_{x \in \overline{\mathbf{r}}_1} \mathbf{C}(Ad_h x e + \phi(Ad_h x) f) + \mathbf{n}_2^- f + \mathbf{l}_2 f = \mathbf{W}(\phi).$$

It follows  $Ad_g \mathbf{W}(\phi) = Ad_r \mathbf{W}(\phi)$  and  $(Ad_g \mathbf{W}(\phi))^{\Gamma} = Ad_r (\mathbf{W}(\phi))^{\Gamma}$ .

Let  $\mathbf{g}(\mathbf{R})$  be a real simple Lie algebra of types L1-L2. We choose  $\sigma$ -invariant subalgebras  $\mathbf{h}$ ,  $\mathbf{n}^{\pm}$  in  $\mathbf{g}(\mathbf{C})$  as above. The  $\mathbf{h}(\mathbf{R})$  is a real form of  $\mathbf{h}(\mathbf{C})$ . The conjugation  $\sigma$  acts onthe set of simple roots as in Satake-diagrams of L1-L5. Let  $\Pi_1^c$  and  $\Pi_2^c$  be  $\sigma$ -invariant subsets in the set of simple roots of  $\mathbf{g}(\mathbf{C})$ . Let the map  $\phi: \Pi_1 \to \Pi_2$  be  $\sigma$ -invariant and satisfy the conditions i), ii), iii).

Denote by  $\mathbf{g}_i(\mathbf{R})$ ,  $\mathbf{h}_i(\mathbf{R})$  the real forms of  $\mathbf{g}_i(\mathbf{C})$ ,  $\mathbf{h}_i(C)$ . Choose two real subspases  $\mathbf{h}_i(\mathbf{R})$  such that  $\overline{\mathbf{h}}_i(\mathbf{R})^{\perp} \subset \mathbf{h}_i(\mathbf{R})$  and  $\mathbf{h}_i(\mathbf{R}) \subset \overline{\mathbf{h}}_i(\mathbf{R})$ .

Denote  $\overline{\mathbf{r}}_i(\mathbf{R}) = \mathbf{g}_i(\mathbf{R}) + \overline{\mathbf{h}}_i(\mathbf{R})$ ,  $\mathbf{l}_i(\mathbf{R}) = \overline{\mathbf{h}}_i(\mathbf{R})^{\perp}$ . One can extend the map  $\phi$  to the isomorphism

$$\phi: \mathbf{g}_1(\mathbf{R}) \mapsto \mathbf{g}_2(\mathbf{R}).$$

Suppose  $\phi$  is extended to the isomorphism

$$\phi: \overline{\mathbf{r}}_1(\mathbf{R})/\mathbf{l}_1(\mathbf{R}) \mapsto \overline{\mathbf{r}}_2(\mathbf{R})/\mathbf{l}_2(\mathbf{R}),$$

preserving Killing form and having no fixed points. We shall refer the above condition as condition  $iv\mathbf{R}$ ) Consider the subalgebra

$$\mathbf{W}(\phi)_{\mathbf{R}} = \mathbf{l}_1(\mathbf{R}) + \mathbf{n}_1(\mathbf{R})^+ e + \sum_{x \in \overline{\mathbf{r}}_1(\mathbf{R})} \mathbf{R}(xe + \phi(x)f) + \mathbf{n}_2(\mathbf{R})^- f + \mathbf{l}_2(\mathbf{R})$$
(3.3)

**Theorem 3.2**. Let  $\phi$  be  $\sigma$ -invariant and satisfy i)-iii) and iv**R**). Let  $\mathbf{g}(\mathbf{R})$  be a simple Lie algebra of type L1-L5. Then

- 1) The triple  $(\mathbf{g}(\mathbf{R}), \mathbf{W}(\phi)_{\mathbf{R}}, \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R}))$  is a Manin triple;
- 2) Every Manin triple  $(\mathbf{g}(\mathbf{R}), V, \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R}))$  is  $Ad_{G(\mathbf{R})}$  equivalent to the triple  $(\mathbf{g}(\mathbf{R}), \mathbf{W}(\phi)_{\mathbf{R}}, \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R}))$ .

**Proof.** Consider the complexifications  $\mathbf{g}_i(\mathbf{C})$ ,  $\overline{\mathbf{r}}_i(\mathbf{C})$ ,  $\mathbf{l}_i(\mathbf{C})$  of  $\mathbf{g}_i(\mathbf{R})$ ,  $\overline{\mathbf{r}}_i(\mathbf{R})$ ,  $\mathbf{l}_i(\mathbf{R})$ . We extend the  $\phi$  to  $\mathbf{C}$ -linear  $\sigma$ -invariant isomorphism of  $\mathbf{g}_1(\mathbf{C})$  onto  $\mathbf{g}_2(\mathbf{C})$  and further to  $\overline{\mathbf{r}}_1(\mathbf{C})/\mathbf{l}_1(\mathbf{C})$  onto  $\overline{\mathbf{r}}_2(\mathbf{C})/\mathbf{l}_2(\mathbf{C})$ . We get  $\phi$ , satisfying conditions i)-iv), and v\*). Therefore,  $\mathbf{W}(\phi)$  is  $\sigma$ -invariant. The algebra  $\mathbf{W}(\phi)^{\Gamma}$  is generated by  $\mathbf{l}_1(\mathbf{R})$ ,  $\mathbf{l}_2(\mathbf{R})$ ,  $\mathbf{n}_1^+(\mathbf{R})$ ,  $\mathbf{n}_2^-(\mathbf{R})$  and the elements

$$xe + \phi(x)f + \sigma(xe + \phi(x)f) = (x + \sigma x)e + (\phi(x)f + \sigma\phi(x)f) = (x + \sigma x)e + \phi(x + \sigma x)f$$

where  $x \in \overline{\mathbf{r}}_1(\mathbf{C})/\mathbf{l}_1(\mathbf{C})$ . We have  $\mathbf{W}(\phi)^{\Gamma} = \mathbf{W}(\phi)_{\mathbf{R}}$  and  $(\mathbf{g}(\mathbf{R}), \mathbf{W}(\phi)_{\mathbf{R}}, \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R}))$  is a Manin triple. this proves 1).

Let  $(\mathbf{g}(\mathbf{R}), W, \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R}))$  be a Manin triple; then  $(\mathbf{g}(\mathbf{C}), W \otimes \mathbf{C}, \mathbf{g}(\mathbf{C}) \oplus \mathbf{g}(\mathbf{C}))$  is also a Manin triple. The algebra  $W \otimes \mathbf{C}$  is  $\sigma$ -invariant and, according to Theorem 1.2,  $W \otimes \mathbf{C} = Ad_g\mathbf{W}(\phi)$ . Then  $\phi$  satisfies  $\mathbf{v}^*$ ) (see Theorem 2.7) and we get the isomorphism  $\mathbf{i}\mathbf{v}\mathbf{R}$ ) of the algebras of invariants. We obtain  $(W \otimes \mathbf{C})^{\Gamma} = (Ad_g\mathbf{W}(\phi))^{\Gamma} = Ad_r(\mathbf{W}(\phi))^{\Gamma} = Ad_r\mathbf{W}(\phi)_{\mathbf{R}}$ . This proves the Theorem.

Corollary 3.3 Let  $\mathbf{g}(\mathbf{R})$  be the split form of a simple complex Lie algebra  $\mathbf{g}(\mathbf{R})$ . Then  $\mathbf{l}_1(\mathbf{R}) = \mathbf{l}_2(\mathbf{R}) = 0$  and

$$\mathbf{W}(\phi)_{\mathbf{R}} = \mathbf{n}_1(\mathbf{R})^+ e + \sum_{x \in \overline{\mathbf{r}}_1(\mathbf{R})} \mathbf{R}(xe + \phi(x)f) + \mathbf{n}_2(\mathbf{R})^- f$$
 (3.4)

All Manin triples  $(\mathbf{g}(\mathbf{R}), W, \mathbf{g}(\mathbf{R}) \oplus \mathbf{g}(\mathbf{R}))$  has the form  $W = Ad_r(\mathbf{W}(\phi)_{\mathbf{R}})$  with  $r \in G(\mathbf{R})$ .

**Proof.** The Killing form  $K(\cdot, \cdot)$  is positive definite on  $\mathbf{h}(\mathbf{R})$ . Therefore,  $\mathbf{l}_1(\mathbf{R}) = \mathbf{l}_2(\mathbf{R}) = 0.\square$ 

Further we shall specify the statement of Theorem 3.2 in the case  $\mathbf{g}(\mathbf{R})$  is a realification of  $\mathbf{g}$ .

**Theorem 3.4** Let  $\mathbf{g}$  be a complex simple Lie algebra and  $\mathbf{g}(\mathbf{R}) = \mathbf{g}_{\mathbf{R}}$  is the realification of  $\mathbf{g}$ . as in section 1 let  $\phi\Pi_1 \mapsto \Pi_2$  satisfy i)-iii). We extend  $\phi$  to  $\mathbf{R}$ -isomorphism  $\phi: \mathbf{g}_1 \mapsto \mathbf{g}_2$  such that  $\phi$  is  $\mathbf{C}$  linear or  $\mathbf{C}$ -antilinear on each simple component of  $\mathbf{g}_1$ . Let  $\overline{\mathbf{h}}_1$ ,  $\overline{\mathbf{h}}_2$  be the  $\mathbf{R}$ -subalgebras of  $\mathbf{h}$  such that  $\overline{\mathbf{h}}_i^{\perp} \subset \overline{\mathbf{h}}_i$  and  $\mathbf{h}_i \subset \overline{\mathbf{h}}_i$ . Denote as above  $\mathbf{l}_i = \overline{\mathbf{h}}_i^{\perp} = KerK(\cdot, \cdot)|_{\overline{\mathbf{h}}_i}$ ,  $\overline{\mathbf{r}}_i = \mathbf{g}_i + \overline{\mathbf{h}}_i$ . Suppose that  $\phi$  is extended to the  $\mathbf{R}$ -isomomorphism

$$\phi: \overline{\mathbf{r}}_1/\mathbf{l}_1 \mapsto \overline{\mathbf{r}}_2/\mathbf{l}_2$$

such that  $\phi$  preserve  $K(\cdot, \cdot)$  and has no fixed points in **h** (condition iv)). Consider the subalgebra  $\mathbf{W}(\phi)$  (see 1.2). Then

- 1)  $(\mathbf{g}_{\mathbf{R}}, \mathbf{W}(\phi), \mathbf{g}_{\mathbf{R}} \oplus \mathbf{g}_{\mathbf{R}}))$  is a Manin triple;
- 2) Every Manin triple  $(\mathbf{g}_{\mathbf{R}}, W', \mathbf{g}_{\mathbf{R}} \oplus \mathbf{g}_{\mathbf{R}})$  is gauge equivalent to  $(\mathbf{g}_{\mathbf{R}}, \mathbf{W}(\phi), \mathbf{g}_{\mathbf{R}} \oplus \mathbf{g}_{\mathbf{R}})$ . **Proof.** The complexification  $\mathbf{g}(\mathbf{C})$  of  $\mathbf{g}_{\mathbf{R}}$  equals to  $\mathbf{g} \oplus \mathbf{g}$ . The set of simple roots of  $\mathbf{g}(\mathbf{C})$  is  $\Pi^c = \Pi \bigcup \Pi'$  where  $\Pi = \{\alpha'_1, \dots, \alpha'_n\}$  and  $\Pi' = \{\alpha_1'', \dots, \alpha_n''\}$  are systems of symple roots of  $\mathbf{g}$ . The conjugation  $\sigma$  of  $\mathbf{g}_{\mathbf{R}}$  acts on  $\Pi^c$  by  $\sigma(\alpha_i) = \alpha'_i$  and  $\sigma(\alpha'_i) = \alpha_i$ .

For the subsets  $\Pi_1$ ,  $\Pi_2 \subset \Pi$  we consider  $\Pi_i^c = \Pi_i \cup \Pi_i'$ . Decompose  $\Pi_1$ ,  $\Pi_2$  in the union of connected components:

$$\Pi_1 = \bigcup_{s=1}^k \Pi_{1s}, \quad \Pi_2 = \bigcup_{s=1}^k \Pi_{2s}$$

Suppose  $\phi(\Pi_{1s}) = \Pi_{2s}$ . We extend  $\phi$  to  $\sigma$ -invariant map  $\phi : \Pi_1^c \mapsto \Pi_2^c$ , putting  $\phi(\Pi_{1s}) = \Pi_{1s}$ , if  $\phi|_{\mathbf{g}_{1s}}$  is C-linear, and  $\phi(\Pi_{1s}) = \Pi'_{1s}$ , if  $\phi|_{\mathbf{g}_{1s}}$  is C-antilinear. We get  $\sigma$ -invariant map  $\phi : \Pi_1^c \mapsto \Pi_2^c$ , satisfying i), ii), iii). We extend  $\phi$  to  $\sigma$ -invariant C-linear map  $\overline{\mathbf{r}}_1(\mathbf{C})/\mathbf{l}_1(\mathbf{C})$  onto  $\overline{\mathbf{r}}_2(\mathbf{C})/\mathbf{l}_2(\mathbf{C})$ , obeying i)-v\*). By Theorem 1.2,  $\mathbf{W}(\phi) \otimes \mathbf{C}$  is complementary subalgebra in Manin triple  $(\mathbf{g}(\mathbf{C}), \mathbf{W}(\phi) \otimes \mathbf{C}, \mathbf{g}(\mathbf{C}) \oplus \mathbf{g}(\mathbf{C}))$ . Considering  $\sigma$ -invariants, we see that  $\mathbf{W}(\phi)$  is complementary subalgebra in real Manin triple  $(\mathbf{g}, \mathbf{W}(\phi), \mathbf{g} \oplus \mathbf{g})$ . This proves 1).

On the other hand, if  $(\mathbf{g}_{\mathbf{R}}, W, \mathbf{g}_{\mathbf{R}} \oplus \mathbf{g}_{\mathbf{R}})$  is a real Manin triple, then W is gauge equivalent to  $\mathbf{W}(\phi)_{\mathbf{R}}$ . The last subalgebra is constructed by the two  $\sigma$ -invariant subsets  $\Pi_i^c = \Pi_i \cup \Pi_i'$  and  $\sigma$  invariant map  $\phi : \Pi_1^c \mapsto \Pi_2^c$ , obeying i)-v\*). If  $\phi(\Pi_{1s}) = \Pi_{1s}$  then the restriction of  $\phi$  to  $\mathbf{g}_{1s}(\mathbf{R}) = (\mathbf{g}_{1s}(\mathbf{C}))^{\Gamma}$  is C-linear. In the case  $\phi(\Pi_{1s}) = \Pi_{1s}'$  the restriction is C-antilinear. The algebra of invariants coinsides with  $Ad_r\mathbf{W}(\phi)$ .

# References

- [VO] E.Vinberg, A.Onishchik, Seminar on Lie Groups and algebraic groups, URSS, Moscow, 1995.
- [GG] M.Goto, F.Grosshans, Semisimple Lie algebras, Marcel Dekker, Inc., New York and Basel, 1978.
- [BD] A.Belavin, V.Drinfel'd, On classical Yang-Baxter equation for simple Lie algebras, Funct.Anal.Appl.16(1982).
- [S] A.Stolin, Some remarks on Lie bialgebra structures on simple complex Lie algebras, Preprint NO 1998-08/ISSN 0347-2809.
- [ST] M.Semenov-Tian-Shanskii, Poisson Lie groups. Quantum principle of duality and twisted quantum double, J.Theor.Math.Phys. 93, N2, 1992, 302-329.
- [CGR] M.Cahen, S.Gutt, J.Rawnsley, Some remarks on the classification of Poisson-Lie groups, Contemporary Math. 179, 1994, 1-15.
- [CH] V.Chloup, Bialgebra structure on a real semisimple Lie algebra, Bull. Belg. Math. Soc. 2, 1995, 265-278.