

# The Classification of Triangular Semisimple and Cosemisimple Hopf Algebras Over an Algebraically Closed Field

Pavel Etingof

Department of Mathematics, Rm 2-165  
Massachusetts Institute of Technology  
77 Massachusetts Avenue  
Cambridge, MA 02139

Shlomo Gelaki <sup>1</sup>

Mathematical Sciences Research Institute  
1000 Centennial Drive  
Berkeley, CA 94720

November 2, 1999

## 1 Introduction

In this paper we classify triangular semisimple and cosemisimple Hopf algebras over *any* algebraically closed field  $k$ . Namely, we construct, for each positive integer  $N$ , relatively prime to the characteristic of  $k$  if it is positive, a bijection between the set of isomorphism classes of triangular semisimple and cosemisimple Hopf algebras of dimension  $N$  over  $k$ , and the set of isomorphism classes of quadruples  $(G, H, V, u)$ , where  $G$  is a group of order  $N$ ,  $H$  is a subgroup of  $G$ ,  $V$  is an irreducible projective representation of  $H$  over  $k$  of dimension  $|H|^{1/2}$ , and  $u \in G$  is a central element of order  $\leq 2$ . This classification implies, in particular, that *any* triangular semisimple and cosemisimple Hopf algebra over  $k$  can be obtained from a group algebra by a twist (it was previously known only in characteristic 0 [EG1, Theorem 2.1]). It also implies that  $(G, H, V, u)$  corresponds to a *minimal* triangular semisimple Hopf algebra over  $k$  if and only if  $G$  is generated by  $H$  and  $u$ . We then answer positively the question from [EG2] whether the group underlying a *minimal* triangular semisimple Hopf

---

<sup>1</sup>This research was supported by THE ISRAEL SCIENCE FOUNDATION founded by the Israel Academy of Sciences and Humanities.

algebra is solvable by proving that the group  $H$  is a quotient of a central type group and hence solvable. We then conclude that any triangular semisimple and cosemisimple Hopf algebra over  $k$  of dimension bigger than 1 contains a non-trivial grouplike element.

The classification uses Deligne's theorem on Tannakian categories [De] and the results of the paper [M] in an essential way. The proof of solvability and existence of grouplike elements relies on a theorem from [HI], which is proved using the classification of finite simple groups. The classification in positive characteristic relies also on the lifting functor from [EG4].

Throughout the paper, the ground field  $k$  is assumed to be algebraically closed.

**Acknowledgments.** We are grateful to Donald Passman for telling us about central type groups and the paper [HI]. We thank Mikhail Movshev for useful comments on his results, and Robert Guralnick for his interest and useful conversations.

## 2 Twists

Let  $A$  be a Hopf algebra over a field  $k$ . Recall [Dr1] that a *twist* for  $A$  is an invertible element  $J \in A \otimes A$  which satisfies

$$(\Delta \otimes I)(J)J_{12} = (I \otimes \Delta)(J)J_{23} \text{ and } (\varepsilon \otimes I)(J) = (I \otimes \varepsilon)(J) = 1, \quad (1)$$

where  $I$  is the identity map of  $A$ .

If  $J$  is a twist for  $A$  and  $x$  is an invertible element of  $A$  then  $J^x = \Delta(x)J(x^{-1} \otimes x^{-1})$  is also a twist for  $A$ . We will call the twists  $J$  and  $J^x$  *gauge equivalent*. The element  $x$  will be called a *gauge transformation*.

Given a twist  $J$  for  $A$ , one can define a Hopf algebra  $(A^J, \Delta^J, \varepsilon)$  as follows:  $A^J = A$  as an algebra, the coproduct is determined by

$$\Delta^J(x) = J^{-1}\Delta(x)J$$

for all  $x \in A$ , and  $\varepsilon$  is the ordinary counit of  $A$ . If  $A$  is triangular with the universal  $R$ -matrix  $R$ , then so is  $A^J$ , with the universal  $R$ -matrix  $R^J = J_{21}^{-1}RJ$ . It is obvious that two gauge equivalent twists, when applied to a fixed (triangular) Hopf algebra, produce two isomorphic (triangular) Hopf algebras.

Let  $A = k[H]$  be the group algebra of a finite group  $H$ . We will say that a twist  $J$  for  $A$  is *minimal* if the right (and left) components of the  $R$ -matrix  $R^J = J_{21}^{-1}RJ$  span  $A$ , i.e. if the corresponding triangular Hopf algebra  $(A^J, J_{21}^{-1}J)$  is minimal [R].

Let  $(A, R)$  be any triangular semisimple and cosemisimple Hopf algebra over  $k$ . Then the Drinfeld element  $u$  of  $A$  is a grouplike element of order  $\leq 2$ . Moreover, by [LR] in characteristic 0, and by [EG4, Theorem 3.1] in positive characteristic, the square of the antipode of  $A$  is the identity map, and hence  $u$  is central.

If  $k$  does not have characteristic 2, set

$$R_u = \frac{1}{2}(1 \otimes 1 + 1 \otimes u + u \otimes 1 - u \otimes u).$$

If  $k$  is of characteristic 2 (in which case  $u = 1$  by semisimplicity), set  $R_u = 1$ . Then  $(A, RR_u)$  is triangular semisimple and cosemisimple with Drinfeld element 1. This observation allows to reduce questions about triangular semisimple and cosemisimple Hopf algebras over  $k$  to the case when the Drinfeld element is 1.

### 3 Twists for Group Algebras

In this section we will prove the following theorem, which will be used later for classification.

**Theorem 3.1** *Let  $G, G'$  be finite groups,  $H, H'$  subgroups of  $G, G'$  respectively, and  $J, J'$  minimal twists for  $k[H], k[H']$  respectively. Suppose that the triangular Hopf algebras  $k[G]^J, k[G']^{J'}$  are isomorphic. Then there exists a group isomorphism  $\phi : G \rightarrow G'$  such that  $\phi(H) = H'$ , and  $(\phi \otimes \phi)(J)$  is gauge equivalent to  $J'$  as twists for  $k[H']$ .*

The rest of the section is devoted to the proof of the theorem.

**Lemma 3.2** *Let  $\mathcal{C}$  be the category of  $k$ -representations of a finite group, and  $F_1, F_2 : \mathcal{C} \rightarrow \text{Vect}(k)$  be two fiber functors (i.e. exact and faithful symmetric tensor functors, see [DM]) from  $\mathcal{C}$  to the category of  $k$ -vector spaces. Then  $F_1$  is isomorphic to  $F_2$ .*

**Proof:** This is a special case of [DM, Theorem 3.2]. ■

The following corollary of this lemma answers positively Movshev's question [M, Remark 1] whether any symmetric twist is trivial.

**Corollary 3.3** *Let  $G$  be a finite group, and  $J$  be a symmetric twist for  $k[G]$  (i.e.  $J_{21} = J$ ). Then  $J$  is gauge equivalent to  $1 \otimes 1$ .*

**Proof:** Let  $\mathcal{C}$  be the category of representations of  $G$ . We have two symmetric tensor structures on the forgetful functor  $F : \mathcal{C} \rightarrow \text{Vect}(k)$ ; namely, the trivial one and the one defined by  $J$ . By Lemma 3.2, the two fiber functors corresponding to these structures are isomorphic. But by definition, an isomorphism between them is an invertible element  $x \in k[G]$  such that  $J = \Delta(x)(x^{-1} \otimes x^{-1})$ . ■

**Remark 3.4** Here is another proof of Corollary 3.3 (in the case when the characteristic of  $k$  is relatively prime to  $|G|$ ) which does not use Lemma 3.2 but uses the results of [M]. Consider the  $G$ -coalgebra  $B_J = k[G]$  with coproduct  $\tilde{\Delta}(x) = (x \otimes x)J$ , and the dual algebra  $B_J^*$ . According to [M], this algebra is semisimple,  $G$  acts transitively on its simple ideals, and  $B_J^*$ , along with the action of  $G$ , completely determines  $J$  up to gauge transformations. Clearly, since  $J$  is symmetric, this algebra is commutative. So, it is isomorphic, as a  $G$ -algebra, to the algebra of functions on a set  $X$  on which  $G$  acts simply transitively. Corollary 3.3 now follows from the fact that such a  $G$ -set is unique up to an isomorphism (the group  $G$  itself with  $G$  acting by left multiplication). ■

**Lemma 3.5** *Let  $G, G'$  be finite groups,  $J, J'$  twists for  $k[G], k[G']$  respectively, and suppose that the triangular Hopf algebras  $k[G]^J, k[G']^{J'}$  are isomorphic. Then there exists a group isomorphism  $\phi : G \rightarrow G'$  such that  $(\phi \otimes \phi)(J)$  is gauge equivalent to  $J'$ .*

**Proof:** Let  $f : k[G]^J \rightarrow k[G']^{J'}$  be an isomorphism of triangular Hopf algebras. Then  $f$  defines an isomorphism of triangular Hopf algebras  $k[G] \rightarrow k[G']^{J'(f \otimes f)(J)^{-1}}$ . This implies that the element  $J'(f \otimes f)(J)^{-1}$  is a symmetric twist for  $k[G']$ . Thus, for some invertible  $x \in k[G']$  one has  $J'(f \otimes f)(J)^{-1} = \Delta(x)(x^{-1} \otimes x^{-1})$ . Let  $\phi = \text{Ad}(x^{-1}) \circ f : k[G] \rightarrow k[G']$ . It is obvious that  $\phi$  is a Hopf algebra isomorphism, so it comes from a group isomorphism  $\phi : G \rightarrow G'$ . We have  $(\phi \otimes \phi)(J) = \Delta(x)^{-1} J'(x \otimes x)$ , as desired. ■

We can now prove Theorem 3.1. By Lemma 3.5, it is sufficient to assume that  $G' = G$ , and that  $J$  is gauge equivalent to  $J'$  as twists for  $k[G]$ , and it is enough to show that there exists an element  $a \in G$  such that  $aHa^{-1} = H'$  and  $(a \otimes a)J(a^{-1} \otimes a^{-1})$  is gauge equivalent to  $J'$  as twists for  $k[H']$ .

So let  $x \in k[G]$  be the invertible element such that  $\Delta(x)J(x^{-1} \otimes x^{-1}) = J'$ . In particular, this implies that  $(x \otimes x)R(x^{-1} \otimes x^{-1}) = R'$ , where  $R, R'$  are the  $R$ -matrices corresponding to  $J, J'$  respectively. By the minimality of  $J, J'$ , we have  $xk[H]x^{-1} = k[H']$ . Thus,

$$J_0 = \Delta(x)(x^{-1} \otimes x^{-1}) = J'(x \otimes x)J^{-1}(x^{-1} \otimes x^{-1}) \in k[H']^{\otimes 2}.$$

It is obvious that  $J_0$  is a symmetric twist for  $k[H']$ , so by Corollary 3.3, it is gauge equivalent to  $1 \otimes 1$ . Thus,  $x = x_0 a$ , for some invertible  $x_0 \in k[H']$ , and  $a \in G$ . It is clear that  $aHa^{-1} = H'$ , and  $\Delta(x_0^{-1})J'(x_0 \otimes x_0) = (a \otimes a)J(a^{-1} \otimes a^{-1})$ . This concludes the proof of Theorem 3.1.

## 4 Construction of Triangular Semisimple and Cosemisimple Hopf Algebras from Group-Theoretical Data

Let  $H$  be a finite group such that  $|H|$  is not divisible by the characteristic of  $k$ . Suppose that  $V$  is an irreducible projective representation of  $H$  over  $k$  satisfying  $\dim V = |H|^{1/2}$ . Following the idea of the proof of Proposition 5 in [M], we will construct a twist  $J \in k[H] \otimes k[H]$ .

Let  $\pi : H \rightarrow PGL(V)$  be the projective action of  $H$  on  $V$ , and let  $\tilde{\pi} : H \rightarrow SL(V)$  be any lifting of this action ( $\tilde{\pi}$  need not be a homomorphism). We have  $\tilde{\pi}(x)\tilde{\pi}(y) = c(x, y)\tilde{\pi}(xy)$ , where  $c$  is a 2-cocycle with coefficients in  $k^*$ . By [M, Proposition 11], this cocycle is nondegenerate (see [M]) and hence by [M, Proposition 12] the representation of  $H$  on  $\text{End}_k(V)$  is isomorphic to the regular representation.

**Remark 4.1** In the paper [M] it is assumed that the characteristic of  $k$  is equal to 0, but all the results generalize in a straightforward way to the case when the characteristic of  $k$  is positive and relatively prime to the order of the group.

Consider the simple coalgebra  $(\text{End}_k V)^*$  with comultiplication  $\Delta$ . Clearly  $H$  acts on this coalgebra, and this representation of  $H$  is also isomorphic to the regular representation. In particular we can choose an element  $\lambda \in (\text{End}_k V)^*$  such that the set  $\{a \cdot \lambda | a \in H\}$  forms a basis of  $(\text{End}_k V)^*$ , and  $\langle \lambda, I \rangle = 1$  where  $I$  is the unit element of  $\text{End}_k V$ . Now, write  $\Delta(\lambda) = \sum_{a,b \in H} \gamma(a,b) a \cdot \lambda \otimes b \cdot \lambda$ , and set  $J = \sum_{a,b \in H} \gamma(a,b) a \otimes b \in k[H] \otimes k[H]$ . We claim that  $J$  is a twist for  $k[H]$ .

Indeed, let  $\tilde{\Delta} : k[H] \rightarrow k[H]^{\otimes 2}$  be determined by  $a \mapsto (a \otimes a)J$ , and let  $f : k[H] \rightarrow (\text{End}_k V)^*$  be determined by  $a \mapsto a \cdot \lambda$ . Clearly,  $f$  is an isomorphism of  $H$ -modules which satisfies  $\Delta(f(a)) = (f \otimes f)\tilde{\Delta}(a)$ . Therefore  $(k[H], \tilde{\Delta}, \varepsilon)$  is a coalgebra where  $\varepsilon = f^*(I)$ , which is equivalent to saying that  $J$  satisfies (1). In order to show that  $J$  is a twist it remains to show that it is invertible. The invertibility of  $J$  is proved in [M, Proposition 13]. We reproduce the proof (in a slightly expanded form) for the convenience of the reader. Suppose on the contrary that  $J$  is not invertible. Then there exists  $0 \neq L \in \text{End}_k V \otimes \text{End}_k V$  such that  $JL = 0$ . Let  $F : (\text{End}_k V)^* \otimes (\text{End}_k V)^* \rightarrow (\text{End}_k V)^* \otimes (\text{End}_k V)^*$  be defined by  $F(x) = xL$ . Clearly,  $F$  is a morphism of  $H \times H$  representations, and  $F \circ \tilde{\Delta} = 0$ . Thus the image  $\text{Im}(F^*)$  of the morphism of  $H \times H$  representations  $F^* : \text{End}_k V \otimes \text{End}_k V \rightarrow \text{End}_k V \otimes \text{End}_k V$  is contained in the kernel of the multiplication map  $m := \tilde{\Delta}^*$ . Let  $U := (\text{End}_k V \otimes 1)\text{Im}(F^*)(1 \otimes \text{End}_k V)$ . Clearly,  $U$  is contained in the kernel of  $m$  too. But, for any  $x \in U$  and  $h \in H$ ,  $(1 \otimes X_h)x(1 \otimes X_h)^{-1} \in U$  (as  $\text{End}_k V = \text{sp}\{X_h | h \in H\}$  with the relations  $X_h X_{h'} = c(h, h')X_{hh'}$ , i.e. conjugation by  $X_h$  is the same as the action of  $h$ ). Thus,  $U$  is a left  $\text{End}_k V \otimes \text{End}_k V$  module under left multiplication. Similarly, it is a right module over this algebra under right multiplication. So, it is a bimodule over  $\text{End}_k V \otimes \text{End}_k V$ . Since  $U \neq 0$ , this implies that  $U = \text{End}_k V \otimes \text{End}_k V$ . This is a contradiction, since we get that  $m = 0$ . Hence  $J$  is invertible.

It is straightforward to see that two different choices of  $\lambda$  produce two gauge equivalent twists  $J$  for  $k[H]$ , so the equivalence class of  $J$  is canonically associated to  $(H, V)$ .

Now, for any group  $G \supseteq H$ , whose order is prime to the characteristic of  $k$ , define a triangular semisimple Hopf algebra  $F(G, H, V) = (k[G]^J, J_{21}^{-1}J)$ . We wish to show that it is also cosemisimple.

**Lemma 4.2** *The Drinfeld element of the triangular semisimple Hopf algebra  $(A, R) = F(G, H, V)$  equals 1.*

**Proof:** The Drinfeld element  $u$  is a grouplike element of  $A$ , and for any finite-dimensional  $A$ -module  $V$  one has  $\text{tr}|_V(u) = \dim_{\text{Rep}(A)} V = \dim V$  (since  $\text{Rep}(A)$  is equivalent to  $\text{Rep}(G)$ , see [EG1, Section 1]). In particular, we can set  $V$  to be the regular representation, and find that  $\text{tr}|_A(u) = \dim(A) \neq 0$  in  $k$ . But it is clear that if  $g$  is a non-trivial grouplike element in any finite-dimensional Hopf algebra  $A$ , then  $\text{tr}|_A(g) = 0$ . Thus,  $u = 1$ . ■

**Corollary 4.3** *The triangular semisimple Hopf algebra  $(A, R) = F(G, H, V)$  is cosemisimple.*

**Proof:** Since  $u = 1$ , one has  $S^2 = I$  and hence  $A$  is cosemisimple (as  $\dim(A) \neq 0$ ). ■

Thus we have assigned a triangular semisimple and cosemisimple Hopf algebra with Drinfeld element  $u = 1$  to any triple  $(G, H, V)$  as above.

## 5 The Classification in Characteristic 0

In this section we assume that  $k$  is of characteristic 0.

**Theorem 5.1** *The assignment  $F : (G, H, V) \mapsto (A, R)$  is a bijection between isomorphism classes of triples  $(G, H, V)$  where  $G$  is a finite group,  $H$  is a subgroup of  $G$ , and  $V$  is an irreducible projective representation of  $H$  over  $k$  satisfying  $\dim V = |H|^{1/2}$ , and isomorphism classes of triangular semisimple Hopf algebras over  $k$  with Drinfeld element  $u = 1$ .*

**Proof:** We need to construct an assignment  $F'$  in the other direction, and check that both  $F' \circ F$  and  $F \circ F'$  are the identity assignments.

Let  $(A, R)$  be a triangular semisimple Hopf algebra over  $k$  whose Drinfeld element  $u$  is 1. It follows from Deligne's theorem on Tannakian categories (see [EG1, Theorem 2.1]) that there exist finite groups  $H \subseteq G$ , and a *minimal* twist  $J \in k[H] \otimes k[H]$ , such that  $(A, R) \cong (k[G]^J, J_{21}^{-1}J)$  as triangular Hopf algebras. As we proved in Section 3, these data are unique up to isomorphism and gauge transformations.

Following Movshev [M], define a coalgebra  $B_J$  which is  $k[H]$  as a vector space, with coproduct  $\hat{\Delta}(x) = (x \otimes x)J$ ,  $x \in H$ , and the usual counit. This coalgebra has a natural  $H$ -action by left multiplication. It follows from [M] that the coalgebra  $B_J$  is simple (see [EG3] for more explanations). Thus, the dual algebra  $B_J^*$  is simple as well, and carries an action of  $H$ . So we see that  $B_J^*$  is isomorphic to  $\text{End}_k(V)$  for some vector space  $V$ , and we have a homomorphism  $\pi : H \rightarrow \text{PGL}(V)$ . Thus  $V$  is a projective representation of  $H$ . It is shown in [M, Proposition 8] that this representation is irreducible, and it is obvious that  $\dim V = |H|^{1/2}$ .

It is clear that the isomorphism class of the representation  $V$  does not change if  $J$  is replaced by a twist  $J'$  which is gauge equivalent to  $J$  as twists for  $k[H]$ . Thus, to any isomorphism class of triangular semisimple Hopf algebras  $(A, R)$  over  $k$  with Drinfeld element 1, we have assigned an isomorphism class of triples  $(G, H, V)$ . Let us write this as  $(G, H, V) = F'(A, R)$ .

Thus, we have constructed the map  $F'$ .

The identity  $F \circ F' = \text{id}$  follows from [M, Proposition 5]. The identity  $F' \circ F = \text{id}$  follows from Theorem 3.1. ■

Now let  $(G, H, V, u)$  be a quadruple, in which  $(G, H, V)$  is as above, and  $u$  is a central element of  $G$  of order  $\leq 2$ . We extend the map  $F$  to quadruples by setting  $F(G, H, V, u) = (A, RR_u)$ , where  $(A, R) = F(G, H, V)$ .

**Theorem 5.2** *The assignment  $F$  is a bijection between isomorphism classes of quadruples  $(G, H, V, u)$  where  $G$  is a finite group,  $H$  is a subgroup of  $G$ ,  $V$  is an irreducible projective representation of  $H$  over  $k$  satisfying  $\dim V = |H|^{1/2}$ , and  $u \in G$  is a central element of order  $\leq 2$ , and isomorphism classes of triangular semisimple Hopf algebras over  $k$ .*

**Proof:** Define  $F'$  by  $F'(A, R) = (F'(A, RR_u), u)$ , where  $F'(A, RR_u)$  is defined in the proof of Theorem 5.1. It is straightforward to see that both  $F' \circ F$  and  $F \circ F'$  are the identity assignments. ■

Theorems 5.1 and 5.2 imply the following classification result for minimal triangular semisimple Hopf algebras over  $k$ .

**Proposition 5.3**  *$F(G, H, V, u)$  is minimal if and only if  $G$  is generated by  $H$  and  $u$ .*

**Proof:** As we have already pointed out, if  $(A, R) = F(G, H, V)$  then the sub Hopf algebra  $k[H]^J \subseteq A$  is minimal triangular. Therefore, if  $u = 1$  then  $F(G, H, V)$  is minimal if and only if  $G = H$ . This obviously remains true for  $F(G, H, V, u)$  if  $u \neq 1$  but  $u \in H$ . If  $u \notin H$  then it is clear that the  $R$ -matrix of  $F(G, H, V, u)$  generates  $k[H']$ , where  $H' = H \cup uH$ . This proves the proposition. ■

**Remark 5.4** As was pointed out already by Movshev, the theory developed in [M] and extended here is an analogue, for finite groups, of the theory of quantization of skew-symmetric solutions of the classical Yang-Baxter equation, developed by Drinfeld [Dr2]. In particular, the operation  $F$  is the analogue of the operation of quantization in [Dr2].

## 6 The Classification in Positive Characteristic

In this section we assume that  $k$  is of positive characteristic  $p$ , and prove an analogue of Theorem 5.2 by using this theorem itself and the lifting techniques from [EG4].

Let  $F$  be the assignment from Section 4. We now have the following.

**Theorem 6.1** <sup>1</sup> *The assignment  $F$  is a bijection between isomorphism classes of quadruples  $(G, H, V, u)$  where  $G$  is a finite group of order prime to  $p$ ,  $H$  is a subgroup of  $G$ ,  $V$  is an irreducible projective representation of  $H$  over  $k$  satisfying  $\dim V = |H|^{1/2}$ , and  $u \in G$  is a central element of order  $\leq 2$ , and isomorphism classes of triangular semisimple and cosemisimple Hopf algebras over  $k$ .*

**Proof:** As in the proof of Theorem 5.2 we need to construct the assignment  $F'$ .

We recall some notation from [EG4]. Let  $\mathcal{O} = W(k)$  be the ring of Witt vectors of  $k$ , and  $K$  the field of fractions of  $\mathcal{O}$  (it is of characteristic 0). Let  $\bar{K}$  be the algebraic closure of  $K$ .

Let  $(A, R)$  be a triangular semisimple and cosemisimple Hopf algebra over  $k$ . Lift it (see [EG4]) to a triangular semisimple Hopf algebra  $(\bar{A}, \bar{R})$  over  $K$ . By Theorem 5.2 we have that  $(\bar{A} \otimes_K \bar{K}, \bar{R}) = F(G, H, V, u)$ . We can now reduce  $V$  “mod  $p$ ” to get  $V_p$  which is an irreducible projective representation of  $H$  over the field  $k$ . This can be done since  $V$  is defined by a

---

<sup>1</sup>The proof is incomplete but a complete proof is given in [E, Remark 4.4].

nondegenerate 2-cocycle  $c$  (see [M]) with values in roots of unity of degree  $|H|^{1/2}$  (as the only irreducible representation of the simple  $H$ -algebra with basis  $\{X_h|h \in H\}$ , and relations  $X_gX_h = c(g, h)X_{gh}$ ). This cocycle can be reduced  $\text{mod } p$  and remains nondegenerate (since the groups of roots of unity of order  $|H|^{1/2}$  in  $k$  and  $K$  are naturally isomorphic), so it defines an irreducible projective representation  $V_p$ . Define  $F'(A, R) = (G, H, V_p, u)$ . It is shown like in characteristic 0 that  $F \circ F'$  and  $F' \circ F$  are the identity assignments. ■

**Corollary 6.2** *Any triangular semisimple and cosemisimple Hopf algebra over  $k$  is obtained from a group algebra by a twist.*

**Remark 6.3** Previously the statement of the corollary was only known in characteristic 0 [EG1, Theorem 2.1], and the best result known to us in positive characteristic was [EG4, Theorem 3.9].

**Proposition 6.4** *Proposition 5.3 holds in positive characteristic.*

**Proof:** As before, if  $(A, R) = F(G, H, V)$  then the sub Hopf algebra  $k[H]^J \subseteq A$  is minimal triangular. This follows from the facts that it is true in characteristic 0, and that the rank of a triangular structure does not change under lifting. Thus, Proposition 5.3 holds in characteristic  $p$ . ■

**Remark 6.5** In view of the results of this section, the results of our previous paper [EG3] generalize, without changes, to cotriangular semisimple and cosemisimple Hopf algebras in positive characteristic.

## 7 The Solvability of the Group Underlying a Minimal Triangular Semisimple Hopf Algebra

A classical fact about complex representations of finite groups is that the dimension of any irreducible representation of a finite group  $K$  does not exceed  $|K : Z(K)|^{1/2}$ , where  $Z(K)$  is the center of  $K$ . Groups of central type are those groups for which this inequality is in fact an equality. More precisely, a finite group  $K$  is said to be of *central type* if it has an irreducible representation  $V$  such that  $(\dim V)^2 = |K : Z(K)|$  (see e.g. [HI]). We shall need the following theorem (conjectured by Iwahori and Matsumoto in 1964) whose proof uses the classification of finite simple groups.

**Theorem 7.1** [HI, Theorem 7.3] *Any group of central type is solvable.*

As corollaries, we have the following results.

**Corollary 7.2** *Let  $A$  be a minimal triangular semisimple Hopf algebra over  $k$ , and let  $G$  be the finite group such that the categories of representations  $\text{Rep}(G)$  and  $\text{Rep}(A)$  are equivalent. Then  $G$  is solvable.*



**Proof:** We may assume that  $k$  has characteristic 0 (otherwise we can lift to characteristic 0), and by Proposition 5.3, that the Drinfeld element of  $A$  is 1. By Theorem 5.1, the corresponding group  $G$  has an irreducible projective representation  $V$  with  $\dim V = |G|^{1/2}$ . Let  $K$  be a finite central extension of  $G$  with central subgroup  $Z$ , such that  $V$  lifts to a linear representation of  $K$ . We have  $(\dim V)^2 = |K : Z|$ . Since  $(\dim V)^2 \leq |K : Z(K)|$  we get that  $Z = Z(K)$  and hence that  $K$  is a group of central type. But by Theorem 7.1,  $K$  is solvable and hence  $G \cong K/Z(K)$  is solvable as well. ■

**Remark 7.3** Movshev conjectures in the introduction to [M] that any finite group with a nondegenerate 2-cocycle is solvable. As explained in the Proof of Corollary 7.2, this result follows from Theorem 7.1.

**Corollary 7.4** *Let  $A$  be a triangular semisimple and cosemisimple Hopf algebra over  $k$  of dimension bigger than 1. Then  $A$  has a non-trivial grouplike element.*

**Proof:** We can assume that the Drinfeld element  $u$  is equal to 1 and that  $A$  is not cocommutative. Let  $A_0$  be the minimal part of  $A$ . By Corollary 7.2,  $A_0 = k[H]^J$  for a solvable group  $H$ ,  $|H| > 1$ . Therefore,  $A_0$  has non-trivial 1-dimensional representations. Since  $A_0 \cong A_0^{*op}$  as Hopf algebras, we get that  $A_0$ , and hence  $A$ , has non-trivial grouplike elements. ■

Corollary 7.4 motivates the following question.

**Question 7.5** *Let  $(A, R)$  be a quasitriangular semisimple and cosemisimple Hopf algebra over  $k$  (e.g. the quantum double of a semisimple and cosemisimple Hopf algebra), and let  $\dim(A) > 1$ . Is it true that  $A$  possesses a non-trivial grouplike element?*

Note that a positive answer to this question would imply that for a semisimple and cosemisimple Hopf algebra  $A$  over  $k$  either  $A$  or  $A^*$  possesses a non-trivial grouplike element. Such a result is very desirable for the problem of the classification of semisimple Hopf algebras.

## 8 Group-Theoretical Data Corresponding to Minimal Triangular Hopf Algebras Constructed from a Bijective 1-Cocycle

In this section we determine the group-theoretical data corresponding, under the bijection of the classification given in Theorem 5.1, to the minimal triangular semisimple Hopf algebras constructed in [EG2, Section 4]. We will use the definitions and notation from [EG2].

Let  $k = \mathbf{C}$ . Let  $G$  be a finite group,  $A$  be a finite abelian group with a left  $G$ -action  $(g, a) \mapsto g \cdot a$ , and  $\pi : G \rightarrow A$  be a bijective 1-cocycle, i.e. a bijective map such that  $\pi(gg') = \pi(g) + g\pi(g')$  (in particular,  $|G| = |A|$ ). Let  $H = G \rtimes A^*$  be the semidirect product

of  $G$  by the dual group  $A^*$  to  $A$ . Following [EG2, Section 4], we can associate to this data the element

$$J = |A|^{-1} \sum_{g \in G, b \in A^*} e^{(\pi(g), b)} b \otimes g$$

(for convenience we use the opposite element to the one from [EG2]). We proved in [EG2] that this element is a minimal twist for  $k[H]$ , so  $k[H]^J$  is a minimal triangular semisimple Hopf algebra with Drinfeld element  $u = 1$ . Now we wish to find the irreducible projective representation  $V$  of  $H$  which corresponds to  $k[H]^J$  under the correspondence of Theorem 5.1.

We will now construct the representation  $V$  and show that it is the one corresponding to  $k[H]^J$ .

Let  $V = \text{Fun}(A, k)$  be the space of  $k$ -valued functions on  $A$ . It has a basis  $\{\delta_a | a \in A\}$  of characteristic functions of points. Define a projective action  $\phi$  of  $H$  on  $V$  by

$$\phi(b)\delta_a = e^{-(a, b)}\delta_a, \quad \phi(g)\delta_a = \delta_{g \cdot a + \pi(g)} \text{ and } \phi(bg) = \phi(b)\phi(g)$$

for  $g \in G$  and  $b \in A^*$ . It is straightforward to verify that this is indeed a projective representation.

**Proposition 8.1** *The representation  $V$  is irreducible, and corresponds to  $k[H]^J$  under the bijection of the classification given in Theorem 5.1.*

**Proof:** Let  $B_J$  be the coalgebra  $(k[H], \tilde{\Delta})$  where  $\tilde{\Delta}(x) = (x \otimes x)J$ ,  $x \in H$ . Let  $B_J^*$  be the dual algebra. It is enough to show that the  $H$ -algebras  $B_J^*$  and  $\text{End}_k(V)$  are isomorphic.

Let us compute the multiplication in the algebra  $B_J^*$ . We have

$$\tilde{\Delta}(bg) = |A|^{-1} \sum_{g' \in G, b' \in B} e^{(\pi(g'), b')} b(g \cdot b')g \otimes bgg'. \quad (2)$$

Let  $\{Y_{bg}\}$  be the dual basis of  $B_J^*$  to the basis  $\{bg\}$  of  $B_J$ . Let  $*$  denote the multiplication law dual to the coproduct  $\tilde{\Delta}$ . Then, dualizing equation (2), we have

$$Y_{b_2g_2} * Y_{b_1g_1} = e^{(\pi(g_1) - \pi(g_2), b_2 - b_1)} Y_{b_1g_2} \quad (3)$$

for  $g_1, g_2 \in G$  and  $b_1, b_2 \in A^*$  (here for convenience we write the operations in  $A$  and  $A^*$  additively). Define  $Z_{bg} := e^{(\pi(g), b)} Y_{bg}$  for  $g \in G$  and  $b \in A^*$ . In the basis  $\{Z_{bg}\}$  the multiplication law in  $B_J^*$  is given by

$$Z_{b_2g_2} * Z_{b_1g_1} = e^{(\pi(g_1), b_2)} Z_{b_1g_2}. \quad (4)$$

Now let us introduce a left action of  $B_J^*$  on  $V$ . Set

$$Z_{bg}\delta_a = e^{(a, b)}\delta_{\pi(g)}. \quad (5)$$

It is straightforward to check using (4) that (5) is indeed a left action. It is also straightforward to compute that this action is  $H$ -equivariant. Thus, (5) defines an isomorphism  $B_J^* \rightarrow \text{End}_k(V)$  as  $H$ -algebras, which proves the proposition. ■

## References

- [Dr1] V. Drinfeld, On Almost Cocommutative Hopf Algebras, *Leningrad Math. J* **1** (1990), 321-342.
- [Dr2] V. Drinfeld, Constant quasiclassical solutions of the quantum Yang-Baxter equation, *Dokl. Acad. Nauk. SSSR* **273 No. 3** (1983), 531-535.
- [De] P. Deligne, Categories Tannakiennes, In The Grothendick Festschrift, Vol. II, Prog. Math. **87** (1990), 111-195.
- [DM] P. Deligne and J. Milne, Tannakian Categories, Lecture Notes in Mathematics **900**, 101-228, 1982.
- [E] P. Etingof, On faithfulness of the lifting for Hopf algebras and fusion categories, *arXiv:1704.07855*.
- [EG1] P. Etingof and S. Gelaki, Some Properties of Finite-Dimensional Semisimple Hopf Algebras, *Mathematical Research Letters* **5** (1998), 191-197.
- [EG2] P. Etingof and S. Gelaki, A Method of Construction of Finite-Dimensional Triangular Semisimple Hopf Algebras, *Mathematical Research Letters* **5** (1998), 551-561.
- [EG3] P. Etingof and S. Gelaki, The Representation Theory of Cotriangular Semisimple Hopf Algebras, *International Mathematics Research Notices* **7** (1999), 387-394.
- [EG4] P. Etingof and S. Gelaki, On Finite-Dimensional Semisimple and Cosemisimple Hopf Algebras In Positive Characteristic, *International Mathematics Research Notices* **16** (1998), 851-864.
- [HI] R. B. Howlett and I. M. Isaacs, On Groups of Central Type, *Mathematische Zeitschrift* **179** (1982), 555-569.
- [LR] R. G. Larson and D. E. Radford, Semisimple Cosemisimple Hopf Algebras, American J. of Mathematics **110** (1988), 187-195.
- [M] M. Movshev, Twisting in group algebras of finite groups, *Func. Anal. Appl.* **27** (1994), 240-244.
- [R] D. E. Radford, Minimal quasitriangular Hopf algebras, J. of Algebra **157** (1993), 285-315.