# CONFIGURATIONS OF LINEAR SUBSPACES AND RATIONAL INVARIANTS

#### DMITRI ZAITSEV

ABSTRACT. We construct a birational equivalence between certain quotients of s-tuples of equidimensional linear subspaces of  $\mathbb{C}^n$  and some quotients of products of square matrices modulo diagonal conjugations. In particular, we prove the rationality of the quotient space of s-tuples of linear 2-planes in  $\mathbb{C}^n$  modulo the diagonal  $\mathbf{GL}_n(\mathbb{C})$ -action . Furthermore, we compute generators of the field of the rational invariants explicitly.

# 1. Introduction

Let  $\mathbf{Gr}_{n,d}(\mathbb{C})$  denote the Grassmannian of all d-dimensional linear subspaces in  $\mathbb{C}^n$  and  $\mathbf{GL}_n(\mathbb{C}) \times (\mathbf{Gr}_{n,d}(\mathbb{C}))^s \to (\mathbf{Gr}_{n,d}(\mathbb{C}))^s$  the canonical diagonal action. I. V. Dolgachev posed the following question:

Is the quotient  $\mathbf{Gr}_{n,2}(\mathbb{C})^s/\mathbf{GL}_n(\mathbb{C})$  (e.g. in the sense of Rosenlicht) always rational?

Recall that a Rosenlicht quotient of an algebraic variety X acted by an algebraic group G is an algebraic variety V together with a rational map  $X \to V$  whose generic fibers coincide with the G-orbits. Such quotients always exist and are unique up to birational isomorphisms ([R56]). In the sequel all quotients will be assumed of this type. An algebraic variety Q is rational, if it is birationally equivalent to  $\mathbb{P}^m$  with  $m = \dim Q$ .

We answer the above question positively by applying the rationality of the quotient  $(\mathbf{GL}_2(\mathbb{C}))^2/\mathbf{GL}_2(\mathbb{C})$ , where  $\mathbf{GL}_2(\mathbb{C})$  acts diagonally by conjugations (see [P67] and surveys [D95, B91]):

**Theorem 1.1.** For all positive integers n and s, the quotient  $(\mathbf{Gr}_{n,2}(\mathbb{C}))^s/\mathbf{GL}_n(\mathbb{C})$  is rational. Equivalently, the field of rational  $\mathbf{GL}_n(\mathbb{C})$ -invariants on  $(\mathbf{Gr}_{n,2}(\mathbb{C}))^s$  is pure transcendental.

The statement of Theorem 1.1 has been recently proved by Megyesi [M98] in the case n=4 and by Dolgachev [DB96] in the case of odd n. Their proofs are independent of the present one.

More generally, we show the birational equivalence between  $(\mathbf{Gr}_{n,d}(\mathbb{C}))^s/\mathbf{GL}_n(\mathbb{C})$  and certain quotients of matrix spaces. Let  $\mathbf{GL}_n(\mathbb{C}) \times (\mathbf{GL}_n(\mathbb{C}))^s \to (\mathbf{GL}_n(\mathbb{C}))^s$  be the action defined by  $(g, M_1, \ldots, M_s) \mapsto (gM_1g^{-1}, \ldots, gM_sg^{-1})$ . The first main result of the present paper consists of the following two statements:

**Theorem 1.2.** 1. Let s and d be arbitrary positive integers and n = rd for some integer r > 1. Then  $(\mathbf{Gr}_{n,d}(\mathbb{C}))^s/\mathbf{GL}_n(\mathbb{C})$  is birationally equivalent to  $(\mathbf{GL}_d(\mathbb{C}))^k/\mathbf{GL}_d(\mathbb{C})$ , where

$$k = \begin{cases} (r-1)(s-r-1) & \text{if } s > r+1\\ 1 & \text{else} \end{cases}$$

2. Let s be arbitrary positive integer and n=(2r+1)e, d=2e for some integers r and e. Then  $(\mathbf{Gr}_{n,d}(\mathbb{C}))^s/\mathbf{GL}_n(\mathbb{C})$  is birationally equivalent to  $(\mathbf{GL}_e(\mathbb{C}))^k/\mathbf{GL}_e(\mathbb{C})$ , where

$$k = \begin{cases} 2r - 4 + (4r - 2)(s - r - 2) & \text{if } r > 1, \ s > r + 1 \\ 2(s - 4) & \text{if } r = 1, \ s > 4 \\ 1 & \text{else} \ . \end{cases}$$

Theorem 1.1 follows from the first part of Theorem 1.2 and from the rationality of the quotient of  $(\mathbf{GL}_2(\mathbb{C}))^s$  by  $\mathbf{GL}_2(\mathbb{C})$  (see Procesi [P67]) in the case n is even and from the second part of Theorem 1.2 in the case n is odd. More recently Formanek [F79, F80] proved the rationality of  $(\mathbf{GL}_n(\mathbb{C}))^s/\mathbf{GL}_n(\mathbb{C})$  for n=3,4. Using this result together with Theorem 1.2 we obtain:

**Theorem 1.3.** For every positive integer s, the following holds:

- 1.  $(\mathbf{Gr}_{n,3}(\mathbb{C}))^s/\mathbf{GL}_n(\mathbb{C})$  is rational for  $n = 0 \pmod{3}$ ;
- 2.  $(\mathbf{Gr}_{n,4}(\mathbb{C}))^s/\mathbf{GL}_n(\mathbb{C})$  is rational for  $n=0 \pmod{2}$ ;
- 3.  $(\mathbf{Gr}_{n,5}(\mathbb{C}))^s/\mathbf{GL}_n(\mathbb{C})$  is rational for  $n=3 \pmod 6$ ;
- 4.  $(\mathbf{Gr}_{n,8}(\mathbb{C}))^s/\mathbf{GL}_n(\mathbb{C})$  is rational for  $n=4 \pmod 8$ ;

We refer the reader to [BS88] for similar equivalences of stable rationalities (an algebraic variety V is stable rational, if  $V \times \mathbb{P}^m$  is rational for some m). We also refer to [GP71, GP72] for the classification of quadruples of linear subspaces of arbitrary dimensions and their invariants. In our situation, however, all rational invariants of the quadruples (i.e. the case s=4) are constant unless r=2 in the first part of Theorem 1.2.

Our method is based on constructing certain normal forms for our algebraic group actions. We call an algebraic variety acted algebraically by an algebraic group G a G-space. A G-subspace is a G-invariant locally closed algebraic subvariety of X. We use the standard notation  $Gs:=\{gs:g\in G\}$  and  $GS:=\{gs:g\in G,s\in S\}$ , where  $S\subset X$  is an arbitrary subset.

**Definition 1.1.** We say that (S, H) is a normal form for (X, G), if  $H \subset G$  is a subgroup and  $S \subset X$  is a H-subspace such that the following holds:

- 1. GS is Zariski dense in X;
- 2.  $Gs \cap S = Hs$  for all  $s \in S$ .

Clearly these conditions guarantee the birational equivalence of the quotients X/G and S/H. In §3 and 4 we construct certain normal forms that are isomorphic to the spaces of matrices as in Theorem 1.2. Then in §5 we use these normal forms for explicit computations of generators of the fields of rational invariants in each case of Theorem 1.2.

This method has also applications to biholomorphic automorphisms of non-smooth bounded domains, where the configurations of linear subspaces appear naturally as collections of tangent subspaces to the so-called characteristic webs. We refer the reader to [Z98] for further details.

# 2. Notation.

For brevity we write  $\mathbf{GL}_n$  and  $\mathbf{Gr}_{n,d}^s$  for  $\mathbf{GL}_n(\mathbb{C})$  and  $(\mathbf{Gr}_{n,d}(\mathbb{C}))^s$  respectively. The actions on the products will be assumed diagonal unless otherwise specified. For E and E' vector spaces,  $\dim E \leq \dim E'$ , denote by  $\mathbf{Gr}_d(E)$  the Grassmannian of all d-dimensional subspaces of E, by  $\mathbf{GL}(E)$  the group of linear automorphisms of E and by  $\mathbf{GL}(E_1, E_2)$  the space of all linear embeddings of  $E_1$  into  $E_2$ . A G-space X is homogeneous (almost homogeneous), if G acts transitively on X (on a Zariski dense subset of X).

## 3. The case d divides n.

In this section we study the diagonal  $\mathbf{GL}_n$ -action on  $\mathbf{Gr}_{n,d}^s$  with n = rd for some integer r. Clearly the set of all r-tuples  $(V_1, \ldots, V_r) \in \mathbf{Gr}_{n,d}^r$  such that  $\mathbb{C}^n = V_1 \oplus \cdots \oplus V_r$  is Zariski open and  $\mathbf{GL}_n$ -homogeneous. This can be reformulated in terms of normal forms:

**Lemma 3.1.** Suppose that  $r \geq 2$ ,  $s \geq r$  and  $(E_1, \ldots, E_r) \in \mathbf{Gr}_{n,d}^r$  is such that  $\mathbb{C}^n = E_1 \oplus \cdots \oplus E_r$ . Define

$$S_1 = S_1(s) := \{ (V_1, \dots, V_s) \in \mathbf{Gr}_{n,d}^s : (V_1, \dots, V_r) = (E_1, \dots, E_r) \}$$

and the group  $H_1 := \mathbf{GL}(E_1) \times \cdots \times \mathbf{GL}(E_r) \subset \mathbf{GL}_n$ . Then  $(S_1, H_1)$  is a normal form for  $(\mathbf{Gr}_{n,d}^s, \mathbf{GL}_n)$ .

Now fix the splitting  $\mathbb{C}^n = E_1 \oplus \cdots \oplus E_r$  as above. Let  $V \in \mathbf{Gr}_{n,d}$  be such that its projection on each  $E_i$ ,  $i=1,\ldots,r$ , is bijective. In this case we say that V is in general position with respect to  $(E_1,\ldots,E_r)$ . Clearly the subset of all V, which are in general position, is Zariski open in  $\mathbf{Gr}_{n,d}$ . Then for every  $i=2,\ldots,r$ , the projection  $V_i$  of V on  $E_1 \times E_i$  is a graph of a linear isomorphism  $\varphi_i \colon E_1 \to E_i$ . We claim that this correspondence between elements  $V \in \mathbf{Gr}_{n,d}$  and (r-1)-tuples of linear isomorphisms

$$(\varphi_2,\ldots,\varphi_r)\in\prod_{i=2}^r\mathbf{GL}(E_1,E_i)$$

is birational.

Lemma 3.2. Under the assumptions of Lemma 3.1 define

$$\Phi \colon \prod_{i=2}^r \mathbf{GL}(E_1, E_i) \to \mathbf{Gr}_{n,d},$$

$$\Phi(\varphi_2,\ldots,\varphi_r):=\{(z\oplus\varphi_2(z)\oplus\cdots\oplus\varphi_r(z)):z\in E_1\}$$

and the  $H_1$ -action on  $\prod_{i=2}^r \mathbf{GL}(E_1, E_i)$  by

$$((g_1,\ldots,g_r),(\varphi_2,\ldots,\varphi_r))\mapsto (g_2\circ\varphi_2\circ g_1^{-1},\ldots,g_r\circ\varphi_r\circ g_1^{-1}).$$

Then  $\Phi$  is birational and  $H_1$ -equivariant. The image of  $\Phi$  consists of all subspaces  $V \in \mathbf{Gr}_{n,d}$  which are in general position with respect to  $(E_1, \ldots, E_r)$ .

*Proof.* Since  $\Phi$  defines a system of standard coordinates on  $\mathbf{Gr}_{n,d}$ , it is birational. The equivariance and the statement on the image are straightforward.

Since  $\prod_{i=2}^r \mathbf{GL}(E_1, E_i)$  is  $H_1$ -homogeneous, we obtain:

**Corollary 3.1.** Under the assumptions of Lemma 3.1 let  $S_1'(s) \subset S_1(s)$  be the Zariski open subset, where  $V_{r+1}$  is in general position with respect to  $(E_1, \ldots, E_r)$ . Then  $H_1$  acts transitively on  $S_1'(r+1)$ .

This proves statement 1 of Theorem 1.2 for  $s \le r+1$ . Suppose now that s > r+1 and  $E_{r+1} \in \mathbf{Gr}_{n,d}$  is in general position with respect to  $(E_1, \ldots, E_r)$ .

**Lemma 3.3.** Define  $S_2 = S_2(s) := \{(V_1, \dots, V_s) \in S_1(s) : V_{r+1} = E_{r+1}\}$  and the group  $H_2 := \mathbf{GL}(E_1)$  with the action on  $S_2$  defined by the homomorphism

$$H_2 \to H_1, \quad g \mapsto (g, \varphi_2 \circ g \circ \varphi_2^{-1}, \dots, \varphi_r \circ g \circ \varphi_r^{-1}),$$

where  $(\varphi_2, \ldots, \varphi_r) := \Phi^{-1}(E_{r+1})$ . Then  $(S_2, H_2)$  is a normal form for  $(\mathbf{Gr}_{n,d}^s, \mathbf{GL}_n)$ .

*Proof.* By Corollary 3.1,  $H_1S_2$  is Zariski open in  $S_1$ . By straightforward calculations,  $(S_2, H_2)$  is a normal form for  $(S_1, H_1)$  and hence, by Lemma 3.1, a normal form for  $(\mathbf{Gr}_{n,d}^s, \mathbf{GL}_n)$ .

Let  $V \in \mathbf{Gr}_{n,d}$  be one more subspace in general position with respect to  $(E_1, \ldots, E_r)$ . Define  $(\psi_2, \ldots, \psi_r) := \Phi^{-1}(V)$ . Clearly the map

$$\prod_{i=2}^{r} \mathbf{GL}(E_1, E_i) \to \mathbf{GL}(E_1)^{r-1}, \quad (\psi_2, \dots, \psi_r) \mapsto (\varphi_2^{-1} \circ \psi_2, \dots, \varphi_r^{-1} \circ \psi_r) \quad (1)$$

is one-to-one, where  $(\varphi_2, \ldots, \varphi_r) = \Phi^{-1}(E_{r+1})$  is fixed. Moreover, we obtain:

**Lemma 3.4.** Define  $K : \mathbf{GL}(E_1)^{r-1} \to \mathbf{Gr}_{n,d}$  by

$$K(\chi_2,\ldots,\chi_r) := \Phi(\varphi_2 \circ \chi_2,\ldots,\varphi_r \circ \chi_r)$$

and the  $H_2$ -action on  $X_2$  by

$$(g,(\chi_2,\ldots,\chi_r))\mapsto (g\circ\chi_2\circ g^{-1},\ldots,g\circ\chi_r\circ g^{-1}).$$

Then K is birational and  $H_2$ -equivariant. The image of K consists of all subspaces  $V \in \mathbf{Gr}_{n,d}$  which are in general position with respect to  $(E_1, \ldots, E_r)$ .

*Proof.* The mapping K is birational as the composition of  $\Phi$  and the map (1). The other statements are straightforward.

Corollary 3.2. Under the assumptions of Lemma 3.3 define

$$\Psi \colon S_2(s) \to (\mathbf{GL}(E_1)^{r-1})^{s-r-1},$$

$$\Psi(E_1,\ldots,E_{r+1},V_{r+2},\ldots,V_s) := (K^{-1}(V_{r+2}),\ldots,K^{-1}(V_s)).$$

Then  $\Psi$  is birational and  $H_2$ -equivariant.

Finally we obtain together with Lemma 3.3:

Corollary 3.3. The space  $\mathbf{Gr}_{n,d}^s$ , where n = rd, is almost  $\mathbf{GL}_n$ -homogeneous if and only if  $s \leq r+1$ . If s > r+1, there is a normal form for  $(\mathbf{Gr}_{n,d}^s, \mathbf{GL}_n)$  which is isomorphic to  $(\mathbf{GL}_d^{(r-1)(s-r-1)}, \mathbf{GL}_d)$ .

This implies the first part of Theorem 1.2.

4. The case 
$$n = (2r + 1)e$$
 and  $d = 2e$ .

We start with an r-tuple  $(V_1, \ldots, V_r) \in \mathbf{Gr}_{n,d}^r$ . Clearly the subset of all r-tuples which form a direct sum is Zariski open and  $\mathbf{GL}_n$ -homogeneous. Fix an r-tuple  $(E_1, \ldots, E_r)$  in this subset and suppose that s > r. A straightforward calculation yields:

Lemma 4.1. Define

$$S_1 = S_1(s) := \{(V_1, \dots, V_s) \in \mathbf{Gr}_{n,d}^s : (V_1, \dots, V_r) = (E_1, \dots, E_r)\}$$

and the group  $H_1 := \{g \in \mathbf{GL}_n : g(E_i) = E_i \text{ for all } i = 1, \dots, r\}$  with the diagonal action on  $S_1$ . Then  $(S_1, H_1)$  is a normal form for  $(\mathbf{Gr}_{n.d}^s, \mathbf{GL}_n)$ .

Now we wish to parametrize the d-dimensional linear subspaces in  $\mathbb{C}^n$  with respect to  $(E_1, \ldots, E_r)$ . We say that V is in general position with respect to  $(E_1, \ldots, E_r)$ , if dim W = e, where  $W := V \cap (E_1 \oplus \cdots \oplus E_r)$ , and the projection of W on each  $E_i$ ,  $i = 1, \ldots, r$ , is injective. Clearly the subset of all  $V \in \mathbf{Gr}_{n,d}$ , which are in general position, is Zariski open.

We first give another description of the subspaces  $W \in \mathbf{Gr}_e(E_1 \oplus \cdots \oplus E_r)$  which are in general position with respect to  $(E_1, \ldots, E_r)$ . For this let  $A_i = A_i(V) \in \mathbf{Gr}_e(E_i)$  be the projections of W and let  $\varphi_i = \varphi_i(V) \in \mathbf{GL}(A_1, E_i)$  be the linear isomorphisms whose graphs are equal to the projections of W on  $E_1 \oplus E_i$ ,  $i = 2, \ldots, r$ . Denote by  $X = X(E_1, \ldots, E_r)$  the space of all tuples  $(A, \varphi_2, \ldots, \varphi_r)$ , where  $A \in \mathbf{Gr}_e(E_1)$  and  $\varphi_i \in \mathbf{GL}(A, E_i)$ , with the standard structure of a quasiprojective variety.

Lemma 4.2. Define

$$\Phi \colon X \to \mathbf{Gr}_e(E_1 \oplus \cdots \oplus E_r),$$

$$\Phi(A, \varphi_2, \dots, \varphi_r) := \{ (z \oplus \varphi_2(z) \oplus \dots \oplus \varphi_r(z)) : z \in A \}$$

and the  $H_1$ -action on X by

$$(g, (A, \varphi_2, \dots, \varphi_r)) \mapsto (g(A), g \circ \varphi_2 \circ g^{-1}, \dots, g \circ \varphi_r \circ g^{-1}).$$

Then  $\Phi$  is birational and  $H_1$ -equivariant. The image of  $\Phi$  consists of all subspaces  $W \in X$  which are in general position with respect to  $(E_1, \ldots, E_r)$ .

*Proof.* The proof is straightforward.

Corollary 4.1. Let  $S'_1(s) \subset S_1(s)$  be the Zariski open subset of all s-tuples  $\mathcal{V} = (V_1, \ldots, V_s)$  such that  $V_{r+1}$  is in general position with respect to  $(E_1, \ldots, E_r)$ . Then  $H_1$  acts transitively on  $S'_1(r+1)$ .

*Proof.* It follows from the general position condition that

$$W(\mathcal{V}) := V_{r+1} \cap (E_1 \oplus \cdots \oplus E_r) \in \mathbf{Gr}_e(E_1 \oplus \cdots \oplus E_r)$$

in the notation of Lemma 4.2. Let  $\mathcal{V}, \mathcal{V}' \in S_1'(r+1)$  be arbitrary elements. Since X is  $H_1$ -homogeneous, there exists  $g_1 \in H_1$  such that  $g_1(W(\mathcal{V})) = W(\mathcal{V}')$ . Then there exists  $g_2 \in \mathbf{GL}_n$  with  $g_2(V_{r+1}) = V_{r+1}'$  and  $g_2|(E_1 \oplus \cdots \oplus E_r) = \mathrm{id}$ . By the construction,  $g_2 \in H_1$  and the proof is finished.

Corollary 4.2. For  $s \leq r+1$ ,  $\mathbf{Gr}_{n,d}^s$  is almost  $\mathbf{GL}_n$ -homogeneous.

Let s > r+1 and  $E_{r+1} \in \mathbf{Gr}_{n,d}$  be in general position with respect to  $(E_1, \ldots, E_r)$ . As a direct consequence of Corollary 4.1 we obtain:

**Lemma 4.3.** Define  $S_2 := \{(V_1, \ldots, V_s) \in S_1 : V_{r+1} = E_{r+1}\}$  and  $H_2 := \{g \in \mathbf{GL}_n : g(E_i) = E_i \text{ for all } i = 1, \ldots, r+1\}$  with the diagonal action on  $S_2$ . Then  $(S_2, H_2)$  is a normal form for  $(\mathbf{Gr}_{n,d}^s, \mathbf{GL}_n)$ .

For the sequel we suppose that  $(E_1, \ldots, E_{r+1})$  as above is fixed. Define

$$(A, \varphi_2, \dots, \varphi_r) := \Phi^{-1}(E_{r+1} \cap (E_1 \oplus \dots \oplus E_r)).$$

Let  $V \in \mathbf{Gr}_{n,d}$  be one more subspace which is in general position with respect to  $(E_1, \ldots, E_r)$ . Clearly V is not uniquely determined by its e-dimensional intersection  $Z_1(V) := V \cap (E_1 \oplus \cdots \oplus E_r)$ . However, using  $E_{r+1}$ , we can consider another intersection  $Z_2(V) := V \cap (E_1 \oplus \cdots \oplus E_{r-1} \oplus E_{r+1})$ . Then the map

$$Z := (Z_1, Z_2) \colon \mathbf{Gr}_{n,d} \to \mathbf{Gr}_e(E_1 \oplus \cdots \oplus E_r) \times \mathbf{Gr}_e(E_1 \oplus \cdots \oplus E_{r-1} \oplus E_{r+1})$$

is birational with the inverse  $Z^{-1}:(W,W')\mapsto W+W'$ .

Using the construction of X and  $\Phi$  for the tuple  $(E_1, \ldots, E_{r-1}, E_{r+1})$  instead of  $(E_1, \ldots, E_r)$ , we obtain  $Y := X(E_1, \ldots, E_{r-1}, E_{r+1})$  and the  $H_2$ -equivariant birational map

$$\Psi: Y \to \mathbf{Gr}_e(E_1 \oplus \cdots \oplus E_{r-1} \oplus E_{r+1})$$

Combining  $\Phi$ ,  $\Psi$  and Z we obtain:

Corollary 4.3. The composition  $\Omega := Z^{-1} \circ (\Phi, \Psi) \colon X \times Y \to \mathbf{Gr}_{n,d}$  is birational and  $H_2$ -invariant.

In the following we consider the cases r > 1 and r = 1 separately.

4.1. The case r > 1. For  $V \in \mathbf{Gr}_{n,d}$ , set

$$(B, \psi_2, \dots, \psi_r) := \Phi^{-1}(V \cap (E_1 \oplus \dots \oplus E_r)) \in X, \tag{2}$$

$$(C, \chi_2, \dots, \chi_{r-1}, \chi_{r+1}) := \Psi^{-1}(V \cap (E_1 \oplus \dots \oplus E_{r-1} \oplus E_{r+1})) \in Y.$$
 (3)

In order to construct a smaller normal form, fix  $(A', \varphi'_2, \dots, \varphi'_r) \in X$ ,  $A'' \in \mathbf{Gr}_e(E_1)$  and  $\chi'_{r+1} \in \mathbf{GL}(A'', E_{r+1})$  such that  $E_1 = A \oplus A'$ ,  $A'' \cap A = A'' \cap A' = \{0\}$ ,  $E_i = \varphi_i(A) \oplus \varphi'_i(A')$  for all  $i = 2, \dots, r$ , and  $\mathbb{C}^n = E_1 \oplus \dots \oplus E_r \oplus \chi'_{r+1}(A'')$ .

**Lemma 4.4.** In the notation of (2,3) define

$$S := \{ V \in \mathbf{Gr}_{n,d} : B = A', C = A'', \psi_2 = \varphi_2', \dots, \psi_r = \varphi_r', \chi_{r+1} = \chi_{r+1}' \}.$$

Then  $H_2S$  is Zariski open in  $\mathbf{Gr}_{n,d}$ .

Proof. For  $V \in \mathbf{Gr}_{n,d}$  generic,  $E_1 = A \oplus B$ . Hence there exists  $g \in \mathbf{GL}(E_1)$  with  $g|A = \mathrm{id}$  and g(B) = A'. Clearly g extends to an isomorphism from the action by  $H_2$ . Without loss of generality, B = A'. Similarly there exists an isomorphism  $g \in H_2$  such that  $g|\varphi_i(A) = \mathrm{id}$  and  $g \circ \psi_i = \varphi_i'$  on A' for all  $i = 2, \ldots, s$ . Thus we may assume that  $\psi_i = \varphi_i'$ . By Lemma 3.1, we may also assume that C = A''. Again, for V generic,  $\mathbb{C}^n = E_1 \oplus \cdots \oplus E_r \oplus \chi_{r+1}(A'')$ , where  $\chi_{r+1}(A'') \subset E_{r+1}$ . Therefore there exists an isomorphism  $g \in H_2$  such that  $g|(E_1 \oplus \cdots \oplus E_r) = \mathrm{id}$  and  $g \circ \chi_{r+1} = \chi_{r+1}'$ . This proves the lemma.

Let  $\alpha \in \mathbf{GL}(A, A')$  be the isomorphism whose graph is A'' with respect to the splitting  $E_1 = A \oplus A'$ . Define  $\beta \colon A \to A''$  by  $\beta(z) := z \oplus \alpha(z)$ . For every  $i = 2, \ldots, r$ , consider

$$\xi_i := (\varphi_i \times \varphi_i')^{-1} \circ \chi_i \circ \beta \in \mathbf{GL}(A, A \oplus A')$$

and its components  $[\xi_i]_1 \in \mathbf{GL}(A)$ ,  $[\xi_i]_2 \in \mathbf{GL}(A, A')$ . Define the rational map

$$\Xi: S \to \mathbf{GL}(A)^{2(r-2)}, \quad \Xi: V \mapsto ([\xi_2]_1, \alpha^{-1} \circ [\xi_2]_2, \dots, [\xi_{r-1}]_1, \alpha^{-1} \circ [\xi_{r-1}]_2),$$

where S is defined in Lemma 4.4.

Let  $g \in \mathbf{GL}(A)$  be arbitrary. Using the isomorphisms  $\alpha \colon A \to A'$ ,  $\varphi_i \colon A \to \varphi_i(A)$ ,  $\varphi_i' \colon A' \to \varphi_i'(A')$ ,  $i = 2, \ldots, r$ , and  $\chi_{r+1}' \circ \beta \colon A \to \chi_{r+1}'(A'')$  we can extend g canonically to an isomorphism from the action by  $H_2$ . This extension defines a canonical  $\mathbf{GL}(A)$ -action on  $\mathbb{C}^n$  and therefore on S (as defined in Lemma 4.4). We compare this action with the diagonal  $\mathbf{GL}(A)$ -action on  $\mathbf{GL}(A)^{2(r-2)}$  by conjugations:

**Lemma 4.5.** The map  $\Xi \colon \mathbf{Gr}_{n,d} \to \mathbf{GL}(A)^{2(r-2)}$  is birational and  $\mathbf{GL}(A)$ -equivariant.

*Proof.* The inverse of  $\Xi$  is given by

$$\Xi^{-1}(\lambda_2,\ldots,\lambda_{r-1},\delta_2,\ldots,\delta_{r-1}) = \Omega(x,y),$$

where  $\Omega: X \times Y \to \mathbf{Gr}_{n,d}$  is birational by Corollary 4.3 and

$$x := (A', \varphi_2', \dots, \varphi_r'), \quad y := (A'', (\varphi_2 \times \varphi_2') \circ (\lambda_2, \alpha \circ \delta_2) \circ \beta^{-1}, \dots, (\varphi_{r-1} \times \varphi_{r-1}') \circ (\lambda_{r-1}, \alpha \circ \delta_{r-1}) \circ \beta^{-1}, \chi_{r+1}').$$

The equivariance is straightforward.

In the following corollary we use the space S as defined in Lemma 4.4.

Corollary 4.4. Define  $S_3(s) := \{(V_1, \ldots, V_s) \in \mathbf{Gr}_{n,d}^s : V_{r+2} \in S\}$  and consider the diagonal  $\mathbf{GL}(A)$ -action on  $S_3$ . Then  $(S_3, \mathbf{GL}(A))$  is a normal form for  $(\mathbf{Gr}_{n,d}^s, \mathbf{GL}_n)$ .

Corollary 4.5. If d = 2e and n = (2r + 1)e, the space  $\mathbf{Gr}_{n,d}^{r+2}$  is almost  $\mathbf{GL}_n$ -homogeneous if and only if r = 2.

Now let  $V \in \mathbf{Gr}_{n,d}$  be one more subspace. Using the data fixed above we associate with V a tuple of 4r-2 linear automorphisms of A as follows. In the notation (2,3) let  $\tau_1 \in \mathbf{GL}(A,A')$  and  $\tau_2 \in \mathbf{GL}(A,A')$  be the linear isomorphisms with the graphs B and C respectively. As above, let  $\alpha \in \mathbf{GL}(A,A')$  be the isomorphism whose graph is A''. For j=1,2, set  $\tau'_j(z):=z+\tau_j(z)$  ( $\tau'_1\colon A\to B, \ \tau'_2\colon A\to C$ ) and  $\sigma_j:=\alpha^{-1}\circ\tau_j\in\mathbf{GL}(A)$ , for  $i=2,\ldots,r$ ,

$$\zeta_i := (\varphi_i \times \varphi_i')^{-1} \circ \psi_i \circ \tau_1' \in \mathbf{GL}(A, A \oplus A')$$

and for i = 2, ..., r - 1,

$$\theta_i := (\varphi_i \times \varphi_i')^{-1} \circ \chi_i \circ \tau_2' \in \mathbf{GL}(A, A \oplus A').$$

We write  $[\zeta_i]_1, [\theta_i]_1 \in \mathbf{GL}(A)$  and  $[\zeta_i]_2, [\theta_i]_2 \in \mathbf{GL}(A, A')$  for the corresponding components with respect to the splitting  $E_1 = A \oplus A'$ . Then we obtain the linear automorphisms of  $A: x_i := [\zeta_i]_1, y_i := \alpha^{-1} \circ [\zeta_i]_2$  for  $i = 2, \ldots, r$  and  $z_i := [\theta_i]_1, t_i := \alpha^{-1} \circ [\theta_i]_2$  for  $i = 2, \ldots, r - 1$ .

In order to deal with the remainder term  $\chi_{r+1}$  we define  $\varphi_{r+1} \in \mathbf{GL}(A, \mathbb{C}^n)$  by

$$\varphi_{r+1}(z) := z \oplus \varphi_2(z) \oplus \cdots \oplus \varphi_r(z).$$

It follows from the construction that  $\varphi_{r+1}(A) \in \mathbf{Gr}_e(E_{r+1})$  and  $E_{r+1} = \varphi_{r+1}(A) \oplus \chi'_{r+1}(A'')$ . Using this splitting we define the remainder isomorphisms

$$\theta_r := (\varphi_{r+1} \times \chi'_{r+1})^{-1} \circ \chi_{r+1} \circ \tau'_2 \in \mathbf{GL}(A, A \oplus A')$$

and 
$$z_r := [\theta_r]_1, t_r := \alpha^{-1} \circ [\theta_r]_2 \in \mathbf{GL}(A)$$
.

Lemma 4.6. Let  $\Theta \colon \mathbf{Gr}_{n,d} \to \mathbf{GL}(A)^{4r-2}$  be given by

$$\Theta: V \mapsto (\sigma_1, \sigma_2, x_2, y_2, z_2, t_2, \dots, x_r, y_r, z_r, t_r).$$

Then  $\Theta$  is birational and GL(A)-equivariant.

*Proof.* In the above notation the inverse  $\Theta^{-1}$  can be calculated as follows:

$$\tau'_j := \alpha \circ \sigma_j + \mathrm{id}, \quad B := \tau'_1(A), \quad C := \tau'_2(A),$$
  
$$\psi_i = (\varphi_i \times \varphi'_i) \circ (x_i, \alpha \circ y_i) \circ (\tau'_1)^{-1}$$

for  $i = 2, \ldots, r$ ,

$$\chi_i = (\varphi_i \times \varphi_i') \circ (z_i, \alpha \circ t_i) \circ (\tau_2')^{-1}$$

for i = 2, ..., r - 2,

$$\chi_{r+1} = (\varphi_{r+1} \times \chi'_{r+1}) \circ (z_r, \alpha \circ t_r) \circ (\tau'_2)^{-1}$$

and finally

$$V = \Omega((B, \psi_2, \dots, \psi_r), (C, \chi_2, \dots, \chi_{r-1}, \chi_{r+1})).$$

The equivariance is straightforward.

Corollary 4.6. If  $s \ge r + 2$ , the space  $(S_3, \mathbf{GL}(A))$  is isomorphic to  $(\mathbf{GL}_e^{2r-4+(s-r-2)(4r-2)}, \mathbf{GL}_e)$ .

*Proof.* The required birational isomorphism is given by

$$(V_1, \dots, V_s) \mapsto (\Xi(V_{r+2}), \Theta(V_{r+3}), \dots, \Theta(V_s)), \tag{4}$$

where  $\Xi$  is birational by Lemma 4.5 and  $\Theta$  is birational by Lemma 4.6.  $\square$ 

This implies the second part of Theorem 1.2 in the case r > 1.

4.2. The case r=1. In this case we have n=3e. Recall that we fixed  $E_1, E_2 \in \mathbf{Gr}_{n,d}$  in general position such that dim A=e, where  $A:=E_1\cap E_2$ . Choose another subspace  $E_3\in \mathbf{Gr}_{n,d}$  such that dim  $A_1=\dim A_2=e$ , where  $A_j:=E_j\cap E_3$  for j=1,2. Recall that we defined  $H_2\subset \mathbf{GL}_n$  to be the stabiliser of both  $E_1$  and  $E_2$ . Denote by  $H_3\subset H_2$  the stabiliser of  $E_3$ . Then  $H_3=\mathbf{GL}(A)\times\mathbf{GL}(A_1)\times\mathbf{GL}(A_2)$  with respect to the splitting  $\mathbb{C}^n=A\oplus A_1\oplus A_2$ . The following is straightforward.

**Lemma 4.7.** Suppose that  $s \geq 3$  and define  $S_3 = S_3(s) := \{(V_1, \ldots, V_s) \in S_2 : V_3 = E_3\}$ . Then  $(S_3, H_3)$  is a normal form for  $(\mathbf{Gr}_{n,d}^s, \mathbf{GL}_n)$ .

Now we choose  $B_j \in \mathbf{Gr}_e(E_j)$  such that  $B_j \cap A = B_j \cap A_j = \{0\}$  for j = 1, 2. Then each  $B_j$  can be seen as the graph of an isomorphism  $\varphi_j \in \mathbf{GL}(A, A_j)$ . On the other hand, for  $V \in \mathbf{Gr}_{n,d}$  generic, the subspaces  $C_j := V \cap E_j$  are graphs of isomorphisms  $\psi_j \in \mathbf{GL}(A, A_j)$ . Define

$$g=(\mathsf{id},\psi_1\circ\varphi_1^{-1},\psi_2\circ\varphi_2^{-1})\in\mathbf{GL}(A\oplus A_1\oplus A_2).$$

Then  $g \in H_3$  and  $g(B_j) = C_j$  for i = 1, 2, in particular,  $V = B_1 + B_2$ . Together with Lemma 3.3 this proves the following:

**Lemma 4.8.** Suppose that  $s \geq 4$  and define

$$S_4 = S_4(s) := \{(V_1, \dots, V_s) \in S_3 : V_4 \cap E_1 = B_1, V_4 \cap E_2 = B_2\}$$

and  $H_4 := \mathbf{GL}(A)$ . Define the  $H_4$ -action on  $S_4$  via the homomorphism

$$H_4 \to H_3$$
,  $g \mapsto (g, \varphi_1 \circ g \circ \varphi_1^{-1}, \varphi_2 \circ g \circ \varphi_2^{-1})$ .

Then  $(S_4, H_4)$  is a normal form for  $(\mathbf{Gr}_{n,d}^s, \mathbf{GL}_n)$ .

As a special case of Corollary 3.3, we obtain:

**Lemma 4.9.** The space  $\mathbf{Gr}_{3e,2e}^s$  is almost  $\mathbf{GL}_{3e}$ -homogeneous if and only if  $s \leq 4$ . If  $s \geq 5$ , there exists a normal form which is isomorphic to  $(\mathbf{GL}_e^{2(s-4)}, \mathbf{GL}_e)$ , where  $\mathbf{GL}_e$  acts diagonally by conjugations.

This implies the second part of Theorem 1.2 in the case r = 1.

## 5. Computation of rational invariants.

5.1. The case n = rd. In order to compute the rational  $\mathbf{GL}_{rd}$ -invariants of  $\mathbf{Gr}^s_{rd,d}$  we represent the elements of  $\mathbf{Gr}^s_{rd,d}$  by the equivalence classes of  $rd \times ds$  matrices  $M \in \mathbb{C}^{rd \times sd}$ , where the equivalence is taken under the right multiplication by  $\mathbf{GL}^s_d$ . Then the diagonal  $\mathbf{GL}_{rd}$ -action on  $\mathbf{Gr}^s_{rd,d}$  corresponds to the left multiplication on  $\mathbb{C}^{rd \times sd}$ . We start with  $2d \times 2d$  matrices. Define

$$D \colon \mathbf{GL}_{2d} \to \mathbf{GL}_d, \quad D \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} := A_{11} A_{21}^{-1} A_{22} A_{12}^{-1}.$$
 (5)

Then D is a rational map that is invariant under the right multiplication by  $\mathbf{GL}_d \times \mathbf{GL}_d$  and the left multiplication by  $\{e\} \times \mathbf{GL}_d$ , where  $e \in \mathbf{GL}_d$  is the unit. Moreover, D is equivariant with respect to the left multiplication by  $\mathbf{GL}_d \times \{e\}$ .

More generally, let  $M \in \mathbb{C}^{rd \times sd}$  be a rectangular matrix, where r and s are arbitrary positive integers. We split M into  $d \times d$  blocks  $A_{ij}$ ,  $i = 1, \ldots, r$ ,  $j = 1, \ldots, s$ . Then for every  $i = 2, \ldots, r$ , and  $j = 2, \ldots, s$ , define

$$D_{ij} : \mathbb{C}^{rd \times sd} \to \mathbf{GL}_d, \quad D_{ij}(M) := D \begin{pmatrix} A_{11} & A_{1j} \\ A_{i1} & A_{ij} \end{pmatrix}.$$

Similarly to D, each  $D_{ij}$  is a rational map that is invariant under the right multiplication by  $\mathbf{GL}_d^s$  and under the left multiplication by  $\{e\} \times \mathbf{GL}_d^{r-1}$ .

Finally, we construct rational maps that are invariant under the left multiplication by the larger group  $\mathbf{GL}_{rd}$ . For this, we assume s > r+1, otherwise the left  $\mathbf{GL}_{rd}$ -multiplication is almost homogeneous (see Corollary 3.3) and, hence, all rational invariants are constant. Then every matrix  $M \in \mathbb{C}^{rd \times sd}$  from a Zariski open subset can be split into two blocks:

$$M = (AB), \quad A \in \mathbf{GL}_{rd}, \quad B \in \mathbb{C}^{rd \times (s-r)d}.$$

Define

$$\varphi(M) := A^{-1}B \in \mathbb{C}^{rd \times (s-r)d}$$

and

$$G_{ij}(M) := D_{ij}(\varphi(M)), \quad i = 2, \dots, r, \quad j = 2, \dots, s - r.$$

In particular, for d = 1, r = 2 and s = 4, we obtain the classical double ratio:

$$G_{22}\begin{pmatrix}1&1&1&1\\z_1&z_2&z_3&z_4\end{pmatrix}=D\begin{pmatrix}z_2-z_3&z_2-z_4\\z_1-z_3&z_1-z_4\end{pmatrix}=\frac{z_2-z_3}{z_1-z_3}:\frac{z_2-z_4}{z_1-z_4}.$$

By the invariance of  $D_{ij}$ , every  $G_{ij}$  is invariant under the left multiplication by  $\mathbf{GL}_{rd}$  and under the right multiplication by  $\{e\} \times \mathbf{GL}_d^{s-1}$ . Moreover, due to the construction,  $G_{ij}$  is equivariant with respect to the subgroup  $\mathbf{GL}_d \times \{e\} \subset$  $\mathbf{GL}_d \times \mathbf{GL}_d^{s-1}$ , where we take the right action on  $\mathbb{C}^{rd \times sd}$  and the conjugation on the image space  $\mathbf{GL}_d$ . Therefore we obtain the following rational  $\mathbf{GL}_{rd} \times \mathbf{GL}_d^s$ -invariants:

$$I_{\alpha\beta} := \mathbf{Tr}(G_{\alpha_1\beta_1} \cdot \ldots \cdot G_{\alpha_k\beta_k}) = \mathbf{Tr}((D_{\alpha_1\beta_1} \circ \varphi) \cdot \ldots \cdot (D_{\alpha_k\beta_k} \circ \varphi)), \tag{6}$$

where  $\alpha \in \{2, ..., r\}^k$  and  $\beta \in \{2, ..., s - r\}^k$  are arbitrary multi-indices.

Using a result of Procesi [P76] we show that the invariant field is actually generated by these traces of monomials.

**Theorem 5.1.** Let d, r, s be arbitrary positive integers such that s > r + 1. Then the field of rational  $\mathbf{GL}_{rd} \times \mathbf{GL}_d^s$ -invariants of  $rd \times sd$  matrices is generated by the functions  $I_{\alpha\beta}$ , where  $\alpha \in \{2, \ldots, r\}^k$  and  $\beta \in \{2, \ldots, s - r\}^k$  and  $k < 2^d$ .

*Proof.* Set n := rd as above. We write the spaces  $E_1, \ldots, E_{r+1}, V_{r+2}, \ldots, V_s \in \mathbf{Gr}_{n,d}$  as in Lemma 3.3 in the form of an  $rd \times sd$  matrix with  $d \times d$  blocks as follow:

$$M := \begin{pmatrix} E & 0 & \cdots & 0 & E & E & \cdots & E \\ 0 & E & \cdots & 0 & E & V_{2,r+2} & \cdots & V_{2,s} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & E & E & V_{r,r+2} & \cdots & V_{r,s} \end{pmatrix}, \tag{7}$$

where E denotes the identity matrix of the size  $d \times d$ .

Let  $\widetilde{S_2} \subset \mathbb{C}^{rd \times sd}$  be subspace of all matrices of the form (7). Clearly  $E_1, \ldots, E_{r+1}$  fulfil the assumptions of Lemma 3.3. Then it follows from Lemma 3.3, that  $(\widetilde{S_2}, \mathbf{GL}_d)$  is a normal form for  $(\mathbb{C}^{rd \times sd}, \mathbf{GL}_{rd} \times \mathbf{GL}_d^s)$ , where  $\mathbf{GL}_d$  acts on  $\widetilde{S_2}$  as the diagonal subgroup of  $\mathbf{GL}_{rd} \times \mathbf{GL}_d^s$ , i.e. each  $d \times d$  block is conjugated by the same matrix.

By the definition of a normal form, it is sufficient to prove that the invariant field of  $(\widetilde{S}_2, \mathbf{GL}_d)$  is generated by the restrictions of  $I_{\alpha\beta}$ 's. Let M be given by (7). By the obvious calculations,

$$\varphi(M) = \begin{pmatrix} E & E & \cdots & E \\ E & V_{2,r+2} & \cdots & V_{2,s} \\ \vdots & \vdots & \ddots & \vdots \\ E & V_{r,r+2} & \cdots & V_{r,s} \end{pmatrix}$$
(8)

and hence

$$I_{\alpha\beta}(M) = \mathbf{Tr}(V_{\alpha_1, r+\beta_1} \cdot \dots \cdot V_{\alpha_k, r+\beta_k}). \tag{9}$$

By a theorem of Procesi [P76], the polynomial invariants of (r-1)(s-r-1)-tuples  $(V_{ij})$  with respect to the diagonal  $\mathbf{GL}_d$ -conjugations are generated by the monomials of the form (9) with  $k < 2^d$ . Since all points of  $(\widetilde{S_2}, \mathbf{GL}_d)$  are semi-stable (see [MFK93]), the categorical quotient  $\widetilde{S_2}//\mathbf{GL}_d$  exists and is given by these monomials. Then the rational invariants on  $(\widetilde{S_2}, \mathbf{GL}_d)$  are pullbacks of rational functions on  $\widetilde{S_2}//\mathbf{GL}_d$  and the proof is finished.

5.2. The case  $n=3e,\ d=2e$ . In this case the elements of  $\mathbf{Gr}_{n,d}^s$  are represented by the equivalence classes of matrices  $M\in\mathbb{C}^{3e\times 2se}$  with  $3e\times 2e$  blocks  $M_1,\ldots,M_s$ . As above we are looking for rational invariants with respect to left multiplications by  $\mathbf{GL}_{3e}$  and right multiplications by  $\mathbf{GL}_d^s$ . We start with the case of 3 blocks. Define the rational map

$$\varphi \colon \mathbb{C}^{3e \times 6e} \to \mathbb{C}^{3e \times 6e}$$
,

$$\varphi \begin{pmatrix} A_1 & A_2 & A_3 \\ B_1 & B_2 & B_3 \end{pmatrix} := \begin{pmatrix} E' & E' & E' \\ B_1 A_1^{-1} & B_2 A_2^{-1} & B_3 A_3^{-1} \end{pmatrix} \\
=: \begin{pmatrix} E & 0 & E & 0 & E & 0 \\ 0 & E & 0 & E & 0 & E \\ c_1 & d_1 & c_2 & d_2 & c_3 & d_3 \end{pmatrix}, (10)$$

where  $A_1, A_2, A_3 \in \mathbf{GL}_d$ ,  $B_1, B_2, B_3 \in \mathbb{C}^{e \times 2e}$  and  $E' \in \mathbf{GL}_{2e}$ ,  $E \in \mathbf{GL}_e$  denote the identity matrices. We see the corresponding subspaces  $E_1, E_2, E_3 \in \mathbf{Gr}_{n,d}$  as graphs of linear maps given by the matrices  $(c_1 d_1), (c_2 d_2), (c_3 d_3)$ . Our first goal will be to compute the intersections  $A := E_1 \cap E_2$  and  $A_j := E_j \cap E_3, j = 1, 2$ . For this, we set

$$\begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} := \begin{pmatrix} \begin{pmatrix} c_1 & d_1 \\ c_2 & d_2 \end{pmatrix}^{-1}, \begin{pmatrix} c_3 & d_3 \\ c_1 & d_1 \end{pmatrix}^{-1}, \begin{pmatrix} c_2 & d_2 \\ c_3 & d_3 \end{pmatrix}^{-1} \end{pmatrix} \begin{pmatrix} E \\ E \end{pmatrix} \in \mathbb{C}^{e \times 3e}.$$

Then  $x_1c_1 + y_1d_1 = x_1c_2 + y_1d_2 = E$  and therefore the  $3e \times e$  matrix

$$\begin{pmatrix} x_1 \\ y_1 \\ E \end{pmatrix}$$

represents the intersection  $E_1 \cap E_2$ . Similarly the  $3e \times e$  matrices

$$\begin{pmatrix} x_2 \\ y_2 \\ E \end{pmatrix}$$
 and  $\begin{pmatrix} x_3 \\ y_3 \\ E \end{pmatrix}$ 

represent the intersections  $E_3 \cap E_1$  and  $E_2 \cap E_3$  respectively. Equivalently, the 3-tuple  $(E_1, E_2, E_3)$  can be represented by the matrix

$$\begin{pmatrix} x_1 & x_2 & x_3 & x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 & y_1 & y_2 & y_3 \\ E & E & E & E & E \end{pmatrix}$$
 (11)

Now take the general matrix  $M=(M_1...M_s)\in\mathbb{C}^{3e\times 2es}$ . Following the construction of §4.2 we bring  $A, A_1$  and  $A_2$ , i.e. the matrix (11) to a normal form. For this, consider the square matrix

$$H(M) := \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ E & E & E \end{pmatrix}$$

and multiply M by  $H(M)^{-1}$ :

$$H(M)^{-1}M := \begin{pmatrix} E & 0 & 0 & E & 0 & 0 & C_{14} & D_{14} & \cdots & C_{1s} & D_{1s} \\ 0 & E & 0 & 0 & E & 0 & C_{24} & D_{24} & \cdots & C_{2s} & D_{2s} \\ 0 & 0 & E & 0 & 0 & E & C_{34} & D_{34} & \cdots & C_{3s} & D_{3s} \end{pmatrix}$$
(12)

Next we normalize the  $3e \times 2e$  blocks:

$$\begin{pmatrix} C_{1i} & D_{1i} \\ C_{2i} & D_{2i} \\ C_{3i} & D_{3i} \end{pmatrix} \begin{pmatrix} C_{1i} & D_{1i} \\ C_{2i} & D_{2i} \end{pmatrix}^{-1} = \begin{pmatrix} E & 0 \\ 0 & E \\ \alpha_{2i-1} & \alpha_{2i} \end{pmatrix}$$

and define

$$\sigma_{2i-1} := \alpha_{2i-1}\alpha_7^{-1}, \quad \sigma_{2i} := \alpha_{2i}\alpha_8^{-1}, \quad i = 5, \dots, s.$$

Comparing this construction with §4.2 we conclude that the matrices  $\sigma_j$ ,  $j = 9, \ldots, 2s$ , represent exactly the 2(s-4) matrices in the isomorphic normal form  $(\mathbf{GL}_e^{2(s-4)}, \mathbf{GL}_e)$ . Using the theorem of Procesi [P76] and the arguments similar to §5.1 we obtain:

**Theorem 5.2.** Let e and s be arbitrary positive integers such that s > 4. Then the field of rational  $\mathbf{GL}_{3e} \times \mathbf{GL}_{2e}^s$ -invariants of  $3e \times 2es$  matrices is generated by the functions

$$J_{\alpha} := \mathbf{Tr}(\sigma_{\alpha_1} \cdot \ldots \cdot \sigma_{\alpha_k}),$$

where  $\alpha \in \{9, \dots, 2s\}^k$  and  $k < 2^e$ .

5.3. The case n=(2r+1)e and d=2e. Here an element  $(V_1,\ldots,V_s)\in\mathbf{Gr}_{n,d}^s$  is represented by a matrix  $M\in\mathbb{C}^{(2r+1)e\times 2se}$  with s blocks  $M_1,\ldots,M_s$  of the size  $(2r+1)e\times 2e$ . Again, the  $\mathbf{GL}_n$ -invariants on  $\mathbf{Gr}_{n,d}^s$  correspond to  $\mathbf{GL}_n\times\mathbf{GL}_{2e}^s$ -invariants on  $\mathbb{C}^{(2r+1)e\times 2se}$ . Similarly to the previous paragraph, we start with a special case of a  $(2r+1)e\times (2r+2)e$  matrix and compute representatives of the intersections  $E_{r+1}\cap(E_1\oplus\cdots\oplus E_r)$  and  $E_{r+2}\cap(E_1\oplus\cdots\oplus E_r)$ 

$$\varphi\begin{pmatrix} A_1 & \dots & A_s \\ B_1 & \dots & B_s \end{pmatrix} := \begin{pmatrix} E' & \dots & E' \\ B_1 A_1^{-1} & \dots & B_s A_s^{-1} \end{pmatrix} =: \begin{pmatrix} E' & \dots & E' \\ C_1 & \dots & C_s \end{pmatrix}. \tag{13}$$

In order to compute the first intersection  $E_{r+1} \cap (E_1 \oplus \cdots \oplus E_r)$  we consider the corresponding system of linear equations:

$$(E' \dots E') \begin{pmatrix} X_1 \\ \vdots \\ X_r \end{pmatrix} = E' X_{r+1}, \quad (C_1 \dots C_r) \begin{pmatrix} X_1 \\ \vdots \\ X_r \end{pmatrix} = C_{r+1} X_{r+1},$$

where each  $X_i$  is a  $2e \times e$  block. Solving from the first equation  $X_{r+1} = X_1 + \ldots + X_r$  and substituting this in the second we obtain:

$$((C_1 - C_{r+1}) \dots (C_r - C_{r+1})) \begin{pmatrix} X_1 \\ \vdots \\ X_r \end{pmatrix} = 0.$$

This is a system of  $(2r + 1)e \times e$  equations with  $2re \times e$  variables. Hence, for  $C_1, \ldots, C_{r+1}$  in general position, it has a solution which can be represented by rational  $2e \times e$  block functions

$$X_i = X_i(M), \quad i = 1, \dots, r.$$

Comparing this with the construction of §4 we see that the blocks

$$\begin{pmatrix} E' \\ C_i \end{pmatrix} (X_i), \quad i = 1, \dots, r,$$

represent the subspaces  $A, \varphi_2(A), \ldots, \varphi_r(A)$ . For our normalization (Lemma 4.4) we also need the subspaces  $A', \varphi_2'(A'), \ldots, \varphi_r'(A')$  and  $\chi_{r+1}'$  which come from the other intersections. For them we can also solve the corresponding linear systems and obtain rational  $2e \times e$  block functions

$$Y_i(M), i = 1, ..., r, \text{ and } Z_{r+1}(M),$$

such that the blocks

$$\begin{pmatrix} E' \\ C_i \end{pmatrix} (Y_i(M)), \quad i = 1, \dots, r, \quad \text{ and } \quad \begin{pmatrix} E' \\ C_{r+1} \end{pmatrix} (Z_{r+1}(M)),$$

represent  $A', \varphi_2'(A'), \ldots, \varphi_r'(A')$  and  $\chi_{r+1}'$  respectively. As above we put these blocks together in the  $n \times n$  matrix:

$$H(M) := \left( \begin{pmatrix} E' & \cdots & E' \\ C_1 & \cdots & C_r \end{pmatrix} \begin{pmatrix} X_1 & Y_1 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & X_r & Y_r \end{pmatrix}, \begin{pmatrix} E' \\ C_{r+1} \end{pmatrix} (Z_{r+1}(M)) \right).$$

The  $n \times e$  columns of H(M) are exactly the representatives of

$$A, A', \varphi(A), \varphi'(A'), \ldots, \varphi_r(A), \varphi'_r(A'), \chi'_{r+1}(A'')$$

in this order. If M is in general position, H(M) is invertible. As above, consider the matrix  $H(M)^{-1}M=$ 

Furthermore, the property  $E_{r+1} = \Phi(A, \varphi_2, \dots, \varphi_r) + \chi'_{r+1}(A'')$  implies

$$a_2 = a_4 = \dots = a_{2r} = a_{2r+1} = 0.$$

Using the property  $W' := \Phi(A', \varphi'_2, \dots, \varphi'_r) \in E_{r+2}$  we may assume that W' is represented by the block

$$\begin{pmatrix} b_{1,r+2} \\ \vdots \\ b_{2r+1,r+2} \end{pmatrix}$$

and therefore

$$b_{1,r+2} = b_{3,r+2} = \dots = b_{2r+1,r+2} = 0.$$

Furthermore, the matrix components of the map (4) can be calculated directly:

$$Z(M) = \left(D\begin{pmatrix} a_1 & c_{1,r+2} \\ a_3 & c_{3,r+2} \end{pmatrix}, D\begin{pmatrix} b_{2,r+2} & c_{2,r+2} \\ b_{4,r+2} & c_{4,r+2} \end{pmatrix}, \cdots, D\begin{pmatrix} a_1 & c_{1,r+2} \\ a_{2r-3} & c_{2r-3,r+2} \end{pmatrix}, D\begin{pmatrix} b_{2,r+2} & c_{2,r+2} \\ b_{2r-2,r+2} & c_{2r-2,r+2} \end{pmatrix}\right)$$
(14)

and

$$\Theta_{i}(M) = \left(D\begin{pmatrix} a_{1} & b_{1,i} \\ a_{3} & c_{3,i} \end{pmatrix}, D\begin{pmatrix} b_{2,r+2} & c_{2,i} \\ b_{4,r+2} & c_{4,i} \end{pmatrix}, \cdots, D\begin{pmatrix} a_{1} & b_{1,i} \\ a_{2r-1} & b_{2r-1,i} \end{pmatrix}, 
D\begin{pmatrix} b_{2,r+2} & c_{2,r+2} \\ b_{2r,r+2} & c_{2r,r+2} \end{pmatrix}, (c_{2r+1,r+2}^{-1}b_{2r+1,i}), (c_{2r+1,r+2}^{-1}c_{2r+1,i}) \right), \quad i = r+3, \dots, s,$$
(15)

where D is the generalized double ratio defined by (5).

Comparing this with the proof of Corollary 4.6 and applying the theorem of Procesi [P76] we obtain:

**Theorem 5.3.** Let e, r and s be arbitrary positive integers such that  $s \ge r + 2$ . Then the field of rational  $\mathbf{GL}_{(2r+1)e} \times \mathbf{GL}_{2e}^s$ -invariants of  $(2r+1)e \times 2es$  matrices is generated by the functions

$$\mathbf{Tr}(\sigma_1 \cdot \ldots \cdot \sigma_k),$$

where each  $\sigma_l$  is either a component of the map Z in (14) or of one of the maps  $\Theta_{r+3}, \ldots, \Theta_s$  in (15) and  $k < 2^e$ .

#### References

- [B91] L. le Bruyn, Centers of generic division algebras, the rationality problem 1965-1990, Isr. J. Math. 76 (1991), 97–111.
- [BS88] L. le Bruyn and A. Schofield, Rational invariants of quivers and the ring of matrix invariants, NATO ASI Ser., Antwerp/Belg., Ser. C 233 (1988), 21–29.
- [D95] I.V. Dolgachev, Rationality of fields of invariants, in Bloch, S. J. (ed.), Algebraic geometry, Proc. Symp. Pure Math. 46 (1987), 3–16.
- [DB96] I.V. Dolgachev and H.U. Boden, Unpublished talk at Kazimierz, 1996.
- [F79] E. Formanek, The center of the ring of  $3 \times 3$  generic matrices, Linear and Multilinear Algebra 7 (1979), 203–212.
- [F80] E. Formanek, The center of the ring of  $4 \times 4$  generic matrices, J. Algebra 62 (1980), 304-319.
- [GP71] I.M. Gelfand and V.A. Ponomarev, Quadruples of subspaces of a finite-dimensional vector space, Sov. Math., Dokl. 12 (1971), 535–539; translation from Dokl. Akad. Nauk SSSR 197 (1971), 762–765.
- [GP72] I.M. Gelfand and V.A. Ponomarev, Problems of linear algebra and classification of quadruples of subspaces in a finite-dimensional vector space, in Hilbert Space Operators Operator Algebras, Colloquia math. Soc. Janos Bolyai 5 (1972), 163–237.
- [M98] G. Megyesi, The rationality of configuration spaces of lines in  $\mathbb{P}^3$ , preprint, 1998.
- [MFK93] D. Mumford, J. Fogarty, and F. Kirwan, Geometric Invariant Theory, Ergebnisse der Mathematik und ihrer Grenzgebiete, Berlin Springer-Verlag, 1993.

- [P67] C. Procesi, Non-commutative affine rings, Atti Accad. Naz. Lincei 8 (1967), 239–255.
- [P76] C. Procesi, The invariant theory of  $n \times n$  matrices, Adv. Math. 19 (1976), 306–381.
- [R56] M. Rosenlicht, Some basic theorems on algebraic groups, Amer. J. Math. 78 (1956), 401–443.
- [Z98] D. Zaitsev, Domains of polyhedral type and boundary extensions of biholomorphisms, to appear in Indiana Univ. Math. J. 1998.

Mathematisches Institut, Eberhard-Karls-Universität Tübingen, 72076 Tübingen, Germany, E-mail address: dmitri.zaitsev@uni-tuebingen.de