

Gerbes of chiral differential operators

Vassily Gorbounov, Fyodor Malikov, Vadim Schechtman

1. In this note we compute the cohomological obstruction to the existence of certain sheaves of vertex algebras on smooth varieties. These sheaves have been introduced and studied in the previous work by A. Vaintrob and two of the authors, cf. [MSV] and [MS1]. Hopefully our result clarifies to some extent the constructions of *op. cit.*

Recall that in [MSV] one discussed two kinds of sheaves on smooth complex algebraic (or analytic) varieties X . First, one defined the sheaf of conformal vertex superalgebras Ω_X^{ch} , called *chiral de Rham algebra*. These sheaves are canonically defined for an arbitrary X . Second, for some varieties X one can define a purely even counterpart of Ω_X^{ch} , a sheaf of graded vertex algebras \mathcal{O}_X^{ch} , called a *chiral structure sheaf*, cf. *op. cit.*, §5. For example, one can define \mathcal{O}_X^{ch} for curves, and for flag spaces G/B . For an arbitrary X , there arises certain cohomological obstruction to the existence of \mathcal{O}_X^{ch} . The infinitesimal incarnation of this obstruction is calculated in *op. cit.*, §5, A.

In the present note we compute explicitly this obstruction, which turns out to admit a very simple expression in terms of characteristic classes of X , cf. Theorem 3 below. From a different viewpoint, this theorem may be regarded as a geometric interpretation of the second component of Chern character. In the last nos. 7 — 9 we give a generalization of Theorem 3, cf. Theorem 8.1.

In particular, Theorem 3, provides a geometric criterion for a Calabi-Yau manifold to be a $BU\langle 6 \rangle$ -manifold: those are precisely the manifolds which admit the above mentioned sheaf \mathcal{O}_X^{ch} .

First, a remark about terminology and notation. As was noted in [MSV], (cf. the last paragraph of §5), the sheaf \mathcal{O}_X^{ch} may be regarded as a chiral counterpart of the sheaf \mathcal{D}_X of differential operators on X . By that reason we prefer in this note to change its notation to \mathcal{D}_X^{ch} and call it a *sheaf of chiral differential operators* on X . From this point of view, the sheaf Ω_X^{ch} is a chiral counterpart of the sheaf \mathcal{D}_{Ω_X} of differential operators on the de Rham algebra Ω_X of differential forms, so it would probably deserve the notation $\mathcal{D}_{\Omega_X}^{ch}$, cf. [MS2].

For a graded vertex algebra V , V_i will denote the component of conformal weight i .

In this note we will live in the analytical category.

We are deeply grateful to A. Beilinson for very enlightening discussions. Our special gratitude goes to him for an important correction in the previous version of the note.

The work was done while the first and the third author visited the University of Glasgow and Max-Planck-Institut für Mathematik; we thank these institutions for the hospitality.

2. Fix an integer $N \geq 1$. Let $O = \mathbb{C}[b^1, \dots, b^N]$ be the polynomial algebra. Following [MSV], define the vertex algebra D as the free bosonic vertex algebra generated by the fields $b^i(z)$ of conformal weight 0 and $a^i(z)$ of conformal weight 1 ($i = 1, \dots, N$), with the OPE

$$a^i(z)b^j(w) \sim \frac{\delta_{ij}}{z-w}, \quad (2.1)$$

the other products being trivial. The algebra D is called *the algebra of (algebraic) chiral differential operators on \mathbb{C}^N* .

The conformal weight zero component D_0 is equal to $\mathbb{C}[b_0^1, \dots, b_0^N]$, and will be identified with O . The component D_1 is a free O -module with the base b_{-1}^i, a_{-1}^i . The submodule $\Omega := \oplus O \cdot b_{-1}^i \subset V_1$ will be identified with the O -module of 1-differentials $\Omega_O^1 = \oplus O \cdot db^i$, by identifying b_{-1}^i with db^i . The submodule $T := \oplus O \cdot a_{-1}^i$ will be identified with the O -module of vector fields $\oplus O \cdot \partial_i$ where $\partial_i := \partial/\partial b^i$, by identifying a_{-1}^i with ∂_i .

Let X be an N -dimensional smooth complex analytic variety. Let $U \subset X$ be an open subset given together with a coordinate system, i.e. an open embedding $b = (b^1, \dots, b^N) : U \rightarrow \mathbb{C}^N$. Let $\mathcal{O}(U)$ denote the algebra of holomorphic functions on U . The map b induces an algebra homomorphism $O \rightarrow \mathcal{O}(U)$. We set $D(U; b) := \mathcal{O}(U) \otimes_O D$. According to [MSV], 6.9, $D(U; b)$ admits a canonical structure of a conformal vertex algebra. It is called the algebra of (holomorphic) chiral differential operators on U , associated with the coordinate system b .

If $V \subset U$ is an open subset, the map b induces a coordinate system on V , and we get an algebra $D(V; b)$. When V varies, we get a sheaf of conformal vertex algebras $\mathcal{D}_{(U; b)}$ on U — the sheaf of holomorphic chiral differential operators on U , associated with the coordinate system b .

The component $\mathcal{D}_{(U; b)0}$ is identified with the sheaf of holomorphic functions \mathcal{O}_U , and the component $\mathcal{D}_{(U; b)1}$ is identified with the direct sum of the sheaf Ω_U^1 of holomorphic 1-differentials and the sheaf \mathcal{T}_U of holomorphic vector fields.

Let us define a two-step filtration on $\mathcal{D}_{(U; b)1}$ by

$$0 = F^2 \mathcal{D}_{(U; b)1} \subset \Omega_U^1 = F^1 \mathcal{D}_{(U; b)1} \subset \mathcal{D}_{(U; b)1} = F^0 \mathcal{D}_{(U; b)1} \quad (2.2)$$

Obviously, $F^0/F^1 = \mathcal{T}_U$. The associated graded space

$$gr_F \mathcal{D}_{(U; b)1} = \Omega_U^1 \oplus \mathcal{T}_U \quad (2.3)$$

does not depend on the choice of a coordinate system b , cf. [MSV], 6.2.

Let us define a groupoid $\mathfrak{D}^{ch}(U)$ as follows. The objects of $\mathfrak{D}^{ch}(U)$ are coordinate systems b on U . Given two coordinate systems $b, 'b$, the set of morphisms $Hom_{\mathfrak{D}^{ch}(U)}(b, 'b)$ consists of isomorphisms of sheaves of graded vertex algebras $\mathcal{D}_{(U; b)} \xrightarrow{\sim} \mathcal{D}_{(U; 'b)}$ which induce the identity on the conformal weight zero components, and on the associated graded (with respect to the filtration (2.2)) of the conformal weight one components. (Note that any morphism of graded vertex algebras necessarily respects the filtration (2.2)). This groupoid will be called

the *groupoid of (holomorphic) chiral differential operators* on U . When U varies, we get a stack (*champ*) of groupoids, i.e. a *gerbe* \mathfrak{D}_X^{ch} on X , called the *gerbe of (holomorphic) chiral differential operators* on X .

Concerning the J.Giraud's language of gerbes, see for example Section 5 of [D]. Now we can formulate our main result.

3. Theorem. *The lien of the gerbe \mathfrak{D}_X^{ch} is canonically isomorphic to the sheaf of closed holomorphic 2-differentials $\Omega_X^{2,cl}$.*

The equivalence class $c(\mathfrak{D}_X^{ch}) \in H^2(X; \Omega_X^{2,cl})$ is equal to

$$c(\mathfrak{D}_X^{ch}) = 2ch_2(\mathcal{T}_X) \quad (3.1)$$

Here \mathcal{T}_X denotes the tangent bundle; for a vector bundle E we set

$$ch_2(E) = \frac{1}{2}c_1^2(E) - c_2(E) \quad (3.2)$$

where $c_i(E) \in H^i(X; \Omega_X^{i,cl})$ are the Hodge Chern classes.

Recall that the set of equivalence classes of gerbes on X bounded by a given abelian sheaf \mathcal{F} is identified with the group $H^2(X; \mathcal{F})$. Namely, if \mathfrak{G} is such a gerbe, choose an open covering $\mathcal{U} = \{U_i\}$ of X such that for each U_i the groupoid $\mathfrak{G}(U_i)$ is nonempty and connected. Choose objects $\mathcal{G}_i \in \mathfrak{G}(U_i)$. On the pairwise intersections there exist isomorphisms $\phi_{ij} : \mathcal{G}_i|_{U_{ij}} \xrightarrow{\sim} \mathcal{G}_j|_{U_{ij}}$. The "discrepancies" $c_{ijk} := \phi_{ij}\phi_{jk}\phi_{ik}^{-1}$ (on triple intersections) define a Čech 2-cocycle in $Z^2(\mathcal{U}; \mathcal{F})$ which represents the class $c(\mathfrak{G})$.

This theorem means that a "sheaf of chiral differential operators" is not a uniquely defined object associated with X , but an object of a canonically defined *groupoid* $\Gamma(X; \mathfrak{D}_X^{ch})$. This groupoid is empty if the class (3.1) is non-trivial.

If (3.1) is trivial then this groupoid is equivalent to the groupoid of $\Omega_X^{2,cl}$ -torsors on X . Thus, in this case the set of isomorphism classes of $\Gamma(X; \mathfrak{D}_X^{ch})$ is isomorphic to $H^1(X; \Omega_X^{2,cl})$, and the automorphism group of an object is isomorphic to $H^0(X; \Omega_X^{2,cl})$.

From this one sees immediately that for curves the obstruction vanishes, and the above groupoid is trivial: one can define canonically "the" sheaf of chiral differential operators \mathcal{D}_X^{ch} . For the explicit construction, see (6.2) below.

The same is true for the flag spaces G/B . This explains the constructions of [MSV]. As a negative example, note that for the projective spaces of dimension greater than 1, the class (3.1) is non-zero.

4. Let us explain a local statement which implies Theorem 3. Let $\widehat{O} \supset O$ be the algebra of the Taylor power series in variables b^i convergent in some neighbourhood of the origin. Let G be the group whose elements are N -tuples $g = (g^1(b), \dots, g^N(b))$ where $g^i \in \widehat{O}$ and the Jacobian matrix $\partial g := (\partial_i g^j)$ is non-degenerate. The composition is defined by

$$(g \circ h)^i(b) = g^i(-1(b)) \circ h^i(b) \quad (4.1)$$

The group G acts in the obvious way from the right on the algebra \widehat{O} .

We have

$$\partial(g_1 g_2) = \partial g_1 \partial g_2 g_1 \quad (4.2)$$

Here we use the notation

$$h_g(b) := h(g(b)) \quad (4.3)$$

We will denote by $\Omega_{\widehat{O}}^1$ (resp., $\mathcal{T}_{\widehat{O}}$) the module of germs of holomorphic 1-differentials (resp., holomorphic vector fields). It is a free \widehat{O} -module with the base $\{db^i\}$ (resp., $\{\partial_i\}$).

Let us consider the vertex algebra $\widehat{D} := \widehat{O} \otimes_O D$. We have $\widehat{D}_0 = O$, and \widehat{D}_1 has a canonical two step filtration defined as in (2.2), with $F^1 \widehat{D}_1 = \Omega_{\widehat{O}}^1$, $\widehat{D}_1 / F^1 \widehat{D}_1 = \mathcal{T}_{\widehat{O}}$.

Let us call an automorphism of \widehat{D} natural if it induces an automorphism $g \in G$ on \widehat{D}_0 , and the induced by g automorphism on $gr_F \widehat{D}_1$.

We want to describe the group G^\sim of all natural automorphisms of \widehat{D} . In coordinates, a natural automorphism is given by a pair (g, h) where $g \in G$ and $h \in M_N(\widehat{O})$ is an $N \times N$ -matrix over \widehat{O} . By definition, such a pair acts on the generating fields as follows

$$(b^i \circ (g, h))(z) = g^i(b)(z) \quad (4.4a)$$

$$(a^i \circ (g, h))(z) = a^j \phi^{ji}(g)(z) + b^k(z)' h^{ki}(z) \quad (4.4b)$$

(this will be the right action). Here the matrix $\phi(g) = (\phi^{ij}(g))$ is defined by

$$\phi(g) = (\partial g)^{-1t} \quad (4.5)$$

where $(a^t)^{ij} = a^{ij}$.

The straightforward computation of OPE shows that a pair (g, h) defines an automorphism of the vertex algebra \widehat{D} if and only if the matrix h satisfies the equations

$$h^t \phi(g) + \phi^t(g) h = \psi(g) \quad (4.6a)$$

and

$$\begin{aligned} & \phi^{pi} \partial_p h^{sj} - \phi^{pj} \partial_p h^{si} + \\ & + \partial_s \phi^{pi} h^{pj} + \partial_s h^{pi} \phi^{pj} - \partial_s \partial_r \phi^{pi} \partial_p \phi^{rj} = 0 \end{aligned} \quad (4.6b)$$

where $\psi(g) = (\psi^{ij}(g))$ is a symmetric matrix

$$\psi^{ij}(g) = \partial_r \phi^{pi}(g) \partial_p \phi^{rj}(g) \quad (4.7)$$

(we imply everywhere the summation over the repeating indices).

The computation similar to that in the proof of Theorem 3.8 from [MSV], shows that the composition on G^\sim is given by the formula

$$(g_1, h_1)(g_2, h_2) = (g_1 g_2, h_1 \phi(g_2)_{g_1} + \partial g_1 h_2 g_1 + \alpha(g_1, g_2)) \quad (4.8)$$

where $\alpha(g_1, g_2) = (\alpha^{ij}(g_1, g_2))$ is given by

$$\alpha^{ij}(g_1, g_2) = \partial_r \phi^{pq}(g_1) \partial_p [\phi^{qi}(g_2) - 1] \quad (4.9)$$

This describes the group G^\sim . The unit is $(1, 0)$. Let us introduce a bigger group G' consisting of all pairs (g, h) as above, with the multiplication (4.8), but with h not necessarily satisfying the conditions (4.6 a, b). One has to check the associativity of multiplication, which is straightforward. Obviously, $G^\sim \subset G'$.

It is convenient to introduce in G' the new coordinates by setting

$$(g, h)^\sim := (g, 0)(1, h) = (g, h\partial g^{-1t}) \quad (4.9)$$

In new coordinates the multiplication in G' looks as follows

$$(g_1, h_1)^\sim (g_2, h_2)^\sim = (g_1 g_2, h_1 + \partial g_1 h_2 g_1 \partial g_1^t + \beta(g_1, g_2))^\sim \quad (4.10)$$

where

$$\beta(g_1, g_2) = \alpha(g_1, g_2) \partial (g_1 g_2)^t \quad (4.11)$$

In particular,

$$(1, h)(g, 0) = (h, \partial g \cdot h_g \cdot \partial g^t) \quad (4.12)$$

We have an obvious epimorphism $G' \rightarrow G$, $(g, h) \mapsto g$, and the formula (4.12) shows that its kernel, as a G -module, may be identified with the G -module of quadratic differentials $\Omega_{\hat{O}}^{1 \otimes 2} := \Omega_{\hat{O}}^1 \otimes_{\hat{O}} \Omega_{\hat{O}}^1$, i.e. we have the group extension

$$0 \rightarrow \Omega_{\hat{O}}^{1 \otimes 2} \rightarrow G' \rightarrow G \rightarrow 1 \quad (4.13)$$

Equation (4.10) means that if we chose a section of the epimorphism $G' \rightarrow G$ as $g \mapsto (g, 0)$ then the extension (4.13) corresponds to the 2-cocycle $\beta(g_1, g_2) \in Z^2(G; \Omega_{\hat{O}}^2)$. The equation of a cocycle (equivalent to the associativity of multiplication in G') reads as

$$\beta(g_1, g_2) - \beta(g_1, g_2 g_3) + \beta(g_1 g_2, g_3) - g_1^* \beta(g_2, g_3) = 0 \quad (4.14)$$

A little computation using the formula $\partial_i(A^{-1}) = -A^{-1} \partial_i A \cdot A^{-1}$ gives the following expression for β :

$$\beta^{ij}(g_1, g_2) = \partial(g_1^{-1})^{qa} \partial_i \partial_a g_1^b \partial^j [(\partial_b g_2^r)_{g_1}] (\partial g_2^{-1})_{g_1}^{rq} \quad (4.15)$$

which may be rewritten in the following compact form:

$$\beta(g_1, g_2) = \text{tr} \{ \partial g_1^{-1} d \partial g_1 \otimes g_1^* (d \partial g_2 \cdot \partial g_2^{-1}) \} \quad (4.16)$$

5. Now let us take care of the subgroup $G'' \subset G'$ given inside G' by the equation (4.6a). Since the matrix $\psi(g)$ is symmetric, and the matrix h is arbitrary, the above equation has $N(N-1)/2$ -dimensional (over \hat{O}) space of solutions.

For example, for $N = 1$, (4.6) has a unique solution, and $G'' = G$, namely

$$h(g) = g''^2 / 2g' \quad (5.1)$$

Now let us treat the case of an arbitrary N . First of all, $\psi(1) = 0$ and $\phi(1) = 1$, and the equation (4.6) means simply that the matrix h is skew symmetric. This means that G'' is included in a group extension

$$0 \rightarrow \Omega_{\hat{O}}^2 \rightarrow G'' \rightarrow G \rightarrow 1 \quad (5.2)$$

It is easy to write down a particular solution of (4.6a), namely

$$h(g) = \phi^t(g)^{-1} \psi(g)/2 \quad (5.3)$$

This gives a section of (5.2), given by

$$g \mapsto (g, h(g)) =: (g, s(g))^\sim \quad (5.4)$$

If $c(g_1, g_2) \in Z^2(G; \Omega_{\hat{O}}^2)$ is the cocycle corresponding to this choice of the section then we have

$$c(g_1, g_2) = \beta(g_1, g_2) + s(g_1) - s(g_1 g_2) + \partial g_1 s(g_2)_{g_1} \partial g_1^t \quad (5.5)$$

In other words, c is cohomologous to β . But as far as we know that the cohomology class of β comes from a cohomology class from $H^2(G; \Omega_{\hat{O}}^2)$, this last class is unique and may be given simply by the skew symmetrization of β , i.e. by

$$\begin{aligned} \gamma(g_1, g_2) &= \text{tr}\{\partial g_1^{-1} d\partial g_1 \wedge g_1^*(d\partial g_2 \cdot \partial g_2^{-1})\} = \\ &= \text{tr}\{d\partial g_1 \cdot \partial g_1^{-1} \wedge \partial g_1 \cdot g_1^*[d\partial g_2 \cdot \partial g_2^{-1}] \cdot \partial g_1^{-1}\} \end{aligned} \quad (5.6)$$

Comparing the expression (5.6) with the formula (4.5) from [TT] we see that the cocycle γ is nothing but the universal cocycle of G representing the characteristic class corresponding to the invariant polynomial $\text{tr}(AB)$ (on pairs of $N \times N$ matrices), i.e. it represents the component of the Chern character $c_1^2 - 2c_2$.

Thus, we arrive at

5.1. Proposition. *The group G'' is included in the group extension*

$$0 \longrightarrow \Omega_{\hat{O}}^2 \longrightarrow G'' \longrightarrow G \longrightarrow 1 \quad (5.7)$$

The cohomology class of this extension is equal to $2ch_2 := c_1^2 - 2c_2 \in H^2(G; \Omega_{\hat{O}}^2)$ where $c_i \in H^i(G; \Omega_{\hat{O}}^i)$ are the universal Chern classes.

6. Finally, group G^\sim is given inside G'' by the equation (4.6b). For $g = 1$ this equation simply means that the form $h^{ij} db^i \wedge db^j$ is closed (we are grateful to A.Beilinson for this remark).

Therefore, the kernel of the obvious map $G^\sim \longrightarrow G$ is isomorphic to the G -module $\Omega_{\hat{O}}^{2,cl}$ of closed 2-forms.

6.1. Lemma. *The function $h(g)$ given by the formula (5.3) satisfies the equation (4.6b).*

In fact, if we symmetrize the equation (4.6b) with respect to i, j , we get $\partial_s(4.6a)$. Therefore, it is enough to check the skew symmetrized part of (4.6b). This is done by a direct computation, using (4.5).

This lemma gives a section of the canonical map $G^\sim \longrightarrow G$; in particular it is surjective.

The above considerations, together with Proposition 5.1, imply

6.2. Theorem. *The group G^\sim of all natural automorphisms of the vertex algebra \widehat{D} is included in the group extension*

$$0 \longrightarrow \Omega^{2,cl} \longrightarrow G^\sim \longrightarrow G \longrightarrow 1 \quad (6.1)$$

The cohomology class of this extension is equal to $2ch_2 := c_1^2 - 2c_2 \in H^2(G; \Omega^{2,cl})$ where $c_i \in H^i(G; \Omega^{i,cl})$ are the universal Chern classes.

Note that the standard cocycles representing the universal Chern classes in $H^i(G; \Omega_{\widehat{O}}^i)$ in fact take values in $\Omega_{\widehat{O}}^{i,cl}$.

Let us sketch another argument proving surjectivity of the map $G^\sim \longrightarrow G$. Note that in [MSV], §5 one has lifted an arbitrary element of the Lie algebra $W_N := Lie(G)$ to a derivation $\pi(\tau)$ of \widehat{D} . Since every $g \in G$ is the exponent of some vector field τ , we can take $\tilde{g} = \exp(\pi(\tau))$ as a lifting of g .

Consider the case $N = 1$. In this case $G^\sim = G'' = G$. Thus, every automorphism $g \in G$ lifts uniquely to a natural automorphism of \widehat{D} given by

$$\tilde{g} = (g, g''^2/2g') \quad (6.2)$$

cf. (5.1). This corresponds to the fact that for curves the sheaf \mathcal{D}_X^{ch} is uniquely defined. Note that the formula [MSV], (5.23b) is a particular case of (6.2) (for $g(b) = b^{-1}$).

Our main Theorem 3 is an immediate consequence of Theorem 6.1.

7. Let E be a vector bundle over X , ΛE its exterior algebra. Proceeding in a similar way as above, we can produce a gerbe $\mathfrak{D}_{\Lambda E}^{ch}$ of chiral differential operators on ΛE , also bounded by the lien $\Omega_X^{2,cl}$.

Let us explain how to do this. Let us describe the local model. In the notations of no. 2, fix an integer $M \geq 0$. Let us define a vertex superalgebra $D_\Lambda \supset D$ which is generated by the free bosonic fields $b^i(z), a^i(z)$ as in *loc. cit.* and free odd fields $\phi^j(z)$ of conformal weight 0 and $\psi^j(z)$ of conformal weight 1 ($j = 1, \dots, M$), with the OPE (2.1) and

$$\psi^i(z)\phi^j(w) \sim \frac{\delta_{ij}}{z-w} \quad (7.1)$$

the other products being trivial. Set $\widehat{D}_\Lambda := \widehat{O} \otimes_O D_\Lambda$. Consider the group H which is the semidirect product of the group of coordinate changes G from no. 4, the group of $M \times M$ matrices $Mat_M(\widehat{O})$, with the obvious action of G on $Mat_M(\widehat{O})$. The group H will be the analogue of the group G in our situation.

Let us call an automorphism of the superalgebra \widehat{D}_Λ *admissible* if it has the form

$$\tilde{b}^i(z) = g^i(b)(z) \quad (7.2a)$$

$$\tilde{a}^i(z) = a^j c^{ji}(g)(z) - \psi^p \phi^q A^{qs} c^{ki} \partial_k A^{-1sp}(z) + b^j(z)' h^{ij}(z) \quad (7.2b)$$

$$\tilde{\phi}^i(z) = \phi^j A^{ji}(z) \quad (7.2c)$$

$$\tilde{\psi}^i(z) = \psi^j A^{-1ji}(z) \quad (7.2d)$$

for some $(g, A) \in H$ and $h \in \text{Mat}_N(\widehat{O})$. Here $c(g) := (\partial g)^{-1t}$ (we have changed the notation from (4.5), since we already have ϕ 's).

We want to compute the group \tilde{H} of all admissible automorphisms of \widehat{D}_Λ . The composition will be

$$\begin{aligned} & (g_1, A_1, h_1)(g_2, A_2, h_2) = \\ & = (g_1 g_2, A_1 A_{2g_1}, h_1 c(g_2)_{g_1} + \partial g_1 h_{2g_1} + \alpha(g_1, g_2) + \gamma(g_1, g_2)) \end{aligned} \quad (7.3)$$

where $\alpha(g_1, g_2)$ is given by (4.9), and

$$\gamma(g_1, A_1; g_2, A_2)^{ij} = \partial_i A_1^{-1pr} A_1^{rq} A_{2g_1}^{qs} c(g_2)_{g_1}^{kj} \partial_k A_{2g_1}^{-1sp} \quad (7.4)$$

Set

$$\delta(g_1, A_1; g_2, A_2) = \gamma(g_1, A_1, g_2, A_2) \partial(g_1 g_2)^t \quad (7.5)$$

cf. (4.11). A little computation gives

7.1. Lemma. *We have*

$$\delta(g_1, A_1, g_2, A_2) = \text{tr}\{dA_1 \cdot A_1^{-1} \otimes A_1 g_1^*(dA_2 \cdot A_2^{-1}) A_1^{-1}\} \quad (7.6)$$

Cf. (4.16).

Let H' be the group of all triples (g, A, h) as above, with the composition given by (7.3). We arrive at

7.2. Proposition. *The group H' is included in the extension*

$$0 \longrightarrow \Omega_{\widehat{O}}^{1 \otimes 2} \longrightarrow H' \longrightarrow H \longrightarrow 1 \quad (7.7)$$

given by the two-cocycle

$$\epsilon(g_1, A_1; g_2, A_2) = \beta(g_1, g_2) - \gamma(g_1, A_1; g_2, A_2) \quad (7.8)$$

where β (resp. γ) is given by (4.16) (resp. by (7.6)).

Now, proceeding in the same way as in nos. 5, 6, we arrive at

7.3. Proposition. *The group \tilde{H} of all admissible automorphisms of the vertex algebra \widehat{D}_Λ is included into the extension*

$$0 \longrightarrow \Omega_{\widehat{O}}^{2, cl} \longrightarrow \tilde{H} \longrightarrow H \longrightarrow 1 \quad (7.9)$$

whose cohomology class is given by the skew symmetrization of the cocycle (7.8), i.e. equal to

$$c_1^2 - 2c_2 - (c_1'^2 - 2c_2') \in H^2(H, \Omega_{\widehat{O}}^{2, cl}) \quad (7.10)$$

where $c_i, c_i' \in H^i(H; \Omega_{\widehat{O}}^{i, cl})$ are the universal Chern classes.

8. Returning to the global situation of the beginning of the preceeding subsection (we assume that $\dim(X) = N$, $rk(E) = M$), we define for a sufficiently small $U \subset X$ the groupoid $\mathfrak{D}_{\Lambda E}^{ch}(U)$ whose objects are pairs

(ϕ coordinate system on U , σ trivialization of E over U)

and morphisms are defined as in no. 2, using the admissible isomorphisms of the corresponding vertex superalgebras. This way we get the gerbe $\mathfrak{D}_{\Lambda E}^{ch}$ over X .

Proposition 7.3 implies

8.1. Theorem. *The gerbe $\mathfrak{D}_{\Lambda E}^{ch}$ is bounded by the lien $\Omega_X^{2,cl}$, and its equivalence class is equal to*

$$c(\mathfrak{D}_{\Lambda E}^{ch}) = 2ch_2(\mathcal{T}_X) - 2ch_2(E) \in H^2(X, \Omega_X^{2,cl}) \quad (8.1)$$

Note that the class (8.1) does not change when E is replaced by the dual bundle.

We see that for $E = \Omega_X^1$, or \mathcal{T}_X we arrive at the neutral gerbe. In fact, the groupoid of global sections $\mathfrak{D}_{\Lambda E}^{ch}(X)$ in both cases admits a canonical global section. For $E = \Omega_X^1$ it is the "chiral de Rham complex" from [MSV], and for $E = \mathcal{T}_X$ it is a sort of a "mirror partner".

References

- [D]
- [
- [D] P. Deligne, Le symbole modéré, *Publ. Math. IHES*, **73** (1991), 147 -181.
- [MSV] F. Malikov, V. Schechtman, A. Vaintrob, Chiral de Rham complex, *Comm. Math. Phys.* (1999), to appear; math.AG/9803041.
- [MS1] F. Malikov, V. Schechtman, Chiral de Rham complex. II, *D.B. Fuchs' 60-th Anniversary volume* (1999), to appear; math.AG/9901065.
- [MS2] F. Malikov, V. Schechtman, Chiral Poincaré duality, math.AG/9905008.
- [TT] D. Toledo, Y.L. Tong, A parametrix for $\bar{\partial}$ and Riemann-Roch in Cech theory, *Topology*, **15** (1976), 273 - 301.

V.G.: Department of Mathematics, University of Kentucky, Lexington, KY 40506, USA; vgorb@ms.uky.edu

F.M.: Department of Mathematics, University of Southern California, Los Angeles, CA 90089, USA; fmalikov@mathj.usc.edu

V.S.: Department of Mathematics, University of Glasgow, 15 University Gardens, Glasgow G12 8QW, UK; vs@maths.gla.ac.uk, vadik@mpim-bonn.mpg.de