On a relation between intrinsic and extrinsic Dirichlet forms for interacting particle systems

José L. da Silva

BiBoS, Bielefeld Univ., D-33615 Bielefeld, Germany CCM, Madeira Univ, P-9000 Funchal, Madeira Portugal

Yuri G. Kondratiev

Inst. Ang. Math., Bonn Univ., D-53115 Bonn, Germany BiBoS, Bielefeld Univ., D-33615 Bielefeld, Germany Inst. Math., NASU, 252601 Kiev, Ukraine

Michael Röckner

Fakultät für Mathematik, Universität Bielefeld, D-33615 Bielefeld, Germany

February 5, 2018

Abstract

In this paper we extend the result obtained in [AKR98a] (see also [AKR96a]) on the representation of the intrinsic pre-Dirichlet form $\mathcal{E}_{\pi_{\sigma}}^{\Gamma}$ of the Poisson measure π_{σ} in terms of the extrinsic one $\mathcal{E}_{\pi_{\sigma},H_{\sigma}^{X}}^{P}$. More precisely, replacing π_{σ} by a Gibbs measure μ on the configuration space Γ_{X} we derive a relation between the intrinsic pre-Dirichlet form $\mathcal{E}_{\mu}^{\Gamma}$ of the measure μ and the extrinsic one $\mathcal{E}_{\mu,H_{\sigma}^{X}}^{P}$. As a consequence we prove the closability of $\mathcal{E}_{\mu}^{\Gamma}$ on $L^{2}(\Gamma_{X},\mu)$ under very general assumptions on the interaction potential of the Gibbs measures μ .

Mathematics Subject Classification: primary 31C25, 60G60; secondary 82C22, 82C21

1 Introduction

In the recent papers [AKR96a], [AKR96b], [AKR98a], and [AKR98b] analysis and geometry on configuration spaces Γ_X over a Riemannian manifold X, i.e,

$$\Gamma_X := \{ \gamma \subset X \mid |\gamma \cap K| < \infty \text{ for any compact } K \subset X \},\$$

was developed. One of the consequences of the discussed approach was a description of the well-known equilibrium process on configuration spaces as the Brownian motion associated with a Dirichlet form of the Poisson measure π_{σ} with intensity measure σ on $\mathcal{B}(X)$. This form is canonically associated with the introduced geometry on configuration spaces and is called *intrinsic Dirichlet form* of the measure π_{σ} .

On the other hand there is a well-known realization (canonical isomorphism) of the Hilbert space $L^2(\Gamma_X, \pi_\sigma)$ and the corresponding Fock space

$$\mathrm{Exp}L^2(X,\sigma) := \bigoplus_{n=0}^{\infty} \mathrm{Exp}_n L^2(X,\sigma),$$

where $\operatorname{Exp}_n L^2(X, \sigma)$ denotes the *n*-fold symmetric tensor product of $L^2(X, \sigma)$ and $\operatorname{Exp}_0 L^2(X, \sigma) := \mathbb{C}$. This isomorphism produces natural operations in $L^2(\Gamma_X, \pi_\sigma)$ as images of the standard Fock space operators, see e.g., [KSSU98] and references therein. In particular, we can consider the image of the annihilation operator from the Fock space as a natural version of a "gradient" operator in $L^2(\Gamma_X, \pi_\sigma)$. The differentiable structure in $L^2(\Gamma_X, \pi_\sigma)$ which appears in this way we consider as *external* because it is produced via transportation from the Fock space.

As was shown in [AKR98a, Section 5] the intrinsic Dirichlet form of the measure π_{σ} can be represented also in terms of the external Dirichlet form $\mathcal{E}^{P}_{\pi_{\sigma},H^{X}_{\sigma}}$ with coefficient H^{X}_{σ} (the Dirichlet operator associated with σ on X) which uses this external differentiable structure, i.e.,

$$\int_{\Gamma} \langle \nabla^{\Gamma} F(\gamma), \nabla^{\Gamma} G(\gamma) \rangle_{T_{\gamma}\Gamma} d\pi_{\sigma}(\gamma) = \int_{\Gamma} (\nabla^{P} F(\gamma), H^{X}_{\sigma} \nabla^{P} G(\gamma))_{L^{2}(X,\sigma)} d\pi_{\sigma}(\gamma).$$

As a result we have a full spectral description of the corresponding Dirichlet operator $H_{\pi_{\pi}}^{\Gamma}$ which is the generator of the equilibrium process on Γ_X .

If we change the Poisson measure π_{σ} to a Gibbs measure μ on the configuration space Γ_X which describes the equilibrium of interacting particle

systems, the corresponding intrinsic Dirichlet form can still be used for constructing the corresponding stochastic dynamics (cf. [AKR98b, Section 5]) or for constructing a quantum infinite particle Hamiltonian in models of quantum fields theory, see [AKR99].

The aim of this paper is to show that even for the interacting case there is a transparent relation between the intrinsic Dirichlet form and the extrinsic one, see Theorem 5.1. The proof is based on the Nguyen-Zessin characterization of Gibbs measure (cf. [NZ79, Theorem 2] or Proposition 5.2 below) which on a heuristic level can be considered as a consequence of the Mecke identity (cf. [Mec67, Satz 3.1]), see Remark 5.3 below for more details.

As a consequence of the mentioned relation we prove the closability of the pre-Dirichlet form $(\mathcal{E}^{\Gamma}_{\mu}, \mathcal{F}C^{\infty}_{b}(\mathcal{D}, \Gamma))$ on $L^{2}(\Gamma_{X}, \mu)$, where μ is a tempered grand canonical Gibbs measure, see Section 2 for this notion. It turns out that this result is obtained as a "lifting" of the closable Dirichlet forms on X. We would like to emphasize that we achieve this result under a general condition (see (6.2) below) on the potential Φ which is not covered by condition (A.6) in [Osa96]. The closability is crucial (for physical reasons, see [AKR99], and) for applying the general theory of Dirichlet forms including the construction of a corresponding diffusion process (cf. [MR92]) which models an infinite particle system with (possibly) very singular interactions (cf. [AKR98b]).

Another motivation for deriving Theorem 5.1 is to use this result for studying spectral properties of Hamiltonians of intrinsic Dirichlet forms associated with Gibbs measures. This will be implemented in a forthcoming paper.

2 Preliminaries and Framework

In this section we describe some facts about probability measures on configuration spaces which are necessary later on.

Let X be a connected, oriented C^{∞} (non-compact) Riemannian manifold. For each point $x \in X$, the tangent space to X at x will be denoted by T_xX ; and the tangent bundle will be denoted by $TX = \bigcup_{x \in X} T_xX$. The Riemannian metric on X associates to each point $x \in X$ an inner product on T_xX which we denote by $\langle \cdot, \cdot \rangle_{T_xX}$ and the associated norm will be denoted by $|\cdot|_{T_xX}$. Let m denote the volume element.

 $\mathcal{O}(X)$ is defined as the family of all open subsets of X and $\mathcal{B}(X)$ denotes the corresponding Borel σ -algebra. $\mathcal{O}_c(X)$ and $\mathcal{B}_c(X)$ denote the systems of all elements in $\mathcal{O}(X)$, $\mathcal{B}(X)$ respectively, which have compact closures.

Let $\Gamma := \Gamma_X$ be the set of all locally finite subsets in X:

$$\Gamma_X := \{ \gamma \subset X \mid |\gamma \cap K| < \infty \text{ for any compact } K \subset X \}.$$

We will identify γ with the positive integer-valued measure $\sum_{x \in \gamma} \varepsilon_x$. Then for any $\varphi \in C_0(X)$ we have a functional $\Gamma \ni \gamma \mapsto \langle \varphi, \gamma \rangle = \sum_{x \in \gamma} \varphi(x) \in \mathbb{R}$. Here $C_0(X)$ is the set of all real-valued continuous functions on X with compact support. The space Γ is endowed with the vague topology. Let $\mathcal{B}(\Gamma)$ denote the corresponding Borel σ -algebra. For $\Lambda \subset X$ we sometimes use the shorthand γ_{Λ} for $\gamma \cap \Lambda$.

For any $B \in \mathcal{B}(X)$ we define, as usual, $\Gamma \ni \gamma \mapsto N_B(\gamma) := \gamma(B) \in \mathbb{Z}_+ \cup \{+\infty\}$. Then $\mathcal{B}(\Gamma) = \sigma(\{N_\Lambda | \Lambda \in \mathcal{O}_c(X)\})$. For any $A \in \mathcal{B}(X)$ we also define $\mathcal{B}_A(\Gamma) := \sigma(\{N_B | B \in \mathcal{B}_c(X), B \subset A\})$.

Let $d\sigma(x) = \rho(x)dm(x)$, where $\rho > 0$ *m*-a.e. such that $\rho^{\frac{1}{2}} \in H^{1,2}_{loc}(X)$ (the Sobolev space of order 1 in $L^2(X,m)$) and $\rho \notin L^1(X,m)$. We recall that the Poisson measure π_{σ} (with intensity measure σ) on $(\Gamma, \mathcal{B}(\Gamma))$ is defined via its Laplace transform by

$$\int_{\Gamma} e^{\langle \gamma, \varphi \rangle} d\pi_{\sigma}(\gamma) = \exp\left(\int_{X} (e^{\varphi(x)} - 1) d\sigma(x)\right), \ \varphi \in C_0(X), \tag{2.1}$$

see e.g. [AKR98a], [GV68], and [Shi94]. Let us mention that if $\rho \in L^1(X, m)$, then we have a finite intensity measure σ on X, and in this case the corresponding measure π_{σ} will be concentrated on finite configurations. The latter can be considered as a degenerated case which can be reduced to finite dimensional analysis on every subset of *n*-particle configurations.

Let us briefly recall the definition of grand canonical Gibbs measures on $(\Gamma, \mathcal{B}(\Gamma))$. We adopt the notation in [AKR98b], and refer the interested reader to the beautiful work by C. Preston, [Pre79], but also [Pre76], and [Geo79].

A function $\Phi : \Gamma \to \mathbb{R} \cup \{+\infty\}$ will be called a *potential* iff for all $\Lambda \in \mathcal{B}_c(X)$ we have $\Phi(\emptyset) = 0$, $\Phi = \mathbb{1}_{\{N_X < \infty\}}\Phi$, and $\gamma \mapsto \Phi(\gamma_\Lambda)$ is $\mathcal{B}_{\Lambda}(\Gamma)$ -measurable.

For $\Lambda \in \mathcal{B}_c(X)$ the conditional energy $E_{\Lambda}^{\Phi} : \Gamma \to \mathbb{R} \cup \{+\infty\}$ is defined by

$$E_{\Lambda}^{\Phi}(\gamma) := \begin{cases} \sum_{\gamma' \subset \gamma, \gamma'(\Lambda) > 0} \Phi(\gamma') & \text{if } \sum_{\gamma' \subset \gamma, \gamma'(\Lambda) > 0} |\Phi(\gamma')| < \infty, \\ +\infty & \text{otherwise,} \end{cases}$$
(2.2)

where the sum of the empty set is defined to be zero.

Later on we will use conditional energies which satisfy an additional assumption, namely, the *stability condition*, i.e., there exists $B \ge 0$ such that for any $\Lambda \in \mathcal{B}_c(X)$ and for all $\gamma \in \Gamma_{\Lambda}$

$$E^{\Phi}_{\Lambda}(\gamma) \ge -B|\gamma|.$$

Definition 2.1 For any $\Lambda \in \mathcal{O}_c(X)$ define for $\gamma \in \Gamma$ the measure $\Pi^{\Phi}_{\Lambda}(\gamma, \cdot)$ by

$$\Pi_{\Lambda}^{\sigma,\Phi}(\gamma,\Delta) := \mathbb{1}_{\{Z_{\Lambda}^{\sigma,\Phi}<\infty\}}(\gamma)[Z_{\Lambda}^{\sigma,\Phi}(\gamma)]^{-1} \int_{\Gamma} \mathbb{1}_{\Delta}(\gamma_{X\setminus\Lambda}+\gamma'_{\Lambda}) \qquad (2.3)$$
$$\cdot \exp[-E_{\Lambda}^{\Phi}(\gamma_{X\setminus\Lambda}+\gamma'_{\Lambda})]d\pi_{\sigma}(\gamma'), \ \Delta \in \mathcal{B}(\Gamma),$$

where

$$Z^{\sigma,\Phi}_{\Lambda}(\gamma) := \int_{\Gamma} \exp[-E^{\Phi}_{\Lambda}(\gamma_{X\backslash\Lambda} + \gamma'_{\Lambda})] d\pi_{\sigma}(\gamma').$$
(2.4)

A probability measure μ on $(\Gamma, \mathcal{B}(\Gamma))$ is called grand canonical Gibbs measure with interaction potential Φ if for all $\Lambda \in \mathcal{O}_c(X)$

$$\mu \Pi_{\Lambda}^{\Phi} = \mu. \tag{2.5}$$

Let $\mathcal{G}_{gc}(\sigma, \Phi)$ denote the set of all such probability measures μ .

- **Remark 2.2** 1. It is well-known that $(\Pi_{\Lambda}^{\sigma,\Phi})_{\Lambda\in\mathcal{O}_{c}(X)}$ is a $(\mathcal{B}_{X\setminus\Lambda}(\Gamma))_{\Lambda\in\mathcal{O}_{c}(X)}$ -specification in the sense of [Pre76, Section 6] or [Pre79].
 - 2. For any $\gamma \in \Gamma$ the measure $\mu \Pi^{\Phi}_{\Lambda}$ in (2.5) is defined by

$$(\mu \Pi_{\Lambda}^{\sigma, \Phi})(\Delta) := \int_{\Gamma} d\mu(\gamma) \Pi_{\Lambda}^{\sigma, \Phi}(\gamma, \Delta), \ \Delta \in \mathcal{B}(\Gamma)$$
(2.6)

and (2.5) are called Dobrushin-Landford-Ruelle (DLR) equations.

3 Intrinsic geometry on Poisson space

We recall some results to be used below from [AKR98a], [AKR96b] to which we refer for the corresponding proofs and more details.

A homeomorphism $\psi: X \to X$ defines a transformation of Γ by

$$\Gamma \ni \gamma \mapsto \psi(\gamma) = \{\psi(x) | x \in \gamma\} = \sum_{x \in \gamma} \varepsilon_{\psi(x)}.$$

Any vector field $v \in V_0(X)$ (i.e., the set of all smooth vector fields on X with compact support) defines (via the exponential mapping) a one-parameter group ψ_t^v , $t \in \mathbb{R}$, of diffeomorphisms of X.

Definition 3.1 For $F : \Gamma \to \mathbb{R}$ we define the directional derivative along the vector field v as (provided the right hand side exists)

$$(\nabla_v^{\Gamma} F)(\gamma) := \frac{d}{dt} F(\psi_t^v(\gamma))|_{t=0}$$

This definition applies to F in the following class $\mathcal{F}C_b^{\infty}(\mathcal{D},\Gamma)$ of so-called smooth cylinder functions. Let $\mathcal{D} := C_0^{\infty}(X)$ (the set of all smooth functions on X with compact support). We define $\mathcal{F}C_b^{\infty}(\mathcal{D},\Gamma)$ as the set of all functions on Γ of the form

$$F(\gamma) = g_F(\langle \gamma, \varphi_1 \rangle, \dots, \langle \gamma, \varphi_N \rangle), \ \gamma \in \Gamma,$$
(3.1)

where $\varphi_1, \ldots, \varphi_N \in \mathcal{D}$ and g_F is from $C_b^{\infty}(\mathbb{R}^N)$. Clearly, $\mathcal{F}C_b^{\infty}(\mathcal{D}, \Gamma)$ is dense in $L^2(\pi_{\sigma}) := L^2(\Gamma, \pi_{\sigma})$. For any $F \in \mathcal{F}C_b^{\infty}(\mathcal{D}, \Gamma)$ we have

$$(\nabla_v^{\Gamma} F)(\gamma) = \sum_{i=1}^N \frac{\partial g_F}{\partial s_i} (\langle \gamma, \varphi_1 \rangle, \dots, \langle \gamma, \varphi_N \rangle) \langle \gamma, \nabla_v^X \varphi_i \rangle, \qquad (3.2)$$

where $x \mapsto (\nabla_v^X \varphi)(x) = \langle \nabla^X \varphi(x), v(x) \rangle_{TX}$ is the usual directional derivative on X along the vector field v and ∇^X denotes the gradient on X.

The logarithmic derivative of the measure σ is given by the vector field $\beta^{\sigma} := \nabla^X \log \rho = \nabla^X \rho / \rho$ (where $\beta^{\sigma} = 0$ on $\{\rho = 0\}$). Then the logarithmic derivative of σ along v is the function $x \mapsto \beta_v^{\sigma}(x) = \langle \beta^{\sigma}(x), v(x) \rangle_{T_x X} + \operatorname{div}^X v(x)$, where div^X denotes the divergence on X w.r.t. the volume element m. Analogously, we define $\operatorname{div}_{\sigma}^X$ as the divergence on X w.r.t. σ , i.e., $\operatorname{div}_{\sigma}^X$ is the dual operator on $L^2(\sigma) := L^2(X, \sigma)$ of ∇^X .

Definition 3.2 For any $v \in V_0(X)$ we define the logarithmic derivative of π_{σ} along v as the following function on Γ :

$$\Gamma \ni \gamma \mapsto B_v^{\pi_\sigma}(\gamma) := \langle \beta_v^{\sigma}, \gamma \rangle = \int_X [\langle \beta^{\sigma}(x), v(x) \rangle_{T_x X} + \operatorname{div}^X v(x)] d\gamma(x). \quad (3.3)$$

Theorem 3.3 For all $F, G \in \mathcal{F}C_b^{\infty}(\mathcal{D}, \Gamma)$ and any $v \in V_0(X)$ the following integration by parts formula for π_{σ} holds:

$$\int_{\Gamma} \nabla_v^{\Gamma} F G d\pi_{\sigma} = -\int_{\Gamma} F \nabla_v^{\Gamma} G d\pi_{\sigma} - \int_{\Gamma} F G B_v^{\pi_{\sigma}} d\pi_{\sigma}, \qquad (3.4)$$

or $(\nabla_v^{\Gamma})^* = -\nabla_v^{\Gamma} - B_v^{\pi_{\sigma}}$, as an operator equality on the domain $\mathcal{F}C_b^{\infty}(\mathcal{D},\Gamma)$ in $L^2(\pi_{\sigma})$.

Definition 3.4 We introduce the tangent space $T_{\gamma}\Gamma$ to the configuration space Γ at the point $\gamma \in \Gamma$ as the Hilbert space of γ -square-integrable sections (measurable vector fields) $V : X \to TX$ with scalar product $\langle V^1, V^2 \rangle_{T_{\gamma}\Gamma} = \int_X \langle V^1(x), V^2(x) \rangle_{T_x X} d\gamma(x), V^1, V^2 \in T_{\gamma}\Gamma = L^2(X \to TX; \gamma)$. The corresponding tangent bundle is denoted by $T\Gamma$.

The intrinsic gradient of a function $F \in \mathcal{F}C_b^{\infty}(\mathcal{D},\Gamma)$ is a mapping $\Gamma \ni \gamma \mapsto (\nabla^{\Gamma} F)(\gamma) \in T_{\gamma}\Gamma$ such that $(\nabla_v^{\Gamma} F)(\gamma) = \langle \nabla^{\Gamma} F(\gamma), v \rangle_{T_{\gamma}\Gamma}$ for any $v \in V_0(X)$. Furthermore, by (3.2), if F is given by (3.1), we have for $\gamma \in \Gamma$, $x \in X$

$$(\nabla^{\Gamma} F)(\gamma; x) = \sum_{i=1}^{N} \frac{\partial g_F}{\partial s_i} (\langle \varphi_1, \gamma \rangle, \dots, \langle \varphi_N, \gamma \rangle) \nabla^X \varphi_i(x).$$
(3.5)

Definition 3.5 For a measurable vector field $V : \Gamma \to T\Gamma$ the divergence $\operatorname{div}_{\pi_{\sigma}}^{\Gamma} V$ is defined via the duality relation for all $F \in \mathcal{F}C_{b}^{\infty}(\mathcal{D},\Gamma)$ by

$$\int_{\Gamma} \langle V_{\gamma}, \nabla^{\Gamma} F(\gamma) \rangle_{T_{\gamma}\Gamma} d\pi_{\sigma}(\gamma) = -\int_{\Gamma} F(\gamma) (\operatorname{div}_{\pi_{\sigma}}^{\Gamma} V)(\gamma) d\pi_{\sigma}(\gamma), \qquad (3.6)$$

provided it exists (i.e., provided

$$F \mapsto \int_{\Gamma} \langle V_{\gamma}, \nabla^{\Gamma} F(\gamma) \rangle_{T_{\gamma}\Gamma} d\pi_{\sigma}(\gamma)$$

is continuous on $L^2(\pi_{\sigma})$).

Proposition 3.6 For any vector field V = Gv, where $G \in \mathcal{F}C_b^{\infty}(\mathcal{D}, \Gamma)$, $v \in V_0(X)$ we have

$$(\operatorname{div}_{\pi_{\sigma}}^{\Gamma}V)(\gamma) = \langle (\nabla^{\Gamma}G)(\gamma), v \rangle_{T_{\gamma}\Gamma} + G(\gamma)B_{v}^{\pi_{\sigma}}(\gamma).$$
(3.7)

For any $F, G \in \mathcal{F}C_b^{\infty}(\mathcal{D}, \Gamma)$ we introduce the *pre-Dirichlet form* which is generated by the intrinsic gradient ∇^{Γ} as

$$\mathcal{E}_{\pi_{\sigma}}^{\Gamma}(F,G) = \int_{\Gamma} \langle (\nabla^{\Gamma} F)(\gamma), (\nabla^{\Gamma} G)(\gamma) \rangle_{T_{\gamma}\Gamma} d\pi_{\sigma}(\gamma).$$
(3.8)

We will also need the *classical pre-Dirichlet* form for the intensity measure σ which is given as $\mathcal{E}^X_{\sigma}(\varphi, \psi) = \int_X \langle \nabla^X \varphi, \nabla^X \psi \rangle_{TX} d\sigma$ for any $\varphi, \psi \in \mathcal{D}$. This form is associated with the *Dirichlet operator* H^X_{σ} which is given on \mathcal{D} by $H^X_{\sigma}\varphi(x) := -\Delta^X\varphi(x) - \langle \beta^{\sigma}(x), \nabla^X\varphi(x) \rangle_{T_xX}$ and which satisfies $\mathcal{E}^X_{\sigma}(\varphi, \psi) = (H^X_{\sigma}\varphi, \psi)_{L^2(\sigma)}, \varphi, \psi \in \mathcal{D}$, see e.g. [BK95] and [MR92].

For any $F \in \mathcal{F}C_b^{\infty}(\mathcal{D},\Gamma)$, $(\nabla^{\Gamma}\nabla^{\Gamma}F)(\gamma,x,y) \in T_{\gamma}\Gamma \otimes T_{\gamma}\Gamma$ and we can define the Γ -Laplacian $(\Delta^{\Gamma}F)(\gamma) := \operatorname{Tr}(\nabla^{\Gamma}\nabla^{\Gamma}F)(\gamma) \in \mathcal{F}C_b^{\infty}(\mathcal{D},\Gamma)$. We introduce a differential operator in $L^2(\pi_{\sigma})$ on the domain $\mathcal{F}C_b^{\infty}(\mathcal{D},\Gamma)$ by the formula

$$(H^{\Gamma}_{\pi_{\sigma}}F)(\gamma) = -\Delta^{\Gamma}F(\gamma) - \langle \operatorname{div}_{\sigma}^{X}(\nabla^{\Gamma}F)(\gamma, \cdot), \gamma \rangle.$$
(3.9)

Theorem 3.7 The operator $H_{\pi_{\sigma}}^{\Gamma}$ is associated with the intrinsic Dirichlet form $\mathcal{E}_{\pi_{\sigma}}^{\Gamma}$, i.e.,

$$\mathcal{E}^{\Gamma}_{\pi_{\sigma}}(F,G) = (H^{\Gamma}_{\pi_{\sigma}}F,G)_{L^{2}(\pi_{\sigma})}, \qquad (3.10)$$

or $H_{\pi_{\sigma}}^{\Gamma} = -\operatorname{div}_{\pi_{\sigma}}^{\Gamma} \nabla^{\Gamma}$ on $\mathcal{F}C_{b}^{\infty}(\mathcal{D},\Gamma)$. We call $H_{\pi_{\sigma}}^{\Gamma}$ the intrinsic Dirichlet operator of the measure π_{σ} .

4 Extrinsic geometry on Poisson space

We recall the extrinsic geometry on $L^2(\pi_{\sigma})$ based on the isomorphism with the Fock space. Our approach is based on [KSS97] but we should also mention [BLL95], [IK88], [KSSU98], [NV95], [Pri95] for related considerations and references therein. For proofs of the results stated below in this section, we refer to [AKR98a, Sect. 5].

Let us define another "gradient" on functions $F : \Gamma \to \mathbb{R}$ This gradient ∇^P has specific useful properties on Poissonian spaces. We will call ∇^P the

Poissonian gradient. To this end we consider as the tangent space to Γ at any point $\gamma \in \Gamma$ the same space $L^2(\sigma)$ and define a mapping $\mathcal{F}C_b^{\infty}(\mathcal{D},\Gamma) \ni$ $F \mapsto \nabla^P F \in L^2(\pi_{\sigma}) \otimes L^2(\sigma)$ by

$$(\nabla^P F)(\gamma, x) := F(\gamma + \varepsilon_x) - F(\gamma), \ \gamma \in \Gamma, x \in X.$$
(4.1)

We stress that the transformation $\Gamma \ni \gamma \mapsto \gamma + \varepsilon_x \in \Gamma$ is π_{σ} -a.e. well-defined because $\pi_{\sigma}(\{\gamma \in \Gamma | x \in \gamma\}) = 0$ for any $x \in X$. The directional derivative is then defined as

$$(\nabla_{\varphi}^{P}F)(\gamma) = (\nabla^{P}F(\gamma, \cdot), \varphi)_{L^{2}(\sigma)} = \int_{X} [F(\gamma + \varepsilon_{x}) - F(\gamma)]\varphi(x)d\sigma(x), \ \varphi \in \mathcal{D}.$$
(4.2)

The Poissonian gradient ∇^P yields (via a corresponding "integration by parts" formula) an orthogonal system of Charlier polynomials on the Poisson space $(\Gamma, \mathcal{B}(\Gamma), \pi_{\sigma})$.

For any $n \in \mathbb{N}$ and all $\varphi \in \mathcal{D}$ we introduce a function in $L^2(\pi_{\sigma})$ by

$$Q_n^{\pi_\sigma}(\gamma;\varphi^{\otimes n}) := ((\nabla_{\varphi}^P)^{*n}1)(\gamma), \tag{4.3}$$

and define $Q_0^{\pi_{\sigma}} := 1$. Due to the kernel theorem [BK95, Chap. 1] these functions have the representation $Q_n^{\pi_{\sigma}}(\gamma; \varphi^{\otimes n}) = \langle Q_n^{\pi_{\sigma}}(\gamma), \varphi^{\otimes n} \rangle$, with generalized symmetric kernels $\Gamma \ni \gamma \mapsto Q_n^{\pi_{\sigma}}(\gamma) \in \operatorname{Exp}_n \mathcal{D}', n \in \mathbb{N}$. Here and below by $\operatorname{Exp}_n E$ we denote the *n*-th symmetric tensor power of a linear space *E*. Then for any smooth kernel $\varphi^{(n)} \in \operatorname{Exp}_n \mathcal{D}^{\otimes n}$ we introduce the function $Q_n^{\pi_{\sigma}}(\gamma; \varphi^{(n)}) := \langle Q_n^{\pi_{\sigma}}(\gamma), \varphi^{(n)} \rangle$ such that for all $\varphi^{(n)} \in \operatorname{Exp}_n \mathcal{D}^{\otimes n}$, $\psi^{(m)} \in \operatorname{Exp}_n \mathcal{D}^{\otimes m}$

$$\int_{\Gamma} Q_n^{\pi_{\sigma}}(\gamma;\varphi^{(n)}) Q_m^{\pi_{\sigma}}(\gamma;\psi^{(m)}) d\pi_{\sigma}(\gamma) = \delta_{nm} n! (\varphi^{(n)},\psi^{(m)})_{L^2(\sigma^{\otimes n})}.$$
(4.4)

Hence (4.3) extends to the case of kernels from the so-called *n*-particle Fock space $\operatorname{Exp}_n L^2(\sigma)$, $n \in \mathbb{N}$, and we set $\operatorname{Exp}_0 L^2(\sigma) := \mathbb{R}$.

As usual the symmetric Fock space over the Hilbert space $L^2(\sigma)$ is defined as $\operatorname{Exp} L^2(\sigma) := \bigoplus_{n=0}^{\infty} \operatorname{Exp}_n L^2(\sigma)$, see e.g. [BK95] and [HKPS93]. The square of the norm of a vector $(f^{(n)})_{n=0}^{\infty} \in \operatorname{Exp} L^2(\sigma)$ is given by $\sum_{n=0}^{\infty} n! \|f^{(n)}\|_{L^2(\sigma^{\otimes n})}^2$. For any $F \in L^2(\pi_{\sigma})$ there exists such a Fock vector, so that we have the following chaos decomposition

$$F(\gamma) = \sum_{n=0}^{\infty} Q_n^{\pi_\sigma}(\gamma; f^{(n)}), \qquad (4.5)$$

and moreover $||F||^2_{L^2(\pi_{\sigma})} = \sum_{n=0}^{\infty} n! ||f^{(n)}||^2_{L^2(\sigma^{\otimes n})}$. And vice versa, by (4.5) any Fock vector generates a function from $L^2(\pi_{\sigma})$. This produces an isomorphism between $L^2(\pi_{\sigma})$ and $\operatorname{Exp} L^2(\sigma)$.

Remark 4.1 In probability theory the functions $Q_n^{\pi_{\sigma}}(\gamma; f^{(n)})$ are called the *n*-multiple stochastic integrals of $f^{(n)}$ with respect to the compensated Poisson process generated by the Poisson measure π_{σ} , see e.g. [NV95].

There is an alternative approach to the chaos decomposition on the Poisson space which uses the concept of generalized Appell systems, see e.g. [KSS97].

The following proposition shows that the operators ∇_{φ}^{P} and ∇_{φ}^{P*} play the role of the annihilation resp. creation operators in the Fock space $\operatorname{Exp} L^{2}(\sigma)$.

Proposition 4.2 For all $\varphi, \psi \in D$, $n \in \mathbb{N}$ the following formulas hold

$$\nabla^P_{\psi} Q_n^{\pi_{\sigma}}(\gamma; \varphi^{\otimes n}) = n(\varphi, \psi)_{L^2(\sigma)} Q_{n-1}^{\pi_{\sigma}}(\gamma; \varphi^{\otimes (n-1)})$$
(4.6)

$$\nabla_{\psi}^{P*}Q_n^{\pi_{\sigma}}(\gamma;\varphi^{\otimes n}) = Q_{n+1}^{\pi_{\sigma}}(\gamma;\varphi^{\otimes n}\hat{\otimes}\psi), \ \gamma \in \Gamma,$$
(4.7)

where $\varphi^{\otimes n} \hat{\otimes} \psi$ means the symmetric tensor product of $\varphi^{\otimes n}$ and ψ .

Next we give an explicit expression for the adjoint of the Poissonian gradient ∇^{P*} .

Proposition 4.3 For any function $F \in L^1(\pi_{\sigma}) \otimes L^1(\sigma)$ we have $F \in D(\nabla^{P*})$ and the following equality holds

$$(\nabla^{P*}F)(\gamma) = \int_X F(\gamma - \varepsilon_x, x) d\gamma(x) - \int_X F(\gamma, x) d\sigma(x), \ \gamma \in \Gamma,$$
(4.8)

provided the right hand side of (4.8) is in $L^2(\pi_{\sigma})$.

Proof. For $X = \mathbb{R}^d$ this proposition was proved in [KSSU98]. Let $G \in \mathcal{F}C_b^{\infty}(\mathcal{D}, \Gamma)$ be given. Then an application of (4.1) gives

$$(\nabla^{P}G, F)_{L^{2}(\pi_{\sigma})\otimes L^{2}(\sigma)} = \int_{X} \int_{\Gamma} G(\gamma + \varepsilon_{x}) F(\gamma, x) d\pi_{\sigma}(\gamma) d\sigma(x) - \int_{X} \int_{\Gamma} G(\gamma) F(\gamma, x) d\pi_{\sigma}(\gamma) d\sigma(x).$$
(4.9)

Now we use the Mecke identity, see e.g., [Mec67, Satz 3.1]

$$\int_{\Gamma} \left(\int_{X} h(\gamma, x) d\gamma(x) \right) d\pi_{\sigma}(\gamma) = \int_{X} \int_{\Gamma} h(\gamma + \varepsilon_{x}, x) d\pi_{\sigma}(\gamma) d\sigma(x), \quad (4.10)$$

where h is any non-negative, $\mathcal{B}(\Gamma) \times \mathcal{B}(X)$ -measurable function. By (4.10) the right hand side of (4.9) transforms into

$$\int_{\Gamma} G(\gamma) \left[\int_{X} F(\gamma - \varepsilon_x, x) d\gamma(x) - \int_{X} F(\gamma, x) d\sigma(x) \right] d\pi_{\sigma}(\gamma),$$

which proves the proposition.

For any contraction B in $L^2(\sigma)$ it is possible to define an operator $\operatorname{Exp} B$ as a contraction in $\operatorname{Exp} L^2(\sigma)$ which in any *n*-particle subspace $\operatorname{Exp}_n L^2(\sigma)$ is given by $B \otimes \cdots \otimes B$ (*n* times). For any positive self-adjoint operator A in $L^2(\sigma)$ (with $\mathcal{D} \subset D(A)$) we have a contraction semigroup e^{-tA} , $t \ge 0$, hence it is possible to introduce the second quantization operator $d\operatorname{Exp} A$ as the generator of the semigroup $\operatorname{Exp}(e^{-tA})$, $t \ge 0$, i.e., $\operatorname{Exp}(e^{-tA}) = \operatorname{exp}(-td\operatorname{Exp} A)$, see e.g., [RS75]. We denote by H_A^P the image of the operator $d\operatorname{Exp} A$ in the Poisson space $L^2(\pi_{\sigma})$ under the described isomorphism.

Proposition 4.4 Let $\mathcal{D} \subset D(A)$. Then the symmetric bilinear form corresponding to the operator H_A^P has the following form

$$(H_A^P F, G)_{L^2(\pi_\sigma)} = \int_{\Gamma} (\nabla^P F(\gamma), A \nabla^P G(\gamma))_{L^2(\sigma)} d\pi_{\sigma}(\gamma), \ F, G \in \mathcal{F}C_b^{\infty}(\mathcal{D}, \Gamma).$$
(4.11)

The right hand side of (4.11) is called the "Poissonian pre-Dirichlet form" with coefficient operator A and is denoted by $\mathcal{E}_{\pi_{\sigma},A}^{P}$.

Let us consider the special case of the second quantization operator $d\operatorname{Exp} A$, where the one-particle operator A coincides with the Dirichlet operator H_{σ}^{X} generated by the measure σ on X. Then we have the following theorem which relates the intrinsic Dirichlet operator $H_{\pi_{\sigma}}^{\Gamma}$ and the operator $H_{H_{\sigma}}^{P}$.

Theorem 4.5 $H_{\pi_{\sigma}}^{\Gamma} = H_{H_{\sigma}}^{P}$ on $\mathcal{F}C_{b}^{\infty}(\mathcal{D},\Gamma)$. In particular, for all $F, G \in \mathcal{F}C_{b}^{\infty}(\mathcal{D},\Gamma)$

$$\int_{\Gamma} \langle \nabla^{\Gamma} F(\gamma), \nabla^{\Gamma} G(\gamma) \rangle_{T_{\gamma}\Gamma} d\pi_{\sigma}(\gamma)$$

$$= \int_{\Gamma} (\nabla^{P} F(\gamma), H_{\sigma}^{X} \nabla^{P} G(\gamma))_{L^{2}(\sigma)} d\pi_{\sigma}(\gamma)$$

$$= \int_{\Gamma} \int_{X} \langle \nabla^{X} \nabla^{P} F(\gamma, x), \nabla^{X} \nabla^{P} G(\gamma, x) \rangle_{T_{x}X} d\sigma(x) d\pi_{\sigma}(\gamma). \quad (4.12)$$

Remark 4.6 Theorem 4.5 gives full information about the spectrum of the intrinsic Dirichlet operator $H^{\Gamma}_{\pi_{\sigma}}$ on the Poisson space in terms of the underlying Dirichlet operator H^{X}_{σ} coming from the intensity measure σ . In the following section we obtain an analogue of Theorem 4.5 for the class of measures $\mathcal{G}^{1}_{ac}(\sigma, \Phi)$ (see (6.5) below) which is the aim of this paper.

5 Relation between intrinsic and extrinsic Dirichlet forms

Here we consider the class of measures $\mathcal{G}_{gc}^1(\sigma, \Phi)$ consisting of all $\mu \in \mathcal{G}_{gc}(\sigma, \Phi)$ such that

$$\int_{\Gamma} \gamma(K) d\mu(\gamma) < \infty \text{ for all compact } K \subset X.$$

We define for any $\mu \in \mathcal{G}^1_{qc}(\sigma, \Phi)$ the pre-Dirichlet form $\mathcal{E}^{\Gamma}_{\mu}$ by

$$\mathcal{E}^{\Gamma}_{\mu}(F,G) := \int_{\Gamma} \langle \nabla^{\Gamma} F(\gamma), \nabla^{\Gamma} G(\gamma) \rangle_{T_{\gamma}\Gamma} d\mu(\gamma), \ F, G \in \mathcal{F}C^{\infty}_{b}(\mathcal{D},\Gamma).$$
(5.1)

After all our preparations we are now going to prove an analogue of (4.12) for $\mu \in \mathcal{G}_{gc}^1(\sigma, \Phi)$. We would like to emphasize that the corresponding formula (5.2) is not obtained from (4.12) by just replacing π_{σ} by $\mu \in \mathcal{G}_{gc}^1(\sigma, \Phi)$. The essential difference is, in addition, an extra factor involving the conditional energy E_{Λ}^{Φ} . **Theorem 5.1** For any $\mu \in \mathcal{G}^1_{gc}(\sigma, \Phi)$, we have for all $F, G \in \mathcal{F}C^{\infty}_b(\mathcal{D}, \Gamma)$

$$\mathcal{E}^{\Gamma}_{\mu}(F,G) = \int_{\Gamma} \langle \nabla^{\Gamma} F(\gamma), \nabla^{\Gamma} G(\gamma) \rangle_{T_{\gamma}\Gamma} d\mu(\gamma)$$

$$= \int_{\Gamma} \int_{X} \langle \nabla^{X} \nabla^{P} F(\gamma,x), \nabla^{X} \nabla^{P} G(\gamma,x) \rangle_{T_{xX}} e^{-E^{\Phi}_{\{x\}}(\gamma+\varepsilon_{x})} d\sigma(x) d\mu(\gamma)$$
(5.2)

Proof. Let us take any $F \in \mathcal{F}C_b^{\infty}(\mathcal{D}, \Gamma)$ of the form (3.1). Then given $\gamma \in \Gamma$ and $x \in X$ (4.1) implies that

$$\nabla^{X} \nabla^{P} F(\gamma, x) = \nabla^{X} F(\gamma + \varepsilon_{x})$$

=
$$\sum_{i=1}^{N} \frac{\partial g_{F}}{\partial s_{i}} (\langle \varphi_{1}, \gamma \rangle + \varphi_{1}(x), \dots, \langle \varphi_{N}, \gamma \rangle + \varphi_{N}(x)) \nabla^{X} \varphi_{i}(x).$$

Let us define $\hat{F}_i(\gamma) := \frac{\partial g_F}{\partial s_i}(\langle \varphi_1, \gamma \rangle, \dots, \langle \varphi_N, \gamma \rangle), i = 1, \dots, N$. Obviously, it is enough to prove the equality (5.2) for F = G. Thus, inserting the result of $\nabla^X \nabla^P F(\gamma, x)$ into the right hand side of (5.2) we obtain

$$\int_{\Gamma} \int_{X} \sum_{i,j=1}^{N} \langle \nabla^{X} \varphi_{i}(x), \nabla^{X} \varphi_{j}(x) \rangle_{T_{xX}} \hat{F}_{i}(\gamma + \varepsilon_{x}) \hat{F}_{j}(\gamma + \varepsilon_{x}) e^{-E_{\{x\}}^{\Phi}(\gamma + \varepsilon_{x})} d\sigma(x) d\mu(\gamma).$$
(5.3)

Then we need the following useful proposition which generalizes the Mecke identity to measures in $\mathcal{G}_{gc}(\sigma, \Phi)$, see [NZ79], [MMW79].

Proposition 5.2 Let $h : \Gamma \times X \to \mathbb{R}_+$ be $\mathcal{B}(\Gamma) \times \mathcal{B}(X)$ -measurable, and let $\mu \in \mathcal{G}_{gc}(\sigma, \Phi)$. Then we have

$$\int_{\Gamma} \left(\int_{X} h(\gamma, x) d\gamma(x) \right) d\mu(\gamma) = \int_{X} \int_{\Gamma} h(\gamma + \varepsilon_x, x) e^{-E_{\{x\}}^{\Phi}(\gamma + \varepsilon_x)} d\mu(\gamma) d\sigma(x).$$
(5.4)

Using this proposition we transform (5.3) into

$$\int_{\Gamma} \sum_{i,j=1}^{N} \hat{F}_i(\gamma) \hat{F}_j(\gamma) \langle \langle \nabla^X \varphi_i(\cdot), \nabla^X \varphi_j(\cdot) \rangle_{TX}, \gamma \rangle d\mu(\gamma).$$

On the other hand using (3.5) we obtain

$$\langle \nabla^{\Gamma} F(\gamma), \nabla^{\Gamma} G(\gamma) \rangle_{T\Gamma} = \sum_{i,j=1}^{N} \hat{F}_{i}(\gamma) \hat{F}_{j}(\gamma) \langle \langle \nabla^{X} \varphi_{i}(\cdot), \nabla^{X} \varphi_{j}(\cdot) \rangle_{TX}, \gamma \rangle.$$

Therefore the equality on the dense $\mathcal{F}C_b^{\infty}(\mathcal{D},\Gamma)$ is valid which proves the theorem.

Remark 5.3 For the reader's convenience let us give a heuristic proof of the Nguyen-Zessin characterization of Gibbs measures in (5.4) which really is a consequence of the Mecke identity (cf. (4.10)). Indeed, let us write (heuristically)

$$d\mu(\gamma) = \frac{1}{Z^{\sigma,\Phi}} e^{-E^{\Phi}(\gamma)} d\pi_{\sigma}(\gamma).$$

Then the function $E_{\{x\}}^{\Phi}(\gamma + \varepsilon_x) = E^{\Phi}(\gamma + \varepsilon_x) - E^{\Phi}(\gamma)$ informally is the variation of the potential energy $E^{\Phi}(\gamma)$ when we add to the configuration γ an additional point $x \in X$. Using this representation we have

$$\int_{X} \int_{\Gamma} h(\gamma + \varepsilon_{x}, x) e^{-E_{\{x\}}^{\Phi}(\gamma + \varepsilon_{x})} d\mu(\gamma) d\sigma(x)$$

= $(Z^{\sigma, \Phi})^{-1} \int_{X} \int_{\Gamma} h(\gamma + \varepsilon_{x}, x) e^{-E_{\{x\}}^{\Phi}(\gamma + \varepsilon_{x})} d\pi_{\sigma}(\gamma) d\sigma(x)$

Then we use the Mecke identity to transform the right hand side of the above equality into

$$(Z^{\sigma,\Phi})^{-1} \int_{\Gamma} \left(\int_{X} h(\gamma, x) e^{-E^{\Phi}(\gamma)} d\gamma(x) \right) d\pi_{\sigma}(\gamma) = \int_{\Gamma} \left(\int_{X} h(\gamma, x) d\gamma(x) \right) d\mu(\gamma).$$

The rigorous proof in [NZ79] is obtained as a formalization of the heuristic computations above.

6 Closability of intrinsic Dirichlet forms

In this section we will prove the closability of the intrinsic Dirichlet form $(\mathcal{E}^{\Gamma}_{\mu}, \mathcal{F}C^{\infty}_{b}(\mathcal{D}, \Gamma))$ on $L^{2}(\mu) := L^{2}(\Gamma, \mu)$ for all $\mu \in \mathcal{G}^{1}_{gc}(\sigma, \Phi)$, using the integral representation (5.2) in Theorem 5.1. The closability of $(\mathcal{E}^{\Gamma}_{\mu}, \mathcal{F}C^{\infty}_{b}(\mathcal{D}, \Gamma))$ over Γ is implied by the closability of an appropriate family of pre-Dirichlet forms over X. Let us describe this more precisely.

We define new intensity measures on X by $d\sigma_{\gamma}(x) := \rho_{\gamma}(x) dm(x)$, where

$$\rho_{\gamma}(x) := e^{-E_{\{x\}}^{\Phi}(\gamma + \varepsilon_x)} \rho(x), \ x \in X, \gamma \in \Gamma$$
(6.1)

It was shown in [AR90, Theorem 5.3] (in the case $X = \mathbb{R}^d$) that the components of the Dirichlet form $(\mathcal{E}_{\sigma_{\gamma}}^X, \mathcal{D}^{\sigma_{\gamma}})$ corresponding to the measure σ_{γ} are closable on $L^2(\mathbb{R}^d, \sigma_{\gamma})$ if and only if σ_{γ} is absolutely continuous with respect to Lebesgue measure on \mathbb{R}^d and the Radon-Nikodym derivative satisfies some condition, see (6.2) below for details. This result allows us to prove the closability of $(\mathcal{E}_{\mu}^{\Gamma}, \mathcal{F}C_b^{\infty}(\mathcal{D}, \Gamma))$ on $L^2(\mu)$. Let us first recall the above mentioned result.

Theorem 6.1 (cf. [AR90, Theorem 5.3]) Let ν by a probability measure on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d), d \in \mathbb{N} \text{ and let } \mathcal{D}^{\nu} \text{ denote the } \nu\text{-classes determined by } \mathcal{D}$. Then the forms $(\mathcal{E}_{\nu,i}, \mathcal{D}^{\nu})$ defined by

$$\mathcal{E}_{\nu,i}(u,v) := \int_{\mathbb{R}^d} \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_i} d\nu, \ u,v \in \mathcal{D},$$

are well-defined and closable on $L^2(\mathbb{R}^d, \nu)$ for $1 \leq i \leq d$ if and only if ν is absolutely continuous with respect to Lebesgue measure λ^d on \mathbb{R}^d , and the Radon-Nikodym derivative $\rho = d\nu/d\lambda^d$ satisfies the condition:

for any $1 \le i \le d$ and λ^{d-1} -a.e. $x \in \left\{ y \in \mathbb{R}^{d-1} | \int_{\mathbb{R}} \rho_y^{(i)}(s) d\lambda^1(s) > 0 \right\},\$ $\rho_x^{(i)} = 0 \ \lambda^1$ -a.e. on $\mathbb{R} \setminus R(\rho_x^{(i)})$, where $\rho_x^{(i)}(s) := \rho(x_1, \dots, x_{i-1}, s, x_i, \dots, x_d),\$ $s \in \mathbb{R}, \text{ if } x = (x_1, \dots, x_{d-1}) \in \mathbb{R}^{d-1}, \text{ and where}$ (6.2)

$$R(\rho_x^{(i)}) := \left\{ t \in \mathbb{R} | \int_{t-\varepsilon}^{t+\varepsilon} \frac{1}{\rho_x^{(i)}(s)} ds < \infty \text{ for some } \varepsilon > 0 \right\}.$$
(6.3)

There is an obvious generalization of Theorem 6.1 to the case where a Riemannian manifold X is replacing \mathbb{R}^d , to be formulated in terms of local charts. Since here we are only interested in the "if part" of Theorem 6.1, we now recall a slightly weaker sufficient condition for closability in the general case where X is a manifold as before.

Theorem 6.2 Suppose $\sigma_1 = \rho_1 \cdot m$, where $\rho_1 : X \to \mathbb{R}_+$ is $\mathcal{B}(X)$ -measurable such that

$$\rho_1 = 0 \ m\text{-a.e. on } X \setminus \left\{ x \in X | \int_{\Lambda_x} \frac{1}{\rho_1} dm < \infty \text{ for some open neighbourhood } \Lambda_x \text{ of } x \right\}$$

$$(6.4)$$

Then $(\mathcal{E}_{\sigma_1}^X, \mathcal{D}^{\sigma_1})$ defined by

$$\mathcal{E}_{\sigma_1}^X(u,v) := \int_X \langle \nabla^X u(x), \nabla^X v(x) \rangle_{T_x X} \, d\sigma_1(x); \ u, v \in \mathcal{D},$$

is closable on $L^2(\sigma_1)$.

The proof is a straightforward adaptation of the line of arguments in [MR92, Chap. II, Subsection 2a] (see also [ABR89, Theorem 4.2] for details). We emphasize that (6.4) e.g. always holds, if ρ_1 is lower semicontinuous, and that neither ν in Theorem 6.1 nore σ_1 in Theorem 6.2 is required to have full support, so e.g. ρ_1 is not necessarily strictly positive *m*-a.e. on *X*.

We are now ready to prove the closability of $(\mathcal{E}^{\Gamma}_{\mu}, \mathcal{F}C^{\infty}_{b}(\mathcal{D}, \Gamma))$ on $L^{2}(\mu)$ under the above assumption.

Theorem 6.3 Let $\mu \in \mathcal{G}_{gc}^1(\sigma, \Phi)$. Suppose that for μ -a.e. $\gamma \in \Gamma$ the function ρ_{γ} defined in (6.1) satisfies (6.4) (resp. (6.2) in case $X = \mathbb{R}^d$). Then the form $(\mathcal{E}_{\mu}^{\Gamma}, \mathcal{F}C_b^{\infty}(\mathcal{D}, \Gamma))$ is closable on $L^2(\mu)$.

Proof. Let $(F_n)_{n\in\mathbb{N}}$ be a sequence in $\mathcal{F}C_b^{\infty}(\mathcal{D},\Gamma)$ such that $F_n \to 0, n \to \infty$ in $L^2(\mu)$ and

$$\mathcal{E}^{\Gamma}_{\mu}(F_n - F_m, F_n - F_m) \xrightarrow[n,m \to \infty]{} 0.$$
(6.5)

We have to show that

$$\mathcal{E}^{\Gamma}_{\mu}(F_{n_k}, F_{n_k}) \xrightarrow[k \to \infty]{} 0 \tag{6.6}$$

for some subsequence $(n_k)_{k\in\mathbb{N}}$. Let $(n_k)_{k\in\mathbb{N}}$ be a subsequence such that

$$\left(\int_{\Gamma} F_{n_k}^2 d\mu\right)^{1/2} + \mathcal{E}_{\mu}^{\Gamma} (F_{n_{k+1}} - F_{n_k}, F_{n_{k+1}} - F_{n_k})^{1/2} < \frac{1}{2^k} \text{ for all } k \in \mathbb{N}.$$

Then

$$\begin{split} \infty &> \sum_{k=1}^{\infty} \mathcal{E}_{\mu}^{\Gamma} (F_{n_{k+1}} - F_{n_k}, F_{n_{k+1}} - F_{n_k})^{1/2} \\ &\geq \sum_{k=1}^{\infty} \int_{\Gamma} \left(\int_{X} |\nabla^X \nabla^P (F_{n_{k+1}} - F_{n_k})(x, \gamma)|^2_{T_{xX}} e^{-E_{\{x\}}^{\Phi}(\gamma + \varepsilon_x)} d\sigma(x) \right)^{1/2} d\mu(\gamma) \\ &= \int_{\Gamma} \sum_{k=1}^{\infty} \left(\int_{X} |\nabla^X \nabla^P (F_{n_{k+1}} - F_{n_k})(x, \gamma)|^2_{T_{xX}} \rho_{\gamma}(x) dm(x) \right)^{1/2} d\mu(\gamma), \end{split}$$

where we used Theorem 5.1 and (6.1). From the last expression we obtain that

$$\sum_{k=1}^{\infty} \mathcal{E}_{\sigma_{\gamma}}^{X} (u_{n_{k+1}}^{(\gamma)} - u_{n_{k}}^{(\gamma)}, u_{n_{k+1}}^{(\gamma)} - u_{n_{k}}^{(\gamma)})^{1/2}$$

$$= \sum_{k=1}^{\infty} \left(\int_{X} |\nabla^{X} \nabla^{P} (F_{n_{k+1}} - F_{n_{k}})(x, \gamma)|^{2}_{T_{x}X} \rho_{\gamma}(x) dm(x) \right)^{1/2} < \infty \text{ for } \mu - \text{a.e. } \gamma \in \Gamma,$$
(6.7)

where for $k \in \mathbb{N}, \gamma \in \Gamma$,

$$u_{n_k}^{(\gamma)}(x) := F_{n_k}(\gamma + \varepsilon_x) - F_{n_k}(\gamma), \ x \in X.$$

Note that $u_{n_k}^{(\gamma)} \in \mathcal{D}$. (6.7) implies that for μ -a.e. $\gamma \in \Gamma$

$$\mathcal{E}^{X}_{\sigma_{\gamma}}(u_{n_{k}}^{(\gamma)}-u_{n_{l}}^{(\gamma)},u_{n_{k}}^{(\gamma)}-u_{n_{l}}^{(\gamma)}) \xrightarrow[k,l\to\infty]{} 0.$$
(6.8)

Let $\Lambda \subset \mathcal{O}_c(X)$.

Claim 1: For μ -a.e. $\gamma \in \Gamma$

$$\int_X (u_{n_k}^{(\gamma)}(x))^2 \mathbb{1}_{\Lambda}(x) d\sigma_{\gamma}(x) \xrightarrow[k \to \infty]{} 0.$$

To prove Claim 1 we first note that for μ -a.e. $\gamma \in \Gamma$

 $\sigma_{\gamma}(\Lambda) < \infty,$

as follows immediately from (5.4) (taking $h(\gamma, x) := \mathbb{1}_{\Lambda}(x)$ for $x \in X, \gamma \in \Gamma$), since $\mu \in \mathcal{G}^{1}_{gc}(\sigma, \Phi)$. Therefore, for μ -a.e. $\gamma \in \Gamma$

$$\int_{X} F_{n_k}^2(\gamma) \mathbb{1}_{\Lambda}(x) d\sigma_{\gamma}(x) = F_{n_k}^2(\gamma) \sigma_{\gamma}(\Lambda) \xrightarrow[k \to \infty]{} 0.$$
(6.9)

Furthermore, by (5.4)

$$\begin{split} &\int_{\Gamma} \int_{X} F_{n_{k}}^{2}(\gamma + \varepsilon_{x}) \mathbb{1}_{\Lambda}(x) d\sigma_{\gamma}(x) (1 + \gamma(\Lambda))^{-1} d\mu(\gamma) \\ &= \int_{\Gamma} F_{n_{k}}^{2}(\gamma) \int_{X} \frac{\mathbb{1}_{\Lambda}(x)}{1 + \gamma(\Lambda) - \mathbb{1}_{\Lambda}(x)} \gamma(dx) d\mu(\gamma) \\ &\leq \int_{\Gamma} F_{n_{k}}^{2}(\gamma) d\mu(\gamma) < \frac{1}{2^{k}}, \end{split}$$

because the integral w.r.t. γ is dominated by 1 for all $\gamma \in \Gamma$. Hence

$$\infty > \sum_{k=1}^{\infty} \left(\int_{\Gamma} \int_{X} F_{n_{k}}^{2}(\gamma + \varepsilon_{x}) \mathbb{1}_{\Lambda}(x) d\sigma_{\gamma}(x) (1 + \gamma(\Lambda))^{-1} d\mu(\gamma) \right)^{1/2}$$

$$\geq \int_{\Gamma} \sum_{k=1}^{\infty} \left(\int_{X} F_{n_{k}}^{2}(\gamma + \varepsilon_{x}) \mathbb{1}_{\Lambda}(x) d\sigma_{\gamma}(x) \right)^{1/2} (1 + \gamma(\Lambda))^{-1} d\mu(\gamma).$$

Therefore, for μ -a.e. $\gamma \in \Gamma$

$$\int_{X} F_{n_k}^2(\gamma + \varepsilon_x) \mathbb{1}_{\Lambda}(x) d\sigma_{\gamma}(x) \xrightarrow[k \to \infty]{} 0.$$
(6.10)

Then Claim 1 follows by (6.9) and (6.10).

Claim 2: For μ -a.e. $\gamma \in \Gamma$

$$|\nabla^X u_{n_k}^{(\gamma)}|_{TX} \xrightarrow[k \to \infty]{} 0 \quad \sigma_\gamma - \text{a.e.}$$

To prove Claim 2 we first note that clearly (6.7) implies that for μ -a.e. $\gamma \in \Gamma$

$$\mathcal{E}^{X}_{\mathbb{1}_{\Lambda}\sigma_{\gamma}}(u_{n_{k}}^{(\gamma)}-u_{n_{l}}^{(\gamma)},u_{n_{k}}^{(\gamma)}-u_{n_{l}}^{(\gamma)}) \xrightarrow[k,l\to\infty]{} 0.$$
(6.11)

Hence we can apply Theorem 6.2 (resp. 6.1) to $\rho_1 := \mathbb{1}_{\Lambda} \rho_{\gamma}$ and conclude by Claim 1 and (6.11) that for μ -a.e. $\gamma \in \Gamma$

$$\mathcal{E}^{X}_{\mathbb{1}_{\Lambda}\sigma_{\gamma}}(u_{n_{k}}^{(\gamma)},u_{n_{k}}^{(\gamma)}) \xrightarrow[k \to \infty]{} 0,$$

hence by (6.7)

$$1\!\!1_{\Lambda} | \nabla^X u_{n_k}^{(\gamma)} |_{TX} \xrightarrow[k \to \infty]{} 0 \quad \sigma_{\gamma} - \text{a.e.}$$

Since Λ was arbitrary, Claim 2 is proven.

From Claim 2 we now easily deduce (6.6) by (5.2) and Fatou's Lemma as follows:

$$\begin{aligned} \mathcal{E}^{\Gamma}_{\mu}(F_{n_k}, F_{n_k}) &\leq \int_{\Gamma} \liminf_{l \to \infty} \int_{X} |\nabla^X (u_{n_k}^{(\gamma)} - u_{n_l}^{(\gamma)})|_{TX}^2 \, d\sigma_{\gamma}(x) d\mu(\gamma) \\ &\leq \liminf_{l \to \infty} \mathcal{E}^{\Gamma}_{\mu}(F_{n_k} - F_{n_l}, F_{n_k} - F_{n_l}), \end{aligned}$$

which by (6.5) can be made arbitrarily small for k large enough.

Remark 6.4 The above method to prove closability of pre-Dirichlet forms on configuration spaces Γ_X extends immediately to the case where X is replaced by an infinite dimensional "manifold" such as the loop space (cf. [MR97]).

Example 6.5 Let $X = \mathbb{R}^d$ with the Euclidean metric and $\sigma := z \cdot m$, $z \in (0, \infty)$. A pair potential is a $\mathcal{B}(\mathbb{R}^d)$ -measurable function $\phi : \mathbb{R}^d \to \mathbb{R} \cup \{\infty\}$ such that $\phi(-x) = \phi(x)$. Any pair potential ϕ defines a potential $\Phi = \Phi_{\phi}$ in the sense of Section 2 as follows: we set $\Phi(\gamma) := 0$, $|\gamma| \neq 2$ and $\Phi(\gamma) := \phi(x - y)$ for $\gamma = \{x, y\} \subset \mathbb{R}^d$. For such pair potentials ϕ the condition in Theorem 6.3 ensuring closability of $(\mathcal{E}_{\mu}^{\Gamma}, \mathcal{F}C_{b}^{\infty}(\mathcal{D}, \Gamma))$ on $L^2(\mu)$ for $\mu \in \mathcal{G}_{gc}^1(\sigma, \Phi)$ can be now easily formulated as follows: for μ -a.e. $\gamma \in \Gamma$ and m-a.e. $x \in \{y \in \mathbb{R}^d | \sum_{y' \in \gamma \setminus \{y\}} |\phi(y - y')| < \infty\}$ it holds that $\int_{V_x} e^{\sum_{y' \in \gamma \setminus \{y\}} \phi(y - y')} m(dy) < \infty$ for some open neighbourhood V_x of x. This condition trivially holds e.g. if supp ϕ is compact, $\{\phi < \infty\}$ is open, and $\phi^+ \in L^{\infty}_{loc}(\{\phi < \infty\}; m)$. If even $\mu \in \mathcal{G}_{gc}^1(z, \phi)$ and ϕ satisfies the assumptions in Proposition 7.1, then it suffices to merely assume that $\{\phi < \infty\}$ is open and $\phi^+ \in L^{\infty}_{loc}(\{\phi < \infty\}; m)$. This follows by an elementary consideration.

- **Remark 6.6** 1. We emphasize that Example 6.5 generalizes the closability result in [Osa96], though an a-priori bigger domain for $\mathcal{E}_{\mu}^{\Gamma}$ is considered there. However, Theorems 6.1-6.3 are also valid for this bigger domain. The proofs are exactly the same.
 - 2. We also like to emphasize that similarly to Example 6.5 one proves the closability of $(\mathcal{E}^{\Gamma}_{\mu}, \mathcal{F}C^{\infty}_{b}(\mathcal{D}, \Gamma))$ (or with a larger domain in [Osa96]) on $L^{2}(\mu)$ for $\mu \in \mathcal{G}^{1}_{gc}(\sigma, \Phi)$ in the case of multi-body potentials ϕ .

Acknowledgments

We would like to thank Tobias Kuna for helpful discussions. Financial support of the INTAS-Project 378, PRAXIS Programme through CITMA, Funchal, and TMR Nr. ERB4001GT957046 are gratefully acknowledged.

References

- [ABR89] S. Albeverio, J. Brasche, and M. Röckner. Dirichlet forms and generalized Schrödinger operators. In H. Holden and A. Jensen, editors, *Schrödinger Operators*, volume 345 of *Lecture Notes in Phys.*, pages 1–42, Berlin, 1989. Springer-Verlag.
- [AKR96a] S. Albeverio, Yu. G. Kondratiev, and M. Röckner. Canonical Dirichlet operator and distorted Brownian motion on Poisson spaces. C. R. Acad. Sci., Paris Sér. I Math., 323:1179–1184, 1996.
- [AKR96b] S. Albeverio, Yu. G. Kondratiev, and M. Röckner. Differential geometry of Poisson spaces. C. R. Acad. Sci. Paris Sér. I Math., 323:1129–1134, 1996.
- [AKR98a] S. Albeverio, Yu. G. Kondratiev, and M. Röckner. Analysis and geometry on configuration spaces. J. Funct. Anal., 154:444–500, 1998.
- [AKR98b] S. Albeverio, Yu. G. Kondratiev, and M. Röckner. Analysis and geometry on configuration spaces: The Gibbsian case. J. Funct. Anal., 157:242–291, 1998.
- [AKR99] S. Albeverio, Yu. G. Kondratiev, and M. Röckner. Diffeomorphism groups and current algebras: configuration spaces analysis in quantum theory. *Rev. Math. Phys.*, 11(97-073):1–23, 1999.
- [AR90] S. Albeverio and M. Röckner. Classical Dirichlet forms on topological vector spaces - closability and a Cameron-Martin formula. J. Func. Anal., 88:395–436, 1990.
- [BK95] Yu. M. Berezansky and Yu. G. Kondratiev. Spectral Methods in Infinite-Dimensional Analysis. Kluwer Academic Publishers, Dordrecht, 1995.
- [BLL95] Yu. M. Berezansky, V. O. Livinsky, and E. W. Lytvynov. A generalization of Gaussian white noise analysis. *Methods of Functional Analysis and Topology*, 1(1):28–55, 1995.

- [Geo79] O. Georgii. Canonical Gibbs Measures, volume 760 of Lectures Notes in Math. Springer-Verlag, Berlin, 1979.
- [GV68] I. M. Gel'fand and N. Ya. Vilenkin. *Generalized Functions*, volume 4. Academic Press, New York and London, 1968.
- [HKPS93] T. Hida, H. H. Kuo, J. Potthoff, and L. Streit. White Noise. An Infinite Dimensional Calculus. Kluwer, Dordrecht, 1993.
- [IK88] Y. Ito and I. Kubo. Calculus on Gaussian and Poisson white noises. Nagoya Math. J., 111:41–84, 1988.
- [KSS97] Yu. G. Kondratiev, J. L. Silva, and L. Streit. Generalized Appell systems. Methods of Functional Analysis and Topology, 3(3):28– 61, 1997.
- [KSSU98] Yu. G. Kondratiev, J. L. Silva, L. Streit, and F. G. Us. Analysis on Poisson and Gamma spaces. *Infinite Dimensional Analysis*, *Quantum Probabilities and Related Topics*, 1(1):91–117, 1998.
- [Mec67] J. Mecke. Stationäre zufällige Maße auf lokalkompakten Abelschen Gruppen. Z. Wahrsch. verw. Gebiete, 9:36–58, 1967.
- [MMW79] K. Matthes, J. Mecke, and W. Warmuth. Bemerkungen zu einer Arbeit von Nguyen Xuan Xanh und Hans Zessin. Math. Nachr., 88:117–127, 1979.
- [MR92] Z.-M. Ma and M. Röckner. Introduction to the Theory of (Non-Symmetric) Dirichlet Forms. Springer-Verlag, Berlin, 1992.
- [MR97] Z.-M. Ma and M. Röckner. Construction of diffusions on configuration spaces. Preprint, 1997.
- [NV95] D. Nualart and J. Vives. A duality formula on the Poisson space and some applications. In Seminar on Stochastic Analysis, Random Fields and Applications, volume 36, pages 205–213, Basel, 1995. Birkhäuser.
- [NZ79] X. X. Nguyen and H. Zessin. Integral and differential characterizations of the Gibbs process. *Math. Nachr.*, 88:105–115, 1979.

- [Osa96] H. Osada. Dirichlet form approach to infinite-dimensional Wiener processes with singular integractions. *Comm. Math. Phys.*, 176(1):117–131, 1996.
- [Osa98] H. Osada. Interacting Brownian motions with measurable potentials. *Japan Academy*, 74, Ser. A(1), 1998.
- [Pre76] C. J. Preston. Random Fields, volume 534 of Lectures Notes in Math. Springer-Verlag, Berlin Heidelberg New York, 1976.
- [Pre79] J. C. Preston. Canonical and microcanonical Gibbs states. Z. Wahrsch. verw. Gebiete, 46:125–158, 1979.
- [Pri95] N. Privault. A transfer principle from Wiener to Poisson space and applications. J. Funct. Anal., 132:335–360, 1995.
- [RS75] M. Reed and B. Simon. *Methods of Modern Mathematical Physics*. Academic Press, New York and London, 1975.
- [Shi94] H. Shimomura. Poisson measures on the configuration space and unitary representations of the group of diffeomorphism. J. Math. Kyoto Univ., 34:599–614, 1994.