

INHOMOGENEOUS MULTIPARAMETER JORDANIAN QUANTUM GROUPS BY CONTRACTION

ROGER J. McDERMOTT AND DEEPAK PARASHAR

*School of Computer and Mathematical Sciences, The Robert Gordon University,
St. Andrew Street, Aberdeen AB25 1HG, U.K.
(e-mail: rm@scms.rgu.ac.uk, deeps@scms.rgu.ac.uk)*

Received XXX

It is known that the inhomogeneous quantum group $IGL_{q,r}(2)$ can be constructed as a quotient of the multiparameter q -deformation of $GL(3)$. We show that a similar result holds for the inhomogeneous Jordanian deformation and exhibit its Hopf structure.

1 Introduction

It is well-known [1] that, analogous to the classical group-theoretical method, the q -deformation of $IGL(2)$ can be constructed by factoring out a certain two-sided Hopf ideal from the multiparameter q -deformation of $GL(3)$. This is an interesting procedure, allowing, for example, the construction of a differential calculus on the quantum plane by a reduction of the differential calculus on the quantum group. In this paper, we apply the same construction to the Jordanian deformation. The multiparameter Jordanian deformation of $GL(3)$ is first produced by a contraction from the corresponding q -deformation and this is then used to construct the inhomogeneous group by factorisation. The Hopf-structure of $IGL_J(2)$ is given explicitly and we show that it is possible to derive from this a coaction of a modified version of $GL_J(2)$ on the Jordanian quantum plane.

Note: In this paper, we denote q -deformed structures using the (multiparameter) subscript Q and structures that have been contracted to the Jordanian form are written with a subscript J (e.g $GL_Q(3)$ and $GL_J(3)$).

2 The R-matrix for $GL_Q(2)$

Following Aschieri and Castellani [1], the R -matrix for $GL_Q(3)$ (where $Q = \{r, s, p, q\}$) can be written as

$$R_Q(3) = \begin{pmatrix} r & & & & \\ & S^{-1} & & & \\ & \Lambda & S & & \\ & & & R_Q(2) & \\ & & & & \end{pmatrix} \quad (1)$$

where $S = \begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix}$, $\Lambda = \begin{pmatrix} r - r^{-1} & 0 \\ 0 & r - r^{-1} \end{pmatrix}$ and

$$R_Q(2) = \begin{pmatrix} r & & & \\ & s & & \\ & r - r^{-1} & s^{-1} & \\ & & & r \end{pmatrix} \quad (2)$$

The matrix indices of $R_Q(3)$ run, in order, through the set (11), (12), (13), (21), (31), (22), (23), (32), (33). This numbering system is chosen to clearly show the embedding of the $R_Q(2)$ matrix in the $R_Q(3)$ matrix which, in turn, allows the Hopf structure of larger algebra to be analysed in terms of the simpler one. The Hopf structure of $GL_Q(3)$ is given by the RTT relations with T -matrix

$$\mathcal{T} = \begin{pmatrix} f & \theta & \phi \\ x & a & b \\ y & c & d \end{pmatrix} \quad (3)$$

and the multiparameter inhomogeneous q -deformation $IGL_Q(2)$ is the quantum homogeneous space

$$IGL_Q(2) = GL_Q(3)/H \quad (4)$$

where H is the two-sided Hopf ideal generated by the T -matrix elements $\{\theta, \phi\}$.

3 The Contraction Procedure

The R -matrix of the Jordanian (or h -deformation) can be viewed as a singular limit of a similarity transformation on the q -deformation R -matrix [2][3]. Let $g(\eta)$ be a matrix dependent on a contraction parameter η which is itself a function of one of the deformation parameters of the q -deformed algebra. This can be used to define a transformed q -deformed R -matrix

$$\tilde{R}_J = (g^{-1} \otimes g^{-1})R_Q(g \otimes g) \quad (5)$$

The R -matrix of the Jordanian deformation is then obtained by taking a limiting value of the parameter η . Even though the contraction parameter η is undefined in this limit, the new R -matrix is finite and gives rise to a new quantum group structure through the RTT -relations. For example, in the contraction process which takes $GL_q(2)$ to $GL_h(2)$, the contraction matrix is

$$g(\eta) = \begin{pmatrix} 1 & 0 \\ \eta & 1 \end{pmatrix} \quad (6)$$

where $\eta = \frac{h}{1-q}$ with h a new free parameter.

It has been shown by Alishahiha [3] that, in the extension of this procedure to the construction of $GL_J(3)$, there are essentially two choices of contraction matrix.

The first has been used in a number of papers, e.g. by Quesne [4] and takes the form

$$G' = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \eta & 0 & 1 \end{pmatrix} \quad (7)$$

There is, however, a second choice (also mentioned in [3] but not pursued there since it gives trivial results for the single-parameter q -deformation)

$$G = \begin{pmatrix} 1 & 0 \\ 0 & g \end{pmatrix} \quad (8)$$

where g is the 2×2 contraction matrix

$$g(\eta) = \begin{pmatrix} 1 & 0 \\ \eta & 1 \end{pmatrix} \quad (9)$$

with $\eta = \frac{r}{1-q}$. In this present work, we take G as our contraction matrix because, unlike the matrix G' , after contraction it allows a non-trivial embedding of $R_J(2)$ in $R_J(3)$ in a manner similar to the q -deformed case. It is then possible to perform the quotient construction for the inhomogeneous quantum group.

If the similarity transformation is made using the matrix G , we obtain

$$R_J(3) = \lim_{r \rightarrow 1} \begin{pmatrix} r & & & & \\ & g^{-1}S^{-1}g & & & \\ & \Lambda & & g^{-1}Sg & \\ & & & & (g^{-1} \otimes g^{-1})R_Q(g \otimes g) \end{pmatrix} \quad (10)$$

$$= \begin{pmatrix} 1 & & & & \\ & K^{-1} & & & \\ & & K & & \\ & & & & R_J(2) \end{pmatrix} \quad (11)$$

where K is the matrix $\begin{pmatrix} p & 0 \\ k & p \end{pmatrix}$ and $R_J(2)$ is the R -matrix for the multiparameter Jordanian deformation of $GL(2)$

$$\begin{pmatrix} 1 & & & & \\ m & 1 & & & \\ -m & 0 & 1 & & \\ mn & n & -n & 1 & \end{pmatrix} \quad (12)$$

The free parameters $\{m, n, k\}$ appear as limits in the contraction process while the parameter $\{p\}$ survives the contraction process. The result is a four parameter Jordanian deformation of $GL(3)$.

4 Multiparameter Jordanian Deformation of $GL(3)$

We denote the T -matrix for the Jordanian deformation by

$$\mathcal{T} = \begin{pmatrix} f & \theta & \phi \\ x & a & b \\ y & c & d \end{pmatrix} = \begin{pmatrix} f & \Theta \\ X & T \end{pmatrix} \quad (13)$$

where $T = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, $X = \begin{pmatrix} x \\ y \end{pmatrix}$, and $\Theta = (\theta, \phi)$.

4.1 Algebra Relations

The algebra structure of the quantum group is obtained through the RTT -procedure. For the commutation relations between the elements of the matrix T , we have the usual relations between the generators of the multiparametric Jordanian deformation of $GL(2)$:

$$\begin{aligned} [a, b] &= nb^2 & [a, c] &= m(\delta - a^2) & [a, d] &= nbd - mba \\ [b, d] &= -mb^2 & [b, c] &= -mba - ndb & [c, d] &= n(d^2 - \delta) \end{aligned} \quad (14)$$

where δ is the quantum determinant of the submatrix T

$$\delta = ad - bc - nbd \quad (15)$$

with commutation relation

$$[\delta, a] = (m - n)\delta b \quad [\delta, b] = 0 \quad [\delta, c] = (m - n)(\delta d - a\delta) \quad [\delta, d] = (n - m)\delta b \quad (16)$$

Thus δ is not central in the (sub-Hopf) algebra generated by elements of T unless $m = n$.

The relations between T and f are given by

$$[a, f] = \frac{k}{p}fb \quad [b, f] = 0 \quad [c, f] = \frac{k}{p}(fd - af) \quad [d, f] = -\frac{k}{p}bf \quad (17)$$

those between T and X are

$$\begin{aligned} [a, x]_p &= kxb & [b, x]_p &= 0 \\ [c, x]_p &= kxd + max & [d, x]_p &= mbx \\ [a, y]_p &= kyb - max & [b, y]_p &= -mbx \\ [c, y]_p &= kyd + ncx - nay - mna & [d, y]_p &= ndx - nby - mnbx \\ \delta x &= p^2x\delta & \delta y &= p^2y\delta + (n - m)\delta x \end{aligned} \quad (18)$$

while those between f and X give

$$[f, x]_p = 0 \quad [f, y]_p = -kxf \quad (19)$$

The commutation relations between the elements of X are the usual relations for the Jordanian quantum plane $C_J(2)$:

$$[x, y] = -mx^2 \tag{20}$$

There are also similar commutation relations between the elements of T , f and Θ , as well as cross-relations between X and Θ .

4.2 Coalgebra Relations and Antipode

The coalgebraic structure of the Hopf algebra is the usual one:

$$\Delta(\mathcal{T}) = \mathcal{T} \dot{\otimes} \mathcal{T} \quad \epsilon(\mathcal{T}) = I_3 \tag{21}$$

with antipode

$$S(\mathcal{T}) = \begin{pmatrix} e & -e\Theta T^{-1} \\ -T^{-1}Xe & T^{-1}Xe\Theta T^{-1} + T^{-1} \end{pmatrix} \tag{22}$$

where we append to the algebra, the element $e = (f - \Theta T^{-1}X)^{-1}$. In terms of these elements, the quantum determinant of the T -matrix \mathcal{T} is

$$\mathcal{D} = \det(\mathcal{T}) = e^{-1}\delta \tag{23}$$

and so, in the usual way, we can add $\xi = \mathcal{D}^{-1}$ to the algebra to obtain the full Hopf algebra.

5 The Inhomogeneous Multiparameter Jordanian Quantum Group $IGL_J(2)$

We define H to be the space of all monomials containing at least one element of Θ . It is straightforward to prove the following :

1. H is a two-sided ideal in $GL_J(3)$.
2. H is a co-ideal i.e. $\Delta(H) \subseteq H \otimes GL_J(3) + GL_J(3) \otimes H$ and $\epsilon(H) = 0$.
3. $S(H) \subseteq H$.

Thus H is a two-sided Hopf ideal and so we can define a canonical projection from $GL_J(3)$ to the quotient space $GL_J(3)/H$ which respects the Hopf-algebraic structure (i.e. the RTT -relations). Consequently the quotient is a Hopf algebra which we denote $IGL_J(2)$.

The algebra sector for this quantum group has commutation relations formally obtained from $GL_J(3)$ by setting the generator set $\Theta = 0$ and this gives rise to the commutation relations explicitly detailed in the previous section. The T -matrix for the coalgebra is given by

$$\mathcal{T} = \begin{pmatrix} f & 0 \\ X & T \end{pmatrix} \tag{24}$$

which gives the coproduct

$$\Delta(\mathcal{T}) = \mathcal{T} \dot{\otimes} \mathcal{T} = \begin{pmatrix} f \otimes f & 0 \\ T \dot{\otimes} X + X \dot{\otimes} f & T \dot{\otimes} T \end{pmatrix} \quad (25)$$

counit $\epsilon(\mathcal{T}) = I_3$ and antipode

$$S(\mathcal{T}) = \begin{pmatrix} f^{-1} & 0 \\ -T^{-1} X f^{-1} & T^{-1} \end{pmatrix} \quad (26)$$

The quantum determinant $\mathcal{D} = f\delta$ is group-like but, since f is not central, it cannot be made simultaneously central with δ unless the whole algebraic structure collapses to a trivial extension of the single-parameter Jordanian deformation of $GL(2)$. This is analogous to the situation in the q -deformed case.

This procedure also shows that it is possible to view the Jordanian quantum plane, $C_J(2)$, as the quantum homogeneous space $IGL_J(2)/GL_J(2)^*$ where $GL_J(2)^*$ is the Hopf algebra formed by appropriately appending the “dilatation element” f to $GL_J(2)$. The comultiplication in $IGL_J(2)$ can then be viewed as a coaction of the quantum group $GL_J(2)^*$ on the quantum plane $C_J(2)$ generated by the elements X . However, unlike the usual case, there is a non-trivial braiding between the elements of the quantum group and quantum plane.

6 Conclusion

We have shown that it is possible to construct the inhomogeneous Jordanian deformation $IGL_J(2)$ as a quotient group by factoring out a Hopf ideal from $GL_J(3)$. It would be of interest to construct the differential calculus on the Jordanian quantum plane by a reduction of the bicovariant differential calculus on $GL_J(3)$. This would allow the investigation of physical models with $GL_J(N)$ symmetry similar to that of Cho *et al* [5] and Madore and Steinacker [6]. Work on this problem is underway.

References

- [1] P. Aschieri and L. Castellani: *Int. J. Mod. Phys.* **A11** (1994) 1019.
- [2] A. Aghomohammadi, M. Khorrami and A. Shariati: *J. Phys.* **A28** (1995) L225.
- [3] M. Alishahiha: *J. Phys.* **A28** (1995) 6187.
- [4] C. Quesne: [math/9810161](#), [math/9903151](#).
- [5] S. Cho, J. Madore, K. S. Park: [q-alg/9709007](#).
- [6] J. Madore, H. Steinacker: [math/9907023](#).