

THE QUANTUM DUALITY PRINCIPLE

FABIO GAVARINI

Università degli Studi di Roma “Tor Vergata” — Dipartimento di Matematica
Via della Ricerca Scientifica 1, I-00133 Roma — ITALY

ABSTRACT. The “quantum duality principle” states that the quantization of a Lie bialgebra — via a quantum universal enveloping algebra (QUEA) — provides also a quantization of the dual Lie bialgebra (through its associated formal Poisson group) — via a quantum formal series Hopf algebra (QFSHA) — and, conversely, a QFSHA associated to a Lie bialgebra (via its associated formal Poisson group) yields a QUEA for the dual Lie bialgebra as well; more precisely, there exist functors $\text{QUEA} \rightarrow \text{QFSHA}$ and $\text{QFSHA} \rightarrow \text{QUEA}$, inverse of each other, such that in either case the Lie bialgebra associated to the target object is the dual of that of the source object. Such a result was claimed true by Drinfel’d, but seems to be unproved in literature: we give here a complete detailed proof of it.

Introduction

The *quantum duality principle* is known in literature with at least two formulations. One claims that quantum function algebras associated to dual Poisson groups can be taken to be dual — in the sense of Hopf algebra duality — of each other; and similarly for quantum enveloping algebras (cf. [FRT] and [Se]). The second one, due to Drinfel’d, states that any quantization of the universal enveloping algebra of a Poisson group can also be “seen” — in a suitable sense — as a quantization of the dual formal Poisson group, and, conversely, any quantization of a formal Poisson group “works” also — in a suitable sense — as a quantization of the universal enveloping algebra of the dual Poisson group: this is the point of view we are interested in. Now we describe more in detail this result.

Let k be a fixed field of characteristic zero. Let $\mathfrak{g}$ be a Lie algebra over k , $U(\mathfrak{g})$ its universal enveloping algebra, and $F[[\mathfrak{g}]]$ the (algebra of regular functions on the) formal group associated to $\mathfrak{g}$: if G is an algebraic (or proalgebraic, if $\dim(\mathfrak{g}) = \infty$) group with tangent Lie algebra $\mathfrak{g}$, and $F[G]$ is the algebra of regular functions on G , then $F[[\mathfrak{g}]]$ can be realized either as the $\mathfrak{m}_e$ -completion of $F[G]$ at the maximal ideal $\mathfrak{m}_e$ of the identity element $e \in G$, or as $F[[\mathfrak{g}]] = U(\mathfrak{g})^*$ (so we do not need G to exist, for having $\mathfrak{g}$ is enough). Then $U(\mathfrak{g})$ is a Hopf algebra, whereas $F[[\mathfrak{g}]]$ is a *topological* Hopf algebra, for it has a

1991 *Mathematics Subject Classification*: Primary 17B37, 81R50.

coproduct which takes values in a suitable topological tensor product of $F[[\mathfrak{g}]]$ with itself; finally, there exist a natural pairing of (topological) Hopf algebras between $U(\mathfrak{g})$ and $F[[\mathfrak{g}]]$.

Now assume $\mathfrak{g}$ is a Lie bialgebra: then $U(\mathfrak{g})$ is a co-Poisson Hopf algebra, $F[[\mathfrak{g}]]$ is a topological Poisson Hopf algebra, and the Hopf pairing mentioned above is compatible with these additional co-Poisson and Poisson structures. Furthermore, the linear dual $\mathfrak{g}^*$ of $\mathfrak{g}$ is a Lie bialgebra as well, hence we can consider also $U(\mathfrak{g}^*)$ and $F[[\mathfrak{g}^*]]$ as above.

Let $\mathfrak{g}$ be a Lie bialgebra. By quantization of $U(\mathfrak{g})$ we mean, roughly speaking, a topological Hopf algebra over $k[[h]]$ which for $h = 0$ is isomorphic, as a co-Poisson Hopf algebra, to $U(\mathfrak{g})$: such an object is called QUEA; similarly, by quantization of $F[[\mathfrak{g}]]$ we mean, roughly speaking, a topological Hopf algebra over $k[[h]]$ which for $h = 0$ is isomorphic, as a Poisson Hopf algebra, to $F[[\mathfrak{g}]]$: such an object is called QFSHA (see Definition 1.2).

The quantum duality principle, in Drinfeld's formulation, states that there exists two functors, namely $(\)': \text{QUEA} \rightarrow \text{QFSHA}$ and $(\)^\vee: \text{QFSHA} \rightarrow \text{QUEA}$, such that:

- *a)* they are inverse of each other;
- *b)* if $U_h(\mathfrak{g})$ is a quantization of $U(\mathfrak{g})$ and $F_h[[\mathfrak{g}]]$ is a quantization of $F[[\mathfrak{g}]]$, then $U_h(\mathfrak{g})'$ is a quantization of $F[[\mathfrak{g}^*]]$, and $F_h[[\mathfrak{g}]]^\vee$ is a quantization of $U(\mathfrak{g}^*)$.

The purpose of this paper is to provide an explicit complete proof of such a result, while also displaying its whole meaning: for instance, we show that Drinfeld's functors establish sort of a "Galois correspondence" within a suitable category of Hopf algebras over $k[[h]]$.

Note that Drinfeld's result is essentially *local* in nature, as it deals with quantizations over the ring of formal series and ends up only with infinitesimal data, i.e. objects attached to Lie bialgebras; a *global* version of the principle, dealing with quantum groups over a ring of Laurent polynomials, which give information on the global data of the underlying Poisson groups will be provided in a forthcoming paper (cf. [Ga]): this is quite useful in applications, e.g. it yields a quantum duality principle for Poisson homogeneous spaces, cf. [CG].

ACKNOWLEDGEMENTS

The author wishes to thank E. Strickland for several useful conversations; he is also indebted with N. Ciccoli and B. Di Blasio for many helpful discussions.

§ 1 Notation and terminology

1.1 The category of topologically free $k[[h]]$ -modules. Let V be a k -vector space; then the space $V[[h]]$ of formal power series in h with coefficients in V has a natural structure of a topological $k[[h]]$ -module. We call a topological $k[[h]]$ -module *topologically free* if it is isomorphic to $V[[h]]$ for some V . We denote by $\mathcal{A}$ the category of topologically free $k[[h]]$ -modules, where morphisms are continuous $k[[h]]$ -linear maps.

For any X in $\mathcal{A}$, we define the *dual* of X to be $X^* := \text{Hom}_{\mathcal{A}}(X, k[[h]])$, which again belongs to $\mathcal{A}$. Furthermore, we set $X_0 := X/hX = k \otimes_{k[[h]]} X$: this is a k -module (via scalar extension $k[[h]] \rightarrow k$), which we call the *specialization* of X at $h = 0$; we shall also

use such a notation as $X \xrightarrow{h \rightarrow 0} \overline{Y}$ to mean that $X_0 \cong \overline{Y}$ (so, for instance, $X \xrightarrow{h \rightarrow 0} X_0$). Finally, we set ${}_F X := k((h)) \otimes_{k[[h]]} X$ (this is a vector space over $k((h))$).

A tensor structure in $\mathcal{A}$ is defined as follows: for X, Y in $\mathcal{A}$, define $X \widehat{\otimes} Y$ to be the projective limit of the $k[[h]]/h^n$ -modules $(X/h^n X) \otimes_{k[[h]]/(h^n)} (Y/h^n Y)$ as $n \rightarrow \infty$. Notice then that $X \widehat{\otimes} Y$ is the h -adic completion of the algebraic tensor product $X \otimes_{k[[h]]} Y$.

Given X in $\mathcal{A}$, when saying that X is a *coalgebra*, resp. a *bialgebra*, resp. a *Hopf algebra*, we mean that X has such a structure with respect to the tensor structure in $\mathcal{A}$: in particular, X has a comultiplication which takes values in $X \widehat{\otimes} X$. We denote by $\mathcal{HA}$ the full subcategory of $\mathcal{A}$ whose objects are the Hopf algebras in $\mathcal{A}$.

Definition 1.2. (cf. [Dr], § 7, or [CP], § 6)

(a) We call *quantized universal enveloping algebra* (in short, *QUEA*) any Hopf algebra $H \in \mathcal{HA}$ such that $H_0 := H/hH$ is isomorphic — as a Hopf algebra over k — to the universal enveloping algebra of a Lie algebra.

(b) We call *quantized formal series Hopf algebra* (in short, *QFSHA*) any Hopf algebra $K \in \mathcal{HA}$ such that $K_0 := K/hK$ is isomorphic — as a topological Hopf algebra over k — to a Hopf algebra of formal series $k[[\{u_i \mid i \in J\}]]$ (for some set J).

1.3 Remark. If H is a QUEA, then its specialization at $h = 0$, that is H_0 , is a co-Poisson Hopf algebra; in particular, this means that, if $\mathfrak{g}$ is the Lie algebra such that $H_0 \cong U(\mathfrak{g})$, then $\mathfrak{g}$ is actually a *Lie bialgebra*; thus we shall write $H = U_h(\mathfrak{g})$. Similarly, if K is a QFSHA, then its specialization at $h = 0$ is a topological Poisson Hopf algebra; in particular, this means that K_0 is (the algebra of regular functions on) a formal Poisson group $F[[\mathfrak{g}]]$: then we shall write $K = F_h[[\mathfrak{g}]]$ (cf. [Dr], § 7, or [CP], § 6, for details). Notice also that K is a local ring, whose maximal ideal is the kernel of the projection map $K \twoheadrightarrow K/hK \cong k[[\{u_i \mid i \in J\}]] \xrightarrow{\bar{\epsilon}} k$ (where $\bar{\epsilon}$ is the counit of $K_0 = K/hK$), or the kernel of $K \xrightarrow{\epsilon} k[[h]] \xrightarrow{ev_0} k[[h]]/hk[[h]] = k$ (ϵ being the counit of K , and ev_0 the k -algebra morphism given by $ev_0(h) := 0$), for the two maps coincide.

The following easy technical results and the definition below will be needed in the sequel:

Lemma 1.4. *Let $X \in \mathcal{A}$. Then $X \cong X_0[[h]]$ as topological $k[[h]]$ -modules; in other words, X and $X_0[[h]]$ are isomorphic in $\mathcal{A}$.*

Proof. Let $\{b_i\}_{i \in I}$ be a basis of X_0 over k . For all $i \in I$, fix a $b'_i \in X$ such that $b_i = b'_i \bmod hX$, and let $\dot{X}$ be the $k[[h]]$ -span of $\{b'_i\}_{i \in I}$: then of course $\dot{X} \cong X_0[[h]]$ as topological $k[[h]]$ -modules. By construction, for any $x \in X$ there exists $\dot{x}_0 \in \dot{X}$ such that $x - \dot{x}_0 \in hX$, that is $x \in \dot{x}_0 + hX$. Iterating, we find $\dot{x}_n \in \dot{X}$ ($\forall n \in \mathbb{N}$) such that $x = \sum_{n=0}^{\infty} \dot{x}_n h^n$; therefore we can conclude that $X = \dot{X}$, whence the claim follows. $\square$

Lemma 1.5. *If H is a QUEA, then H^* is a QFSHA.*

Proof. Let $\mathfrak{g}$ be the Lie algebra over k such that $H_0 \cong U(\mathfrak{g})$; then $H \xrightarrow{h \rightarrow 0} U(\mathfrak{g})$ implies $H^* \xrightarrow{h \rightarrow 0} U(\mathfrak{g})^* = F[[\mathfrak{g}]]$, whence the claim follows. $\square$

Definition 1.6. Let H, K be Hopf algebras (in any category) over a ring R . A pairing $\langle \cdot, \cdot \rangle : H \times K \longrightarrow R$ is called perfect if it is non-degenerate; it is called a Hopf pairing if for all $x, x_1, x_2 \in H, y, y_1, y_2 \in K$, we have

$$\begin{aligned} \langle x, y_1 \cdot y_2 \rangle &= \langle \Delta(x), y_1 \otimes y_2 \rangle, & \langle x_1 \cdot x_2, y \rangle &= \langle x_1 \otimes x_2, \Delta(y) \rangle \\ \langle x, 1 \rangle &= \epsilon(x), & \langle 1, y \rangle &= \epsilon(y), & \langle S(x), y \rangle &= \langle x, S(y) \rangle. \end{aligned}$$

For instance, if H is a Hopf algebra in $\mathcal{A}$, then H^* is a Hopf algebra (in $\mathcal{A}$) as well, and the natural pairing between H and H^* is a perfect Hopf pairing (see [CP] for details).

§ 2 The quantum duality principle

2.1 Drinfeld's functors. Let H be an object of $\mathcal{HA}$. For every $n \in \mathbb{N}$, define $\Delta^n : H \longrightarrow H^{\otimes n}$ by $\Delta^0 := \epsilon$ (where ϵ denotes the counit), $\Delta^1 := id_H$, and $\Delta^n := (\Delta \otimes id_H^{\otimes(n-2)}) \circ \Delta^{n-1}$ if $n > 2$. For any ordered subset $\Sigma = \{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$ with $i_1 < \dots < i_k$, define the morphism $j_\Sigma : H^{\otimes k} \longrightarrow H^{\otimes n}$ by $j_\Sigma(a_1 \otimes \dots \otimes a_k) := b_1 \otimes \dots \otimes b_n$ with $b_i := 1$ if $i \notin \Sigma$ and $b_{i_m} := a_m$ for $1 \leq m \leq k$; then set $\Delta_\Sigma := j_\Sigma \circ \Delta^k$, $\Delta_\emptyset := \Delta^0$, and

$$\delta_\Sigma := \sum_{\Sigma' \subset \Sigma} (-1)^{n-|\Sigma'|} \Delta_{\Sigma'}, \quad \delta_\emptyset := \epsilon. \quad (2.1)$$

By the inclusion-exclusion principle, this relation is equivalent to

$$\Delta_\Sigma = \sum_{\Psi \subseteq \Sigma} \delta_\Psi. \quad (2.2)$$

Moreover, (2.1) also implies: $\delta_\Sigma(1) = 1$ if $\Sigma = \emptyset$, $\delta_\Sigma(1) = 0$ if $\Sigma \neq \emptyset$. We shall also use the notation $\delta_n := \delta_{\{1,2,\dots,n\}}$ for all $n \in \mathbb{N}_+$, $\delta_0 := \delta_\emptyset$. One also has the useful formula

$$\delta_n = (id_H - \epsilon)^{\otimes n} \circ \Delta^n, \quad \forall n \in \mathbb{N}_+. \quad (2.3)$$

Now, H splits as $H = k[[h]] \cdot 1_H \oplus Ker(\epsilon_H) = k[[h]] \cdot 1_H \oplus J_H$, and $(id_H - \epsilon_H)$ is the projector of H onto $J_H := Ker(\epsilon_H)$: so $(id_H - \epsilon_H)^{\otimes n}$ is the projector of $H^{\otimes n}$ onto $J_H^{\otimes n}$; therefore, if $a \in H$ formula (2.3) means that $\delta_n(a) = (id_H - \epsilon_H)^{\otimes n}(\Delta^n(a))$ is nothing but the (unique!) component of $\Delta^n(a)$ in $J_H^{\otimes n}$. Finally, define

$$H' := \{a \in H \mid \delta_n(a) \in h^n H^{\otimes n}\} \quad (\subseteq H)$$

and endow H' with the induced topology.

Now let $I := Ker(H \xrightarrow{\epsilon} k[[h]] \longrightarrow k[[h]]/hk[[h]] = k) = Ker(H \longrightarrow H/hH \xrightarrow{\bar{\epsilon}} k)$, a Hopf ideal of H , and consider the subalgebra $H^\times := \sum_{n \geq 0} h^{-n} I^n = \sum_{n \geq 0} (h^{-1} I)^n$ inside ${}_F H$: then define

$$H^\vee := h\text{-adic completion of } H^\times.$$

Note also that $I = J + h \cdot H$, with $J := \text{Ker}(\epsilon : H \rightarrow k[[h]])$, so $H^\times = \sum_{n \geq 0} h^{-n} J^n$ and

$$H^\vee = h\text{-adic completion of } H^\times.$$

We are now ready to state the main result we are interested in:

Theorem 2.2. (*"The quantum duality principle"; cf. [Dr], §7*)

(a) *The assignment $H \mapsto H'$, resp. $H \mapsto H^\vee$, defines a functor $(\)' : \mathcal{HA} \rightarrow \mathcal{HA}$, resp. $(\)^\vee : \mathcal{HA} \rightarrow \mathcal{HA}$, whose image lies in QFSHA, resp. in QUEA.*

(b) *For all $H \in \mathcal{HA}$, we have $H \subseteq (H^\vee)'$ and $H \supseteq (H')^\vee$. Moreover, we have*

$$H = (H^\vee)' \iff H \in \text{QFSHA} \quad \text{and} \quad H = (H')^\vee \iff H \in \text{QUEA}.$$

Thus the restricted functors $(\)' : \text{QUEA} \rightarrow \text{QFSHA}$ and $(\)^\vee : \text{QFSHA} \rightarrow \text{QUEA}$ are inverse of each other.

(c) (*"Quantum Duality Principle"*) *Letting $\mathfrak{g}$ and $\mathfrak{g}^*$ be as in § 1.1, and using notation of §1.3, we have*

$$U_h(\mathfrak{g})' = F_h[[\mathfrak{g}^*]], \quad F_h[[\mathfrak{g}]]^\vee = U_h(\mathfrak{g}^*),$$

that is to say $U_h(\mathfrak{g})' / h U_h(\mathfrak{g})' = F[[\mathfrak{g}^]]$, $F_h[[\mathfrak{g}]]^\vee / h F_h[[\mathfrak{g}]]^\vee = U(\mathfrak{g}^*)$. $\square$*

Remark 2.3: due to parts (a) and (b) of the theorem above, we can think at Drinfeld's functors as establishing sort of a "Galois correspondence" from $\mathcal{HA}$ to itself. In particular, for any $H \in \mathcal{HA}$ there exists a (unique) minimal QFSHA containing H , namely $(H^\vee)'$, which we might call "the QFSHA-closure of H ", and there exists a (unique) maximal QUEA contained in H , namely $(H')^\vee$, which we might call "the QUEA-interior of H ".

§ 3 Properties of Drinfeld's functors

The rest of this paper will be devoted to prove Theorem 2.2. A key fact is the following:

Proposition 3.1. *Let $H, K \in \mathcal{HA}$, and let $\pi = \langle \ , \ \rangle : H \times K \rightarrow k[[h]]$ be a Hopf pairing:*

(a) *π induces pairings $\langle \ , \ \rangle : H^\vee \times K' \rightarrow k[[h]]$, $\langle \ , \ \rangle : H' \times K^\vee \rightarrow k[[h]]$;*

(b) *if the pairing π is perfect, and the k -valued pairing $\pi_0 = \langle \ , \ \rangle : H_0 \times K_0 \rightarrow k$ naturally induced on the specialized k -algebras is still perfect, then we have (with respect to the natural $k((h))$ -valued pairings induced by scalar extension)*

$$\begin{aligned} K' &= \left\{ \psi \in {}_F K \mid \langle H^\vee, \psi \rangle \subseteq k[[h]] \right\}, & H' &= \left\{ \eta \in {}_F H \mid \langle \eta, K^\vee \rangle \subseteq k[[h]] \right\}, \\ H^\times &= \left\{ \vartheta \in {}_F H \mid \langle \vartheta, K' \rangle \subseteq k[[h]] \right\}, & K^\times &= \left\{ \varphi \in {}_F K \mid \langle H', \varphi \rangle \subseteq k[[h]] \right\}. \end{aligned}$$

Moreover, the k -valued pairings given by specialization of $\langle \cdot, \cdot \rangle : H^\vee \times K' \longrightarrow k[[h]]$ and $\langle \cdot, \cdot \rangle : H' \times K^\vee \longrightarrow k[[h]]$ at $h = 0$ are still perfect.

Proof. Of course, by symmetry, it is enough to prove only one half of the statement, e.g. the parts about the pairing $H^\vee \times K' \longrightarrow k[[h]]$, its specialization, and the identities involving K' and $H^\times$. For later use, set $(H^\vee)^\dagger := \left\{ \psi \in {}_F K \mid \langle H^\vee, \psi \rangle \subseteq k[[h]] \right\}$.

Let $I = I_H$ be the ideal of H considered in §2.1: then $H^\vee$ is the h -adic completion of the space $H^\times := \sum_{n=0}^\infty h^{-n} I^n$; so any $\varphi \in H^\times$ can be uniquely written as a sum $\varphi = \sum_{n=0}^N h^{-n} \varphi_n$ (for some $N \in \mathbb{N}$) with $\varphi_n \in I^n$ for all $n \in \mathbb{N}$, and any $f \in H^\vee$ can be written as a series $f = \sum_{n=0}^{+\infty} s_n h^n$ with $s_n \in H^\times$ for all $n \in \mathbb{N}$.

First, the Hopf pairing $\langle \cdot, \cdot \rangle : {}_F H \times {}_F K \longrightarrow k((h))$ given by scalar extension clearly restricts to a similar pairing $\langle \cdot, \cdot \rangle : H^\times \times K' \longrightarrow k((h))$: we have to prove that the latter takes values in $k[[h]]$, that is $\langle H^\times, K' \rangle \subseteq k[[h]]$, therefore it extends by continuity to a pairing $\langle \cdot, \cdot \rangle : H^\vee \times K' \longrightarrow k[[h]]$; in addition, this will also imply $K' \subseteq (H^\vee)^\dagger$.

To begin with, take $c_1, \dots, c_n \in I$; then $\langle c_i, 1 \rangle = \epsilon(c_i) \in h \cdot k[[h]]$, because $c_i \in I := \text{Ker} \left(H \xrightarrow{\epsilon} k[[h]] \xrightarrow{ev_0} k \right)$ ($i = 1, \dots, n$). Given $y \in K'$, consider

$$\left\langle \prod_{i=1}^n c_i, y \right\rangle = \left\langle \otimes_{i=1}^n c_i, \Delta^{(n)}(y) \right\rangle = \left\langle \otimes_{i=1}^n c_i, \sum_{\Psi \subseteq \{1, \dots, n\}} \delta_\Psi(y) \right\rangle = \sum_{\Psi \subseteq \{1, \dots, n\}} \left\langle \otimes_{i=1}^n c_i, \delta_\Psi(y) \right\rangle$$

(where we make use of (2.2)) and look at the generic summand in the last term above. Let $|\Psi| = t$ ($t \leq n$): then $\langle \otimes_{i=1}^n c_i, \delta_\Psi(y) \rangle = \langle \otimes_{i \in \Psi} c_i, \delta_t(y) \rangle \cdot \prod_{j \notin \Psi} \langle c_j, 1 \rangle$, by definition of δ_Ψ . Thanks to the previous analysis, we have $\prod_{j \notin \Psi} \langle c_j, 1 \rangle \in h^{n-t} k[[h]]$, and $\langle \otimes_{i \in \Psi} c_i, \delta_t(y) \rangle \in h^t k[[h]]$ because $y \in K'$; thus we get $\langle \otimes_{i \in \Psi} c_i, \delta_t(y) \rangle \cdot \prod_{j \notin \Psi} \langle c_j, 1 \rangle \in h^n k[[h]]$, whence $\langle \prod_{i=1}^n c_i, y \rangle \in h^n k[[h]]$. We conclude that

$$\langle \psi, y \rangle \in h^n k[[h]] \quad \forall \quad \psi \in I^n, y \in K'. \quad (2.4)$$

Now take $\varphi = \sum_{n=0}^N h^{-n} \varphi_n \in H^\times$ and $x \in K'$. Since $\varphi_n \in I^n$ for all $n \in \mathbb{N}$, we have (thanks to (2.4)): $\langle \varphi, x \rangle = \sum_{n=0}^N h^{-n} \cdot \langle \varphi_n, x \rangle \in \sum_{n=0}^N h^{-n} \cdot h^n k[[h]] = k[[h]]$, q.e.d.

Therefore the restriction of $\langle \cdot, \cdot \rangle : {}_F H \times {}_F K \longrightarrow k((h))$ to $H^\times \times K'$ does take values in $k[[h]]$: since it is $k[[h]]$ -bilinear and $H^\vee$ is the h -adic completion of $H^\times$, this gives by continuous extension a pairing on $H^\vee \times K'$. This proves (a), which implies $K' \subseteq (H^\vee)^\dagger$ too.

Now we go for (b). First, we prove $(H^\vee)^\dagger \subseteq K'$, by reverting the previous argument.

Let $\psi \in (H^\vee)^\dagger$: then $\langle h^{-s} I^s, \psi \rangle \in k[[h]]$ hence $\langle I^s, \psi \rangle \in h^s k[[h]]$, for all $s \in \mathbb{N}$. In particular, for $s = 0$ this gives $\langle H, \psi \rangle \in k[[h]]$, whence — thanks to the non-degeneracy

of the specialized pairing — we get $\psi \in K$. Now let $n \in \mathbb{N}$ and $i_1, \dots, i_n \in I$; then

$$\begin{aligned} \langle \otimes_{k=1}^n i_k, \delta_n(\psi) \rangle &= \left\langle \otimes_{k=1}^n i_k, \sum_{\Psi \subseteq \{1, \dots, n\}} (-1)^{n-|\Psi|} \cdot \Delta_\Psi(\psi) \right\rangle = \\ &= \sum_{\Psi \subseteq \{1, \dots, n\}} (-1)^{n-|\Psi|} \cdot \left\langle \prod_{k \in \Psi} i_k, \psi \right\rangle \cdot \prod_{k \notin \Psi} \langle i_k, 1 \rangle \in \\ &\in \sum_{\Psi \subseteq \{1, \dots, n\}} \left\langle I^{|\Psi|}, \psi \right\rangle \cdot h^{n-|\Psi|} k[[h]] \subseteq \sum_{s=0}^n h^s \cdot h^{n-s} k[[h]] = h^n k[[h]] \end{aligned}$$

therefore $\langle I^{\otimes n}, \delta_n(\psi) \rangle \subseteq h^n k[[h]]$. In addition, H splits as $H = k \cdot 1_H \oplus \text{Ker}(ev_0 \circ \epsilon_H) = k \cdot 1_H \oplus I$, so $H^{\otimes n}$ splits into the direct sum of $I^{\otimes n}$ plus other direct summands which are tensor products on their own in which at least one tensor factor is $k \cdot 1_H$. Since

$$J_K := \text{Ker}(\epsilon_K) = \{ y \in K \mid \langle 1_H, y \rangle = 0 \} = \langle k \cdot 1_H \rangle^\perp$$

(the orthogonal space of $\langle k \cdot 1_H \rangle$), we have $\langle H^{\otimes n}, j \rangle = \langle I^{\otimes n}, j \rangle$ for any $j \in J_K^{\otimes n}$. Now recall — cf. §2.1 — that $\delta_n(\psi) \in J_K^{\otimes n}$; this and the previous analysis together give $\langle H^{\otimes n}, \delta_n(\psi) \rangle \subseteq h^n k[[h]]$, whence — because of the non-degeneracy of the specialized pairing — we get $\delta_n(\psi) \in h^n K^{\otimes n}$. Therefore $\psi \in K'$; hence we get $(H^\vee)^\dagger \subseteq K'$, q.e.d.

Second, we have to prove that the pairing $\langle \cdot, \cdot \rangle : H^\vee \times K' \longrightarrow k[[h]]$ is still perfect.

This pairing is $k[[h]]$ -bilinear and continuous (with respect to the h -adic topology), hence it induces a natural (continuous) morphism of topological Hopf algebras

$$H^\vee \xrightarrow{\mu} (K')^* := \text{Hom}_{\mathcal{A}}(K', k[[h]]) \quad , \quad f \mapsto \langle f, - \rangle .$$

The non-degeneracy of the pairing is equivalent, by definition, to the triviality of both the right and left kernel of the pairing: the first condition is trivially verified, and the second is equivalent to the injectivity of μ ; hence our task is to prove that $\text{Ker}(\mu) = 0$.

Now, if $f \in H^\vee$ is given by the series $f = \sum_{n=0}^\infty s_n h^n = \lim_{N \rightarrow \infty} \sum_{n=0}^N s_n h^n$, then

$$\mu(f) = \mu \left(\lim_{N \rightarrow \infty} \sum_{n=0}^N s_n h^n \right) = \lim_{N \rightarrow \infty} \sum_{n=0}^N \mu(s_n) h^n = \sum_{n=0}^\infty \mu(s_n) h^n .$$

In particular,

$$\mu(f) = 0 \iff \sum_{n=0}^\infty \mu(s_n) h^n = 0 \iff \mu(s_n) = 0 \quad \forall n \in \mathbb{N} ,$$

where the last equivalence holds because $H^\vee$ is Hausdorff. As a consequence, we have

$$\mu(f) = \sum_{n=0}^\infty \mu(s_n) h^n = 0 \iff \mu(s_n) = 0 \quad \forall n \in \mathbb{N} ,$$

so $\text{Ker}(\mu)$ is the closure of $\text{Ker}(\mu) \cap H^\times = \text{Ker}(\mu|_{H^\times})$. Therefore we simply have to prove that the latter space is zero.

Let $s \in H^\times := \sum_{n=0}^\infty h^{-n} I^n$, $s \neq 0$: then there exists $N \in \mathbb{N}$ such that $s = h^{-N} \cdot s_+$, with $s_+ = \sum_{k=0}^N s_{N-k} h^k \in H$, $s_+ \neq 0$; so $\mu(s) = 0 \iff \mu(s_+) = 0$. Therefore, for the injectivity of μ to be proved it is enough for us to show that $\mu(s_+) \neq 0$.

Since $s_+ \in H \subseteq {}_F H$ and the latter vector space is in perfect pairing with ${}_F K$, there exist $\kappa \in {}_F K$ and a vector subspace $H_+ \leq {}_F H$ such that ${}_F H = k((h)) \cdot s_+ \oplus H_+$ and $\langle s_+, \kappa \rangle \neq 0$, $\langle H_+, \kappa \rangle = 0$; moreover, there exists $t \in \mathbb{N}$ such that $\langle s_+, h^t \kappa \rangle = h^t \langle s_+, \kappa \rangle \in k[[h]]$, so that $\kappa_+ := h^t \kappa \in (H^\vee)^\dagger$. But we have already proved the equality $(H^\vee)^\dagger = K'$, hence $\kappa_+ \in K'$, and $\langle s_+, \kappa_+ \rangle \neq 0$: therefore $\mu(s_+) \neq 0$, q.e.d.

Now we prove that $H^\times = \left\{ \vartheta \in {}_F H \mid \langle \vartheta, K' \rangle \subseteq k[[h]] \right\}$. We already know that $\langle H^\vee, K' \rangle \subseteq k[[h]]$: since $H^\times \subset H^\vee$, this gives $H^\times \subseteq \left\{ \vartheta \in {}_F H \mid \langle \vartheta, K' \rangle \subseteq k[[h]] \right\}$. For the reverse inclusion, pick $\theta \in \left\{ \vartheta \in {}_F H \mid \langle \vartheta, K' \rangle \subseteq k[[h]] \right\}$: then there exists $m \in \mathbb{N}$ such that $h^m \theta \in H$ (by the very definition of ${}_F H$). Thus there exists $n \in \mathbb{N}$ such that $h^m \theta \in I^n \setminus I^{n+1}$: then (2.4) gives $\langle h^m \theta, K' \rangle \in h^n k[[h]]$. Moreover, $\langle \theta, K' \rangle \subseteq k[[h]]$, by hypothesis, so we must have $m \leq n$, whence we conclude: $\theta \in h^{-m} I^n \subseteq h^{-n} I^n \subseteq H^\times$. Therefore $\left\{ \vartheta \in {}_F H \mid \langle \vartheta, K' \rangle \subseteq k[[h]] \right\} \subseteq H^\times$, q.e.d.

Finally, proving that the specialization of the pairing $\langle \cdot, \cdot \rangle : H^\vee \times K' \longrightarrow k[[h]]$ (at $h = 0$) is still perfect amounts to show that

$$(a) \quad \kappa \in K', \quad \langle H^\vee, \kappa \rangle \subseteq h k[[h]] \Rightarrow \kappa \in h K', \quad (b) \quad \eta \in H^\vee, \quad \langle \eta, K' \rangle \subseteq h k[[h]] \Rightarrow \eta \in h H^\vee.$$

Let κ be as in case (a): then $\kappa_- := h^{-1} \kappa \in (H^\vee)^\dagger = K'$, hence $\kappa = h \kappa_- \in h K'$, q.e.d. As for case (b), we can assume $\eta \in H^\times$ (by truncating the h -adic expansion of the initial η): this allows us to resort to the identity $H^\times = \left\{ \vartheta \in {}_F H \mid \langle \vartheta, K' \rangle \subseteq k[[h]] \right\}$. In fact, our hypotheses $\eta \in H^\times$ and $\langle \eta, K' \rangle \subseteq h k[[h]]$ jointly give $\langle h^{-1} \eta, K' \rangle \subseteq k[[h]]$, therefore by the identity we get $h^{-1} \eta \in H^\times$, whence $\eta \in h H^\times \subset H^\vee$, q.e.d. $\square$

Lemma 3.2. *Let H be in $\mathcal{HA}$. Let $a, b \in H$, and let Φ be a finite subset of $\mathbb{N}$. Then*

$$(a) \quad \delta_\Phi(ab) = \sum_{\substack{\Lambda, Y \subseteq \Phi \\ \Lambda \cup Y = \Phi}} \delta_\Lambda(a) \delta_Y(b);$$

$$(b) \quad \text{if } \Phi \neq \emptyset, \text{ then } \delta_\Phi(ab - ba) = \sum_{\substack{\Lambda, Y \subseteq \Phi \\ \Lambda \cup Y = \Phi \\ \Lambda \cap Y \neq \emptyset}} (\delta_\Lambda(a) \delta_Y(b) - \delta_Y(b) \delta_\Lambda(a)). \quad \square$$

Proof. (cf. [KT], Lemma 3.2) In order to prove (a), notice first that (2.2) gives $\sum_{\Psi \subseteq \Phi} \delta_\Psi(ab) = \Delta_\Phi(ab) = \Delta_\Phi(a) \Delta_\Phi(b) = \sum_{\Lambda, Y \subseteq \Phi} \delta_\Lambda(a) \delta_Y(b)$; this can be rewritten as

$$\sum_{\Psi \subseteq \Phi} \delta_\Psi(ab) = \sum_{\Psi \subseteq \Phi} \sum_{\substack{\Lambda, Y \subseteq \Psi \\ \Lambda \cup Y = \Psi}} \delta_\Lambda(a) \delta_Y(b). \quad (2.5)$$

We prove (a) by induction on the cardinality of Φ (to be denoted with $|\Phi|$).

If $\Phi = \emptyset$ then $\delta_\Phi = j_\emptyset \circ \epsilon$, which is a morphism of algebras, so (a) does hold. Now assume (a) is valid for all sets of cardinality less than $|\Phi|$, hence also for all proper subsets of Φ : then the right-hand-side of (2.5) equals

$$\sum_{\substack{\Psi \subseteq \Phi \\ \Psi \neq \Phi}} \delta_\Psi(ab) + \sum_{\substack{\Lambda, Y \subseteq \Phi \\ \Lambda \cup Y = \Phi}} \delta_\Lambda(a) \delta_Y(b).$$

Therefore (a) follows by subtracting from both sides of (2.5) the summands corresponding to the proper subsets Ψ of Φ . Finally, from definitions we have that $\delta_\Lambda(a) \delta_Y(b) = \delta_Y(b) \delta_\Lambda(a)$ whenever $\Lambda \cap Y = \emptyset$, thus (b) follows at once from this and from (a). $\square$

Lemma 3.3. *Let H be a Hopf algebra in $\mathcal{A}$. Then $H^\vee$ is a Hopf algebra in $\mathcal{A}$ as well.*

Proof. Since $H^\vee$ is the h -adic completion of $H^\times := \sum_{n=0}^\infty h^{-n} I^n$ ($\subseteq {}_F H$) (cf. §2.1), it will be enough to show that $H^\times$ is a Hopf subalgebra (over $k[[h]]$) of ${}_F H$, for then the claim will follow by h -adic continuity.

Recall (see §2.1) that I is a Hopf ideal of H : in particular, $S(I) = I$ and $\Delta(I) \subseteq H \widehat{\otimes} I + I \widehat{\otimes} H$, whence also $S(I^n) = I^n$ and $\Delta(I^n) \subseteq \sum_{r+s=n} I^r \widehat{\otimes} I^s$ for all $n \in \mathbb{N}$. Therefore $S(h^{-n} I^n) = h^{-n} I^n$ for all $n \in \mathbb{N}$, so $S(H^\times) = H^\times$ follows at once; similarly

$$\Delta(h^{-n} I^n) \subseteq h^{-n} \sum_{r+s=n} I^r \widehat{\otimes} I^s = \sum_{r+s=n} (h^{-r} I^r) \widehat{\otimes} (h^{-s} I^s) \subseteq H^\times \widehat{\otimes} H^\times$$

for all $n \in \mathbb{N}$, whence $\Delta(H^\times) \subseteq H^\times \otimes H^\times$: so $H^\times$ is a Hopf subalgebra of ${}_F H$, q.e.d. $\square$

Lemma 3.4. *Let H be a Hopf algebra in $\mathcal{A}$. Then H' is a Hopf algebra in $\mathcal{A}$ as well.*

Proof. We shall show that H' is a Hopf subalgebra of H . First, by definition H' is a $k[[h]]$ -submodule of H , for the maps δ_n ($n \in \mathbb{N}$) are $k[[h]]$ -linear; moreover, H' is multiplicatively closed, thanks to the first part of Lemma 3.2. Thus, H' is a $k[[h]]$ -subalgebra of H .

Second, it is clear by construction that

$$(H \otimes H)' = H' \otimes H', \quad (K \otimes K)^\vee = K^\vee \otimes K^\vee \quad (2.6)$$

where K is any other Hopf algebra in $\mathcal{A}$. Now take $a \in H'$, and let $\varphi, \psi \in (H^\star)^\vee$: then

$$\langle \Delta(a), \varphi \otimes \psi \rangle = \langle a, \varphi \cdot \psi \rangle \in k[[h]] \quad (2.7)$$

thanks to Proposition 3.1 (recalling that $(H^\star)^\vee$ is an algebra, by Lemma 3.3 — applied to the Hopf algebra $H^\star$). Even more, again because of Proposition 3.1 — now applied to the Hopf algebras $H \otimes H$ and $H^\star \otimes H^\star$ in the role of H and K , and taking care of (2.6) — (2.7) yields $\Delta(a) \in H' \otimes H'$. Therefore H' is also a subcoalgebra of H .

Third, if S is the antipode of H , then $\Delta \circ S = S^{\otimes 2} \circ \Delta$, so $\Delta^{(n)} \circ S = S^{\otimes n} \circ \Delta^{(n)}$, whence $\delta_n \circ S = S^{\otimes n} \circ \delta_n$, for all $n \in \mathbb{N}$; therefore, for any $a \in H'$ one has $\delta_n(S(a)) = S^{\otimes n}(\delta_n(a)) \in S^{\otimes n}(h^n H^{\otimes n}) = h^n H^{\otimes n}$, for all $n \in \mathbb{N}$, hence $S(a) \in H'$.

Finally, the unit of H obviously does belong to H' , and the counit of H restricts to a counit for H' . Thus H' is a Hopf subalgebra (over $k[[h]]$) of H , q.e.d. $\square$

Corollary 3.5. *Let H, K be Hopf algebras in $\mathcal{A}$, and let $\langle \cdot, \cdot \rangle: H \times K \longrightarrow k[[h]]$ be a Hopf pairing. Then the induced pairings $\langle \cdot, \cdot \rangle: H^\vee \times K'$ and $\langle \cdot, \cdot \rangle: H' \times K^\vee \longrightarrow k[[h]]$ (see Proposition 3.1(a)) are Hopf pairings.*

Proof. Lemmas 3.3–4 tell that $H^\vee, K', H'$, and $K^\vee$ are all Hopf algebras, so the statement is meaningful. Then the proof trivially follows from definitions and the fact that the pairings are induced by a Hopf pairing. $\square$

Proposition 3.6. *$H \mapsto H'$ and $H \mapsto H^\vee$ give well-defined functors from $\mathcal{HA}$ to $\mathcal{HA}$, and for any $H \in \mathcal{HA}$ we have $H \subseteq (H^\vee)'$ and $H \supseteq (H')^\vee$.*

Proof. To define the functors, due to lemmas 3.3–4 we only have to set them on morphisms.

Let $\varphi \in \text{Mor}_{\mathcal{HA}}(H, K)$ be a morphism in $\mathcal{HA}$: by scalar extension it gives a morphism ${}_F H \longrightarrow {}_F K$ of $k((h))$ -Hopf algebras, which maps $h^{-1}I_H$ into $h^{-1}I_K$, hence $H^\times$ into $K^\times$: by continuous extension this gives the morphism $\varphi^\vee \in \text{Mor}_{\mathcal{HA}}(H^\vee, K^\vee)$ we were looking for. On the other hand, we have $\delta_n \circ \varphi = \varphi^{\otimes n} \circ \delta_n$ (for all $n \in \mathbb{N}$), so that $\varphi(H') \subseteq K'$: therefore as $\varphi' \in \text{Mor}_{\mathcal{HA}}(H', K')$ we take simply the restriction $\varphi|_{H'}$ of φ to H' .

Now take $H \in \mathcal{HA}$. For any $n \in \mathbb{N}$, we have $\delta_n(H) \subseteq J_H^{\otimes n}$ (see §2.1); this can be read as $\delta_n(H) \subseteq J_H^{\otimes n} = h^n(h^{-1}J_H)^{\otimes n} \subseteq h^n(H^\vee)^{\otimes n}$, which gives $H \subseteq (H^\vee)'$, q.e.d. On the other hand, let $I' := \text{Ker}(H' \xrightarrow{\epsilon} k[[h]] \xrightarrow{ev_0} k)$; since $(H')^\vee$ is the h -adic completion of $H^\times := \sum_{n=0}^{\infty} (h^{-1}I')^n$, in order to show that $H \supseteq (H')^\vee$ it is enough to check that $H \supseteq h^{-1}I'$. So let $x' \in I'$: then there exists $n \in \mathbb{N}$ such that $x := h^{-n}x' \in H \setminus hH$. Now $x' \in I'$ implies $\delta_1(x') \in hH$, hence $x' = \delta_1(x') + \epsilon(x') \in hH$: therefore $n > 0$; then $h^{-1}x' = h^{-1} \cdot h^n x \in h^{n-1}H \subseteq H$, so $h^{-1}x' \in H$, q.e.d. $\square$

§ 4 Proof of the main theorem

We are now ready to prove Theorem 2.2. We proceed in steps:

Proposition 4.1. *Let H be a Hopf algebra in $\mathcal{A}$. Then H' is a QFSHA.*

Proof. First we prove (following [KT], Proposition 3.5) that H'_0 is commutative.

The very definitions and (2.2) give $id_H = \Delta^{(1)} = \delta_1 + \delta_0 = \delta_1 + \epsilon$; therefore $x = \delta_1(x) + \epsilon(x)$ for all $x \in H$. If $x \in H'$ we have $\delta_1(x) \in hH$, hence there exists $x_1 \in H$ such that $\delta_1(x) = hx_1$. Now let $a, b \in H'$: then

$$a = ha_1 + \epsilon(a), \quad b = hb_1 + \epsilon(b) \implies ab - ba = hc \quad \text{with} \quad c = h(a_1b_1 - b_1a_1);$$

so we are left to show that $c \in H'$. To this end, we have to check that $\delta_\Phi(c)$ is divisible by $h^{|\Phi|}$ for any nonempty finite subset Φ of $\mathbb{N}_+$: but multiplication by h is injective (for H is topologically free), so it is enough to show that $\delta_\Phi(ab - ba)$ is divisible by $h^{|\Phi|+1}$.

Let Λ and Y be subsets of Φ such that $\Lambda \cup Y = \Phi$ and $\Lambda \cap Y \neq \emptyset$: then $|\Lambda| + |Y| \geq |\Phi| + 1$. Now, $\delta_\Lambda(a)$ is divisible by $h^{|\Lambda|}$ and $\delta_Y(b)$ is divisible by $h^{|\Lambda|}$. Then from this and from Lemma 3.2(b) it follows that $\delta_\Phi(ab - ba)$ is divisible by $h^{|\Phi|+1}$.

Now consider $J := \text{Ker}(H \xrightarrow{\epsilon} k[[h]])$ and $J' := \text{Ker}(H' \xrightarrow{\epsilon} k[[h]]) = H' \cap J$. We claim that H' is complete in the J' -adic topology. To begin with, take $j' \in J'$: then $j' = id(j') = \delta_1(j') + \epsilon(j') = \delta_1(j') \in hH \cap J = hJ$, so $J' \subseteq hJ$. Therefore $\prod_{n=0}^{\infty} J'^n \subseteq \prod_{n=0}^{\infty} h^n J^n \subseteq H$, so completing H' with respect to the J' -adic topology does not lead out of H . Thus we only have to show the following: $\delta_n(j'_\infty) \in h^n H^{\otimes n}$ for any $n \in \mathbb{N}$ and $j'_\infty \in \prod_{s=0}^{\infty} J'^s$. Let $j'_\infty = \sum_{s=0}^{\infty} j'_s$ with $j'_s \in J'^s$ for all $s \in \mathbb{N}$. Then $\gamma'_N := \sum_{s=0}^N j'_s \in H'$ hence $\delta_n(\gamma'_N) \in h^n H^{\otimes n}$ for any $n, N \in \mathbb{N}$. Now, $h^n H^{\otimes n}$ is (both open and) closed in the h -adic topology of $H^{\otimes n}$; moreover, the series $j'_\infty = \sum_{s=0}^{\infty} j'_s$ is convergent in the h -adic topology of H as well — for $j'_s \in J'^s \subseteq h^s J^s$ for all $s \in \mathbb{N}$ — and the maps δ_n ($n \in \mathbb{N}$) are continuous with respect to the h -adic topology: therefore we have

$$\delta_n(j'_\infty) = \delta_n\left(\lim_{N \rightarrow \infty} \gamma'_N\right) = \lim_{N \rightarrow \infty} \delta_n(\gamma'_N) \in h^n H^{\otimes n} \quad \text{for all } n \in \mathbb{N}, \quad \text{q.e.d.}$$

As H' is complete in the J' -adic topology, H'_0 is complete in the J'_0 -adic topology, where $J'_0 := \text{Ker}(H'_0 \xrightarrow{\bar{\epsilon}} k)$; moreover, H'_0 is a topological Hopf algebra (for H' is), commutative by the first part of the proof: thus it is a formal group, so H' is a QFSHA. $\square$

Proposition 4.2. *Let H be a Hopf algebra in $\mathcal{A}$. Then $H^\vee$ is a QUEA.*

Proof. Let us consider $K := H^*$: this Hopf algebra is in perfect pairing with H , so Proposition 3.1 tells us that the same is true for K' and $H^\vee$, and also for the k -Hopf algebras K'_0 and $H^\vee_0$. But K'_0 is commutative, by Proposition 4.1, thus the existence of a perfect Hopf pairing among K'_0 and $H^\vee_0$ forces the latter to be cocommutative. As a further consequence, $H^\vee_0$ is a k -Hopf subalgebra of $(K'_0)^*$, the (full) linear dual of K'_0 .

The very definitions imply that $H^\vee_0 := H^\vee / hH^\vee = H^\times / hH^\times =: H^\times_0$: now $H^\times := \sum_{n \geq 0} h^{-n} J_H^n = \sum_{n \geq 0} (h^{-1} J_H)^n$ is generated, as a unital $k[[h]]$ -algebra, by the (maximal) ideal $J_H^\times := h^{-1} J_H$, therefore $H^\times_0$ is generated, as a unital k -algebra, by the (maximal) ideal $(J_H^\times)_0 := J_H^\times \bmod hH^\times$; here $J_H := \text{Ker}(\epsilon: H \rightarrow k[[h]])$, so the (non-unital) $k[[h]]$ -algebra generated by $J_H^\times$ is $\text{Ker}(\epsilon^\times: H^\times \rightarrow k[[h]])$, and the (non-unital) k -algebra generated by $(J_H^\times)_0$ is $\text{Ker}(\epsilon_0: H^\times_0 \rightarrow k)$. Recall also that J_H is a Hopf ideal, which implies that $\Delta^\ell(J_H^n) \subseteq \sum_{\sum_k \nu_k = n} J_H^{\nu_1} \otimes \cdots \otimes J_H^{\nu_\ell}$ for all $\ell, n \in \mathbb{N}$: in particular, we have $\Delta^\ell(J_H^n) \subseteq \sum_{\sum_k \nu_k = n} J_H^{\nu_1} \otimes \cdots \otimes J_H^{\nu_\ell} \subseteq \sum_{r+s+1=\ell} H^r \otimes 1_H \otimes H^s$ for all $\ell > n$. The like holds for $J_H^\times$, so that

$$\Delta^\ell\left((J_H^\times)^n\right) \subseteq \sum_{\sum_k \nu_k = n} (J_H^\times)^{\nu_1} \otimes \cdots \otimes (J_H^\times)^{\nu_\ell} \subseteq \sum_{r+s+1=\ell} H^r \otimes 1_H \otimes H^s \quad \text{for all } \ell > n. \quad (2.8)$$

Now consider $I_K := \text{Ker}(K \xrightarrow{\epsilon} k[[h]] \xrightarrow{ev_0} k)$ and $I'_K := I_K \cap K'$, that is $I'_K = \text{Ker}(\epsilon: K' \xrightarrow{\epsilon} k[[h]] \xrightarrow{ev_0} k)$: these are maximal ideals respectively of K and K' , and

$(I'_K)_0 := I'_K \bmod hK'$ is a maximal ideal of K'_0 . For all $\ell, n \in \mathbb{N}$ with $\ell > n$ we have

$$\begin{aligned} \langle (J_H^\times)^n, (I'_K)^\ell \rangle &= \langle \Delta^{(\ell)}((J_H^\times)^n), (I'_K)^{\otimes \ell} \rangle \subseteq \left\langle \sum_{\sum_k \nu_k = n} J_H^{\nu_1} \otimes \cdots \otimes J_H^{\nu_\ell}, (I'_K)^{\otimes \ell} \right\rangle \subseteq \\ &\subseteq \left\langle \sum_{r+s+1=\ell} H^r \otimes 1_H \otimes H^s, (I'_K)^{\otimes \ell} \right\rangle \in h k[[h]] \end{aligned}$$

due to (2.8) and the fact that $\langle 1_H, I'_K \rangle = \epsilon(I'_K) \subseteq h k[[h]]$. Thus $\langle (J_H^\times)_0^n, (I'_K)_0^\ell \rangle = 0$ for all $\ell > n$, so, since $H^\vee_0$ is generated, as a k -algebra, by $(J_H^\times)_0$, we can conclude that

$$H^\vee_0 \subseteq \left\{ f \in (K'_0)^* \mid f((I'_K)_0^\ell) = 0 \text{ for some } \ell \in \mathbb{N} \right\}. \quad (2.9)$$

Now, by Proposition 4.1 K' is a QFSHA, hence K'_0 is (the algebra of regular functions on) a formal Poisson group, say $K'_0 = F[[\mathfrak{g}]]$; the latter is a local algebra, with (unique) maximal ideal $\mathfrak{m}_{[[e]]} = (I'_K)_0$. Thus (2.9) reads

$$H^\vee_0 \subseteq \left\{ f \in F[[\mathfrak{g}]]^* \mid f(\mathfrak{m}_{[[e]]}^\ell) = 0 \text{ for some } \ell \in \mathbb{N} \right\} = U(\mathfrak{g}) \quad (2.10)$$

where the last equality is one of the possible characterizations of $U(\mathfrak{g})$ (cf. for instance [OV], Ch. 4, Theorem 2.2).

Now, $H^\vee_0$ is a Hopf algebra which, by the first part of the proof, is cocommutative: hence it is automatically a co-Poisson topological Hopf algebra, and the inclusion in (2.10) is clearly a monomorphism of co-Poisson topological Hopf algebras; since $U(\mathfrak{g})$ is a *standard* — i.e. non-topological — Hopf algebra, the same holds true for $H^\vee_0$. Thus $H^\vee_0$ is a co-Poisson cocommutative (standard) Hopf algebra, hence it is the universal enveloping algebra of some Lie bialgebra $\mathfrak{j}$ (namely, the set of its primitive elements), i.e. $H^\vee_0 = U(\mathfrak{j})$; then (2.10) implies that $\mathfrak{j}$ is a Lie sub-bialgebra of $\mathfrak{g}$. But $H^\vee_0 = U(\mathfrak{j})$ is in perfect pairing with $K'_0 = F[[\mathfrak{g}]]$, whence we get $H^\vee_0 = U(\mathfrak{j}) = U(\mathfrak{g})$ and $\mathfrak{j} = \mathfrak{g}$. $\square$

Remarks 4.3: (a) The last part of the previous proof is not strictly necessary: in fact, for the claim to be proved it is enough to show that $H^\vee_0 = U(\mathfrak{j})$, for some Lie bialgebra $\mathfrak{j}$, even without knowing that $\mathfrak{j} = \mathfrak{g}$ with $F[[\mathfrak{g}]]$ being the semiclassical limit of $K' = (H^*)'$.

(b) When the Hopf algebra H in Proposition 4.2 is a QFSHA, say $H = F_h$, we can provide the following alternative proof, which elucidates what's going on.

From the first part of the proof of Proposition 4.1 we know that the topological Hopf algebra $(F_h^\vee)_0 := F_h^\vee / hF_h^\vee$ is cocommutative; if we prove that it is a "standard" (i.e. "non-topological") Hopf algebra, it will necessarily be a universal enveloping algebra, namely $(F_h^\vee)_0 = U(\mathfrak{g})$ for $\mathfrak{g} = \{ \bar{x} \in (F_h^\vee)_0 \mid \Delta(\bar{x}) = \bar{x} \otimes 1 + 1 \otimes \bar{x} \}$.

Since F_h is a QFSHA, we have $(F_h)_0 \cong k[[\{x_j\}_{j \in J}]]$ for some set J ; therefore, letting $\pi: F_h \twoheadrightarrow (F_h)_0$ be the natural projection, if we pick a $y_j \in \pi^{-1}(x_j)$ for any $j \in J$, then (use Lemma 1.4) F_h is generated by $\{y_j\}_{j \in J}$ as a topological $k[[h]]$ -algebra, that is to say

$$F_h = k[[h]][[\{y_j\}_{j \in J}]].$$

Now let $I := \text{Ker}\left(F_h \twoheadrightarrow F_h/hF_h \cong k[[\{y_j\}_{j \in J}]] \xrightarrow{\bar{\epsilon}} k\right) = \text{Ker}\left(F_h \xrightarrow{\epsilon} k[[h]] \xrightarrow{ev_0} k\right)$: we have $I = \langle \{y_j\}_{j \in J} \cup \{h\} \rangle = \langle \{y_j\}_{j \in J} \rangle + hF_h$. By definition $F_h^\vee$ is the h -adic completion of $F_h^\times := \sum_{n=0}^\infty h^{-n}I^n$, so $F_h^\vee/hF_h^\vee = F_h^\times/hF_h^\times$, that is $(F_h^\vee)_0 = (F_h^\times)_0$, hence we go and study the latter space.

Let $h^d y_{j_1}^{d_1} y_{j_2}^{d_2} \cdots y_{j_s}^{d_s}$ be any monomial in h and the y_j 's of degree $\partial = d + \sum_{i=1}^s d_i$; the ideal I^∂ is the set of all the series (in h and the y_j 's) whose degree — i.e., the degree of the "lowest" monomials occurring in the series — is at least ∂ , that is φ is a series of type $\varphi = \sum_{\underline{j}, \underline{d}} c_{\underline{j}, \underline{d}} h^{d_0} y_{j_1}^{d_1} y_{j_2}^{d_2} \cdots y_{j_s}^{d_s} + \text{terms of higher degree}$ (for some $c_{\underline{j}, \underline{d}} \in k$, where the sum runs over all $\underline{j} = (j_1, \dots, j_s)$, $\underline{d} = (d_0, d_1, \dots, d_s)$ — for any $s \in \mathbb{N}$ — such that $\sum_k d_k = \partial$), hence of type $\varphi = \sum_{\underline{j}, \underline{d}} c_{\underline{j}, \underline{d}} h^{d_0} y_{j_1}^{d_1} y_{j_2}^{d_2} \cdots y_{j_s}^{d_s} + \varphi_+$ with $\varphi_+ \in I^{\partial+1}$. Thus the typical element of $(h^{-1}I)^\partial = h^{-\partial}I^\partial$ is of type $h^{-\partial}\varphi = \sum_{\underline{j}, \underline{d}} c_{\underline{j}, \underline{d}} h^{-\partial} h^{d_0} y_{j_1}^{d_1} y_{j_2}^{d_2} \cdots y_{j_s}^{d_s} + h^{-\partial}\varphi_+$. But $\varphi_+ \in I^{\partial+1}$ implies $h^{-\partial}\varphi_+ \in h^{-\partial}I^{\partial+1} = h \cdot h^{-(\partial+1)}I^{\partial+1} \subseteq hF_h^\times$, therefore $h^{-\partial}\varphi \equiv h^{-\partial} \sum_{\underline{j}, \underline{d}} c_{\underline{j}, \underline{d}} h^{d_0} y_{j_1}^{d_1} y_{j_2}^{d_2} \cdots y_{j_s}^{d_s} \pmod{hF_h^\times}$: this means that $(F_h^\vee)_0 = (F_h^\times)_0$ is generated

by the set $\{h^{-1}y_j \pmod{hF_h^\times}\}_{j \in J} =: \{\overline{h^{-1}y_j}\}_{j \in J}$ as a k -algebra (non-topological!). In other words, any $\bar{\varphi} \in (F_h^\vee)_0 = (F_h^\times)_0$ can be written as a polynomial in the $\overline{h^{-1}y_j}$'s. In particular, this implies that the coproduct of $(F_h^\vee)_0$ takes values in the algebraic tensor product $(F_h^\vee)_0 \otimes (F_h^\vee)_0$, so that $(F_h^\vee)_0$ is a standard Hopf algebra, q.e.d.

The next step is just a technical lemma. Hereafter, if $\mathfrak{g}$ is any Lie algebra and $\bar{x} \in U(\mathfrak{g})$, we denote by $\partial(\bar{x})$ the degree of $\bar{x}$ with respect to the standard filtration of $U(\mathfrak{g})$.

Lemma 4.4. *Let U_h be a QUEA, let $x' \in U_h'$, and let $x \in U_h \setminus hU_h$, $n \in \mathbb{N}$, be such that $x' = h^n x$. Set $\bar{x} := x \pmod{hU_h} \in (U_h)_0$. Then $\partial(\bar{x}) \leq n$.*

Proof. (cf. [EK], Lemma 4.12) By hypothesis $\delta_{n+1}(x') \in h^{n+1}U_h^{\otimes(n+1)}$, hence $\delta_{n+1}(x) \in hU_h^{\otimes(n+1)}$, therefore $\delta_{n+1}(\bar{x}) = 0$, i.e. $\bar{x} \in \text{Ker}\left(\delta_{n+1}: U(\mathfrak{g}) \longrightarrow U(\mathfrak{g})^{\otimes(n+1)}\right)$, where $\mathfrak{g}$ is the Lie bialgebra such that $(U_h)_0 := U_h/hU_h = U(\mathfrak{g})$. But the latter kernel equals the subspace $U(\mathfrak{g})_n := \{\bar{y} \in U(\mathfrak{g}) \mid \partial(\bar{y}) \leq n\}$ (cf. [KT], §3.8), whence the claim follows. $\square$

Proposition 4.5. *Let H be a QFSHA. Then $(H^\vee)' = H$.*

Proof. Let $x' \in (H^\vee)'$ be given; let $n \in \mathbb{N}$, $x \in H^\vee \setminus hH^\vee$ be such that $x' = h^n x$, and let $x_0 \in H^\times$ be such that $x = x_0 + hx_{\langle 1 \rangle}$ for some $x_{\langle 1 \rangle} \in H^\vee$, so that $x \equiv x_0 \pmod{hH^\vee}$. By Proposition 4.2 $H^\vee$ is a QUEA, with semiclassical limit $U(\mathfrak{g})$ for some Lie bialgebra $\mathfrak{g}$; then Lemma 4.4 gives $\partial(\bar{x}_0) = \partial(\bar{x}) \leq n$.

Fix an ordered basis $\{b_\lambda\}_{\lambda \in \Lambda}$ of $\mathfrak{g}$, and a subset $\{x_\lambda\}_{\lambda \in \Lambda}$ of $H^\vee$ such that $x_\lambda \pmod{hH^\vee} = b_\lambda$ for all λ ; in particular, since $\mathfrak{g} \subset \text{Ker}(\epsilon_{U(\mathfrak{g})})$ we can choose the x_λ 's inside $J^\vee := h^{-1}J$, where $J := \text{Ker}(\epsilon: H \longrightarrow k[[h]])$: so $x_\lambda = h^{-1}x'_\lambda$ for some $x'_\lambda \in J$, for all $\lambda \in \Lambda$.

Since $\partial(\bar{x}_0) \leq n$, that is $\bar{x}_0 \in U(\mathfrak{g})_n := \{\bar{y} \in U(\mathfrak{g}) \mid \partial(\bar{y}) \leq n\}$, we can write $\bar{x}_0$ as a polynomial $P(\{b_\lambda\}_{\lambda \in \Lambda})$ in the b_λ 's of degree $d \leq n$; then $P(\{x_\lambda\}_{\lambda \in \Lambda}) \equiv x_0 \pmod{hH^\vee}$,

so we can assume $x_0 = P(\{x_\lambda\}_{\lambda \in \Lambda})$. Now we can write x_0 as $x_0 = \sum_{s=0}^d j_s h^{-s}$, where $j_s \in J^s$ is a homogeneous polynomial in the x'_λ 's of degree s , and $j_d \neq 0$; but then $h^n x_0 = \sum_{s=0}^d j_s h^{n-s} \in H$ because $d \leq n$. Since $H \subseteq (H^\vee)'$ — thanks to Proposition 3.6 — we get also $h^{n+1} x_{\langle 1 \rangle} = x' - h^n x_0 \in (H^\vee)'$: therefore we have

$$x' = h^n x_0 + h^{n+1} x_{\langle 1 \rangle}, \quad \text{with } h^n x_0 \in H \quad \text{and} \quad h^{n+1} x_{\langle 1 \rangle} \in (H^\vee)'.$$

So now we can repeat the argument for $x_{(1)} := h^{n+1} x_{\langle 1 \rangle}$ in the role of $x' =: x_{(0)}$: this will provide us with an $x_1 \in H^\times$ and an $x_{\langle 2 \rangle} \in H^\vee$ such that

$$x_{(1)} = h^{n+1} x_1 + h^{n+2} x_{\langle 2 \rangle}, \quad \text{with } h^{n+1} x_1 \in H \quad \text{and} \quad h^{n+2} x_{\langle 2 \rangle} \in (H^\vee)'.$$

Iterating, we eventually find a sequence $\{x_\ell\}_{\ell \in \mathbb{N}} \subset H^\times$ such that $h^{n+\ell} x_\ell \in H$ for all $\ell \in \mathbb{N}$, and $x' = \sum_{\ell=0}^{\infty} h^{n+\ell} x_\ell$; to be precise, the series $\sum_{\ell=0}^{\infty} h^{n+\ell} x_\ell$ does converge to x' in $H^\vee$, hence also in $(H^\vee)'$, for the topology of the latter space is induced from that of $H^\vee$: furthermore, the partial sums of the series belong to H , whose topology coincides with the one given by restriction from $H^\vee (\supset H)$, hence $x' \in H$ as well. $\square$

Proposition 4.6. *Let H be a QUEA. Then $(H')^\vee = H$.*

Proof. Consider $K := H^*$: this is a QFSHA (cf. Lemma 1.5), hence Proposition 4.5 gives $(K^\vee)' = K$. Applying twice Proposition 3.1 we find that both H and $(H')^\vee$ are in perfect pairing with K ; moreover, by Proposition 3.6 we have $(H')^\vee \subseteq H$. Since $K := H^*$, if $(H')^\vee \subsetneq H$ then there exists $\phi \in K$ such that $\langle \phi, H \rangle \neq 0$, i.e. $\phi \neq 0$, whilst $\langle \phi, (H')^\vee \rangle = 0$: but this is in contrast with the non-degeneracy of the pairing between $(H')^\vee$ and K , thus $(H')^\vee = H$. $\square$

4.7 Remark. The explicit description of $U(\mathfrak{g})^* = F[[\mathfrak{g}]]$ (see for instance [Dx], Ch. 2, §7) shows that it can be realized as follows; consider $\mathfrak{g}^*$, its tensor powers $(\mathfrak{g}^*)^{\otimes n}$ ($n \in \mathbb{N}$) and their direct product $\prod_{n=0}^{\infty} (\mathfrak{g}^*)^{\otimes n}$ (the completion, in a suitable topology, of the direct sum $\bigoplus_{n=0}^{\infty} (\mathfrak{g}^*)^{\otimes n} = S(\mathfrak{g}^*)$): this has a natural structure of topological associative commutative algebra with 1 (extending the structure of the symmetric algebra $S(\mathfrak{g}^*)$). In addition, this algebra structure can be completed to make $\prod_{n=0}^{\infty} (\mathfrak{g}^*)^{\otimes n}$ into a topological Hopf algebra, isomorphic to $U(\mathfrak{g})^*$. If $\{b_j\}_{j \in J}$ is any basis of $\mathfrak{g}$ over k , let $\{x_j\}_{j \in J} \in \mathfrak{g}^*$ be such that $\langle x_i, b_\ell \rangle = \delta_{i,\ell}$ for all $i, \ell \in J$. If $\dim(\mathfrak{g}) < +\infty$, then $\{x_j\}_{j \in J}$ is a basis of $\mathfrak{g}^*$, hence the monomials of degree n in the x_j 's form a basis of $(\mathfrak{g}^*)^{\otimes n}$ ($n \in \mathbb{N}$) and the set of all monomials in the x_j 's is a topological basis of $\prod_{n=0}^{\infty} (\mathfrak{g}^*)^{\otimes n} = U(\mathfrak{g})^*$. On the other hand, if $\dim(\mathfrak{g}) = +\infty$, then $\{x_j\}_{j \in J}$ is no longer a basis, but it is a *pseudobasis*

of $\mathfrak{g}^*$, so that any element of $\mathfrak{g}^*$ can be uniquely written as a k -linear combination of the x_j 's (cf. for instance [Di, Ch. I]): but then the monomials of degree n in the x_j 's form a pseudobasis of $(\mathfrak{g}^*)^{\otimes n}$ ($n \in \mathbb{N}$) and the set of *all* monomials in the x_j 's is a pseudobasis of $\coprod_{n=0}^{\infty} (\mathfrak{g}^*)^{\otimes n} = U(\mathfrak{g})^*$. The upset is that $U(\mathfrak{g})^*$ can still be viewed as an algebra of formal power series, namely $k[[\{x_j\}_{j \in J}]]$, which contains also all series in which infinitely many monomials in the x_j 's of the same degree can occur with non-zero coefficient.

Proposition 4.8. *Let $F_h = F_h[[\mathfrak{g}]]$ be a QFSHA. Then $F_h[[\mathfrak{g}]]^\vee = U_h(\mathfrak{g}^*)$, that is to say*

$$F_h[[\mathfrak{g}]]^\vee / h F_h[[\mathfrak{g}]]^\vee = U(\mathfrak{g}^*).$$

Proof. We first recall some standard facts about "classical" objects. Consider the formal group $F_h/hF_h = F[[\mathfrak{g}]] = U(\mathfrak{g})^*$: this is a local ring, with maximal ideal $J_F := \text{Ker}(\epsilon: F[[\mathfrak{g}]] = U(\mathfrak{g})^* \rightarrow k)$. Let $\rho: U(\mathfrak{g})^* \twoheadrightarrow \mathfrak{g}^*$ be the restriction map. Then — cf. Remark 4.7 — we can take a linear map $\nu: \mathfrak{g}^* \hookrightarrow U(\mathfrak{g})^*$ which is a section of ρ , i.e. $\rho \circ \nu = \text{id}_{\mathfrak{g}^*}$; moreover, we can choose such a map that has image in J_F , so $\nu: \mathfrak{g}^* \hookrightarrow J_F$. In addition, restricting ρ to J_F induces an isomorphism $J_F/J_F^2 \xrightarrow{\cong} \mathfrak{g}^*$; in particular, if $\nu': \mathfrak{g}^* \hookrightarrow J_F$ is another section as above, then for any $f \in \mathfrak{g}^*$ we have $(\nu - \nu')(f) \in J_F^2$. From now on, we fix a section $\nu: \mathfrak{g}^* \hookrightarrow J_F$ of the previous type.

Now consider $J_{F_h} := \text{Ker}(\epsilon: F_h[[\mathfrak{g}]] \rightarrow k[[h]])$: then the specialization map $\pi: F_h \twoheadrightarrow F_h/hF_h$ restricts to $\pi': J_{F_h} \twoheadrightarrow J_{F_h}/(J_{F_h} \cap hF_h) = J_{F_h}/hJ_{F_h} = J_F$. We fix a section $\gamma: J_F \hookrightarrow J_{F_h}$ of π' . Multiplication by h^{-1} yields a $k[[h]]$ -module isomorphism $\mu: J_{F_h} \xrightarrow{\cong} h^{-1}J_{F_h}$. Furthermore, $h^{-1}J_{F_h} \subset F_h^\times \subset F_h^\vee$, and the specialization map $\pi^\times: F_h^\times \twoheadrightarrow F_h^\times/hF_h^\times = F_h^\vee/hF_h^\vee$ restricts — taking into account that $(h^{-1}J_{F_h}) \cap (hF_h^\times) = h^{-1}J_{F_h}^2$ — to a similar map $\eta: h^{-1}J_{F_h} \twoheadrightarrow h^{-1}J_{F_h}/h^{-1}J_{F_h}^2$ (surjective again). By Proposition 4.2 we know that $F_h^\vee$ is a QUEA, so $F_h^\vee/hF_h^\vee$ is the universal enveloping algebra $U(\mathfrak{h})$ of some Lie bialgebra $\mathfrak{h}$; our purpose is to prove that $\mathfrak{h} \cong \mathfrak{g}^*$ as Lie bialgebras. Setting $J_U := \text{Ker}(\epsilon: U(\mathfrak{h}) \rightarrow k)$, we remark also that $\eta(h^{-1}J_{F_h}) \subseteq J_U$, and $J_U/J_U\mathfrak{h} = \mathfrak{h}$; moreover, what we saw in Remark 4.3(b) shows also that the composition map $\vartheta := p \circ \eta: h^{-1}J_{F_h} \xrightarrow{\eta} J_U \xrightarrow{p} J_U/J_U\mathfrak{h} = \mathfrak{h}$ (where the map $p: J_U \twoheadrightarrow J_U/J_U\mathfrak{h} = \mathfrak{h}$ is the natural projection) is onto as well, with $\text{Ker}(\vartheta) = h^{-1}J_{F_h}^2 + J_{F_h}$. In fact, resuming notation of Remark 4.3(b), one sees that the y_j 's may be chosen in J_{F_h} and then the cosets $\vartheta(h^{-1}y_j)$ ($j \in J$) span $J_U/J_U\mathfrak{h} = \mathfrak{h}$.

Now look at the composition map $\sigma := \vartheta \circ \mu \circ \gamma \circ \nu$, i.e.

$$\sigma: \mathfrak{g}^* \xrightarrow{\nu} J_F \xrightarrow{\gamma} J_{F_h} \xrightarrow{\mu} h^{-1}J_{F_h} \xrightarrow{\vartheta} J_U/J_U\mathfrak{h} = \mathfrak{h};$$

we are going to show that this is an isomorphism of Lie bialgebras.

We begin by proving that σ is injective. Let $\sigma(\phi) = 0$ for some $\phi \in \mathfrak{g}^*$: then we have

$$\begin{aligned} \sigma(\phi) = 0 &\implies \vartheta(\mu(\gamma(\nu(\phi)))) = 0 \implies \mu(\gamma(\nu(\phi))) \in \text{Ker}(\vartheta) = h^{-1}J_{F_h}^2 + J_{F_h} \implies \\ &\implies \gamma(\nu(\phi)) \in J_{F_h}^2 + hJ_{F_h} \implies \nu(\phi) = \pi'(\gamma(\nu(\phi))) \in \pi'(J_{F_h}^2 + hJ_{F_h}) = J_F^2 \implies \\ &\implies \phi = \rho(\nu(\phi)) = 0 \end{aligned}$$

where the last identity follows from the (above mentioned fact) that restricting ρ to J_F induces an isomorphism $J_F/J_F^2 \xrightarrow{\cong} \mathfrak{g}^*$ (so in particular $\rho(J_F^2) = 0$). Thus σ is injective.

Now we prove that σ is surjective. Let $y \in \mathfrak{h}$ be given. Since ϑ and μ are onto, there exists $j \in J_{F_h}$ such that $\vartheta(\mu(j)) = y$; then, since γ is a section of $\pi': J_{F_h} \twoheadrightarrow J_F$, there exists $j \in J_F$ such that $\gamma(j) = j + hj'$ for some $j' \in J_{F_h}$. Second, since ν is a section of $J_F \twoheadrightarrow J_F/J_F^2 = \mathfrak{g}^*$, there exists $\phi \in \mathfrak{g}^*$ such that $\nu(\phi) = j + j_2$ for some $j_2 \in J_F^2$. As above, since $\pi'(J_{F_h}^2) = J_F^2$ and γ is a section of π' we have $\gamma(j_2) = j'' + hj_1$ for some $j'' \in J_{F_h}^2$, $j_1 \in J_{F_h}$. Therefore

$$\begin{aligned} \sigma(\phi) &= \vartheta(\mu(\gamma(\nu(\phi)))) = \vartheta(\mu(\gamma(j + j_2))) = \vartheta(\mu(\gamma(j))) + \vartheta(\mu(\gamma(j_2))) = \\ &= \vartheta(\mu(j)) + \vartheta(\mu(hj')) + \vartheta(\mu(j'')) + \vartheta(\mu(hj_1)) = y + \vartheta(j') + \vartheta(h^{-1}j'') + \vartheta(j_1) = y \end{aligned}$$

for $j', j_1 \in J_{F_h} \subseteq \text{Ker}(\vartheta)$ and $h^{-1}j'' \in h^{-1}J_{F_h}^2 \subseteq \text{Ker}(\vartheta)$. So $\sigma(\phi) = y$, hence σ is onto.

Actually, we can prove also that σ is independent of the choice of ν and γ . Indeed, if $\gamma, \gamma': J_F \hookrightarrow J_{F_h}$ are two sections of π' , and σ, σ' are defined correspondingly (with the same ν for both), we have $\gamma(f) - \gamma'(f) \in hJ_{F_h} \subseteq \text{Ker}(\vartheta \circ \mu)$, thus we have also $\vartheta(\mu(\gamma(f))) = \vartheta(\mu(\gamma'(f)))$: thus $\vartheta \circ \mu \circ \gamma = \vartheta \circ \mu \circ \gamma'$, hence $\sigma = \sigma'$, that is σ is independent of the choice of γ . Similarly, if $\nu, \nu': \mathfrak{g}^* \hookrightarrow J_F$ are two sections of ρ , and σ, σ' are defined correspondingly (with some fixed γ for both), we noticed that for any $f \in \mathfrak{g}^*$ we have $\nu(f) - \nu'(f) \in J_F^2$: therefore the analysis above gives $\vartheta(\mu(\gamma(\nu(f)))) = \vartheta(\mu(\gamma(\nu'(f))))$, so $\vartheta \circ \mu \circ \gamma \circ \nu = \vartheta \circ \mu \circ \gamma \circ \nu'$, whence $\sigma = \sigma'$, that is σ is independent of the choice of ν .

In other words, what we proved is that σ is equal to the composition map

$$\mathfrak{g}^* \xrightarrow{\bar{\nu}} J_F/J_F^2 \xrightarrow{\bar{\gamma}} J_{F_h}/(J_{F_h}^2 + hJ_{F_h}) \xrightarrow{\bar{\mu}} h^{-1}J_{F_h}/(h^{-1}J_{F_h}^2 + J_{F_h}) \xrightarrow{\bar{\vartheta}} J_U/J_U^2 = \mathfrak{h}$$

where the maps $\bar{\nu}, \bar{\gamma}, \bar{\mu}, \bar{\vartheta}$, are the ones canonically induced by $\nu, \gamma, \mu, \vartheta$, and $\bar{\nu}$, resp. $\bar{\gamma}$, does not depend on the choice of ν , resp. γ , as it is the inverse of the isomorphism $\bar{\rho}: J_F/J_F^2 \xrightarrow{\cong} \mathfrak{g}^*$, resp. $\bar{\pi}': J_{F_h}/(J_{F_h}^2 + hJ_{F_h}) \xrightarrow{\cong} J_F/J_F^2$, induced by ρ , resp. by π' . We use this remark to show that σ is also an isomorphism of the Lie bialgebra structure.

Take $\phi_1, \phi_2 \in \mathfrak{g}^*$, and let $f_i := \nu(\phi_i)$, $\varphi_i := \gamma(f_i)$, for $i = 1, 2$. First, we have $\bar{\nu}([\phi_1, \phi_2]) = \overline{\{f_1, f_2\}}$ (hereafter for any space X with a subspace X_+ and an $x \in X$, we denote by $\bar{x} \in X/X_+$ the coset $x + X_+$), because $\pi'(\{f_1, f_2\}) = [\phi_1, \phi_2]$ — i.e. π' is a Lie algebra morphism between $(J_F, \{, \})$ and $(\mathfrak{g}^*, [,])$ — since $\langle x, [\phi_1, \phi_2] \rangle = \langle \delta(x), \phi_1 \otimes \phi_2 \rangle = \langle \delta(x), \rho(f_1) \otimes \rho(f_2) \rangle = \langle \delta(x), f_1 \otimes f_2 \rangle = \langle x, \{f_1, f_2\} \rangle$ for all $x \in \mathfrak{g}$. Similarly, $\bar{\gamma}(\overline{\{f_1, f_2\}}) = \overline{h^{-1}[\varphi_1, \varphi_2]}$ because $h^{-1}[\varphi_1, \varphi_2] \in J_{F_h}$ and $\pi'(h^{-1}[\varphi_1, \varphi_2]) = \{f_1, f_2\}$. Then we have $\mu(h^{-1}[\varphi_1, \varphi_2]) = h^{-2}[\varphi_1, \varphi_2] = [h^{-1}\varphi_1, h^{-1}\varphi_2] = [\mu(\varphi_1), \mu(\varphi_2)]$ hence $\bar{\mu}(\overline{h^{-1}[\varphi_1, \varphi_2]}) = \overline{[\mu(\varphi_1), \mu(\varphi_2)]}$, and $\eta([\mu(\varphi_1), \mu(\varphi_2)]) = [\eta(\mu(\varphi_1)), \eta(\mu(\varphi_2))]$ because η is the restriction of a morphism of algebras. Finally, the last term is mapped by p onto $p([\eta(\mu(\varphi_1)), \eta(\mu(\varphi_2))]) = [p(\eta(\mu(\varphi_1))), p(\eta(\mu(\varphi_2)))]$, so $\vartheta([\mu(\varphi_1), \mu(\varphi_2)]) = [\vartheta(\mu(\varphi_1)), \vartheta(\mu(\varphi_2))]$. Summing up, we get

$$\begin{aligned} \sigma([\phi_1, \phi_2]) &= \bar{\vartheta}(\bar{\mu}(\bar{\gamma}(\bar{\nu}([\phi_1, \phi_2])))) = \bar{\vartheta}(\bar{\mu}(\bar{\gamma}(\overline{\{f_1, f_2\}}))) = \bar{\vartheta}(\bar{\mu}(\overline{h^{-1}[\varphi_1, \varphi_2]}))) = \\ &= \bar{\vartheta}(\overline{[\mu(\varphi_1), \mu(\varphi_2)]}) = \vartheta([\mu(\varphi_1), \mu(\varphi_2)]) = [\vartheta(\mu(\varphi_1)), \vartheta(\mu(\varphi_2))] = \\ &= [\vartheta(\mu(\gamma(\nu(\phi_1)))), \vartheta(\mu(\gamma(\nu(\phi_2))))] = [\sigma(\phi_1), \sigma(\phi_2)] \end{aligned}$$

that is $\sigma([\phi_1, \phi_2]) = [\sigma(\phi_1), \sigma(\phi_2)]$, so that σ is a morphism of Lie algebras.

The Lie cobracket can be handled similarly; but it takes values respectively into $\mathfrak{g}^* \otimes \mathfrak{g}^*$ and $\mathfrak{h} \otimes \mathfrak{h}$, so the same arguments are to be applied to such maps as $\bar{\nu}^{\otimes 2}$, $\bar{\gamma}^{\otimes 2}$, etc.

Take $\phi \in \mathfrak{g}^*$, and let $f := \nu(\phi)$, $\varphi := \gamma(f)$. First, we have $\bar{\nu}^{\otimes 2}(\delta(\phi)) = \overline{\nabla(f)}$ (hereafter we set $\nabla(x) := (\Delta - \Delta^{op})(x)$ where $\Delta^{op} := \tau_{1,2} \circ \Delta$ is the opposite co-product of Δ , and $\tau_{1,2}$ is the flip), because $\pi'^{\otimes 2}(\nabla(f)) = \delta(f)$ since $\langle x \otimes x_2, \delta(\phi) \rangle = \langle [x_1, x_2], \phi \rangle = \langle [x_1, x_2], \rho(f) \rangle = \langle [x_1, x_2], f \rangle = \langle x_1 \otimes x_2, \nabla(f) \rangle$ for all $x_1, x_2 \in \mathfrak{g}$. Similarly, $\bar{\gamma}^{\otimes 2}(\nabla(\bar{f})) = \nabla(\bar{\gamma}^{\otimes 2}(\bar{f})) = \nabla(\bar{\varphi})$ because $\pi'^{\otimes 2}(\nabla(\bar{\varphi})) = \nabla(\pi'(\bar{\varphi}))$ for π' is the restriction of a morphism of Hopf algebras. Then $\bar{\mu}^{\otimes 2}(\nabla(\bar{\varphi})) = \overline{h^{-2}\nabla(\bar{\varphi})} = \overline{h^{-1}\nabla(\overline{h^{-1}\varphi})} = \overline{h^{-1}\nabla(\bar{\mu}(\bar{\varphi}))}$, and $\eta^{\otimes 2}(h^{-1}\nabla(\mu(\varphi))) = \delta(\eta(\mu(\varphi)))$; finally, the latter element is mapped by $p^{\otimes 2}$ onto $p^{\otimes 2}(\delta(\eta(\mu(\varphi)))) = \delta(p(\eta(\mu(\varphi)))) = \delta(\vartheta(\mu(\varphi)))$. Therefore

$$\begin{aligned} \sigma^{\otimes 2}(\delta(\phi)) &= \bar{\vartheta}^{\otimes 2}(\bar{\mu}^{\otimes 2}(\bar{\gamma}^{\otimes 2}(\bar{\nu}^{\otimes 2}(\delta(\phi))))) = \bar{\vartheta}^{\otimes 2}(\bar{\mu}^{\otimes 2}(\bar{\gamma}^{\otimes 2}(\nabla(\bar{f})))) = \\ &= \bar{\vartheta}^{\otimes 2}(\bar{\mu}^{\otimes 2}(\nabla(\bar{\varphi}))) = \bar{\vartheta}^{\otimes 2}(\overline{h^{-1}\nabla(\bar{\mu}(\bar{\varphi}))}) = \vartheta^{\otimes 2}(h^{-1}\nabla(\bar{\mu}(\bar{\varphi}))) = p^{\otimes 2}(\eta^{\otimes 2}(h^{-1}\nabla(\mu(\varphi)))) = \\ &= p^{\otimes 2}(\delta(\eta(\mu(\varphi)))) = \delta(p(\eta(\mu(\varphi)))) = \delta(\vartheta(\mu(\varphi))) = \delta(\vartheta(\mu(\gamma(\nu(\phi)))) = \delta(\sigma(\phi)) \end{aligned}$$

that is $\sigma^{\otimes 2}(\delta(\phi)) = \delta(\sigma(\phi))$, so that σ is a morphism of Lie coalgebras too. $\square$

Proposition 4.9. *Let $U_h(\mathfrak{g})$ be a QUEA. Then $U_h(\mathfrak{g})' = F_h[[\mathfrak{g}^*]]$, that is to say*

$$U_h(\mathfrak{g})' / h U_h(\mathfrak{g})' = F[[\mathfrak{g}^*]].$$

Proof. By Lemma 1.5, $F_h[[\mathfrak{g}]] := U_h(\mathfrak{g})^*$ is a QFSHA, whose semiclassical limit is $F[[\mathfrak{g}]]$. Then Proposition 4.2 applies, so $F_h[[\mathfrak{g}]]^\vee$ is a QUEA, whose semiclassical limit, thanks to Proposition 4.8, is $U(\mathfrak{g}^*)$. On the other hand, $U_h(\mathfrak{g})'$ is a QFSHA, due to Proposition 4.1, so $U_h(\mathfrak{g})' = F_h[[\mathfrak{h}]]$ for some Lie bialgebra $\mathfrak{h}$. Now, $U_h(\mathfrak{g})$ and $F_h[[\mathfrak{g}]]$ are Hopf algebras in perfect pairing; thanks to Proposition 3.1, the same holds as well for $U_h(\mathfrak{g})'$ and $F_h[[\mathfrak{g}]]^\vee$, and when specializing h to zero we get a perfect Hopf pairing between $(U_h(\mathfrak{g})')_0 = F[[\mathfrak{h}]]$ and $(F_h[[\mathfrak{g}]]^\vee)_0 = U(\mathfrak{g}^*)$: this forces $\mathfrak{h} = \mathfrak{g}^*$, q.e.d. $\square$

Finally, our efforts are rewarded:

Proof of Theorem 2.2. We just have to collect together the previous results. Part (a) is proved by the first part of Proposition 3.6 and by Propositions 4.1–2; part (b) is proved by the second part of Proposition 3.6 and by Propositions 4.5–6 (the latter give one half of the double implications, whereas the other half follows from part (a)); finally, part (c) is proved by Propositions 4.8–9. $\square$

REFERENCES

- [CG] N. Ciccoli, F. Gavarini, *A quantum duality principle for Poisson homogeneous spaces*, Preprint.
- [CP] V. Chari, A. Pressley, *A guide to Quantum Groups*, Cambridge University Press, Cambridge, 1994.
- [Di] J. Dieudonné, *Introduction to the theory of formal groups*, Pure and Appl. Mathem. **20**, M. Dekker inc., New York, 1973.
- [Dx] J. Dixmier, *Enveloping Algebras*, American Mathematical Society, Providence, RI, 1996.
- [Dr] V. G. Drinfeld, *Quantum groups*, Proc. Intern. Congress of Math. (Berkeley, 1986), 1987, pp. 798–820.
- [EK] P. Etingof, D. Kazhdan, *Quantization of Lie bialgebras, I*, Selecta Math. (New Series) **2** (1996), 1–41.
- [FRT] L. D. Faddeev, N. Yu. Reshetikhin, L. A. Takhtajan, *Quantum groups*, in: M. Kashiwara, T. Kawai (eds.), *Algebraic Analysis*, (1989), Academic Press, Boston, 129–139.
- [KT] C. Kassel, V. Turaev, *Biquantization of Lie bialgebras*, Preprint math.QA/9810009 (1998).
- [Ga] F. Gavarini, *The global quantum duality principle*, Preprint.

UNIVERSITÀ DEGLI STUDI DI ROMA “TOR VERGATA”
DIPARTIMENTO DI MATEMATICA
VIA DELLA RICERCA SCIENTIFICA, 1
I-00133 ROMA — ITALY
E-MAIL: GAVARINI@MAT.UNIROMA2.IT