EXACT ESTIMATES FOR MOMENTS OF RANDOM BILINEAR FORMS

By R. Ibragimov¹, Sh. Sharakhmetov² and A. Cecen³

Running head: Exact estimates for moments of random bilinear forms

Abstract. The present paper concentrates on the analogues of Rosenthal's inequalities for ordinary and decoupled bilinear forms in symmetric random variables. More specifically, we prove the exact moment inequalities for these objects in terms of moments of their individual components. As a corollary of these results we obtain the explicit expressions for the best constant in the analogues of Rosenthal's inequality for ordinary and decoupled bilinear forms in identically distributed symmetric random variables in the case of the fixed number of random variables.

Key words: random bilinear forms, moment inequalities, decoupling, symmetric statistics.

AMS 1991 Subject Classification: Primary 60E15, 60F25, 60G50

Mailing address: Department of Economics, Yale University, 28 Hillhouse Ave., New Haven, CT 06511; Phone: (203) 7772750; E-Mail: rustam.ibragimov@yale.edu

¹ Department of Economics, Yale University, 28 Hillhouse Ave., New Haven, CT 06511. E-Mail: rustam.ibragimov@yale.edu. Supported in part by a grant from Central Michigan University Graduate Studies and by a grant from the American Council of Teachers of Russian/American Council for Collaboration in Education and Language Study, with funds provided by the United States Information Agency

² Department of Probability Theory, Tashkent State Economics University, Tashkent, Uzbekistan
 ³ Department of Economics, Central Michigan University, Mt. Pleasant, Michigan, USA
 Supported in part by a grant from Central Michigan University Department of Economics

1. Introduction. In recent years, several studies have focused on moment and probability inequalities for multilinear forms and symmetric statistics (see, in particular, Serfling (1980), Krakowiak and Szulga (1986), McConell and Taqqu (1986), de la Pena (1992), de la Pena and Klass (1994), Koroljuk and Borovskikh (1994), de la Pena and Montgomery-Smith (1995), Sharakhmetov (1995, 1997), Ibragimov and Sharakhmetov (1996a, 1999), Borovskikh and Korolyuk (1997), Ibragimov (1997), Klass and Nowicki (1997a, b) and Gine et. al. (2000)). Interest in such inequalities is motivated by their applications in limit theorems, multiple stochastic integration, harmonic analysis, operator theory, quantum mechanics, theory of income inequality and species' diversity measurement, etc. (see, in addition to the above-mentioned papers, Bonami (1970), Rosinski and Szulga (1982), Sjorgen (1982), Rosinski and Woyczynski (1984, 1986), Cambanis et al. (1985) and Kwapien and Woyczinski (1992)). Furthermore, the bounds on moments for symmetric statistics can also be applied in investment theory and in testing for chaos in time series data based on the notion of correlation integral, which has the form of symmetric statistics (see Cecen and Erkal (1996a, b)).

In the case of linear statistics (sums of independent random variables (r.v.'s)), the exact moment estimates are given by the well-known Khintchine, Marcinkiewicz-Zygmund and Rosenthal inequalities (see Khintchine (1923), Marcinkiewicz and Zygmund (1937), Rosenthal (1970)). Let us remind the latter ones $(A_i(\cdot), B_i(\cdot))$ denote constants depending on parameters in parentheses only).

Theorem 1. If $\xi_1, ..., \xi_n$ are independent mean zero r.v.'s with finite *t*th moment, $2 < t < \infty$, then

$$A_{1}(t)max\left(\sum_{i=1}^{n} E|\xi_{i}|^{t}, \left(\sum_{i=1}^{n} E\xi_{i}^{2}\right)^{t/2}\right) \leq E|\sum_{i=1}^{n} \xi_{i}|^{t} \leq B_{1}(t)max\left(\sum_{i=1}^{n} E|\xi_{i}|^{t}, \left(\sum_{i=1}^{n} E\xi_{i}^{2}\right)^{t/2}\right).$$
(1)

The exact upper constants in inequality (1) (case t=2m) and in its analogue for nonnegative r.v.'s were found in Ibragimov and Sharakhmetov (1996b, 1998a, b). The best constant in inequality (1) for symmetric r.v.'s was independently found by Figiel et al. (1997) and Ibragimov and Sharakhmetov (1995, 1997). The results obtained by Ibragimov and Sharakhmetov (1996b, 1997, 1998a, b) and their proofs were presented in Ibragimov (1997). Concerning refinements and extensions of Rosenthal's inequalities and related problems see also Prokhorov (1962), Nagaev and Pinelis (1977), Pinelis (1980, 1994), Pinelis and Utev (1984), Johnson et al. (1985), Utev (1985), Talagrand (1989), Hitczenko (1990, 1994), Nagaev (1990, 1998), Kwapien and Szulga (1991) and Peshkir and Shiryaev (1995).

Recently, Sharakhmetov (1995, 1997), Ibragimov and Sharakhmetov (1996a, 1998a, 1999, 2000) (see also Ibragimov (1997)), Klass and Nowicki (1997a, b) and Gine et. al. (2000) obtained analogues of Rosenthal's inequality (1) and its analogue for nonnegative r.v.'s in the case of symmetric statistics. Ibragimov and Sharakhmetov (2000) also showed the significance of each term in the analogues of Rosenthal's bounds for symmetric statistics of arbitrary order. Ibragimov (1997) showed that the best constants in the analogues of Rosenthal's inequalities grow not slower than $(t/\ln t)^m$, as $t \rightarrow \infty$, where *m* is the order of a symmetric statistic. Gine et. al. (2000) showed that the actual rate of growth of the above constants is $(t/\ln t)^m$.

The qualitative difference of the results on Rosenthal's inequalities for nonlinear statistics from the linear case is the exact constants in them are unknown yet. The main goal of the present paper is to fill partially this gap in the case of bilinear forms. More specifically, we obtain the explicit expressions for the best constant in the analogues of Rosenthal's inequalities for ordinary and decoupled bilinear forms in identically distributed symmetric r.v.'s in the case of fixed number of r.v.'s. The proof of the expressions for the best constants in the non-linear analogues of Rosenthal inequalities is based on a theorem, which extends the extremal results obtained in Utev (1985) and Ibragimov and Sharakhmetov (1996b, 1997) in the case of bilinear forms and gives the exact estimates for moments of random bilinear forms in terms of moment characteristics of their particular components. To our knowledge, this theorem and its proof are the first attempt to apply methods which were used to investigate the extremal problems in moment inequalities for sums of independent r.v.'s for non-linear statistics. The results obtained in the present paper can be extended to the case of nonnegative random variables, multilinear forms of arbitrary order and generalized moments; these extensions will be presented elsewhere.

2. Main results. Let t>2, $X_1, Y_1, X_2, Y_2, ..., X_n, Y_n$ be independent symmetric r.v.'s with finite *t*th moment. Let $a_i \ge 0$, $b_i \ge 0$, $c_i \ge 0$, $d_i \ge 0$, $a_i^t \le b_i$, $c_i^t \le d_i, i = 1, ..., n$. Set

$$(X, n) = (X_1, ..., X_n), (Y, n) = (Y_1, ..., Y_n)$$

$$M_1(n,a,b) = \{(X,n): EX_i^2 = a_i^2, E|X_i|^t = b_i, i = 1,...,n\}$$

$$M_1(n,c,d) = \{(Y,n) : EY_i^2 = c_i^2, E|Y_i|^t = d_i, i = 1,...,n\}$$

$$M_{2}(n,a,b) = \{(X,n) : EX_{i}^{2} \le a_{i}^{2}, E | X_{i} |^{t} \le b_{i}, i = 1,...,n\}$$

$$M_{2}(n,c,d) = \{(Y,n): EY_{i}^{2} \le c_{i}^{2}, E|Y_{i}|^{t} \le d_{i}, i = 1,...,n\}$$

Let $U_i(a_i, b_i, t)$, $V_i(c_i, d_i, t)$, i = 1, ..., n, be independent r.v.'s such that

$$P(U_i(a_i, b_i, t) = 0) = 1 - (a_i^t / b_i)^{2/(t-2)}$$

$$P(U_i(a_i, b_i, t) = \pm (b_i / a_i^2)^{1/(t-2)}) = (1/2)(a_i^t / b_i)^{2/(t-2)}$$

$$P(V_i(c_i, d_i, t) = 0) = 1 - (c_i^t / d_i)^{2/(t-2)}$$

$$P(V_i(c_i, d_i, t) = \pm (d_i / c_i^2)^{1/(t-2)}) = (1/2)(c_i^t / d_i)^{2/(t-2)}$$

and let U_i , V_i , i = 1, ..., n, be independent r.v.'s with distribution

$$P(U_i = \pm 1) = P(V_i = \pm 1) = 1/2, i = 1, ..., n$$

The following theorem extends the results obtained in Utev (1985) and Ibragimov and Sharakhmetov (1997) on the non-linear case and gives the explicit bounds for moments of random bilinear forms in terms of moment characteristics of their particular components.

Theorem 4. If 2 < *t* < 4, then

$$\sup_{\substack{(X,n)\in M_{k}(n,a,b)}} E \left| \sum_{1\leq i< j\leq n} X_{i}X_{j} \right|^{t} = \sum_{1\leq i< j\leq n} (b_{i} - a_{i}^{t})(b_{j} - a_{j}^{t}) + \sum_{i=1}^{n} (b_{i} - a_{i}^{t})E \left| \sum_{j\neq i}^{n} a_{j}U_{j} \right|^{t} + E \left| \sum_{1\leq i< j\leq n} a_{i}a_{j}U_{i}U_{j} \right|^{t}$$

$$(2)$$

$$\sup_{\substack{(X,n)\in M_{k}(n,a,b),\\(Y,n)\in M_{l}(n,c,d)}} E \left| \sum_{\substack{1\leq i,j\leq n,\ i\neq j}} X_{i}Y_{j} \right|^{t} = \sum_{\substack{1\leq i,j\leq n\\i\neq j}} (b_{i} - a_{i}^{t})(d_{j} - c_{j}^{t}) + \sum_{\substack{1\leq i,j\leq n\\i\neq j}}^{n} (d_{j} - c_{j}^{t})E \left| \sum_{\substack{i=1,\\i\neq j}}^{n} a_{i}U_{i} \right|^{t} + \sum_{i=1}^{n} (b_{i} - a_{i}^{t})E \left| \sum_{j=1,\ j\neq i}^{n} c_{j}V_{j} \right|^{t} + \left| E \left| \sum_{\substack{1\leq i,j\leq n,\ i\neq j}}^{n} a_{i}c_{j}U_{i}V_{j} \right|^{t},\ k, l = 1, 2$$

$$(3)$$

If $3 \le t < 4$, then

$$\inf_{(X,n)\in M_1(n,a,b)} E \left| \sum_{1 \le i < j \le n} X_i X_j \right|^t = E \left| \sum_{1 \le i < j \le n} U_i(a_i, b_i, t) U_j(a_j, b_j, t) \right|^t$$
(4)

$$\inf_{\substack{(X,n)\in M_1(n,a,b),\\(Y,n)\in M_1(n,c,d)}} E\left|\sum_{\substack{1\leq i,j\leq n,\\i\neq j}} X_i Y_j\right|^t = E\left|\sum_{\substack{1\leq i,j\leq n,\\i\neq j}} U_i(a_i,b_i,t)V_j(c_j,d_j,t)\right|^t$$
(5)

If $t \ge 4$, then

$$\sup_{(X,n)\in M_{k}(n,a,b)} E\left|\sum_{1\leq i< j\leq n} X_{i}X_{j}\right|^{t} = E\left|\sum_{1\leq i< j\leq n} U_{i}(a_{i},b_{i},t)U_{j}(a_{j},b_{j},t)\right|^{t}$$
(6)

$$\sup_{\substack{(X,n)\in M_{k}(n,a,b),\\(Y,n)\in M_{l}(n,c,d)}} E\left|\sum_{\substack{1\leq i,\ j\leq n,\\i\neq j}} X_{i}Y_{j}\right|^{t} = E\left|\sum_{\substack{1\leq i,\ j\leq n,\\i\neq j}} U_{i}(a_{i},b_{i},t)V_{j}(c_{j},d_{j},t)\right|^{t}, k, l = 1,2$$
(7)

$$\inf_{\substack{(X,n)\in M_1(n,a,b)}} E \left| \sum_{1\leq i< j\leq n} X_i X_j \right|^t = \sum_{1\leq i< j\leq n} (b_i - a_i^t)(b_j - a_j^t) + \\
+ \sum_{i=1}^n (b_i - a_i^t) E \left| \sum_{j=1, \ j\neq i}^n a_j U_j \right| + E \left| \sum_{1\leq i< j\leq n} a_i a_j U_i U_j \right|^t$$
(8)

$$\inf_{\substack{(X,n)\in M_{1}(n,a,b),\\(Y,n)\in M_{1}(n,c,d)}} E\left|\sum_{\substack{1\leq i,\ j\leq n,\\i\neq j}} X_{i}Y_{j}\right|^{t} = \sum_{\substack{1\leq i,\ j\leq n,\\i\neq j}} (b_{i} - a_{i}^{t})(d_{j} - c_{j}^{t}) + \sum_{\substack{1\leq i,\ j\leq n,\\i\neq j}}^{n} (d_{j} - c_{j}^{t})E\left|\sum_{\substack{i=1,\\i\neq j}}^{n} a_{i}U_{i}\right|^{t} + \sum_{\substack{i=1\\i\neq j}}^{n} (b_{i} - a_{i}^{t})E\left|\sum_{\substack{j=1,\\j\neq i}}^{n} c_{j}V_{j}\right|^{t} + E\left|\sum_{\substack{1\leq i,\ j\leq n,\\i\neq j}}^{n} a_{i}c_{j}U_{i}V_{j}\right|^{t} \right|$$
(9)

<u>Remark.</u> The expressions in relations (2)-(9) are of a simple structure and their values can be easily calculated for given sequences a_i , b_i , c_i , d_i , i = 1,...,n.

Let us fix t>2 and $n \ge 1$. From the results obtained in Ibragimov and Sharakhmetov (1999) and decoupling theorems for symmetric statistics (see McConell and Taqqu (1986) and de la Pena and Montgomery-Smith (1995)) it follows that for all independent identically distributed symmetric r.v.'s $X_1,...,X_n$, $\overline{X}_1,...,\overline{X}_n$ with finite *t*th moment the following Rosenthal-type inequalities are true ($C_n^2 = n(n-1)/2$):

$$E\left|\sum_{1\leq i< j\leq n} X_{i} X_{j}\right|^{t} \leq B_{4}(t,n) \max(C_{n}^{2}(E|X_{1}|^{t})^{2}, (C_{n}^{2})^{t/2}(EX_{1}^{2})^{t})$$
(10)

$$E\left|\sum_{1\leq i< j\leq n} X_{i} X_{j}\right|^{t} \leq B_{5}(t,n) \max\left(n^{2} \left(E\left|X_{1}\right|^{t}\right)^{2}, n^{t} \left(EX_{1}^{2}\right)^{t}\right)$$
(11)

$$E\left|\sum_{1\leq i\neq j\leq n} X_i \overline{X}_j\right|^t \leq B_6(t,n) \max(C_n^2(E|X_1|^t)^2, (C_n^2)^{t/2}(EX_1^2)^t)$$
(12)

$$E\left|\sum_{1\leq i\neq j\leq n} X_{i}\overline{X}_{j}\right|^{t} \leq B_{7}(t,n)\max(n^{2}(E|X_{1}|^{t})^{2},n^{t}(EX_{1}^{2})^{t})$$
(13)

The following theorems give the explicit expressions for the best constants in inequalities (10) and (11).

Theorem 5. The exact constant in inequality (10) is given by

$$B_{4}^{*}(t,n) = C_{n}^{2} (1/(C_{n}^{2})^{1/2} - 1/(C_{n}^{2})^{t/2})^{2} + (1/(C_{n}^{2})^{1/2} - 1/(C_{n}^{2})^{t/2})n/(C_{n}^{2})^{t/4} E \left| \sum_{i=2}^{n} U_{i} \right|^{t} + E \left| \sum_{1 \le i < j \le n} U_{i} U_{j} / (C_{n}^{2})^{1/2} \right|^{t}, 2 < t < 4$$

$$B_{4}^{*}(t,n) = E \left| \sum_{1 \le i < j \le n} U_{i} (1/(C_{n}^{2})^{1/4}, 1/(C_{n}^{2})^{1/2}, t) + U_{j} (1/(C_{n}^{2})^{1/4}, 1/(C_{n}^{2})^{1/2}, t) \right|^{t}, t \ge 4$$

$$(14)$$

Theorem 6. The exact constant in inequality (11) is given by

$$B_{5}^{*}(t,n) = C_{n}^{2}(1/n - 1/n^{t})^{2} + (1/n^{t/2} - 1/n^{3t/2 - 1})E\left|\sum_{i=2}^{n} U_{i}\right|^{t} + E\left|\sum_{1 \le i < j \le n} U_{i}U_{j} / n\right|^{t}, 2 < t < 4$$
(16)

$$B_{5}^{*}(t,n) = E \left| \sum_{1 \le i < j \le n} U_{i}(1/n^{1/2}, 1/n, t) U_{j}(1/n^{1/2}, 1/n, t) \right|^{t}, t \ge 4$$
(17)

Theorems 7 and 8 below give the explicit expressions for the exact constants in inequalities (12) and (13).

Theorem 7. The exact constant in inequality (12) is given by

$$B_{6}^{*}(t,n) = 2C_{n}^{2}(1/(C_{n}^{2})^{1/2} - 1/(C_{n}^{2})^{t/2})^{2} + +2(1/(C_{n}^{2})^{1/2} - 1/(C_{n}^{2})^{t/2})n/(C_{n}^{2})^{t/4}E\left|\sum_{i=2}^{n}U_{i}\right|^{t} + E\left|\sum_{1\leq i, j\leq n, i\neq j}U_{i}V_{j}/(C_{n}^{2})^{1/2}\right|^{t}, 2 < t < 4$$

$$(18)$$

 $B_6^*(t,n) =$

$$E \left| \sum_{1 \le i \ne j \le n} U_i (1/(C_n^2)^{1/4}, 1/(C_n^2)^{1/2}, t) \right|^{t} \times V_j (1/(C_n^2)^{1/4}, 1/(C_n^2)^{1/2}, t) \right|^{t}, t \ge 4$$
(19)

Theorem 8. The exact constant in inequality (13) is given by

$$B_{7}^{*}(t,n) = 2C_{n}^{2}(1/n - 1/n^{t})^{2} + 2(1/n^{t/2} - 1/n^{3t/2 - 1})E\left|\sum_{i=2}^{n} U_{i}\right|^{t} + E\left|\sum_{\substack{1 \le i, j \le n, \\ i \ne j}} U_{i}V_{j}/n\right|^{t}, 2 < t < 4$$

$$B_{7}^{*}(t,n) = E\left|\sum_{\substack{1 \le i, j \le n, \\ i \ne j}} U_{i}(1/n^{1/2}, 1/n, t)V_{j}(1/n^{1/2}, 1/n, t)\right|^{t},$$

$$t \ge 4$$

$$(20)$$

3. Preliminaries. Let us formulate some auxiliary steps needed for the proof of the theorems.

Lemma 1. If 2 < t < 4, $z_1 \ge 0$, $z_2 \in \mathbf{R}$, $a \ge 0$, $b \ge 0$, $a^t \le b$, X is a symmetric r.v. with $EX^2 \le a^2$, $E[X]^t \le b$, then

$$E |z_1 X + z_2|^t - bz_1^t \le E |az_1 U + z_2|^t - a^t z_1^t$$
(22)

<u>Proof.</u> It suffices to consider the case $z_1 \neq 0$. From Lemma 5 in Ibragimov and Sharakhmetov (1997) it follows that

$$E \left| X + z_2 / z_1 \right|^t - b \le E \left| aU + z_2 / z_1 \right|^t - a^t$$
(23)

Multiplying (23) by z_1^t we obtain (22).

Applying Lemma 7 in Ibragimov and Sharakhmetov (1997) and Lemmas 7.3 and 7.4 in Utev (1985) analogously to the proof of Lemma 1 above we easily obtain the following Lemmas 2-4.

Lemma 2. If $3 \le t < 4$, $z_1, z_2 \in \mathbf{R}$, $a \ge 0$, $b \ge 0$, $a^t \le b$, X is a symmetric r.v. with $EX^2 = a^2$, $E|X|^t = b$, then

$$E |z_1 X + z_2|^t \ge E |z_1 U(a, b, t) + z_2|^t$$

Lemma 3. If $t \ge 4$, $z_1, z_2 \in \mathbf{R}$, $a \ge 0$, $b \ge 0$, $a^t \le b$, X is a symmetric r.v. with $EX^2 \le a^2$, $E|X|^t \le b$, then

$$E |z_1 X + z_2|^t \le E |z_1 U(a, b, t) + z_2|^t$$

Lemma 4. If $t \ge 4$, $z_1 \ge 0$, $z_2 \in \mathbf{R}$, $a \ge 0$, $b \ge 0$, $a^t \le b$, X is a symmetric r.v. with $EX^2 = a^2$, $E|X|^t = b$, then

$$E |z_1 X + z_2|^t - bz_1^t \ge E |az_1 U + z_2|^t - a^t z_1^t.$$

Lemma 5. Let $1 \le k \le n$, $X_1, ..., X_{k-1}, U_k, X_{k+1}, ..., X_n$ be independent r.v.'s with $E|X_i|^t < \infty$, i = 1, ..., n, $i \ne k$, $a_k, b_k \ge 0$, $a_k^t \le b_k$, $c_i \in \mathbf{R}$, i = 1, ..., k-1, and let F_1 be the set of symmetric r.v.'s X_k independent of $X_1, ..., X_{k-1}, X_{k+1}, ..., X_n$ and satisfying the conditions $EX_k^2 \le a_k^2$, $E|X_k|^t \le b_k$, F_2 be the subset of F_1 consisting of r.v.'s X_k such that $EX_k^2 = a_k^2$, $E|X_k|^t = b_k$. If 2 < t < 4, then

$$\sup_{X_k \in F_l} \left(\sum_{i=1}^{k-1} c_i E \left| \sum_{\substack{j=1\\j \neq i}}^n X_j \right|^t + E \left| \sum_{\substack{1 \le i < j \le n}}^n X_i X_j \right|^t \right)$$

$$= \sum_{i=1}^{k-1} c_i E \left| a_k U_k + \sum_{\substack{j=1\\j\neq i,k}}^n X_j \right|^t + \sum_{i=1}^{k-1} c_i (b_k - a_k^t) +$$

$$+(b_{k}-a_{k}^{t})E\left|\sum_{\substack{j=1\\j\neq k}}^{n}X_{j}\right|^{t}+E\left|a_{k}U_{k}(\sum_{\substack{j=1\\j\neq k}}^{n}X_{j})+\sum_{\substack{1\leq i< j\leq n\\i,j\neq k}}X_{i}X_{j}\right|^{t},\ l=1,2$$

If $t \ge 4$, then

$$\inf_{\substack{X_k \in F_2 \\ X_k \in F_2}} \left(\sum_{i=1}^{k-1} c_i E \left| \sum_{\substack{j=1 \\ j \neq i}}^n X_j \right|^t + E \left| \sum_{1 \le i < j \le n} X_i X_j \right|^t \right)$$

$$= \sum_{i=1}^{k-1} c_i E \left| a_k U_k + \sum_{\substack{j=1\\j \neq i,k}}^n X_j \right|^t + \sum_{i=1}^{k-1} c_i (b_k - a_k^t) + (b_k - a_k^t) E \left| \sum_{\substack{j=1\\j \neq k}}^n X_j \right|^t + E \left| a_k U_k (\sum_{\substack{j=1\\j \neq k}}^n X_j) + \sum_{\substack{1 \le i < j \le n\\i, j \neq k}}^n X_i X_j \right|^t$$

<u>Proof.</u> From Lemmas 1 and 4 above and Lemma 5 in Ibragimov and Sharakhmetov (1997) it follows that it suffices to find a sequence of r.v.'s X_{mk} , m = 1, 2, ..., independent of $X_1, ..., X_{k-1}, X_{k+1}, ..., X_n$ and satisfying the conditions $EX_{mk}^2 = a_k^2, E \left| X_{mk} \right|^t = b_k$, $\lim_{m \to \infty} E \left| X_{mk} + \sum_{\substack{j=1\\j \neq i, k}}^n X_j \right|^t = E \left| a_k U_k + \sum_{\substack{j=1\\j \neq i, k}}^n X_j \right|^t + b_k - a_k^t, i = 1, ..., k-1$ (24)

$$\lim_{m \to \infty} E \left| X_{mk} (\sum_{\substack{j=1\\j \neq k}}^{n} X_j) + \sum_{\substack{1 \le i < j \le n\\i, j \neq k}}^{n} X_i X_j \right|^t = (b_k - a_k^t) E \left| \sum_{\substack{j=1\\j \neq k}}^{n} X_j \right|^t + E \left| a_k U_k (\sum_{\substack{j=1\\j \neq k}}^{n} X_j) + \sum_{\substack{1 \le i < j \le n\\i, j \neq k}}^{n} X_i X_j \right|^t$$
(25)

If
$$b_k = a_k^t$$
, then one can take $X_{mk} = a_k U_k$. Let $a_k^t < b_k$. Set $\delta_m = 1/m$,
 $P(X_{mk} = \pm a_k) = 1/2(1 - \delta_m)$, $P(X_{mk} = \pm b_{mk}) = 1/2\delta_{mk}^*$, $\delta_{mk}^* = a_k^2 \delta_m / b_{mk}^2$,
 $P(X_{mk} = 0) = \delta_m - \delta_{mk}^*$, $b_{mk} = ((b_k - a_k^t (1 - \delta_m))/a_k^2 \delta_m)^{1/(t-2)}$, $m = 1, 2, ...$

Then

$$b_{mk} \ge a_k, \ 0 \le \delta_{mk}^* \le \delta_m,$$

$$EX_{mk}^2 = a_k^2, \ E |X_{mk}|^t = b_k, \ m = 1, 2, \dots$$

$$\delta_m \to 0, \ b_{mk} \to \infty, \ b_{mk}^t \delta_{mk}^* \to b_k - a_k^t, \ m \to \infty$$
(26)

From (26) and the proof of Lemma 7.6 in Utev (1985) it follows that relations (24) are valid.

Let us prove that (25) is true. We have

$$E \left| X_{mk} \left(\sum_{\substack{j=1, \\ j \neq k}}^{n} X_{j} \right) + \sum_{\substack{1 \le i < j \le n, \\ i, j \neq k}}^{n} X_{i} X_{j} \right|^{t} = E \left| a_{k} U_{k} \left(\sum_{\substack{j=1, \\ j \neq k}}^{n} X_{j} \right) + \sum_{\substack{1 \le i < j \le n, \\ i, j \neq k}}^{n} X_{i} X_{j} \right|^{t} (1 - \delta_{m}) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k} \left(\sum_{\substack{j=1, \\ i, j \neq k}}^{n} X_{j} \right) + C \left| A_{k} U_{k}$$

$$+E\left|\sum_{\substack{1\leq i< j\leq n, \\ i, j\neq k}} X_{i}X_{j}\right|^{t} (\delta_{m} - \delta_{mk}^{*}) + (E\left|b_{mk}U_{k}(\sum_{\substack{j=1, \\ j\neq k}}^{n} X_{j}) + \sum_{\substack{1\leq i< j\leq n, \\ i, j\neq k}} X_{i}X_{j}\right|^{t} - b_{mk}^{t}E\left|\sum_{\substack{j=1, \\ j\neq k}}^{n} X_{j}\right|^{t})\delta_{mk}^{*} + b_{mk}^{t}\delta_{mk}^{*}E\left|\sum_{\substack{j=1, \\ j\neq k}}^{n} X_{j}\right|^{t}$$

From (26) it follows that for the proof of (25) it suffices to check that

$$(E \left| b_{mk} U_k \left(\sum_{\substack{j=1, \\ j \neq k}}^n X_j \right) + \sum_{\substack{1 \leq i < j \leq n, \\ i, j \neq k}} X_i X_j \right|^t - b_{mk}^t E \left| \sum_{\substack{j=1 \\ j \neq k}}^n X_j \right|^t) \delta_{mk}^* \to 0, \ m \to \infty$$

This follows from the fact that $b_{mk}^{t} \delta_{mk}^{*}$ converges and that, on the strength of the inequality $||x+y|^{t} - |x|^{t}| \le 2^{t} t(|x|^{t-1}|y|+|y|^{t}), x, y \in \mathbf{R}, t \ge 1$ (see Lemma 7.5. in Utev (1984)), and the dominated convergence principle,

$$\lim_{m \to \infty} E \left| U_k \left(\sum_{\substack{j=1, \\ j \neq k}}^n X_j \right) + \sum_{\substack{1 \le i < j \le n, \\ i, j \neq k}}^n X_i X_j / b_{mk} \right|^t = E \left| U_k \left(\sum_{\substack{j=1, \\ j \neq k}}^n X_j \right) \right|^t = E \left| \sum_{\substack{j=1, \\ j \neq k}}^n X_j \right|^t$$

Arguing analogously with the proof of Lemma 5, we easily obtain the following

Lemma 6. Let $1 \le k \le n$, $X_1, ..., X_{k-1}, U_k, X_{k+1}, ..., X_n, Y_1, ..., Y_n$ be independent r.v.'s with $E|X_i|^t < \infty$, i = 1, ..., n, $i \ne k$, $E|Y_i|^t < \infty$, i = 1, ..., n, $a_k, b_k \ge 0$, $a_k^t \le b_k$, $c_i \in \mathbf{R}$, i = 1, ..., k-1, and let G_1 be the set of symmetric r.v.'s X_k independent of $X_1, ..., X_{k-1}, X_{k+1}, ..., X_n, Y_1, ..., Y_n$ and satisfying the conditions $EX_k^2 \le a_k^2$, $E|X_k|^t \le b_k$, G_2 be the subset of G_1 consisting of r.v.'s X_k such that $EX_k^2 = a_k^2$, $E|X_k|^t = b_k$. If 2 < t < 4, then

$$\begin{split} \sup_{\substack{X_k \in G_l \ i=1}} & \left(\sum_{i=1}^n c_i E \left| \sum_{j=1, \ j \neq i}^n X_j \right|^t + E \left| \sum_{1 \le i \neq j \le n} X_i Y_j \right|^t \right) = \\ &= \sum_{i=1}^n c_i E \left| a_k U_k + \sum_{j=1, \ j \neq i, k}^n X_j \right|^t + \sum_{i=1}^n c_i (b_k - a_k^t) + \\ &+ (b_k - a_k^t) E \left| \sum_{j=1, \ j \neq k}^n Y_j \right|^t + E \left| a_k U_k (\sum_{j=1, \ j \neq k}^n Y_j) + \sum_{1 \le i \neq j \le n, \ i \neq k} X_i Y_j \right|^t, \ l=1, 2 \end{split}$$

If $t \ge 4$, then

$$\begin{split} \inf_{\substack{X_k \in G_2 \ i=1}} c_i E \left| \sum_{j=1, \ j \neq i}^n X_j \right|^t + E \left| \sum_{\substack{1 \le i \neq j \le n}}^n X_i Y_j \right|^t) = \\ &= \sum_{i=1}^n c_i E \left| a_k U_k + \sum_{j=1, \ j \neq i,k}^n X_j \right|^t + \sum_{i=1}^n c_i (b_k - a_k^t) + \\ &+ (b_k - a_k^t) E \left| \sum_{\substack{j=1 \ j \neq k}}^n Y_j \right|^t + E \left| a_k U_k (\sum_{j=1, \ j \neq k}^n Y_j) + \sum_{\substack{1 \le i \neq j \le n, \ i, j \neq k}}^n X_i Y_j \right|^t \end{split}$$

4. Proofs of the theorems.

Proof of theorem 3. Relations (4)-(7) easily follow from Lemmas 2 and 3 by induction. Let us prove (2). Let 2 < t < 4, $1 \le k \le n$, $U_1, ..., U_{k-1}, X_{k+1}, ..., X_n$ be independent symmetric r.v.'s, $E|X_i|^t < \infty$, i = k + 1, ..., n, $a_i \ge 0$, $b_i \ge 0$, $a_i^t \le b_i$, i = 1, ..., k. Denote by H_1 the set of symmetric r.v.'s X_k independent of $U_1, ..., U_{k-1}, X_{k+1}, ..., X_n$ and satisfying the conditions $EX_k^2 \le a_k^2$, $E|X_k|^t \le b_k$, and by H_2 the subset of H_1 consisting of r.v.'s X_k such that $EX_k^2 = a_k^2$, $E|X_k|^t = b_k$. On the strength of Lemma 5 we have

Applying (27) n times we get (2).

Let us show that (3) is valid. Let 2 < t < 4, $1 \le k \le n$, $U_1, ..., U_{k-1}, X_{k+1}, ..., X_n$, $Y_1, ..., Y_n$ be independent symmetric r.v.'s, $E|X_i|^t < \infty$, i = k + 1, ..., n, $E|Y_i|^t < \infty$, $i = 1, ..., n, a_i \ge 0, b_i \ge 0, a_i^t \le b_i, i = 1, ..., k$. Denote by K_1 the set of symmetric r.v.'s X_k independent of $U_1,..., U_{k-1}, X_{k+1},..., X_n, Y_1,..., Y_n$ and satisfying the conditions $EX_k^2 \le a_k^2, E |X_k|^t \le b_k$, and by K_2 the subset of K_1 consisting of r.v.'s X_k such that $EX_k^2 = a_k^2, E |X_k|^t = b_k$. From Lemma 6 with $c_i = 0, i = 1,...,n$, it follows that

$$\sup_{X_{k} \in K_{l}} \left(\sum_{i=1}^{k-1} (b_{i} - a_{i}^{t}) E \left| \sum_{\substack{j=1 \ j \neq i}}^{n} Y_{j} \right|^{t} + E \left| \sum_{i=1}^{k-1} a_{i} U_{i} \left(\sum_{\substack{j=1 \ j \neq i}}^{n} Y_{j} \right) + \sum_{i=k}^{n} X_{i} \left(\sum_{\substack{j=1 \ j \neq i}}^{n} Y_{j} \right) \right|^{t} \right) =$$

$$=\sum_{i=1}^{k} (b_{i} - a_{i}^{t}) E \left| \sum_{\substack{j=1\\j\neq i}}^{n} Y_{j} \right|^{t} + E \left| \sum_{i=1}^{k} a_{i} U_{i} (\sum_{\substack{j=1\\j\neq i}}^{n} Y_{j}) + \sum_{\substack{i=k+1\\j\neq i}}^{n} X_{i} (\sum_{\substack{j=1\\j\neq i}}^{n} Y_{j}) \right|^{t}, \ l = 1, 2$$
(28)

Using (28) n times we obtain

$$\sup_{\substack{(X,n) \in M_{k}(n,a,b) \\ i = 1}} E \left| \sum_{1 \le i \ne j \le n} X_{i} Y_{j} \right|^{t}} = \sum_{i=1}^{n} (b_{i} - a_{i}^{t}) E \left| \sum_{j=1, \ j \ne i}^{n} Y_{j} \right|^{t} + E \left| \sum_{1 \le i \ne j \le n} a_{i} U_{i} Y_{j} \right|^{t}, \quad k = 1, 2$$

$$(29)$$

Let $1 \le k \le n$, $U_1, \dots, U_n, V_1, \dots, V_{k-1}, Y_{k+1}, \dots, Y_n$ be independent symmetric

r.v.'s,
$$E|Y_i|^t < \infty$$
, $i = k + 1, ..., n$, $a_i \ge 0$, $b_i \ge 0$, $a_i^t \le b_i$, $i = 1, ..., n$, $c_i \ge 0$, $d_i \ge 0$,
 $c_i^t \le d_i$, $i = 1, ..., k$. Denote by B_1 the set of symmetric r.v.'s Y_k independent of

 $U_1, \dots, U_n, V_1, \dots, V_{k-1}, Y_{k+1}, \dots, Y_n$ and satisfying the conditions $EY_k^2 \le c_k^2$,

 $E|Y_k|^t \le d_k$, and by B_2 the subset of B_1 consisting of r.v.'s Y_k such that $EY_k^2 = c_k^2$, $E|Y_k|^t = d_k$.

Applying Lemma 6 again with $c_i = b_i - a_i^t$ we obtain

$$\sup_{Y_k \in B_l} \left(\sum_{i=1}^n (b_i - a_i^t) \left(\sum_{j=1, j \neq i}^{k-1} (d_j - c_j^t) \right) + \sum_{i=1}^n (b_i - a_i^t) E \left| \sum_{j=1, j \neq i}^{k-1} c_j V_j + \sum_{j=k, j \neq i}^n Y_j \right|^t + \sum_{j=1}^n \left(\sum_{j=1}^{k-1} (d_j - c_j^t) \right) + \sum_{i=1}^n (b_i - a_i^t) E \left| \sum_{j=1, j \neq i}^{k-1} (d_j - c_j^t) \right|^t + \sum_{i=1}^n \left(\sum_{j=1}^{k-1} (d_j - c_j^t) \right) + \sum_{i=1}^n (d_j - a_i^t) E \left| \sum_{j=1, j \neq i}^{k-1} (d_j - c_j^t) \right|^t + \sum_{i=1}^n \left(\sum_{j=1}^{k-1} (d_j - c_j^t) \right) + \sum_{i=1}^n (d_j - a_i^t) E \left| \sum_{j=1, j \neq i}^{k-1} (d_j - c_j^t) \right|^t + \sum_{i=1}^n (d_j - c_j^t) + \sum_{i=1}^n (d_$$

$$+ E \left| \sum_{j=1}^{k-1} c_j V_j \left(\sum_{i=1, i \neq j}^n a_i U_i \right) + \sum_{j=k}^n Y_j \left(\sum_{i=1, i \neq j}^n a_i U_i \right) \right|^t \right) =$$

$$=\sum_{i=1}^{n} (b_{i} - a_{i}^{t}) \left(\sum_{j=1, \ j \neq i}^{k} (d_{j} - c_{j}^{t})\right) + \sum_{i=1}^{n} (b_{i} - a_{i}^{t}) E \left|\sum_{j=1, \ j \neq i}^{k} c_{j} V_{j} + \sum_{j=k+1, \ j \neq i}^{n} Y_{j}\right|^{t} + E \left|\sum_{j=1}^{k} c_{j} V_{j} \left(\sum_{i=1, \ i \neq j}^{n} a_{i} U_{i}\right) + \sum_{j=k+1}^{n} Y_{j} \left(\sum_{i=1, \ i \neq j}^{n} a_{i} U_{i}\right)\right|^{t}, \ l=1, 2$$
(30)

Using (30) n times we get (3).

Relations (8) and (9) might be proven in the same way.

Proofs of theorems 4-8. Let us prove (14). Let $2 < t < 4, D \ge 0$, and let L(D) be a class of independent identically distributed r.v.'s X_1, \dots, X_n , for which

$$max(C_n^2(E|X_1|^t)^2, (C_n^2)^{t/2}(EX_1^2)^t) = D.$$

It is evident that

$$\sup_{\substack{(X,n) \in M_{1}(n,D^{1/2t} / (C_{n}^{2})^{1/4},D^{1/2} / (C_{n}^{2})^{1/2})}} E \left| \sum_{1 \le i < j \le n} X_{i} X_{j} \right|^{t}$$

$$\leq \sup_{\substack{(X,n) \in L(D) \\ (X,n) \in M_{2}(n,D^{1/2t} / (C_{n}^{2})^{1/4},D^{1/2} / (C_{n}^{2})^{1/2})}} E \left| \sum_{1 \le i < j \le n} X_{i} X_{j} \right|^{t}$$

$$(31)$$

From relation (2) and its proof it follows that

$$\sup_{(X,n) \in M_k(n,D^{1/2t} / (C_n^2)^{1/4},D^{1/2} / (C_n^2)^{1/2})} E \left| \sum_{1 \le i < j \le n} X_i X_j \right|^t =$$

$$= \left(C_{n}^{2}\left(1/(C_{n}^{2})^{1/2} - 1/(C_{n}^{2})^{t/2}\right)^{2} + \left(1/(C_{n}^{2})^{1/2} - 1/(C_{n}^{2})^{t/2}\right)n/(C_{n}^{2})^{t/4}E\left|\sum_{i=2}^{n}U_{i}\right|^{t} + E\left|\sum_{1\leq i< j\leq n}U_{i}U_{j}/(C_{n}^{2})^{1/2}\right|^{t}\right)D, \ k = 1, 2$$

$$(32)$$

(14) now follows from (31), (32) and the equality

$$B_{4}^{*}(t,n) = \sup_{D>0} (\sup_{(X,n) \in L(D)} E \left| \sum_{1 \le i < j \le n} X_{i} X_{j} \right|^{t} / D)$$

The remaining relations (15)-(21) might be proven in the similar way.

Acknowledgements. We are grateful to an anonymous referee and Victor de la Pena for many useful suggestions, which led to the improvement of the paper. We would like also to thank Jaksa Cvitanic, Victor de la Pena and other participants of the probability seminar in the Department of Statistics at Columbia University for an opportunity to present and discuss the results.

REFERENCES

Bonami, A. (1970). Etude des coefficients de Fourier des fonctions de $L_p(G)$. Ann. Inst. Fourier 20, 335-402.

Borovskikh, , Yu. V. and Korolyuk, V. S. (1997). *Martingale approximation*. VSP, Utrecht, 322 pp.

Cambanis, S., Rosinski, J. and Woyczynski, W.A. (1985). Convergence of quadratic forms in *p*-stable random variables and θ_p -radonifying operators. *Ann. Probab.* **13**, 885-897.

Cecen, A. A. and Erkal, C. (1996a). Distinguishing between stochastic and deterministic behavior in high frequency foreign exchange rate returns: Can non-linear dynamics help forecasting? *International J. Forecast.* **12**, 465-473.

Cecen, A. A. and Erkal, C. (1996b). Distinguishing between stochastic and deterministic behavior in high frequency exchange rate returns: Further evidence. *Economics Letters* **51**, 323-329.

De la Pena, V. H. (1992). Decoupling and Khintchine's inequalities for *U*-statistics. *Ann. Probab.* **20**, 1877-1892.

De la Pena, V. H. and Klass, M. J. (1994). Order of magnitude bounds for expectations involving quadratic forms. *Ann. Probab.* **22**, 1044-77.

De la Pena, V. H. and Montgomery-Smith (1995). Decoupling inequalities for the tail probabilities of multivariate *U*-statistics. *Ann. Probab.* **23**, 806-816.

Figiel, T., Hitczenko, P., Johnson, W. B., Schechtman, G. and Zinn, J. (1997). Extremal properties of Rademacher functions with applications to the Khintchine and Rosenthal inequalities. *Trans. Amer. Math. Soc.*, **349**, 997-1027.

Gine, E., Latala, R. and Zinn, J. (2000). Exponential and moment inequalities for Ustatistics. High Dimensional Probability II - Progress in Probability, Birkhauser, 13-35.

Hitczenko, P. (1990). Best constants in martingale version of Rosenthal's inequality. *Ann. Probab.* **18**, 1656-1668.

Hitczenko, P. (1994). On a domination of sums of random variables by sums of conditionally independent ones. *Ann. Probab.* **22**, 453-468.

Ibragimov, R. (1997). *Estimates for the moments of symmetric statistics*. Ph.D. Dissertation.
Institute of Mathematics of Uzbek Academy of Sciences, Tashkent, 127 pp. (in Russian).
Ibragimov, R. and Sharakhmetov, Sh. (1995). On the best constant in Rosenthal's inequality.
In: *Theses of reports of the conference on probability theory and mathematical statistics dedicated to the 75th anniversary of Academician S. Kh. Sirajdinov (Fergana, Uzbekistan).*Tashkent, 43-44 (in Russian).

Ibragimov, R. and Sharahmetov, Sh. (1996a). Bounds for the moments of symmetric statistics. Submitted to "Studia Scientiarum Mathematicarum Hungarica".

Ibragimov, R. and Sharakhmetov, Sh. (1996b). Some extremal problems in moment inequalities. To be published in *Theory Probab. Appl.*

Ibragimov, R. and Sharakhmetov, Sh. (1997). On an exact constant for the Rosenthal inequality. *Teor. Veroyatnost. i Primen.* **42**, 341-350 (translation in *Theory Probab. Appl.* **42** (1997), 294-302 (1998)).

Ibragimov, R. and Sharakhmetov, Sh. (1998a). Exact bounds on the moments of symmetric statistics. *7th Vilnius Conference on Probability Theory and Mathematical Statistics*. *22nd European Meeting of Statisticians*. *Abstracts of communications*. Vilnius, Lithuania, 243-244.

Ibragimov, R. and Sharakhmetov, Sh. (1998b). The best constant in Rosenthal's inequality for random variables with zero mean. To be published in *Theory Probab. Appl.* Ibragimov, R. and Sharakhmetov, Sh. (1999). Analogues of Khintchine, Marcinkiewicz-Zygmund and Rosenthal inequalities for symmetric statistics. *Scand. J. Statist.* 26, 621-633. Ibragimov, R. and Sharakhmetov, Sh. (2000). Moment inequalities for symmetric statistics. In: "Modern Problems of Probability Theory and Mathematical Statistics. Proceedings of the 4th Fergana International Colloquium on Probability Theory and Mathematical Statistics held 27-29 September, 1995," Tashkent, pp. 184-193 (in Russian); also available at http://front.math.ucdavis.edu/math.PR/0005004. Johnson, W. B., Schechtman, G. and Zinn, J. (1985). Best constants in moment inequalities

for linear combinations of independent and exchangeable random variables. *Ann. Probab.* **13**, 234-253. Khintchine, A. (1923). Uber dyadische Bruche. Math. Z. 18, 109-116.

Klass, M. J. and Nowicki, K. (1997a). Order of magnitude bounds for expectations of Δ_2 -functions of generalized bilinear forms. To appear in *Probab. Theory Related Fields*.

Klass, M. J. and Nowicki, K. (1997b). Order of magnitude bounds for expectations of Δ_2 -functions of nonnegative random bilinear forms and generalized *U*-statistics. *Ann. Probab.* **25**, 1471-1501.

Koroljuk, V. S. and Borovskich, Yu. V. (1994). *Theory of U-statistics*. Mathematics and its Applications, **273**. Kluwer Academic Publishers Group, Dordrecht, 552 pp.

Krakowiak, W. and Szulga, J. (1986) Random multilinear forms. *Ann. Probab.* **14**, 955-973.

Kwapien, S. and Szulga, J. (1991). Hypercontraction methods in moment inequalities for series of independent random variables in normed spaces. *Ann. Probab.* **19**, 1-8.

Kwapien, S. and Woyczynski, W. (1992). *Random series and Stochastic Integrals: Single and Multiple*. Burkhauser, Boston, 360 pp.

Marcinkiewicz, J. and Zygmund, A. (1937). Sur les fonction independantes. *Fund. Math.* **29**, 60-90.

McConnell, T.R. and Taqqu, M. (1986). Decoupling inequalities for multilinear forms in independent symmetric random variables. *Ann. Probab.* **14**, 943-954.

Nagaev, S. V. (1990). On a new approach to the study of the distribution of a norm of a random element in Hilbert space. *Probability Theory and Mathematical Statistics, Proceedings of the Fifth Vilnius Conference,* Mosklas/VSP, Vilnius/Utrecht, 214-226.

Nagaev, S. V. (1998). Some refinements of probabilistic and moment inequalities. *Theory Probab. Appl.* **42**, 707-713.

Nagaev, S. V. and Pinelis, I. F. (1977). Some inequalities for the distributions of sums of independent random variables. *Theory of Probab. Appl.* **22**, 248-256.

Peshkir, G. and Shiryaev, A. N. (1995). Khintchine's inequalities and a martingale extension of the area of their action. *Uspekhi Mat. Nauk* **50**, No. 5, 3-62 (in Russian).

Pinelis, I. F. (1980). Estimates for moments of infinite-dimensional martingales. *Math. Notes* **27**, 459-462.

Pinelis, I. (1994). Extremal probabilistic problems and Hotteling's T^2 test under a symmetry condition. *Ann. Probab.* **22**, 357-368.

Pinelis, I. F. and Utev, S. A. (1984). Estimates of moments of sums of independent random variables. *Theory Probab. Appl.* **29**, 574-577.

Prokhorov, Yu. V. (1962). Extremal problems in limit theorems. In *Proc. VI All-Union Conference on Probability Theory and Mathematical Statistics*, Vilnius, 77-84 (in Russian).

Rosenthal, H. P. (1970). On the subspaces of L^p (p>2) spanned by sequences of independent random variables. *Israel J. Math.* **8**, 273-303.

Rosinski, J. and Szulga, J. (1982). Product random measures and double stochastic

integrals. Lecture Notes in Math. 939, 181-199. Springer, Berlin-New York.

Rosinski, J. and Woyczynski, W. A. (1984). Products of random measures, multilinear forms and multiple stochastic integrals. *Lecture Notes in Math.* **1089**, 294-315. Springer, Berlin-New York.

Rosinski, J. and Woyczynski, W. A. (1986). On Ito stochastic integration with respect to *p*-stable motion: Inner clock, integrability of sample paths, double and multiple integrals. *Ann. Probab.* **14**, 271-286.

Serfling, R. J. (1980). Approximation theorems of mathematical statistics. New York: Wiley, 371 p.

Sharakhmetov, Sh. (1995). Estimates for moments of symmetric statistics. In: *Theses of reports of the conference on probability theory and mathematical statistics dedicated to the* 75th anniversary of Academician S. Kh. Sirajdinov (Fergana, Uzbekistan). Tashkent, p. 119 (in Russian).

Sharakhmetov, Sh. (1997). *General representations for a joint distribution of random variables and their applications*. Doctor of Sciences Dissertation. Institute of Mathematics of Uzbek Academy of Sciences, 229 pp. (in Russian).

Sjorgen, P. (1982). On the convergence of bilinear and quadratic forms in independent random variables. *Studia Math.* **71**, 285-296.

Talagrand (1989). Isoperimetry and integrability of the sum of independent Banach-space valued random variables. *Ann. Probab.* **17**, 1546-1570.

Utev, S. A. (1985). Extremal problems in moment inequalities. In *Proc. Mathematical Institute of the Siberian Branch of the USSR Academy of Sciences*, **5**, 56-75 (in Russian).