

# AN INFINITE ANALOGUE OF RINGS WITH STABLE RANK ONE

PERE ARA, GERT K. PEDERSEN  
and FRANCESC PERERA

October 1999

**ABSTRACT.** Replacing invertibility with quasi-invertibility in Bass' first stable range condition we discover a new class of rings, the  $QB$ -rings. These constitute a considerable enlargement of the class of rings with stable rank one ( $B$ -rings), and include examples like  $\text{End}_{\mathbb{F}}(V)$ , the ring of endomorphisms of a vector space  $V$  over some field  $\mathbb{F}$ , and  $\mathbb{B}(\mathbb{F})$ , the ring of all row- and column-finite matrices over  $\mathbb{F}$ .

We show that the category of  $QB$ -rings is stable under the formation of corners, ideals and quotients, as well as matrices and direct limits. We also give necessary and sufficient conditions for an extension of  $QB$ -rings to be a  $QB$ -ring, and show that extensions of  $B$ -rings often lead to  $QB$ -rings. Specializing to the category of exchange rings we characterize the subset of exchange  $QB$ -rings as those in which every von Neumann regular element extends to a maximal regular element, i.e. a quasi-invertible element. Finally we show that the  $C^*$ -algebras that are  $QB$ -rings are exactly the extremally rich  $C^*$ -algebras studied by L.G. Brown and the second author.

## 1. Introduction

According to Bass, see e.g. [6], a unital ring  $R$  will have  $n$  in its *stable range* if whenever  $(a_0, \dots, a_n)$  is a left unimodular row, i.e.  $Ra_0 + \dots + Ra_n = R$ , there is a unimodular row  $(b_1, \dots, b_n)$  of the form  $b_k = a_k + y_k a_0$  for some elements  $y_k$  in  $R$ ,  $1 \leq k \leq n$ . The smallest  $n$  in the Bass range is – through a fortuitous error in translation from English to Russian and back – called the *stable rank* of  $R$ . This number, denoted by  $\text{bsr}(R)$ , is important in the non-stable  $K$ -theory of  $R$ , since the natural morphisms  $GL_n(R)/E_n(R) \rightarrow K_1(R)$ , where  $E_n(R)$  denotes the

---

1991 *Mathematics Subject Classification.* Primary 16A12; Secondary 16A30, 16A32.

*Key words and phrases.* Bass stable rank, quasi-invertible element, von Neumann regularity, exchange ring, semi-prime ring, extremally rich  $C^*$ -algebras.

Research supported by the European Community (HCM Programme), the Danish Research Council (SNF), the DGESIC (Spain) and the Comissionat per Universitats i Recerca de la Generalitat de Catalunya.

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

multiplicative group generated by the elementary  $n \times n$ -matrices, will be surjective whenever  $n \geq \text{bsr}(R) + 1$  and injective whenever  $n \geq \text{bsr}(R) + 2$ , cf. [6, Theorem 2.1].

Specializing to rings with  $\text{bsr}(R) = 1$  we see that they are characterized by the condition that whenever  $Ra + Rb = R$ , then  $R(a + yb) = R$  for some  $y$  in  $R$ . But in such a  $B$ -ring (to use a terminology suggested by Vaserstein, [29]) any left (or right) invertible element is actually invertible, cf. [29, Theorem 2.6], so the definition of a  $B$ -ring can be simplified to the demand:

- (\*) Given  $a, b$  and  $x$  in  $R$  such that  $xa + b = 1$ ,  
there is a  $y$  in  $R$  such that  $a + yb \in R^{-1}$ .

$B$ -rings have many pleasant properties, notably the cancellation property for finitely generated projective  $R$ -modules, which states that if  $M, N$  and  $P$  are  $R$ -modules and  $P$  is projective and finitely generated, then  $M \oplus P$  is isomorphic to  $N \oplus P$  if and only if  $M$  is isomorphic to  $N$ . Also it is clear that the unit can not be equivalent to any other idempotent, so these rings are “finite”. Through the work of Rieffel, cf. [28], the Bass stable range was introduced in  $C^*$ -algebra theory, and linked with Čech’s covering dimension; and  $C^*$ -algebras having stable rank one were identified with those unital  $C^*$ -algebras  $A$  for which the set  $A^{-1}$  of invertible elements is dense in  $A$ . In particular, the algebra  $C(X)$  of complex functions on a compact Hausdorff space  $X$  is a  $B$ -ring if and only if  $\dim X \leq 1$ .

Replacing the set of units  $A^{-1}$  in a  $C^*$ -algebra  $A$  with the union of left and right invertible elements as in [24], and more generally (and technically much more demanding) with the set  $A_q^{-1}$  of quasi-invertible elements in  $A$ , a theory of so-called *extremally rich*  $C^*$ -algebras has been developed in [8], [9], [10], [11] and [12]. These algebras are characterized topologically by the fact that the quasi-invertible elements form a dense set, but one of the equivalent conditions is a version of Bass’ first stable range condition, cf. [8, Theorem 3.3]. This we show to be equivalent with condition (\*) above, when  $R^{-1}$  replaced by the set  $R_q^{-1}$  of quasi-invertible elements, see Proposition 9.1.

In the present work we systematically build a theory for rings that satisfy the condition

- (\*\*) Given  $a, b$  and  $x$  in  $R$  such that  $xa + b = 1$ ,  
there is a  $y$  in  $R$  such that  $a + yb \in R_q^{-1}$ .

These we call  $QB$ -rings. Our aim, as in the  $C^*$ -algebra project mentioned above, is to extend as much as possible of the theory of  $B$ -rings to the much larger class of  $QB$ -rings. In this paper we concentrate on the categorical properties of  $QB$ -rings. In subsequent papers we plan to show that the class of  $QB$ -rings is the proper carrier for non-stable  $K$ -theory and for an index theory, cf. [27], including a generalized index set, cf. [10].

In §2 we define quasi-invertibility, and prove that it implies a strong form of von Neumann regularity. In fact, every quasi-invertible element is maximal in the natural order (by extension) on von Neumann regular elements. These relations are treated in some detail in order, we hope, to convince the reader that quasi-invertibility is an important concept linking von Neumann regularity with invertibility. We then in §3 define  $QB$ -rings by condition (\*\*) above, and show that it

is left-right symmetric. It is easy to see that the  $QB$ -property passes to quotients, but in order to show that it passes to ideals we need to reformulate the concept of quasi-invertibility to make sense for non-unital rings. This is done in Section 4 by introducing quasi-adversibility in exactly the same manner as Kaplansky used adversibility instead of invertibility (although not quite using that name).

In Section 5 we study “skew corners”, i.e. subsets of the form  $pRq$ , where  $p$  and  $q$  are idempotents in  $R$ . We develop a suitable notion of  $QB$ -corner in order to prove in Section 6 that the class of  $QB$ -rings is stable under matrix formation and under Morita equivalence.

We establish in Section 7 necessary and sufficient conditions for an extension of  $QB$ -rings to be again a  $QB$ -ring. In particular we obtain easily verifiable sufficient conditions, when one of the rings in an extension is a  $B$ -ring. One of the motivating examples here is, of course, the algebra of compact perturbations of Toeplitz operators, which is known to be an extension of two  $B$ -rings, viz. the algebra of compact operators and the algebra of functions on the unit circle, but which fails to be a  $B$ -ring itself. However, it is a  $QB$ -ring. Some algebraic analogues of the Toeplitz algebra are also presented.

Specializing in Section 8 to exchange rings we show how the quasi-invertible elements are precisely the maximally extended von Neumann regular elements. This leads to a characterization of  $QB$ -exchange rings as exactly those exchange rings in which every von Neumann regular element can be extended to a maximal regular element. Finally in Section 9 we prove that a  $C^*$ -algebra is a  $QB$ -ring if and only if it is *extremally rich* as defined in [8]. This means that we can use all the examples described in that and subsequent papers, in particular we can show that our extension results are best possible.

## 2. Quasi-Invertibility

**2.1. Definitions.** In this section  $R$  will denote a unital ring. We say that two elements  $x$  and  $y$  in  $R$  are *centrally orthogonal*, in symbols  $x \perp y$ , if  $xRy = 0$  and  $yRx = 0$ . Similarly, two subsets  $I$  and  $J$  of  $R$  are centrally orthogonal if  $IRJ = 0 = JRI$ . If  $I$  and  $J$  are actually ideals in  $R$ , then  $I \perp J$  simply means that  $IJ = 0 = JI$ . We shall refer to this situation by saying that  $I$  and  $J$  are *orthogonal ideals*. Note that this does not necessarily mean that  $I \cap J = 0$ , only that  $I \cap J$  is contained in the prime radical of  $R$ .

We have defined our orthogonality relations above so that they will apply in any ring. In important examples  $R$  will be at least semi-prime if not semi-primitive. In that case  $xRx = 0$  implies that  $x = 0$  for any element  $x$  in  $R$ . Semi-primeness of  $R$  is also equivalent to the demand that  $I^2 = 0$  implies  $I = 0$  for any ideal  $I$  in  $R$ , so that for any pair of ideals  $I$  and  $J$ , the three conditions  $IJ = 0$ ,  $JI = 0$  and  $I \cap J = 0$  are all equivalent. In particular,  $xRy = 0 \iff yRx = 0$  for any pair of elements  $x, y$  in a semi-prime ring  $R$ .

An element  $u$  in an arbitrary ring  $R$  is said to be *quasi-invertible* if there exist elements  $a, b$  in  $R$  such that

$$(*) \quad (1 - ua) \perp (1 - bu) .$$

The set of quasi-invertible elements in  $R$  will be denoted by  $R_q^{-1}$ . Similarly, the sets of left invertible, right invertible and two-sided invertible elements will be denoted by  $R_\ell^{-1}$ ,  $R_r^{-1}$  and  $R^{-1}$ , respectively.

If  $u \in R_q^{-1}$ , and if  $I$  and  $J$  denote the ideals of  $R$  generated by  $1 - ua$  and  $1 - bu$ , respectively, then evidently  $u + I \in (R/I)_r^{-1}$  and  $u + J \in (R/J)_\ell^{-1}$ , whereas  $I \perp J$  by (\*).

Conversely, if  $I$  and  $J$  are orthogonal ideals in  $R$ , and  $u \in R$  such that  $u + I \in (R/I)_r^{-1}$  and  $u + J \in (R/J)_\ell^{-1}$ , then  $1 - ua \in I$  and  $1 - bu \in J$  for some elements  $a, b$  in  $R$ , whence  $u \in R_q^{-1}$ . These conditions, therefore, furnish an equivalent definition of quasi-invertibility. In particular we see that if  $R$  is a prime ring, then  $R_q^{-1} = R_\ell^{-1} \cup R_r^{-1}$ .

Recall that an element  $a$  in a ring  $R$  is *von Neumann regular* if  $a = axa$  for some  $x$  in  $R$ . We say that  $x$  is a *partial inverse* for  $a$ . Replacing if necessary  $x$  with  $xax$  we may assume that  $xax = x$ , so that  $a$  is also a partial inverse for  $x$ . Note that  $p = ax$  and  $q = xa$  are idempotents in  $R$  satisfying  $pR = aR$  and  $Rq = Ra$ . Conversely, if  $p$  is an idempotent in  $R$  such that  $aR = pR$ , then  $a$  is von Neumann regular, and if  $p = ax$  then  $x$  is a partial inverse for  $a$ .

If  $u \in R_q^{-1}$ , then by (\*) we have the equation  $(1 - ua)u(1 - bu) = 0$ . Taking  $v = a + b - aub$  this implies that  $u = uvu$ , so that  $u$  is von Neumann regular in  $R$  with partial inverse  $v$ . However, being quasi-invertible is much more than just having a partial inverse, i.e. being von Neumann regular. For this reason (and with due apologies to previous authors) we will in this paper reserve the name *quasi-inverse* for an element satisfying the stronger conditions in Proposition 2.2. By computation  $1 - uv = (1 - ua)(1 - ub)$  and  $1 - vu = (1 - au)(1 - bu)$ , so that we have the relation

$$(**) \quad (1 - uv) \perp (1 - vu) .$$

Moreover, replacing if necessary  $v$  by  $vuv$ , we see that  $v$  is also von Neumann regular with partial inverse  $u$ . This replacement will not affect the orthogonality relations, since  $1 - u(vuv) = (1 - uv)(1 + uv)$ .

We summarize our observations in the following statement:

**2.2. Proposition.** *Each element  $u$  in  $R_q^{-1}$  is von Neumann regular, and we may choose a quasi-inverse  $v$  for  $u$  such that  $u$  and  $v$  are partial inverses for each other and the two idempotents  $1 - uv$  and  $1 - vu$  are centrally orthogonal in the sense of (\*\*). In particular,  $v \in R_q^{-1}$ .  $\square$*

Note that if  $u \in R_\ell^{-1}$  with left inverses  $v$  and  $v'$ , then  $v' = v + v'(1 - uv)$ , and any element of the form  $v + a(1 - uv)$  will be a left inverse for  $u$ . In particular, we need not have  $v' = v$ . We do not, therefore, expect any unicity for the quasi-inverses of elements in  $R_q^{-1}$ , so the relations described in the following result are actually much more powerful than one might have expected.

**2.3. Theorem.** *If  $u$  and  $v$  are elements in  $R_q^{-1}$  and quasi-inverses for each other, so that  $(1 - uv) \perp (1 - vu)$ , then each element of the form*

$$(*) \quad v' = v + a(1 - uv) + (1 - vu)b ,$$

with  $a, b$  in  $R$ , will be a quasi-inverse for  $u$  in  $R_q^{-1}$  and satisfy the relations

$$(1 - uv') \perp (1 - v'u), \quad (1 - uv') \perp (1 - vu), \quad (1 - uv) \perp (1 - v'u).$$

Conversely, if  $v'$  is any partial inverse for  $u$ , then  $v' \in R_q^{-1}$  and has the form  $(*)$  with  $a = b = v'$ . Moreover,

$$(**) \quad 1 - uv = (1 - uv)(1 - uv') \quad \text{and} \quad 1 - uv' = (1 - uv')(1 - uv),$$

and similarly for the  $vu$  and  $v'u$  products. In particular, the idempotents  $1 - uv$  and  $1 - uv'$  are Murray-von Neumann equivalent, as are  $1 - vu$  and  $1 - v'u$ , so there are orthogonal ideals  $I$  and  $J$  in  $R$ , such that  $uv = 1 = uv'$  modulo  $I$  and  $vu = 1 = v'u$  modulo  $J$ .

*Proof.* Evidently each  $v'$  in  $R$  of the prescribed form  $(*)$  will be a partial inverse for  $u$ . Moreover,

$$1 - uv' = (1 - ua)(1 - uv) \quad \text{and} \quad 1 - v'u = (1 - vu)(1 - bu),$$

whence  $(1 - uv') \perp (1 - v'u)$ , so that  $v' \in R_q^{-1}$  and  $v'$  is a quasi-inverse for  $u$ .

Conversely, if  $uv'u = u$  for some  $v'$  in  $R$  then

$$0 = (1 - vu)v'(1 - uv) = v' - vuv' - v'uv + v.$$

Consequently,

$$v' = v + 2v' - vuv' - v'uv = v + (1 - vu)v' + v'(1 - uv),$$

as desired, and  $v' \in R_q^{-1}$  with  $(1 - uv') \perp (1 - v'u)$  by the first part of the proof.

The equations in  $(**)$  follow by straightforward computations. The equivalence between the idempotents  $1 - uv$  and  $1 - uv'$  is evident from  $(**)$ , and therefore  $1 - uv$  and  $1 - uv'$  generate the same ideal  $I$ . Similarly,  $1 - vu$  and  $1 - v'u$  generate the same ideal  $J$ . These ideals are orthogonal since  $(1 - uv) \perp (1 - vu)$ , and evidently the desired relations are satisfied in  $R/I$  and  $R/J$ , respectively.  $\square$

**2.4. Definitions.** If  $a$  and  $b$  are von Neumann regular elements in  $R$  we say that  $b$  extends  $a$ , and write  $a \leq b$ , if

$$(*) \quad a = axb = bxa = axa$$

for some  $x$  in  $A$ . Thus  $x$  is a partial inverse for  $a$ , and we may assume that  $a$  is also a partial inverse for  $x$ .

Taking  $p = ax$  and  $q = xa$  in the equations above we see from  $(*)$  that  $p$  and  $q$  are idempotents in  $R$  such that  $pb = bq = a$ . Moreover,  $pR = aR$  and  $Rq = Ra$ , since  $x$  is a partial inverse for  $a$ . Conversely, if  $a \in R$  and if we can find idempotents  $p$  and  $q$ , and a von Neumann regular element  $b$ , such that

$$(**) \quad pR = aR \quad \text{and} \quad Rq = Ra, \quad \text{and moreover} \quad pb = bq = a,$$

then  $a$  is von Neumann regular and  $a \leq b$ . Thus,  $(**)$  furnishes an alternative description of the relation  $\leq$ . Observe from this that if  $a \leq b$  and  $aR = bR$  (or if  $Ra = Rb$ ), then  $a = b$  (because then  $pR = bR$ , so  $b = pb = a$ ).

The idempotents  $p$  and  $q$ , above, depend not only on  $a$  but also on  $x$ . Nevertheless we may think of them as the “range” and the “source” of  $a$ , noting that  $a$ , as a left multiplier on  $R$ , is a bijection from  $qR$  onto  $pR$  with kernel  $(1 - q)R$  and cokernel  $(1 - p)R$ . In this setting  $a \leq b$  expresses an ordinary extension of operators.

If  $y$  is a partial inverse for  $b$ , with  $b$  as its partial inverse, then from  $(*)$  we derive the simple relations

- (i)  $ax = bx, xa = xb$ .
- (ii)  $a = ayb = bya$ .
- (iii)  $a = bxb = aya$ .

Now put  $x' = yay$  and check that

$$ax'a = a, \quad x'ax' = x', \quad x' = x'ay = yax'.$$

Thus, replacing if necessary  $x$  with  $x'$ , we may assume that also  $x \leq y$ .

This implies that  $\leq$  is a transitive relation. For if  $a \leq b$  and  $b \leq c$  we may assume that  $a$  and  $b$  have partial inverses  $x$  and  $y$ , respectively, such that  $x \leq y$  as above. Then

$$cxa = c(yax)a = (cy)(axa) = (by)a = a,$$

and similarly  $axc = a$ , so  $a \leq c$ .

If  $a \leq b$  and  $b \leq a$  then in particular  $aR = bR$ , whence  $a = b$ . We summarize our observations in the following result:

**2.5. Proposition.** *The relation  $\leq$  defined above on the set  $R^r$  of von Neumann regular elements in a ring  $R$  gives a partial order on  $R^r$ .  $\square$*

**2.6. Lemma.** *Let  $R$  be a unital ring, and let  $a$  be a regular element in  $R$ . Write  $aR = pR$  and  $Ra = Rq$  for some idempotents  $p$  and  $q$ . If  $b$  is a regular element such that  $pb = a = bq$ , and if  $q'$  is an idempotent satisfying  $Rq = Rq'$ , then there is an element  $b'$  such that  $Rb = Rb'$ ,  $bR = b'R$ , and  $pb' = a = b'q'$ .*

*Proof.* Let  $b' = a + (1 - p)b(1 - q')$ . Clearly then,  $pb' = pa = a$  and  $b'q' = aq' = a$ . Note that we can also write  $b' = a + b(1 - q') - pb(1 - q') = a + b(1 - q') = bq + b(1 - q')$ . Since  $qq' = q$  and  $q'q = q'$ , it follows that  $w = q + (1 - q')$  is invertible with  $w^{-1} = q' + (1 - q)$ . From above we have that  $b' = bw$ , whence  $b = b'w^{-1}$ , and thus  $bR = b'R$ .

Since  $Rq' = Ra \subset Rb$ , we have that  $b' = a + (1 - p)b - (1 - p)bq' \in Rb$ , whence  $Rb' \subset Rb$ . Write  $q' = ta$ , with  $q'tp = t$ , and compute that

$$\begin{aligned} (bt + (1 - p))b' &= (bt + (1 - p))(a + (1 - p)b(1 - q')) \\ &= bta + (1 - p)b(1 - q') = bq' + b(1 - q') = b, \end{aligned}$$

so  $Rb \subset Rb'$ , hence  $Rb = Rb'$ .  $\square$

**2.7. Lemma.** *If  $a \leq b$  in  $R^r$  there exist idempotents  $p$  and  $q$  in  $R$  such that  $pb = a = bq$  and  $b - a \in (1 - p)R(1 - q)$ . Conversely, if  $a$  and  $c$  are elements in  $R^r$ , where  $a \in pRq$  and  $c \in (1 - p)R(1 - q)$  for some idempotents  $p$  and  $q$  in  $R$ , then  $a \leq a + c$  in  $R^r$ .*

*Proof.* By  $(**)$  in 2.4 we have idempotents  $p$  and  $q$  such that  $pb = a = bq$  and  $(1 - p)a = 0 = a(1 - q)$ . Consequently,

$$b = (p + 1 - p)b(q + 1 - q) = a + (1 - p)a + a(1 - q) + (1 - p)b(1 - q) = a + (1 - p)b(1 - q).$$

Conversely, if  $a$  and  $c$  are regular elements in  $pRq$  and  $(1 - p)R(1 - q)$ , respectively, we can find partial inverses  $x$  and  $z$  for them. We may assume that  $x \in qRp$  and  $z \in (1 - q)R(1 - p)$ , replacing them if necessary with  $qxp$  and  $(1 - q)z(1 - p)$ . Then with  $b = a + c$  we have  $bx a = a = axb$ , so  $a \leq b$ . Moreover,  $y = x + z$  will be a partial inverse for  $b$  (even satisfying  $x \leq y$ ).  $\square$

**2.8. Proposition.** *For a unital ring  $R$ , each element in  $R_q^{-1}$  is maximal in  $R^r$  with respect to the ordering  $\leq$*

*Proof.* If  $u \in R_q^{-1}$  and  $u \leq a$  for some  $a$  in  $R^r$ , then  $u = uvu = avu = uva$  for some  $v$  in  $R$ . But then  $v$  is a quasi-inverse for  $u$  by Theorem 2.3. In particular,

$$a = a - (1 - uv)a(1 - vu) = uva + avu - uvavu = u + u - uvu = u.$$

$\square$

**2.9. Proposition.** *For a von Neumann regular element  $a$  in a unital ring  $R$  the following conditions are equivalent:*

- (i)  $a \leq u$  for some  $u$  in  $R_q^{-1}$ ;
- (ii)  $a = ava$  for some  $v$  in  $R_q^{-1}$ .

*Proof.* (i)  $\implies$  (ii) If  $a \leq u$  in  $R^r$ , choose a quasi-inverse  $v$  for  $u$ . By (iii) in 2.4 this implies that  $a = ava$ . Indeed,  $a$  has a partial inverse  $x$  such that  $a = axu = xua$ , whence

$$a = a(xa) = (axu)(xa) = (ax)(uvu)(xa) = (axu)v(uxa) = ava.$$

(ii)  $\implies$  (i) If  $a = ava$  with  $v$  in  $R_q^{-1}$ , consider the idempotents  $p = va$  and  $q = av$ . Let  $w$  be a partial inverse for  $v$  and consider the idempotents  $e = vw$  and  $f = wv$ . Then  $e' = (1 - p)e$  is an idempotent because  $ep = vwva = va = p$ . Evidently  $e' \leq 1 - p$ . Moreover,  $(p + e')v = pv + ev - pev = v$ , so  $vR \subset (p + e')R = pR + e'R$ . On the other hand, both  $pR \subset vR$  and  $e'R \subset vR$ , so we have equality. By symmetry we can find an idempotent  $f' \leq 1 - q$ , such that  $v(q + f') = v$  and  $Rv = Rq + Rf'$ . Write  $p + e' = vs$  and  $q + f' = tv$  for some elements  $s, t$  in  $R$ . Then put  $u = tvs$  and check that  $vu = vtv s = v(q + f')s = vs = p + e'$  and similarly  $uv = q + f'$ . In particular,  $vuv = v$ , so  $u \in R_q^{-1}$  by Theorem 2.3. Finally,  $avu = a(p + e') = ava + 0 = a$  and  $uva = (q + f')a = ava + 0 = a$ , so that  $a \leq u$ , as desired.  $\square$

**2.10. Remark.** Notice the affinities between the statements in Theorem 2.3 and Proposition 2.9: If  $uvu = u$  for some  $u$  in  $R_q^{-1}$ , then necessarily  $v \in R_q^{-1}$ . But if  $ava = a$  for some  $v$  in  $R_q^{-1}$ , then at least  $a$  extends to some  $u$  in  $R_q^{-1}$ .

### 3. QB-Rings

**3.1. Definition.** For each subset  $A$  of a unital ring  $R$  we define  $\text{cl}(A)$  to be the set of elements  $a$  in  $R$ , such that whenever  $xa + b = 1$  for some elements  $x$  and  $b$  in  $R$ , there is an element  $y$  in  $R$  such that  $a + yb \in A$ . Equivalently,  $a \in \text{cl}(A)$  if  $(a + Rb) \cap A \neq \emptyset$  whenever  $Ra + Rb = R$ .

**3.2. Lemma.** *The operation  $\text{cl}$  defined in 3.1 has the following properties relative to any subsets  $A$  and  $B$  of  $R$ :*

- (i)  $\text{cl}(\emptyset) = \emptyset$  and  $\text{cl}(R) = R$ ;
- (ii)  $A \subset B$  implies  $\text{cl}(A) \subset \text{cl}(B)$ ;
- (iii)  $A \subset \text{cl}(A) = \text{cl}(\text{cl}(A))$ ;
- (iv) If  $A \neq \emptyset$  then  $\mathcal{J}(R) \subset \text{cl}(A)$ , where  $\mathcal{J}(R)$  is the Jacobson radical of  $R$ ;
- (v)  $\mathcal{J}(R) = \text{cl}(0)$ ;
- (vi)  $\text{cl}(A) \cap R_\ell^{-1} = A \cap R_\ell^{-1}$ ;
- (vii)  $B \text{cl}(A) \subset \text{cl}(BA)$ ;
- (viii) If  $B \subset R^{-1}$  then  $\text{cl}(A)B \subset \text{cl}(AB)$ ;
- (ix)  $\text{cl}(A) + B \subset \text{cl}(A + RB)$ ;
- (x) If  $RB \subset B$  and  $A + B \subset \text{cl}(A)$ , then  $\text{cl}(A) + B \subset \text{cl}(A)$ ;
- (xi) If  $\pi: R \rightarrow S$  is any quotient morphism, then  $\pi(\text{cl}(A)) \subset \text{cl}(\pi(A))$ .

*Proof.* Properties (i) and (ii) are trivial to verify, as well as the first part of (iii). To complete the argument take any  $a$  in  $\text{cl}(\text{cl}(A))$ , and assume that  $xa + b = 1$ . Then by assumption  $a + yb \in \text{cl}(A)$  for some  $y$  in  $R$ . But since  $x(a + yb) + (1 - xy)b = 1$  this implies that  $a + yb + z(1 - xy)b \in A$  for some  $z$  in  $R$ . Consequently  $a + (y + z - zxy)b \in A$ , whence  $a \in \text{cl}(A)$ .

(iv) If  $A \neq \emptyset$  and  $z \in \mathcal{J}(R)$ , then any equation  $xz + b = 1$  implies that  $1 - b \in \mathcal{J}(R)$ , whence  $b \in R^{-1}$ . Evidently then,  $A \subset z + Rb$ , so  $z \in \text{cl}(A)$ .

(v) From (iv) we know that  $\mathcal{J}(R) \subset \text{cl}(0)$ . But if  $a \notin \mathcal{J}(R)$  then  $a \notin L$  for some maximal left ideal  $L$  of  $R$ . As  $Ra + L = R$  by maximality, we have  $xa + l = 1$  for some  $l$  in  $L$ . If  $a \in \text{cl}(0)$  this would imply that  $a + yl = 0$  for some  $y$  in  $R$ , whence  $a \in R$ , a contradiction. Therefore  $(R \setminus \mathcal{J}(R)) \cap \text{cl}(0) = \emptyset$ , so  $\text{cl}(0) = \mathcal{J}(R)$ .

(vi) If  $a \in \text{cl}(A) \cap R_\ell^{-1}$  we may consider the trivial decomposition  $Ra + R0 = R$ . By assumption  $a + y0 \in A$ , i.e.  $a \in A$ .

(vii) If  $e \in B$  and  $a \in \text{cl}(A)$ , consider any equation  $xea + b = 1$ . Then  $a + yb \in A$  for some  $y$ , whence  $ea + eyb \in eA \subset BA$ , and so  $ea \in \text{cl}(BA)$ .

(viii) Consider now an equation  $xae + b = 1$ . Since  $B \subset R^{-1}$  this rewrites as  $exa + ebe^{-1} = 1$ . Therefore  $a + yebe^{-1} \in A$  for some  $y$  in  $R$ , whence  $ae + yeb \in Ae \subset AB$ , showing that  $ae \in \text{cl}(AB)$ .



(ix) Suppose that  $e \in B$  and  $a \in \text{cl}(A)$ . For any equation  $x(e + a) + b = 1$  we can then find  $y$  such that  $a + y(xe + b) \in A$ . But then  $e + a + yb \in A + (1 - yx)e \subset A + RB$ , whence  $e + a \in \text{cl}(A + RB)$ .

(x) By (ix), (ii) and (iii) we have

$$\text{cl}(A) + B \subset \text{cl}(A + RB) \subset \text{cl}(A + B) \subset \text{cl}(\text{cl}(A)) = \text{cl}(A).$$

(xi) If  $a \in \text{cl}(A)$  and  $\pi(x)\pi(a) + \pi(b) = \pi(1)$  in  $S$ , then  $xa + b + t = 1$  in  $R$  for some  $t$  in  $\ker \pi$ . But then  $a + y(b + t) \in A$  for some  $y$  in  $R$ , whence  $\pi(a) + \pi(y)\pi(b) \in \pi(A)$ , proving that  $\pi(a) \in \text{cl}(\pi(A))$ .  $\square$

**3.3. Remark.** If  $R$  is commutative and  $Ra + Rb_i = R$  for  $i = 1, 2$ , then also  $Ra + Rb_1b_2 = R$ . Moreover,

$$a + Rb_1b_2 \subset (a + Rb_1) \cap (a + Rb_2).$$

This means that the sets of “neighbourhoods” of  $a$ , each of the form  $\mathcal{O}_a(b) = a + Rb$  for some  $b$  such that  $Ra + Rb = R$ , is directed by inclusion. Observe also that if  $c = a + yb \in \mathcal{O}_a(b)$ , then  $Rc + Rb = R$  and

$$\mathcal{O}_c(b) = c + Rb = a + Rb = \mathcal{O}_a(b).$$

Thus the sets  $\mathcal{O}_a(b)$  form the neighbourhood basis in a topology on  $R$  for which  $\text{cl}$  is the closure operation. In particular,  $\text{cl}(A \cup B) = \text{cl}(A) \cup \text{cl}(B)$ .

For non-commutative rings this fails already when  $R = \mathbb{M}_2(\mathbb{R})$ . Nevertheless the operation  $\text{cl}$  may with advantage be compared to a closure. A striking case occurs in  $C^*$ -algebra theory, cf. Proposition 9.1.

**3.4. Definition.** We shall be (almost) exclusively concerned with applying the operation  $\text{cl}$  to the set  $R_q^{-1}$  of quasi-invertible elements in a unital ring  $R$ . Since  $R^{-1}R_q^{-1} = R_q^{-1}$  and  $R_q^{-1}R^{-1} = R_q^{-1}$  we see from (vii) and (viii) in Lemma 3.2 that  $R^{-1}\text{cl}(R_q^{-1}) = \text{cl}(R_q^{-1})$  and  $\text{cl}(R_q^{-1})R^{-1} = \text{cl}(R_q^{-1})$ .

If  $\text{cl}(R_q^{-1}) = R$  we say that  $R$  is a  $QB$ -ring.

As mentioned in 3.1 the condition that  $a \in \text{cl}(R_q^{-1})$  is equivalent to the demand that whenever  $(a, b)$  is a left unimodular row, i.e.  $Ra + Rb = R$ , then  $a + yb \in R_q^{-1}$  for some  $y$  in  $R$ . Replacing quasi-invertibility by honest invertibility we are back at the definition of Bass stable rank 1, which in this setting says that  $\text{cl}(R^{-1}) = R$ . Thus,  $QB$ -rings are a generalization of  $B$ -rings, and actually a substantial weakening of this concept in the non-commutative case. Of course, if  $R$  is commutative,  $R_q^{-1} = R^{-1}$ .

Evidently the definition of  $\text{cl}(R_q^{-1})$  is not left-right symmetric, so we define  $\text{cr}(R_q^{-1})$  to be the set of elements  $a$  in  $R$ , such that whenever  $ax + b \in R^{-1}$  for some elements  $x, b$  in  $R$ , then  $a + by \in R_q^{-1}$  for some  $y$  in  $R$ . We have no reason to believe that  $\text{cl}(R_q^{-1}) = \text{cr}(R_q^{-1})$  in general. (But see Corollary 9.2.) However, just as for  $B$ -rings we do have complete symmetry in the definition of a  $QB$ -ring. This is proved by adapting [13, Lemma 1] to our present situation.

**3.5. Lemma.** *If  $a \in R$  such that  $ax + b = 1$  for some  $x$  in  $\text{cl}(R_q^{-1})$  and  $b$  in  $R$ , then  $a + by \in R_q^{-1}$  for some  $y$  in  $R$ .*

*Proof.* Since  $x \in \text{cl}(R_q^{-1})$  we have  $x + cb \in R_q^{-1}$  for some  $c$  in  $R$ . Thus

$$(1 - (x + cb)z) \perp (1 - z(x + cb))$$

for some  $z$  in  $R$ .

Define  $y = z(1 - ca)$  and  $d = x + (1 - xa)c$ . By straightforward, albeit lengthy computations, using that  $ax + b = 1$ , we then see that

$$\begin{aligned} 1 - (a + by)d &= 1 - (a + bz(1 - ca))(x + (1 - xa)c) \\ &= \dots = b(1 - z(x + cb))(1 - ac) . \end{aligned}$$

Similarly,

$$\begin{aligned} 1 - d(a + by) &= 1 - (x + (1 - xa)c)(a + bz(1 - ca)) \\ &= (1 - xa)(1 - (x + cb)z)(1 - ca) . \end{aligned}$$

It follows that

$$(1 - (a + by)d) \perp (1 - d(a + by)) ,$$

which shows that  $a + by \in R_q^{-1}$ , as desired.  $\square$

**3.6. Theorem.** *In any unital ring  $R$  we have*

$$\text{cl}(R_q^{-1}) = R \quad \Longleftrightarrow \quad \text{cr}(R_q^{-1}) = R .$$

*Proof.* By symmetry it suffices to show that  $\text{cr}(R_q^{-1}) = R$ , assuming that  $\text{cl}(R_q^{-1}) = R$ . But that is immediate from Lemma 3.5.  $\square$

**3.7. Proposition.** *If  $I$  is an ideal in a unital ring  $R$  and  $\pi: R \rightarrow R/I$  denotes the quotient morphism, then*

$$\pi(R_q^{-1}) \subset (R/I)_q^{-1} \quad \text{and} \quad \pi(\text{cl}(R_q^{-1})) \subset \text{cl}((R/I)_q^{-1}) .$$

*Proof.* Since central orthogonality is preserved under quotient morphisms it is evident that  $\pi(R_q^{-1}) \subset (R/I)_q^{-1}$ . The other inclusion follows from Conditions (xi) and (ii) in Lemma 3.2.  $\square$

**3.8. Corollary.** *If  $I$  is an ideal in a unital ring  $R$  such that  $I + \text{cl}(R_q^{-1}) = R$  then  $R/I$  is a  $QB$ -ring. In particular, the property of being a  $QB$ -ring passes to quotients.*  $\square$

The next result, a generalization of [11, Proposition 2.6], indicates the position of the  $B$ -rings as the “finite” members in the category of  $QB$ -rings.

**3.9. Proposition.** *Let  $R$  be a unital  $QB$ -ring. Then the following conditions are equivalent:*

- (i)  $R$  is a  $B$ -ring ,
- (ii)  $R_q^{-1} = R^{-1}$  .

*Proof.* (i)  $\implies$  (ii) Consider  $u$  in  $R_q^{-1}$  with quasi-inverse  $v$ , and let  $I$  and  $J$  denote the orthogonal ideals generated by  $1 - uv$  and  $1 - vu$ , respectively. Then  $u$  is right invertible in  $R/I$ , and since  $\text{bsr}(R/I) = 1$  because  $\text{bsr}(R) = 1$ , this means that  $u$  is actually invertible in  $R/I$  with (the image of)  $v$  as its inverse. It follows that  $1 - uv \in I$  and  $1 - vu \in I$ . Similarly  $1 - uv \in J$  and  $1 - vu \in J$ . Since both elements are idempotents and  $IJ = 0$ , the elements must be zero, and we conclude that  $uv = vu = 1$ , so that  $u \in R^{-1}$ .

(ii)  $\implies$  (i) If  $R$  is a  $QB$ -ring, then  $\text{cl}(R_q^{-1}) = R$ . If in addition  $R_q^{-1} = R^{-1}$  then by definition  $R$  is a  $B$ -ring.  $\square$

Recall that a simple, unital ring  $R$  is said to be *purely infinite* if  $R$  is not a division ring, but for any non-zero element  $x$  in  $R$  there are  $s, t$  in  $R$  such that  $sxt = 1$ . It is an open problem whether such a ring must be an exchange ring, cf. §8, but it is certainly well supplied with idempotents.

**3.10. Proposition.** *Every simple, purely infinite ring  $R$  is a  $QB$ -ring, but has infinite Bass stable rank.*

*Proof.* Take  $a$  in  $R$  and assume that  $xa + b = 1$  for some elements  $x, b$  in  $R$ . If  $b = 0$  then  $a$  is left invertible, so  $a \in R_q^{-1}$ . Otherwise, since  $R$  is purely infinite, we can write  $sbt = 1$  and compute

$$(a + (1 - at)sb)t = at + (1 - at) = 1.$$

With  $y = (1 - at)s$  this proves that  $a + yb$  is right invertible. Thus  $a \in \text{cl}(R_q^{-1})$ , and  $R$  is a  $QB$ -ring.

Since  $R$  is not a division ring, there is a non-zero element  $z$  in  $R$  which is not left invertible. Write  $czd = 1$  for some elements  $c, d$  in  $R$ . Let  $e = zdc$ , and note that  $e$  is a non-trivial idempotent in  $R$ , equivalent to 1. Set  $f_1 = 1 - e$ . Take any non-zero element  $z_1$  in  $f_1 R f_1$ , which is neither right nor left invertible. Since  $R$  is purely infinite, there are elements  $c_1$  and  $d_1$  in  $R$  such that  $c_1 z_1 d_1 = 1$ . We may assume that  $c_1 \in R f_1$ . Let  $f = z_1 d_1 c_1$ . Hence  $e$  and  $f$  are two orthogonal idempotents in  $R$ , both equivalent to 1. Now an argument similar to [28, Proposition 6.5] shows that  $R$  has infinite stable rank.  $\square$

**3.11. Example.** Let  $V$  be a vector space over a field  $\mathbb{F}$  and consider the ring  $R = \text{End}_{\mathbb{F}}(V)$  of all endomorphisms of  $V$ , which is known to be a unital, prime ring. Then  $R$  is a  $QB$ -ring (whereas the Bass stable rank is  $\infty$  if  $V$  is infinite dimensional).

To show this, consider  $a, b$  and  $x$  in  $R$  such that  $xa + b = 1$ . Evidently, then,  $\ker a \cap \ker b = 0$ , so we can choose a subspace  $U$  of  $V$  such that  $V = \ker a \oplus \ker b \oplus U$ . We can also choose subspaces  $W$  and  $W'$  of  $V$  such that  $V = b(V) \oplus W = a(V) \oplus W'$ .

Define  $c$  in  $R$  by  $c|W = 0$  and  $c|b(V) = (b|\ker a \oplus U)^{-1}$ . We can also define  $d$  in  $R$  by choosing an injective morphism  $z : \ker a \rightarrow W'$  if  $\dim(\ker a) \leq \dim W'$  or a surjective morphism  $z : \ker a \rightarrow W'$  if  $\dim(\ker a) \geq \dim W'$ , and then take  $d|\ker b \oplus U = 0$  and  $d|\ker a = z$ . Finally, let  $y = dc$ . Then  $a + yb$  gives a bijective morphism from  $\ker b \oplus U$  onto  $a(V)$ , whereas  $a + yb|\ker a = dcb|\ker a = z$ . It follows that  $a + yb$  is either injective or surjective on  $V$ , whence  $a + yb \in R_\ell^{-1} \cup R_r^{-1} = R_q^{-1}$ , since  $R$  is prime. Consequently  $a \in \text{cl}(R_q^{-1})$ , whence  $R = \text{cl}(R_q^{-1})$ , so  $R$  is a  $QB$ -ring.

In later sections, notably §7, §8 and §9, we shall present many more examples of  $QB$ -rings, as well as some counterexamples. For the time being we just notice that when trying to generalize the above example to the ring  $\text{End}(G)$  of an abelian group  $G$  we see that if  $G$  contains  $\mathbb{Z}$  as a direct summand, then  $\text{End}(G)$  has an idempotent  $p$  such that  $p\text{End}(G)p = \mathbb{Z}$ . Using Corollary 5.8 it follows that  $\text{End}(G)$  is not a  $QB$ -ring.

## 4. Rings Without Unit

**4.1. Adversibility.** Let  $R$  be a not necessarily unital ring. There is then a canonical unitization  $\tilde{R} = R \oplus \mathbb{Z}$ , with the obvious multiplication

$$(a, n)(b, m) = (ab + ma + nb, nm)$$

for  $a, b$  in  $R$  and  $n, m$  in  $\mathbb{Z}$ . Evidently  $\tilde{R}$  contains  $R$  as an ideal with  $\tilde{R}/R = \mathbb{Z}$ .

The fact that  $\mathbb{Z}$  is not a  $B$ -ring and thus, being commutative, neither a  $QB$ -ring, coupled with the fact that the  $QB$ -property passes to quotients (Corollary 3.8) shows that  $\tilde{R}$  will never be a  $QB$ -ring. We shall therefore seek a definition for the  $QB$ -property which is intrinsic for the non-unital case, even though it may seemingly borrow some structure from  $\tilde{R}$  or other unitizations.

Following ideas going back to Kaplansky we say that an element  $x$  in  $R$  is *left* (respectively *right*) *adversible* if  $a + x = ax$  (respectively  $x + a = xa$ ) for some  $a$  in  $R$ , which we call a *left* (respectively *right*) *adverse* to  $x$ . If  $x$  has both a left adverse  $a$  and a right adverse  $b$  it is called *adversible*, in which case

$$\begin{aligned} a &= a + (x + b - xb) - a(x + b - xb) \\ &= b + (x + a - ax) - (x + a - ax)b = b. \end{aligned}$$

The subsets of left, right and two-sided adversible elements in  $R$  are denoted  $R_\ell^\circ$ ,  $R_r^\circ$  and  $R^\circ$ , respectively.

We say that an element  $x$  in  $R$  is *quasi-adversible*, in symbols  $x \in R_q^\circ$ , if there exist elements  $b$  and  $c$  in  $R$  such that

$$(*) \quad (x + b - xb) \perp (x + c - cx).$$

Here  $s \perp t$  in  $R$  means that  $s\tilde{R}t = 0$  and  $t\tilde{R}s = 0$ . We say in this case that  $s$  and  $t$  are *centrally orthogonal*. If therefore  $I$  and  $J$  denote the ideals in  $R$  generated by  $x + b - xb$  and  $x + c - cx$ , respectively, then  $I \perp J$  in the sense that  $s \perp t$  for any pair of elements  $s$  in  $I$  and  $t$  in  $J$ . Moreover,  $x + I$  is right adversible in  $R/I$  and  $x + J$  is left adversible in  $R/J$ . Conversely, if we can find an orthogonal pair  $I, J$  of ideals in  $R$  (i.e.  $I\tilde{R}J = 0$  and  $J\tilde{R}I = 0$ ) such that  $x + I \in (R/I)_r^\circ$  and  $x + J \in (R/J)_\ell^\circ$ , then  $(*)$  is satisfied, so that  $x \in R_q^\circ$ .

**4.2. Proposition.** *Let  $S$  be any unital ring containing  $R$  as a subring, and let  $R_1$  denote the unital subring of  $S$  generated by  $S$  and  $1$ . Then we have the equalities*

$$\begin{aligned} 1 - R_\ell^\circ &= (R_1)_\ell^{-1} \cap (1 - R), & 1 - R_r^\circ &= (R_1)_r^{-1} \cap (1 - R), \\ 1 - R^\circ &= (R_1)^{-1} \cap (1 - R), & 1 - R_q^\circ &= (R_1)_q^{-1} \cap (1 - R). \end{aligned}$$

*Proof.* Straightforward computations based on the equation

$$1 - (1 - x)(1 - a) = x + a - xa.$$

□

**4.3. Corollary.** *If  $x \in R_q^\circ$  we can find a single element  $y$  in  $R$ , called the quasi-adverse for  $x$ , such that  $x + y - xy$  and  $x + y - yx$  are centrally orthogonal idempotents in  $R$ .*

*Proof.* Combine Propositions 2.2 and 4.2. □

**4.4. Definition.** For any ring  $R$  we define  $\text{cl}^\circ(R_q^\circ)$  to be the set of elements  $a$  in  $R$ , such that whenever  $xa - x - a + b = 0$  for some  $x, b$  in  $R$ , there is an element  $y$  in  $R$  such that  $a - yb \in R_q^\circ$ . Note that  $R_q^\circ \subset \text{cl}^\circ(R_q^\circ)$ . If  $\text{cl}^\circ(R_q^\circ) = R$  we say that  $R$  is a  $QB$ -ring. Similarly, one defines  $\text{cr}^\circ(R_q^\circ)$ . This prompts the question whether the notion of  $QB$ -ring in the non-unital case is also left-right symmetric. We shall address this problem in Remark 4.8.

**4.5. Remarks.** If  $R$  is unital then  $R_q^{-1} = 1 - R_q^\circ$  by Proposition 4.2. Moreover,  $(1 - x)(1 - a) + b = 1$  if and only if  $xa - x - a + b = 0$ . It follows that  $\text{cl}(R_q^{-1}) = 1 - \text{cl}^\circ(R_q^\circ)$ . Thus,  $\text{cl}(R_q^{-1}) = R$  if and only if  $\text{cl}^\circ(R_q^\circ) = R$ , so that our two definitions for  $QB$ -rings coincide.

In the non-unital case  $\text{cl}^\circ(R_q^\circ) = R$  means that  $1 - R \subset \text{cl}(\tilde{R}_q^{-1})$ , which is not at all the same as  $\tilde{R} = \text{cl}(\tilde{R}_q^{-1})$ .

If  $R$  is an algebra over a field  $\mathbb{F}$ , we redefine  $\tilde{R} = R \oplus \mathbb{F}$ , which is the minimal unitization of  $R$  as an algebra over  $\mathbb{F}$ . Now the discrepancies above vanish, and we see that  $R$  is a  $QB$ -algebra if and only if  $\tilde{R}$  is a  $QB$ -algebra.

To utilize the new definition in the non-unital case we shall need the following reformulation of [29, Lemma 3.5].

**4.6. Lemma.** *Let  $I$  be a right ideal in a unital ring  $R$ . Then for any pair of elements  $a$  in  $I$  and  $b$  in  $R$  the following conditions are equivalent:*

- (i)  $1 \in R(1 - a) + Rb$ ;
- (ii)  $I = I(1 - a) + Ib$ ;
- (iii)  $1 - I \subset (1 - I)(1 - a) + Ib$ ;
- (iv)  $xa - x - a + yb = 0$  for some  $x, y$  in  $I$ .

*Proof.* (i)  $\implies$  (ii) If  $1 = c(1 - a) + db$  for some  $c, d$  in  $R$  and if  $t \in I$ , then  $t = tc(1 - a) + tdb$ , and  $tc, td$  belong to  $I$ .

(ii)  $\implies$  (iii) Given  $t$  in  $I$  choose  $x, y$  in  $I$  such that  $a - t = -x(1 - a) + yb$ . Then

$$1 - t = 1 - a - x(1 - a) + yb = (1 - x)(1 - a) + yb.$$

(iii)  $\implies$  (iv) Choose  $x, y$  in  $I$  such that  $(1 - x)(1 - a) + yb = 1$ , whence  $xa - x - a + yb = 0$ .

(iv)  $\implies$  (i) If  $xa - x - a + yb = 0$ , then  $1 = (1 - x)(1 - a) + yb$ .  $\square$

**4.7. Proposition.** *If  $I$  is an ideal in a unital ring  $R$ , and  $t \in I$ , then  $t \in I_q^\circ$  if and only if  $1 - t \in R_q^{-1}$ . Moreover,  $t \in \text{cl}^\circ(I_q^\circ)$  if and only if  $1 - t \in \text{cl}(R_q^{-1})$ .*

*Proof.* If  $t \in I_q^\circ$ , then by Corollary 4.3 we can find centrally orthogonal idempotents in  $I$  of the form  $p = s + t - ts$  and  $q = s + t - st$  for some  $s$  in  $I$ . Since  $I$  is an ideal in  $R$  this implies that

$$pRq = p^2Rq^2 \subset pIq = 0.$$

Similarly  $qRp = 0$ . It follows that  $1 - t \in R_q^{-1}$  with partial inverse  $1 - s$ .

Conversely, if  $1 - t \in R_q^{-1}$  we can find a quasi-inverse in  $R$ , written in the form  $1 - s$ , such that

$$(1 - (1 - t)(1 - s)) \perp (1 - (1 - s)(1 - t)).$$

The equation  $1 - t = (1 - t)(1 - s)(1 - t)$  shows that  $s = -t + t^2 + ts + st - tst \in I$ , whence  $t \in I_q^\circ$  with quasi-adverse  $s$ .

If now  $t \in \text{cl}^\circ(I_q^\circ)$  and  $a(1 - t) + b = 1$  for some  $a, b$  in  $R$ , then  $xt - x - t + yb = 0$  for some  $x, y$  in  $I$  by Lemma 4.6, whence  $t - ryb \in I_q^\circ$  for some  $r$  in  $I$ . From the first part of the proof it follows that  $1 - t + ryb \in R_q^{-1}$ , which shows that  $1 - t \in \text{cl}(R_q^{-1})$ .

Conversely, if  $1 - t \in \text{cl}(R_q^{-1})$ , and if  $st - t - s + r = 0$  for some  $s, r$  in  $I$ , then evidently  $(1 - s)(1 - t) + r = 1$  in  $R$ . By Lemma 4.6 this implies that  $1 - 0 = (1 - x)(1 - t) + yr$  for some  $x, y$  in  $I$ . Since  $1 - t \in \text{cl}(R_q^{-1})$  it follows that  $1 - t + cyr \in R_q^{-1}$  for some  $c$  in  $R$ , whence  $t - cyr \in I_q^\circ$  by the first part of the proof. As  $cy \in I$  this proves that  $t \in \text{cl}^\circ(I_q^\circ)$ , as desired.  $\square$

**4.8. Remark.** We are now in a position to show that the notion of  $QB$ -ring is also symmetric in the non-unital case. Indeed, let  $I$  be a non-unital ring and let  $R = \tilde{I}$  (as in 4.1). We identify  $I$  with a two-sided ideal of  $R$ . It is enough to see that  $\text{cr}^\circ(I_q^\circ) = I$  provided that  $\text{cl}^\circ(I_q^\circ) = I$ . Let  $t$  in  $I$ , and assume that  $tx - t - x + b = 0$  for some  $x, b$  in  $I$ . In  $R$  this reads as  $(1 - t)(1 - x) + b = 1$ .

By the right-handed version of Lemma 4.6 there are elements  $x_1$  and  $y$  in  $I$  such that  $(1-t)(1-x_1) + by = 1$ . Since  $x_1 \in \text{cl}^\circ(I_q^\circ)$  it follows by Proposition 4.7 that  $1-x_1 \in \text{cl}(R_q^{-1})$ . Now, by Lemma 3.5, there exists  $z$  in  $R$  such that  $1-t+byz \in R_q^{-1}$ . Again Proposition 4.7 implies that  $t - b(yz) \in I_q^\circ$ . Since  $yz \in I$ , we see that  $t \in \text{cr}^\circ(I_q^\circ)$ .

**4.9. Theorem.** *An ideal  $I$  in a unital ring  $R$  is a  $QB$ -ring if and only if*

$$1 - I \subset \text{cl}(R_q^{-1}).$$

*Proof.* By Proposition 4.7 the condition  $1 - I \subset \text{cl}(R_q^{-1})$  is equivalent to  $I \subset \text{cl}^\circ(I_q^\circ)$ , which by definition means that  $I$  is a  $QB$ -ring  $\square$

**4.10. Corollary.** *Every ideal in a  $QB$ -ring (unital or not) will be a  $QB$ -ring.*

*Proof.* Only the non-unital case deserves an argument. But if  $I$  is an ideal in the non-unital  $QB$ -ring  $R$ , then  $I$  is also an ideal of  $\tilde{R}$ , so by Proposition 4.7

$$\text{cl}^\circ(I_q^\circ) = (1 - \text{cl}(\tilde{R}_q^{-1})) \cap I = \text{cl}^\circ(R_q^\circ) \cap I = R \cap I = I.$$

$\square$

**4.11. Example.** The ring  $\mathbb{Z}[[x]]$  of formal power series in one variable over  $\mathbb{Z}$  is not a  $B$ -ring, but the ideal  $x\mathbb{Z}[[x]]$  is. To see this, recall first that  $\mathbb{Z}[[x]]$  is well supplied with units, in fact  $a = \sum \alpha_n x^n \in (\mathbb{Z}[[x]])^{-1}$  if and only if  $\alpha_0 = \pm 1$ . It follows that for any equation

$$(1-y)(1-a) + b = 1,$$

with  $a, b$  and  $y$  in  $x\mathbb{Z}[[x]]$ , we can find an element  $z$  in  $x\mathbb{Z}[[x]]$ , such that  $1-a+zb \in (\mathbb{Z}[[x]])^{-1}$ . Indeed,  $z = 0$  will do.

We claim that there is no unital  $B$ -ring or even  $QB$ -ring  $R$  that contains  $x\mathbb{Z}[[x]]$  as an ideal, and we mention this fact to show that our specific  $QB$ -definitions are needed for the non-unital case.

To establish the claim, assume that  $R$  was such a  $QB$ -ring. The trivial equation  $3 \cdot 2 - 5 = 1$  in  $R$  shows that  $2 - 5y \in R_q^{-1}$  for some  $y$  in  $R$ , so that

$$(1 - (2 - 5y)v)R(1 - v(2 - 5y)) = 0$$

for some  $v$  in  $R$ . Multiplying left and right with  $x$ , and replacing  $R$  with  $x^2$  we obtain the equation

$$(x^2 - (2x - 5xy)vx)(x^2 - xv(2x - 5yx)) = 0$$

in  $x\mathbb{Z}[[x]]$ . However, this is a prime ring, so one of the factors must be zero, say the one to the left. Write  $xy = \sum \alpha_n x^n$  and  $vx = \sum \beta_n x^n$ , and note that the equation

$$x^2 = (2x - 5xy)vx = (2x - 5 \sum_{n=1}^{\infty} \alpha_n x^n) (\sum_{n=1}^{\infty} \beta_n x^n)$$

forces  $1 = (2 - 5\alpha_1)\beta_1$  in  $\mathbb{Z}$ , which is plainly impossible.  $\square$

## 5. Skew Corners in QB-Rings

**5.1. Definition.** Let  $p$  and  $q$  be idempotents in a ring  $R$  such that  $pRq \neq 0$ . We say that an element  $x$  in  $pRq$  is *quasi-invertible*, and write  $x \in (pRq)_q^{-1}$ , if there exist elements  $a, b$  in  $qRp$  such that

$$(p - xa) \perp (q - bx) .$$

As in 2.1 we can take  $y = a + b - axb$  in  $qRp$ , to obtain the equations  $x = xyx$  and

$$(p - xy) \perp (q - yx) .$$

In particular,  $qRp \neq 0$  and  $y \in (qRp)_q^{-1}$ . Replacing if necessary  $y$  by  $xyx$  we may also assume that  $y = yxy$ , so that  $x$  and  $y$  are von Neumann regular elements and partial inverses for each other.

**5.2. Definitions.** With  $p, q$  and  $R$  as in 5.1 we define the subset  $\text{cl}^\sim((pRq)_q^{-1})$  to be the set of elements  $a$  in  $pRq$  such that whenever  $xa + b = q$  for some elements  $x$  in  $qRp$  and  $b$  in  $qRq$  we have  $a + yb \in (pRq)_q^{-1}$  for some  $y$  in  $pRq$ . Symmetrically we define the subset  $\text{cr}^\sim((pRq)_q^{-1})$  as the elements  $a$  in  $pRq$  such that whenever  $ax + b = p$  for some  $x$  in  $qRp$  and  $b$  in  $pRp$  then  $a + by \in (pRq)_q^{-1}$  for some  $y$  in  $pRq$ .

We leave it to the reader to verify, using the computations in Lemma 3.5, that if  $ax + b = q$  for some  $a$  in  $qRp$ ,  $b$  in  $qRq$  and  $x$  in  $\text{cl}^\sim((pRq)_q^{-1})$ , then  $a + by \in (qRp)_q^{-1}$  for some  $y$  in  $qRp$ . In particular,  $\text{cl}^\sim((pRq)_q^{-1}) = pRq$  if and only if  $\text{cr}^\sim((qRp)_q^{-1}) = qRp$ .

We shall be exclusively concerned with the case where both  $\text{cl}^\sim((pRq)_q^{-1}) = pRq$  and  $\text{cr}^\sim((pRq)_q^{-1}) = pRq$ , and will refer to this situation as  $pRq$  being a *QB-corner*. It follows that in this case we also have that  $qRp$  is a *QB-corner*.

For an idempotent  $p$  in  $R$  we now seemingly have two notions of *QB-structure*, one regarding  $pRp$  as a unital ring in its own right, and one regarding it as a corner in  $R$ . Fortunately these coincide, as will be seen from Theorem 5.5, cf. Corollary 5.7.

As already mentioned in Theorem 2.3 we say that two idempotents are (*Murray-von Neumann*) *equivalent*, in symbols  $p \sim q$ , if  $p = uv$  and  $q = vu$  for some elements  $u, v$  in  $R$ . Replacing if necessary  $u$  and  $v$  with  $uvu$  and  $vuv$ , we may assume that  $u$  and  $v$  are von Neumann regular elements and partial inverses for each other, and that

$$(1 - p)u = u(1 - q) = 0 = (1 - q)v = v(1 - p) .$$

We shall tacitly make these assumptions when we write  $p \sim q$ .

**5.3. Lemma** (Cf. [8, 2.1]). *Let  $p$  and  $q$  be idempotents in a unital ring  $R$  such that  $1 - p \sim 1 - q$ , i.e.  $1 - p = uv$  and  $1 - q = vu$ . If  $x \in Rq$  such that  $u + x$*



is left or right invertible in  $R$ , then the same holds for any element of the form  $u + x + (1 - p)yq$ . In particular for  $u + px$ .

*Proof.* Using the matrix decompositions  $R = (p + (1 - p))R \begin{pmatrix} q \\ 1 - q \end{pmatrix}$  for  $u + x$  and  $R = (q + (1 - q))R \begin{pmatrix} p \\ 1 - p \end{pmatrix}$  for its inverses we can write  $u + x = \begin{pmatrix} x_1 & 0 \\ x_2 & u \end{pmatrix}$ . If this element has a left inverse  $a$  we have a matrix equation

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} x_1 & 0 \\ x_2 & u \end{pmatrix} = \begin{pmatrix} q & 0 \\ 0 & 1 - q \end{pmatrix}.$$

Thus,  $a_{22}u = 1 - q$ , whence

$$a_{22} = a_{22}(1 - p) = a_{22}uv = (1 - q)v = v.$$

Moreover,  $a_{12}u = 0$ , whence

$$a_{12} = a_{12}(1 - p) = a_{12}uv = 0.$$

This means that

$$q = a_{11}x_1 + a_{12}x_2 = a_{11}x_1.$$

Thus, for any  $y$  in  $(1 - p)Rq$  we have

$$\begin{pmatrix} a_{11} & 0 \\ -vya_{11} & v \end{pmatrix} \begin{pmatrix} x_1 & 0 \\ y & u \end{pmatrix} = \begin{pmatrix} q & 0 \\ 0 & 1 - q \end{pmatrix},$$

so  $u + px + y$  is left invertible.

Assume now that  $u + x$  has a right inverse, i.e.

$$\begin{pmatrix} x_1 & 0 \\ x_2 & u \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} p & 0 \\ 0 & 1 - p \end{pmatrix}.$$

This implies that  $x_1a_{11} = p$ , so for any  $y$  in  $(1 - p)Rq$  we have

$$\begin{pmatrix} x_1 & 0 \\ y & u \end{pmatrix} \begin{pmatrix} a_{11} & 0 \\ -vya_{11} & v \end{pmatrix} = \begin{pmatrix} p & 0 \\ 0 & 1 - p \end{pmatrix},$$

so that  $u + px + y$  is right invertible.  $\square$

**5.4. Corollary.** *If  $p$  and  $q$  are idempotents in a unital ring  $R$  with  $1 - p \sim 1 - q$ , i.e.  $1 - p = uv$  and  $1 - q = vu$ , and if  $u + x \in R_q^{-1}$  for some  $x$  in  $Rq$ , then for any  $y$  in  $(1 - p)Rq$  we also have  $u + x + y \in R_q^{-1}$ . In particular,  $u + px \in R_q^{-1}$ .  $\square$*

**5.5. Theorem.** *Let  $p$  and  $q$  be idempotents in a unital ring  $R$  such that  $1 - p = uv$  and  $1 - q = vu$ , and consider a non-zero element  $x$  in  $pRq$ . Then  $x \in (pRq)_q^{-1}$  if and only if  $u + x \in R_q^{-1}$ . Moreover,  $x \in \text{cl}^\sim((pRq)_q^{-1})$  if and only if  $u + x \in \text{cl}(R_q^{-1})$ .*

*Proof.* If  $x \in (pRq)_q^{-1}$  we can find an element  $y$  in  $qRp$  such that  $p - xy \perp q - yx$ . Set  $u' = u + x$  and  $v' = v + y$ . Then by computation  $1 - u'v' = 1 - uv - xy = p - xy$ , and similarly  $1 - v'u' = q - yx$ , so that  $1 - u'v' \perp 1 - v'u'$  and  $u + x \in R_q^{-1}$ .

Conversely, if  $u' = u + x \in R_q^{-1}$  and  $v'$  is a quasi-inverse for  $u'$  in  $R$ , so that  $1 - u'v' \perp 1 - v'u'$ , then evidently  $p - pu'v'p \perp q - qv'u'q$ . Since  $pu' = x = u'q$  we can use  $y = qv'p$  in  $qRp$  to write  $p - xy \perp q - yx$ , i.e.  $x \in (pRq)_q^{-1}$ .

Assume now that  $x \in \text{cl}^\sim((pRq)_q^{-1})$ , and with  $a = u + x$  consider an equation  $za + b = 1$  in  $R$ . Multiplying left and right with  $q$  this leads to the equation  $qzpx + qbq = q$ , since  $aq = x = px$ . By assumption we therefore have  $x + ybq \in (pRq)_q^{-1}$  for some  $y$  in  $pRq$ . From what we proved above this implies that  $u + x + ybq \in R_q^{-1}$ . However,  $x + ybq \in pR$ , so by (the symmetric version of) Corollary 5.4 it follows that also  $u + x + yb \in R_q^{-1}$ . Consequently  $u + x \in \text{cl}(R_q^{-1})$ .

Finally assume that  $u + x \in \text{cl}(R_q^{-1})$ , and consider an equation  $zx + b = q$  for some elements  $z$  in  $qRp$  and  $b$  in  $qRq$ . Then

$$(v + z)(u + x) + b = 1 - q + zx + b = 1.$$

Since  $u + x \in \text{cl}(R_q^{-1})$  this implies that

$$u + x + yb \in R_q^{-1}, \quad y \in R.$$

As  $x + yb \in Rq$  we conclude from Corollary 5.4 that also  $u + x + pyb \in R_q^{-1}$ , i.e. we may assume that  $y \in pRq$ . By definition there is an element  $c$  in  $R$  with

$$(1 - (u + x + yb)c) \perp (1 - c(u + x + yb)).$$

Multiplying the two elements from left and right with  $p$  and with  $q$ , respectively, we see that

$$(p - (x + yb)qcp) \perp (q - qcp(x + yb)),$$

which shows that  $x + yb \in (pRq)_q^{-1}$ , whence  $x \in \text{cl}^\sim((pRq)_q^{-1})$ .  $\square$

**5.6. Corollary.** *If  $p$  and  $q$  are idempotents in a unital ring  $R$  such that  $1 - p = uv$  and  $1 - q = vu$ , and if  $pRq \neq 0$ , then  $pRq$  is a  $QB$ -corner if and only if*

$$u + pRq \subset \text{cl}(R_q^{-1}) \cap \text{cr}(R_q^{-1}). \quad \square$$

**5.7. Corollary.** *For any non-zero idempotent  $p$  in  $R$  the ring  $pRp$  is a  $QB$ -corner, hence also a  $QB$ -ring, if and only if  $1 - p + pRp \subset \text{cl}(R_q^{-1})$ .  $\square$*

**5.8. Corollary.** *If  $R$  is a  $QB$ -ring, then  $pRp$  is a  $QB$ -corner for any non-zero idempotent  $p$  in  $R$ , and for any pair of idempotents  $p$  and  $q$  such that  $1 - p \sim 1 - q$  we either have  $p \perp q$  or else  $pRq$  is a  $QB$ -corner.*  $\square$

Using the preceding results we can take up again the discussion from §2 about the rôle of  $R_q^{-1}$  as the maximally extended von Neumann regular elements. In  $QB$ -rings the solution is optimal:

**5.9. Theorem** (Cf. [8, 2.6]). *Let  $a$  be a von Neumann regular element in a unital ring  $R$ . If  $a \in \text{cl}(R_q^{-1})$ , then  $a \leq u$  for some  $u$  in  $R_q^{-1}$ .*

*Proof.* By assumption  $a = axa$  and  $x = xax$  for some  $x$  in  $R$ . Define the idempotents  $p = 1 - ax$  and  $q = 1 - xa$ , where  $pa = aq = 0$ . Thus,  $xa + q = 1$ , and since  $a \in \text{cl}(R_q^{-1})$  this means that  $a + yq \in R_q^{-1}$  for some  $y$  in  $R$ . However,  $1 - p \sim 1 - q$ , so we can apply Corollary 5.4 to conclude that also  $a + pyq \in R_q^{-1}$ .

If we now define  $u = a + pyq$  then it follows from Lemma 2.7 that  $a \leq u$ , as desired.  $\square$

**5.10. Corollary** (Cf. [8, 4.3]). *In a unital  $QB$ -ring every von Neumann regular element extends to a quasi-invertible element.*  $\square$

**5.11. Corollary.** *In a unital  $QB$ -ring  $R$  an element in  $R^r$  is maximal with respect to the order  $\leq$  if and only if it belongs to  $R_q^{-1}$ .*

*Proof.* In view of Proposition 2.8 we need only consider an element  $a$  in  $R^r$  which is maximally extended, and prove that  $a \in R_q^{-1}$ . But that is evident from Corollary 5.10.  $\square$

**5.12. Morita Contexts.** The concrete concept of a skew corner  $pRq$  in a ring  $R$  and its transposed corner  $pRq$  can be developed abstractly as a theory of bimodules in Morita contexts.

If  $R$  and  $S$  are unital rings and if  $M$  is an  $R - S$ -bimodule and  $N$  is an  $S - R$ -bimodule, we say that the pair  $M, N$  is in a *Morita context* if we can find surjective bimodule maps

$$\varphi: M \otimes_S N \rightarrow R \quad \text{and} \quad \psi: N \otimes_R M \rightarrow S.$$

Moreover these maps should satisfy the compatibility relations

$$\varphi(x \otimes y)x' = x\psi(y \otimes x') \quad \text{and} \quad y'\varphi(x \otimes y) = \psi(y' \otimes x)y$$

for all  $x, x'$  in  $M$  and  $y, y'$  in  $N$ .

Given a pair  $M, N$  of bimodules in a Morita context for the rings  $R$  and  $S$  we can define the *Morita ring* (or *linking ring*)  $L = \begin{pmatrix} R & M \\ N & S \end{pmatrix}$  with pointwise addition and “matrix product” given by

$$\begin{pmatrix} a_1 & x_1 \\ y_1 & b_1 \end{pmatrix} \begin{pmatrix} a_2 & x_2 \\ y_2 & b_2 \end{pmatrix} = \begin{pmatrix} a_1a_2 + \varphi(x_1 \otimes y_2) & a_1x_2 + x_1b_2 \\ y_1a_2 + b_1y_2 & \psi(y_1 \otimes x_2) + b_1b_2 \end{pmatrix}.$$

In  $L$  we find the orthogonal pair of idempotents

$$p = \begin{pmatrix} 1_R & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad q = \begin{pmatrix} 0 & 0 \\ 0 & 1_S \end{pmatrix}$$

and the identifications

$$R = pLp, \quad S = qLq, \quad M = pLq, \quad N = qLp.$$

The conditions that  $\varphi$  and  $\psi$  are surjective prove that  $R$  and  $S$  are Morita equivalent. Without this restriction we see that  $\varphi(M \otimes_S N)$  and  $\psi(N \otimes_R M)$  will generate ideals  $R_0$  and  $S_0$  in  $R$  and  $S$ , respectively, such that  $M$  and  $N$  will be in a Morita context for  $R_0$  and  $S_0$ . We can still form the large Morita ring  $L$ , but we can also form the smaller Morita ring  $L_0$ , using  $R_0$  and  $S_0$ , which will then be an ideal in  $L$  with quotient  $L/L_0 = R/R_0 \oplus S/S_0$ .

## 6. Matrices over QB-Rings

We shall prove, with considerable effort, that  $M_n(R)$  is a QB-ring whenever  $R$  is. Our argument is an amalgamation of an argument attributed to Kaplansky for the fact that Bass stable rank one passes to matrices, and the series of reductions found in the proof of [8, Theorem 4.5].

First we show that [8, Proposition 4.4] generalizes to the purely algebraic setting.

**6.1. Lemma.** *Consider elements  $u, v$  in  $R_q^{-1}$  with quasi-inverses  $x, y$  so that we have  $(1 - ux) \perp (1 - xu)$  and  $(1 - vy) \perp (1 - yv)$ . If moreover*

$$(*) \quad uv + (1 - ux)R(1 - yv) \subset \text{cl}(R_q^{-1}) \cap \text{cr}(R_q^{-1}),$$

*in particular if  $R$  is a QB-ring, then either  $(1 - ux)R(1 - yv) = 0$  or it is a QB-corner.*

*Proof.* Let  $I$  denote the ideal generated by the two idempotents  $1 - xu$  and  $1 - vy$ . Since  $((1 - ux)R(1 - yv)) \cap I = 0$  it follows that  $(1 - ux)R(1 - yv)$  is isomorphic to its image in  $R/I$ . Moreover, by Proposition 3.7 condition  $(*)$  is still valid in  $R/I$ . To establish the lemma we may therefore assume that  $I = 0$ , i.e.  $1 = xu = vy$ . Consequently

$$(uv)(yx) = ux \quad \text{and} \quad (yx)(uv) = yv,$$

so  $ux \sim yv$ . We can therefore apply Corollary 5.6 to conclude that  $(1 - ux)R(1 - yv)$  is a QB-corner if it is non-zero.  $\square$

**6.2. Remark.** It is perhaps instructive to realize that Lemma 6.1 also has a non-unital version: Consider elements  $a, b$  in  $R_q^\circ$  with quasi-adverses  $x, y$  so that

$$(ax - a - x) \perp (xa - a - x) \quad \text{and} \quad (by - b - y) \perp (yb - b - y).$$

If we can show that

$$ab - a - b + (ax - a - x)R(yb - b - y) \subset \text{cl}^\circ(R_q^\circ) \cap \text{cr}^\circ(R_q^\circ),$$

in particular if  $R$  is a  $QB$ -ring, then  $(ax - a - x)R(yb - b - y)$  is a  $QB$ -corner if it is non-zero.

The proof is a tedious check of the fact that when we replace  $a, b, x$  and  $y$  by  $1 - a, 1 - b, 1 - x$  and  $1 - y$  in  $\tilde{R}$ , we can still apply Corollary 5.6.

To facilitate the arguments in the following we say that a right unimodular row  $(a, b)$  in  $R$  (i.e.  $aR + bR = R$ ) is *right reducible* if  $a + by \in R_q^{-1}$  for some  $y$  in  $R$ . Note that we deliberately choose a non-symmetric version, favouring the first coordinate in the row. In the applications we shall consider a pair  $(a, b)$  such that  $ax + b \in R^{-1}$ , and right reducibility therefore means that  $a \in \text{cr}(R_q^{-1})$ .

The next lemma is a special case of the fact that multiplication by invertible matrices preserve unimodular rows.

**6.3. Lemma.** *Let  $(a, b)$  be a right unimodular row in a unital ring  $R$ . Then for any pair of units  $u, v$  in  $R^{-1}$  and  $c$  in  $R$  the pair  $(vau + vbc, vb)$  is also right unimodular, and it is right reducible if and only if  $(a, b)$  is right reducible.*

*Proof.* If  $ax + by = 1$  then  $(vau + vbc)u^{-1}xv^{-1} + vb(y - cu^{-1}x)v^{-1} = 1$ , so the new row is still right unimodular.

Assume now that  $(a, b)$  is right reducible, more specifically,  $a + by \in R_q^{-1}$ . Then  $vau + vbc + vb(yu - c) = v(a + by)u \in R_q^{-1}$ , as desired. The converse implications are similar.  $\square$

**6.4. Theorem.** *If  $R$  is a unital  $QB$ -ring, then  $\mathbb{M}_n(R)$  is also a  $QB$ -ring for any natural number  $n$ .*

*Proof.* It suffices to prove the theorem for  $n = 2$ , since by iteration this would give the result for all numbers  $n = 2^k$ ; and since  $\mathbb{M}_n(R)$  is a corner in  $\mathbb{M}_{2^k}(R)$  for  $n \leq 2^k$ , the general case follows from Corollary 5.8.

Consider therefore a right unimodular row  $\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} e & f \\ g & h \end{pmatrix}\right)$  and assume that

$$(*) \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} + \begin{pmatrix} e & f \\ g & h \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Since  $aa' + bc' + e = 1$  and  $R$  is a  $QB$ -ring, we have  $a + (bc' + e)z_1 \in R_q^{-1}$  for some  $z_1$  in  $R$ . By Lemma 6.3 our original row is right reducible if and only if this is so for the row with elements

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ c'z_1 & 1 \end{pmatrix} + \begin{pmatrix} e & f \\ g & h \end{pmatrix} \begin{pmatrix} z_1 & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} e & f \\ g & h \end{pmatrix};$$

and now the first matrix has the  $(1, 1)$ -element  $a + bc'z_1 + ez_1$ , which is quasi-invertible, whereas its second column is unchanged. Without loss of generality we may therefore consider the equation  $(*)$  under the assumption that  $a \in R_q^{-1}$ .

Choose a quasi-inverse  $x$  for  $a$ , and apply Lemma 6.3 to obtain the new right unimodular row of matrices with elements

$$\begin{pmatrix} 1 & 0 \\ -cx & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & -xb \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 0 \\ -cx & 1 \end{pmatrix} \begin{pmatrix} e & f \\ g & h \end{pmatrix}.$$

Now the first matrix has the form

$$\begin{pmatrix} a & (1-ax)b \\ c(1-xa) & d_0 \end{pmatrix},$$

so we may assume, without loss of generality, that we have equation (\*) under the further restriction that  $axb = 0$  and  $cxa = 0$ .

Computing the  $(2,2)$ -element in (\*) we obtain the equation  $cb' + dd' + h = 1$ , and since  $R$  is a  $QB$ -ring this implies that  $d + (cb' + h)z_2 \in R_q^{-1}$  for some  $z_2$  in  $R$ . Using Lemma 6.3 we pass to the right unimodular row with elements

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & b'z_2 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} e & f \\ g & h \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & z_2 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} e & f \\ g & h \end{pmatrix};$$

where now the first matrix has the form

$$\begin{pmatrix} a & b + (ab' + f)z_2 \\ c & d + (cb' + h)z_2 \end{pmatrix},$$

so its  $(2,2)$ -element is quasi-invertible. For the  $(1,2)$ -position we compute

$$ax(b + (ab' + f)z_2) = 0 + ax(ab' + bd' + f)z_2 = 0$$

using the  $(1,2)$ -equation from (\*) and the fact that  $axb = 0$ . Without loss of generality we may therefore assume that we have equation (\*) under the further restriction that  $d \in R_q^{-1}$ .

Choose now a quasi-inverse  $y$  for  $d$  and apply Lemma 6.3 to obtain a new right unimodular row, where the first matrix has the form

$$\begin{pmatrix} 1 & -by \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -yc & 1 \end{pmatrix} = \begin{pmatrix} a & b(1-yd) \\ (1-dy)c & d \end{pmatrix}.$$

Without loss of generality we may therefore assume that we have equation (\*) with the elements  $a$  and  $d$  in  $R_q^{-1}$  with quasi-inverses  $x$  and  $y$ , such that

$$axb = 0 = byd \quad \text{and} \quad cxa = 0 = dyc.$$

The  $(1,1)$ -equation in (\*) followed by multiplication left and right by  $1 - ax$ , yields

$$(1 - ax)b(1 - yd)c'(1 - ax) + (1 - ax)e(1 - ax) = 1 - ax.$$

Either  $(1 - ax)R(1 - yd) = 0$ , in which case also the element  $b$  is zero, and we do nothing. Otherwise  $(1 - ax)R(1 - yd)$  is a  $QB$ -corner by Lemma 6.1, whence

$$b + (1 - ax)e(1 - ax)z_3 \in ((1 - ax)R(1 - yd))_q^{-1}$$

for some  $z_3$  in  $(1 - ax)R(1 - yd)$ . Applying Lemma 6.3 we pass to the new right unimodular row with elements

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & -xez_3 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} e & f \\ g & h \end{pmatrix} \begin{pmatrix} 0 & z_3 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} e & f \\ g & h \end{pmatrix};$$

in which the first matrix has the form

$$\begin{pmatrix} a & b + (1 - ax)ez_3 \\ c & d + g(1 - ax)z_3 \end{pmatrix} = \begin{pmatrix} a & b_1 \\ c & d_1 \end{pmatrix},$$

where  $b_1 \in ((1 - ax)R(1 - yd))_q^{-1}$ . Since  $z_3 \in R(1 - yd)$  it follows from Theorem 2.3 that  $d_1 \in R_q^{-1}$  with quasi-inverse  $y$  such that

$$(1 - d_1y) \perp (1 - yd_1), \quad (1 - d_1y) \perp (1 - yd), \quad (1 - dy) \perp (1 - yd_1).$$

Moreover,

$$d_1y = (d + g(1 - ax)z_3)y = dy$$

since  $z_3y = 0$ , so  $d_1 = d_1yd_1 = dy(d + g(1 - ax)z_3) \in d + dyR$ . Replacing  $b$  and  $d$  with  $b_1$  and  $d_1$  in  $(*)$  we therefore still have the equations

$$d_1yc = 0 \quad \text{and} \quad axb_1 = 0.$$

Furthermore,

$$b_1yd_1 = (b + (1 - ax)ez_3)yd_1 = byd_1 = bydyd_1 = 0.$$

By Lemma 6.3 we may therefore assume, on top of the previous conditions, that we have  $(*)$  either with  $b = 0$  or with  $b$  in  $((1 - ax)R(1 - yd))_q^{-1}$ .

A symmetric argument, using the  $(2, 2)$ -equation in  $(*)$  will transform our right unimodular row to one with elements of the form

$$\begin{pmatrix} a_1 & b \\ c_1 & d \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} e & f \\ g & h \end{pmatrix},$$

where now either  $c_1 = 0$  and  $a_1 = a$  (in case  $(1 - dy)R(1 - xa) = 0$ ); or else  $a_1 \in R_q^{-1}$  with  $xa_1 = xa$ , such that  $a_1$  has  $x$  as its quasi-inverse and

$$c_1 \in ((1 - dy)R(1 - xa))_q^{-1} = ((1 - dy)R(1 - xa_1))_q^{-1}.$$

Applying Lemma 6.3 for the last time we may therefore assume that we have equation  $(*)$ , but such that  $a$  and  $b$  are both quasi-invertible with quasi-inverses  $x$  and  $y$  so that

$$(1 - ax) \perp (1 - xa) \quad \text{and} \quad (1 - by) \perp (1 - yb),$$

and such that also

$$b \in ((1 - ax)R(1 - yd))_q^{-1} \cup \{0\} \quad \text{and} \quad c \in ((1 - dy)R(1 - xa))_q^{-1} \cup \{0\}$$

with quasi-inverses  $s$  in  $(1 - yd)R(1 - ax)$  and  $t$  in  $(1 - xa)R(1 - dy)$  so that

$$(1 - ax - bs) \perp (1 - yd - sb) \quad \text{and} \quad (1 - dy - ct) \perp (1 - xa - tc).$$

Straightforward computations show that

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x & t \\ s & y \end{pmatrix} = \begin{pmatrix} ax + bs & 0 \\ 0 & dy + ct \end{pmatrix},$$

$$\begin{pmatrix} x & t \\ s & y \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} xa + tc & 0 \\ 0 & yd + sb \end{pmatrix},$$

from which we deduce that  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbb{M}_2(R)_q^{-1}$  with quasi-inverse  $\begin{pmatrix} x & t \\ s & y \end{pmatrix}$ . Evidently this means that the row is right reducible, and thus the original row from (\*) is also right reducible, which proves that  $\mathbb{M}_2(R) = \text{cr}(\mathbb{M}_2(R))_q^{-1}$ , as desired.  $\square$

**6.5. Remark.** Theorem 6.4 remains true also in the case where  $R$  is non-unital. Now instead of equation (\*) we must by Definition 4.4 consider an equation

$$(**) \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} - \begin{pmatrix} a & b \\ c & d \end{pmatrix} - \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix} + \begin{pmatrix} e & f \\ g & h \end{pmatrix} = 0,$$

where all the matrices belong to  $\mathbb{M}_2(R)$ . Working in the unital ring  $\mathbb{M}_2(\tilde{R}) = \mathbb{M}_2(R) + \mathbb{M}_2(\mathbb{Z})$ , but using only matrices of the form  $1 - x$ , where  $x \in \mathbb{M}_2(R)$ , we can rewrite (\*\*) in the form (\*). Now all the matrix elements in (\*) belong to  $R$ , except  $a, a', d$  and  $d'$ , which are of the form  $1 - R$ , where, of course, 1 denotes the unit in  $\tilde{R}$ . Each of the reduction steps in the proof of Theorem 6.4 will respect this structure. At the point where we invoke Lemma 6.1 to transform  $b$  (and later  $c$ ) to a quasi-invertible element in a skew corner, it is well to recall that Lemma 6.1 has a non-unital version, cf. Remark 6.2.

**6.6. Corollary.** *Let  $R$  and  $S$  be unital rings. If  $R$  is a QB-ring and  $S$  is Morita equivalent to  $R$ , then  $S$  is also a QB-ring.*

*Proof.* If  $R$  and  $S$  are Morita equivalent, then there is a positive integer  $n$  and an idempotent  $e$  in  $M_n(R)$  such that  $S \simeq eM_n(R)e$  (see, e.g., [1, Corollary 22.7]). Then the result follows using Theorem 6.4 and Corollary 5.8.  $\square$

**6.7. Prime Embeddings.** We say that a subring  $S$  of a ring  $R$  is *primely embedded* if  $p \perp q$  in  $S$  implies  $p \perp q$  in  $R$  for any pair of idempotents  $p$  and  $q$  in  $S$ .

We have already used the fact that any ideal  $I$  in a ring  $R$  is primely embedded. Indeed, if  $p \perp q$  in  $I$  for two idempotents, then  $pRq = p^2Rq^2 \subset pIq = 0$ , and similarly  $qRp = 0$ , so that  $p \perp q$  in  $R$ .

It is clear that if  $R$  is a unital ring and  $S$  is a primely embedded subring containing 1, then  $S_q^{-1} \subset R_q^{-1}$ . Similarly we have  $S_q^\circ \subset R_q^\circ$  in the non-unital case.



**6.8. Proposition.** *Let  $(R_n)$  be a sequence of  $QB$ -rings, and assume that we have homomorphisms  $\varphi_n : R_n \rightarrow R_{n+1}$ , such that  $\varphi_n(R_n)$  is primely embedded in  $R_{n+1}$  for every  $n$ . Then  $R = \varinjlim R_n$  is a  $QB$ -ring.*

*Proof.* The elements in  $R$  may be realized as (equivalence classes of) sequences  $x = (x_n)$ , where  $x_n \in R_n$  and  $\varphi_n(x_n) = x_{n+1}$  eventually (i.e. from a certain  $n_0$  onwards). (Two sequences being equivalent if they agree eventually.)

If  $a = (a_n) \in R$  and  $ax - a - x + b = 0$  for some elements  $x = (x_n)$  and  $b = (b_n)$  in  $R$ , then  $\varphi_n(a_n) = a_{n+1}$ ,  $\varphi_n(b_n) = b_{n+1}$  and  $\varphi_n(x_n) = x_{n+1}$  for all  $n \geq m$  for some  $m$ . Since  $R_m$  is a  $QB$ -ring we can find  $y_m$  in  $R_m$  such that  $a_m - y_m b_m = c_m \in (R_m)_q^\circ$ . Define  $c = (c_n)$  for  $n \geq m$  recursively by  $c_{n+1} = \varphi_n(c_n)$ . If  $c_n \in (R_n)_q^\circ$  then  $\varphi_n(c_n) \in (\varphi_n(R_n))_q^\circ$  by Proposition 3.7. Since  $\varphi_n(R_n)$  is primely embedded in  $R_{n+1}$  it follows that  $c_{n+1} \in (R_{n+1})_q^\circ$ . As  $c_m \in (R_m)_q^\circ$  we see by induction that  $c_n \in (R_n)_q^\circ$  for all  $n$ , whence  $c \in R_q^\circ$ . Defining  $y = (y_n)$  in  $R$  starting with  $y_m$  and inductively setting  $y_{n+1} = \varphi_n(y_n)$  we obtain the equation  $a - yb = c$  in  $R$ , whence  $a \in \text{cl}^\circ(R_q^\circ)$ . Since  $a$  was arbitrary,  $R$  is a  $QB$ -ring.  $\square$

**6.9. Remark.** For each ring  $R$  and each natural number  $n$  there is a canonical embedding  $\iota : \mathbb{M}_n(R) \rightarrow \mathbb{M}_{n+1}(R)$ , where  $\iota(a)_{ij} = a_{ij}$  if  $1 \leq i, j \leq n$ , but  $\iota(a)_{ij} = 0$  if either  $i = n+1$  or  $j = n+1$ . This embedding is not unital, so the direct limit  $\mathbb{M}_\infty(R) = \varinjlim \mathbb{M}_n(R)$  is a non-unital ring consisting of those matrices  $a = (a_{ij})$  over  $R$  such that  $a_{ij} = 0$  if  $i + j \geq m$  for some  $m$  (depending on  $a$ ).

Combining Theorem 6.4 (maybe in its non-unital version described in Remark 6.5) and Proposition 6.8 (which is allowed, since each embedding  $\iota$  certainly is prime) we therefore obtain the following result:

**6.10. Corollary.** *If  $R$  is a  $QB$ -ring, then so is  $\mathbb{M}_\infty(R)$ .*  $\square$

## 7. Extensions of $QB$ -Rings

**7.1. Proposition.** *If  $I$  is an ideal in a unital  $QB$ -ring  $R$ , then*

$$(R_q^{-1} + I)/I = (R/I)_q^{-1}.$$

*Proof.* Let  $\pi : R \rightarrow R/I$  denote the quotient morphism. From Proposition 3.7 we know that  $\pi(R_q^{-1}) \subset (R/I)_q^{-1}$ , so it only remains to show that every quasi-invertible element in  $R/I$  lifts to a quasi-invertible element in a  $QB$ -ring  $R$ .

For this we may consider  $a$  and  $b$  in  $R$  such that  $\pi(a)$  and  $\pi(b)$  are elements in  $(R/I)_q^{-1}$  with  $(1 - \pi(b)\pi(a)) \perp (1 - \pi(a)\pi(b))$  in  $\pi(R)$ . Thus, upstairs we have the relations

$$(1 - ba)R(1 - ab) \subset I \quad \text{and} \quad (1 - ab)R(1 - ba) \subset I.$$

Since  $R$  is a  $QB$ -ring the trivial equation  $ab + (1 - ab) = 1$  shows that we have

$$v = b + y(1 - ab) \in R_q^{-1}$$

for some  $y$  in  $R$ . Now choose a quasi-inverse  $u$  for  $v$  in  $R_q^{-1}$ . By (\*) in Theorem 2.3 we then also have

$$w = u + a(1 - vu) + (1 - uv)a \in R_q^{-1}.$$

Passing to  $R/I$  we evidently get  $\pi(v)\pi(u)\pi(v) = \pi(v)$ . But we also have

$$\begin{aligned} \pi(v)\pi(a)\pi(v) &= \pi((b + y(1 - ab))a(b + y(1 - ab))) \\ &= \pi(ba(b + y(1 - ab))) = \pi(b + y(1 - ab)) = \pi(v), \end{aligned}$$

since  $(1 - \pi(ba))\pi(y)(1 - \pi(ab)) = 0$ . Thus, both  $\pi(a)$  and  $\pi(u)$  are partial inverses for  $\pi(v)$ , whence by Theorem 2.3

$$\pi(a) = \pi(u) + \pi(a)(1 - \pi(vu)) + (1 - \pi(uv))\pi(a).$$

It follows that  $\pi(a) = \pi(w)$ , so that  $w$  is the required lift of  $\pi(a)$  in  $R_q^{-1}$ .  $\square$

**7.2. Theorem** (Cf. [8, 6.1]). *Let  $I$  be an ideal in a unital ring  $R$ . Then  $R$  is a  $QB$ -ring if and only if the following conditions are satisfied:*

- (i)  $R/I$  is a  $QB$ -ring ;
- (ii)  $(R_q^{-1} + I)/I = (R/I)_q^{-1}$ ;
- (iii)  $I + R_q^{-1} \subset \text{cl}(R_q^{-1})$ .

*Proof.* If  $R$  is a  $QB$ -ring then the first two conditions are satisfied by Corollary 3.8 and Proposition 7.1, while Condition (iii) is trivially true.

Conversely, if the three conditions are satisfied, take an arbitrary element  $a$  in  $R$  and assume that  $xa + b = 1$  for some  $x, b$  in  $R$ . If  $\pi: R \rightarrow R/I$  denotes the quotient morphism, then  $\pi(x)\pi(a) + \pi(b) = 1$  in  $R/I$ , and since  $R/I$  is a  $QB$ -ring this implies that  $\pi(a) + \pi(yb) \in (R/I)_q^{-1}$  for some  $y$  in  $R$ . By Condition (ii) we can find an element  $t$  in  $I$  such that  $a + yb - t \in R_q^{-1}$ .

Using Condition (iii) this implies that  $a + yb \in \text{cl}(R_q^{-1})$ , whence  $a \in \text{cl}(\text{cl}(R_q^{-1}))$  by the definition of  $\text{cl}$ . But then  $a \in \text{cl}(R_q^{-1})$  by Condition (iii) in Lemma 3.2, and thus  $R = \text{cl}(R_q^{-1})$ , as desired.  $\square$

**7.3. Remark.** With obvious modifications the results in 7.1 and 7.2 remain true also in the non-unital case. If  $R$  is a non-unital ring with an ideal  $I$  and quotient morphism  $\pi: R \rightarrow R/I$ , we consider the unitization  $\tilde{R} = R \oplus \mathbb{Z}$  as in 3.1 and extend  $\pi$  to a unital morphism  $\tilde{\pi}: \tilde{R} \rightarrow R/I$  (if  $R/I$  has a unit) or  $\tilde{\pi}: \tilde{R} \rightarrow (R/I)^\sim$  (if  $R/I$  has no unit) by setting  $\tilde{\pi}(1) = 1$ .

The argument in Proposition 7.1 now shows that  $\pi(R_q^\circ) = (R/I)_q^\circ$  if  $R$  is a  $QB$ -ring, and in Theorem 7.2 we just have to replace the last two conditions with

- (ii')  $(R_q^\circ + I)/I = (R/I)_q^\circ$ ;
- (iii')  $I + R_q^\circ \subset \text{cl}^\circ(R_q^\circ)$ .

**7.4. Remark.** Condition (iii) in Theorem 7.2 is not easy to verify directly, so it is fortunate that it is vacuously satisfied in a number of interesting cases, cf. Theorems 7.11 & 7.14.

On the other hand, the condition is of independent interest and we shall devote some attention to it. Note first that by Theorem 4.9 the condition implies that the ideal is a  $QB$ -ring. Secondly observe from Condition (x) in Lemma 3.2 that if an ideal  $I$  satisfies Condition (iii), then actually  $I + \text{cl}(R_q^{-1}) \subset \text{cl}(R_q^{-1})$ .

Having identified the extremally rich  $C^*$ -algebras with those  $C^*$ -algebras which are  $QB$ -rings, cf. Proposition 9.1, it follows from [8, Example 6.12] that we can not in Theorem 7.2 replace Condition (iii) with the weaker condition:  $I$  is a  $QB$ -ring. By necessity this means that the extension theory for  $QB$ -rings is somewhat more complicated than that governing  $B$ -rings and exchange rings.

**7.5. Lemma.** *If  $I$  is a  $QB$ -ideal in a unital ring  $R$ , then*

$$I + R^{-1} \subset \text{cl}(R_q^{-1}).$$

*Proof.* Take  $a$  in  $R^{-1}$  and  $t$  in  $I$  and assume that  $x(a - t) + b = 1$  for some  $x$  and  $b$  in  $R$ . Then  $xa(1 - a^{-1}t) + b = 1$ , so by Lemma 4.6 there are elements  $r, r'$  in  $I$  such that  $(1 - r')(1 - a^{-1}t) + rb = 1$ .

Since  $I$  is a  $QB$ -ring  $a^{-1}t \in \text{cl}(I_q^\circ)$ , whence  $1 - a^{-1}t \in \text{cl}(R_q^{-1})$  by Proposition 4.7, so that  $a - t \in \text{cl}(R_q^{-1})$ .  $\square$

**7.6. Lemma.** *If  $I$  is a  $B$ -ideal in a unital ring  $R$ , then*

$$I + R_q^{-1} \subset \text{cl}(R_q^{-1}).$$

*Proof.* Take  $u$  in  $R_q^{-1}$  and  $t$  in  $I$ , and assume that

$$(*) \quad x(u - t) + b = 1$$

for some  $x$  and  $b$  in  $R$ . Choose a quasi-inverse  $v$  for  $u$  so that the two idempotents  $p = uv$  and  $q = vu$  satisfy  $(1 - p) \perp (1 - q)$ . Now rewrite the equation  $(*)$  as

$$1 = xu(1 - vt) - x(1 - p)t + b.$$

Using Lemma 4.6 (with  $vt$  and  $b - x(1 - p)t$  in place of  $a$  and  $b$ ) and that  $I$  is a  $B$ -ideal, it follows that  $w_1 = 1 - vt + s(b - x(1 - p)t) \in R^{-1}$  for some  $s$  in  $I$  (and actually  $w_1 - 1 \in I$ ). It follows that

$$\begin{aligned} uw_1 &= u - pt + usb - usx(1 - p)t \\ &= u - t + usb + (1 - usx(1 - p))(1 - p)t. \end{aligned}$$

Since  $(usx(1 - p))^2 = 0$  the element  $w_2 = 1 + usx(1 - p)$  is invertible with  $w_2^{-1} = 1 - usx(1 - p)$ . Moreover,  $w_2u = w_2^{-1}u = u$ . Therefore, with  $t' = tw_1^{-1}$  we have

$$w_2(u - t + usb)w_1^{-1} = u - (1 - p)t' \in R_q^{-1}$$

by Theorem 2.3. It follows that also  $u - t + usb \in R_q^{-1}$ , so  $u - t \in \text{cl}(R_q^{-1})$ .  $\square$

**7.7. Remark.** Inspection of the preceding proof shows that if  $I$  is a  $B$ -ideal in  $R$ , then we also have the relations

- (i)  $I + R^{-1} \subset \text{cl}(R^{-1})$ ;
- (ii)  $I + R_r^{-1} \subset \text{cl}(R_r^{-1})$ ;
- (iii)  $I + R_\ell^{-1} \subset \text{cl}(R_\ell^{-1})$ .

For the first two it is actually easier to use a direct argument, but the proof of relation (iii) needs the full force of the argument in Lemma 7.6.

**7.8. Proposition.** *In any unital ring  $R$  there is a largest ideal, denoted by  $I_{qb}$ , such that  $I_{qb} + R_q^{-1} \subset \text{cl}(R_q^{-1})$ .*

*Proof.* If  $I_1$  and  $I_2$  both satisfy the condition  $I_i + R_q^{-1} \subset \text{cl}(R_q^{-1})$ , then by Condition (x) in Lemma 3.2 they also satisfy  $I_i + \text{cl}(R_q^{-1}) \subset \text{cl}(R_q^{-1})$  for  $i = 1, 2$ , whence

$$I_1 + I_2 + R_q^{-1} \subset I_1 + \text{cl}(R_q^{-1}) \subset \text{cl}(R_q^{-1}).$$

Therefore  $I_{qb}$  is simply the sum of all the ideals that satisfy the desired condition.  $\square$

**7.9. Proposition** (Cf. [11, 2.14]). *If  $R$  is a unital ring such that  $R$  is additively generated by its units ( $R = R^{-1} + R^{-1} + \dots$ ) then*

$$I_{qb} = \{x \in R \mid x + \text{cl}(R_q^{-1}) \subset \text{cl}(R_q^{-1})\}.$$

*Proof.* Obviously  $I_{qb}$  is contained in the set  $S$  defined by the right side of the equation, so by maximality it suffices to show that  $S$  is an ideal. From the definition we see  $S + S \subset S$  and also  $R^{-1}SR^{-1} = S$ , because  $R^{-1}\text{cl}(R_q^{-1})R^{-1} = \text{cl}(R_q^{-1})$ . Since every element  $a$  in  $R$  has a representation  $a = \sum a_i$  with  $a_i$  in  $R^{-1}$  it follows that  $aS \subset S$ , and  $Sa \subset S$ , whence  $S$  is an ideal.  $\square$

**7.10. Remark.** Lemma 7.6 shows that the  $QB$ -ideal  $I_{qb}$  defined above contains every ideal in  $R$  of Bass stable rank one. Since the sum of  $B$ -ideals is again a  $B$ -ideal,  $I_{qb}$  therefore contains the maximal  $B$ -ideal  $I_b$  in  $R$ . Unfortunately the sum of  $QB$ -ideals may fail to be a  $QB$ -ideal, cf. [8, Example 6.12], so we can not hope to describe  $I_{qb}$  as “the maximal  $QB$ -ideal”. In  $C^*$ -algebra theory one may instead characterize  $I_{qb}$  as the “largest well-behaved  $QB$ -ideal”, in the sense that  $I_{qb}$  is the maximal ideal such that  $I_{qb} + B$  is a  $QB$ -algebra for any  $QB$ -subalgebra  $B$  of  $A$  containing 1, such that  $B_q^{-1} \subset A_q^{-1}$ , cf. [11, Theorem 2.14]. The proof, however, depends heavily on topological arguments.

**7.11. Theorem** (Cf. [8, 6.3]). *If  $I$  is a  $B$ -ideal in a unital ring  $R$ , then  $R$  is a  $QB$ -ring if and only if the following conditions are satisfied:*

- (i)  $R/I$  is a  $QB$ -ring;
- (ii)  $(R_q^{-1} + I)/I = (R/I)_q^{-1}$ .

*Proof.* By Theorem 7.2 we only need to show that the two conditions are sufficient. However, given the conditions it follows from Lemma 7.6 that Condition (iii) in Theorem 7.2 is satisfied, whence  $R$  is a  $QB$ -ring.  $\square$

**7.12. Corollary.** *If  $I$  is a  $B$ -ideal in a unital ring  $R$ , and  $S$  is a  $QB$ -subring of  $R$  containing 1, such that  $S$  is primely embedded in  $R$  and  $R = I + S$ , then  $R$  is a  $QB$ -ring.*

*Proof.* Since  $R/I$  is isomorphic to  $S/I \cap S$  we know that  $R/I$  is a  $QB$ -ring by Corollary 3.8. By Theorem 7.11 we therefore only need to verify that  $(R_q^{-1} + I)/I = (R/I)_q^{-1}$ . However, if  $v \in (R/I)_q^{-1}$  there is by Proposition 7.1 an element  $u$  in  $S_q^{-1}$  such that  $u + I = v$ . Since  $S$  is primely embedded in  $R$  this means that  $u \in R_q^{-1}$  cf. 6.7.  $\square$

**7.13. Remark.** For each unital ring  $R$  its Jacobson radical  $\mathcal{J}(R)$  is a  $B$ -ideal, since  $1 - x \in R^{-1}$  for any  $x$  in  $\mathcal{J}(R)$ . Moreover, any lift of a left, respectively right invertible element in  $R/\mathcal{J}(R)$  will be left, respectively right invertible in  $R$ . It follows from Theorem 7.11 that if  $R/\mathcal{J}(R)$  is a prime ring, then  $R$  is a  $QB$ -ring if and only if  $R/\mathcal{J}(R)$  is a  $QB$ -ring.

**7.14. Theorem** (Cf. [8, 6.6]). *If  $I$  is a  $QB$ -ideal in a unital ring  $R$ , such that  $R/I$  is a  $B$ -ring, then  $R$  is a  $QB$ -ring provided that  $(R^{-1} + I)/I = (R/I)^{-1}$ .*

*Proof.* Take any  $a$  in  $R$  and assume that  $xa + b = 1$  for some elements  $x$  and  $b$  in  $R$ . Then with  $\pi: R \rightarrow R/I$  the quotient morphism we also have  $\pi(x)\pi(a) + \pi(b) = 1$  in  $R/I$ , and since  $R/I$  is a  $B$ -ring this implies that  $\pi(a) + \pi(y)\pi(b) = \pi(u) \in (R/I)^{-1}$  for some  $y$  and  $u$  in  $R$ . Since  $\pi(R^{-1}) = (R/I)^{-1}$  we may assume that  $u \in R^{-1}$ , so that we have the equation  $a - t + yb = u$  for some  $t$  in  $I$ . By Lemma 7.5 this implies that  $a + yb \in \text{cl}(R_q^{-1})$ . Therefore  $a \in \text{cl}(\text{cl}(R_q^{-1})) = \text{cl}(R_q^{-1})$  (see Condition (iii) in Lemma 3.2).  $\square$

**7.15. Corollary.** *If  $I$  is an  $QB$ -ideal in a unital ring  $R$ , and  $S$  is a  $B$ -subring of  $R$  containing 1, such that  $R = I + S$ , then  $R$  is a  $QB$ -ring.*

*Proof.* Since  $R/I$  is isomorphic to  $S/I \cap S$  we know that  $R/I$  is a  $B$ -ring, because Bass stable rank one is preserved under quotients. To apply Theorem 7.14 we therefore need only to verify that invertibles lift from  $R/I$ . But invertibles certainly lift from quotients of  $B$ -rings, so if  $x \in (R/I)^{-1} (= (S/(I \cap S))^{-1})$  there is an element  $u$  in  $S^{-1}$  such that  $u + I = v$ . As  $S^{-1} \subset R^{-1}$  we conclude that  $\pi(R^{-1}) = (R/I)^{-1}$ , as desired.  $\square$

**7.16. Toeplitz-like Examples.** We present an example of a unital  $QB$ -ring  $S$  which is von Neumann regular and is an extension of two  $B$ -rings, but has Bass stable rank two. As we shall see, this example can be thought of as an algebraic analogue of the Toeplitz algebra, since it is generated, in a suitable sense, by a unilateral shift. Other examples of non-regular (even not exchange) rings will be given later (cf. 8.9). Our construction is modelled after the example given in [15]. We provide some details for the convenience of the reader.

Let  $\mathbb{F}$  be a countable field and let  $t$  be an indeterminate. Let  $\mathbb{F}(t)$  be the field of rational functions on  $\mathbb{F}$ , and let  $\delta$  be the valuation on  $\mathbb{F}(t)$  associated with the maximal ideal  $(t)$  of  $\mathbb{F}[t]$ , i.e.  $\delta(0) = +\infty$  and  $\delta(t^n f(t)/g(t)) = n$  where  $t$  does not

divide  $f(t)g(t)$ . Let  $V = \{x \in \mathbb{F}(t) \mid \delta(x) \geq 0\}$  be the valuation ring associated with  $\delta$ . Note that  $V$  is a local ring with maximal ideal  $\{x \in \mathbb{F}(t) \mid \delta(x) > 0\}$ .

We claim that the vector space  $W = \mathbb{F}(t)$  has an  $\mathbb{F}$ -basis  $\{v_i\}_{i \in \mathbb{Z}}$  such that  $\delta(v_i) = i$  for all  $i$  in  $\mathbb{Z}$ . First note that  $W$  is of countable dimension over  $\mathbb{F}$  because  $\mathbb{F}$  is countable. Take an  $\mathbb{F}$ -basis  $\{w_n\}_{n \geq 0}$  for  $W$ . We can modify this basis in order to get  $\delta(w_i) \neq \delta(w_j)$  for all  $i \neq j$ . In fact, if  $\delta(w_n) = \delta(w_i)$  for  $i < n$  there is an element  $\alpha_i$  in  $\mathbb{F}$  such that  $w_n/w_i - \alpha_i \in tV$ , which implies that  $\delta(w_n - \alpha_i w_i) > \delta(w_n)$ . If  $\delta(w_n - \alpha_i w_i) = \delta(w_j)$  for some  $j < n$ , then the same argument shows that there is an element  $\alpha_j$  in  $\mathbb{F}$  such that  $\delta(w_n - \alpha_i w_i - \alpha_j w_j) > \delta(w_j) > \delta(w_i)$ . Thus we get an element  $w_n - \alpha_i w_i - \dots - \alpha_k w_k$ , such that  $\delta(w_n - \alpha_i w_i - \dots - \alpha_k w_k) \neq \delta(w_t)$  for all  $t < n$ . Then we substitute  $w_n$  by  $w_n - \alpha_i w_i - \dots - \alpha_k w_k$ .

Now assume that  $\delta(w_i) \neq \delta(w_j)$  for  $i \neq j$ . Writing an arbitrary element  $v$  of  $W$  as  $v = \lambda_{i_1} w_{i_1} + \dots + \lambda_{i_k} w_{i_k}$  with  $\delta(w_{i_1}) < \delta(w_{i_2}) < \dots < \delta(w_{i_k})$ , we have  $\delta(v) = \delta(w_{i_1})$ . Since  $\delta(t^i) = i$  for all  $i$  in  $\mathbb{Z}$  we have a bijective correspondence  $\varphi : \mathbb{Z} \rightarrow \mathbb{Z}^+$  such that  $\delta(w_{\varphi(i)}) = i$ . Take  $v_i = w_{\varphi(i)}$  for all  $i$  in  $\mathbb{Z}$ . Note that  $V = \langle \{v_i\}_{i \geq 0} \rangle$ , and obviously we may assume that  $v_0 = 1$ .

Consider the representation  $\lambda : \mathbb{F}(t) \rightarrow \text{End}_{\mathbb{F}}(W)$  given by multiplication. Let  $\pi : W \rightarrow W$  be the projection onto  $V$  with kernel  $\langle \{v_i\}_{i < 0} \rangle$ . Identifying  $\pi \text{End}_{\mathbb{F}}(W) \pi$  with  $\text{End}_{\mathbb{F}}(V)$ , we may regard  $\pi \lambda(\mathbb{F}(t)) \pi$  as a subring of  $\text{End}_{\mathbb{F}}(V)$ . Each endomorphism of  $V$  has a column-finite matrix associated to the basis  $\{v_i\}_{i \geq 0}$  of  $V$ . Since  $\lambda(v)(v_i) = vv_i \in \langle v_{i+\delta(v)}, v_{i+\delta(v)+1}, \dots \rangle$  (we use that  $\delta(vv_i) = i + \delta(v)$ ), it follows that  $\pi \lambda(\mathbb{F}(t)) \pi \subset \mathbb{B}(\mathbb{F})$ , where  $\mathbb{B}(\mathbb{F})$  is the algebra of row-and-column finite matrices over  $\mathbb{F}$  (and we identify an element of  $\pi \lambda(\mathbb{F}(t)) \pi$  with its matrix with respect to the basis  $\{v_i\}_{i \geq 0}$  of  $V$ ). Let  $\mathbb{M}_{\infty}(\mathbb{F})$  be the ideal of  $\mathbb{B}(\mathbb{F})$  consisting of matrices with only a finite number of non-zero entries, cf. 6.9, and consider the ring  $S = \pi \lambda(\mathbb{F}(t)) \pi + \mathbb{M}_{\infty}(\mathbb{F})$ . Since  $\pi \lambda(u) \pi \lambda(v) \pi - \pi \lambda(uv) \pi \in \mathbb{M}_{\infty}(\mathbb{F})$  for all  $u, v$  in  $\mathbb{F}(t)$ , we see that  $S$  is an  $\mathbb{F}$ -subalgebra of  $\mathbb{B}(\mathbb{F})$ . There is a surjective homomorphism  $\rho : S \rightarrow \mathbb{F}(t)$  defined by  $\rho(\pi \lambda(u) \pi + m) = u$  for all  $u$  in  $\mathbb{F}(t)$  and all  $m$  in  $\mathbb{M}_{\infty}(\mathbb{F})$ . Since  $\mathbb{M}_{\infty}(\mathbb{F})$  is regular and  $\mathbb{F}(t)$  is also regular this shows that  $S$  is a regular ring.

Now let  $a = \pi \lambda(t) \pi$  and  $b = \pi \lambda(t^{-1}) \pi$ . Since  $\lambda(t) \pi = \pi \lambda(t) \pi$  we get  $ba = 1$ . However,  $ab$  is not equal to 1, and  $1 - ab$  is a one-dimensional idempotent in  $\mathbb{M}_{\infty}(\mathbb{F})$ . It follows that  $S$  is not directly finite. In particular it is not a  $B$ -ring, and it is an extension of the  $B$ -rings  $\mathbb{M}_{\infty}(\mathbb{F})$  and  $\mathbb{F}(t)$ . Notice that  $\text{bsr}(S) \leq \max\{\text{bsr}(\mathbb{F}(t)), \text{bsr}(\mathbb{M}_{\infty}(\mathbb{F})) + 1\} = 2$ , whence  $\text{bsr}(S) = 2$ . Also we see that  $a, b \in S_q^{-1}$ , and since every non-zero element in  $\mathbb{F}(t)$  has the form  $t^i v$ , where  $i \in \mathbb{Z}$  and  $v$  is an invertible in  $V$ , we get that every non-zero element in  $\mathbb{F}(t)$  lifts to a quasi-invertible element of  $S$ . (Note that the map  $\pi \lambda(-) \pi$  provides an isomorphism from  $V$  onto a subalgebra of  $S$ .)

It follows from Theorem 7.11 that  $S$  is a  $QB$ -ring.

The next example is based on a construction due to Bergman (see [16, Example 5.10], and also [20]).

**7.17. Example.** In the setting of Example 7.16, consider the map  $\rho : S \rightarrow \mathbb{F}(t)$ . Let  $S^o$  be the opposite ring of  $S$ . Since  $\mathbb{F}(t)$  is commutative we have an

induced surjection  $\rho : S^o \rightarrow \mathbb{F}(t)$ . Taking the pullback of both maps we get the ring  $T = \{(x, \bar{y}) \mid \rho(x) = \rho(y)\}$ . The ring  $T$  has a unique maximal ideal, viz.  $\mathbb{M}_\infty(\mathbb{F}) \times \mathbb{M}_\infty(\mathbb{F})^o$  and  $T/(\mathbb{M}_\infty(\mathbb{F}) \times \mathbb{M}_\infty(\mathbb{F})^o) \simeq \mathbb{F}(t)$ , which proves that  $T$  is a regular ring. The ring  $T$  is directly finite by [20, Lemma 13], but has a quotient isomorphic to  $S$  and a quotient isomorphic to  $S^o$ , so it is not a  $B$ -ring. The elements  $(a, \bar{a})$  and  $(b, \bar{b})$  are quasi-invertible in  $T$  (though they are not right or left invertible) and it follows again from Theorem 7.11 that  $T$  is a  $QB$ -ring.

## 8. Exchange Rings

**8.1. Definitions.** A unital ring  $R$  is called an *exchange ring* if for every element  $a$  in  $R$  there is an idempotent  $p$  in  $aR$  such that  $1 - p \in (1 - a)R$ . This is not the original definition (which concerns a finite exchange property in  $R$ -modules, cf. [30]), but is an equivalent description found by Goodearl and Nicholson, see [17] and [21]. Rewriting the condition  $1 - p \in (1 - a)R$  as an equation  $1 - p = (1 - a)(1 - y)$  for some  $y$  in  $R$ , i.e.  $p = a + y - ay$ , we obtain a definition of an exchange ring suitable for the non-unital case, cf. [2].

The class of exchange rings is extensive and includes all von Neumann regular rings, all  $\pi$ -regular rings, the semi-perfect rings (which are exactly the exchange rings that are semi-local), right self-injective rings and  $C^*$ -algebras of real rank zero. (In fact, the  $C^*$ -algebras which are exchange rings are precisely those of real rank zero, by [4, Theorem 7.2].) The class of exchange rings is stable under ideals and quotients as well as corners and matrix tensoring; and if  $0 \rightarrow I \rightarrow R \rightarrow Q \rightarrow 0$  is a short exact sequence of rings, then  $R$  is exchange if and only if both  $I$  and  $Q$  are exchange and idempotents lift from  $Q$  to  $R$ , cf. [2, Theorem 2.2]. Evidently the class is also stable under direct limits.

**8.2. Proposition.** *In a unital, semi-primitive exchange ring  $R$  an element in  $R^r$  is maximal with respect to the order  $\leq$  if and only if it belongs to  $R_q^{-1}$ .*

*Proof.* By Proposition 2.8 we need only consider a maximally extended element  $a$  in  $R^r$ , and prove that  $a \in R_q^{-1}$ .

Towards this end choose a partial inverse  $x$  for  $a$  and set  $p = 1 - ax$  and  $q = 1 - xa$ . If  $pRq \neq 0$  we can find a non-zero element  $y$  in  $pRq$ . Since  $R$  is semi-primitive there is a maximal right ideal  $J$  of  $R$  such that  $y \notin J$ . Hence  $1 - yc \notin R_r^{-1}$  for some  $c$  in  $R$ . Since  $R$  is an exchange ring we can find an idempotent  $e$  in  $ycR$  such that  $1 - e \in (1 - yc)R$ . We have made sure that  $1 - e \neq 1$ , so  $e \neq 0$ . Since  $e = eycd$  for some  $d$  in  $R$ , the element  $ey \in pRq \setminus \{0\}$  and  $(ey)(cd)(ey) = e^2ey = ey$  so  $ey$  is von Neumann regular with  $cd$  as partial inverse.

By Lemma 2.7 the element  $b = a + ey$  is von Neumann regular in  $R$  and properly extends  $a$ , contradicting the maximality. Thus, after all,  $pRq = 0$  (and also  $qRp = 0$ ), whence  $a \in R_q^{-1}$ .  $\square$

**8.3. Examples.** Corollary 5.11 and Proposition 8.2 have different implications, despite their similarity. In a  $QB$ -ring every von Neumann regular element extends to a maximal one, but maybe they are all maximal to begin with (except zero). This happens e.g. if  $R = C([0, 1])$ , where each non-zero regular element is invertible.

In an exchange ring, by contrast, there is an abundance of idempotents, and therefore also a great variety of von Neumann regular elements. The problem here is that they might not all extend to quasi-invertible elements. Of course, for the idempotents there are no problems: each extends to 1.

To construct an example of an exchange ring with a regular element that does not extend, take  $A = \mathbb{B}(\mathcal{H}) \oplus \mathbb{B}(\mathcal{H})$ . Represent  $A$  with infinite multiplicity on the Hilbert space  $\mathcal{K} = (\mathcal{H} \oplus \mathcal{H}) \otimes \ell^2$  (so every operator in  $A$  is repeated infinitely often along the diagonal), and put  $R = A + \mathbb{K}(\mathcal{K})$ . Thus,  $R$  is a split extension of the algebra of compact operators on  $\mathcal{K}$  and  $A$ . Evidently both  $\mathbb{K}(\mathcal{K})$  and  $A$  are exchange rings (they are  $C^*$ -algebras of real rank zero), and projections lift since the extension splits, so  $R$  is an exchange ring. Let  $s$  denote the unilateral shift in  $\mathbb{B}(\mathcal{H})$  (or any other non-unitary isometry). Then  $u = s \oplus s^* \in A_q^{-1}$  with quasi-inverse  $u^* = s^* \oplus s$ . However, since  $R$  is a primitive algebra its quasi-invertible elements are either left or right invertible, so  $u \notin R_q^{-1}$ . If  $u$  could be extended to a left or right invertible element  $w$  in  $R$ , then in the quotient  $R/\mathbb{K}(\mathcal{K}) = A$  we would have  $w - u + \mathbb{K}(\mathcal{K}) = 0$ , since  $u$  is maximally extended in  $A$ . But since both the kernel and the co-kernel of  $u$  on  $\mathcal{K}$  are infinite dimensional, no compact perturbation can make  $u$  left or right invertible.

As the following result shows, the global obstruction to extension of regular elements in an exchange ring  $R$  is exactly that  $R$  fails to be a  $QB$ -ring.

**8.4. Theorem.** *If  $R$  is a unital exchange ring the following conditions are equivalent:*

- (i)  $R$  is a  $QB$ -ring;
- (ii) Every element in  $R^r$  extends to an element in  $R_q^{-1}$ ;
- (iii) For every element  $x$  in  $R^r$  there is a  $v$  in  $R_q^{-1}$  such that  $x = xvx$ .

*Proof.* (i)  $\implies$  (ii) This is immediate from Corollary 5.10.

(ii)  $\implies$  (i) Given an equation  $xa + b = 1$  in  $R$  we use the exchange property to find an idempotent  $p$  in  $Rxa$ , such that  $1 - p \in Rb$ . Specifically,  $p = rxa$  and  $1 - p = sb$ . Then  $ap$  is a von Neumann regular element with partial inverse  $rx$ , and by assumption it extends to an element  $u$  in  $R_q^{-1}$ . Thus by definition, and using Lemma 2.6 if necessary,  $ap = u(rx)(ap) = up$ . It follows that with  $y = (u - a)s$  we can write

$$u = up + u(1 - p) = ap + u(1 - p) = a + (u - a)sb = a + yb,$$

as desired.

(ii)  $\iff$  (iii) This is Proposition 2.9. □

The next lemma is an adaptation of [31, Theorem 2.1] to our situation.



**8.5. Lemma.** *Let  $R$  be a unital ring. Then the following conditions are equivalent:*

- (i)  $R$  is a  $QB$ -ring;
- (ii) *Given any  $R$ -module  $M$  and decompositions  $M = A_1 \oplus H = A_2 \oplus K$  with  $A_1 \simeq A_2 \simeq (R_R)^n$  for some  $n \geq 1$ , there exists a pair of orthogonal ideals  $I$  and  $J$  and decompositions  $M = E \oplus B \oplus H = E \oplus C \oplus K$  such that  $BI = B$  and  $CJ = C$ ;*
- (iii) *Given any  $R$ -module  $M$  and decompositions  $M = A_1 \oplus H = A_2 \oplus K$  with  $A_1 \simeq A_2 \simeq (R_R)^n$  for some  $n \geq 1$ , there exists a pair of orthogonal ideals  $I$  and  $J$  and decompositions  $M = E_1 \oplus N = E_2 \oplus N$  such that  $A_1 = E_1 \oplus B$  and  $A_2 = E_2 \oplus C$  with  $BI = B$  and  $CJ = C$ .*

*Proof.* (i)  $\implies$  (ii) For  $i = 1, 2$  let  $\rho'_i: M \rightarrow A_i$  and  $\rho_1: M \rightarrow H$  and  $\rho_2: M \rightarrow K$  denote the projections and  $\tau'_i: A_i \rightarrow M$  and  $\tau_1: H \rightarrow M$  and  $\tau_2: K \rightarrow M$  the corresponding injections. Fix isomorphisms  $\iota_i: A_i \rightarrow R^n$  and put  $\pi_i = \iota_i \rho'_i$  and  $\sigma_i = \tau'_i \iota_i^{-1}$ . Then  $\pi_i \sigma_i = 1_{R^n}$  and  $\sigma_i \pi_i + \tau_i \rho_i = 1_M$ . It follows that

$$1_{R^n} = \pi_2 \sigma_2 = \pi_2 (\sigma_1 \pi_1 + \tau_1 \rho_1) \sigma_2 = (\pi_2 \sigma_1) (\pi_1 \sigma_2) + (\pi_2 \tau_1 \rho_1 \sigma_2) = ax + b.$$

Identifying  $\mathbb{M}_n(R)$  with  $\text{End}_R(R^n)$ , and using that  $\mathbb{M}_n(R)$  is a  $QB$ -ring, cf. Theorem 6.4, we can find  $y$  in  $\mathbb{M}_n(R)$  and  $u$  in  $(\mathbb{M}_n(R))_q^{-1}$ , such that  $u = a + by$ , i.e.

$$u = \pi_2 (\sigma_1 + \tau_1 \rho_1 \sigma_2 y).$$

Choose a quasi-inverse  $v$  for  $u$  and define the idempotents  $p = vu$  and  $q = uv$  in  $\mathbb{M}_n(R)$ . Moreover, let  $\varphi = \sigma_1 + \tau_1 \rho_1 \sigma_2 y$  in  $\text{Hom}(R^n, M)$ . Then from the equation above we see that  $\pi_2 \varphi = u$ . We also compute  $\pi_1 \varphi = \pi_1 (\sigma_1 + \tau_1 \rho_1 \sigma_2 y) = 1_{R^n}$ , since  $\pi_1 \tau_1 = 0$ .

Let  $D_1 = \ker p \pi_1$  and  $D_2 = \ker q \pi_2$ , and put  $E = \varphi p(R^n)$  in  $M$ . If  $m \in E \cap D_1$ , then  $m = \varphi p(a)$  for some  $a$  in  $R^n$ , where

$$0 = p \pi_1(m) = p \pi_1 \varphi p(a) = p 1_{R^n} p(a) = p(a).$$

Consequently  $m = 0$ , and so  $E \cap D_1 = 0$ . Similarly,  $E \cap D_2 = 0$ . Now take  $m$  in  $M$  and write

$$m = \varphi p \pi_1(m) + (m - \varphi p \pi_1(m)).$$

Then  $p \pi_1(m - \varphi p \pi_1(m)) = p \pi_1(m) - p 1_{R^n} p \pi_1(m) = 0$ , so  $M = E \oplus D_1$ . Similarly  $M = E \oplus D_2$ .

Now we decompose further: Let  $B = \sigma_1(1 - p)(R^n)$  and  $C = \sigma_2(1 - q)(R^n)$  in  $M$ . Then

$$D_1 = \ker p \pi_1 = \ker \pi_1 \oplus \sigma_1(1 - p)(R^n) = H \oplus B.$$

Similarly,

$$D_2 = \ker q \pi_2 = \ker \pi_2 \oplus \sigma_2(1 - q)(R^n) = K \oplus C.$$

Finally, let  $\mathbb{M}_n(I)$  and  $\mathbb{M}_n(J)$  be the ideals in  $\mathbb{M}_n(R)$  generated by  $1 - p$  and  $1 - q$ , respectively, where  $I$  and  $J$  are ideals in  $R$ . Since  $(1 - p) \perp (1 - q)$  it follows that

$I \perp J$ , and evidently  $BI = B$  and  $CJ = C$ . Moreover,  $E \oplus B \oplus H = E \oplus D_1 = M = E \oplus D_2 = E \oplus C \oplus K$ , as desired.

(ii)  $\implies$  (i) Take  $(a_1, a_2)$  in  $R^2$  such that  $a_1R + a_2R = R$ . Then  $(a_1, a_2)$  provides a split epimorphism  $\varphi : M \rightarrow R$ , where  $M = R^2$ , so we have decompositions  $M = R \oplus R = A_2 \oplus K$ , where  $A_2 \simeq R_R$  and  $K$  is the kernel of  $\varphi$ . Therefore there exists a pair of orthogonal ideals  $I$  and  $J$  and a decomposition

$$M = E \oplus B \oplus R = E \oplus C \oplus K$$

with  $BI = B$  and  $CJ = C$ . Write  $E_1 = \varphi(E)$  and  $C_1 = \varphi(C)$ . Then  $R = E_1 \oplus C_1$ . Let  $e$  in  $R = \text{End}(R_R)$  be the projection onto  $E_1$  with kernel  $C_1$ , so that  $E_1 = eR$ . Notice that  $1 - e \in J$ . Let  $\psi_1 : R \rightarrow M$  be the inverse of  $\varphi$  restricted to  $E \oplus C$  and let  $\psi = \psi_1 \circ e$ . Then  $\psi$  is given by a pair  $(x_1, x_2)$  in  $M$  such that  $x_i e = x_i$  for  $i = 1, 2$  and  $a_1 x_1 + a_2 x_2 = e$ . Let  $\pi : M = R \oplus R \rightarrow R$  be the projection onto the first factor. Write  $E_2 = \pi(E)$  and  $B_1 = \pi(B)$ . As before,  $R = E_2 \oplus B_1$ . Let  $f$  in  $R = \text{End}(R_R)$  be the projection onto  $E_2$  with kernel  $B_1$ , so that  $E_2 = fR$ . Notice that  $1 - f \in I$ . Now we have that  $\pi \circ \psi$  provides an isomorphism from  $E_1$  onto  $E_2$ . Let  $y_1$  in  $R = \text{End}(R_R)$  be an element such that  $y_1 f = y_1 = e y_1$  and implements the inverse of the isomorphism  $(\pi \circ \psi)|_{E_1}$ . Observe that  $y_1 x_1 = e$  and  $x_1 y_1 = f$  so that  $x_1$  and  $y_1$  are quasi-invertible and quasi-inverses for each other (because  $1 - e \in J$  and  $1 - f \in I$  and  $I$  and  $J$  are orthogonal). Since  $(a_1 + a_2 x_2 y_1)x_1 = y_1 x_1 = e$  this implies that  $a_1 + a_2 x_2 y_1$  is quasi-invertible (see, for example, Theorem 2.3), showing that  $R$  is a  $QB$ -ring.

The proof of (i)  $\iff$  (iii) is dual to the one of (i)  $\iff$  (ii), cf. [31, Theorem 1.6]. Since we know that the notion of  $QB$ -ring is symmetric (Theorem 3.6), we are done.  $\square$

**8.6. Definitions.** For a unital ring  $R$  we denote by  $\mathcal{V}(R)$  the monoid of isomorphism classes of finitely generated, projective, right  $R$ -modules. (The standard notation for the category of such modules is  $\mathcal{FP}(R)$ .) The addition in  $\mathcal{V}(R)$  is direct sum of representatives, and  $\mathcal{V}(R)$  is ordered by  $x \leq y$  if  $M \oplus P \simeq N$  for some representatives  $[M] = x$  and  $[N] = y$ . The order unit is  $\mathbf{1} = [R_R]$ . An *order-ideal* in  $\mathcal{V}(R)$  is a submonoid  $S$  of  $\mathcal{V}(R)$  that is order-hereditary (i.e. if  $x \leq y$  and  $y \in S$ , then  $x \in S$ ). We say that two order-ideals  $S$  and  $T$  of  $\mathcal{V}(R)$  are *orthogonal* provided that  $S \cap T = 0$ . We denote by  $\mathcal{L}(R)$  the lattice of (two-sided) ideals of  $R$ , and by  $\mathcal{L}(\mathcal{V}(R))$  the lattice of order-ideals of  $\mathcal{V}(R)$ . If  $I$  is a (proper) two-sided ideal of  $R$ , we denote by  $\mathcal{FP}(I, R)$  the class of modules  $P$  in  $\mathcal{FP}(R)$  such that  $PI = P$ , and by  $\mathcal{V}(I)$  the set of isomorphism classes of elements from  $\mathcal{FP}(I, R)$ . It can be seen that  $\mathcal{V}(I)$  depends only on the structure of  $I$  as a ring without unit. (In fact, we can also describe  $\mathcal{V}(I)$  as the monoid of equivalence classes of idempotents in  $\mathbb{M}_\infty(I)$ .) By construction,  $\mathcal{V}(I)$  is an order-ideal of  $\mathcal{V}(R)$ . If  $I$  and  $J$  are ideals of  $R$ , then it is clear that  $\mathcal{V}(I \cap J) = \mathcal{V}(IJ) = \mathcal{V}(JI) = \mathcal{V}(I) \cap \mathcal{V}(J)$ . In particular, if  $I$  and  $J$  are orthogonal ideals of  $R$  we observe that  $\mathcal{V}(I)$  and  $\mathcal{V}(J)$  are orthogonal order-ideals of  $\mathcal{V}(R)$ .

If  $R$  is an exchange ring, then the map  $\phi : \mathcal{L}(R) \rightarrow \mathcal{L}(\mathcal{V}(R))$  given by  $I \rightarrow \mathcal{V}(I)$  is a surjective lattice homomorphism. Indeed, if  $S$  is an order-ideal of  $\mathcal{V}(R)$ , then if

we denote by  $I(S)$  the ideal of  $R$  generated by the set  $\{e = e^2 \mid [eR] \in S\}$ , we have that  $\phi(I(S)) = \mathcal{V}(I(S)) = S$ . This correspondence is an isomorphism if we restrict the domain to the lattice  $\mathcal{L}_{\mathcal{J}}(R)$  of *semi-primitive* ideals, i.e. those ideals  $I$  of  $R$  such that  $\mathcal{J}(R/I) = 0$ , which form a lattice with infima given by intersections and suprema by Jacobson radicals of sums (see [22]).

**8.7. Theorem.** *Let  $R$  be a unital ring, and consider the three conditions:*

- (i)  *$R$  is a QB-ring ;*
- (ii) *If  $n \cdot \mathbf{1} + b_1 = n \cdot \mathbf{1} + b_2$  in  $\mathcal{V}(R)$  for some  $n \geq 1$ , then we can find orthogonal order-ideals  $S_1$  and  $S_2$  in  $\mathcal{V}(R)$  and elements  $x, c_1, c_2$ , such that  $c_i \in S_i$  for  $i = 1, 2$ , and moreover  $b_1 + c_1 = b_2 + c_2$  and  $x + c_1 = n \cdot \mathbf{1} = x + c_2$  ;*
- (iii) *If  $a + b_1 = a + b_2$  in  $\mathcal{V}(R)$  then there exist orthogonal order-ideals  $S_1$  and  $S_2$  in  $\mathcal{V}(R)$  and elements  $c_1, c_2$ , such that  $c_i \in S_i$  for  $i = 1, 2$ , and  $b_1 + c_1 = b_2 + c_2$ .*

*We always have (i)  $\implies$  (ii)  $\implies$  (iii), but if  $R$  is a semi-primitive exchange ring all three conditions are equivalent.*

*Proof.* (i)  $\implies$  (ii). Choose representatives  $B_1$  and  $B_2$  for  $b_1$  and  $b_2$  such that  $A_1 \oplus B_1 = A_2 \oplus B_2$  for some  $A_i \simeq R_R^n$ . Applying Condition (ii) in Lemma 8.5 we find decompositions  $A_i = N \oplus C_i$  and  $B_1 \oplus C_1 \simeq B_2 \oplus C_2$  for some finitely generated projective modules  $N, C_1, C_2$ , such that  $C_i I_i = C_i$  for a pair of orthogonal ideals  $I_1, I_2$  in  $R$ . Let  $S_i = \mathcal{V}(I_i)$  and put  $x = [N]$  and  $c_i = [C_i]$ . (Note that  $S_1$  and  $S_2$  are orthogonal order-ideals of  $\mathcal{V}(R)$ .) Then  $b_1 + c_1 = b_2 + c_2$  and  $x + c_1 = n \cdot \mathbf{1} = x + c_2$ , and evidently  $c_i \in S_i$ .

(ii)  $\implies$  (iii). This follows from the fact that  $\mathbf{1}$  is an order-unit in  $\mathcal{V}(R)$ .

Assume now that  $R$  is a semi-primitive exchange ring satisfying (iii). Consider an arbitrary von Neumann regular element  $a$  in  $R$  with partial inverse  $b$ , and put  $p = ab$  and  $q = ba$ , so that  $aR = pR$  and  $Ra = Rq$ . Then

$$\begin{aligned} R \oplus (1 - p)R &= qR \oplus (1 - q)R \oplus (1 - p)R \\ &\simeq pR \oplus (1 - p)R \oplus (1 - q)R = R \oplus (1 - q)R. \end{aligned}$$

By (iii) there exist  $c_1, c_2$  in  $\mathcal{V}(R)$ , such that

$$[(1 - p)R] + c_1 = [(1 - q)R] + c_2.$$

Moreover,  $c_i \in S_i$ , where  $S_1$  and  $S_2$  are orthogonal order-ideals in  $\mathcal{V}(R)$ .

Since  $R$  is an exchange ring  $\mathcal{V}(R)$  is a refinement monoid, cf. [4, Corollary 1.3], so we can find decompositions

$$\begin{aligned} [(1 - p)R] &= m_{11} + m_{12}, & c_1 &= m_{21} + m_{22}, \\ [(1 - q)R] &= m_{11} + m_{21}, & c_2 &= m_{12} + m_{22}. \end{aligned}$$

Since  $S_1 \cap S_2 = 0$ , we have  $m_{22} = 0$ . Corresponding to this we have pairs of orthogonal idempotents  $e_1, f_1$  and  $e_2, f_2$  in  $R$ , such that  $1 - p = e_1 + f_1$ ,  $1 - q = e_2 + f_2$ ,  $e_1 R \simeq e_2 R$  and  $[f_1 R] = c_2$  and  $[f_2 R] = c_1$ . For  $i = 1, 2$  let  $I_i$  be the ideals of  $R$  generated by the idempotents  $e$  in  $R$  such that  $[eR] \in S_i$ , respectively. Then

$I_1 \cap I_2$  contains no non-zero idempotents, and so  $I_1 \cap I_2 \subset \mathcal{J}(R)$  because  $R$  is an exchange ring. Since  $R$  is semi-primitive  $I_1 \cap I_2 = 0$ .

Choose  $c$  and  $z$  in  $R$  such that  $e_1 = cz$  and  $e_2 = zc$ . Then  $c$  is a regular element in  $(1-p)R(1-q)$ , so  $a \leq a+c$  by Lemma 2.7. Moreover,

$$1 - (a+c)(b+z) = 1 - ab - cz = 1 - p - e_1 = f_1 \in I_2,$$

and similarly  $1 - (b+z)(a+c) = f_2 \in I_1$ . Since  $I_1 \cap I_2 = 0$  it follows that  $a+c \in R_q^{-1}$ .

We have shown that every regular element in  $R$  extends to an element in  $R_q^{-1}$ , whence  $R$  is a  $QB$ -ring by Theorem 8.4.  $\square$

**8.8. Examples.** Theorem 8.7 provides us with more interesting examples of  $QB$ -rings.

(A) Let  $\mathbb{F}$  be a field and define  $\mathbb{M}_\infty(\mathbb{F}) = \varinjlim \mathbb{M}_n(\mathbb{F})$  as in 6.9 and 7.16. This is a well-known example of a simple (non-unital)  $B$ -ring. Now let  $\mathbb{B}(\mathbb{F})$  denote the algebra of all row- and column-finite matrices over  $\mathbb{F}$ . This algebra contains  $\mathbb{M}_\infty(\mathbb{F})$  as its only non-trivial ideal. In fact, if  $V = \mathbb{F}^{(\omega)}$  denotes the countably infinite dimensional vector space over  $\mathbb{F}$ , and  $\mathbb{B}(\mathbb{F})$  is regarded as a subalgebra of  $\text{End}_{\mathbb{F}}(V)$ , then  $\mathbb{B}(\mathbb{F})$  is the idealizer of  $\mathbb{M}_\infty(\mathbb{F})$  in  $\text{End}_{\mathbb{F}}(V)$ .

It is known that  $\mathbb{B}(\mathbb{F})$  is not a regular ring, but O'Meara has proved to us that  $\mathbb{B}(\mathbb{F})$  is an exchange ring. (Private communication.)

Since idempotents in  $\mathbb{M}_\infty(\mathbb{F})$  are equivalent if and only if they have the same (finite) rank, and moreover all idempotents in  $\mathbb{B}(\mathbb{F}) \setminus \mathbb{M}_\infty(\mathbb{F})$  are equivalent to 1, we see that  $\mathcal{V}(\mathbb{B}(\mathbb{F})) = \{0, 1, \dots, \infty\}$ , where  $\infty = [1]$ . Then it is easy to check that Condition (ii) in Theorem 8.7 is satisfied, and hence  $\mathbb{B}(\mathbb{F})$  is a  $QB$ -ring.

(B) Replacing the field with a  $QB$ -ring  $A$ , whose structure is not too intricate, say  $A$  being a simple exchange ring with prescribed  $\mathcal{V}(A)$ , it seems safe to predict that  $\mathbb{B}(A)$  will often be a  $QB$ -ring. We already know from Corollary 6.10 that  $\mathbb{M}_\infty(A)$  is a  $QB$ -ring, and since this is an ideal in  $\mathbb{B}(A)$  we just have to control the quotient. However, the ideal structure of  $\mathbb{B}(A)/\mathbb{M}_\infty(A)$  can be quite complicated (see [3], [19], [25], [26]).

(C) Following Chen, [14], we say that a unital ring  $R$  satisfies *related comparability* if whenever we have two idempotents  $p$  and  $q$  such that  $1-p \sim 1-q$ , then there is a central idempotent  $z$  in  $R$ , such that  $zpR$  is isomorphic to a direct summand in  $zqR$  and  $(1-z)qR$  is isomorphic to a direct summand in  $(1-z)pR$ . It follows from [14, Theorem 2] that if  $R$  has related comparability, and if  $a$  is any von Neumann regular element in  $R$ , then we can find a partial inverse  $v$  for  $a$ , and a central idempotent  $z$  such that  $zv$  is left invertible and  $(1-z)v$  is right invertible in  $R$ . Evidently  $v \in R_q^{-1}$ . Thus, if  $R$  is a semi-primitive exchange ring with related comparability, then  $R$  is a  $QB$ -ring by Theorem 8.4.

(D) An exchange ring  $R$  is said to satisfy *general comparability* provided that for any modules  $A$  and  $B$  in  $\mathcal{FP}(R)$ , there exists a central idempotent  $e$  in  $R$  such that  $Ae$  is isomorphic to a direct summand of  $Be$  and  $B(1-e)$  is isomorphic to a direct summand of  $A(1-e)$ . Thus,  $R$  satisfies the fundamental comparison lemma for Murray-von Neumann equivalence of projections in a von Neumann algebra. Any exchange ring satisfying general comparability also satisfies related comparability.

Consequently, every regular, self-injective ring is a  $QB$ -ring. This class contains  $\text{End}(V_D)$ , where  $D$  is a division ring and  $V$  is a right  $D$ -vector space. More generally, it contains  $\text{End}(A_R)$ , where  $A$  is any non-singular, quasi-injective right module over a ring  $R$ , cf. [16, Corollary 1.23].

(E) We finally consider the rings constructed in [5, §3]. Let  $R(p, q)$  be the ring construction in Section 3 of [5] based on a simple unit-regular ring  $L$  and idempotents  $p$  and  $q$  in  $L$ . Then  $R(p, q)$  is a prime regular ring with a unique maximal ideal  $N(p, q)$ . It is proved in [5, Lemma 3.1] that  $R(p, q)$  has stable rank one if and only if  $p \sim q$ . By using the same techniques and the fact that  $R(p, q)_q^{-1} = R(p, q)_r^{-1} \cup R(p, q)_l^{-1}$  (because  $R(p, q)$  is a prime ring), one can show that  $R(p, q)$  is a  $QB$ -ring if and only if either  $p \lesssim q$  or  $q \lesssim p$ . In particular, Example 3.2 (in [5]) provides an example of a stably finite, regular ring  $U$  which is not a  $QB$ -ring, but satisfies 2-comparability (see [16, p. 275]). In fact,  $U$  satisfies the stronger property of almost comparability. This contrasts heavily with the situation of regular rings satisfying the comparability axiom (in the sense of [16]), which are all  $QB$ -rings since they satisfy general comparability.

**8.9. Example.** Let  $S$  be the ring constructed in Example 7.16. Recall that  $S$  is an extension of the two  $B$ -rings  $\mathbb{M}_\infty(\mathbb{F})$  and  $\mathbb{F}(t)$ , the field of rational functions on a countable field  $\mathbb{F}$ . Denote by  $\rho : S \rightarrow \mathbb{F}(t)$  the quotient map. Selecting appropriate subrings of  $\mathbb{F}(t)$  with stable rank one it is possible to produce other examples of  $QB$ -rings which are neither  $B$ -rings nor exchange rings. For example, consider the subring  $\mathbb{F}(t)_{1,2}$  of  $\mathbb{F}(t)$  of those rational functions  $f(t)/g(t)$  such that  $g(1)$  is non-zero and  $g(2)$  is non-zero (assuming that  $2 \neq 0$  in  $\mathbb{F}$ ). The subring  $S_{1,2}$  of  $S$  which is the inverse image of  $\mathbb{F}(t)_{1,2}$  through  $\rho$  will be a  $QB$ -ring, by the same argument as in Example 7.16. Since  $\mathbb{F}(t)_{1,2}$  is semilocal but not local we get that  $S_{1,2}$  is not an exchange ring. Again  $S_{1,2}$  is an extension of two  $B$ -rings,  $\mathbb{M}_\infty(\mathbb{F})$  and  $\mathbb{F}(t)_{1,2}$ , and  $\text{bsr}(S_{1,2}) = 2$ . We can think of  $S$  as a nice localization of  $S_{1,2}$ .

## 9. Extremally Rich $C^*$ -Algebras

Let  $A$  be a  $C^*$ -algebra, i.e. an algebra of bounded operators on a complex Hilbert space, which is closed under taking adjoints and under norm convergence. Then  $A$  is a semi-primitive ring, and the  $*$ -operation makes it particularly simple to link the left and the right structure of  $A$ .

The concept of quasi-invertibility for elements in a unital  $C^*$ -algebra was explored in [8, Theorem 1.1], and it was shown that  $A_q^{-1}$  is the open subset of  $A$  consisting of elements of the form  $A^{-1}\mathcal{E}(A)A^{-1}$ . Here  $\mathcal{E}(A)$  denotes the set of extreme points in the unit ball of  $A$ , identified by Kadison as the partial isometries  $v$  in  $A$  such that

$$(1 - vv^*) \perp (1 - v^*v),$$

cf. [23, 1.4.7].

In [8, §3] a unital  $C^*$ -algebra  $A$  was defined to be *extremally rich* if the set of quasi-invertible elements in  $A$  was dense in  $A$ , i.e. if  $(A_q^{-1})^\circ = A$ . This seemed a

most appropriate generalization of Rieffel's notion of  $C^*$ -algebras with topological stable rank one, since by results of Rieffel [28], and of Herman and Vaserstein [18] a  $C^*$ -algebra has Bass stable rank one if and only if  $(A^{-1})^= = A$ , i.e.  $A$  has topological stable rank one.

To show that our notion of  $QB$ -ring reduces to extremal richness in the case of  $C^*$ -algebras we offer the following result, of which the second half essentially is a modification of the argument from [18].

**9.1. Proposition.** *For an element  $x$  in a unital  $C^*$ -algebra  $A$  the following conditions are equivalent:*

- (i)  $x \in (A_q^{-1})^=$  ;
- (ii)  $x \in \text{cl}(A_q^{-1})$  .

*Proof.* (i)  $\implies$  (ii) If  $ax + b = 1$  for some  $x, b$  in  $A$ , then

$$1 = (ax + b)^*(ax + b) \leq 2x^*a^*ax + 2b^*b \leq 2\|a\|^2x^*x + 2b^*b .$$

Thus,  $x^*x + b^*b \in A^{-1}$ . By Theorem 3.3 in [8] there is an extreme point  $u$  in  $\mathcal{E}(A)$  such that  $x + ub \in A_q^{-1}$ . In particular,  $x \in \text{cl}(A_q^{-1})$ .

(ii)  $\implies$  (i) If  $x \in \text{cl}(A_q^{-1})$  and  $\varepsilon > 0$  is given, define  $b = (1 - \varepsilon^{-2}x^*x)_+$ , the positive part of the element  $1 - \varepsilon^{-2}x^*x$ . Then by spectral theory

$$x^*x + b = x^*x + (1 - \varepsilon^{-2}x^*x)_+ \geq \varepsilon^2 1 ,$$

since  $t + (1 - \varepsilon^{-2}t)_+ \geq \varepsilon^2$  for every real  $t \geq 0$ . Consequently  $x^*x + b \in A^{-1}$ . By assumption we therefore have  $x + yb \in A_q^{-1}$  for some  $y$  in  $A$ . Take a natural number  $n$  and define

$$z = (x + yb)(1 + nb)^{-1} \quad \text{in } A_q^{-1} .$$

Then with  $|x| = (x^*x)^{\frac{1}{2}}$  we have

$$\begin{aligned} \|x - z\| &\leq \|x(1 - (1 + nb)^{-1})\| + \|y\|\|b(1 + nb)^{-1}\| \\ &= \| |x| nb(1 + nb)^{-1} \| + \frac{1}{n} \|y\|\|nb(1 + nb)^{-1}\| . \end{aligned}$$

Regarding  $b$  as a function of  $|x|$  it vanishes on  $\text{sp}(|x|)$  if  $t \geq \varepsilon$ , so  $\||x|b\| \leq \varepsilon$ . Moreover,  $\|nb(1 + nb)^{-1}\| \leq 1$  by spectral theory, whence

$$\|x - z\| \leq \varepsilon + \frac{1}{n} \|y\| < 2\varepsilon$$

for  $n$  large enough. Thus,  $x \in (A_q^{-1})^=$ . □

**9.2. Corollary.** *In a  $C^*$ -algebra  $A$  we always have*

$$\text{cl}(A_q^{-1}) = \text{cr}(A_q^{-1}) \quad (= (A_q^{-1})^=) . \quad \square$$

A (unital)  $C^*$ -algebra  $A$  has *real rank zero* provided that every self-adjoint element can be approximated arbitrarily well by self-adjoint, invertible elements.

This concept was introduced and explored in [7], where a number of equivalent characterizations were also provided.

To appreciate the following result, which was originally proved by L.G. Brown and the second author by a direct argument [unpublished], note that if  $v$  and  $w$  are partial isometries such that  $w$  extends  $v$  in the sense of 2.4, then we actually have  $v = vv^*w = ww^*v$ , which are the relations usually employed to describe the relation  $v \leq w$ . To see this, assume that  $v = vxv = vxw = wxv$  for some  $x$ . Then  $v^*v \leq w^*w$ , and therefore

$$\begin{aligned} v^*v &= v^*x^*w^*wxv = v^*x^*(v^*v + (1 - v^*v))w^*w((1 - v^*v) + v^*v)xv \\ &= (v^*v + v^*x^*(1 - v^*v))w^*w((1 - v^*v)xv + v^*v) \\ &= v^*v + v^*x^*(1 - v^*v)w^*w(1 - v^*v)xv. \end{aligned}$$

It follows that  $w(1 - v^*v)xv = 0$ , i.e.  $v = wxv = ww^*vxv = ww^*v$ .

**9.3. Proposition.** *A unital  $C^*$ -algebra  $A$  of real rank zero is extremally rich if and only if every partial isometry in  $A$  extends to an extreme partial isometry.*

*Proof.* The necessity of the condition is proved in [8, Proposition 2.6], cf. [8, Corollary 4.3], so it only remains to show sufficiency.

Towards this end let  $a$  be a von Neumann regular element in  $A$ . Then  $a$  has a polar decomposition  $a = v|a|$ , and the spectrum of  $|a|$  has a gap  $]0, \varepsilon[$  for some  $\varepsilon > 0$ . Therefore  $e = 1 - v^*v + |a|$  is an invertible, positive element in  $A$ , and  $x = e^{-1}v^*$  is a partial inverse for  $a$ , since  $axa = v|a|e^{-1}v^*v|a| = vv^*v|a| = v|a| = a$ .

By assumption  $v$  extends to an extreme partial isometry  $u$  in  $A$ , so that  $(1 - uu^*)A(1 - u^*u) = 0$ , and  $v = uv^*v = vv^*u$ . Set  $b = ue$  and  $y = e^{-1}u^*$ . Then  $1 - by = 1 - uu^*$  and  $1 - yb = e^{-1}(1 - u^*u)e$ , so  $b \in A_q^{-1}$  with quasi-inverse  $y$ . Moreover,

$$\begin{aligned} axb &= v|a|e^{-1}v^*ue = vv^*ue = ve = v|a| = a, \\ bxa &= uee^{-1}v^*v|a| = uv^*v|a| = v|a| = a, \end{aligned}$$

so  $a \leq b$ . Since  $a$  was arbitrary, it follows from Theorem 8.4 and [4, Theorem 7.2] that  $A$  is a  $QB$ -ring, and therefore extremally rich by Proposition 9.1.  $\square$

**9.4. Remarks.** Using examples from the theory of extremally rich  $C^*$ -algebras we can exhibit several phenomena in the theory of  $QB$ -rings. Thus, it is possible to have a unital ring  $R$ , with an idempotent  $p$ , such that both  $pRp$  and  $(1 - p)R(1 - p)$  are  $QB$ -rings, but  $R$  is not, cf. [8, Example 6.12].

In the example above the ideals  $I_1$  and  $I_2$  of  $R$ , generated by  $p$  and  $1 - p$ , respectively, will still be  $QB$ -rings, and of course  $I_1 + I_2 = R$ . Since these are complex algebras we also have that  $\tilde{I}_1 = I_1 + \mathbb{C}1$  is a  $QB$ -ring. This shows that we can not in Corollary 7.15 relax the condition on the subring from being a  $B$ -ring to being a  $QB$ -ring.

In [8, Example 1.3] we find a  $C^*$ -algebra  $A$  which is not a  $QB$ -ring although it is a direct limit of algebras isomorphic to  $\mathbb{B}(\mathcal{H})$ . However, the embedding of one copy of  $\mathbb{B}(\mathcal{H})$  into the next is not prime. This shows that our condition for prime embeddings in Proposition 5.7 is necessary.

The algebra  $A$  contains an element  $a$  such that  $\pi(a)$  is left or right invertible in  $\pi(A)$  for each primitive (i.e. irreducible) representation  $(\pi, \mathcal{H})$  of  $A$ . However,  $a \notin \text{cl}(A_q^{-1})$  by [8, Corollary 1.10]. We can therefore not in general hope to simplify the definition of quasi-invertibility to just a question of one-sided invertibility in prime or primitive quotients.

## REFERENCES

- [1] Frank W. Anderson & Kent R. Fuller, *Rings and categories of modules*, Graduate Texts in Mathematics, **13**, Springer-Verlag, Berlin, 1974.
- [2] Pere Ara, *Extensions of exchange rings*, Journal of Algebra **197** (1997), 409–423.
- [3] Pere Ara & Francesc Perera, *Multipliers of von Neumann regular rings*, Communications in Algebra, to appear.
- [4] Pere Ara, Ken R. Goodearl, Kevin C. O’Meara & Enric Pardo, *Separative cancellation for projective modules over exchange rings*, Israel Journal of Mathematics **105** (1998), 105–137.
- [5] Pere Ara, Kevin C. O’Meara & Dmitry V. Tyukavkin, *Cancellation of projective modules over regular rings with comparability*, Journal of Pure and Applied Algebra **107** (1996), 19–38.
- [6] Hyman Bass, *Introduction to some Methods of Algebraic K-Theory*, Regional Conference Series in Mathematics, **21**, American mathematical Society, Providence R.I. 1974.
- [7] Lawrence G. Brown & Gert K. Pedersen,  *$C^*$ -algebras of real rank zero*, Journal of Functional Analysis **99** (1991), 131–149.
- [8] Lawrence G. Brown & Gert K. Pedersen, *On the geometry of the unit ball of a  $C^*$ -algebra*, Journal für die reine und angewandte Mathematik **469** (1995), 113–147.
- [9] Lawrence G. Brown & Gert K. Pedersen, *Approximation and convex decomposition by extremals in  $C^*$ -algebras*, Mathematica Scandinavica **81** (1997), 69–85.
- [10] Lawrence G. Brown & Gert K. Pedersen, *Extremal K-theory and index for  $C^*$ -algebras*, K-Theory, to appear.
- [11] Lawrence G. Brown & Gert K. Pedersen, *Extremally rich ideals in  $C^*$ -algebras*, Preprint 1999.
- [12] Lawrence G. Brown & Gert K. Pedersen, *Non-stable K-theory and extremally rich  $C^*$ -algebras*, In preparation.
- [13] Huanyin Chen, *Comparability of modules over regular rings*, Communications in Algebra **25** (1997), 3531–3543.
- [14] Huanyin Chen, *Exchange rings, related comparability and power-substitution*, Communications in Algebra **26** (10) (1998), 3383–3401.
- [15] Chen-Lian Chuang and Pjek-Hwee Lee, *On regular subdirect products of simple artinian rings*, Pacific Journal of Mathematics **142** (1990), 17–21.
- [16] Ken R. Goodearl, *Von Neumann Regular Rings*, Pitman, London, 1979, second edition Krieger, Malabar, FL, 1991.
- [17] Ken R. Goodearl & Robert B. Warfield, Jr, *Algebras over zero-dimensional rings*, Mathematische Annalen **223** (1976), 157–168.
- [18] Richard H. Herman & Leonid N. Vaserstein, *The stable range of  $C^*$ -algebras*, Inventiones Mathematicæ **77** (1984), 553–555.
- [19] Nadia S. Larsen & Hiroyuki Osaka, *Extremal richness of multiplier algebras and corona algebras of simple  $C^*$ -algebras*, Journal of Operator Theory **38** (1997), 131–149.
- [20] Pere Menal & Jaume Moncasi, *On regular rings with stable range 2*, Journal of Pure and Applied Algebra **24** (1982), 25–40.
- [21] W. Keith Nicholson, *Lifting idempotents and exchange rings*, Transactions of the American Mathematical Society **229** (1977), 269–278.
- [22] Enric Pardo, *Monoides de refinament i anells d’intercanvi*, Ph.D. Thesis, Universitat Autònoma de Barcelona, 1995.
- [23] Gert K. Pedersen,  *$C^*$ -Algebras and their Automorphism Groups*, LMS Monographs **14**, Academic Press, London, 1979.



- [24] Gert K. Pedersen, *The  $\lambda$ -function in operator algebras*, Journal of Operator Theory **26** (1991), 345–381.
- [25] Francesc Perera, *Ideal structure of multiplier algebras of simple  $C^*$ -algebras with real rank zero*, Preprint.
- [26] Francesc Perera, *Extremal richness of multiplier and corona algebras of simple  $C^*$ -algebras with real rank zero*, Journal of Operator Theory, to appear.
- [27] Francesc Perera, *Lifting units modulo exchange ideals and  $C^*$ -algebras with real rank zero*, Journal für die reine und angewandte Mathematik, to appear.
- [28] Marc A. Rieffel, *Dimension and stable rank in the  $K$ -theory of  $C^*$ -algebras*, Proceedings of the London Mathematical Society **46** (1983), 301–333.
- [29] Leonid N. Vaserstein, *Bass' first stable range condition*, Journal of Pure and Applied Algebra **34** (1984), 319–330.
- [30] Robert B. Warfield, Jr, *Exchange rings and decompositions of modules*, Mathematische Annalen **199** (1972), 31–36.
- [31] Robert B. Warfield, Jr, *Cancellation of modules and groups and stable rank of endomorphism rings*, Pacific Journal of Mathematics **91** (1980), 457–485.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193, BEL-LATERRA (BARCELONA), SPAIN

*E-mail address:* para@mat.uab.es, perera@mat.uab.es

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARKEN 5, DK-2100 COPENHAGEN Ø, DENMARK

*E-mail address:* gkped@math.ku.dk, perera@math.ku.dk