HARMONIC MORPHISMS AND THE JACOBI OPERATOR

STEFANO MONTALDO AND JOHN C. WOOD

ABSTRACT. We prove that harmonic morphisms preserve the Jacobi operator along harmonic maps. We apply this result to prove infinitesimal and local rigidity (in the sense of Toth) of harmonic morphisms to a sphere.

1. HARMONIC MORPHISMS

Harmonics maps $\phi : (M, g) \to (N, h)$ between two smooth Riemannian manifolds are critical points of the energy functional $E(\phi, \Omega) = \frac{1}{2} \int_{\Omega} |d\phi|^2 dv_g$ for any compact domain $\Omega \subseteq M$ [4], i.e. the first variation of the energy vanishes for any smooth variation of ϕ . The Euler-Lagrange equation for the energy is the vanishing of the tension field $\tau_{\phi} = \text{trace } \nabla d\phi$, where ∇ denotes the connection on $T^*M \otimes \phi^{-1}TN$ induced from the Levi-Civita connections ∇^M on M and ∇^N on N. If $\{e_i\}_{i=1}^m$ is a local orthonormal frame on M we have $\tau_{\phi} = \sum_{i=1}^m \{\nabla_{e_i}^{\phi}(d\phi(e_i)) - d\phi(\nabla_{e_i}^M e_i)\}$ where ∇^{ϕ} denotes the pull-back connection on $\phi^{-1}TN$.

Let $\phi : (M,g) \to (N,h)$ be a smooth map between two Riemannian manifolds. The tangent space at a point $x \in M$ can be decomposed as $T_x M = H_x \oplus V_x$ where $V_x = \ker(d\phi_x)$ and $H_x = V_x^{\perp}$. The spaces V_x and H_x are called the *vertical* and *horizontal space* at the point $x \in M$, respectively.

Date: October 1999.

¹⁹⁹¹ Mathematics Subject Classification. 58E20.

Key words and phrases. Harmonic maps, harmonic morphisms, Jacobi operator, rigidity.

Let $C_{\phi} = \{x \in M \mid \operatorname{rank}(d\phi_x) \text{ is not maximal}\}$; points of C_{ϕ} are called *critical* points of ϕ .

Definition 1.1. A map $\phi : (M, g) \to (N, h)$ is called *horizontally (weakly)* conformal if, for every $x \in M$, either $d\phi_x|_{H_x}$ is conformal and surjective or $d\phi_x = 0$.

In particular, either $x \in M \setminus C_{\phi}$ and ϕ is submersive at x or $x \in C_{\phi}$ and the differential $d\phi_x$ has rank 0. If $\phi: M \to N$ is horizontally (weakly) conformal, then there exists a function $\lambda: M \setminus C_{\phi} \to \mathbb{R}^+$ such that $h_{\phi(x)}(d\phi_x(X), d\phi_x(Y))$ - $= \lambda^2 g_x(X, Y)$, for all $X, Y \in H_x$, and $x \in M \setminus C_{\phi}$. The function λ can be extended continuously to the whole of M by setting $\lambda|_{C_{\phi}} = 0$; the extended function is called the *dilation function* of ϕ . Note that λ^2 is smooth.

Harmonic morphisms between Riemannian manifolds are defined to be maps which pull back harmonic functions to harmonic functions; they are characterized as the harmonic maps which are horizontally (weakly) conformal [6, 10]. Note that non-constant harmonic morphisms can only exist if dim $M \ge \dim N$. For a bibliography of papers on harmonic morphisms and an 'atlas' of the known examples, see [7]; for the general theory, see [2].

2. Second variation and the Jacobi operator

Let $\phi : M \to N$ be a harmonic map, for simplicity we assume that M is compact. Then the first variation $D_V E(\phi) = 0$ for all vector fields along ϕ , where by a vector field along ϕ we mean a smooth section $V \in \Gamma(\phi^{-1}TN)$ of $\phi^{-1}TN$. Given a vector field along ϕ we consider a smooth one-parameter variation ϕ_t ($-\epsilon < t < \epsilon$) of $\phi = \phi_0$ such that $V = \frac{d}{dt}(\phi_t)|_{t=0}$. Let $\nabla^2 = \nabla \circ \nabla^{\phi}$ so that

$$\nabla_{X,Y}^2 W = \nabla_X^{\phi}(\nabla_Y^{\phi} W) - \nabla_{\nabla_X^M Y}^{\phi} W \qquad \left(X, Y \in TM, \ W \in \Gamma(\phi^{-1}TN)\right);$$

note that this is tensorial in X and Y but not, in general, symmetric. We have the following second variation formula for the energy [8, 12]; we use Milnor's convention for the curvature as in [5].

$$\begin{aligned} H_{\phi}(V,V) &= \left. \frac{d^2 E(\phi_t)}{dt^2} \right|_{t=0} &= \int_M \langle \Delta^{\phi} V - \operatorname{trace} R^N(d\phi,V) d\phi,V \rangle \, v_g \\ &= \left. \int_M \langle J^{\phi} V,V \rangle \, v_g \right. \end{aligned}$$

Here Δ^{ϕ} is the Laplacian on sections of $\phi^{-1}TN$ given in a local orthonormal frame $\{e_i\}$ on M by

$$\Delta^{\phi} = -\operatorname{trace} \nabla^2 = -\sum_{i=1}^m \nabla^2_{e_i, e_i} = -\sum_{i=1}^m \left\{ \nabla^{\phi}_{e_i} \nabla^{\phi}_{e_i} - \nabla^{\phi}_{\nabla_{e_i} e_i} \right\}$$

The operator $J^{\phi} = \Delta^{\phi} - \operatorname{trace} R^{N}(d\phi, \cdot)d\phi$ is called the *Jacobi* operator of ϕ ; the elements of its kernel are called *Jacobi fields* and form a subspace $J(\phi) \subset \Gamma(\phi^{-1}TN)$.

Definition 2.1. The *index* of ϕ is the dimension of the largest subspace of $\Gamma(\phi^{-1}TN)$ on which H_{ϕ} is negative definite. The *nullity* of ϕ is the dimension of the kernel of J^{ϕ} . Alternatively, the index of ϕ is the sum of the multiplicities of the negative eigenvalues λ of J^{ϕ} and the nullity of ϕ is the multiplicity of the eigenvalue 0 of J^{ϕ} .

Note that the Jacobi operator is a linear elliptic self-adjoint operator with positive principal part Δ^{ϕ} . It follows from standard elliptic theory (cf. [8]) that the index and the nullity are both finite.

Our main result is that harmonic morphisms preserve the Jacobi operator along harmonic maps as follows:

Theorem 2.2. Let $\phi : M \to N$ be a harmonic morphism, $\psi : N \to P$ a harmonic map and V a vector field along ψ . Then the Jacobi operator for the

vector field $V \circ \phi$ along $\psi \circ \phi$ is given by

$$J^{\psi \circ \phi}(V \circ \phi) = \lambda^2 J^{\psi}(V) \circ \phi \,,$$

where λ is the dilation function of ϕ . In particular, if V is a Jacobi field along ψ , then $V \circ \phi$ is a Jacobi field along $\psi \circ \phi$.

Proof. We first remark that the composition $\psi \circ \phi$ is harmonic. Set $W = V \circ \phi \in \Gamma(\phi^{-1}\psi^{-1}TP)$. By definition of pull-back connection, for any $X \in \Gamma(TM)$, we have

(2.1)
$$\nabla_X^{\psi \circ \phi} W = \left\{ \nabla_{d\phi(X)}^{\psi} V \right\} \circ \phi.$$

Let γ be a smooth curve in M tangent to X and extend X to a vector field along γ . Then, by (2.1), both sides of the next equation are well-defined and we have

$$\nabla_X^{\psi \circ \phi}(\nabla_X^{\psi \circ \phi}W) = \left\{ \nabla_{d\phi(X)}^{\psi} \left(\nabla_{d\phi(X)}^{\psi}V \right) \right\} \circ \phi \,.$$

Again, by (2.1),

$$\nabla_{\nabla_X^M X}^{\psi \circ \phi} W = \left\{ \nabla_{d\phi(\nabla_X^M X)}^{\psi} V \right\} \circ \phi$$

so that

$$\begin{aligned} \nabla_{X,X}^2 W &= \nabla_X^{\psi \circ \phi} \left(\nabla_X^{\psi \circ \phi} W \right) - \nabla_{\nabla_X X}^{\psi \circ \phi} W \\ &= \left\{ \nabla_{d\phi(X)}^{\psi} \left(\nabla_{d\phi(X)}^{\psi} V \right) - \nabla_{d\phi(\nabla_X^M X)}^{\psi} V \right\} \circ \phi \,. \end{aligned}$$

Since ϕ is harmonic, for any orthonormal frame $\{e_i\}$ on M we have

$$\sum_{i=1}^{m} d\phi \left(\nabla_{e_i}^M e_i \right) = \sum_{i=1}^{m} \nabla_{d\phi(e_i)}^N d\phi(e_i)$$

so that

$$\begin{aligned} \operatorname{trace} \nabla^2 W &= \sum_{i=1}^m \left\{ \nabla^{\psi}_{d\phi(e_i)} \nabla^{\psi}_{d\phi(e_i)} V - \nabla^{\psi}_{\nabla^N_{d\phi(e_i)} d\phi(e_i)} V \right\} \circ \phi \\ &= \sum_{i=1}^m \left\{ \nabla^2_{d\phi(e_i), d\phi(e_i)} V \right\} \circ \phi \,. \end{aligned}$$

Now choosing the bases $\{e_i\}$, $\{f_i\}$ at the points x, $\phi(x)$ such that $d\phi(e_i) = \lambda f_i$ for i = 1, ..., n and $d\phi(e_i) = 0$ otherwise, we have

trace
$$\nabla^2 W = \sum_{i=1}^n \nabla^2_{d\phi(e_i), d\phi(e_i)} V \circ \phi$$

= $\lambda^2 \operatorname{trace} \nabla^2 V \circ \phi$.

Next, an easy calculation gives

trace
$$R^P(d(\psi \circ \phi), W)d(\psi \circ \phi) = \lambda^2$$
 trace $R^P(d\psi, V)d\psi$

so that

$$J^{\psi \circ \phi}(W) = J^{\psi \circ \phi}(V \circ \phi) = \lambda^2 \operatorname{trace} \left\{ -\nabla^2 V - R^P(d\psi, V) d\psi \right\} \circ \phi$$
$$= \lambda^2 J^{\psi}(V) \circ \phi \,.$$

Remark 2.3. A version of Theorem 2.2 with the assumption that the fibres of ϕ are minimal was proved by the first author in [9].

Corollary 2.4. Let $\phi : M \to N$ be a non-constant harmonic morphism between compact manifolds and let $\psi : N \to P$ be a harmonic map. Then

- (a) (i) $\operatorname{index}(\psi \circ \phi) \ge \operatorname{index}(\psi);$
 - (ii) $index(\phi) \ge index(Id^N)$, in particular, if Id^N is unstable, then so is ϕ ;

- (b) (iii) nullity $(\psi \circ \phi) \ge$ nullity (ψ) ;
 - (iv) nullity(ϕ) \geq nullity(Id^N).

Proof. (a) For part (i), let $V \in \Gamma(\psi^{-1}TP)$ be an eigenvector with negative eigenvalue, thus

(2.2)
$$J^{\psi}(V) = \alpha V$$

for some constant $\alpha < 0$. Set $W = V \circ \phi \in \Gamma((\psi \circ \phi)^{-1}TP)$. Then, by Theorem 2.2,

$$J^{\psi \circ \phi}(W) = \lambda^2 J^{\psi}(V) \circ \phi$$
$$= \lambda^2 \alpha \left(V \circ \phi \right) = \lambda^2 \alpha W$$

Now W cannot be identically zero; if it were, V would be zero on $\phi(M)$; this is an open set since any non-constant harmonic morphism is an open mapping [6]. But then, by the unique continuation theorem [1, 3] applied to (2.2), V would be identically zero on N. Further, by unique continuation for harmonic morphisms [6], λ cannot be zero on an open set. Hence

$$H_{\phi}(W,W) = \int_{M} \lambda^2 \alpha \langle W,W \rangle < 0.$$

Now let V_1, \ldots, V_s be linearly independent eigenvectors with negative eigenvalues. Then, again by unique continuation, $W_1 = V_1 \circ \phi, \ldots, W_s = V_s \circ \phi$ are linearly independent. Further, if V_i and V_j correspond to different eigenvalues α_i and α_j ,

$$\alpha_i \int_M \lambda^2 \langle W_i, W_j \rangle = \int_M \langle J^{\psi \circ \phi} W_i, W_j \rangle = \int_M \langle W_i, J^{\psi \circ \phi} W_j \rangle = \alpha_j \int_M \lambda^2 \langle W_i, W_j \rangle$$

so that $H_{\psi \circ \phi}(W_i, W_j) = 0$. It follows that H_{ϕ} is negative definite on the span of the W_i and the estimate on the index follows. Part (ii) follows by putting $\psi = \mathrm{Id}^N$.

 $\mathbf{6}$

(b) Let V be a Jacobi vector field along ψ . Then, by Theorem 2.2, so is $W = V \circ \phi$ and we argue as in part (a).

3. Toth rigidity of harmonic morphisms

Let $\phi: M \to N$ be a harmonic map. A section $V \in \Gamma(\phi^{-1}TN)$ is said to be a harmonic variation of ϕ if $\phi_t = \exp(tV) : M \to N$ is harmonic for all $t \in \mathbb{R}$. Let $H(\phi) \subset \Gamma(\phi^{-1}TN)$ denote the set of all harmonic variations of a given map $\phi: M \to N$.

Note that, if ϕ_t is a variation of ϕ through harmonic maps, then $V = \frac{\partial \phi_t}{\partial t}|_{t=0}$ is a Jacobi field along ϕ [5]. The converse is not always true, in fact there are examples of Jacobi fields along a harmonic map which do not arise as variations through harmonic maps (see, e.g., [11]). Nevertheless we have the following.

Theorem 3.1 ([13]). Let $\phi : (M, g) \to N = N(c)$ be a harmonic map from a compact Riemannian manifold to a real space form of constant curvature $c \neq 0$. Then a section $V \in \Gamma(\phi^{-1}TN)$ is a harmonic variation if and only if V is a Jacobi field with ||V|| constant and trace $\langle d\phi, \nabla^{\phi}V \rangle = 0$.

Let $\phi: M \to \mathbb{S}^n$ be a harmonic map to the Euclidean sphere of dimension $n \ (n \in \{1, 2, ...\})$ and set

$$K(\phi) = \{ V \in J(\phi) : \operatorname{trace} \langle d\phi, \nabla^{\phi} V \rangle = 0 \}.$$

Then $K(\phi) \subset J(\phi)$ is a linear subspace and, by Theorem 3.1,

 $H(\phi) = \{ V \in K(\phi) : \|V\| \text{ is constant} \}.$

Recall that a variation $V \in \Gamma(\phi^{-1}TN)$ is called *projectable* if $\phi(x) = \phi(x')$ implies that V(x) = V(x').

Definition 3.2 ([13]). A harmonic map $\phi : M \to \mathbb{S}^n$ is said to be

- 1. infinitesimally rigid if, for every projectable $V \in K(\phi)$, there exists $X \in so(n+1)$ such that the equation $V = X \circ \phi$ holds;
- 2. locally rigid if, for every projectable harmonic variation V, there exists a 1-parameter subgroup $(g_t) \subset O(n+1)$ such that $\phi_t = \exp(tV) = g_t \circ \phi$ for all $t \in \mathbb{R}$.

Lemma 3.3 ([13]). Let $\mathrm{Id}^{\mathbb{S}^n} : \mathbb{S}^n \to \mathbb{S}^n$ be the identity map. Then

$$K(\mathrm{Id}^{\mathbb{S}^n}) = so(n+1).$$

We extend a theorem of Toth [13] on rigidity of harmonic Riemannian submersions to submersive harmonic morphisms.

Theorem 3.4. (i) Any surjective and submersive harmonic morphism ϕ : $M \to \mathbb{S}^n$ is infinitesimally rigid.

(ii) Let n be odd. Then any surjective and submersive harmonic morphism $\phi: M \to \mathbb{S}^n$ is locally rigid.

Proof. (i) Let $V \in K(\phi)$ be projectable. Then since ϕ is surjective and submersive we have $V = X \circ \phi$ for some $X \in \Gamma(T\mathbb{S}^n)$ and so, from Theorem 2.2,

(3.1)
$$J^{\phi}(V) = \lambda^2 J^{\mathbb{S}^n}(X) \circ \phi,$$

where $J^{\mathbb{S}^n}$ denotes the Jacobi operator along $\mathrm{Id}^{\mathbb{S}^n}$. Moreover, choosing orthonormal bases $\{e_i\}$ at a point $x \in M$ and $\{f_i\}$ at $\phi(x) \in N$ such that $d\phi(e_i) = \lambda f_i \circ \phi$ for $i = 1, \ldots, n$ and $d\phi(e_i) = 0$ otherwise, we have

(3.2)
$$0 = \operatorname{trace} \langle d\phi, \nabla^{\phi} V \rangle = \sum_{i=1}^{m} \langle d\phi(e_i), \nabla^{\phi}_{e_i}(X \circ \phi) \rangle$$
$$= \lambda^2 \sum_{i=1}^{n} \langle f_i, \nabla_{f_i} X \rangle \circ \phi$$
$$= \lambda^2 \operatorname{trace} \langle d(\operatorname{Id}^{\mathbb{S}^n}), \nabla X \rangle \circ \phi$$

where ∇ denoted the Levi-Civita connection on S^n . Equations (3.1) and (3.2) imply that $X \in K(\mathrm{Id}^{\mathbb{S}^n})$ and by Lemma 3.3, $X \in so(n+1)$. This proves that ϕ is infinitesimally rigid.

(ii) Suppose that $V \in H(\phi)$ so that ||V|| is constant. As in part (i), $V = X \circ \phi$ for some $X \in \Gamma(TS^n)$. Since ϕ is surjective, ||X|| is also constant and, since, as in the first part of the proof, X is Killing, for any vector field Y on \mathbb{S}^n we have

$$\langle Y, \nabla_X X \rangle = -\langle X, \nabla_Y X \rangle = -\frac{1}{2} Y ||X||^2 = 0$$

It follows that $\nabla_X X = 0$ on \mathbb{S}^n which implies that every integral curve $t \to g_t(x)$ $(t \in \mathbb{R}, x \in \mathbb{S}^n)$ of X is a geodesic. Hence

$$\phi_t(x) = \exp(tV_x) = \exp(tX_{\phi(x)}) = g_t(\phi(x))$$

for all $x \in M$, and, since $X \in so(n+1)$, we have $(g_t) \subset O(n+1)$.

We remark that part (ii) is true for n even but says nothing new as there are no non-trivial projectable harmonic variations $V \in K(\phi)$ in that case.

References

- N. ARONSZAJN, 'A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order', J. Math. Pures Appl. (9) 36 (1957) 235–249.
- [2] P. BAIRD and J.C. WOOD, Harmonic morphisms between Riemannian manifolds, London Math. Soc. Monogr. (N.S.) (Oxford Univ. Press) (to appear).
- [3] H.O. CORDES, 'Über die eindeutige Bestimmtheit der Lösungen elliptischer Differentialgleichungen durch Anfangsvorgaben', Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II 1956 (1956) 239–258.

- [4] J. EELLS and J.H. SAMPSON, 'Harmonic mappings of Riemannian manifolds', Amer. J. Math. 86 (1964) 109–160.
- [5] J. EELLS and L. LEMAIRE, 'A report on harmonic maps', Bull. London Math. Soc. 10 (1978) 1–68.
- [6] B. FUGLEDE, 'Harmonic morphisms between Riemannian manifolds', Ann. Inst. Fourier (Grenoble) 28 (1978) 107–144.
- S. GUDMUNDSSON, 'The bibliography of harmonic morphisms', http://www.maths.lth.se/matematiklu/personal/sigma/harmonic/biblio--graphy.html
- [8] E. MAZET, 'La formule de la variation seconde de l'énergie au voisinage d'une application harmonique', J. Differential Geom. 8 (1973) 279–296.
- [9] S. MONTALDO, 'Stability of harmonic morphisms', Algebras, Groups and Geometries (to appear).
- [10] T. ISHIHARA, 'A mapping of Riemannian manifolds which preserves harmonic functions', J. Math. Kyoto Univ. 19 (1979) 215–229.
- [11] R.T. SMITH, 'Harmonic mappings of spheres', doctoral thesis, Warwick University, 1972.
- [12] R.T. SMITH, 'Harmonic mappings of spheres', Amer. J. Math. 97 (1975) 364–385.
- [13] G. TOTH, 'On rigidity of harmonic mappings into spheres', J. London Math. Soc. 26 (1982) 475–486.

UNIVERSITÀ DEGLI STUDI DI CAGLIARI, DIPARTIMENTO DI MATEMATICA, VIA OS-PEDALE 72, 09124 CAGLIARI

 $E\text{-}mail\ address: \texttt{montaldo@unica.it}$

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, GB E-mail address: j.c.wood@leeds.ac.uk