Projections in free product C*-algebras, II

Kenneth J. Dykema and Mikael Rørdam

22 August, 1999

Abstract

Let (A, φ) be the reduced free product of infinitely many C^{*}-algebras $(A_{\iota}, \varphi_{\iota})$ with respect to faithful states. Assume that the A_{ι} are not too small, in a specific sense. If φ is a trace then the positive cone of $K_0(A)$ is determined entirely by $K_0(\varphi)$. If, furthermore, the image of $K_0(\varphi)$ is dense in \mathbb{R} , then A has real rank zero. On the other hand, if φ is not a trace then A is simple and purely infinite.

Introduction

Let I be a set having at least two elements and, for every $\iota \in I$, let A_{ι} be a unital C^{*}-algebra with a state, φ_{ι} , whose GNS representation is faithful. Their reduced free product,

$$(A,\varphi) = \underset{\iota \in I}{*} (A_{\iota},\varphi_{\iota}) \tag{1}$$

was introduced by Voiculescu [20] and independently (in a more restricted way) by Avitzour [1]. Thus A is a unital C^{*}-algebra with canonical, injective, unital ^{*}-homomorphisms, $\pi_{\iota}: A_{\iota} \to A$, and φ is a state on A such that $\varphi \circ \pi_{\iota} = \varphi_{\iota}$ for all ι . It is the natural construction in Voiculescu's free probability theory (see [21]), and Voiculescu's theory has been vital to the study of these C^{*}-algebras.

In [12], for reduced free product C^{*}-algebras A as in (1), when all the φ_{ι} are faithful, we investigated projections in A and the related topic of positive elements in $K_0(A)$. The behaviour we discovered, under mild conditions specifying that the A_{ι} are not too small, depended broadly on whether φ is a trace, (i.e. on whether all the φ_{ι} are traces). If φ is a not trace then by [12] A is properly infinite. It remained open whether A must be purely infinite. (Some special classes of reduced free product C^{*}-algebras have in [13] and [9] been shown to be purely infinite.) When φ is a trace, then it follows from [12] that for every element, x, of the subgroup, G, of $K_0(A)$ generated by $\bigcup_{\iota \in I} K_0(\pi_\iota)(K_0(A_\iota))$, if $K_0(\varphi)(x) > 0$ then $x \ge 0$ and if $0 < K_0(\varphi)(x) < 1$ then there is a projection $p \in A$ such that $x = [p]_0$. By work of E. Germain [14], [15], [16], if each A_ι is an amenable C^{*}-algebra then $K_0(A) = G$ and G can be found from the groups $K_0(A_\iota)$ by using exact sequences, (and by taking inductive limits if I is infinite); hence under the hypothesis of amenability, we used Germain's results to give a complete characterization of the positive cone of $K_0(A)$ and of its elements corresponding to projections in A.

In the present paper we investigate similar questions for reduced free product C^{*-} algebras, (1), when I is infinite and when, for infinitely many $\iota \in I$, there is a unitary, $u \in A_{\iota}$ such that $\varphi_{\iota}(u) = 0$. We show that in this case, if φ is not a trace then A is purely infinite and simple. If φ is a trace then, although we do not know in general if the subgroup G described above exhausts $K_0(A)$, we nonetheless show that for every $x \in K_0(A)$, if $K_0(\varphi)(x) > 0$ then $x \ge 0$; furthermore, we show that if $x \in K_0(A)$ and if $0 < K_0(\varphi)(x) < 1$ then there is a projection $p \in A$ such that $x = [p]_0$. We also show that if the image of $K_0(\varphi)$ is dense in \mathbb{R} then A has real rank zero.

The real rank of a C^{*}-algebra, A, is denoted RR(A) and was invented by L.G. Brown and G.K. Pedersen [4]. Of particular interest is the case RR(A) = 0, which is defined, for a unital C^{*}-algebra A, to mean that the invertible self-adjoint elements are dense in the set of all self-adjoint elements of A. If φ is a faithful state on an infinite dimensional, simple C^{*}-algebra A, then a necessary condition for RR(A) = 0 is that there be projections, p, in A such that $\varphi(p)$ is arbitrarily small and positive; hence in particular, the image of $K_0(\varphi)$ must be dense in \mathbb{R} . We show that this condition is sufficient when A is a reduced free product of infinitely many algebras, as above, and when φ is a trace. (Moreover, when φ is not a trace then A is purely infinite, so by a result of S. Zhang [22], A has real rank zero.)

1 Comparison between positive elements and projections

Most of this section is a reformulation of results from [19]. The proof of Theorem 1.5 below is almost identical to the proof of [19, Theorem 7.2], but the statements of these two theorems are quite different.

We recall the notion of comparison of positive elements as introduced by J. Cuntz in [5] and [6] (see also [19]). Let A be a C^{*}-algebra, and let a, b be positive elements in A.

Then $a \leq b$ will mean that there exists a sequence $\{x_n\}_{n \in \mathbb{N}}$ in A with

$$\lim_{n \to \infty} \|a - x_n b x_n^*\| = 0.$$

If $p, q \in A$ are projections, then the definition above of $p \leq q$ agrees with the usual definition: $p = vv^*$ and $v^*v \leq q$ for some partial isometry $v \in A$.

If A is unital, and if φ is a state on A, then define $D_{\varphi} \colon A^+ \to [0, 1]$ by

$$D_{\varphi}(a) = \lim_{\varepsilon \to 0^+} \varphi(f_{\varepsilon}(a)),$$

where $f_{\varepsilon} \colon \mathbb{R}^+ \to [0, 1]$ is the continuous function, which is zero on $[0, \varepsilon/2]$, linear on $[\varepsilon/2, \varepsilon]$, and equal to 1 on $[\varepsilon, \infty)$. If φ is a trace, then D_{φ} is a dimension function (in the sense of Cuntz, [6]). Notice that $D_{\varphi}(p) = \varphi(p)$ for all projections $p \in A$.

We shall use the following facts:

$$f_{2\varepsilon}(a) \le f_{\delta}(f_{\varepsilon}(a)) \le f_{\varepsilon/2}(a), \qquad f_{\varepsilon/2}(a)f_{\varepsilon}(a) = f_{\varepsilon}(a),$$
(2)

when $\varepsilon > 0$ and $0 < \delta \leq 1/2$, and, consequently, $D_{\varphi}(f_{\varepsilon}(a)) \leq \varphi(f_{\varepsilon/2}(a))$. Moreover, if $0 \leq a \leq 1$, then $\varphi(a) \leq D_{\varphi}(a)$. Recall from [18] that the *stable rank* of a unital C^{*}-algebra A is equal to 1 if and only if the set invertible elements of A is dense in A.

Lemma 1.1 Let A be a C^{*}-algebra of stable rank one, let B be a hereditary subalgebra of A, let a be a positive element in B, and let q be a projection in B such that $a \leq q$. Then for each $\varepsilon > 0$ there is a projection $p \in B$ such that $f_{\varepsilon}(a) \leq p \sim q$.

Proof: Observe first that the comparisons $a \leq p$ and $p \sim q$ are independent of whether they are relative to A, B, or \tilde{B} , where \tilde{B} denotes the C*-algebra obtained by adjoining a unit to B. If follows from [18] and [3] that if A has stable rank one, then so do B and \tilde{B} . By [19, Proposition 2.4], there is for each $\varepsilon > 0$ a unitary u in \tilde{B} with $uf_{\varepsilon}(a)u^* \in q\tilde{B}q$ (= qBq). Put $p = u^*qu$. Then p is as desired. \Box

Lemma 1.2 Let A be a C^* -algebra, let a be a positive element in A, and let p be a projection in A. Then the following are equivalent:

- (i) $p \lesssim a$,
- (ii) $p = xax^*$ for some $x \in A$,

(iii) p is equivalent to some projection in the hereditary subalgebra of A generated by a.

Proof: (i) \Rightarrow (ii). If $p \leq a$, then $||p - yay^*|| < 1/2$ for some $y \in A$. Hence $pyay^*p$ is invertible (and positive) in pAp, and therefore $p = zpyay^*pz^*$ for some $z \in pAp$.

(ii) \Rightarrow (iii). Put $u = xa^{1/2}$. Then $uu^* = p$, and hence u is a partial isometry. Put $q = u^*u = a^{1/2}x^*xa^{1/2}$. Then q is a projection in the hereditary subalgebra generated by a, and $q \sim p$.

(iii) \Rightarrow (i). Assume q is a projection in the hereditary subalgebra generated by a, and that $q \sim p$. Then there exists an $n \in \mathbb{N}$ such that $||q - a^{1/n}qa^{1/n}|| < 1/2$. It follows that $qa^{1/n}qa^{1/n}q$ and, consequently, $qa^{2/n}q$, are invertible in qAq. Therefore $q = rqa^{2/n}qr^*$ for some $r \in qAq$. This shows that $p \sim q \leq a^{2/n} \leq a$. \Box

Lemma 1.3 ([19, Proposition 2.2]) Let A be a unital C*-algebra, let a, b be positive elements in A, and let $\varepsilon > 0$. If $||a - b|| < \varepsilon$, then $f_{\varepsilon}(a) \leq b$.

Lemma 1.4 Let A be a unital C^{*}-algebra, and let \mathfrak{A} be a dense unital *-subalgebra of A. Suppose that for each positive element $a \in \mathfrak{A}$ and each $\varepsilon > 0$ there is a projection $p \in A$ and $0 < \delta < \varepsilon$ such that $f_{\varepsilon}(a) \leq p \leq f_{\delta}(a)$. Then $\operatorname{RR}(A) = 0$.

Proof: To show that RR(A) = 0 it will suffice (by [4]) to show that all self-adjoint elements in the dense *-subalgebra \mathfrak{A} can be approximated by invertible self-adjoint elements.

Let a be a self-adjoint element in \mathfrak{A} , and write $a = a_+ - a_-$. For each $n \in \mathbb{N}$ find $\delta_n > 0$ and projections p_n, q_n in A such that

$$f_{1/n}(a_+) \le p_n \le f_{\delta_n}(a_+), \qquad f_{1/n}(a_-) \le q_n \le f_{\delta_n}(a_-).$$

Then $p_n \perp q_n$, $p_n a_+ p_n \rightarrow a_+$, and $q_n a_- q_n \rightarrow a_-$. Set

$$b_n = (p_n a_+ p_n + \frac{1}{n} p_n) - (q_n a_- q_n + \frac{1}{n} q_n) + \frac{1}{n} (1 - p_n - q_n).$$

Then each b_n is invertible and self-adjoint, and $b_n \to a$. \Box

Let \mathcal{Q} be a compact convex subset of the state space of a unital C*-algebra A, and let $\operatorname{Aff}(\mathcal{Q})$ denote the real vector space of all affine continuous functions $\mathcal{Q} \to \mathbb{R}$. Equip this space with the strict ordering, i.e., $f \geq 0$ if f = 0 or if $f(\varphi) > 0$ for all $\varphi \in \mathcal{Q}$, and, in turn, with the topology induced by this ordering. All self-adjoint elements $a \in A$ induce

an element $\hat{a} \in \operatorname{Aff}(\mathcal{Q})$ through the formula $\hat{a}(\varphi) = \varphi(a)$. We will consider the interval [0, 1] of $\operatorname{Aff}(\mathcal{Q})$, defined by

$$[0,1] = \{ f \in \operatorname{Aff}(\mathcal{Q}) \mid 0 \le f \le 1 \}.$$

Theorem 1.5 Let A be a unital C^* -algebra, let Q be a compact convex subset of the state space of A, let Π be a subset of of the set of projections in A, and let \mathfrak{A} be a dense *-subalgebra of A, which is closed under continuous function calculus (on its normal elements).

Assume that each state in Q is faithful on \mathfrak{A} , and that the following comparison properties hold for all positive elements $a \in \mathfrak{A}$ and for all projections $p \in \Pi$:

- (α) if $D_{\varphi}(a) < \varphi(p)$ for all $\varphi \in \mathcal{Q}$, then $a \leq p$,
- (β) if $\varphi(p) < D_{\varphi}(a)$ for all $\varphi \in \mathcal{Q}$, then $p \leq a$, and
- (γ) the subset of Aff(\mathcal{Q}) induced by Π is dense in the interval [0,1] of Aff(\mathcal{Q}).

It follows that

- (i) if sr(A) = 1, then RR(A) = 0, and
- (ii) if all nonzero projections in Π are infinite and full, then A is simple and purely infinite.

Proof: (i). We show that the conditions in Lemma 1.4 are satisfied. So let $a \in \mathfrak{A}$ be a positive element, and let $\varepsilon > 0$. We must find $0 < \delta < \varepsilon$ and a projection $p \in A$ with $f_{\varepsilon}(a) \leq p \leq f_{\delta}(a)$.

If $\operatorname{sp}(a) \cap (\varepsilon/8, \varepsilon/4) = \emptyset$, then $p = f_{\varepsilon/8}(a)$ and $\delta = \varepsilon/8$ will be as desired.

Assume now that $\operatorname{sp}(a) \cap (\varepsilon/8, \varepsilon/4) \neq \emptyset$. Then $0 \leq \varphi(f_{\varepsilon/4}(a)) < \varphi(f_{\varepsilon/8}(a)) \leq 1$ for all $\varphi \in \mathcal{Q}$ because each such φ is assumed to be faithful on \mathfrak{A} . Since Π is dense in the interval [0, 1] of $\operatorname{Aff}(\mathcal{Q})$ there is $q \in \Pi$ with $\varphi(f_{\varepsilon/4}(a)) < \varphi(q) < \varphi(f_{\varepsilon/8}(a))$ for all $\varphi \in \mathcal{Q}$. By (2),

$$D_{\varphi}(f_{\varepsilon/2}(a)) \le \varphi(f_{\varepsilon/4}(a)) < \varphi(q) < \varphi(f_{\varepsilon/8}(a)) \le D_{\varphi}(f_{\varepsilon/8}(a))$$

By assumptions (α) and (β) this implies that $f_{\varepsilon/2}(a) \leq q \leq f_{\varepsilon/8}(a)$.

From Lemma 1.2 there is a projection r in the hereditary subalgebra, B, generated by $f_{\varepsilon/8}(a)$ such that $q \sim r$. By Lemma 1.1 there is a projection $p \in B$ such that $f_{1/2}(f_{\varepsilon/2}(a)) \leq c_{1/2}(a)$

 $p \text{ (and } p \sim r)$. By (2), this entails that $f_{\varepsilon}(a) \leq p \leq f_{\varepsilon/16}(a)$. The claim is therefore proved with $\delta = \varepsilon/16$.

(ii). If each non-zero hereditary subalgebra of A contains a full element, then A must be simple. If, moreover, each such hereditary subalgebra contains an infinite projection, then A is purely infinite and simple (c.f. Cuntz' definition of purely infinite simple C*-algebras in [7]). It therefore suffices to show that each non-zero hereditary subalgebra of A contains an infinite full projection.

Let *B* be a non-zero hereditary subalgebra of *A*, and let *b* be a positive element in *B* with ||b|| = 1. Find a positive element $a \in \mathfrak{A}$ with ||a - b|| < 1/2. Since each $\varphi \in \mathcal{Q}$ is faithful, since $f_{1/2}(a) \neq 0$, and since Π is dense in the interval [0,1] of Aff (\mathcal{Q}) , there is $q \in \Pi$ with $\varphi(q) < \varphi(f_{1/2}(a))$ for all $\varphi \in \mathcal{Q}$. This implies that $\varphi(q) < D_{\varphi}(f_{1/2}(a))$ for all $\varphi \in \mathcal{Q}$, and by assumption (β) we get $q \leq f_{1/2}(a)$. Using Lemma 1.3 we obtain that $q \leq b$, and Lemma 1.2 finally implies that there is a projection *p* in the hereditary subalgebra of *A* generated by *b* (which is contained in *B*, so that $p \in B$) such that $p \sim q$. Since *q* is infinite and full, so is *p*, and the proof is complete. \Box

2 Application to reduced free products of C^* -algebras

Throughout this section, we consider a reduced free product of C^{*}-algebras,

$$(A,\varphi) = \underset{\iota \in I}{*} (A_{\iota},\varphi_{\iota}), \tag{3}$$

where I is an infinite set, where each φ_{ι} is a faithful state and where for infinitely many $\iota \in I$ there is a unitary $u \in A_{\iota}$ with $\varphi_{\iota}(u) = 0$. It follows from [8] that φ is faithful on A.

Avitzour's result [1] gives that A is simple if, for example, φ is a trace. Indeed, by partitioning the set I into two suitable subsets, A can be viewed as a reduced free product,

$$(A, \varphi) = (B_1, \psi_1) * (B_2, \psi_2),$$

such that there are unitaries, $u \in B_1$ and $v, w \in B_2$ satisfying that $\psi_1(u) = 0 = \psi_2(v) = \psi_2(w)$ and that v and w are *-free; hence also $\psi_2(v^*w) = 0$. (Avitzour's result also applies in somewhat more general instances.) In addition, if φ is a trace then by [10] the stable rank of A is equal to 1.

The K_0 -group, $K_0(D)$, of a C^{*}-algebra D is equipped with a positive cone and a scale

defined respectively by

$$K_0(D)^+ = \{ [p]_0 \mid p \in \operatorname{Proj}(D \otimes \mathcal{K}) \},$$

$$\Sigma(D) = \{ [p]_0 \mid p \in \operatorname{Proj}(D) \},$$

where $\operatorname{Proj}(D)$ is the set of projection in D, and where $[\cdot]_0: \operatorname{Proj}(D \otimes \mathcal{K}) \to K_0(D)$ is the canonical map from which K_0 is defined. The positive cone gives rise to an ordering on $K_0(D)$ by $x \leq y$ if $y - x \in K_0(D)^+$, and x < y if $y - x \in K_0(D)^+ \setminus \{0\}$. Each (positive) trace φ on D induces a positive group-homomorphism $K_0(\varphi): K_0(D) \to \mathbb{R}$ which satisfies $K_0(\varphi)([p]_0) = \varphi(p)$ for $p \in \operatorname{Proj}(D)$, and $K_0(\varphi)([p]_0) = (\varphi \otimes \operatorname{Tr}_n)(p)$ for $p \in \operatorname{Proj}(D \otimes M_n(\mathbb{C}))$, where Tr_n is the (unnormalized) trace on $M_n(\mathbb{C})$. The ordered abelian group $(K_0(D), K_0(D)^+)$ is called *weakly unperforated* if nx > 0 for some $n \in \mathbb{N}$ and some $x \in K_0(D)$ implies that $x \geq 0$.

Theorem 2.1 Let

$$(A,\varphi) = \underset{\iota \in I}{*} (A_{\iota},\varphi_{\iota})$$

be the reduced free product C^* -algebra, where each A_{ι} is a unital C^* -algebra, φ_{ι} is a faithful state on A_{ι} , the index set I is infinite, and infinitely many A_{ι} contain a unitary in the kernel of φ_{ι} .

If φ is a trace (which is the case if all φ_{ι} are traces), then

- (i) whenever $p, q \in A \otimes M_n(\mathbb{C})$ are projections such that $(\varphi \otimes \operatorname{Tr}_n)(p) < (\varphi \otimes \operatorname{Tr}_n)(q)$, it follows that $p \leq q$;
- (ii) the positive cone and the scale of $K_0(A)$ are given by

$$K_0(A)^+ = \{0\} \cup \{x \in K_0(D) \mid 0 < K_0(\varphi)(x)\},$$

$$\Sigma(A) = \{0,1\} \cup \{x \in K_0(D) \mid 0 < K_0(\varphi)(x) < 1\},$$

and, as a consequence, $(K_0(A), K_0(A)^+)$ is weakly unperforated;

(iii) $\operatorname{RR}(A) = 0$ if and only if $K_0(\varphi)(K_0(A))$ is dense in \mathbb{R} .

If φ is not a trace (i.e., if at least one φ_{ι} is not a trace), then A is simple and purely infinite.

Proof: We consider, for every finite subset $F \subseteq I$, the C^{*}-subalgebra, \mathfrak{A}_F , of A generated by $\bigcup_{\iota \in F} \pi_\iota(A_\iota)$, and we let $\mathfrak{A} = \bigcup_{F \ll I} \mathfrak{A}_F$, where the union is over all finite subsets of

I. Note that \mathfrak{A} is a dense, unital *-subalgebra of A that is closed under the continuous functional calculus.

Suppose that φ is a trace, let $n \in \mathbb{N}$ and let $p, q \in A \otimes M_n(\mathbb{C})$ be projections with $(\varphi \otimes \operatorname{Tr}_n)(p) < (\varphi \otimes \operatorname{Tr}_n)(q)$. Using the density of \mathfrak{A} in A and continuous functional calculus, we find a finite subset F of I and projections $\tilde{p}, \tilde{q} \in \mathfrak{A}_F \otimes M_n(\mathbb{C})$ such that $\|\tilde{p}-p\| < 1$ and $\|\tilde{q}-q\| < 1$. This implies $\tilde{p} \sim p$ and $\tilde{q} \sim q$. There are n^2 distinct elements $\iota(1), \iota(2), \ldots, \iota(n^2) \in I \setminus F$ with unitaries $u_k \in A_{\iota(k)}$ such that $\varphi_{\iota(k)}(u_k) = 0$. Let B be the C^{*}-algebra generated by $\{u_1, u_2, \ldots, u_{n^2}\}$. Note that B and \mathfrak{A}_F are free. Then as in the proof of Proposition 3.3 of [12], from the unitaries $u_1, u_2, \ldots, u_{n^2}$ we can construct a Haar unitary, $v \in B \otimes M_n(\mathbb{C})$ such that $\{v\}$ and $\mathfrak{A}_F \otimes M_n(\mathbb{C})$ are *-free (with respect to the tracial state $\varphi \otimes (\frac{1}{n} \operatorname{Tr}_n)$). Now $\tilde{q} \sim v^* \tilde{q}v$ and the pair \tilde{p} and $v^* \tilde{q}v$ is free; moreover, $(\varphi \otimes \operatorname{Tr}_n)(v^* \tilde{q}v) = (\varphi \otimes \operatorname{Tr}_n)(\tilde{q})$. So by Proposition 1.1 of [12], $v^* \tilde{q}v$ is equivalent to a subprojection, r, of \tilde{p} ; hence $q \leq p$. We have thus proved (i).

The inclusions \subseteq in (ii) are easy consequences of the fact that φ is faithful. Assume $x \in K_0(A)$ and that $K_0(\varphi)(x) > 0$. Since A is unital, there are $n \in \mathbb{N}$ and projections $p, q \in A \otimes M_n(\mathbb{C})$ such that $x = [p]_0 - [q]_0$. Now,

$$(\varphi \otimes \operatorname{Tr}_n)(p) - (\varphi \otimes \operatorname{Tr}_n)(q) = K_0(\varphi)(x) > 0.$$

Hence, by (i), q is equivalent to a subprojection \tilde{q} of p. Thus $x = [p - \tilde{q}]_0 \in K_0(A)^+$.

Assume next that $x \in K_0(A)$ and that $0 < K_0(\varphi)(x) < 1$. Then, by the argument above, $x = [p]_0$ for some projection $p \in A \otimes M_n(\mathbb{C})$. Let 1_A denote the unit of A, and let $e \in A \otimes M_n(\mathbb{C})$ be the diagonal projection whose upper left corner is 1_A and with all other entries equal to 0. Then $(\varphi \otimes \operatorname{Tr}_n)(p) = K_0(\varphi)(x) < 1 = (\varphi \otimes \operatorname{Tr}_n)(e)$. By (i), this implies that p is equivalent to a subprojection \tilde{p} of e. Hence $x = [\tilde{p}]_0$, and it is easily seen that $[\tilde{p}]_0 \in \Sigma(A)$.

Finally, to see that $(K_0(A), K_0(A)^+)$ is weakly unperforated, assume that $x \in K_0(A)$ and that nx > 0 for some $n \in \mathbb{N}$. Then $K_0(\varphi)(x) = \frac{1}{n}K_0(\varphi)(nx) > 0$. Hence x > 0. We have thus shown (ii).

Let

$$\Pi = \bigcup_{F \ll I} \operatorname{Proj}(\mathfrak{A}_F).$$

For the set \mathcal{Q} used in Theorem 1.5, we take the singleton $\{\varphi\}$. We now show that, regardless of whether φ is a trace or not, conditions (α) and (β) of Theorem 1.5 hold for every $p \in \Pi$ and every positive element, $a \in \mathfrak{A}$. Given a positive element $a \in \mathfrak{A}$ and given $p \in \Pi$, there is a finite subset F of I such that $a, p \in \mathfrak{A}_F$. Let $u \in A_\iota$, for some $\iota \in I \setminus F$, be a unitary such that $\varphi_\iota(u) = 0$. Then $\{a, p\}$ and $\{u\}$ are *-free with respect to φ . Now it follows that u^*pu is a projection with $\varphi(u^*pu) = \varphi(p)$, and that u^*pu and a are free. Hence by Lemma 5.3 of [12] it follows that $a \leq u^*pu$ if $D_{\varphi}(a) < \varphi(p)$ and $u^*pu \leq a$ if $\varphi(p) < D_{\varphi}(a)$. But $u^*pu \sim p$, so (α) and (β) hold.

Suppose now that φ is a trace and that the image of $K_0(\varphi)$ is dense in \mathbb{R} , and let us show that $\operatorname{RR}(A) = 0$. We will show that $\{\varphi(p) \mid p \in \Pi\}$ is dense in [0, 1], which will imply that condition (γ) of Theorem 1.5 holds. Since the image of $K_0(\varphi)$ is dense in \mathbb{R} , the intersection of this image with [0, 1] is dense in [0, 1]. By (ii), it follows that $\{\varphi(p) \mid p \in \operatorname{Proj}(A)\}$ is dense in [0, 1]. Since \mathfrak{A} is dense in A, and using continuous functional calculus, we find for every $p \in \operatorname{Proj}(A)$, a projection, $\tilde{p} \in \mathfrak{A}$ such that $\varphi(\tilde{p}) = \varphi(p)$. But $\tilde{p} \in \Pi$. We have shown that condition (γ) of Theorem 1.5 holds, and we have already shown that conditions (α) and (β) hold. Now using the fact that $\operatorname{sr}(A) = 1$, we get from Theorem 1.5(i) that $\operatorname{RR}(A) = 0$. This implies one direction of (iii), but the other direction follows from more general results. Indeed, the image of $K_0(\varphi)$ will be dense in \mathbb{R} if A contains at least one projection and if A has no minimal projections. Both of these conditions hold if $\operatorname{RR}(A) = 0$, and if A is simple and infinite dimensional, as in our case.

Now suppose that φ is not a trace, and let us show that A is purely infinite and simple. Let F be a finite subset of I such that for at least three distinct $\iota \in F$ there is a unitary $u \in A_{\iota}$ satisfying $\varphi_{\iota}(u) = 0$, and such that for some $\iota \in F$, φ_{ι} is not a trace. Then by Theorem 4 of [12], the unit is a properly infinite projection in \mathfrak{A}_{F} . Let $\Pi' \subseteq \Pi$ be the set of all full, properly infinite projections in \mathfrak{A} . We have already shown that conditions (α) and (β) are satisfied for every $a \in \mathfrak{A}$ and every $p \in \Pi'$. Since 1 is a properly infinite projection in some \mathfrak{A}_{F} , using Lemma 2.2 below we get that $\{\varphi(p) \mid p \in \Pi'\}$ is dense in [0, 1], so condition (γ) is satisfied. Tautologically, each $p \in \Pi'$ is infinite and full. Hence by Theorem 1.5(ii), A is purely infinite and simple. \Box

Lemma 2.2 Let A be a unital C^{*}-algebra in which 1 is properly infinite and let φ be a state on A. Then for every $t \in \mathbb{R}$, $0 < t \leq 1$, there is a projection $p \in A$ such that $p \sim 1$ and $\varphi(p) = t$.

Proof: Using that 1 is properly infinite, we find isometries, v_1, v_2, \ldots in A whose range projections are mutually orthogonal. These generate a unital C^{*}-subalgebra of A isomorphic to the Cuntz algebra \mathcal{O}_{∞} . By Cuntz's paper [7], it follows that if $p, q \in \operatorname{Proj}(\mathcal{O}_{\infty}) \setminus \{0, 1\}$ and if $[p]_0 = [q]_0$ in $K_0(\mathcal{O}_{\infty})$ then p is homotopic to q. (Indeed, it follows that $p \sim q$ and $1-p \sim 1-q$, hence p is unitarily similar to q. But the unitary group of \mathcal{O}_{∞} is connected.) Now let $\varepsilon > 0$. For some *n* we must have $\varphi(v_n v_n^*) < \varepsilon$; let $p = v_n v_n^*$. Then $q \stackrel{\text{def}}{=} 1 - v_n(1-p)v_n^*$ is a projection in \mathcal{O}_{∞} with $[q]_0 = [1]_0$, and $\varphi(q) > 1 - \varepsilon$. Thus there is a continuous path r_t in $\operatorname{Proj}(\mathcal{O}_{\infty})$ such that $r_0 = p$ and $r_1 = q$. We have that $r_t \sim 1$ for all t and $\{\varphi(r_t) \mid t \in [0,1]\} \supseteq (\varepsilon, 1-\varepsilon)$. \Box

Let us now state a straightforward application of Theorem 2.1 to reduced group C^{*}– algebras. For a group, G, taken with the discrete topology, the reduced group C^{*}–algebra of G, denoted $C^*_{\text{red}}(G)$, is the C^{*}–algebra generated by the left regular representation of G. The canonical tracial state, τ_G , is the vector state for the characteristic function of the identity element of G. The following corollary was cited in [11], where also a partial converse was included.

Corollary 2.3 Let I be an infinite set and let

$$G = \underset{\iota \in I}{*} G_{\iota}$$

be the free product of nontrivial groups, G_{ι} . Suppose that G has finite subgroups of arbitrarily large order. Then

$$\operatorname{RR}(C^*_{\operatorname{red}}(G)) = 0.$$

Proof: We have

$$(C^*_{\mathrm{red}}(G), \tau_G) \cong \underset{\iota \in I}{*} (C^*_{\mathrm{red}}(G_\iota), \tau_{G_\iota}).$$

If x is a nontrivial element of G_{ι} then the left translation operator, $\lambda_x \in C^*_{\text{red}}(G_{\iota})$ is a unitary and $\tau_{G_{\iota}}(\lambda_x) = 0$. In order to apply Theorem 2.1, it is thus sufficient to show that $C^*_{\text{red}}(G)$ contains projections whose traces (under τ_G) are arbitrarily small and positive. But this is clear, since for a finite group H, $C^*_{\text{red}}(H)$ contains projections of trace 1/|H|. \Box

It follows easily from the Kurosh Subgroup Theorem for free products of groups, (see page 178 of [17]), that G has finite subgroups of arbitrarily high order only if for every positive integer n there is $\iota \in I$ such that G_{ι} has a finite subgroup of order greater than n.

Example 2.4 If

$$G = \overset{\infty}{\underset{n=2}{\ast}} (\mathbb{Z}/n\mathbb{Z}),$$

then $C^*_{\text{red}}(G)$ has real rank zero. Moreover, this C^* -algebra is simple, has unique tracial state, has stable rank one and is not approximately divisible.

Proof: It has real rank zero by the above corollary. It is simple and has unique tracial state by Avitzour [1]. It has stable rank one by [10]. That it is not approximately divisible follows from the argument in Example 4.8 of [2], because $C^*_{\text{red}}(G)$ is weakly dense in the group von Neumann algebra L(G), which does not have central sequences. \Box

References

- D. AVITZOUR, Free products of C*-algebras, Trans. Amer. Math. Soc. 271 (1982), 423-465.
- [2] B. BLACKADAR, A. KUMJIAN AND M. RØRDAM, Approximately central matrix units and the structure of noncommutative tori *K*-theory **6** (1992), 267-284.
- [3] L.G. BROWN, Stable isomorphism of hereditary subalgebras of C*-algebras, *Pacific J. Math.* **71** (1977), 335-348.
- [4] L.G. BROWN, G.K. PEDERSEN, C*-algebras of real rank zero, J. Funct. Anal. 99 (1991), 131-149.
- [5] J. CUNTZ, The structure of multiplication and addition in simple C*-algebras, Math. Scand. 40 (1977), 215–233.
- [6] J. CUNTZ, Dimension functions on simple C*-algebras, Math. Ann. 233 (1978), 145– 153.
- [7] J. CUNTZ, K-theory for certain C*-algebras, Ann. of Math. 113 (1981), 181–197.
- [8] K.J. DYKEMA, Faithfulness of free product states, J. Funct. Anal. 154 (1998), 223– 229.
- [9] K.J. DYKEMA, Purely infinite simple C*-algebras arising from free product constructions, II, *preprint* (1997).
- [10] K.J. DYKEMA, U. HAAGERUP, M. RØRDAM, The stable rank of some free product C*-algebras Duke Math. J. 90 (1997), 95-121, correction 94 (1998), 213.
- [11] K.J. DYKEMA, P. DE LA HARPE, Some groups whose reduced C^{*}-algebras have stable rank one, *J. Math. Pures Appl.* (to appear).

- [12] K.J. DYKEMA, M. RØRDAM, Projections in free product C^{*}-algebras, Geom. Funct. Anal. 8 (1998), 1-16.
- [13] K.J. DYKEMA, M. RØRDAM, Purely infinite simple C*-algebras arising from free product constructions, *Canad. J. Math*, **50** (1998), 323-342.
- [14] E. GERMAIN, KK-theory of reduced free product C*-algebras Duke Math. J. 82 (1996), 707-723.
- [15] E. GERMAIN, KK-theory of the full free product of unital C*-algebras J. reine angew. Math. 485 (1997), 1-10.
- [16] E. GERMAIN, Amalgamated free product of C*-algebras and KK-theory Fields Inst. Commun. 12, D. Voiculescu, ed., (1997), 89-103.
- [17] R.C. LYNDON, P.E. SCHUPP, Combinatorial Group Theory, Springer-Verlag, 1977.
- [18] M.A. RIEFFEL, Dimension and stable rank in the K-theory of C*-algebras, Proc. London Math. Soc. (3), 46 (1983), 301–333.
- [19] M. RØRDAM, On the structure of simple C*-algebra tensored with a UHF-algebra, II, J. Funct. Anal. 107 (1992), 255–269.
- [20] D. VOICULESCU, Symmetries of some reduced free product C*-algebras, Operator Algebras and Their Connections with Topology and Ergodic Theory, Lecture Notes in Mathematics, vol. 1132, Springer-Verlag, 1985, 556–588.
- [21] D. VOICULESCU, K.J. DYKEMA, A. NICA, Free Random Variables, CRM Monograph Series vol. 1, American Mathematical Society, 1992.
- [22] S. ZHANG, Certain C*-algebras with real rank zero and their corona and multiplier algebras. I. Pacific J. Math. 155 (1992), 169–197.

Department of Mathematics and Computer Science, Odense University, Campusvej 55, 5230 Odense M, Denmark

E-mail address: dykema@imada.ou.dk Internet home page: www.imada.ou.dk/~dykema/ Department of Mathematics, University of Copenhagen, Universitets parken 5, 2100 Copenhagen \emptyset , Denmark

E-mail address: rordam@math.ku.dk Internet home page: www.math.ku.dk/~rordam/