A NOTE ON A THEOREM OF H. CARTAN

José M. Isidro

Universidad de Santiago de Compostela

ABSTRACT. We prove that if $D \subset \mathbb{C}^n$ is a bounded domain with real analytic boundary, and D is either pseudoconvex or it satisfies condition R, then the the compact open topology in Aut (D), the group of holomorphic automorphisms of D is the the topology of uniform convergence on D.

0 Introduction

Let D be a bounded domain in \mathbb{C}^n with $n \ge 1$, and let Aut (D) denote the group of holomorphic automorphisms of D. It was shown by H. Cartan [2, Chap. 9 th. 4] that Aut (D), endowed with the compact open topology (that is, the topology of local uniform convergence on D), is a real Lie group whose Lie algebra $\mathfrak{aut}(D)$ consists of all complete holomorphic vector fields $X: D \to \mathbb{C}^n$, and that the natural action

Aut $(D) \times D \to D$, $(f, z) \mapsto f(z)$

is real analytic in joint variables. This result is no longer valid if \mathbb{C}^n is replaced by an arbitrary complex Banach space E; however, it is still valid if we assume that $D \subset E$ is a bounded symmetric domain ([4], [5]).

It is remarkable that no assumptions on the smoothness of the boundary ∂D is needed either in the \mathbb{C}^n setting or in the infinite dimensional case. This leads naturally to the question of whether every f in Aut (D) extends holomorphically beyond ∂D . Affirmative answers to this question have been found, in the \mathbb{C}^n setting, by S. Chen S. Jang in [1, th.1.2] assuming that Dhas real analytic boundary and that either D is pseudoconvex or that condition (R) holds on D. By condition (R) we mean that the Bergmann projector (that is, the orthogonal projection from $L^2(D)$ onto the the closed subspace $\mathcal{O}^2(D)$ of square integrable holomorphic functions) maps $\mathcal{C}^{\infty}(\overline{D})$ continuously into itself). In the infinite dimensional context, an affirmative answer has been established in [6] for bounded circular domains with no restrictions on the boundary.

In turn, the possibility of extending every element f in Aut (D) to a neighbourhood of \overline{D} gives new information about the topology of Aut (D). Indeed, it has been proved in [6] that for bounded circular domains D in a Banach space E, the topology on Aut (D) of local uniform convergence over D is actually the topology of uniform convergence on D. In this note we use the results in [1] to prove the following theorem:

Theorem Let $D \subset \mathbb{C}^n$, where $n \geq 1$, be a bounded domain with real analytic boundary, and suppose that either D is pseudoconvex or it satisfies condition (R). Then the compact open topology on Aut (D) is the same as the topology of uniform convergence over D.

¹⁹⁹¹ Mathematics Subject Classification. 48 G 20, 72 H51.

Key words and phrases. Holomorphic automorphisms, Lie groups..

Supported by Comisión Hispano-Húngara de Cooperación Científica y Tecnológica

JOSÉ M. ISIDRO

For a domain $D \subset \mathbb{C}^n$, we let Hol (D, \mathbb{C}^n) denote the vector space of all holomorphic mappings $h: D \to \mathbb{C}^n$, and the subset Hol (D) consists of the mappings such that $f(D) \subset D$. A set $S \subset D$ is a vanishing set for D if the relations $f \in \text{Hol}(D, \mathbb{C}^n)$ and $f_{|S} = 0$ entail f = 0. We say that ∂D is an algebraically determining set for Aut (D) if the relations $f, g \in \text{Aut}(D)$ and f = g on ∂D entail f = g. Similarly, the expression ∂D is a topologically determining set for Aut (D) means that whenever $(f_n)_{n\in\mathbb{N}}$ and f are respectively a sequence and an element in Aut (D) such that $f_n \to f$ uniformly over ∂D we have $f_n \to f$ in Aut (D). Obvious changes give us the meaning of the expressions $S \subset D$ is an algebraically or topologically determining set for $\mathfrak{aut}(D)$, see [3]

1 The main result.

1.1 Theorem. Let $D \subset \mathbb{C}^n$, where $n \geq 1$, be a bounded domain with real analytic boundary, and suppose that either D is pseudoconvex or it satisfies condition (R). Then for every sequence $(f_n)_{n\in\mathbb{N}}$ in Aut (D) and every f in Hol (D) the following conditions are equivalent:

- (1) $f \in Aut(D)$ and there is a neighbourhood Ω of \overline{D} in \mathbb{C}^n such that $(f_n)_{n \in \mathbb{N}}$ converges to f uniformly on Ω .
- (2) $f \in Aut$ (D) and $(f_n)_{n \in \mathbb{N}}$ converges to f in the group Aut (D).
- (3) There are a non void open subset $U \subset D$ and a point $a \in U$ such that $(f_n(z))_{n \in \mathbb{N}}$ converges to f(z) pointwise on U and $f(a) \notin \partial D$.

Proof. We only need to prove $3 \Longrightarrow 1$. By the identity principle, U is a vanishing set for D. Also the family $(f_n)_{n \in \mathbb{N}}$ is uniformly bounded in D. Therefore by Vitali's theorem [2, Chap. 1 prop. 7] the sequence $(f_n)_{n \in \mathbb{N}}$ converges to f uniformly on each compact subset $K \subset D$. Since $f(a) \notin \partial D$, we have $f \in \text{Aut}(D)$ by Cartan's theorem [2, Chap. 5 th. 4]. On the other hand, since D either is pseudoconvex or it satisfies condition (R), by [1, th. 1.2] there are a neighbourhood \mathcal{V}_f of f in Aut (D) and a neighbourhood Ω_f of \overline{D} in \mathbb{C}^n such that every $g \in \mathcal{V}_f$ extends to some $\tilde{g} \in \text{Hol}(\Omega_f)$ and the action

$$\mathcal{V}_f \times \Omega_f \to \mathbb{C}^n, \qquad (g, z) \mapsto \tilde{g}(z)$$

is real analytic on $\mathcal{V}_f \times \Omega_f$. By restricting ourselves to smaller neighbourhoods $\mathcal{W} \subset \mathcal{V}_f$ and $\Omega \subset \Omega_1 \subset \Omega_f$ we may assume that $\overline{\mathcal{W}}$ and $\overline{\Omega}_1$ are compact and that the latter is connected. Since $f_n \to f$ in Aut (D), we have $f_n \in \mathcal{W}$ for n large enough, say $n \ge n_0$ and so

$$\sup_{n \ge n_0, z \in \Omega} \|f_n(z)\| \le M < \infty,$$

and a new application of Vitali's theorem yields that $f_n \to f$ uniformly on each compact subset of Ω_1 , in particular on Ω . \Box

1.2 Corollary. Let the domain $D \subset \mathbb{C}^n$ satisfy the assumptions in (1.1). Then

- (1) The compact open topology on Aut (D) coincides with the topology of uniform convergence over \overline{D} and with the topology of uniform convergence over ∂D .
- (2) ∂D is a determining set for the topological group Aut (D).
- (3) If D is a bounded convex domain with real analytic boundary, then every element in Aut (D) has a fixed point in \overline{D} .

Proof. (1): It is clear from the above proof that on Aut (D) uniform convergence over the compact subsets of D is the same as uniform convergence over \overline{D} , which in turn is the same as uniform convergence over ∂D as a consequence of the maximum modulus theorem. (2) and (3): Extend to \overline{D} the elements in Aut (D) and apply respectively the maximum modulus theorem and Brower's fixed point theorem. \Box

We now consider the similar problems at the Lie algebra level.

1.3 Theorem. Let $D \subset \mathbb{C}^n$, where $n \geq 1$, be a bounded domain with real analytic boundary, and suppose that either D is pseudoconvex or it satisfies condition (R). Then there are a neighbourhood \mathcal{M}_0 of 0 in aut (D) and a neighbourhood Ω_0 of \overline{D} in \mathbb{C}^n such that every vector field $X \in \mathcal{M}_0$ extends to some $\tilde{X} \in \text{Hol}(\Omega_0, \mathbb{C}^n)$ that is the infinitesimal generator of a local one-parameter group of holomorphic transformations on Ω_0 .

Proof. Choose a neighbourhood V_{ld} of ld in the group $\mathsf{Aut}(D)$ and a neighbourhood Ω_0 of \overline{D} in \mathbb{C}^n such that every element $f \in V_{\mathsf{ld}}$ extends to a holomorphic mapping \tilde{f} in Ω_0 and the action

$$\mathcal{V}_{\mathsf{Id}} \times \Omega_0 \to \mathbb{C}^n, \qquad (f, z) \mapsto \tilde{f}(z)$$

is real analytic on $\mathcal{V}_{\mathsf{Id}} \times \Omega_0$. The exponential mapping exp defines a homeomorphism of a neighbourhood $\mathcal{W}_{\mathsf{Id}}$ of Id in $\mathsf{Aut}(D)$ onto a neighbourhood of 0 in $\mathfrak{aut}(D)$. Let $\mathcal{U}_0 := \mathcal{V}_{\mathsf{Id}} \cap \mathcal{W}_{\mathsf{Id}}$ and set $\mathcal{M}_0 := \exp^{-1}\mathcal{U}_0$. We may assume that \mathcal{M}_0 is balanced hence connected. Let $X \in \mathcal{M}_0$ and let $f_t := \exp tX \in \mathsf{Aut}(D)$ denote the one-parameter group corresponding to X. For small values of t, say $|t| < \tau$, we have $f_t \in \mathcal{U}_0$, hence f_t extends to a holomorphic mapping $f_t \in \mathsf{Hol}(\Omega_0, \mathbb{C}^n)$. By the identity principle we have $f_t \circ f_s = f_{t+s}$ whenever $|t+s| < \tau$. Thus $t \mapsto f_t$ is a local one- parameter group of holomorphic transformations on Ω_0 and so its infinitesimal generator $\tilde{X} = \frac{d}{dt}_{|t=0} f_t$ is a holomorphic mapping on Ω_0 that extends X. \Box

1.4 Corollary. Let the domain $D \subset \mathbb{C}^n$ satisfy the assumptions in (1.3). Then ∂D is a determining set for aut (D), and for every sequence $(X_n)_{n \in \mathbb{N}}$ in aut (D) and every function $X \in \mathsf{Hol}(D, \mathbb{C}^n)$ the following conditions are equivalent:

- (1) $X \in \mathfrak{aut}$ (D) and there is a neighbourhood Ω of \overline{D} in \mathbb{C}^n such that $X_n \to X$ uniformly on Ω .
- (2) $X \in \mathfrak{aut}(D)$ and $(X_n)_{n \in \mathbb{N}}$ converges to X in the Lie algebra $\mathfrak{aut}(D)$.
- (3) $(X_n(z))_{n\in\mathbb{N}}$ is uniformly bounded in D and $X_n \to X$ pointwise on some open non void subset $U \subset D$

Proof. Clearly $1 \implies 2$. Suppose that (2) holds; then a repetition of the arguments made in the proof of (1.3) show that X and all X_n extend holomorphically to a suitable neighbourhood Ω of \overline{D} and $X_n \to X$ uniformly over Ω , hence in particular $(X_n)_{n \in \mathbb{N}}$ is uniformly bounded on D. Now suppose (3) holds; by Vitali's theorem we have $X_n \to X$ uniformly over the compact subset of D, hence $X \in \mathfrak{aut}(D)$, and by (1.3) all X_n and X extend holomorphically to some neighbourhood Ω of \overline{D} . A new application of Vitali's theorem gives the result. The other claim is now obvious. \Box

References

- Chen So-Chin & Jang Shin-Biau, An extension of H. Cartan theorem, Proc. Amer. Math. Soc. 127 (1999), 2265-2271.
- Narasimhan, R., Several Complex Variables, Chicago Lectures in Mathematics, The University of Chicago Press, 1970.
- Isidro, J. M. & Kaup, W., Determining Boundary Sets of Bounded Symmetric Domains, Manuscripta Math. 81 (1993), 149-159.
- Upmeier, H., Über the Automorphismengruppen von Banachmannigfeltigkeiten mit invarianter Metrik, Math. Ann. 223 (1976), 279-288.
- Vigué, J. P., Le groupe des automorphismes analytiques d'un domaine borné d'un espace de Banach complexe, Ann. Sci. Ecole Norm. Sup. s'erie IV 9 (1976), 203-282.
- Vigué, J. P., Le groupe des automorphismes analytiques d'un domaine borné cerclé, Bull. Sc. Math. 2 s'erie 106 (1982), 417-426.

 $E\text{-}mail\ address:\ \texttt{jmisidro@zmat.usc.es}$