## LATTICES IN R<sup>2</sup> AND FINITE SUBSETS OF A CIRCLE

JACOB MOSTOVOY

ABSTRACT. An elementary geometric construction is used to relate the space of lattices in  $\mathbb{R}^2$  to the space  $\exp_3 S^1$  of the subsets of a circle of cardinality at most 3. As a consequence we obtain a new proof of known statements about the space  $\exp_3 S^1$ .

**Spaces of finite subsets.** For X a topological space let  $\exp_k X$  be the set of all finite subsets of X of cardinality at most k. There is a map from the Cartesian product of k copies of X with itself to  $\exp_k X$  which sends  $(x_1, \ldots, x_k)$  to  $\{x_1\} \cup \ldots \cup \{x_k\}$ . The quotient topology gives  $\exp_k X$  the structure of a topological space. Clearly,  $\exp_1 X = X$  for any X.

The simplest non-trivial example is provided by the space  $\exp_2 S^1$  which is easily seen to be homeomorphic to the Möbius band. The space  $\exp_3 S^1$  is described by a well-known theorem of R. Bott:

**Theorem 1.** (Bott, [1]) The space  $\exp_3 S^1$  is homeomorphic to a 3-sphere  $S^3$ .

The homeomorphism between  $\exp_3 S^1$  and  $S^3$  is rather non-obvious. This is illustrated by the following result which is, apparently, due to E. Shchepin. Consider the canonical embedding  $\Delta : S^1 \to \exp_3 S^1 = S^3$  which sends a point  $x \in S^1$  to the subset  $\{x\} \in \exp_3 S^1$ .

**Theorem 2.** (Shchepin, [unpublished]) The map  $\Delta : S^1 \to S^3$  is a trefoil knot.

In his proof of Theorem 1 Bott used a "cut-and-paste" argument. Shchepin's proof of Theorem 2 was based on a direct calculation of the fundamental group of  $S^3 \setminus \Delta(S^1)$ . The purpose of this note is to show that the above theorems can be deduced from well-known facts about lattices in a plane.

**Spaces of lattices.** A lattice in  $\mathbf{R}^2$  is a subgroup of the vector space  $\mathbf{R}^2$  generated by two linearly independent vectors. The set  $\mathcal{L}$  of all lattices considered up to multiplication by a non-zero real number can be identified with  $SL(2, \mathbf{R})/SL(2, \mathbf{Z})$  and will be considered with this topology.

The space  $\mathcal{L}$  can be compactified by adding points which correspond to degenerate lattices, that is, subgroups of  $\mathbb{R}^2$  generated by one non-zero vector. Degenerate lattices modulo multiplication by a non-zero real number form a circle  $S^1$ . The topology on the union  $\widehat{\mathcal{L}} = \mathcal{L} \cup S^1$  can be described as follows. Denote by  $\widehat{T}$  the space of triangles of perimeter 1 in  $\mathbb{R}^2$  each of whose angles does not exceed  $\pi/2$ and is possibly equal to zero. The space  $\widehat{T}$  contains degenerate triangles which have one side equal to 0 and both angles adjacent to it equal to  $\pi/2$ . Let  $T \subset \widehat{T}$  be the subspace of non-degenerate triangles. There is a map  $p: \widehat{T} \to \widehat{\mathcal{L}}$  which sends

Date: August 3, 2018.

<sup>1991</sup> Mathematics Subject Classification. Primary 11H06, 54B20; Secondary 57M25.

## J.MOSTOVOY

a triangle to the lattice generated by any two of its sides considered as vectors. Notice that the restriction of p to T is a continuous map from T onto  $\mathcal{L}$ . We define the topology on  $\widehat{\mathcal{L}}$  as the quotient topology with respect to the map p.

**Theorem 3.** The compactification  $\widehat{\mathcal{L}}$  of the space of lattices  $\mathcal{L}$  is homeomorphic to a 3-sphere  $S^3$ . The inclusion  $\widehat{\mathcal{L}} \setminus \mathcal{L} \to \widehat{\mathcal{L}}$  is a trefoil knot.

The proof of the above theorem (due to D.Quillen) can be found in [2] on page 84 where a homeomorphism between  $\mathcal{L}$  and the complement of a trefoil is constructed. This map can be easily seen to extend to a homeomorphism between  $\widehat{\mathcal{L}}$  and  $S^3$ .

Main theorem. Our main result relates Theorem 3 to Theorems 1 and 2.

**Theorem 4.** There is a homeomorphism  $\Phi : \widehat{\mathcal{L}} \to \exp_3 S^1$  which identifies the circle  $\widehat{\mathcal{L}} \setminus \mathcal{L}$  with  $\Delta(S^1)$ .

*Remark.* The circle  $S^1$  acts as a subgroup of  $PSL(2, \mathbf{R})$  on  $\mathcal{L}$  (and, in fact, on  $\widehat{\mathcal{L}}$ ) rotating the lattices. It will be clear from the construction below that  $\Phi$  sends the action of  $S^1$  on  $\widehat{\mathcal{L}}$  to the action of SO(2) by rotations on  $\exp_3 S^1$ .

*Proof.* For any non-degenerate lattice L there is a finite number of triangles  $V_i^L$  with vertices in L whose sides are generators of L and whose angles do not exceed  $\pi/2$ . There are 12 such triangles for a rectangular lattice and 6 for any other lattice, as shown in the figure. For each non-degenerate lattice L choose one of the triangles



 $V_i^L$  and trace 3 lines which connect the vertices of the triangle with the centre of its circumscribed circle. These lines are distinct unless L is rectangular in which case two of them coincide. Each line determines a point in  $\mathbf{RP}^1 = S^1$  so we obtain a point  $\Phi(L)$  in  $\exp_3 S^1 \setminus \Delta(S^1)$  for each  $L \in \mathcal{L}$ . It is clear that  $\Phi(L)$  depends only on L and not on a particular choice of a triangle. Now, for a degenerate lattice L define  $\Phi(L)$  as the point of  $\mathbf{RP}^1$  which corresponds to the line containing L. A straightforward check shows that  $\Phi$  is a homeomorphism.

Acknowledgments. I would like to thank Evgenii Shchepin, Abdelghani Mouhallil and Alberto Verjovsky for useful discussions.

## References

- [1] R. Bott, On the third symmetric potency of  $S_1$ , Fund. Math. **39** (1952), 264–268.
- [2] J. Milnor, Introduction to algebraic K-theory, Princeton, 1971.

INSTITUTO DE MATEMÁTICAS (UNIDAD CUERNAVACA), UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, A.P. 273 ADMON. DE CORREOS # 3, C.P. 62251, CUERNAVACA, MORELOS, MEXICO *E-mail address*: jacob@matcuer.unam.mx