

Integrable discretization of sine- and sinh-Gordon equations: Novelty and open problems

M. Boiti, F. Pempinelli, B. Prinari and A. Spirez

Dipartimento di Fisica dell'Università and Sezione INFN, 73100 Lecce, Italy

Physique Mathématique et Théorique, CNRS-UMR 5825, Université

Montpellier 2, 34095 Montpellier, France

Abstract. The inverse scattering theory for the discretized sine- and sinh-Gordon equations is considered. In the case in which only the space is discretized some difficulties of the inverse scattering theory related to the traditional Lax pair proposed by Hirota and Okamoto are overcome by introducing a Lax triplet and by using the inverse scattering transform of the so-called exactly discretized Schrödinger operator. The method works also in the case in which both space and time are discretized but only for the sinh-Gordon equation. The problem of building an inverse scattering theory for the doubly discrete sine-Gordon equation in the form proposed by Hirota remains open.

1. Introduction

In the framework of integrable nonlinear evolution equations, Hirota recovered almost 25 years ago the Bianchi superposition formula for the solutions of the sine-Gordon equation, showing that it can be considered as an integrable doubly discrete (both in space and time) version of the sine-Gordon equation [1]. He found also its Lax pair, Backlund transformations and N -soliton solutions. In the following this study was generalized to include the discrete (in space or time) case [2], soliton solutions were obtained by using the dressing method [3] and the related inverse spectral transform was studied [4].

Here we are interested in the spectral transform for the discrete and doubly discrete sine-Gordon and sinh-Gordon equations with solutions decaying at space infinity.

We show that the inverse spectral transform for the semi-discrete case studied in [4] has an unexpected drawback. In order to overcome it we use as principal spectral problem the "exact discretization" of the Schrödinger operator introduced by Shabat

$$g_{n+2} = g_n g_{n+1} (1 + \epsilon) g_n \quad (1.1)$$

and obtained iterating the Darboux transformations [5].

The spectral theory for this operator was exhaustively studied in [6] in the case of a potential g_n , not necessarily real, satisfying

$$g_{n+1} = \epsilon g_n \quad (1 + \epsilon) g_n = 1 \quad \text{for } j < 1 : \quad (1.2)$$

and applied to a discrete version of the KdV.

Following a procedure analogous to that used in [7] for getting a 2+1 generalization of the sine- and sinh-Gordon equations, we do not consider a Lax pair but a Lax triplet, i.e. we derive the discrete sine- and sinh-Gordon equations as compatibility conditions of (1.1) with a pair of auxiliary spectral problems.

The evolution of spectral data is easily derived and, therefore, according to the usual inverse scattering scheme, the Cauchy initial value problem is solved and the drawback in [4] is overcome. However, one must be advised that the solutions of the discrete sine- and sinh-Gordon equations are related to the potential g_n via a continued fraction, which can be considered a discretized version of a Riccati equation.

In the sinh-Gordon case the potential g_n is real and the characterization equations for the spectral data are easily obtained in analogy with the continuous case. In the sine-Gordon case the potential g_n is complex and the problem (most surely not trivial) of characterizing the spectral data is left open.

We are also able to find the couple of doubly discrete auxiliary spectral operators that in triplet with the doubly discrete Schrodinger operator gives a difference-difference evolution equation in a field ψ_{nm} . The reduction ψ_{nm} real is the only evident admissible reduction and in this case in the limit of continuous time we recover the differential-difference sinh-Gordon equation. The equation is of sixth order and we were not able to verify if it can be integrated and, then, if another reduction is admissible such that one recovers the Hirota-Bianchi fourth order doubly discrete sine-Gordon equation.

Therefore the problem of solving the Hirota-Bianchi doubly discrete equation by using the inverse scattering method is left open.

Finally let us remark that the origin of the difficulties both in our approach and in the approach of [4] may be related to the fact that we are trying to describe solutions of the Hirota-Bianchi equation, that have instabilities as discussed in [8, 9], by using standard methods of inverse scattering that are not able to describe such instabilities.

2. Standard Lax Pair

In the following we found convenient to use, for any discrete function $f(n; m)$, the notation

$$f_{ij} = f(n+i; m+j) \quad (2.1)$$

when n and m are considered generic but fixed.

Let us consider the Lax pair proposed by Orfanidis [2]

$$\psi_{10} = M \psi_{00} \quad (2.2)$$

$$\psi_{01} = N \psi_{00} \quad (2.3)$$

with

$$M_{00} = \begin{pmatrix} e^{i(\psi_{10} - \psi_{00})=2k} & k \\ k & e^{i(\psi_{10} - \psi_{00})=2} \end{pmatrix} \quad (2.4)$$

$$N_{00} = \begin{pmatrix} 1 & \bar{k} e^{i(\psi_{01} - \psi_{00})=2} \\ \bar{k} e^{i(\psi_{01} - \psi_{00})=2} & 1 \end{pmatrix} \quad (2.5)$$

with $(n; m)$ real, k the spectral parameter and α a real constant. The compatibility condition

$$F_{10} = E_{01}$$

where E shifts n and F shifts m gives by inserting in it (2.2) and (2.3)

$$M_{01}N_{00} = N_{10}M_{00}:$$

From it we have the Bianchi-Hirota equation

$$^2 \sin \frac{1}{4} (\varphi_{11} + \varphi_{10} + \varphi_{01} + \varphi_{00}) = \sin \frac{1}{4} (\varphi_{11} - \varphi_{10} - \varphi_{01} + \varphi_{00}):$$

If we introduce

$$\begin{aligned} i_j &= (x + h; t + j) \\ x &= nh; \quad t = m; \end{aligned}$$

choose the constant h as follows

$$^2 = \frac{h}{4} \quad (2.6)$$

and take the limit $h \rightarrow 0$, $\varphi \rightarrow 0$ and $n \rightarrow 1, m \rightarrow 1$ at x and t fixed we get the sine-Gordon equation in the light cone coordinates

$$\varphi_{xt} = \sin \varphi: \quad (2.7)$$

If we make only the limit $h \rightarrow 0$ we have

$$\varphi_{1t} - \varphi_{0t} = h \sin \frac{1}{2} (\varphi_{11} + \varphi_{00}) \quad (2.8)$$

and an analogous equation if $h \rightarrow 0$.

Notice that in both cases, doubly discrete and space discrete cases, the principal spectral problem (2.2) is the same.

If in (2.2) we perform the following k dependent transformation

$$(n) = c(k)^n (1 - i_2) (n) \quad (2.9)$$

where the dependence on m is understood, i_2 is the Pauli matrix and

$$c(k) = \frac{p}{1 - k^2} \quad (2.10)$$

we recover the spectral problem considered in [4]

$$(n+1) = [U + A(n)] (n) \quad (2.11)$$

with

$$U = \begin{pmatrix} z & 0 \\ 0 & 1/z \end{pmatrix}; \quad A(n) = \begin{pmatrix} z & \frac{1}{z} \\ \frac{1}{z} & z \end{pmatrix} \mathcal{A}(n) \quad (2.12)$$

$$\mathcal{A}(n) = \frac{1}{2} \begin{pmatrix} \cos \varphi(n) & 1 \\ i \sin \varphi(n) & \cos \varphi(n) \end{pmatrix} \quad (2.13)$$

and where

$$z = \frac{p}{1 - k} \quad (2.14)$$

is the new spectral parameter and

$$(n; m) = \frac{1}{2} (\varphi(n+1) - \varphi(n)): \quad (2.15)$$

In terms of

$$(n) = (n) U^{-n} \quad (2.16)$$

(2.11) becomes

$$(n+1)U = (U + A(n)) \phi(n); \quad (2.17)$$

We consider the case when $\phi(n)$ is going rapidly to zero as $n \rightarrow \pm 1$. Then $\phi(n) \rightarrow 0$ as $n \rightarrow \pm 1$.

As usual, the vector Jost solutions are defined by fixing their behaviour for $n \rightarrow +1$ or for $n \rightarrow -1$. It is easy to check that the matrices Φ and Ψ defined by the integralequations

$$\Phi(n) = 1 + \sum_{j=n}^{\infty} U^{n-j-1} A(j) \phi(j) U^{j-n} \quad (2.18)$$

$$\Psi(n) = 1 + \sum_{j=-1}^{\infty} U^{n-j-1} A(j) \phi(j) U^{j-n} \quad (2.19)$$

solve (2.17) and their columns have the required behaviours, respectively, for $n \rightarrow +1$ and for $n \rightarrow -1$.

If we write the solution to (2.19) in the form of the following Neumann series

$$\Psi(n) = \sum_{l=0}^{\infty} \Psi^{(l)}(n) \quad (2.20)$$

$$\Psi^{(0)}(n) = 1 \quad (2.21)$$

$$\Psi^{(l+1)}(n) = \sum_{j=-1}^{\infty} U^{n-j-1} A(j) \Psi^{(l)}(j) U^{j-n} \quad (2.22)$$

we have

$$\Psi^{(l)}(n) = \sum_{j_1=-1}^{\infty} U^{n-j_1-1} A(j_1) \sum_{j_2=j_1-1}^{\infty} U^{j_1-j_2-1} A(j_2) \cdots \sum_{j_l=-1}^{\infty} U^{j_{l-1}-j_l-1} A(j_l) U^{j_l-n};$$

This means that

$$\Psi^{(l)}(n) = \sum_{j_1=-1}^{\infty} \mathcal{A}(j_1) \sum_{j_2=j_1-1}^{\infty} \mathcal{A}(j_2) \cdots \sum_{j_l=-1}^{\infty} \mathcal{A}(j_l)$$

where we introduced

$$\mathcal{A}(n; z) = U^{-1}(z) A(n; z) = \begin{pmatrix} 1+z^2 & 0 \\ 0 & z^2+1 \end{pmatrix} \phi(n);$$

We consider the Neumann series in the circular annulus $D_R = \{z \in \mathbb{C} : 1/R < |z| < R\}$ with $R > 1$. We have

$$\mathcal{A}(n; z) = R^2 q(n)$$

with

$$q(n) = 2 \phi(n)$$

and we require

$$\sum_{j=-1}^{\infty} q(j) < +1;$$

Then we have

$$S^{(l)}(n) = S^{(l)}(n)$$

where

$$S^{(l)}(n) = R^{2l} \prod_{j_1=1}^{X-1} q(j_1) \prod_{j_1=1}^{X-1} q(j_1-1) \cdots \prod_{j_1=1}^{X-1} q(j_1):$$

One can prove by induction that

$$S^{(l)}(n) = \frac{1}{l!} R^{2l} Q(n)^{-1}$$

where

$$Q(n) = \prod_{j=1}^{X-1} q(j):$$

Therefore

$$S^{(l)}(n) = \frac{1}{l!} R^{2l} Q(n)^{-1};$$

that is

$$k(n)k = e^{R^2 Q(n)}$$

and we conclude that in D_R the Neumann series (2.20) is uniformly convergent.

Analogously, one can prove that the following Neumann series for

$$\begin{aligned} (n) &= \prod_{l=0}^{X-1} (l-1)^{l-1} (n) \\ (0) &= 1 \\ (l+1) &= \prod_{j=n}^{X-1} U^{n-j-1} A(j)^{(l)}(j) U^{j-n} \end{aligned}$$

is uniformly convergent in the same region with

$$k(n)k = e^{R^2 P(n)}$$

where

$$P(n) = \prod_{j=n}^{X-1} q(j):$$

We conclude that the matrix Jost solutions and are analytic in the complex plane with two singularities at $z = 0$ and at $z = 1$.

Writing (2.18)-(2.19) explicitly we have

$$\begin{aligned} \begin{pmatrix} 11(n) \\ 21(n) \end{pmatrix} &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ &+ \prod_{j=1}^{n-1} \left(1 + \frac{1}{z^2} (A_{11}(j) 11(j) + A_{12}(j) 21(j)) \right) \\ &+ \prod_{j=1}^{n-1} \frac{1}{z^{2(j-n+1)}} \left(1 + \frac{1}{z^2} (A_{21}(j) 11(j) + A_{22}(j) 21(j)) \right) \end{aligned} \quad (2.23)$$

$$\begin{aligned} \begin{pmatrix} 12(n) \\ 22(n) \end{pmatrix} &= \begin{pmatrix} 0 \\ 1 \end{pmatrix} \\ &+ \prod_{j=1}^{n-1} \frac{1}{z^{2(n-j-1)}} \left(1 + z^2 (A_{11}(j) 12(j) + A_{12}(j) 22(j)) \right) \\ &+ \prod_{j=1}^{n-1} \left(1 + z^2 (A_{21}(j) 12(j) + A_{22}(j) 22(j)) \right) \end{aligned} \quad (2.24)$$

and

$$\begin{aligned} \begin{pmatrix} 1 \\ 2 \end{pmatrix}(n) &= \begin{pmatrix} 11(n) \\ 21(n) \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ &+ \sum_{j=n}^{P+1} \begin{pmatrix} 1 + \frac{1}{z^2} & (\Phi_{11}(j) \quad 11(j) + \Phi_{12}(j) \quad 21(j)) \\ z^{2(j-n)} & 1 + z^2 (\Phi_{21}(j) \quad 11(j) + \Phi_{22}(j) \quad 21(j)) \end{pmatrix} \end{aligned} \quad (2.25)$$

$$\begin{aligned} \begin{pmatrix} 2 \\ 2 \end{pmatrix}(n) &= \begin{pmatrix} 12(n) \\ 22(n) \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \\ &+ \sum_{j=n}^{P+1} \begin{pmatrix} z^{2(n-j)} & 1 + \frac{1}{z^2} (\Phi_{11}(j) \quad 12(j) + \Phi_{12}(j) \quad 22(j)) \\ 1 + z^2 & (\Phi_{21}(j) \quad 12(j) + \Phi_{22}(j) \quad 22(j)) \end{pmatrix} \end{aligned} \quad (2.26)$$

Taking into account the result obtained for the analytic properties of Φ and Ψ , just by looking to the explicit form of the above discrete integral equations defining the columns of Φ and Ψ we deduce that the Jost solutions ψ_1 and ψ_2 have, respectively, an essential singularity at $z = 1$ and at $z = 0$, while the Jost solutions ϕ_1 and ϕ_2 have an essential singularity both at $z = 0$ and at $z = 1$. Therefore one cannot use the usual procedure for building the inverse scattering theory.

One could come back to the original Lax operator (2.4) and use the spectral parameter k , but then the potential at large n is going to the constant unity matrix. Therefore, we think that the encountered difficulty is intrinsic to the problem, since it is well known that when the potential in the spectral problem is not going to zero at large distances the usual scheme of the inverse scattering method must be accordingly modified.

In the following section we try to overcome the difficulty by using a different principal spectral problem with a potential going to zero at large n .

3. Lax triplet

3.1. Semi-discrete case

Let us consider the following semi-discrete spectral problems

$$L = g_0 L_1 (1 + \epsilon) L_2 \quad (3.1)$$

$$L_1 t = L_0 L_1 t + L_2 L_2 \quad (3.2)$$

where ϵ plays the role of spectral parameter. The compatibility condition $L_1 L_2 = E L_1 t$; where E is the shift operator, $E v_n = v_{n+1}$, gives

$$(L_1 L_0 - L_0 g_0 + 1 + \epsilon) L_1 t = (L_0 g_0 - L_1 L_2) L_1 t + (g_0 L_2 - L_1) L_1 t; \quad (3.3)$$

which can be regarded a third Lax operator to be added to (3.1) and (3.2). In contrast with the usual case we have not a Lax pair but a Lax triplet. The compatibility conditions for this triplet of spectral problems are closed by requiring that the spectral problem

$$(L_2 L_1 - L_1 g_1 + 1 + \epsilon) L_1 t = (L_1 g_1 - L_2 L_1) L_1 t + (g_1 L_2 - L_1) L_1 t \quad (3.4)$$

obtained by shifting (3.3) and using (3.1) is compatible with (3.2). We have

$$\begin{aligned} (L_1 L_0 - L_0 g_0 + 1 + \epsilon) f[L_1 g_1 - L_2 L_1 + g_0 (g_1 L_2 - L_1)] L_1 t &= (L_2 L_1 - L_1 g_1 + 1 + \epsilon) \\ &\quad f(L_0 g_0 - L_1 L_2) L_1 t + (g_0 L_2 - L_1) L_1 t + (L_1 L_0 - L_0 g_0 + 1 + \epsilon) L_1 t \end{aligned}$$

Assuming g_0 and g_1 to be linearly independent, and equating the coefficients of the different powers in t we get the following relations

$$g_{1t} = 2g_0 \quad (3.5)$$

$$2g_1 = g_0 \quad (3.6)$$

$$(g_0 + 1) = (2g_1 + 1)(g_0 + 1) + (g_0 + 1)(2g_1 + 1) \quad (3.7)$$

and

$$g_{1t} = 2g_0 \quad (3.8)$$

$$(g_0 + 1)(g_{1t} - 2g_0) = 0 \quad (3.9)$$

where we used (3.5) in (3.6) and (3.8) in (3.9). Note that, taking into account (3.6) and (3.8), (3.7) and (3.9) are identically satisfied and we are left with

$$g_{1t} = 2g_0 \quad (3.10)$$

$$g_0 = 1 \quad (3.11)$$

$$g_0 = 2g_1 + 1 \quad (3.12)$$

where, without loss of generality, the constant of integration of the second equation was scaled to 1. The last equation can be rewritten as

$$g_1 = 2g_0 - 1$$

and, then, integrated to

$$g_1 = 2g_0 - 1 = c \quad (3.13)$$

In order to get the discrete sine-Gordon equation the choice of the constant c of integration in (3.13) is crucial. The study of the continuous limit suggests to choose this constant to be zero. From (3.11) we get

$$g_0 = 1 + \frac{1}{g_0} \quad (3.14)$$

Notice that this equation, which can be considered as a discrete Riccati equation, can be solved for g_0 in terms of a continuous fraction of g (see [10]).

Then we substitute g_0 into (3.13) with $c = 0$ and we get

$$g_0 = \frac{0}{1} \quad (3.15)$$

If we put

$$g_0 = e^{q_0}; \quad (3.16)$$

we get from (3.15) and (3.14)

$$g_0 = e^{\frac{1}{2}(q_0 - q_{-1})} \quad (3.17)$$

$$g_0 = e^{\frac{1}{2}(q_{-1} - q_0)} + e^{\frac{1}{2}(q_1 - q_0)} \quad (3.18)$$

and the evolution equation (3.10) takes the form

$$\frac{1}{2}(q_1 - q_0)_t e^{\frac{1}{2}(q_1 - q_0)} + \frac{1}{2}(q_2 - q_1)_t e^{\frac{1}{2}(q_2 - q_1)} = e^{q_2} - e^{q_0} :$$

We multiply both sides by e^{q_1} getting

$$\frac{1}{2}(q_1 - q_0)_t e^{\frac{1}{2}(q_1 + q_0)} + e^{q_1 + q_0} = \frac{1}{2}(q_2 - q_1)_t e^{\frac{1}{2}(q_1 + q_2)} + e^{q_1 + q_2}$$

and then it can be integrated with respect to the spatial variable giving

$$\frac{1}{2} (q_1 - q_0)_t e^{\frac{1}{2} (q_1 + q_0)} = e^{q_1 + q_0} + \text{const.}$$

so that if we choose the constant of integration equal to 1 we finally get

$$(q_1 - q_0)_t = 4 \sinh \frac{q_1 + q_0}{2}$$

which for q real is the discrete sinh-Gordon equation and for q purely imaginary is the discrete sine-Gordon equation.

Note that for $q(n)$ vanishing at large n we have from (3.18) that $g(n) \rightarrow 2$ and the spectral theory developed in [6] is applicable.

3.2. Evolution of spectral data

The evolution of the spectral data can be determined by considering the first and the third Lax operators, i.e. the spectral problem (3.1) and (3.4).

In order to meet the notations used in [6] we introduce the new spectral k and the new eigenfunction as follow

$$k = k^2 \quad (3.19)$$

$$(n; k) = (1 + ik)^n (n; k): \quad (3.20)$$

The principal spectral problem (3.1) reads

$$(1 + ik) (n + 2) + (1 - ik) (n) = g(n) (n + 1) \quad (3.21)$$

while the auxiliary spectral problem (3.4) thanks to (3.10) and (3.11) becomes

$$k^2 \psi_t = A_{1-1} + (1 - ik) \psi_0 \quad (3.22)$$

where

$$A_1 = \begin{pmatrix} g_1 & 0 \\ 0 & g_0 \end{pmatrix}.$$

We need the Jost solution $\psi^+(n; k)$ introduced in [6], which is analytic in the upper half plane and is defined via the following discrete integral equation

$$\psi^+(n; k) = 1 + \frac{1}{2ik} \sum_{j=n+1}^{\infty} \frac{1 + ik}{1 - ik} (g(j-1) - 2) \psi^+(j; k) \quad (3.23)$$

and which has the following asymptotic behaviour

$$\lim_{n \rightarrow +\infty} \psi^+(n; k) = 1; \quad k = 0 \quad (3.24)$$

at large positive n and

$$\psi^+(n; k) \sim a^+(k) + \frac{1 - ik}{1 + ik} b^+(k); \quad k = 0 \quad (3.25)$$

at large negative n , where $a^+(k)$ is the inverse of the transmission coefficient and $b^+(k)$ is the reflection coefficient.

Then we look for a solution of (3.22) of the form

$$\psi(t; n; k) = C(t; k) \psi^+(t; n; k) \quad (3.26)$$

If we substitute (3.26) into (3.22), we get

$$k^2 C_t C^{-1} \psi^+ + \psi^+_{1t} = A_{1-1} \psi^+ + (1 - ik) \psi^+_0: \quad (3.27)$$

As $j \rightarrow \infty$, $g \rightarrow 0$; $1 \rightarrow 0$ and then $A_1 \rightarrow 1$ so that from (3.27), taking into account the asymptotic behavior of ψ for $n \rightarrow \infty$ we get, first,

$$C_t C^{-1} = \frac{i}{k}$$

and, then, by considering the limit for $n \rightarrow \infty$ ($k_- = 0$) the evolution equation for the spectral data

$$\begin{aligned} a_t(k) &= 0 \\ k^2 b_t(k) &= 2ikb(k); \end{aligned}$$

which can be trivially integrated to

$$\begin{aligned} a(t; k) &= a_0(k) \\ b(t; k) &= b_0(k) \exp \frac{2i}{k} t : \end{aligned} \quad (3.28)$$

3.3. Doubly discrete case

Let us consider now the doubly discrete version of (3.1)–(3.2), that is

$$g_{00} = g_{00} (1 + \epsilon) \quad (3.29)$$

$$g_{11} = g_{00} g_{01} + (g_{00} - g_{01}) g_{00} + g_{10} : \quad (3.30)$$

The compatibility condition $E g_{11} = F g_{20}$, where E is the shift operator with respect to the space variable, that is $E v_{n,m} = v_{n+1,m}$; and F is the time-shift operator, $F v_{n,m} = v_{n,m+1}$, gives, thanks to (3.30),

$$[A_{00} + \epsilon] g_{01} = [g_{01} (g_{00} - g_{10}) + A_{00} + \epsilon] g_{00} [g_{10} + g_{00} - g_{01}] g_{10} \quad (3.31)$$

where we introduced the notation

$$A_{00} = g_{10} g_{00} - g_{01} g_{00} + 1 : \quad (3.32)$$

This spectral problem can be considered defining the third Lax operator. The compatibility condition for the Lax triplet is obtained by imposing $g_{11} = E g_{01}$, that is, taking into account (3.29)–(3.31)

$$\begin{aligned} [A_{00} + \epsilon] [g_{11} (g_{00} - g_{20}) + A_{10} + \epsilon] g_{10} \\ = [A_{00} + \epsilon] [g_{10} (g_{01} - g_{11}) + g_{00} g_{10} (1 + \epsilon) g_{00}] \\ = g_{00} [A_{10} + \epsilon] [g_{01} (g_{00} - g_{10}) + A_{00} + \epsilon] g_{00} \\ = g_{00} [A_{10} + \epsilon] [g_{10} + g_{00} - g_{01}] g_{10} + [A_{00} + \epsilon] [A_{10} + \epsilon] g_{10} \\ + (g_{00} - g_{01}) [A_{00} + \epsilon] [A_{10} + \epsilon] g_{00} \end{aligned}$$

which involves only g_{00} and g_{10} . We assume that they are linearly independent and equate the coefficients of the different powers of ϵ . We get from the coefficient of ϵ^0

$$g_{11} - g_{10} = g_{20} - g_{00} \quad (3.33)$$

$$(g_{20} + g_{10} - g_{11}) (A_{00} + 1) = g_{00} (A_{10} + 1) \quad (3.34)$$

$$(g_{20} + g_{10} - g_{11}) A_{00} = g_{00} A_{10} : \quad (3.35)$$

The first equation is the searched evolution equation. Inserting it into the two following equations we get $A_{00} = A_{10}$ which can be integrated to

$$A_{00} = d^2 \quad (3.36)$$

with d an arbitrary constant. From the coefficient of g_{10} we get

$$g_{11} - 2g_{00} + g_{10} - g_{11}) + g_{00} (g_{10} + g_{00} - g_{01}) = 0 \quad (3.37)$$

$$(g_{11} - 2g_{00})A_{00} - g_{00}A_{00} (g_{10} + g_{11}) + A_{10} (g_{10} + g_{00} - g_{01}) = 0 \quad (3.38)$$

which, thanks to (3.33) and (3.36), reduce to only one equation that can be written as

$$g_{00} + g_{11} - 2g_{00} - 2g_{10} = g_{00} + g_{01} - g_{00} - g_{10} - g_{00}: \quad (3.39)$$

It can be integrated to

$$g_{00} + g_{01} - g_{00} - g_{10} - g_{00} = c \quad (3.40)$$

with c an arbitrary constant.

In conclusion from (3.36) we get

$$g_{01} = g_{10} + \frac{1}{g_{00}} d^2 \quad (3.41)$$

and inserting it into (3.40)

$$g_{00} = \frac{1}{2} g_{00} - \frac{c}{2g_{10}} + \frac{B_{00}}{2g_{10}} \quad (3.42)$$

where

$$B_{00} = \frac{p}{(g_{00} - g_{10} - c)^2 + 4(1 - d^2)g_{00} - g_{10}}: \quad (3.43)$$

From (3.33), (3.41) and (3.42) noting that

$$\frac{1}{g_{00}} d^2 = \frac{1}{2} g_{10} + \frac{c}{2g_{00}} + \frac{B_{00}}{2g_{00}} \quad (3.44)$$

we get

$$g_{01} = \frac{1}{2} (g_{10} - g_{10}) + \frac{B_{10} + B_{00}}{2g_{00}} \quad (3.45)$$

and, therefore, the following evolution equation in the doubly discrete field

$$g_{20} - g_{00} + g_{21} - g_{01} = \frac{B_{20} + B_{10}}{g_{10}} - \frac{B_{21} + B_{11}}{g_{11}} \quad (3.46)$$

In analogy with the semi-discrete case we can put

$$g(n; m) = e^{g(n; m)} \quad (3.47)$$

and rewrite the evolution equation in terms of $q(n; m)$. The real parameter d takes into account that the time can be scaled and it is necessary for performing the continuous limit in time. We consider solutions $q(n; m)$ decaying at large n . In order to be able to apply the spectral theory of the discrete Schrodinger equation the constants of integration c and d must be chosen in such a way that for $n \rightarrow 1$ the potential $g(n; m) \rightarrow 2$. Precisely, we get

$$c = \frac{1}{2} - 2d: \quad (3.48)$$

The evolution equation (3.46) admits the reduction $q = \bar{q}$ but not the reduction $q = \bar{q}$ admitted in the semi-discrete case. If we choose $d = 0$ and let $q(n; t) = q(n; m)$, in the limit $m \rightarrow 0$ we recover the semi-discrete sinh-Gordon equation.

3.4. Evolution of spectral data

Let us rewrite the auxiliary spectral problem (3.31) by using (3.33), (3.41) and the spectral parameter $k^2 = \lambda$ introduced in (3.19). We have

$$d^2 + k^2 \psi_{01} = \psi_{00} \frac{1}{\psi_{00}} d^2 + d^2 + k^2 \psi_{00} \quad (10 \quad 10):$$

It is convenient now to introduce the doubly discrete $(n; m)$ equivalent to (3.20) and write

$$d^2 + k^2 \psi_{01} = \psi_{00} \frac{1}{\psi_{00}} d^2 + d^2 + k^2 \psi_{00} \quad (1 + ik) \quad (10 \quad 10):$$

We look for a solution of the form

$$\psi(n; m; k) = \psi(m; k) + \psi(n; m; k) \quad (3.49)$$

where $\psi(m; k)$ is the Jost solution introduced in [6]. We have

$$\begin{aligned} (d^2 + k^2) \psi(m+1; k) &= \psi(m; k) + \psi(n; m+1; k) \\ &= \psi(n; m) \frac{1}{\psi(n; m)} d^2 + d^2 + k^2 \psi(n; m; k) \\ &\quad (1 + ik) \psi(n-1; m) + \psi(n+1; m; k): \end{aligned} \quad (3.50)$$

Let us first note that thanks to (3.43), (3.44) and (3.48) we have

$$\lim_{|n| \rightarrow \infty} \psi(n; m) \frac{1}{\psi(n; m)} d^2 = (1 - d):$$

Then we proceed in the usual way evaluating the limit of (3.50) for $n \rightarrow \infty$ and $n \rightarrow -\infty$. Taking into account the asymptotic behaviours (3.24) and (3.25) we obtain, respectively, for $n \rightarrow \infty$ the time evolution of

$$\psi(m+1; k) = \psi(m; k) = 1 \quad \frac{1}{d - ik} \quad (3.51)$$

and for $n \rightarrow -\infty$ the time evolution of the spectral data ($k_- = 0$)

$$a(m+1; k) = a(m; k) \quad (3.52)$$

$$b(m+1; k) = \frac{(d - ik)(d + ik)}{(d + ik)(d - ik)} b(m; k): \quad (3.53)$$

Notice that for $d = 0$ in the limit $\lambda \rightarrow 0$ we recover as expected the time evolution of the spectral data for the sinh-Gordon equation.

Acknowledgments

This work was partially supported by COFIN 2000 "Sintesi"

References

- [1] R. Hirota, J. Phys. Soc. of Japan 43, 2079 (1977)
- [2] S. J. Orfanidis, Phys. Review D 18, 3828 (1978)
- [3] D. Levi, O. Ragnisco and M. Bruschi, Nuovo Cimento 58A, 56 (1980)
- [4] L. Pilloni, D. Levi, Phys. Lett. 92A, 5 (1982)
- [5] A. Shabat, in Nonlinearity, Integrability and All That. Twenty Years After NEEDS '79, M. Boiti, L. Martini, F. Pempinelli, B. Prinari and G. Soliani eds, pp.331-342, World Scientific Pub. Co., Singapore (2000).

- [6] M. Boiti, F. Pempinelli, B. Prinari and A. Spire, "An integrable discretization of the KdV at large times", *nlm SI/0101023*, January 2001
- [7] M. Boiti, J. Leon and F. Pempinelli, *Inverse Problems* 3, 37 (1987)
- [8] M. J. Ablowitz, B. M. Herbst and C. Schober, *J. of Comput. Phys.* 126, 299 (1996)
- [9] M. J. Ablowitz, B. M. Herbst and C. Schober, *J. of Comput. Phys.* 131, 354 (1997)
- [10] H. S. Wall, "Analytic Theory of Continued Fractions" (New York: Chelsea, 1967)