

TOPOLOGY OF THE REAL PART OF HYPERELLIPTIC JACOBIAN ASSOCIATED WITH THE PERIODIC TODA LATTICE

YUJI KODAMA[†]

ABSTRACT. This paper concerns the topology of the isospectral *real* manifold of the $\mathfrak{sl}(N)$ periodic Toda lattice consisting of 2^{N-1} different systems. The solutions of those systems contain blow-ups, and the set of those singular points defines a divisor of the manifold. Then adding the divisor, the manifold is compactified as the real part of the $(N-1)$ -dimensional Jacobi variety associated with a hyperelliptic Riemann surface of genus $g = N-1$. We also study the real structure of the divisor, and then provide conjectures on the topology of the affine part of the real Jacobian and on the gluing rule over the divisor to compactify the manifold based upon the sign-representation of the Weyl group of $\mathfrak{sl}(N)$.

1. PRELIMINARY

Let us start with an infinite Toda lattice which is defined as a hamiltonian system with the hamiltonian,

$$H = \frac{1}{2} \sum_{k \in \mathbb{Z}} P_k^2 + \sum_{k \in \mathbb{Z}} \epsilon_k \exp(Q_k - Q_{k+1}),$$

where $\epsilon_k \in \{\pm 1\}$. Because of the signs ϵ_k in the potential term, the solution blows up if at least one sign is negative [10]. With Flaschka's coordinates, $a_k := \epsilon_k \exp(Q_k - Q_{k+1})$ and $b_k := -P_k$, Hamilton's equation with the hamiltonian H is given by

$$(1.1) \quad \begin{cases} \frac{da_k}{dt} = a_k(b_{k+1} - b_k), \\ \frac{db_k}{dt} = a_k - a_{k-1}. \end{cases}$$

Remark 1.1. *The sign ϵ_k is invariant under the Toda flow, and the affine part of the isospectral manifold of the Toda flow consists of 2^{N-1} disconnected pieces labeled by the signs of a_k 's (note $\prod_{k=1}^N a_k$ is invariant).*

The infinite Toda lattice in $\{(a_k, b_k) : k \in \mathbb{Z}\}$ can be put in a matrix equation, the Lax formulation,

$$(1.2) \quad \frac{dL}{dt} = [L, A],$$

[†]) Department of Mathematics, Ohio State University, Columbus, OH 43210
E-mail address: kodama@@math.ohio-state.edu.

We assume that the discriminant of (1.8), $F(\lambda) := P(\lambda)^2 - 4 \prod_{j=1}^N a_j$, has all distinct roots, i.e.

$$(1.10) \quad F(\lambda) = \prod_{k=1}^{2N_R} (\lambda - \lambda_k) \prod_{l=1}^{N_I} ((\lambda - \nu_l)(\lambda - \bar{\nu}_l)).$$

with all different roots $\lambda_j \in \mathbb{R}, \nu_l \in \mathbb{C}$, and $N_R + N_I = N$. This implies that the algebraic curve having this condition is a smooth affine variety. We say that a matrix L_P is in the class (P) if the corresponding periodic matrix L satisfies this condition. The phase space for the periodic lattice is given by

$$Z_{\mathbb{R}}^P := \left\{ (a_1, \dots, a_N, b_1, \dots, b_N) \in \mathbb{R}^{2N} : L_P \text{ is in the class (P)} \right\}.$$

Since the $\mathfrak{sl}(N)$ Toda lattice is a hamiltonian system of degree N , its integrability can be shown by finding N independent and involutive integrals. Those integrals I_k are given by the coefficients of the characteristic equation (1.8) where $P(\lambda)$ is defined as

$$(1.11) \quad P(\lambda) = (-1)^N \left(\lambda^N + \sum_{k=1}^N (-1)^k I_k(L_P) \lambda^{N-k} \right),$$

and there is an additional integral obtained from the residue with respect to the spectral parameter z , $\prod_{i=1}^N a_i$, i.e.

$$\left\{ \begin{array}{l} I_1(L_P) = \sum_{i=1}^N b_i, \\ I_2(L_P) = \sum_{i>j} b_i b_j - \sum_{i=1}^N a_i, \\ \dots \\ I_N(L_P) = \Delta_{1,N}(0) - a_N \Delta_{2,N-1}(0) = \prod_{i=1}^N b_i + \dots, \\ I_{N+1}(L_P) = \prod_{i=1}^N a_i. \end{array} \right.$$

Then the isospectral set is defined by

$$Z_{\mathbb{R}}^P(\gamma) := \left\{ (a_1, \dots, a_N, b_1, \dots, b_N) \in Z_{\mathbb{R}}^P : I_k(L_P) = \gamma_k \in \mathbb{R}, k = 1, \dots, N+1 \right\}$$

which is the affine part of the compactified manifold $\hat{Z}_{\mathbb{R}}^P(\gamma)$ of $\dim Z_{\mathbb{R}}^P(\gamma) = N-1$, and with a divisor Θ associated to the blow-ups of $a_k, k = 1, \dots, N$, we have [2],

$$(1.12) \quad Z_{\mathbb{R}}^P(\gamma) = \hat{Z}_{\mathbb{R}}^P(\gamma) \setminus \Theta, \quad \text{with } \Theta = \bigcup_{k=1}^N \{a_k^{-1} = 0\}.$$

It turns out that the compact manifold $\hat{Z}_{\mathbb{R}}^P(\gamma)$ can be identified as the real part of the Jacobian \mathbb{C}^g/Γ with the lattice Γ defined by the period matrix Ω associated with the Riemann surface of genus g (see the next section for the details). The Riemann surface is determined by the spectral curve, and the genus is just $g = N-1$. We will discuss some details of this statement in the following sections. The main purpose in this paper is to find the topological structure of the divisor Θ in the real part of the Jacobian and to determine the gluing rule on the divisor to compactify the affine variety $Z_{\mathbb{R}}^P(\gamma)$. To study this, we start with a brief review of algebraic curve and the solution of periodic Toda lattice in the next section.

2. ALGEBRAIC CURVES AND PERIODIC TODA LATTICES

2.1. Riemann surface and Jacobian. The characteristic equation $\det(L_P - \lambda I) = 0$ determines a real plane curve \mathcal{C} (1-dimensional affine curve on \mathbb{C}^2);

$$(2.1) \quad y^2 = P(\lambda)^2 - 4 = \prod_{k=1}^{2N} (\lambda - \lambda_k),$$

where we have made a coordinate change from (λ, z) to (λ, y) with $y = 2z + P(\lambda)$. Also the complex conjugate pairs of the roots are denoted as $(\lambda_{N_R+2k-1} = \nu_k, \lambda_{N_R+2k} = \bar{\nu}_k)$ for $k = 1, \dots, N_I$ as in (1.10). Then a compact Riemann surface associated with the plane curve \mathcal{C} is obtained by adding points at infinity by introducing a second chart \mathcal{C}' ;

$$(y')^2 = \prod_{k=1}^{2N} (1 - \lambda_k \lambda')$$

which is glued to the chart \mathcal{C} with the isomorphism, $\lambda' = 1/\lambda, y' = y/\lambda^N$. This Riemann surface $\mathcal{R} = \mathcal{C} \cup \mathcal{C}'$ is called a hyperelliptic Riemann surface of genus $g = N - 1$, and it is topologically a compact surface with g handles. On the \mathcal{R} one can define a morphism $\pi : \mathcal{R} \rightarrow \mathbb{C}P^1$ by $(\lambda, y) \mapsto \lambda$ on the chart \mathcal{C} and similarly on the chart \mathcal{C}' . Note that the morphism π is a two-fold covering map on $\mathbb{C}P^1$ except at the branch points λ_k 's. The points at ∞ of \mathcal{R} , which are denoted as p_∞^\pm , are given by the points $(\lambda' = 0, y' = \pm 1)$, i.e.

$$(2.2) \quad \mathcal{R} = \mathcal{R}_0 \cup \{p_\infty^+, p_\infty^-\}.$$

where the \mathcal{R}_0 is the affine part of \mathcal{R} . On the \mathcal{R} , we also define a symplectic basis of $H_1(\mathcal{R}, \mathbb{Z})$ denoted as $\{(\alpha_j, \beta_j) : j = 1, \dots, g\}$, in which the paths α_j 's are disjoint from each other as are the paths β_j 's, and α_i, β_j intersect only if $i = j$ and then in one point.

Now let us define the *real* part of the Riemann surface, denoted as $\mathcal{R}_\mathbb{R}$. It is defined as the set of fixed points of the antiholomorphic involution obtained by conjugation, $(\lambda, y) \mapsto (\bar{\lambda}, \bar{y})$,

$$\mathcal{R}_\mathbb{R} := \left\{ (\lambda, y) \in \mathcal{R} : \lambda = \bar{\lambda}, y = \bar{y} \right\}.$$

Then it is easy to see:

Proposition 2.1. *Let \mathcal{R} be a compact (smooth) Riemann surface of genus g . Then the real part $\mathcal{R}_\mathbb{R}$ is isomorphic to N_R disjoint union of circles, $\mathbb{R}P^1 \cong S^1$,*

$$\mathcal{R}_\mathbb{R} \cong \overbrace{S^1 \sqcup \dots \sqcup S^1}^{N_R} \quad (\text{disjoint union}),$$

where $2N_R$ is the number of real zeros in (2.1), i.e. the real ramification points of \mathcal{R} .

On the Riemann surface \mathcal{R} of genus g , one can also have g holomorphic differentials (the Riemann-Roch theorem), which we denote $\omega_1, \dots, \omega_g$. We normalize those differentials with

$$(2.3) \quad \oint_{\alpha_i} \omega_j = \delta_{i,j}, \quad \text{for } i, j = 1, \dots, g.$$

Then the Jacobian $\text{Jac}(\mathcal{R})$ associated with the \mathcal{R} is defined as a complex torus with a lattice Γ of rank $2g$;

$$\text{Jac}(\mathcal{R}) = \mathbb{C}^g / \Gamma, \quad \Gamma := \{ \Omega m : m = (m_1, \dots, m_{2g})^t \in \mathbb{Z}^{2g} \},$$

where Ω is a $g \times 2g$ matrix, the period matrix, defined by

$$(2.4) \quad \Omega = (I \ B), \quad \text{with } B = \left(\oint_{\beta_i} \omega_j \right)_{1 \leq i, j \leq g},$$

where I is the $g \times g$ identity matrix. It turns out that our purpose of studying the topological structure of the isospectral manifold of periodic Toda lattice is just to study the topological structure of the real part of the Jacobian and its divisors corresponding to the blow-ups of the Toda flow.

2.2. Periodic Toda lattice. The Toda flow can be described as a linear flow on the Jacobian $\text{Jac}(\mathcal{R})$ of the Riemann surface \mathcal{R} . Let us first illustrate the solution of a simplest case with $N = 2$ ($g = 1$), and the general case will be discussed in the next section.

The Lax matrix L_P in this case is given by

$$L_P = \begin{pmatrix} b_1 & 1 + a_2 z^{-1} \\ z + a_1 & b_2 \end{pmatrix}$$

whose spectral curve defines an elliptic curve,

$$\det(L_P - \lambda I) = -\left(z + \frac{a_1 a_2}{z} - P(\lambda)\right) = 0.$$

The polynomial $P(\lambda)$ is given by

$$(2.5) \quad P(\lambda) = \lambda^2 + I_2, \quad \text{with} \quad I_2 = -b_1^2 - a_1 - \frac{1}{a_1},$$

where we have assumed $I_1 = b_1 + b_2 = 0$ and $I_3 = a_1 a_2 = 1$. Then the affine part of the isospectral manifold is given by

$$Z_{\mathbb{R}}^P(\gamma) = \left\{ (a_1, b_1) \in \mathbb{R}^2 : b_1^2 + a_1 + \frac{1}{a_1} = \gamma_2 \in \mathbb{R} \right\} \cong \begin{cases} S^1 \sqcup \mathbb{R} \sqcup \mathbb{R} & \text{if } \gamma_2 > 2, \\ \mathbb{R} \sqcup \mathbb{R} & \text{if } \gamma_2 < 2. \end{cases}$$

Thus we have

$$(2.6) \quad Z_{\mathbb{R}}^P(\gamma) \cong \mathcal{R}_{\mathbb{R}} \setminus \{p_{\infty}^+, p_{\infty}^-\} = \mathcal{R}_{\mathbb{R}} \cap \mathcal{R}_0.$$

Two disconnected pieces of \mathbb{R} are compactified to make a circle S^1 , that is, the corresponding solution blows up once in each point p_{∞}^+ or p_{∞}^- and total twice in one cycle. Thus the compactified manifold $\hat{Z}_{\mathbb{R}}^P(\gamma)$ is just $\mathcal{R}_{\mathbb{R}}$ (see below for more precise statement).

The Toda flow is determined by the equations,

$$\frac{da_1}{dt} = -2a_1 b_1, \quad \frac{db_1}{dt} = a_1 - \frac{1}{a_1}.$$

whose solution can be obtained by eliminating a_1 from (2.5),

$$\frac{db_1}{dt} = \pm \sqrt{(b_1^2 + I_2)^2 - 4}.$$

This can be also written in the form,

$$\frac{d\lambda}{y(\lambda)} \Big|_{\lambda=b_1} = \pm dt,$$

Here $y(\lambda) = \sqrt{(\lambda^2 + I_2)^2 - 4}$, and the differential $\omega_1 = \frac{d\lambda}{y}$ is the holomorphic differential on \mathcal{R} . Then the solution can be written in the *elliptic* integral,

$$(2.7) \quad \int_{b_0}^{b_1} \omega_1 = \pm t + \delta_1$$

where b_0 is a fixed point, and δ_1 is an arbitrary constant. Since the integral (2.7) depends on the path from b_0 to b_1 , it should be defined on the Jacobian $\text{Jac}(\mathcal{R})$, a complex torus \mathbb{C}/Γ with the lattice Γ generated by the period matrix $\Omega = (1 \ B_{11})$ with $B_{11} = \oint_{\beta_1} \omega_1$.

Furthermore the elliptic integral (2.7) should be on the real part of the Jacobian. Namely we have the following maps with an Abel-Jacobi map v to describe the Toda flow;

$$\begin{array}{ccccc} Z_{\mathbb{R}}^P(\gamma) & \longrightarrow & \mathcal{R}_{\mathbb{R}} & \xrightarrow{v} & \text{Jac}(\mathcal{R})_{\mathbb{R}} \\ b_1 & \longmapsto & \mu_1 & \longmapsto & \left[\int_{\mu_*}^{\mu_1} \omega_1 \right] \end{array}$$

where μ_* is a fixed point on $\mathcal{R}_{\mathbb{R}}$, the real part of the Riemann surface. This implies that the Toda flow is *linear* on the Jacobian, and the solution $b_1(t)$ is given by solving the Jacobi inversion problem (just the Jacobi elliptic function in this case), which is defined as the inverse map of v ;

$$\psi : \text{Jac}(\mathcal{R}) \longrightarrow \mathcal{R}.$$

Then the solution of the Jacobi inversion problem is obtained by the zeros of Riemann theta function (see the next section for an explicit form of the solution of Toda flow for genus $g > 1$). Now the question is to determine the structure of the real part of $\text{Jac}(\mathcal{R}) \setminus \Theta$ which is isomorphic to $Z_{\mathbb{R}}^P(\gamma)$. In this example, the Θ is given by

$$\Theta = \{v(p_{\infty}^+), v(p_{\infty}^-)\},$$

and the compactified manifold is given by $\hat{Z}_{\mathbb{R}}^P(\gamma) \cong \mathcal{R}_{\mathbb{R}} \cong \text{Jac}(\mathcal{R})_{\mathbb{R}}$, i.e.

$$(2.8) \quad \text{Jac}(\mathcal{R})_{\mathbb{R}} \cong \begin{cases} S^1 \times \mathbb{Z}/2\mathbb{Z}, & \text{for } \gamma_2 > 2, \\ S^1, & \text{for } \gamma_2 < 2. \end{cases}$$

where each S^1 is labeled by the signs (ϵ_1, ϵ_2) , i.e. $(++)$ or $(--)$ for $\gamma_2 > 2$, and $(--)$ for $\gamma_2 < 2$. Thus one of S^1 for $\gamma_2 > 2$ is obtained by gluing two pieces of \mathbb{R} on the Θ . Finding the gluing structure of the affine pieces of the real part of the Jacobian, $\text{Jac}(\mathcal{R})_{\mathbb{R}} \setminus \Theta \cong Z_{\mathbb{R}}^P(\gamma)$ is one of the main purpose of the present study.

For the case of arbitrary genus, it has been known for quite a while that the number of connected components in the real part of the Jacobian, denoted as $n(\text{Jac}_{\mathbb{R}}) := \text{rank}H_0(\text{Jac}(\mathcal{R})_{\mathbb{R}}, \mathbb{Z})$, is completely determined by the number of connected ones in the real part of Riemann surface, $N_R = \text{rank}H_0(\mathcal{R}_{\mathbb{R}}, \mathbb{Z})$, that is, we have:

Proposition 2.2. ([14], see also [5, 9]) *Let \mathcal{R} be a genus g compact Riemann surface with non-empty real component, $N_R \neq 0$. Then each connected component of $\text{Jac}(\mathcal{R})_{\mathbb{R}}$ is isomorphic to g -dimensional real torus, $\mathbb{T}^g = \mathbb{R}^g/\mathbb{Z}^g$, and*

$$\text{Jac}(\mathcal{R})_{\mathbb{R}} \cong \mathbb{T}^g \times (\mathbb{Z}/2\mathbb{Z})^m,$$

where $m = N_R - 1 \leq g$, and thus $n(\text{Jac}_{\mathbb{R}}) = 2^m$.

We will give the proof based on the solution structure of the periodic Toda lattice in the next section. Then we will study the topological structure of the affine parts of $\text{Jac}(\mathcal{R})_{\mathbb{R}}$ and the divisor Θ , and find the gluing rule among the affine parts labeled by the signs $(\epsilon_1, \dots, \epsilon_N)$ for the compactification. It was shown in [2] that the divisor Θ consists of $g+1$ irreducible components Θ_k which corresponds to the blow-up of a_k , (see (1.12)). We then expect that each torus \mathbb{T}^g in Proposition 2.2 is divided into $g+1$ affine varieties through those divisors Θ_k . In order to study the case with arbitrary genus, we will give a brief review of the solutions of periodic Toda lattice in the next section.

3. SOLUTION OF THE PERIODIC TODA LATTICE

Here we give a brief summary of the theory of periodic Toda lattice and the solution formula in terms of the Riemann theta function $\vartheta(x|B)$. This will be used in the next

section to study the geometry of the divisors which are given by the zeros of $\vartheta(x|B)$ with some translates in x .

3.1. Isospectral manifold and symmetric products of Riemann surfaces. Let a pair of ∞ -vectors $\{\phi^{(1)}, \phi^{(2)}\}$ be a fundamental set of solutions for the difference equation $L\phi = \lambda\phi$ with the periodic condition (1.5). The $\phi^{(j)} = (\dots, \phi_{-1}^{(j)}, \phi_0^{(j)}, \phi_1^{(j)}, \dots)$ are assumed to satisfy

$$(3.1) \quad \begin{cases} \phi_0^{(1)} = 1, \\ \phi_1^{(1)} = 0, \end{cases} \quad \text{and} \quad \begin{cases} \phi_0^{(2)} = 0, \\ \phi_1^{(2)} = 1. \end{cases}$$

(Note here that the choice of the index “0” can be shifted by an arbitrary number $k \pmod{N+1}$.) Then one can define a monodromy matrix M by setting

$$(3.2) \quad \begin{pmatrix} \phi_{k+N}^{(1)} \\ \phi_{k+N}^{(2)} \end{pmatrix} = M \begin{pmatrix} \phi_k^{(1)} \\ \phi_k^{(2)} \end{pmatrix}, \quad \text{with} \quad M = \begin{pmatrix} \phi_N^{(1)} & \phi_{N+1}^{(1)} \\ \phi_N^{(2)} & \phi_{N+1}^{(2)} \end{pmatrix}.$$

First we have:

Lemma 3.1. *With the condition $\prod_{k=1}^N a_k = 1$, we have $\det M = 1$.*

Proof. One can easily show that the Wronskian $W(n) := \phi_n^{(1)}\phi_{n+1}^{(2)} - \phi_n^{(2)}\phi_{n+1}^{(1)}$ satisfies

$$W(n) = a_{n-1}W(n-1) = \left(\prod_{k=0}^{n-1} a_k \right) W(0).$$

Then put $n = N$, i.e. $W(N) = \det M$, and use $W(0) = 1$ to complete the proof. \square

We can also see from the periodic condition $\phi_{k+N} = z\phi_k$ in (1.6) that the spectral parameter z is the eigenvalue of M , and we have,

$$z + \frac{1}{z} - \text{tr}M = 0.$$

Then we have

$$(3.3) \quad \text{tr}M = \phi_N^{(1)} + \phi_{N+1}^{(2)} = P(\lambda),$$

where the N -th polynomial $P(\lambda)$ is given by (1.11).

Let us now find the explicit form of the fundamental solutions $\phi^{(j)}$. By a direct computation, one can easily find:

Lemma 3.2. *The $\phi_{n+1}^{(j)}$ for $n \geq 1$ are given by*

$$(3.4) \quad \phi_{n+1}^{(1)} = -a_N(-1)^{n-1}\Delta_{2,n}, \quad \phi_{n+1}^{(2)} = (-1)^n\Delta_{1,n},$$

where $\Delta_{m,n}$ are defined by (1.9) and $\Delta_{2,1} := 1$.

Then we consider the so-called auxiliary spectrum μ_n 's which are defined by the roots of the equation $\phi_{N+1}^{(1)}(\lambda) = 0$, and we denote

$$(3.5) \quad u_g(\lambda) := (-1)^{N-1}\Delta_{2,N}(\lambda) = \prod_{k=1}^{N-1} (\lambda - \mu_k) = 0, \quad \text{with } g = N - 1.$$

Note that some of the roots are complex, and they form a pair of complex conjugate roots $(\mu_k, \bar{\mu}_k)$. With (3.5), the solution of the Toda lattice a_k 's and b_k 's are expressed

by the symmetric functions of the roots μ_j 's under the isospectral conditions, $I_k = \gamma_k$ for $k = 1, \dots, N+1$. For example, we have from (3.5) and $I_1(L_P) = 0$

$$(3.6) \quad b_1 = -\sum_{j=2}^N b_j = -\sum_{n=1}^{N-1} \mu_n.$$

This implies:

Proposition 3.1. *The map φ from $Z_{\mathbb{R}}^P(\gamma)$ to the real part of the g -fold symmetric product of the affine part \mathcal{R}_0 , denoted as $\mathcal{R}_0^{(g)} := \mathcal{R}_0^g/S_g$,*

$$\varphi : Z_{\mathbb{R}}^P(\gamma) \longrightarrow \left(\mathcal{R}_0^{(g)} \right)_{\mathbb{R}}$$

is surjective. Here S_g is the symmetry group of order $g = N - 1$.

The formulation of the Toda lattice with the variable μ_k 's is called the Mumford system [11] which parametrizes the moduli space of the hyperelliptic Riemann surface in terms of the triples of polynomials, $(u_g(\lambda), v_g(\lambda), w_g(\lambda))$, defined by the factorization of the curve (2.1),

$$(3.7) \quad y^2 = u_g(\lambda)w_g(\lambda) + v_g^2(\lambda).$$

Here $u_g(\lambda)$ is given by (3.5), $v_g(\lambda)$ is a polynomial of degree $g - 1 = N - 2$, and $w_g(\lambda)$ is a monic polynomial of degree $g + 2$. Note here that if all the roots μ_k 's are distinct, the $v_g(\lambda)$ can be expressed in terms of μ_k 's with the condition $v_g(\mu_k) = y(\mu_k)$, i.e.

$$(3.8) \quad v_g(\lambda) = \sum_{k=1}^g y(\mu_k) \prod_{j \neq k} \frac{\lambda - \mu_j}{\mu_k - \mu_j}.$$

The Mumford systems of the triples $(u_d(\lambda), v_d(\lambda), w_d(\lambda))$ with $0 < d < g$ can be also used to parametrize the divisors (see the next section).

Now we discuss the topology of the set of the roots μ_k 's: We first note that $\pi(\mathcal{R}_{\mathbb{R}})$ with $\pi : \mathcal{R} \rightarrow \pi(\mathcal{R})$, $P = (\lambda, y) \mapsto \pi(P) = \lambda$, consists of some intervals,

$$\pi(\mathcal{R}_{\mathbb{R}}) = \bigcup_{m=0}^{N_R} \Lambda_m, \quad 1 \leq N_R \leq N,$$

where $\Lambda_m = [\lambda_{2m}, \lambda_{2m+1}]$ with $\lambda_0 = \lambda_{2N_R+1} = \infty$, (i.e. $[\lambda_0, \lambda_1]$ and $[\lambda_{2N_R}, \lambda_{2N_R+1}]$ are connected through $\pi(p_{\infty}^{\pm})$). Then we first note:

Lemma 3.3. *If the root μ_k is real, then μ_k belongs to one of the intervals Λ_m .*

Proof. Since $\phi_{N+1}^{(1)}(\mu_k) = 0$, we have

$$\text{tr}M = \phi_N^{(1)} + \frac{1}{\phi_N^{(1)}}.$$

This implies $|\text{tr}M| \geq 2$, and therefore $\mu_k \in \pi(\mathcal{R}_{\mathbb{R}})$. \square

Since the set of the points $P_k \in \mathcal{R}$ consists of real ones and/or pairs of complex conjugates, we have the following Lemma on the topology of those sets:

Lemma 3.4. [11] *The set of points $P_k \in \mathcal{R}$ with $\pi(P_k) = \mu_k$, $k = 1, \dots, g$ can be classified as*

- a) $\mu_k \in \Lambda_{m_k}$ for some $m_k \in \{0, 1, \dots, N_R\}$, and the set of those points for each k is isomorphic to S^1 , or
- b) $\overline{P}_{s+1} = P_{s+2}$, or both μ_{s+1}, μ_{s+2} in the same Λ_{m_s} for some m_s , and the set of those points is determined by $V = \{(P, \overline{P}) \in \mathcal{R}^{(2)} = \mathcal{R}^2/S_2\}$.

The later case b) is obtained by considering two points $P, \bar{P} \in \mathcal{R}$, which we can move continuously on \mathcal{R} until $P = \bar{P}$ with $\pi(P) \in \Lambda_{m_s}$, and then move the two points independently on the real loop over Λ_{m_s} . The set of points in this case includes p_∞^\pm . From this Lemma, we can see that the solution corresponding to the case a) except the case $\mu_k \in \Lambda_0 \cup \Lambda_{N_R}$ is just a periodic cycle, and the solutions to all other cases blow up.

Now we can give a proof of Propositions 2.2 (see also [5, 9]):

Proof. Let us first denote $\mathcal{R}_\mathbb{R} = C_0 \sqcup \cdots \sqcup C_m$ with $m = N_R - 1$, where each C_i gives a set corresponding to the case a) in Lemma 3.4, that is, $P \in C_i$ implies $\pi(P) \in \Lambda_i$. Then Lemma 3.4 implies that the symmetric product $\mathcal{R}_\mathbb{R}^{(g)} = (\mathcal{R}^g/S_g)_\mathbb{R}$ is covered by the sets,

$$U(i_1, \dots, i_j) := C_{i_1} \times \cdots \times C_{i_j} \times V^k, \quad \text{with } j + 2k = g.$$

Then the number of disconnected sets can be computed as follows: Let (n_1, \dots, n_g) be an increasing sequence of numbers with $n_k \in \{0, 1, \dots, m\}$ and $m = N_R - 1$, such that if $s \in \{1, \dots, g\}$ is the minimum number for the first nonzero n_k , then n_k 's in the sequence assume,

$$0 = n_1 = \cdots = n_{s-1} < n_s < n_{s+1} < \cdots < n_g \leq m.$$

Then we associate the number n_k in the sequence with the cases in Lemma 3.4 as follows:

- i) $n_k = 0$ implies that either μ_k belongs to $\Lambda_0 \cup \Lambda_M$, or it is in the case b), i.e. $P_k \in C_0$ or $(P_k, P_l) \in V$ for some $P_l = \bar{P}_k$,
- ii) $n_k \neq 0$ implies that μ_k belongs to a bounded interval $\Lambda_{n_k} = [\lambda_{2n_k}, \lambda_{2n_k+1}]$.

Then counting the number of sequences (n_1, \dots, n_g) with $n_k \in \{0, 1, \dots, m\}$ for each case of $\mathcal{R}_\mathbb{R}$ having $N_R = m + 1$ connected components, we obtain the number of connected components of $\mathcal{R}_\mathbb{R}^{(g)}$ as

$$n(\mathcal{R}_\mathbb{R}^{(g)}) = \sum_{j=0}^m \binom{m}{j} = 2^m.$$

On the other hand, writing the anti-holomorphic involution in a matrix form on the lattice given by the period matrix $\Omega = (I, B)$ in (2.4), one can show that each connected component of $(\mathbb{C}^g/\Gamma)_\mathbb{R} \cong \text{Jac}(\mathcal{R})_\mathbb{R}$ is diffeomorphic to the real torus $\mathbb{R}^g/\mathbb{Z}^g = \mathbb{T}^g$.

This then completes the proof by noticing $n(\text{Jac}_\mathbb{R}) = n(\mathcal{R}_\mathbb{R}^{(g)})$ (see also below). \square

Note here that the number $\binom{m}{j}$ represents the total number of the sequences having the form which is identified as the set $U(i_1, \dots, i_j)$ as

$$(3.9) \quad \underbrace{(0, \dots, 0)}_{g-j=s-1}, \underbrace{(n_s, \dots, n_g)}_j \equiv \begin{cases} V^k \times C_{n_s} \times \cdots \times C_{n_g} & \text{if } s = 2k + 1 \\ V^k \times C_0 \times C_{n_s} \times \cdots \times C_{n_g} & \text{if } s = 2k + 2, \end{cases}$$

with $n_s \neq 0$. Thus each connected component can be characterized by the number of components C_{n_i} with nonzero n_i .

3.2. Toda flow on the Jacobian. Let us now find the Toda flow on the Riemann surface which is determined by the image of the dynamics of μ_n 's under the g -hold Abel-Jacobi map $v_g : \mathcal{R}^{(g)} \rightarrow \text{Jac}(\mathcal{R})_\mathbb{R}$. First we note that the time evolution of $\phi_n^{(1)}$ satisfies

$$\frac{d\phi_n^{(1)}}{dt} + a_{n-1}\phi_{n-1}^{(1)} = a_{N-1}\phi_{-1}^{(1)}\phi_n^{(1)} + a_N\phi_n^{(2)}.$$

which can be shown by computing the n -th element of the equation, $\frac{d\phi^{(1)}}{dt} - A\phi^{(1)} = c_1\phi^{(1)} + c_2\phi^{(2)}$. Then the dynamics of μ_n can be obtained by evaluating the equation with (3.4) at

$\lambda = \mu_j$ and $n = N + 1$, i.e.

$$\frac{d}{dt} \left(-a_N \prod_{k=1}^{N-1} (\lambda - \mu_k) \right) \Big|_{\lambda=\mu_j} + a_N \phi_N^{(1)}(\mu_j) = a_N \phi_{N+1}^{(2)}(\mu_j).$$

Noting $\phi_{N+1}^{(2)} = 1/\phi_N^{(1)}$ and using (3.3), we obtain

$$(3.10) \quad \frac{1}{y(\mu_j)} \frac{d\mu_j}{dt} = \pm \frac{1}{\prod_{k \neq j} (\mu_j - \mu_k)},$$

where $y(\lambda) = \sqrt{P(\lambda)^2 - 4}$. Using the Lagrange interpolation formula,

$$\sum_{j=1}^{N-1} \frac{\mu_j^n}{\prod_{k \neq j} (\mu_j - \mu_k)} = \begin{cases} 0 & \text{if } n < N - 2, \\ 1 & \text{if } n = N - 2. \end{cases}$$

we obtain, after the integration,

$$(3.11) \quad \sum_{j=1}^{N-1} \int_{\mu_*}^{\mu_j} \frac{\lambda^n d\lambda}{y(\lambda)} = \begin{cases} \delta_n & n < N - 2, \\ \pm t + \delta_{g-1} & n = N - 2. \end{cases}$$

These equations can be also written in terms of the holomorphic differentials ω_k normalized as (2.3), and we now have the well-known result (see for example [12]):

Proposition 3.2. *The periodic Toda flow can be expressed as a linear flow on the Jacobian $\text{Jac}(\mathcal{R}) \cong \mathbb{C}^g/\Gamma$ associated with the Riemann surface \mathcal{R} of genus $g = N - 1$, i.e.*

$$\sum_{j=1}^g \int_{\mu_*}^{\mu_j} \omega_k = \pm C_{k,g-1} t + \delta'_k, \quad (\text{mod } \Gamma) \quad k = 1, \dots, g,$$

where δ' is a constant, and the $g \times g$ matrix $(C_{k,n})$ is determined by (2.3) with

$$(3.12) \quad \omega_k = \sum_{n=0}^{g-1} C_{k,n} \frac{\lambda^n d\lambda}{y}.$$

With Propositions 3.1 and 3.2, we have constructed the maps,

$$(3.13) \quad \begin{array}{ccccc} \mathbb{Z}_{\mathbb{R}}^P(\gamma) & \xrightarrow{\varphi} & \mathcal{R}_{\mathbb{R}}^{(g)} & \xrightarrow{v_g} & \text{Jac}(\mathcal{R})_{\mathbb{R}} \\ (b_1, \dots, b_N) & \mapsto & [\mu_1, \dots, \mu_g] & \mapsto & \left[\sum_{j=1}^g \int_{\mu_*}^{\mu_j} \omega \right] \end{array}$$

where $\omega = (\omega_1, \dots, \omega_g)$, and the g -fold Abel-Jacobi map v_g is surjective [11].

Since the preimage of each connected component in $\text{Jac}(\mathcal{R})_{\mathbb{R}}$ is given by (3.9) in $\mathcal{R}_{\mathbb{R}}^{(g)}$, we denote

$$(3.14) \quad (\mathbb{T}^g)_j \cong (0, \dots, 0, n_{g-j+1}, \dots, n_g), \quad \text{for } 0 \leq j \leq m \leq g,$$

so that we have

$$(3.15) \quad \text{Jac}(\mathcal{R})_{\mathbb{R}} \cong \bigsqcup_{j=0}^m \mathcal{M}_j, \quad \mathcal{M}_j = \overbrace{(\mathbb{T}^g)_j \sqcup \dots \sqcup (\mathbb{T}^g)_j}^{\binom{m}{j}}.$$

Remark 3.1. *Since $(\mathbb{T}^g)_j$ is a real part of a complex abelian group $\text{Jac}(\mathcal{R}) \cong \mathbb{C}^g/\Gamma$, $(\mathbb{T}^g)_j$ is a real abelian group diffeomorphic to a g -dimensional real torus $\mathbb{R}^g/\mathbb{Z}^g$. We will later introduce $(\mathbb{T}^k)_j$ with*

$$(\mathbb{T}^k)_j \cong \overbrace{(0, \dots, 0, n_{g-j+1}, \dots, n_g)}^k, \quad \text{for } j \leq k < g.$$

However this may not be a k -dimensional real torus except the case with $k = j$. In this paper, we will not discuss the details of $(\mathbb{T}^k)_j$, but its topology can be determined by the set $V = \{(P, \bar{P}) \in \mathcal{R}^{(2)}\}$ (see [13]). The details will be given elsewhere.

Remark 3.2. Equation (3.10) can be expressed as an integrable system of the Mumford system (3.7) [13]:

$$(3.16) \quad \frac{dM_g(\lambda)}{dt} = [M_g(\lambda), B_g(\lambda)],$$

where the matrices $M_g(\lambda)$ and $B_g(\lambda)$ are given by the triples (u_g, v_g, w_g) of the Mumford system,

$$M_g(\lambda) = \begin{pmatrix} v_g(\lambda) & u_g(\lambda) \\ w_g(\lambda) & -v_g(\lambda) \end{pmatrix}, \quad B_g(\lambda) = \frac{1}{2} \begin{pmatrix} 0 & 1 \\ b_g(\lambda) & 0 \end{pmatrix}.$$

Here $b_g(\lambda)$ is a monic polynomial of degree $g + 2$ which is uniquely determined by the consistency of the system (3.16). The equation can be generalized to describe the dynamics on the divisor (see below).

3.3. Riemann theta function and Θ divisor. Here we give the solution of the periodic Toda lattice in terms of the Riemann theta function. We also discuss the geometry of the set of blow-up points determined by the Θ divisor, the zeros of the Riemann theta function.

Let us first define the Siegel upper half plane \mathfrak{S}_g as the set of symmetric $g \times g$ complex matrices with positive definite imaginary part, i.e.

$$\mathfrak{S}_g := \{B \in \mathfrak{M}(g, \mathbb{C}) : B^t = B, \text{Im } B > 0\}.$$

Then on the product $\mathbb{C}^g \times \mathfrak{S}_g$ we can define the Riemann theta function

$$\vartheta(z, B) = \sum_{m \in \mathbb{Z}^g} \mathbf{e} \left(m^t z + \frac{1}{2} m^t B m \right).$$

where $\mathbf{e}(x) = e^{2\pi\sqrt{-1}x}$. Since $\text{Im } B > 0$, the above series converges uniformly on compact sets on $\mathbb{C}^g \times \mathfrak{S}_g$ to a holomorphic function. It is easy to see that the theta function satisfies the quasi-periodic relations,

$$\begin{cases} \vartheta(z + E_k, B) &= \vartheta(z, B) \\ \vartheta(z + B_k, B) &= \mathbf{e}(-z_k - B_{k,k})\vartheta(z, B) \end{cases}$$

where $E_1, \dots, E_g, B_1, \dots, B_g$ are the $2g$ column vectors in the period matrix $\Omega = (I \ B)$. It follows from those equations that for each fixed $B \in \mathfrak{S}_g$ the divisor of the Riemann theta function is invariant under translations by elements of the lattice Γ generated by E_1, \dots, B_g and it induces a divisor Θ_0 on $\text{Jac}(\mathcal{R})$, which is called the theta divisor,

$$\Theta_0 := \{[z] \in \text{Jac}(\mathcal{R}) : \vartheta(z, B) = 0\}.$$

Let us consider a function $f : \mathcal{R} \rightarrow \mathbb{C}$ defined by

$$f(P) := \vartheta(\tilde{v}(P) - c, B)$$

where $c \in \mathbb{C}^g$ is an arbitrary constant vector, and \tilde{v} is a lift of the Abel-Jacobi map v , i.e.

$$\tilde{v} : \mathcal{R}' \longrightarrow \mathbb{C}^g; \quad P \longmapsto \left(\int_{P_0}^P \omega \right)$$

with \mathcal{R}' the $4g$ -sided polygon obtained by cutting \mathcal{R} along the symplectic basis, $\{(\alpha_k, \beta_k) : k = 1, \dots, g\}$, and $\omega = (\omega_1, \dots, \omega_g)$ the holomorphic differential. Then the following Lemma is standard (e.g. see [7] for the proof):

Lemma 3.5. *The followings are satisfied for the zeros of the function $f(P)$:*

- a) $f(P)$ takes exactly g zeros on \mathcal{R} including their multiplicities.
b) If P_1, \dots, P_g are the zeros of $f(P)$, then

$$\sum_{j=1}^g u(P_j) + [\kappa] = [c].$$

where κ is the Riemann constant.

This implies

Proposition 3.3. *Let Q_1, \dots, Q_g be a set of arbitrary points on \mathcal{R} . Then the function $\vartheta\left(\tilde{v}(P) - \sum_{j=1}^g \tilde{v}(Q_j) - \kappa\right)$ has g zeros precisely at $P = Q_1, \dots, Q_g$.*

Geometrically, Proposition 3.3 provides the well-known fact on the theta divisor Θ_0 ,

Corollary 3.1. *The theta divisor Θ_0 of $\text{Jac}(\mathcal{R})$ satisfies*

$$(3.17) \quad \Theta_0 = W^{g-1} + [\kappa]$$

where W^{g-1} is the image of the r -fold Abel-Jacobi map v_r with $r = g - 1$, i.e.

$$(3.18) \quad \begin{array}{ccc} v_r : & \mathcal{R}^{(r)} & \longrightarrow \text{Jac}(\mathcal{R}) \\ & [Q_1, \dots, Q_r] & \longmapsto \left[\sum_{j=1}^r \int_{Q_0}^{Q_j} \omega \right] \end{array}$$

Example 3.1. $g = 2$ Riemann surface and the divisor Θ_0 : In the case of $g = 2$ compact Riemann surface \mathcal{R} , the image W^1 is isomorphic to the Riemann surface, and thus the Jacobian $\text{Jac}(\mathcal{R})$ contains a $g = 2$ Riemann surface as a closed submanifold, $\Theta_0 \cong \mathcal{R}$. Thus we have $\text{Jac}(\mathcal{R})_{\mathbb{R}} \cap \Theta_0 \cong \mathcal{R}_{\mathbb{R}}$.

Now we can find the solution of the periodic Toda lattice by solving the Jacobi inversion problem for μ_j 's in (3.2). Then the solution $\mu_j(t) = \pi(P_j)$ is given by the zero of the theta function,

$$f(P) = \vartheta(\tilde{v}(P) - c, B) = 0, \quad \text{with } c = \pm C_{g-1}t + \kappa',$$

where $\kappa' = \kappa + \delta'$, and $C_{g-1} = (C_{1,g-1}, \dots, C_{g,g-1})$ is the column vector of the matrix $(C_{k,n})$ in (3.12). It follows from (3.6) that we need just the sum of μ_n 's for the solution $b_k(t)$. The sum $\sum_{j=1}^g \mu_j$ can be obtained by the integral, that is,

$$\frac{1}{2\pi\sqrt{-1}} \int_{\partial\mathcal{R}'} \pi(P) d \log f(P) = \sum_{j=1}^g \mu_j + \pi(p_{\infty}^+) + \pi(p_{\infty}^-),$$

where $\pi(p_{\infty}^{\pm})$ are the contributions from the infinity. Finally we obtain (see [12] for the details of the proof):

Proposition 3.4. *The solution $(a_1, \dots, a_N, b_1, \dots, b_N)$ of the Toda lattice is given by*

$$(3.19) \quad \begin{cases} a_k(t) &= \frac{\tau_{k+1}(t)\tau_{k-1}(t)}{(\tau_k(t))^2} \\ b_k(t) &= \frac{d}{dt} \log \frac{\tau_k(t)}{\tau_{k-1}(t)} \end{cases}$$

where $\tau_k(t) = \vartheta(c't + ck + \delta_0, B)$ with $c' = \pm C_{g-1}$, $c = \tilde{v}(p_{\infty}^-) - \tilde{v}(p_{\infty}^+)$ and $\delta_0 = \tilde{v}(p_{\infty}^+) - \kappa - \delta'$.

Now it is clear that if a $\tau_k(t)$ vanishes for some k and t , the functions $a_k(t)$ and $b_{k\pm 1}(t)$ blow up. Thus the divisor Θ is given by the sum of the zero sets of τ -functions, which are the theta divisor Θ_0 and its translates Θ_k , i.e.

$$(3.20) \quad \Theta = \bigcup_{k=0}^g \Theta_k, \quad \text{with} \quad \Theta_k = \Theta_0 + k(\tilde{v}(p_\infty^-) - \tilde{v}(p_\infty^+)).$$

Thus the isospectral set $Z_{\mathbb{R}}^P(\gamma)$ can be compactified by gluing on the Θ divisors. One can also observe that the multiplicity of the zero of $\tau_k(t)$ gives the intersection multiplicity of the divisors. The complex version of the geometry of the divisors has been worked out in [2]. In this paper, we are concerned with the real version of this study, and our goal is to give a Lie theoretic study on the gluing structure in terms of a representation of the affine Weyl group W .

Remark 3.3. *With the solution form (3.19), one can easily show that the tau-functions $\tau_k(t)$ satisfy the Hirota bilinear form,*

$$(3.21) \quad \tau_k \tau_k'' - (\tau_k')^2 = \tau_{k+1} \tau_{k-1},$$

where $\tau_k' = d\tau_k/dt$ and $\tau_k'' = d^2\tau_k/dt^2$. This formula will be useful for the study of degeneracy of the zeros, which characterizes the intersection of the divisors.

4. CONJECTURES ON THE TOPOLOGY OF $\text{Jac}(\mathcal{R})_{\mathbb{R}}$

Here we provide several conjectures on the geometry of the divisor on $\text{Jac}(\mathcal{R})_{\mathbb{R}}$ and the topology of the affine part of $\text{Jac}(\mathcal{R})_{\mathbb{R}}$.

4.1. Geometry of the divisor Θ . Recall that Θ_k is the translate of the theta divisor Θ_0 by $k(Q - P)$ with $Q = v(p_\infty^-), P = v(p_\infty^+)$, so that $\Theta_k \cong \Theta_0 \cong W^{g-1}$. Then following [2], we define the sets Θ_J for each subset $J \subsetneq \{0, 1, \dots, g\}$ as the intersections with Θ_k ,

$$(4.1) \quad \Theta_J := \bigcap_{k \in J} \Theta_k, \quad \text{and} \quad \dim_{\mathbb{C}} \Theta_J = g - |J|.$$

Note here that $\Theta_J \supset \Theta_{J'}$ if $J \subset J'$, and $\Theta_J \cong \mathcal{R}$ if $|J| = g - 1$. We then have a stratification of the Jacobian,

$$\text{Jac}(\mathcal{R}) \supset \Theta^{(1)} \supset \Theta^{(2)} \supset \dots \supset \Theta^{(g)}, \quad \text{with} \quad \Theta^{(k)} := \bigcup_{|J|=k} \Theta_J.$$

We also have a cell decomposition of the divisor Θ in terms of those sets,

$$(4.2) \quad \Theta = \bigsqcup_{J \subsetneq \{0, 1, \dots, g\}} D_J, \quad \text{with} \quad D_J := \Theta_J \bigcap_{k \notin J} \Theta_k^c,$$

where Θ_k^c is the complement set of Θ_k in $\text{Jac}(\mathcal{R})$. In order to compute the intersections $\text{Jac}(\mathcal{R})_{\mathbb{R}} \cap \Theta_J$, we first state:

Lemma 4.1. *Let $(\mathbb{T}^g)_j$ be a g -dimensional torus defined by (3.14). Then we have*

$$(\mathbb{T}^g)_j \cap \Theta_J = \begin{cases} (\mathbb{T}^{g-|J|})_j & \text{if } |J| \leq g - j, \\ \emptyset & \text{if } |J| > g - j. \end{cases}$$

Proof. Let us first consider the case of $J = \{k\}$ for $k = 0, 1, \dots, g$. Since Θ_k is isomorphic to the image of the $(g - 1)$ -hold symmetric product of the Riemann surfaces, $\Theta_k \cong W^{g-1}$

(Proposition 3.1), $\Theta_k \cap (\mathbb{T}^g)_j$ consists of the elements denoted by $(\overbrace{0, \dots, 0}^{g-j-1}, n_{g-j+1}, \dots, n_g)$, that is, one of the first $(g - j)$ elements in $(\mathbb{T}^g)_j$ is fixed at a point of infinity. This then

implies the Lemma. In the case of $|J| > 1$, it is not hard to see that a similar argument can lead to the proof. \square

Remark 4.1. *Lemma 4.1 is a consequence of the Poincaré's formula,*

$$\Theta_J \cong |J|! W^{g-|J|} \quad (\text{homologous})$$

and the variety $(\mathbb{T}^{g-|J|})_j$ is singular in general. It seems however that the $(\mathbb{T}^{g-|J|})_j$ is smooth if J consists of a consecutive numbers $J = \{s+1, \dots, s+|J|\}, (\text{mod } N)$ (see Theorem 5. and Corollary 1. in [2]).

Since the divisor Θ is the sum of the zeros of tau-functions, $\tau_k(t) = \theta(k(Q-P) + R(t))$, the cell D_J is also defined by

$$R(t_J) \in D_J \cap \text{Jac}(\mathcal{R})_{\mathbb{R}} \stackrel{\text{def}}{\iff} \tau_k(t_J) = 0, \text{ iff } k \in J.$$

On D_J , one can determine the multiplicity m_k of the zero of $\tau_k(t)$ with $k \in J$, which describes the intersection geometry of the divisors Θ_k with $k \in J$.

Lemma 4.2. *Let J be a set of consecutive numbers given by $\{i+1, \dots, i+s\} \pmod{N}$. Then $\tau_k(t)$ has the following form near its zero $t = t_J$,*

$$(4.3) \quad \tau_{i+k}(t) \simeq (t - t_J)^{m_k} + \dots, \quad \text{with } m_k = k(s+1-k), \quad 1 \leq k \leq s.$$

Proof. Substituting (4.3) into (3.21), and using $\tau_i(t_J) \neq 0$, we have $m_k = k(m_1 + 1 - k)$. Then $\tau_{i+s+1}(t_J) \neq 0$ implies $m_1 = s$. \square

As was shown in [3], the divisor D_J can be parametrized by the so-called *limit matrix*, say L_J , defined by

$$(4.4) \quad L_J := \lim_{t \rightarrow t_J} n_J^{-1}(t) L(t) n_J(t), \quad \text{for some } n_J(t) \in N_-,$$

where L_J has $2N - |J|$ variables, and N_- is the set of lower triangular matrices with 1's on the diagonals. The matrix n_J is determined uniquely by the structure of the blow-up given in Lemma 4.2. For the set $J = \{i_1, \dots, i_n\}$, the limit matrix L_J can be constructed from a chain of matrices starting from the original matrix L_P ,

$$L_P \rightarrow L_{J_1} \rightarrow L_{J_2} \rightarrow \dots \rightarrow L_J,$$

where $J_k = \{i_1, \dots, i_k\}$ with $k < n = |J|$ and $J_n = J$. The matrices $n_{J_k} \in N_-$ can be found easily in terms of L_{J_k} , and then we have the limit $n_{J_k}^{-1} L_{J_k} n_{J_k} \rightarrow L_{J_{k+1}}$. Then the L_{J_k} parametrizes the D_{J_k} divisor of $\text{codim}_{\mathbb{C}} D_{J_k} = k$ (we will report the details elsewhere).

Now with the limit matrix L_J , one can also define an integrable system on D_J in terms of the Mumford triples $(u_d(\lambda), v_d(\lambda), w_d(\lambda))$ with $d = \dim_{\mathbb{C}} D_J = g - |J|$ [11, 13]. Here the triples satisfy

$$y^2 = P_J(\lambda)^2 - 4 = u_d w_d + v_d^2, \quad \det(L_J - \lambda I) = -(z + z^{-1} - P_J(\lambda)),$$

where $u_d(\lambda)$ is a monic polynomial of degree d , $v_d(\lambda)$ is a polynomial of degree $d-1$ with $v_d(\mu_j) = y(\mu_j)$ whenever $u_d(\mu_j) = 0$, and $w_d(\lambda)$ is a monic polynomial of degree $2g+2-d$. The zeros of $u_d(\lambda)$ parametrize the variety $W^d \subset \text{Jac}(\mathcal{R})$, the image of the symmetric products $\mathcal{R}^{(d)}$ under the k -hold Abel-Jacobi map (see [11] for this statement in terms of the divisor $\text{Div}_0^{+,d}(\mathcal{R})$). In particular, if J is given by a set of $(g-d)$ consecutive numbers, say $J = \{1, \dots, g-d\}$, then $u_d(\lambda)$ can be expressed as

$$\begin{aligned} u_d(\lambda) &= (-1)^d \Delta_{N-d+1, N}(\lambda) = \lambda^d - \left(\sum_{i=N-d+1}^N b_i \right) \lambda^{d-1} + \dots \\ &= \prod_{i=1}^d (\lambda - \mu_i), \end{aligned}$$

where $\delta_{m,n}(\lambda)$ is given by (1.9) (see the Example 4.1 below). The integrable system on D_J is then given by the same form as (3.16) which gives (3.10) with the index j taking

$1 \leq j \leq d = g - |J|$. This system can be used to show the smoothness of Θ_J for this type of J . Note that the Jacobi inversion problem for such system is not complete, and the solution manifold cannot be completed into Abelian variety (in this sense the system is *not* algebraically integrable) [1]. We will study the details of the solutions of those integrable systems elsewhere.

Example 4.1. $D_J \subset \text{Jac}(\mathcal{R})$ of the Riemann surface of $g = 3$:

i) $J = \{1, 2\}$: The limit matrix $L_{\{1,2\}}$ is given by

$$L_{\{1,2\}} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \xi_3 & -\xi_2 & \xi_1 & 1 - \zeta_1 z^{-1} \\ z - \zeta_2 & 0 & 0 & \mu_1 \end{pmatrix},$$

where the 3×3 submatrix at the top left corner is the companion matrix of the corresponding submatrix, i.e. $\xi_1 = b_1 + b_2 + b_3, \xi_2 = b_1 b_2 + b_2 b_3 + b_3 b_1 - a_1 - a_2, \xi_3 = b_1 b_2 b_3 - a_1 b_3 - a_2 b_1$, and $\zeta_1 = -a_1 a_3, \zeta_2 = -a_2 a_4, \mu_1 = b_4$. The spectral curve associated with $L_{\{1,2\}}$ is given by

$$\det(L_{\{1,2\}} - \lambda I) = -(z + I_5 z^{-1} - P_{\{1,2\}}(\lambda)), \quad P_{\{1,2\}}(\lambda) = \lambda^4 + \sum_{k=1}^4 (-1)^k I_k \lambda^{4-k}.$$

Here the invariants I_1, \dots, I_4 and I_5 are given by

$$\begin{cases} I_1 = \xi_1 + \mu_1 = 0, & I_2 = \xi_1 \mu_1 + \xi_2, & I_3 = \xi_2 \mu_1 + \xi_3, \\ I_4 = \xi_3 \mu_1 + \zeta_1 + \zeta_2, & I_5 = \zeta_1 \zeta_2 = 1, \end{cases}$$

The polynomials $u_1(\lambda), v_1(\lambda)$ of the Mumford triples are just $u_1(\lambda) = \lambda - \mu_1, v_1(\lambda) = \pm y(\mu_1)$. Then the isospectral variety in $\text{Jac}(\mathcal{R})_{\mathbb{R}}$ with the constraints $\tau_1 = \tau_2 = 0$ is given by

$$\begin{aligned} D_{\{1,2\}} \cap \text{Jac}(\mathcal{R})_{\mathbb{R}} &= \{(\xi_1, \xi_2, \xi_3, \zeta_1, \zeta_2, \mu_1) \in \mathbb{R}^6 : I_k = \gamma_k \in \mathbb{R}, k = 1, \dots, 5\} \\ &\cong \{(\zeta_1, \mu_1) \in \mathbb{R}^2 : \zeta_1 + \zeta_1^{-1} - P_{\{1,2\}}(\mu_1) = 0\} \\ &\cong \mathcal{R}_{\mathbb{R}} \cap \mathcal{R}_0. \end{aligned}$$

The integrable system on this variety, $du_1(\lambda)/dt = v_1(\lambda)$, is given by

$$\frac{d\mu_1}{dt} = \pm y(\mu_1) = \pm \sqrt{P_{\{1,2\}}(\mu_1)^2 - 4}.$$

which describes the dynamics on the curve $\mathcal{R}_{\mathbb{R}}$.

ii) $J = \{1, 3\}$: The limit matrix in this case is given by

$$L_{\{1,3\}} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -\xi_2 & \xi_1 & 1 - \zeta_1 z^{-1} & 0 \\ 0 & 0 & 0 & 1 \\ z - \zeta_2 & 0 & -\eta_2 & \eta_1 \end{pmatrix},$$

where $\xi_1 = b_1 + b_2, \xi_2 = b_1 b_2 - a_1, \eta_1 = b_3 + b_4, \eta_2 = b_3 b_4 - a_3, \zeta_1 = a_4 b_1 b_2$ and $\zeta_2 = a_2 b_1 b_3$. The invariants are obtained as

$$\begin{cases} I_1 = \xi_1 + \eta_1 = 0, & I_2 = \xi_1 \eta_1 + \xi_2 + \eta_2, & I_3 = \xi_1 \eta_2 + \xi_2 \eta_1, \\ I_4 = \xi_2 \eta_2 + \zeta_1 + \zeta_2, & I_5 = \zeta_1 \zeta_2 = 1, \end{cases}$$

from which we have the isospectral variety with constraints $\tau_1 = \tau_3 = 0$,

$$\begin{aligned} D_{\{1,3\}} \cap \text{Jac}(\mathcal{R})_{\mathbb{R}} &= \{(\xi_1, \xi_2, \eta_1, \eta_2, \zeta_1, \zeta_2) \in \mathbb{R}^6 : I_k = \gamma_k \in \mathbb{R}, k = 1, \dots, 5\} \\ &\cong \{(\zeta_1, \xi_1) \in \mathbb{R}^2 : \zeta_1 + \zeta_1^{-1} + Q(\xi_1) = 0\}, \end{aligned}$$

where $Q(\xi_1)$ is a rational function of ξ_1 given by

$$Q(\xi_1) = \frac{1}{4}(\xi_1^2 + I_2)^2 - \frac{I_3^2}{4\xi_1^2} - I_4.$$

Then the corresponding divisor $\Theta_{\{1,3\}}$ in $\text{Jac}(\mathcal{R})_{\mathbb{R}}$ contains a 1-dimensional disconnected singular variety consisting of two circles transversally intersecting at two distinct points of infinity. Thus the divisors Θ_J have quite different topological structure depending on the set J , and the difference may be characterized by the sub-Dynkin diagram associated with $\tau_k = 0$ for $k \in J$ (see [2], and we will report the details elsewhere).

4.2. Affine part of the Jacobian. Let us first state the following Conjecture on the structure of the affine parts of $\text{Jac}(\mathcal{R})_{\mathbb{R}}$, which is isomorphic to $Z_{\mathbb{R}}^P(\gamma)$ (see (2.6) for the case of $g = 1$);

Conjecture 4.1. *The affine part of $\text{Jac}(\mathcal{R})_{\mathbb{R}}$ has a decomposition,*

$$\text{Jac}(\mathcal{R})_{\mathbb{R}} \setminus \Theta \cong \bigsqcup_{j=0}^m \mathcal{M}_j^{\circ}, \quad m = N_{\mathbb{R}} - 1 \leq g,$$

with \mathcal{M}_j° consisting of $\binom{m}{j}$ disconnected components $(\mathbb{T}^g)_j \setminus \Theta$ in which each torus $(\mathbb{T}^g)_j$ is divided into $g + 1$ cylinders, $\mathbb{R}^{g-j} \times \mathbb{T}^j$, through the divisors Θ_k in (3.20),

$$(\mathbb{T}^g)_j \setminus \Theta = (\mathbb{R}^{g-j} \times \mathbb{T}^j) \sqcup \cdots \sqcup (\mathbb{R}^{g-j} \times \mathbb{T}^j).$$

This is motivated as follows: Recall that $(\mathbb{T}^g)_j$ is labeled by $(\overbrace{0, \dots, 0, n_{g-j+1}, \dots, n_g}^g)$ where each element with the label 0 contains the divisors corresponding to $p_{\infty}^{\pm} \in \mathcal{R}$. Then removing those points through $(g + 1)$ divisor Θ_k , $\Theta = \cup_{k=0}^g \Theta_k$, we expect to have the $(g + 1)$ disconnected cylinders. To prove the above Conjecture, one needs to determine the structure of the intersections of the divisors on $(\mathbb{T}^g)_j$.

We then consider the gluing pattern of those disconnected cylinders in $(\mathbb{T}^g)_j \setminus \Theta$ which are marked by the set of signs $(\epsilon_1, \dots, \epsilon_N)$ with $\epsilon_k = \text{sign}(a_k)$: Let Σ be the set of periodic sequences of signs,

$$\Sigma(N) := \{(\cdots, \sigma_{k-1}, \sigma_k, \sigma_{k+1}, \cdots) : \sigma_k \in \{\pm\}, \sigma_{k+N} = \sigma_k, \forall k \in \mathbb{Z}\}.$$

We also denote by $\sigma^0 \in \Sigma(N)$ the special element consisting of all + signs. We then mark each cylinder $\mathbb{R}^{g-j} \times \mathbb{T}^j$ with a periodic sequence of signs, σ , and we relate the sign $\epsilon_k = \text{sign}(a_k)$ with $\epsilon_k = \sigma_k \sigma_{k+1}$.

We now define a permutation s_k on $\Sigma(N)$,

$$\begin{aligned} s_k : \quad \Sigma(N) &\longrightarrow \Sigma(N) \\ (\cdots, \sigma_k, \sigma_{k+1}, \cdots) &\longmapsto (\cdots, \sigma_{k+1}, \sigma_k, \cdots) \end{aligned}$$

where s_k also acts on the periodic ones σ_{k+mN} and σ_{k+1+mN} for all $m \in \mathbb{Z}$, that is, $k \in \mathbb{Z}/N\mathbb{Z}$. The set of all permutations forms an affine Weyl group W associated with $A_{N-1}^{(1)}$, the affine Kac-Moody algebra of $\mathfrak{sl}(N)$,

$$W := \left\langle s_1, \dots, s_N : \begin{array}{l} s_k^2 = e \text{ (identity)}, (s_k s_{k+1})^3 = e \text{ mod } N \\ (s_k s_j)^2 = e, \text{ if } k \neq j, j \pm 1 \text{ mod } N \end{array} \right\rangle$$

Then one can show that the action of W on signs has the following periodic property:

Lemma 4.3. *For each $\sigma \in \Sigma(N) \setminus \{\sigma^0\}$, there exists a periodic action of length N , $w_N^{(\sigma)} := s_{i_N} \cdots s_{i_1} \in W$ with $i_j \neq i_k$ if $j \neq k$, such that*

- i) $w_N^\sigma \sigma = \sigma$
- ii) $s_{i_j} \cdots s_{i_1} \sigma \neq s_{i_k} \cdots s_{i_1} \sigma$, for $0 \leq j < k < N$.
- iii) N is the minimum number, i.e. if there is $w (\neq e) \in W$ such that $w\sigma = \sigma$, then $l(w) \geq N$,

where $l(w)$ is the length of the element $w \in W$.

Proof. Consider a sequence $(\overbrace{+\cdots+}^{n_1} \overbrace{-\cdots-}^{n_2} \overbrace{+\cdots+}^{n_3} \overbrace{-\cdots-}^{n_4})$ with $\sum_{k=1}^4 n_k = N$, and assume σ be given by its periodic extension of period N . First move the $+$ at the n_1 -th place to the $(n_1 + n_2)$ -th place by applying $w_1 = s_{n_1+n_2-1} \cdots s_{n_1+1} s_{n_1}$ with $l(w_1) = n_2$. Also move the $+$ at the $(n_1 + n_2 + n_3)$ -th place to the N -th place by $w_2 = s_{N-1} \cdots s_{n_1+n_2+n_3}$ with $l(w_2) = n_4$. Then we obtain $(\overbrace{+\cdots+}^{n_1-1} \overbrace{-\cdots-}^{n_2} \overbrace{+\cdots+}^{n_3} \overbrace{-\cdots-}^{n_4} +)$. Now move the $-$ at the n_1 -th place to the 0-th place by $w_3 = s_N s_1 \cdots s_{n_1-2} s_{n_1-1}$ with $l(w_3) = n_1$, and move the $-$ at the $(n_1 + n_2 + n_3 - 1)$ -th place to the $(n_1 + n_2)$ -th place by $w_4 = s_{n_1+n_2+1} \cdots s_{n_1+n_2+n_3}$ with $l(w_4) = n_3$. Thus applying $w = w_4 w_3 w_2 w_1$ with $l(w) = N$, the sequence σ returns to the original one. Notice that the sequences in the middle steps are all different. It is easy to extend the procedure to the general case. \square

Let us consider the sign changes in a_k 's through a blow-up, which is determined by $\tau_k(t_*) = 0$ for some k and $t_* \in \mathbb{R}$. Then from (3.19), if $\tau_k(t) \sim (t - t_*)$ near $t = t_*$, the solutions $a_k(t)$ and $b_k(t)$ near $t = t_*$ has the following form,

$$\begin{aligned} a_k(t) &\sim \frac{1}{(t - t_*)^2}, & a_{k-1}(t) &\sim (t - t_*), & a_{k+1}(t) &\sim (t - t_*), \\ b_k(t) &\sim \frac{1}{t - t_*}, & b_{k+1}(t) &\sim \frac{1}{t - t_*}, \end{aligned}$$

and all the other variables a_j and b_j are regular near $t = t_*$. Observe that the a_k remains the same sign (negative) before and after the blow-up point, and $a_{k\pm 1}$ change their signs. This implies that the sign changes of $a_{k\pm 1}$ through the blow-up of a_k correspond to the permutation s_k . Recall $\epsilon_k = \text{sign}(a_k) = \sigma_k \sigma_{k+1}$. Then Lemma 4.3 suggests that the chain of σ 's appearing in the periodic permutation $w_N^{(\sigma)}$ provides the labels of the cylinders in each torus $(\mathbb{T}^g)_j$, and the gluing rule is given by the sign-representation of the affine Weyl group W . It may be convenient to use the ϵ -sign for the label. Then we have the general formula for the sign change in terms of the extended Cartan matrix $(C_{i,j})_{1 \leq i,j \leq N}$ for $A_{N-1}^{(1)}$,

$$(4.5) \quad s_k : \epsilon_j \longmapsto \epsilon_j \epsilon_k^{-C_{k,j}}$$

which was also obtained in [6] for the case of the nonperiodic Toda lattice. We summarize those as the following Conjecture:

Conjecture 4.2. *Each connected piece (cylinder) in the affine parts of the isospectral level set $\mathbb{Z}_R^P(\gamma)$ can be marked by the signs $(\epsilon_1, \dots, \epsilon_N)$ with $\epsilon_k = \text{sign}(a_k)$, and they are glued together according to the action of a length N -element of the affine Weyl group W on those signs (sign-representation of W).*

Example 4.2. $N = 3$ periodic Toda lattice [4]:

- i) $m = 2$ ($N_R = 3$, three connected components in $\mathcal{R}_{\mathbb{R}}$): The real part of the Jacobian $\text{Jac}(\mathcal{R})_{\mathbb{R}}$ consists of four tori, and the affine part is given by

$$\text{Jac}(\mathcal{R})_{\mathbb{R}} \setminus \Theta \cong \mathcal{M}_0^{\circ} \sqcup \mathcal{M}_1^{\circ} \sqcup \mathcal{M}_2^{\circ},$$

where

$$\begin{cases} \mathcal{M}_2^\circ &= (\mathbb{T}^2)_2 = \mathbb{T}^2, \\ \mathcal{M}_1^\circ &= (\mathbb{T}^2 \setminus \Theta)_1 \sqcup (\mathbb{T}^2 \setminus \Theta)_1, \\ \mathcal{M}_0^\circ &= (\mathbb{T}^2 \setminus \Theta)_0 = \mathbb{R}^2 \sqcup \mathbb{R}^2 \sqcup \mathbb{R}^2, \end{cases}$$

Each $(\mathbb{T}^2 \setminus \Theta)_1$ consists of three cylinders $\mathbb{R} \times S^1$. The four \mathbb{T}^2 's as compactified connected level sets are marked by the signs $(\epsilon_1, \epsilon_2, \epsilon_3)$ with $\prod_{k=1}^3 \epsilon_k = 1$. For examples, \mathbb{T}^2 in the \mathcal{M}_2° is marked by $(+++)$, and each connected component in $(\mathbb{T}^2 \setminus \Theta)_j$ for other \mathcal{M}_j 's is marked by either $(+- -)$ or $(- + -)$ or $(- - +)$. Those components are glued together by the W -action on the signs,

$$(+, -, -) \xrightarrow{s_2} (-, -, +) \xrightarrow{s_1} (-, +, -) \xrightarrow{s_3} (+, -, -).$$

ii) $m = 1$ ($N_R = 2$): The $\text{Jac}(\mathcal{R})_{\mathbb{R}}$ consists of two tori, and the affine part is

$$\text{Jac}(\mathcal{R})_{\mathbb{R}} \setminus \Theta \cong \mathcal{M}_0^\circ \sqcup \mathcal{M}_1^\circ,$$

where

$$\begin{cases} \mathcal{M}_1^\circ &= (\mathbb{T}^2 \setminus \Theta)_1 = (\mathbb{R} \times S^1) \sqcup (\mathbb{R} \times S^1) \sqcup (\mathbb{R} \times S^1), \\ \mathcal{M}_0^\circ &= (\mathbb{T}^2 \setminus \Theta)_0 = \mathbb{R}^2 \sqcup \mathbb{R}^2 \sqcup \mathbb{R}^2, \end{cases}$$

iii) $m = 0$ ($N_R = 1$): The $\text{Jac}(\mathcal{R})_{\mathbb{R}}$ is just one torus, and the affine part is

$$\text{Jac}(\mathcal{R})_{\mathbb{R}} \setminus \Theta \cong \mathcal{M}_0^\circ = (\mathbb{T}^2 \setminus \Theta)_0 = \mathbb{R}^2 \sqcup \mathbb{R}^2 \sqcup \mathbb{R}^2$$

Acknowledgement The author is grateful to N. Ercolani, H. Flaschka and J. Gibbons for valuable discussions and suggestions. He would also like to thank L. Casian who has provided several useful ideas on the beginning stage of this study through their fruitful collaboration. This work is partially supported by the NSF grant DMS0071523 and the Newton Institute Program ‘‘Integrable Systems’’ for the period of July-September, 2001.

REFERENCES

1. S. Abenda and Yu. Fedorov, On the weak Kowalevski-Painlevé Property for hyperelliptically separable systems, *Acta Appl. Math.* **60** (2000) 137-178.
2. M. Adler and P. van Moerbeke, The Toda lattice, Dynkin diagrams, singularities and Abelian varieties, *Invent. math.* **103** (1991) 223-278.
3. M. Adler, L. Haine and P. van Moerbeke, Limit matrices for the Toda flow and periodic flags for loop groups, *Math. Ann.* **296** (1993) 1-33.
4. M. Audin, Vecteurs propres de matrices de Jacobi, *Ann. Inst. Fourier, Grenoble* **44**, 5 (1994) 1505-1517.
5. M. Audin, *Spinning Top*, (Cambridge University Press, 1996).
6. L. Casian and Y. Kodama, Toda lattice and toric varieties for real split semisimple Lie algebras. Preprint OSU-MRI 99-17 (math-sg/9912021), *Pacific J. of Math.* (2001) in press.
7. H. Farkas and I. Kra, *Riemann surfaces*, Graduate Texts in Math. **71** (Springer-Verlag, 1980).
8. H. Flaschka and L. Haine. Varieties de drapeaux et reseaux de Toda. *Mathematische Zeitschrift*, **208** (1991), 545-556.
9. B. H. Gross and J. Harris, Real algebraic curves, *Ann. scient. Ec. Norm. Sup.*, **14** (1981), 157-182.
10. Y. Kodama and J. Ye, Toda Lattices with indefinite metric II: Topology of the iso-spectral manifolds. *Physica D*, **121** (1998), 89-108.
11. D. Mumford, *Tata Lectures on Theta II*, Progress in Mathematics **43** (Birkhäuser, Boston 1984).
12. M. Toda, *Theory of Nonlinear Lattices*, (Springer-Verlag, New York, 1981).
13. P. Vanhaecke, Integrable systems and symmetric products of curves. *Math. Z.*, **227** (1998) 93-127.
14. G. Weichold, Über symmetrische Riemannsche Flächen, *Zeit. J. Math. Phys.* **28** (1883), 321-351.