

Proton asymmetry in non-mesonic weak decay of light hypernuclei

T. Maruta^a, S. Ajimura^b, K. Aoki^c, A. Banu^d, H. Bhang^e, T. Fukuda^f, O. Hashimoto^g, J. I. Hwang^e, S. Kameoka^g, B. H. Kang^e, E. H. Kim^e, J. H. Kim^e, M. J. Kim^e, Y. Miura^g, Y. Miyake^a, T. Nagae^c, M. Nakamura^a, S. N. Nakamura^g, H. Noumi^c, S. Okada^h, Y. Okayasu^g, H. Outaⁱ, H. Park^j, P. K. Saha^f, Y. Sato^c, M. Sekimoto^c, T. Takahashi^g, H. Tamura^g, K. Tanidaⁱ, A. Toyoda^c, K. Tsukada^g, T. Watanabe^g, H. J. Yim^e

^aDepartment of Physics, University of Tokyo, Tokyo 113-0033, Japan

^bDepartment of Physics, Osaka University, Osaka 560-0043, Japan

^cHigh Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan

^eDepartment of Physics, Seoul National University, Seoul 151-742, Korea

^dGSI, Darmstadt D-64291, Germany

^fOsaka Electro-Communication University, Osaka 572-8530, Japan

^gPhysics Department, Tohoku University, Sendai 980-8578, Japan

^hDepartment of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan.

ⁱRIKEN, Wako, Saitama 351-0198, Japan

^jKorea Research Institute of Standards and Science (KRISS), Daejeon, 305-600, Korea

We have obtained the decay asymmetry parameters in non-mesonic weak decay of polarized Λ -hypernuclei by measuring the proton asymmetry. The polarized Λ -hypernuclei, ${}^5_{\Lambda}\text{He}$, ${}^{12}_{\Lambda}\text{C}$, and ${}^{11}_{\Lambda}\text{B}$, were produced in high statistics via the (π^+, K^+) reaction at 1.05 GeV/c in the forward angles. Preliminary analysis shows that the decay asymmetry parameters are very small for these s-shell and p-shell hypernuclei.

1. INTRODUCTION

The non-mesonic weak decay (NMWD : $\Lambda N \rightarrow NN$) of Λ -hypernuclei gives us a very unique opportunity to study baryon-baryon weak interaction. There exist several experimental observables; life times (total decay rates), branching ratios ($\Gamma(\Lambda p \rightarrow np)$, $\Gamma(\Lambda n \rightarrow nn)$), etc. The asymmetry parameter of decay proton from the $\Lambda p \rightarrow np$ process, α_p^{NM} , is another important observable to investigate the reaction mechanism of NMWD, because it comes from the interference between the parity-conserving and parity-violating amplitudes.

The decay angular distribution of protons from the NMWD of polarized hypernuclei, $W(\theta)$, is expressed as

$$W(\theta) = 1 + A \cos \theta = 1 + \alpha P_{\Lambda} \cos \theta \quad (1)$$

, where A is the asymmetry, P_{Λ} denotes the polarization of a Λ -hypernucleus, and θ means the emission angle of protons with respect to the polarization axis.

So far, two experiments reported the asymmetry parameters of the NMWD's of ${}^{12}_{\Lambda}\text{C}$ [1] and ${}^5_{\Lambda}\text{He}$ [2]. In KEK-PS E160, a large negative α_p^{NM} of -1.3 ± 0.4 was observed for p-shell

hypernuclei[1], while the α_p^{NM} of ${}^5_{\Lambda}\text{He}$ was measured to be 0.24 ± 0.22 by KEK-PS E278[2]. There is a large discrepancy between two values, which might suggest the difference of the reaction mechanisms of the NMWD's between s-shell and p-shell hypernuclei. However, it would be too early to draw a conclusion in taking account of the statistical significance of the data and the systematic errors between two different measurements. Thus, a new measurement with improved statistics and better systematic errors had been awaited.

2. EXPERIMENTS: KEK-PS E462 AND E508

From year 2000 to 2002, we performed two series of experiments to measure the NMWD of ${}^5_{\Lambda}\text{He}$ (E462) and ${}^{12}_{\Lambda}\text{C}$ (E508) in the K6 beamline at the KEK 12-GeV PS with the high-resolution and large-acceptance SKS spectrometer. The (π^+, K^+) reactions at 1.05 GeV/c were used to produce highly-polarized Λ hypernuclei with respect to the horizontal reaction plane. Two decay counter systems were installed symmetrically above and below the target to detect decay protons from NMWD. Each system consisted of a set of drift chambers, two sets of timing counters and a neutron counter array with six layers of plastic counters.




Figure 1. Excitation energy spectra of ${}^5_{\Lambda}\text{He}$ (left) and ${}^{12}_{\Lambda}\text{C}$ (right). From top to bottom, inclusive spectra, those with a pion coincidence, and with a proton coincidence, respectively.

Figure 1 shows excitation energy spectra of ${}^5_{\Lambda}\text{He}$ (left) and ${}^{12}_{\Lambda}\text{C}$ (right) in three different conditions: from top to bottom, inclusive spectra, those with a pion coincidence, and with a proton coincidence. We succeeded to detect 52,000 events of the ${}^5_{\Lambda}\text{He}$ ground state and 67,000 events of the ${}^{12}_{\Lambda}\text{C}$ ground state, whose statistics are several times higher than the previous measurements. The substitutional states of ${}^{12}_{\Lambda}\text{C}$ at around 10 MeV excitation

region, whose selection gate is shown in the figure, are open for proton emission, i.e. it goes to ${}^1_1\text{B}$.

3. PRELIMINARY RESULTS

The asymmetry, A , in Eq. (1) is obtained from the ratio, R , of numbers of decay particles emitted to the parallel and anti-parallel directions with respect to that of the hypernuclear polarization. Owing to the large acceptance of SKS spectrometer, we can polarize the hypernucleus upward(downward) by selecting the (π^+, K^+) scattering direction on the left(right) at one setup. Then, the R is expressed as

$$A = \frac{1 - R}{1 + R}, \quad R = \left(\frac{N_{Up}^+ \times N_{Down}^-}{N_{Up}^- \times N_{Down}^+} \right)^{\frac{1}{2}} \quad (2)$$

, where $N_{Up(Down)}^{+(-)}$ denotes the yield of the up (down) counter system at the scattering angle to the left (right). In this expression, systematic errors coming from the differences of detection efficiencies and acceptances between the up and down decay counter systems are canceled out in the first order. In fact, we found the asymmetries of pions and protons from the (π^+, p) reaction, in which we don't expect any asymmetries, were less than 0.3%.

As shown in Eq. (1), we need to know the polarization of a Λ hypernucleus to obtain the asymmetry parameter of proton. In the case of ${}^5_\Lambda\text{He}$, assuming the asymmetry parameter of the mesonic weak decay, α^M , is same as that in free space (-0.642 ± 0.013), we can measure the polarization of the ${}^5_\Lambda\text{He}$ from the pion decay asymmetry. In the case of p-shell hypernuclei, however, it is hard to estimate the polarization from the experimental data, because mesonic decay mode is suppressed due to Pauli blocking, and we don't have a good estimate of α^M . Therefore, we needed to estimate the polarization of p-shell hypernuclei with a help of a theoretical calculation in Ref. [3].

Figure 2 shows a preliminary result of proton asymmetries in the decay of ${}^5_\Lambda\text{He}$ measured in different scattering angles, $6^\circ < |\theta_K| < 15^\circ$, as a function of the polarization of ${}^5_\Lambda\text{He}$ obtained from the mesonic decay. The slope of the linear fit corresponds to the α_p^{NM} , and we found it to be $0.07 \pm 0.08_{-0.00}^{+0.08}$. As a systematic error, we took into account the effect of a small amount of pion contamination into protons, which tends to reduce the proton asymmetry to the negative side. We confirmed the very small positive value of α_p^{NM} reported by Ajimura *et al.*[2] with improved accuracy. In addition, we found α_p^{NM} of p-shell hypernuclei obtained from a weighted average of each hypernucleus to be almost zero ($-0.24 \pm 0.26_{-0.00}^{+0.08}$) as shown in Fig. 2. Note that the present result is twice more accurate than the previous result. The α_p^{NM} (p-shell) of our result is rather consistent with that of ${}^5_\Lambda\text{He}$. It suggests that the decay mechanism of p-shell hypernuclei is similar to that of s-shell hypernuclei.

4. SUMMARY

We have measured the decay asymmetry parameters of proton in the NMWD's of the polarized hypernuclei, ${}^5_\Lambda\text{He}$, ${}^{12}_\Lambda\text{C}$, and ${}^{11}_\Lambda\text{B}$. For the s-shell hypernucleus, ${}^5_\Lambda\text{He}$, it was found to be $0.07 \pm 0.08_{-0.00}^{+0.08}$, which is consistent to the previous value of 0.24 ± 0.22 [2] with smaller errors. For the p-shell hypernuclei, ${}^{12}_\Lambda\text{C}$ and ${}^{11}_\Lambda\text{B}$, we found a similarly small value




Figure 2. (up) Preliminary result of ${}^5_{\Lambda}\text{He}$. The horizontal axis denotes polarization, P_{Λ} , and the vertical one denotes asymmetry, A , for the scattering angles from 6° to 15° . α_p^{NM} is obtained from the slope of a linear fitting. (down) Preliminary result of ${}^{12}_{\Lambda}\text{C}$ and ${}^{11}_{\Lambda}\text{B}$. each axis is the same as that of upper one. Circle points for ${}^{11}_{\Lambda}\text{B}$ and triangle ones for ${}^{12}_{\Lambda}\text{C}$. The final value is derived from a weighted average of two hypernuclei.

of $-0.24 \pm 0.26^{+0.08}_{-0.00}$, which largely deviates from the old value of -1.3 ± 0.4 [1]. The new values are different from those, $-0.6 \sim -0.7$, obtained in the recent theoretical calculations [4, 5]. We might need to reconsider the reaction mechanism of the NMWD.

REFERENCES

1. S.Ajimura *et al.*, Phys. Lett. B282 (1992) 293
2. S.Ajimura *et al.*, Phys. Rev. Lett. 84 (2000) 4052
3. K.Itonaga, T.Motoba, O.Richter and M.Sotona, Phys. Rev. C49 (1994) 1045.
4. K.Sakaki, T.Inoue and M.Oka, Nucl. Phys. A691 (2001) 201c
5. A.Parreño and A.Ramos, Phys. Rev. C65 (2002) 015204