

# Subthreshold $\phi$ meson production in heavy-ion collisions

M Zétényi<sup>†</sup>, H W Barz<sup>‡</sup>, Gy Wolf<sup>†</sup> and B Kämpfer<sup>‡</sup>

<sup>†</sup> KFKI Research Institute for Particle and Nuclear Physics, POB 49, H-1525 Budapest, Hungary

<sup>‡</sup> Forschungszentrum Rossendorf, Pf 510119, 01314 Dresden, Germany


**Abstract.** Within a BUU type transport model we study  $\phi$  meson production in subthreshold Ni+Ni and Ru+Ru reactions. For the first time we included in our model the elementary reaction channels  $\rho + N, \Delta \rightarrow \phi + N, \pi + N(1520) \rightarrow \phi + N$  and  $\pi\rho \rightarrow \phi$ . In spite of a substantial increase of the  $\phi$  multiplicities by these channels our results stay significantly below the preliminary experimental data.

PACS numbers: 25.75.-q, 25.75.Dw

## 1. Introduction

Meson production is a sensitive probe of the reaction dynamics of heavy-ion collisions. This is particularly true for the production of heavy mesons in subthreshold reactions. Open strangeness production in the threshold region has been studied extensively both experimentally and theoretically. On the contrary, only a few measurements have been reported on hidden strangeness (i.e.  $\phi$  meson) production. Recent preliminary results for the reactions Ni + Ni at 1.93 A·GeV and Ru + Ru at 1.69 A·GeV by the FOPI collaboration [1] point to a surprisingly large production cross section. Earlier transport model calculations [2] underestimate these data substantially. In those calculations  $\phi$  mesons are produced in the elementary reactions  $BB \rightarrow \phi BB, \pi B \rightarrow \phi B$  and  $K\bar{K} \rightarrow \phi$ , where B stands for a nucleon or a  $\Delta$  baryon. The discrepancy of the calculations and the data raises the question of whether further elementary channels might essentially contribute to the  $\phi$  meson yield.

The relatively large branching ratio of the  $\phi \rightarrow \pi\rho$  decay channel suggests that processes with a  $\phi\rho\pi$  vertex and hence the  $\rho$  meson can play an important role. Therefore in this contribution we consider  $\phi$  meson production in elementary processes with a  $\rho$  meson in the entrance channel, namely  $\rho B \rightarrow \phi B$  and  $\rho\pi \rightarrow \phi$ . We also test the effect of higher resonances by studying the elementary reaction  $\pi N(1520) \rightarrow N\phi$  via  $\rho$  meson exchange. The reason for choosing the N(1520) resonance is its large partial decay width to  $\rho N$ . Diagrams representing these elementary reaction channels are shown in Fig. 1.



**Figure 1.** Diagrammatic representation of processes contributing to the reactions (a)  $\rho(N, \Delta) \rightarrow \phi N$ , (b)  $\pi N^* \rightarrow \phi N$  and (c)  $\pi \rho \rightarrow \phi$ .

## 2. Cross sections of elementary reactions

Since the cross sections of the elementary reactions are not well known experimentally we employ a one-boson exchange model to estimate them. For the  $BB \rightarrow BB\phi$  and the  $\pi B \rightarrow \phi B$  channels we use the same cross sections as in [2]. The calculations of the  $\rho N \rightarrow \phi N$  and  $\rho \Delta \rightarrow \phi N$  cross sections (Fig. 1a) are based on the effective interaction Lagrangians

$$\mathcal{L}_{\pi NN} = \frac{f_{\pi NN}}{m_\pi} \bar{\psi} \gamma_5 \gamma^\mu \vec{\tau} \psi \cdot \partial_\mu \vec{\pi}, \quad \mathcal{L}_{\pi N\Delta} = -\frac{f_{\pi N\Delta}}{m_\pi} \bar{\psi} \vec{T} \psi^\mu \cdot \partial_\mu \vec{\pi} + \text{h.c.},$$

$$\mathcal{L}_{\pi \rho \phi} = \frac{f_{\pi \rho \phi}}{m_\phi} \epsilon_{\mu\nu\alpha\beta} \partial^\mu \phi^\nu \partial^\alpha \vec{\rho}^\beta \cdot \vec{\pi}.$$


Here  $\psi$  and  $\psi_\mu$  are the nucleon field and the Rarita-Schwinger field for the  $\Delta$  resonance, respectively, and  $\pi$ ,  $\vec{\rho}_\mu$ ,  $\phi_\mu$  stand for the meson fields. We include multipole form factors for off-shell pions and  $\rho$  mesons. Coupling constants and cutoff parameters are taken from [2, 3]. The resulting isospin averaged cross sections are shown in Fig. 2. It should be noted that the cross section of the  $\rho B$  channels also depends strongly on the actual mass of the incoming  $\rho$  meson (and  $\Delta$  particle).

To calculate the cross section of the  $\pi N(1520)$  channel (Fig. 1b) we tested several possibilities for the interaction Lagrangian for the  $\rho NN(1520)$  vertex such as

$$\mathcal{L}_{\rho NN(1520)} = \frac{f_1}{m_\rho} \bar{\psi} \vec{T} \gamma^\nu \psi_{1520}^\mu (\partial_\nu \vec{\rho}_\mu - \partial_\mu \vec{\rho}_\nu) + \text{h.c.}, \quad (1)$$

$$\mathcal{L}_{\rho NN(1520)} = \frac{f_2}{m_\rho^2} \bar{\psi} \vec{T} \sigma^{\alpha\beta} \psi_{1520}^\mu \partial_\alpha \partial_\mu \vec{\rho}_\beta + \text{h.c.} . \quad (2)$$

We fitted the coupling constants to the  $\rho(\text{two-pion})$  decay channel of the  $N(1520)$  and obtained the values  $f_1 = 10.5$  and  $f_2 = 68$ . We observe that Lagrangian (2) gives a much larger cross section for  $\phi$  meson production than Lagrangian (1) (see Fig. 2). However, the value of the coupling  $f_2$  extracted from the  $N(1520)$  decay differs significantly from the value estimated within the quark model. As the relevant energies in the decay and the production process are very different the assumption of a constant coupling might fail. In our BUU calculations reported below we use the cross sections obtained using



**Figure 2.** Production cross sections for  $\phi$  mesons for several binary reactions as a function of the relative kinetic centre-of-mass energy in the entrance channel. The cross sections are calculated for the nominal masses of the resonances. Especially cross sections for  $\rho\Delta$  collisions having different masses can considerably deviate from the curve shown.

Lagrangian (1) as standard. We also carried out some calculations using Lagrangian (2) to test the uncertainties of the  $\phi$  meson yield from the  $\pi N(1520)$  channel.

### 3. BUU calculations of the $\phi$ meson yield

We used the above elementary cross sections to calculate the  $\phi$  meson production within a BUU model. In our model nucleonic and  $\Delta$  resonances are included up to 2.2 GeV. The resonance properties are fitted to one- and two-pion production data in  $\pi N$  scattering while the production cross sections of resonances are derived from meson production data in NN collisions. We employed a soft momentum dependent potential for nucleons and  $\Delta(1232)$ . The in-medium mass of the  $\phi$  meson is treated according to [4]. We included also the absorption of  $\phi$  mesons via the  $\phi N \rightarrow K\Lambda$  [5] and the  $\phi N$  elastic scattering with a cross section of 0.56 mb.

The resulting  $\phi$  meson yields are shown in Table 1 together with the data obtained by extrapolating the experimental yields to the entire phase space. The inclusion of the new  $\rho$  channels results in a substantial increase of the  $\phi$  meson yields by a factor of 2.5. The dominant contribution comes from the  $\rho B$  channel. The  $\pi N(1520)$  channel calculated with the Lagrangian (1) does not contribute essentially to the  $\phi$  meson yield. However, using the Lagrangian (2) the yields of this channel rise by a factor of about 10

yields from	Ni + Ni (1.93 GeV)	Ru + Ru (1.69 GeV)
B + B	$3.5 \cdot 10^{-4}$	$3.1 \cdot 10^{-4}$
$\pi$ + B	$2.9 \cdot 10^{-4}$	$3.2 \cdot 10^{-4}$
$\rho$ + B	$8.9 \cdot 10^{-4}$	$11.8 \cdot 10^{-4}$
$\pi$ + $\rho$	$1.6 \cdot 10^{-4}$	$1.5 \cdot 10^{-4}$
$\pi$ + N(1520)	$0.5 \cdot 10^{-4}$	$0.6 \cdot 10^{-4}$
total yield	$1.7 \cdot 10^{-3}$	$2.0 \cdot 10^{-3}$
experiment [1]	$(8.7 \pm 3.6) \cdot 10^{-3}$	$(6.4 \pm 2.5) \cdot 10^{-3}$

**Table 1.** Multiplicities of  $\phi$  mesons per central event in Ni + Ni and Ru + Ru reactions. The first 5 lines give the results from the special channels indicated. The symbol B comprises the nucleon and the  $\Delta$  particle. The cited experimental yields are to be considered very preliminary.

pointing to the importance of this channel for  $\phi$  meson production. The experimental yields are still underestimated by our calculations, this statement holds independently of the uncertainties of the  $\pi$ N(1520) channel.

#### 4. Conclusions

We have presented a calculation of the  $\phi$  meson production in heavy-ion collisions including in our BUU model the elementary channels  $\rho$ B $\rightarrow \phi$ N,  $\pi$ N(1520) $\rightarrow \phi$ N and  $\pi\rho \rightarrow \phi$  for the first time. As a result the  $\phi$  meson yields are substantially increased but the preliminary experimental data are still underestimated. A dedicated measurement with enlarged statistics would be highly welcome to resolve the  $\phi$  puzzle. For a more detailed discussion of our results see [6].

#### Acknowledgments

We thank R. Kotte for continuous information on his analyses of the FOPI data. This work was supported in part by the German BMBF grant 06DR921, the DAAD scientific exchange program with Hungary, and the National Fund for Scientific Research of Hungary, OTKA T30171, T30855 and T32038.

#### References

- [1] Kotte R (FOPI collaboration) 2000 *Proc. Int. Workshop XXVIII on Gross Properties of Nuclei and Nuclear Excitations, Hirschegg, Austria, January 16 - 22, 2000* p 112
- [2] Chung W S, Li G Q and Ko C M 1997 *Nucl. Phys. A* **625** 347
- [3] Machleidt R 1989 *Adv. Nucl. Phys.* **19** 189
- [4] Hatsuda T and Lee S H 1992 *Phys. Rev. C* **46** R34  
Hatsuda T and Shiomi H 1995 *Nucl. Phys. A* **590** 545c
- [5] Smith W and Haglin K L 1998 *Phys. Rev. C* **57** 1449
- [6] Barz H W, Zétényi M, Wolf Gy and Kämpfer B *Nucl. Phys. A* in press, *Preprint* nucl-th/0110013