

An Introduction to the Interacting Boson Model of the Atomic Nucleus Part II

Walter Pfeifer

87 pp, 17 line figures

1998

Part I of this publication is available in lanl.arXiv.org

Dr. Walter Pfeifer :

Street address:

Stapfenackerweg 9
CH 5034 Suhr
Switzerland

email:

wpfeifer@swissonline.ch

Contents

Contents	2
13 Lie algebras	4
13.1 Definition	4
13.2 The $u(N)$ - and the $su(N)$ -algebra	5
13.3 The $so(N)$ algebra	7
13.4 Dimensions of three classical algebras	8
13.5 Operators constituting Lie algebras, their basis functions and their Casimir operators	9
13.6 Properties of operators	10
14 Group theoretical aspects of the IBM1	13
14.1 Basis operators of the Lie algebras in the IBM1	13
14.2 Subalgebras within the IBM1	15
14.3 The spherical basis as function basis of Lie algebras	18
14.4 Casimir operators of the $u(5)$ - or vibrational limit	20
14.5 Casimir operators of the $su(3)$ - or rotational limit	23
14.6 Casimir operators of the $so(6)$ - or γ -instable limit	28
15 The proton-neutron interacting boson model IBM2	35
15.1 The complete Hamiltonian of the IBM2	35
15.2 Basis states and the angular momentum operator of the IBM2	37
15.3 A reduced Hamiltonian of the IBM2	38
15.4 Eigenenergies and electromagnetic transitions within the IBM2	42
15.5 Comparisons with experimental data	43
15.6 Chains of Lie algebras in the IBM2	54
16 The interacting boson-fermion model IBFM	58
16.1 The Hamiltonian of the IBFM	58
16.2 The $u(5)$ -limit of the IBFM	61
16.3 Numerical treatment of the IBFM. Comparison with experimental data	64
Appendix	69
A1 Clebsch-Gordan coefficients and 3- j symbols	69
A2 Symmetry properties of coupled spin states of identical objects	72
A3 Racah-coefficients and 6- j symbols	73
A4 The 9- j symbol	75
A5 The Wigner-Eckart theorem	76
A6 A further multipole representation of the Hamiltonian	78
A7 The program package PHINT	80
A8 Commutators of operators such as $[d^+ \times d^-]^{(j)}_M$	81
A9 The commutator $[[d^+ \times d^-]^{(j')}_{M'}, [d^+ \times d^+]^{(j)}_M]$	82
References	83

Part I of this publication contains the chapters

- Preface
- 1 Introduction
- 2 characteristics of the IBM
- 3 Many-body configurations
- 4 Many-body states with undefined angular momentum
- 5 Operators and matrix elements
- 6 Applications of the creation and annihilation operators
- 7 The Hamilton operator of the IBM1
- 8 The angular momentum operator of the IBM1
- 9 The Hamiltonian expressed in terms of Casimir operators
- 10 The $u(5)$ - or vibrational limit
- 11 Electromagnetic transitions in the $u(5)$ -limit
- 12 The treatment of the complete Hamiltonian of the IBM1.

It is available in lanl.arXiv.org.

13 Lie algebras

In chapter 9 the importance of the Lie algebras for interpreting and evaluating the IBM-Hamiltonian has been stressed. In the present chapter the essential characteristics of these algebras will be outlined starting from well-known examples. In the first section the rules are described which hold for the elements of a Lie algebra. Looking into special matrices in sections 2 and 3 three classical, real Lie algebras are identified and their dimensions are put together in section 4. In the last section above all Casimir operators (chapter 9 and 14) will be treated.

13.1 Definition

A Lie algebra \mathcal{L} comprises infinitely many elements a, b, c, \dots which show common characteristics. For instance Lie algebras can be represented by matrices with certain symmetry properties or by operators. In particular the elements of a Lie algebra must be able to form Lie products or commutators $[a, b]$. For quadratic matrices a and b the well-known relation

$$[a, b] = ab - ba \quad (13.1)$$

holds, which results again in a matrix of the same type. If the elements of a Lie algebra are operators instead of matrices the commutator $[a, b]$ is still defined but the right hand side of (13.1) need not be satisfied.

The definition of a Lie algebra demands the following properties from its elements a, b, c

a) the commutator of two elements is again an element of the algebra

$$[a, b] \in \mathcal{L} \text{ for all } a, b \in \mathcal{L}, \quad (13.2)$$

b) a linear combination $\alpha a + \beta b$ of two elements with the numbers α and β is again an element of the algebra and the relation

$$[\alpha a + \beta b, c] = \alpha [a, c] + \beta [b, c] \quad (13.3)$$

holds (therefore the element 0 (zero) belongs also to the algebra),

c) interchanging both elements of a commutator results in

$$[a, b] = -[b, a], \quad (13.4)$$

d) finally "Jacobi's identity" has to be satisfied as follows

$$[a, [b, c]] + [b, [c, a]] + [c, [a, b]] = 0. \quad (13.5)$$

Anyhow, other kinds of coupling the elements are possible in principle. As an

example the Casimir operators (13.32) can be mentioned, which result from multiplicative couplings of elements.

Furthermore, the definition demands that a Lie algebra has a finite dimension n i. e. it comprises a set of n elements a_1, a_2, \dots, a_n which act as a basis, by which every element x of the algebra can be represented like this

$$x = \sum_{i=1}^n \xi_i a_i. \quad (13.6)$$

In other words the algebra constitutes a n -dimensional vector space. The structure of the algebra can be formulated in terms of the basis. Because the commutator of two basis elements a_p and a_q belongs also to the algebra, according to (13.6) the relation

$$[a_p, a_q] = \sum_{r=1}^n c_{pq}^{(r)} a_r \quad (13.7)$$

holds. The numbers $c_{pq}^{(r)}$ are named structure constants. One formulates the commutator of the arbitrary elements x (see 13.6) and $y (= \sum_j \zeta_j a_j)$ in terms of the structure constants like this

$$[x, y] = \sum_{i,j=1}^n \xi_i \zeta_j [a_i, a_j] = \sum_{i,j,r=1}^n \xi_i \zeta_j c_{ij}^{(r)} a_r. \quad (13.8)$$

If an algebra corresponds element by element to an other both are isomomorphic and we mark them with the same symbol. In the following we will restrict ourselves to real algebras which are characterised by real coefficients ξ_i in the linear combinations (13.6). With regard to the interacting boson model we will deal with the following types of real algebras: $u(N)$, $su(N)$ and $so(N)$.

13.2 The $u(N)$ - and the $su(N)$ -algebra

We maintain that quadratic and antihermitian matrices of the rank N ($N \times N$ -matrices) form a Lie algebra. Antihermiticity means that the adjoint of the matrix a i. e. the conjugate complex and transposed (reflected on the diagonal) matrix a^\dagger is identical with $-a$ that is

$$a^\dagger = -a. \quad (13.9)$$

Providing real coefficients α and β , for two antihermitian matrices a and b with the same rank naturally the following relation holds

$$(\alpha a + \beta b)^\dagger = \alpha a^\dagger + \beta b^\dagger = -\alpha a - \beta b = -(\alpha a + \beta b). \quad (13.10)$$

Thus linear combinations of antihermitian matrices are again of this type. We make use of the well-known relation

$$(ab)^\dagger = b^\dagger a^\dagger \quad (13.11)$$

in order to look into the commutator of antihermitian matrices as follows

$$[a, b]^\dagger = (ab)^\dagger - (ba)^\dagger = b^\dagger a^\dagger - a^\dagger b^\dagger = ba - ab = -[a, b]. \quad (13.12)$$

Thus the commutator of two antihermitian matrices is again antihermitian.

Now it is not difficult to show that the conditions (13.3) up to (13.5) are met. This means that the antihermitian matrices of the rank N constitute a Lie algebra. It is named $u(N)$ algebra because it is closely related to the Lie group of the unitary matrices.

A matrix is antihermitian if every element of the matrix equals the negative, conjugate complex value of the element which is in the transposed position. Any $N \times N$ -matrix of this kind can be composed linearly by the following quadratic matrices

$$\begin{pmatrix} i & 0 & . \\ 0 & 0 & . \\ . & . & . \end{pmatrix}, \begin{pmatrix} 0 & 0 & . \\ 0 & i & 0 \\ . & 0 & 0 \end{pmatrix}, \dots, \begin{pmatrix} 0 & i & 0 \\ i & 0 & . \\ 0 & . & . \end{pmatrix}, \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & . \\ i & 0 & . & . \\ 0 & . & . & . \end{pmatrix}, \\
 \dots, \begin{pmatrix} 0 & 1 & 0 & . \\ -1 & 0 & . & . \\ 0 & . & . & . \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & . \\ -1 & 0 & . & . \\ 0 & . & . & . \end{pmatrix}, \dots \quad (13.13)$$

Simple counting reveals that there are N^2 matrices in (13.13). As just mentioned they represent the basis of the Lie algebra $u(N)$. The number of basis elements is named dimension n . Thus for the Lie algebra $u(N)$ we have

$$n = N^2. \quad (13.14)$$

Now we will investigate antihermitian matrices of the rank N with vanishing traces (sum of the diagonal elements). We expect that they constitute a Lie algebra as well. Obviously a linear combination of these matrices again has a vanishing trace. Furthermore, for quadratic matrices a and b the relation

$$\text{trace}(ab) = \text{trace}(ba). \quad (13.15)$$

holds. Thus the trace of a commutator vanishes always. This means that the conditions (13.2) up to (13.5) are satisfied and that we have a Lie algebra.

The trace of the basis matrices is always zero if in (13.13) all N diagonal matrices are replaced by $(N-1)$ matrices of the following form

$$\begin{pmatrix} i & 0 & . \\ 0 & -i & 0 \\ . & 0 & . \end{pmatrix}, \begin{pmatrix} 0 & 0 & . \\ 0 & i & 0 \\ . & 0 & -i & 0 \\ . & 0 & . \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & . \\ 0 & . & i \\ . & . & -i \end{pmatrix}, \dots \quad (13.16)$$

Every diagonal, antihermitian matrix with trace 0 can be built by linearly combining matrices from (13.16). The elements (13.16) and the non-diagonal elements of (13.13) constitute the basis of the new Lie algebra. It comprises one matrix less than the one of the $u(N)$ algebra (13.13). Hence its dimension amounts to

$$n = N^2 - 1. \quad (13.17)$$

This algebra represents a special kind of the $u(N)$ algebra and is therefore named $su(N)$.

13.3 The $so(N)$ algebra

The third Lie algebra, which we are investigating here, is constituted by real, antisymmetric and quadratic matrices of the rank N . The transposed a^{tr} of such a matrix a equals to $-a$

$$a^{tr} = -a.$$

Obviously each linear combination of these matrices is still antisymmetric. With the help of

$$(a b)^{tr} = b^{tr} a^{tr} \quad (13.18)$$

one shows in the same way as in the previous section that real, antisymmetric $N \times N$ matrices constitute a Lie algebra. As it is closely related to the Lie group of the orthogonal matrices it is named $o(N)$. Each diagonal element of an antisymmetric matrix is zero, i.e. its trace vanishes. Therefore its Lie algebra is named $so(N)$ as well in analogy to $su(N)$.

The basis elements of this algebra show the following structure

$$\begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & \\ 0 & 0 & \ddots \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & \ddots \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \dots \quad (13.19)$$

and their number is $N(N - 1)/2$. The dimension n of the Lie algebra $so(N)$ is thus

$$n = N(N - 1)/2. \quad (13.20)$$

Specially for $N = 3$ the $so(3)$ basis elements read

$$a_1 = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad a_2 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad a_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}. \quad (13.21)$$

By calculating one obtains the following commutator relation

$$[a_1, a_2] = -a_3 \text{ with cyclic permutations.} \quad (13.22)$$

The structure constants (13.7) of this algebra are therefore 1, 0 or -1. The minus sign in (13.22) disappears if one changes the sign of every basis matrix. The substitution

$$e_1 = (a_1 + ia_2)/\sqrt{2}, \quad e_{-1} = (a_1 - ia_2)/\sqrt{2}, \quad e_0 = -ia_3 \quad \text{yields} \quad (13.23)$$

$$[e_{-1}, e_1] = e_0, \quad [e_{-1}, e_0] = e_{-1}, \quad [e_0, e_1] = e_1. \quad (13.24)$$

Obviously the elements e_1 , e_{-1} and e_0 constitute a Lie algebra with the dimension 3. It is $so(3)$ as can be shown. The commutation relations (13.24) agree completely with the rules (8.7) of the angular momentum operators J_μ . Thus they constitute the Lie algebra $so(3)$.

13.4 Dimensions of three classical algebras

In table 13.1 the lowest dimensions n of the three Lie algebras mentioned above are put together. Group theory shows that there is only a small number of simple and real Lie algebras. Therefore, a simple algebra can be identified directly by means of the dimension. For the IBM1 the n -values of table 13.1 are sufficient. They facilitate to project a given Lie algebra on one of the algebras $u(N)$, $su(N)$ or $so(N)$. The corresponding matrix algebra is a representation of the given Lie algebra. It is possible to transform the basis of one algebra (see 13.23 and 13.24) in order to obtain a corresponding algebra.

Table 13.1. The lowest dimensions n of the Lie algebras $u(N)$, $su(N)$ and $so(N)$ according to (13.14), (13.17) and (13.20).

Algebra : $u(N)$	$su(N)$	$so(N)$	dimension n
Rank : N	N	N	
	2	3	3
2			4
		4	6
	3		8
3			9
		5	10
	4	6	15
4			16
		7	21
	5		24
5			25
		8	28
	6		35
6		9	36

13.5 Operators constituting Lie algebras, their basis functions and their Casimir operators

In the IBM Lie algebras constituted by operators prevail. A very important property of the operators is the existence of sets of basis functions $\psi_k^{(j)}$ which are affiliated to the operators. For example such basis functions are frequently represented by eigenstates of a physical system. They constitute a basis vector, the dimension of which can be chosen. Its elements can be combined linearly and generate a multidimensional vector space. They are characterised by ordering numbers.

In order to clarify the situation we consider the angular momentum operators \mathbf{J}_1 , \mathbf{J}_{-1} and \mathbf{J}_0 (8.6). They obey the commutation rules (8.7)

$$[\mathbf{J}_{-1}, \mathbf{J}_0] = \mathbf{J}_{-1}, \quad [\mathbf{J}_{-1}, \mathbf{J}_1] = \mathbf{J}_0, \quad [\mathbf{J}_0, \mathbf{J}_1] = \mathbf{J}_1. \quad (13.25)$$

In section 13.3 we have seen that they constitute the Lie algebra $so(3)$. On the other hand, from quantum mechanics we know that the operators \mathbf{J}_1 , \mathbf{J}_{-1} and \mathbf{J}_0 generate functions or quantum states φ_{jm} , which are characterised by the quantity j (an integer for bosons). For a given j there are $2j + 1$ functions, which are labelled by m (projection of the spin or of the angular momentum). If an operator \mathbf{J}_i acts on φ_{jm} the result is not an entirely new function but as we know from quantum mechanics the relations

$$\begin{aligned}
J_1 \varphi_{jm} &= -2^{-1/2} [j(j+1) - m(m+1)]^{1/2} \varphi_{j, m+1}, \\
J_{-1} \varphi_{jm} &= 2^{-1/2} [j(j+1) - m(m-1)]^{1/2} \varphi_{j, m-1}, \\
J_0 \varphi_{jm} &= m \varphi_{jm}
\end{aligned} \tag{13.26}$$

hold. Thus when acting on a function φ_{jm} the operators J_1 , J_{-1} or J_0 generate again a function of this kind with the same j . In general an operator of a Lie algebra creates a linear combination of basis functions when it acts on such a function.

We turn to the operator

$$J^2 = J_x^2 + J_y^2 + J_z^2 = -J_1 J_{-1} - J_{-1} J_1 + J_0^2 \tag{13.27}$$

defined in (8.8). With the aid of (13.25) one shows that the following relation holds

$$[J^2, J_i] = 0 \text{ for } i = -1, 0, 1. \tag{13.28}$$

We see that the operator J^2 , which does not belong to the Lie algebra (J_1, J_{-1}, J_0), commutes with the basis elements and therefore with all elements of the algebra. Operators which behave like J^2 are named **Casimir operators**. In physics they are important because they generate eigenvalues. In the case of J^2 we have according to (8.5)

$$J^2 \varphi_{jm} = j(j+1) \varphi_{jm}. \tag{13.29}$$

Thus, when the Casimir operator J^2 acts on the basis function φ_{jm} , this is reproduced and supplied with a factor, which depends only on j .

Now we can put together the properties of the Lie algebras, which we need for the IBM. We refer to the situation of the angular momentum and abstain from group theoretical proofs.

13.6 Properties of operators

We start with basis operators $\Phi^{(i)}$ with $i = 1, \dots, n$ which are linearly independent, constitute a Lie algebra according to (13.2) up to (13.5) and are connected by the structure constants $c^{(r)}_{p,q}$ (13.7). There exist sets of basis functions $\psi^{(j)}_k$ (j stands for the set) on which the operators $\Phi^{(i)}$ act like this

$$\Phi^{(i)} \psi^{(j)}_k = \sum_{l=1}^n \Gamma^{(ij)}_{kl} \psi^{(j)}_l \tag{13.30}$$

analogously to (13.26). The Casimir operators Ξ , which commute with all operators $\Phi^{(i)}$ according to (13.28) as follows

$$[\Xi, \Phi^{(i)}] = 0, \tag{13.31}$$

are formed in terms of the elements $\Phi^{(i)}$. We need the quadratic type Ξ_2 of that operator. The last has the following form

$$\Xi_2 = \sum_{i,j} X_{ij} \Phi^{(i)} \Phi^{(j)}. \quad (13.32)$$

The coefficients X_{ij} of the quadratic Casimir operator depend from the structure constants $c^{(r)}_{p,q}$ (13.7). For example the coefficients of the Casimir operator \mathbf{J}^2 read according to (13.27) : $X_{1,-1} = X_{-1,1} = -1$, $X_{00} = 1$ and all other vanish.

The sets of functions $\psi^{(j)}_k$, which have so-called irreducible matrices $\Gamma^{(ij)}_{kl}$ (13.30), generate eigenvalues $C^{(j)}_2$ under the influence of the Casimir operator Ξ_2 like this

$$\Xi_2 \psi^{(j)}_k = C^{(j)}_2 \psi^{(j)}_k. \quad (13.33)$$

According to (13.29) the eigenvalue of the quadratic Casimir operator \mathbf{J}^2 of the Lie algebra $so(3)$ has the form $j(j+1)$.

Using creation and annihilation operators for bosons, Frank and Van Isacker (1994, p. 310) show that the eigenvalues of the Casimir operators of the $so(N)$ -algebras have the following form

$$C^{(\rho)}_{2, so(N)} = \rho(\rho + N - 2). \quad (13.34)$$

The quantity N is the matrix dimension of the so -algebra and ρ is an integer which is limited by a function of the boson number. For instance for $N = 3$ (13.34) agrees with (13.29), provided that ρ is replaced by the angular momentum J of the state. We know that in the $u(5)$ -limit of the IBM the quantity J has the values $2n_d, 2n_d - 2, 2n_d - 3, \dots, 1, 0$. In the next chapter we will meet further applications of formula (13.34).

For the Lie algebra $su(3)$ Cornwell (1990, p. 596) derives the following eigenvalue of the quadratic Casimir operator (apart from a factor $1/9$)

$$C^{(n,m)}_{2, su(3)} = n^2 + m^2 + nm + 3n + 3m. \quad (13.35)$$

The integers n and m characterise the irreducible matrices $(\Gamma^{(i,n,m)}_{kl})$ (13.30) of the algebra and the affiliated functions $\psi^{(n,m)}_l$. The expression (13.35) will arise in the next chapter in connection with the second special case of the IBM1.

The basis operators of the IBM (chapter 14) not only constitute a Lie algebra in all but they contain subsets which are subalgebras. This means for instance that commutators of operators of a subalgebra are linear combinations of elements of this subset (13.2). The Casimir operator of the higher algebra commutes with the Casimir operator of the subalgebra, which follows from $\Xi \Phi^{(i)} = \Phi^{(i)} \Xi$ (13.31) and from (13.32). The Casimir operators of the subalgebras, which we will meet in the next chapter, are summands in the Casimir operators of the higher algebra.

Furthermore, it is true that the basis function $\psi^{(j)}_k$ (13.30) of the higher algebra is also the basis function of the subalgebra. Therefore this function is labelled both with the index of the Casimir eigenvalue C_2 of the higher algebra and with the corresponding index of the subalgebra. For example in the basis function

φ_{jm} (13.29) the quantity m is the characteristic parameter of the subalgebra $so(2)$ (consisting only of the operator $\mathbf{J}_z = \mathbf{J}_0$) and j is the index of the higher algebra $so(3)$, which comprises \mathbf{J}_{-1} , \mathbf{J}_1 and \mathbf{J}_0 .

14 Group theoretical aspects of the IBM1

In this chapter first we will show that the IBM-Hamiltonian is composed of elements which constitute a Lie algebra. In the second section the affiliated subalgebras are looked into. Three chains of subalgebras can be formulated. The third section is rather illustrative and shows how the basis operators of the IBM act on the accompanying basis functions. In sections 4, 5 and 6 special cases of the IBM are treated. The eigenvalues of the Casimir operators which belong to these cases, determine the energy eigenvalues. Comparisons with measured level schemes will be performed.

14.1 Basis operators of the Lie algebras in the IBM1

In this section we show first that the Hamilton operator of the IBM1 is composed exclusively of the following operators

$$[\mathbf{d}^+ \times \mathbf{d}^-]^{(J)}_M, [\mathbf{d}^+ \times \mathbf{s}]^{(2)}_M, [\mathbf{s}^+ \times \mathbf{d}^-]^{(2)}_M \text{ and } [\mathbf{s}^+ \times \mathbf{s}]^{(0)}_0 \quad (14.1)$$

Owing to (9.37) we can write the Hamiltonian like this

$$H = \varepsilon_n \mathbf{N} + \varepsilon_d' n_d + v_n \mathbf{N}^2 + v_{nd} \mathbf{N} n_d + v_d n_d^2 + v_r \mathbf{R}^2 + v_t \mathbf{T}^2 + v_j \mathbf{J}^2 + v_q \mathbf{Q}^2. \quad (14.2)$$

The cited operators - apart from \mathbf{R}^2 and \mathbf{T}^2 - have the following structure (9.1), (9.4)

$$\begin{aligned} n_d &= \sqrt{5} [\mathbf{d}^+ \times \mathbf{d}^-]^{(0)}_0, \\ \mathbf{N} &= \mathbf{s}^+ \mathbf{s} + n_d = [\mathbf{s}^+ \times \mathbf{s}]^{(0)}_0 + \sqrt{5} [\mathbf{d}^+ \times \mathbf{d}^-]^{(0)}_0, \\ \mathbf{J}^2 &= -\sqrt{3} \cdot 10 [[\mathbf{d}^+ \times \mathbf{d}^-]^{(1)} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(1)}]^{(0)}_0 = \\ &= 10 \sum_{\mu} (-1)^{\mu} [\mathbf{d}^+ \times \mathbf{d}^-]^{(1)}_{\mu} [\mathbf{d}^+ \times \mathbf{d}^-]^{(1)}_{-\mu}, \\ \mathbf{Q}^2 &= \sum_{\mu} (-1)^{\mu} \mathbf{Q}_{\mu} \mathbf{Q}_{-\mu} \end{aligned} \quad (14.3)$$

$$\text{with } \mathbf{Q}_{\mu} = [\mathbf{d}^+ \times \mathbf{s}]^{(2)}_{\mu} + [\mathbf{s}^+ \times \mathbf{d}^-]^{(2)}_{\mu} - (\sqrt{7}/2) [\mathbf{d}^+ \times \mathbf{d}^-]^{(2)}_{\mu}.$$

Thus these operators are built by elements of the type (14.1). This is also true for both remaining operators in (14.2), which we will show by remodelling. We write the seniority operator $\mathbf{T}^2 = n_d (n_d + 3) - 5 [\mathbf{d}^+ \times \mathbf{d}^-]^{(0)}_0 [\mathbf{d}^- \times \mathbf{d}^+]^{(0)}_0$ (9.2) with the aid of (9.12) this way

$$\mathbf{T}^2 = n_d^2 + 4n_d - \sum_{J'} \sqrt{(2J' + 1)} \cdot [[\mathbf{d}^+ \times \mathbf{d}^-]^{(J')} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(J')}]^{(0)}.$$

We replace n_d^2 making use of (9.20)

$$\begin{aligned} n_d^2 &= 5 [[\mathbf{d}^+ \times \mathbf{d}^-]^{(0)} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(0)}]^{(0)} = \\ &= \sum_{J''} (-1)^{J''} \sqrt{(2J'' + 1)} [[\mathbf{d}^+ \times \mathbf{d}^-]^{(J'')} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(J'')}]^{(0)} - 4 n_d \end{aligned} \quad (14.4)$$

and obtain

$$T^2 = -2 \sum_{J'=1,3} \sqrt{(2J'+1)} \cdot [[d^+ \times d^-]^{(J')} \times [d^+ \times d^-]^{(J')}]^{(0)}. \quad (14.5)$$

The remaining operator R^2 (9.3) reads

$$\begin{aligned} R^2 = & N(N+4) - (\sqrt{5} \cdot [d^+ \times d^+]^{(0)} - s^+ s^+) (\sqrt{5} [d^- \times d^-]^{(0)} - ss) = \\ & N(N+4) - 5 \cdot [[d^+ \times d^+]^{(0)} \times [d^- \times d^-]^{(0)}]^{(0)} + \sqrt{5} [d^+ \times d^+]^{(0)} ss + \\ & \sqrt{5} s^+ s^+ [d^- \times d^-]^{(0)} - s^+ s^+ ss. \end{aligned} \quad (14.6)$$

With the help of (9.2) we obtain

$$\begin{aligned} R^2 = & N(N+4) - T^2 - n_d(n_d+3) + \sqrt{5} [d^+ \times d^+]^{(0)} ss + \\ & \sqrt{5} s^+ s^+ [d^- \times d^-]^{(0)} - s^+ s^+ ss, \end{aligned}$$

insert T^2 from (14.5) and transform like this

$$\begin{aligned} R^2 = & N(N+4) - n_d(n_d+3) - 2 \sum_{J'=1,3} \sqrt{(2J'+1)} \cdot [[d^+ \times d^-]^{(J')} \times [d^+ \times d^-]^{(J')}]^{(0)} + \\ & \sqrt{5} [[d^+ \times s]^{(2)} \times [d^+ \times s]^{(2)}]^{(0)} + \sqrt{5} [[s^+ \times d^-]^{(2)} \times [s^+ \times d^-]^{(2)}]^{(0)} - \\ & [[s^+ \times s]^{(0)} \times [s^+ \times s]^{(0)}]^{(0)} + [s^+ \times s]^{(0)}. \end{aligned} \quad (14.7)$$

Thus, the entire Hamilton operator is built by operators of the type (14.1).

Now we show that the operators (14.1) constitute a Lie algebra. For the moment we are restricting ourselves to operators such as $[d^+ \times d^-]^{(J)}_M$. According to (A8.4) the commutation relation

$$\begin{aligned} & [[d^+ \times d^-]^{(J')}_{M'}, [d^+ \times d^-]^{(J'')_{M''}}] = \\ & \sum_{\kappa \kappa'} [d^+ \times d^-]^{(k)}_{\kappa} \cdot \sqrt{(2J'+1)} \sqrt{(2J''+1)} \cdot \\ & (J' M' J'' M'' | k \kappa) \left\{ \begin{matrix} J' & J'' & k \\ 2 & 2 & 2 \end{matrix} \right\} (-1)^k (1 - (-1)^{J'+J''+k}) \end{aligned} \quad (14.8)$$

holds. That is, the commutator of two operators $[d^+ \times d^-]^{(J)}_M$ is a linear combination of such operators. Therefore the defining conditions (13.2) and (13.7) for Lie algebras are satisfied. Interchanging both operators in the commutator (14.8) results in altering the sign of the Clebsch-Gordan coefficient (consult (A1.4) and the relation $J' + J'' + k = \text{odd}$). Consequently the condition (13.4) is met and one can show also that the relations (13.3) and (13.5) are satisfied. Therefore the operators of the type $[d^+ \times d^-]^{(J)}_M$ constitute a Lie algebra.

The operators in (14.1) which contain s -operators satisfy the following commutation relations

$$\begin{aligned}
[[\mathbf{s}^+ \times \mathbf{s}]^{(0)}_0, [\mathbf{s}^+ \times \mathbf{d}^\sim]^{(2)}_\mu] &= [\mathbf{s}^+ \times \mathbf{d}^\sim]^{(2)}_\mu, \\
[[\mathbf{d}^+ \times \mathbf{s}]^{(2)}_\mu, [\mathbf{s}^+ \times \mathbf{s}]^{(0)}_0] &= [\mathbf{d}^+ \times \mathbf{s}]^{(2)}_\mu, \\
[[\mathbf{s}^+ \times \mathbf{d}^\sim]^{(2)}_\kappa, [\mathbf{s}^+ \times \mathbf{d}^\sim]^{(2)}_\mu] &= [[\mathbf{d}^+ \times \mathbf{s}]^{(2)}_\kappa, [\mathbf{d}^+ \times \mathbf{s}]^{(2)}_\mu] = 0, \\
[[\mathbf{d}^+ \times \mathbf{s}]^{(2)}_\kappa, [\mathbf{s}^+ \times \mathbf{d}^\sim]^{(2)}_\mu] &= \\
\sum_{JM} (2 \kappa 2 \mu | J M) [\mathbf{d}^+ \times \mathbf{d}^\sim]^{(J)}_M - \delta_{\kappa, -\mu} (-1)^\mu [\mathbf{s}^+ \times \mathbf{s}]^{(0)}_0, \\
[[\mathbf{s}^+ \times \mathbf{d}^\sim]^{(2)}_\mu, [\mathbf{d}^+ \times \mathbf{d}^\sim]^{(J)}_M] &= (2, -\mu, 2, \mu + M | J M) (-1)^\mu [\mathbf{s}^+ \times \mathbf{d}^\sim]^{(2)}_{\mu + M}, \\
[[\mathbf{d}^+ \times \mathbf{s}]^{(2)}_\mu, [\mathbf{d}^+ \times \mathbf{d}^\sim]^{(J)}_M] &= - (2, \mu + M, 2, -\mu | J M) (-1)^\mu [\mathbf{d}^+ \times \mathbf{s}]^{(2)}_{\mu + M}
\end{aligned} \tag{14.9}$$

which is shown similarly as in (A8.4). From (14.8) and (14.9) follows that all operators listed in (14.1) constitute a Lie algebra.

14.2 Subalgebras within the IBM1

We now will investigate how this algebra can be subdivided in subalgebras. We start with the elements $[\mathbf{d}^+ \times \mathbf{d}^\sim]^{(1)}_\mu$. According to (8.9) these operators represent the ladder operators J_μ of the angular momentum apart from the factor $\sqrt{10}$. Owing to (13.24) they constitute the Lie algebra $so(3)$.

We turn to elements of the type

$$[\mathbf{d}^+ \times \mathbf{d}^\sim]^{(J)}_\mu \text{ with } J = 1, 3. \tag{14.10}$$

According to (14.8) the commutator of two elements of this kind creates a linear combination of operators $[\mathbf{d}^+ \times \mathbf{d}^\sim]^{(J)}_\mu$ (with $J = 1, 3$). Obviously they also meet the other definition conditions (section 13.1) and constitute a Lie algebra of the dimension (= number of basis elements) $n = 3 + 7 = 10$. Table 13.1 contains only one Lie algebra with $n = 10$ namely the algebra $so(5)$.

The totality of the elements $[\mathbf{d}^+ \times \mathbf{d}^\sim]^{(J)}_M$ with $0 \leq J \leq 4$ also makes up a Lie algebra. By means of its dimension $n = 1 + 3 + 5 + 6 + 9 = 25$ we learn from table 13.1 that it is the algebra $u(5)$.

If one adds the elements $[\mathbf{d}^+ \times \mathbf{s}]^{(2)}_M$, $[\mathbf{s}^+ \times \mathbf{d}^\sim]^{(2)}_M$ and $[\mathbf{s}^+ \times \mathbf{s}]^{(0)}_0$, the resulting 36 elements constitute a Lie algebra, which has been mentioned in the last section. Table 13.1 reveals that it is the algebra $u(6)$.

Consequently the basis elements (14.1) constitute the following chain of Lie algebras

$$I. \quad u(6) \supset u(5) \supset so(5) \supset so(3). \tag{14.11}$$

It defines the vibrational or $u(5)$ limit (chapter 10), which will be reviewed in section 14.4 once more.

Now we are claiming that the elements

$$[\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(1)} \text{ and } \mathbf{Q}_{\mu} = [\mathbf{d}^+ \times \mathbf{s}]_{\mu}^{(2)} + [\mathbf{s}^+ \times \mathbf{d}^-]_{\mu}^{(2)} - (\sqrt{7}/2) [\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(2)} \quad (14.12)$$

make up a Lie algebra. First we investigate the commutator

$$\begin{aligned} [\mathbf{Q}_{\mu}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(1)}] &= [[\mathbf{s}^+ \times \mathbf{d}^-]_{\mu}^{(2)}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(1)}] + \\ & [[\mathbf{d}^+ \times \mathbf{s}]_{\mu}^{(2)}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(1)}] - (\sqrt{7}/2) [[\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(2)}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(1)}]. \end{aligned} \quad (14.13)$$

With the aid of (14.8), (14.9), (A1.4) up to (A1.6) and (A3.14) one obtains finally

$$[\mathbf{Q}_{\mu}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(1)}] = (-1)^{\mu} (2, -\mu, 2, \mu + \kappa | 1 \kappa) \mathbf{Q}_{\mu + \kappa}. \quad (14.14)$$

The Clebsch-Gordan coefficient prevents that $|\mu + \kappa|$ exceeds 2. Furthermore, we need the commutator $[\mathbf{Q}_{\mu}, \mathbf{Q}_{\kappa}]$, for which we write the non vanishing terms

$$\begin{aligned} [\mathbf{Q}_{\mu}, \mathbf{Q}_{\kappa}] &= [[\mathbf{d}^+ \times \mathbf{s}]_{\mu}^{(2)}, [\mathbf{s}^+ \times \mathbf{d}^-]_{\kappa}^{(2)}] - [[\mathbf{d}^+ \times \mathbf{s}]_{\kappa}^{(2)}, [\mathbf{s}^+ \times \mathbf{d}^-]_{\mu}^{(2)}] - \\ & (\sqrt{7}/2) [[\mathbf{s}^+ \times \mathbf{d}^-]_{\mu}^{(2)}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(2)}] + (\sqrt{7}/2) [[\mathbf{s}^+ \times \mathbf{d}^-]_{\kappa}^{(2)}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(2)}] - \\ & (\sqrt{7}/2) [[\mathbf{d}^+ \times \mathbf{s}]_{\mu}^{(2)}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(2)}] + (\sqrt{7}/2) [[\mathbf{d}^+ \times \mathbf{s}]_{\kappa}^{(2)}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(2)}] + \\ & (7/4) [[\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(2)}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(2)}]. \end{aligned} \quad (14.15)$$

The four terms with the factor $\sqrt{7}/2$ in (14.15) cancel out in pairs because of (14.9), (A1.4) and (A1.6). By means of the explicit values of the 6- j symbols (A3.14)

$$\left\{ \begin{matrix} 2 & 2 & 1 \\ 2 & 2 & 2 \end{matrix} \right\} = -1/10, \quad \left\{ \begin{matrix} 2 & 2 & 3 \\ 2 & 2 & 2 \end{matrix} \right\} = 4/35$$

and of (14.8) and (14.9) one obtains the following result

$$[\mathbf{Q}_{\mu}, \mathbf{Q}_{\kappa}] = (15/4) (2 \mu 2 \kappa | 1, \kappa + \mu) [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa + \mu}^{(1)}. \quad (14.16)$$

The Clebsch-Gordan coefficient prevents that $|\mu + \kappa|$ exceeds 1. In this way the choice of the factor $\sqrt{7}/2$ is explained. It causes the term $[\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(3)}$ in (14.16) to vanish. Clearly the factor $\sqrt{7}/2$ might have also a positive sign in (14.12) because in (14.15) it appears only in squared form.

Thus the operators (14.12) constitute a Lie algebra with the dimension $n = 3 + 5 = 8$, which is in fact the $su(3)$ algebra according to table 13.1. It includes the elements $[\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(1)}$, which make up the algebra $so(3)$, as we know.

If we add the operators $[\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(k)}$ ($k = 0, 3, 4$), $[\mathbf{s}^+ \times \mathbf{d}^-]_{\mu}^{(2)}$, $[\mathbf{d}^+ \times \mathbf{s}]_{\mu}^{(2)}$ and $[\mathbf{s}^+ \times \mathbf{s}]_0^{(0)}$ to the 8 elements of (14.12) all possible commutators create linear combinations of these 36 elements. In case single elements $[\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(2)}$ are generated, they have to be replaced by $([\mathbf{s}^+ \times \mathbf{d}^-]_{\mu}^{(2)} + [\mathbf{d}^+ \times \mathbf{s}]_{\mu}^{(2)} - \mathbf{Q}_{\mu}) 2/\sqrt{7}$. The totality constitutes the algebra $u(6)$, which is subdivided in subalgebras like this

$$\text{II.} \quad u(6) \supset su(3) \supset so(3). \quad (14.17)$$

This chain represents the $su(3)$ limit, which is named rotational limit as well. In section 14.5 it will be looked into.

We come to the third special case of the IBM by the following set of elements

$$[\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(J)} (J = 1, 3) \text{ and } \mathbf{S}_{\lambda} = [\mathbf{d}^+ \times \mathbf{s}]_{\lambda}^{(2)} + [\mathbf{s}^+ \times \mathbf{d}^-]_{\lambda}^{(2)}. \quad (14.18)$$

The first group of operators in (14.18) makes up the algebra $so(5)$ as we know from (14.11). According to (14.9) the relation

$$[\mathbf{S}_{\lambda}, \mathbf{S}_{\mu}] = 2 \sum_{J=1,3} (2 \lambda \ 2 \ \mu | J, \lambda + \mu) [\mathbf{d}^+ \times \mathbf{d}^-]_{\lambda + \mu}^{(J)} \quad (14.19)$$

holds and for odd J we have

$$[\mathbf{S}_{\lambda}, [\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(J)}] = -((2J+1)/5)^{1/2} (2 \lambda \ J \ \mu | 2, \lambda + \mu) ([\mathbf{d}^+ \times \mathbf{s}]_{\lambda + \mu}^{(2)} + [\mathbf{s}^+ \times \mathbf{d}^-]_{\lambda + \mu}^{(2)}). \quad (14.20)$$

Therefore, the elements of (14.18) constitute a Lie algebra of the dimension $n = 3 + 7 + 5 + = 15$. In table 13.1 the algebras $su(4)$ and $so(6)$ have this dimension. They are isomorphic as can be shown.

Adding the elements $[\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(J)} (J = 0, 3, 4)$, $[\mathbf{s}^+ \times \mathbf{d}^-]_{\mu}^{(2)}$ and $[\mathbf{s}^+ \times \mathbf{s}]_0^{(0)}$ to the set (14.18) one obtains 36 elements constituting the $u(6)$ algebra, which is shown analogously to the derivation of (14.17). Thus we obtain the following chain of algebras

$$\text{III.} \quad u(6) \supset so(6) \supset so(5) \supset so(3). \quad (14.21)$$

This is the third special case of the IBM, which is named $so(6)$ or γ -instable limit.

In table 14.1 the chains of algebras of the IBM1 and the affiliated basis elements are put together.

Table 14.1. Chains of algebras for the three special cases of the IBM1

Basis elements	dimension	algebra	
$[s^+ \times d^-]_{\mu}^{(2)}, [d^+ \times s]_{\mu}^{(2)}, [s^+ \times s]_0^{(0)}$	$\left. \begin{array}{c} n \\ 36 \end{array} \right\}$	$u(6)$	vibrational limit
$[d^+ \times d^-]_0^{(0)}, [d^+ \times d^-]_{\mu}^{(2)}, [d^+ \times d^-]_{\mu}^{(4)}$	$\left. \begin{array}{c} 25 \end{array} \right\}$	$u(5)$	
$[d^+ \times d^-]_{\mu}^{(3)}$	$\left. \begin{array}{c} 10 \end{array} \right\}$	$so(5)$	
$[d^+ \times d^-]_{\mu}^{(1)}$	$\left. \begin{array}{c} 3 \end{array} \right\}$	$so(3)$	
$[s^+ \times d^-]_{\mu}^{(2)}, [d^+ \times s]_{\mu}^{(2)}, [s^+ \times s]_0^{(0)}, [d^+ \times d^-]_0^{(0)}, [d^+ \times d^-]_{\mu}^{(3)}, [d^+ \times d^-]_{\mu}^{(4)}$	$\left. \begin{array}{c} 36 \end{array} \right\}$	$u(6)$	rotational limit
$[s^+ \times d^-]_{\mu}^{(2)} + [d^+ \times s]_{\mu}^{(2)} - (\sqrt{7/2})[d^+ \times d^-]_{\mu}^{(2)}$	$\left. \begin{array}{c} 8 \end{array} \right\}$	$su(3)$	
$[d^+ \times d^-]_{\mu}^{(1)}$	$\left. \begin{array}{c} 3 \end{array} \right\}$	$so(3)$	
$[s^+ \times d^-]_{\mu}^{(2)}, [s^+ \times s]_0^{(0)}, [d^+ \times d^-]_0^{(0)}, [d^+ \times d^-]_{\mu}^{(2)}, [d^+ \times d^-]_{\mu}^{(4)}$	$\left. \begin{array}{c} 36 \end{array} \right\}$	$u(6)$	γ -instable limit
$[s^+ \times d^-]_{\mu}^{(2)} + [d^+ \times s]_{\mu}^{(2)}$	$\left. \begin{array}{c} 15 \end{array} \right\}$	$so(6)$	
$[d^+ \times d^-]_{\mu}^{(3)}$	$\left. \begin{array}{c} 10 \end{array} \right\}$	$so(5)$	
$[d^+ \times d^-]_{\mu}^{(1)}$	$\left. \begin{array}{c} 3 \end{array} \right\}$	$so(3)$	

14.3 The spherical basis as function basis of Lie algebras

In chapter 6 we have treated the seniority scheme and represented its boson states like this (6.18)

$$|n_s n_d \tau n_{\Delta} J M\rangle = ((2\tau + 3)!! / (n_s! (n_d + \tau + 3)!! (n_d - \tau)!!))^{1/2} (s^+)^{n_s} (P^+)^{(n_d - \tau)/2} | \tau \tau n_{\Delta} J M \rangle \quad (14.22)$$

with $P^+ = \sqrt{5} [d^+ \times d^+]_0^{(0)}$ (6.12).

These so-called spherical basis states are assigned to the $u(5)$ - or vibrational limit (chapter 10) and we know that they are eigenfunctions of the operators J^2 and T^2 , which are components of the corresponding Hamiltonian. As an

exercise we check if these states behave like basis functions of the $u(5)$ limit and if they satisfy the relation (13.30). We restrict ourselves to the $so(5)$ subalgebra. It comprises the elements

$$[\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(1)} \text{ and } [\mathbf{d}^+ \times \mathbf{d}^-]_{\nu}^{(3)}. \quad (14.23)$$

According to (13.30) we expect the following behaviour: when an operator of the type (14.23) acts on a basis function (14.22) a linear combination of such functions arises.

The equations (13.26) show that the first operator in (14.23) satisfies this condition because it agrees with the angular momentum operator \mathbf{J}_{μ} (8.9) apart from a factor.

Now we let the operator $[\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(3)}$ act on the state (14.22) and move it to the right in front of the operator \mathbf{P}^+ . Then we maintain that these operators commute. In order to show it, according to (A9.3) and (A1.16) we can write for odd k

$$\begin{aligned} & [[\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(k)}, [\mathbf{d}^+ \times \mathbf{d}^-]_0^{(0)}] = \\ & 2 \sum_{\mu} (2 \mu 2, -\mu | 0 0) (2, \kappa + \mu, 2, -\mu | k \kappa) (-1)^{\mu} \mathbf{d}_{\kappa+\mu}^+ \mathbf{d}_{-\mu}^+ = \\ & (2/\sqrt{5}) \sum_{\mu} (2, \kappa + \mu, 2, -\mu | k \kappa) \mathbf{d}_{\kappa+\mu}^+ \mathbf{d}_{-\mu}^+ = (2/\sqrt{5}) [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(k)}. \end{aligned} \quad (14.24)$$

Because k is odd the last expression vanishes owing to (6.6). Therefore $[\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(3)}$ ($k = 1, 3$) can be interchanged with \mathbf{P}^+ . Generally the relation

$$[\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(k)} (\mathbf{P}^+)^{(n_d - \tau)/2} = (\mathbf{P}^+)^{(n_d - \tau)/2} [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(k)}, \quad k = 1, 3, \dots, \quad (14.25)$$

holds. This means that the $so(5)$ operator (14.23) can be moved in front of the state $|n_s \tau \tau n_d J M\rangle$ with maximal seniority in (14.22). For the time being we are restricting ourselves to states with $n_d = \tau = 2$. They read according to (6.5)

$$|n_d = \tau = 2, J M\rangle = (2)^{-1/2} [\mathbf{d}^+ \times \mathbf{d}^-]_{M}^{(J)} | \rangle \text{ with } J = 2, 4. \quad (14.26)$$

Acting on the state (14.26) the $so(5)$ operator $[\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(3)}$ generates the expression

$$\begin{aligned} & [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(3)} (2)^{-1/2} [\mathbf{d}^+ \times \mathbf{d}^-]_{M}^{(J)} | \rangle = \\ & \sqrt{2} \sum_{\mu} (2 \mu 2, M - \mu | J M) (2, \kappa + \mu, 2, -\mu | 3 \kappa) (-1)^{\mu} \mathbf{d}_{\kappa+\mu}^+ \mathbf{d}_{M-\mu}^+, \end{aligned} \quad (14.27)$$

which is shown by means of (9.3) and the relation $\mathbf{d}_{\mu} | \rangle = 0$. With the aid of (A1.14) the pair of \mathbf{d}^+ -operators is transformed to expressions such as $[\mathbf{d}^+ \times \mathbf{d}^-]_{M'}^{(J')}$. Because we are interested in the question if $J' = 0$ arises, we investigate situations with $M' = 0$. That is we put $M = -\kappa$ in (14.27) and obtain

$$\begin{aligned}
& [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(3)} (2)^{-1/2} [\mathbf{d}^+ \times \mathbf{d}^+]_{-\kappa}^{(J)} | \rangle = \\
& \sqrt{2} \sum_{\mu} (2 \mu 2, -\kappa - \mu | J, -\kappa) (2, \kappa + \mu, 2, -\mu | 3 \kappa) (-1)^{\mu} \mathbf{d}^+_{\kappa+\mu} \mathbf{d}^+_{-\kappa-\mu} = \\
& \sqrt{2} \sum_{\mu} (2 \mu 2, -\kappa - \mu | J, -\kappa) (2, \kappa + \mu, 2, -\mu | 3 \kappa) (-1)^{\mu} \cdot \\
& \sum_{J'} (2, \kappa + \mu, 2, -\kappa - \mu | J' 0) [\mathbf{d}^+ \times \mathbf{d}^+]_{0}^{(J')} | \rangle. \quad (14.28)
\end{aligned}$$

With the help of (A1.16), (A1.4) and (A1.5) we write the coefficient for $J' = 0$ as follows

$$\sqrt{(2/5)} (-1)^{\kappa} \sum_{\mu} (2 \mu 2, -\kappa - \mu | J, -\kappa) (2 \mu 2, -\kappa - \mu | 3, -\kappa) = 0. \quad (14.29)$$

Owing to the orthogonality relation (A1.10) this expression vanishes because J has only the values 2 or 4 (14.26).

Thus we have shown that the operator $[\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(3)}$ acting on the state (14.26) which has maximal seniority creates a linear combination of states of the same kind. In other words, when the operator $[\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(3)}$ acts on a state with seniority 2, the resulting states have the same seniority. That is, the rule (13.30) for Lie algebras is satisfied.

It's obvious that this statement holds also for states with $\tau = 1$, i. e. $\mathbf{d}^+ | \rangle$. For $\tau = 3$ it can be checked with the aid of (6.9). The proof of the general relation

$$\begin{aligned}
& [\mathbf{d}^+ \times \mathbf{d}^-]_{\kappa}^{(3)} | N n_d \tau n_{\Delta} J M \rangle = \\
& \sum_{n'_{\Delta} J' M'} \Gamma^{(3, \kappa)}_{N n_d \tau n_{\Delta} J M n'_{\Delta} J' M'} | N n_d \tau n'_{\Delta} J' M' \rangle \quad (14.30)
\end{aligned}$$

is laborious.

14.4 Casimir operators of the $u(5)$ - or vibrational limit

In chapter 9 the Hamilton operator was transformed in a combination of operators which we named Casimir operators. In sections 14.4 up to 14.6 we will show now that this naming is correct. We begin with the special form (10.1) of the Hamiltonian characterising the $u(5)$ limit

$$H^{(1)} = \varepsilon_n \mathbf{N} + v_n \mathbf{N}^2 + (\varepsilon_d' + v_{nd} \mathbf{N}) \mathbf{n}_d + v_d \mathbf{n}_d^2 + v_t \mathbf{T}^2 + v_j \mathbf{J}^2. \quad (14.31)$$

We will investigate the properties of the operators in (14.31), confront it with the rules (13.31) up to (13.33) and make visible how they are related to the subalgebras of the $u(5)$ limit.

According to (13.28) the operator $\mathbf{J}^2 = -10\sqrt{3} [[\mathbf{d}^+ \times \mathbf{d}^-]^{(1)} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(1)}]^{(0)}$ (8.14) commutes with the components $\mathbf{J}_{\mu} = \sqrt{10} [\mathbf{d}^+ \times \mathbf{d}^-]_{\mu}^{(1)}$, $\mu = 1, 0, -1$ (8.9), which constitute the Lie algebra $so(3)$ (13.24). Clearly \mathbf{J}^2 must be its quadratic Casimir operator. It satisfies not only (13.31) and (13.32) but it generates also the eigenvalue $J(J+1)$ and therefore it meets the conditions (13.34).

We expect that

$$\begin{aligned} \mathbf{T}^2 = & -2 \sum_{J=1,3} \sqrt{(2J+1)} \cdot [[\mathbf{d}^+ \times \mathbf{d}^-]^{(J)} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(J)}]^{(0)} = \\ & 2 \sum_{J=1,3} \sum_M (-1)^M [\mathbf{d}^+ \times \mathbf{d}^-]^{(J)}_M [\mathbf{d}^+ \times \mathbf{d}^-]^{(J)}_{-M} \end{aligned}$$

is the quadratic Casimir operator of the higher algebra $so(5)$. Firstly it is composed of products of operators $[\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}_\kappa$, $k = 1, 3$ of this algebra and obeys therefore the relation (13.32). According to (10.13) \mathbf{T}^2 meets the condition (13.33) and creates the eigenvalue $\tau(\tau + 3)$, which has the form (13.34), because the rank N amounts to 5. According to (10.13) and (14.30) we have

$$\begin{aligned} [\mathbf{d}^+ \times \mathbf{d}^-]^{(3)}_\kappa \mathbf{T}^2 |N n_d \tau n_\Delta J M\rangle &= \tau(\tau + 3) [\mathbf{d}^+ \times \mathbf{d}^-]^{(3)}_\kappa |N n_d \tau n_\Delta J M\rangle = \\ \tau(\tau + 3) \sum_{n'_\Delta J' M'} \Gamma^{(3, \kappa)}_{N n_d \tau n_\Delta J M n'_\Delta J' M'} |N n_d \tau n'_\Delta J' M'\rangle &= \\ \mathbf{T}^2 \sum_{n'_\Delta J' M'} \Gamma^{(3, \kappa)}_{N n_d \tau n_\Delta J M n'_\Delta J' M'} |N n_d \tau n'_\Delta J' M'\rangle &= \\ \mathbf{T}^2 [\mathbf{d}^+ \times \mathbf{d}^-]^{(3)}_\kappa |N n_d \tau n_\Delta J M\rangle. \end{aligned} \quad (14.32)$$

Thus the condition (13.31) is met, which is valid also for the basis operator $[\mathbf{d}^+ \times \mathbf{d}^-]^{(1)}_\kappa$.

The next higher algebra $u(5)$ in this special case of the IBM1 comprises all elements $[\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}_\kappa$, $k = 0$ up to 4. According to (14.4) the relation

$$\begin{aligned} n_d^2 + 4 n_d = \sum_{k=0}^4 (-1)^k \sqrt{(2k+1)} [[\mathbf{d}^+ \times \mathbf{d}^-]^{(k)} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}]^{(0)} = \\ \sum_{k=0}^4 \sum_\kappa (-1)^\kappa [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}_\kappa [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}_{-\kappa} \end{aligned} \quad (14.33)$$

holds. This operator must be the quadratic Casimir operator of the algebra $u(5)$ because it satisfies (13.31) up to (13.33). We guess that the number operator $n_d = \sum_\mu (-1)^\mu \mathbf{d}^+_\mu \mathbf{d}^-_{-\mu}$ is the linear Casimir operator of the $u(5)$ algebra.

The highest algebra in the $u(5)$ chain, $u(6)$, comprises all 36 elements of the type $[\mathbf{b}^+_{l'} \times \mathbf{b}^-_{l''}]^{(k)}_\kappa$ ($l, l' = 0, 2$; $\mathbf{b}_0 = \mathbf{s}$, $\mathbf{b}_{2,\mu} = \mathbf{d}_\mu$). We maintain that its quadratic Casimir operator reads as follows in agreement with (13.32)

$$\begin{aligned} \mathbf{C}_{u(6)} &= \sum_{l, l', l'', l'''=0,2} \sum_{k, \kappa} (-1)^\kappa [\mathbf{b}^+_{l'} \times \mathbf{b}^-_{l''}]^{(k)}_\kappa [\mathbf{b}^+_{l'''} \times \mathbf{b}^-_{l'''}]^{(k)}_{-\kappa} \quad \text{i. e.} \quad (14.34) \\ \mathbf{C}_{u(6)} &= \sum_{k=0}^4 \sum_\kappa (-1)^\kappa [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}_\kappa [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}_{-\kappa} + \\ &\quad \sum_\kappa (-1)^\kappa ([\mathbf{d}^+ \times \mathbf{s}]^{(2)}_\kappa [\mathbf{s}^+ \times \mathbf{d}^-]^{(2)}_{-\kappa} + [\mathbf{s}^+ \times \mathbf{d}^-]^{(2)}_\kappa [\mathbf{d}^+ \times \mathbf{s}]^{(2)}_{-\kappa}) + \\ &\quad [\mathbf{s}^+, \mathbf{s}]^{(0)} [\mathbf{s}^+, \mathbf{s}]^{(0)}. \end{aligned} \quad (14.35)$$

With the aid of (14.33), (6.35), (6.36), (5.36) and little conversions one obtains

$$\begin{aligned} \mathbf{C}_{u(6)} &= n_d^2 + 4 n_d + n_d(1 + n_s) + n_s(n_d + 5) + n_s^2 = \\ &= (n_d + n_s)(n_d + n_s + 5) = N(N + 5). \end{aligned} \quad (14.36)$$

Because \mathbf{N} is the number operator for bosons, the operator $\mathbf{C}_{2, u(6)}$ has the eigenvalue $N(N + 5)$ and it meets the conditions (13.31) up to (13.33). According to (14.36) one can write $\mathbf{N}^2 = \mathbf{C}_{u(6)} - 5\mathbf{N}$. The operator \mathbf{N} is the linear Casimir operator of $u(6)$.

In table 14.2 the Casimir operators of $\mathbf{H}^{(1)}$ (14.31) are put together.

Table 14.2

Operators included in the Hamilton operator of the vibrational limit

Symbols for Eigenvalues these Casimir operators

$\mathbf{J}^2 + 1)$	$= -10\sqrt{3} \cdot [[\mathbf{d}^+ \times \mathbf{d}^-]^{(1)} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(1)}]^{(0)}$	$\mathbf{C}_{2,so(3)}$	$J(J + 1)$
\mathbf{T}^2	$= -2 \sum_{J=1,3} \sqrt{(2J+1)} \cdot [[\mathbf{d}^+ \times \mathbf{d}^-]^{(J)} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(J)}]^{(0)}$	$\mathbf{C}_{2,so(5)}$	$\tau(\tau + 3)$
$n_d(n_d + 4)$	$= \sum_{k=0}^4 (-1)^k \sqrt{(2k+1)} [[\mathbf{d}^+ \times \mathbf{d}^-]^{(k)} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}]^{(0)}$	$\mathbf{C}_{2,u(5)}$	$n_d(n_d + 4)$
n_d	$= \sqrt{5} [\mathbf{d}^+ \times \mathbf{d}^-]^{(0)}$	$\mathbf{C}_{1,u(5)}$	n_d
$\mathbf{N}(\mathbf{N} + 5)$	$= \sum_{k=0}^4 (-1)^k \sqrt{(2k+1)} [[\mathbf{d}^+ \times \mathbf{d}^-]^{(k)} \times [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}]^{(0)} + \sqrt{5} [[\mathbf{d}^+ \times \mathbf{s}]^{(2)} \times [\mathbf{s}^+ \times \mathbf{d}^-]^{(2)}]^{(0)} + \sqrt{5} [[\mathbf{s}^+ \times \mathbf{d}^-]^{(2)} \times [\mathbf{d}^+ \times \mathbf{s}]^{(2)}]^{(0)} + [[\mathbf{s}^+, \mathbf{s}]^{(0)} \times [\mathbf{s}^+, \mathbf{s}]^{(0)}]^{(0)}$	$\mathbf{C}_{2,u(6)}$	$N(N + 5)$
\mathbf{N}	$= [\mathbf{s}^+, \mathbf{s}]^{(0)} + \sqrt{5} [\mathbf{d}^+ \times \mathbf{d}^-]^{(0)}$	$\mathbf{C}_{1,u(6)}$	N

The operator $\mathbf{H}^{(1)}$ (14.31) can be written like this as well

$$\mathbf{H}^{(1)} =$$

$$\alpha_1 \mathbf{C}_{1,u(6)} + \alpha_2 \mathbf{C}_{2,u(6)} + \beta \mathbf{C}_{1,u(6)} \mathbf{C}_{1,u(5)} + \gamma_1 \mathbf{C}_{1,u(5)} + \gamma_2 \mathbf{C}_{2,u(5)} + \delta \mathbf{C}_{2,so(5)} + \varepsilon \mathbf{C}_{2,so(3)}$$

$$\text{with } \alpha_1 = \varepsilon_n - 5v_n, \alpha_2 = v_n, \quad \beta = v_{nd}, \quad \gamma_1 = \varepsilon'_d - 4v_d, \quad (14.37)$$

$$\gamma_2 = v_d, \quad \delta = v_t, \quad \varepsilon = v_j.$$

With the help of table 14.2 the eigenenergy can be represented this way

$$E^{(1)} = \quad (14.38)$$

$$\alpha_1 N + \alpha_2 N(N + 5) + \beta N n_d + \gamma_1 n_d + \gamma_2 n_d(n_d + 4) + \delta \tau(\tau + 3) + \varepsilon J(J + 1).$$

In section 10.3 eigenvalues of the Hamilton operator $\mathbf{H}^{(1)}$ have been calculated with (10.17) and compared with measured level energies.

Whenever it is possible to represent energy eigenvalues in terms of eigenvalues of Casimir operators there is talk of **dynamical symmetry**.

14.5 Casimir operators of the $su(3)$ - or rotational limit

According to (14.12) the operators

$$\begin{aligned} \mathbf{J}_\kappa &= [\mathbf{d}^+ \times \mathbf{d}^-]^{(1)}_\kappa \text{ and} \\ \mathbf{Q}_\mu &= [\mathbf{d}^+ \times \mathbf{s}]^{(2)}_\mu + [\mathbf{s}^+ \times \mathbf{d}^-]^{(2)}_\mu - (\sqrt{7}/2) [\mathbf{d}^+ \times \mathbf{d}^-]^{(2)}_\mu \end{aligned} \quad (14.39)$$

are basis elements of the $su(3)$ algebra in the rotational limit ((14.17) and table 14.1). In order to find its Casimir operator we look into the commutator of $\mathbf{Q}^2 = \sum_\lambda (-1)^\lambda \mathbf{Q}_\lambda \mathbf{Q}_{-\lambda}$ (9.4) and \mathbf{Q}_μ and maintain

$$[\mathbf{Q}^2, \mathbf{Q}_\mu] = 3\sqrt{(3/32)} \sum_{\mu' \mu''} (1 \mu' 2 \mu'' | 2 \mu) (\mathbf{J}_{\mu'} \mathbf{Q}_{\mu''} + \mathbf{Q}_{\mu''} \mathbf{J}_{\mu'}). \quad (14.40)$$

Owing to (14.16) and (14.14) the relations

$$\begin{aligned} [\mathbf{Q}_\kappa, \mathbf{Q}_\mu] &= 3\sqrt{(5/32)} (2 \kappa 2 \mu | 1, \kappa + \mu) \mathbf{J}_{\kappa + \mu} \\ [\mathbf{Q}_\kappa, \mathbf{J}_\mu] &= -\sqrt{6} (2 \kappa 1 \mu | 2, \kappa + \mu) \mathbf{Q}_{\kappa + \mu} \end{aligned} \quad (14.41)$$

are valid. We insert them in succession in $[\mathbf{Q}^2, \mathbf{Q}_\mu]$ and obtain the equation (14.40) making use of (A1.4) and (A1.6). Analogously one obtains

$$[\mathbf{J}^2, \mathbf{Q}_\mu] = -\sqrt{6} \sum_{\mu' \mu''} (1 \mu' 2 \mu'' | 2 \mu) (\mathbf{J}_{\mu'} \mathbf{Q}_{\mu''} + \mathbf{Q}_{\mu''} \mathbf{J}_{\mu'}). \quad (14.42)$$

One multiplies this relation by $3/8$, adds it to (14.40) and obtains

$$[\mathbf{Q}^2 + (3/8)\mathbf{J}^2, \mathbf{Q}_\mu] = 0. \quad (14.43)$$

$$\text{The operator } (1/2)\mathbf{C}_{2, su(3)} = \mathbf{Q}^2 + (3/8)\mathbf{J}^2 \quad (14.44)$$

commutes not only with \mathbf{Q}_μ but also with \mathbf{J}_μ , because on the one hand \mathbf{J}^2 commutes anyway with \mathbf{J}_μ (13.28) and on the other hand one can show that \mathbf{Q}^2 commutes with \mathbf{J}_μ . That is, with the aid of (14.41), (A1.4) and (A1.6) one obtains

$$\begin{aligned} [\mathbf{Q}^2, \mathbf{J}_\mu] &= -\sqrt{6} \sum_\lambda (-1)^\lambda (2, -\lambda, 1 \mu | 2, -\lambda + \mu) \mathbf{Q}_\lambda \mathbf{Q}_{-\lambda + \mu} - \\ &\quad \sqrt{6} \sum_\lambda (-1)^\lambda (2 \lambda 1 \mu | 2, \lambda + \mu) \mathbf{Q}_{\lambda + \mu} \mathbf{Q}_{-\lambda} = \\ &\quad -\sqrt{6} \sqrt{(5/3)} (\sum_{\lambda' \lambda''} (2, -\lambda', 2 - \lambda'' | 1, -\mu) \mathbf{Q}_{\lambda'} \mathbf{Q}_{-\lambda''} + \\ &\quad \sum_{\lambda' \lambda''} (2, -\lambda'', 2 - \lambda' | 1, -\mu) \mathbf{Q}_{\lambda'} \mathbf{Q}_{-\lambda''}) = 0. \end{aligned} \quad (14.45)$$

Thus the operator (14.44) commutes with all elements (14.39) of the $su(3)$ algebra and represents its Casimir operator. Its structure meets the condition (13.32). The factor $1/2$ in (14.44) gives the Casimir operator the usual form. According to (13.35) the operator $\mathbf{C}_{2, su(3)}$ has the following eigenvalues

$$\lambda^2 + \mu^2 + \lambda\mu + 3(\lambda + \mu), \quad \lambda, \mu \text{ integers}$$

when it acts on the corresponding eigenstates (provided that they are in a so-called irreducible representation) which are themselves characterised by the parameters λ and μ . Naturally these states are also eigenfunctions of \mathbf{J}^2 with the eigenvalue $J(J+1)$. A detailed treatment of these eigenstates reveals that there exist restrictions for λ and μ . They are positive integers and obey the following rules

$$\mu = 0, 2, 4, \dots$$

$$\lambda = 2N - 6l - 2\mu \text{ with } l = 0, 1, \dots \text{ and } N = \text{number of bosons.} \quad (14.46)$$

Therefore, the doublets (λ, μ) contain the following values: for $N = 1$ they amount to $(2, 0)$; for $N = 2$ we have $(4, 0)$, $(0, 2)$; for $N = 3$ consequently $(6, 0)$, $(2, 2)$, $(0, 0)$ and so forth. Further doublets can be taken from figure 14.1.




Figure 14.1. Graph of the doublets (λ, μ) according to the equations (14.46). They represent descending straight lines. At their highest point the corresponding doublets (N, l) are given.

A given doublet (λ, μ) admits only selected values for J . They depend on a ordering number K , for which the following values are permitted

$$K = 0, 2, 4, \dots, \min(\lambda, \mu). \quad (14.47)$$

For $K = 0$ the values $J = 0, 2, 4, \dots, \max(\lambda, \mu)$ are allowed

$$\text{and for } K > 0 \quad J = K, K + 1, K + 2, \dots, K + \max(\lambda, \mu). \quad (14.48)$$

We look at the simplest values. For $\mu = l = 0$ we have $(\lambda, \mu) = (2N, 0)$ and the series of the ground state shows the values $J = 0, 2, 4, \dots, 2N$. The case $\mu = 2, l = 0$ is the next higher irreducible representation with $(\lambda, \mu) = (2N - 4, 2)$ where the values $K = 0$ with $J = 0, 2, 4, \dots, 2N - 2$ and $K = 2$ with $J = 2, 3, 4, \dots, 2N - 2$ arise. It happens that J -values which belong to the same doublet (λ, μ) appear many times. We will show that such states have the same energy i. e. they are degenerated. The parameter K is needed for describing the eigenstates.

The $su(3)$ - or rotational limit is realised by deleting the following coefficients in the Hamiltonian (9.39)

$$\varepsilon_{d'} = v_{nd} = v_d = v_r = v_t = 0. \quad (14.49)$$

The remaining Hamiltonian consists of the Casimir operators of the second chain of algebras ((14.17) and table 14.1) as follows

$$H^{(II)} = \varepsilon_n N + v_n N^2 + v_j J^2 + v_q Q^2. \quad (14.50)$$

Using (14.44) we write the last two terms of (14.50) like this

$$v_j J^2 + v_q Q^2 = \quad (14.51)$$


$$v_j J^2 - (3/8)v_q J^2 + v_q Q^2 + (3/8)v_q J^2 = (v_j - (3/8)v_q)J^2 + (1/2)v_q C_{su(3)}.$$

The eigenstates of the Casimir operator of the $su(3)$ chain are eigenstates of the Hamilton operator $H^{(II)}$ as well. The operators N and $N(N + 5) = C_{u(6)}$ have already been discussed (14.36). Thus the eigenvalues of $H^{(II)}$ read

$$E^{(II)} = \quad (14.52)$$

$$\varepsilon_n N + v_n N^2 + (v_j - (3/8)v_q)J(J + 1) + (1/2)v_q(\lambda^2 + \mu^2 + \lambda\mu + 3(\lambda + \mu)).$$

They don't depend directly on the ordering number K (14.47). Figure 14.2 shows a typical energy spectrum of a nucleus of the $su(3)$ limit, which is calculated with the help of (14.52). Because of $v_q < 0$ the level energies of every series with given (λ, μ) ascend according to the term $(v_j - (3/8)v_q)J(J + 1)$. This behaviour is typical for the rotator and that's why the $su(3)$ limit is named rotational limit as well.



Extensive investigations (Arima and Iachello, 1978, 1987) reveal that the nuclei which behave like the $su(3)$ model have proton- and neutron numbers deviating clearly from the magic numbers 82 and 126 (figure 14.3).




Figure 14.3. Card of even-even nuclei. Z = number of protons, N = number of neutrons. The grey circles denote nuclei of the $su(3)$ limit and the black ones are representations of the $so(6)$ limit (Iachello and Arima, 1987, p.42).

Figure 14.4 shows that the rare earth ^{170}Er corresponds well to the energy scheme of the $su(3)$ limit.




Figure 14.4. Comparison between the measured and the calculated $su(3)$ -spectrum of ^{170}Er (Arima and Iachello, 1978, p. 215). The parameters $N = 17$, $v_j = 9.19$ keV and $v_q = -8.8$ keV have been used with formula (14.52). Broken lines represent the calculated energies.

14.6 Casimir operators of the $so(6)$ - or γ -instable limit

The chain of Lie algebras (14.21) defining the $so(6)$ limit comprises the subalgebras $so(5) \supset so(3)$. Their Casimir operators have been dealt with in

the elements (14.18), has the following Casimir operator (Talmi, 1993, p.778)

$$\mathbf{C}_{so(6)} = 2 \sum_{k=1,3} \sum_{\kappa} (-1)^{\kappa} [\mathbf{d}^+ \times \mathbf{d}]_{\kappa}^{(k)} [\mathbf{d}^+ \times \mathbf{d}]_{-\kappa}^{(k)} + \sum_{\lambda} (-1)^{\lambda} \mathbf{S}_{\lambda} \mathbf{S}_{-\lambda} \text{ with} \\ \mathbf{S}_{\lambda} = [\mathbf{d}^+ \times \mathbf{s}]_{\lambda}^{(2)} + [\mathbf{s}^+ \times \mathbf{d}]_{\lambda}^{(2)}. \quad (14.53)$$

Its form meets the condition (13.32). We have to check if $\mathbf{C}_{so(6)}$, which is identical with \mathbf{R}^2 (14.7), commutes with the operators (14.18). Because the first part of $\mathbf{C}_{so(6)}$ is identical with \mathbf{T}^2 according to (14.5), this part commutes with the affiliated elements $[\mathbf{d}^+ \times \mathbf{d}]_{\kappa}^{(k)}$ ($k = 1, 3$) (see 14.32). But these operators commute also with the second part of $\mathbf{C}_{so(6)}$, i. e. with $\sum_{\lambda} (-1)^{\lambda} \mathbf{S}_{\lambda} \mathbf{S}_{-\lambda}$, which can be shown with the aid of (14.20) and roughly with the procedure employed for (14.45). With the same methods one proves that \mathbf{S}_{λ} commutes with $\mathbf{C}_{so(6)}$ and in doing so, one finds that the following expressions cancel out

$$[\mathbf{T}^2, \mathbf{S}_{\kappa}] + \sum_{\lambda} (-1)^{\lambda} [\mathbf{S}_{\lambda} \mathbf{S}_{-\lambda}, \mathbf{S}_{\kappa}] = 0, \text{ i. e. } [\mathbf{C}_{so(6)}, \mathbf{S}_{\kappa}] = 0. \quad (14.54)$$

Consequently the operator $\mathbf{C}_{so(6)}$ meets the condition (13.31). It is written in the standard form and therefore its eigenvalue reads according to (13.34) (rank $N = 6$)

$$\rho(\rho + 4). \quad (14.55)$$

A detailed investigation of the eigenstates of the $so(6)$ algebra reveals that the admissible values for ρ are

$$\rho = N, N - 2, N - 4, \dots, 1 \text{ or } 0 \quad (N : \text{number of bosons}). \quad (14.56)$$

Moreover, for the parameter τ of the Casimir operator \mathbf{T}^2 the following values are admitted

$$\tau = 0, 1, \dots, \rho. \quad (14.57)$$

According to (10.5) J can amount to

$$J = \lambda, \lambda + 1, \dots, 2\lambda - 3, 2\lambda - 2, 2\lambda \quad (14.58)$$

with $\lambda = \tau - 3n_{\Delta}$ and $n_{\Delta} = 0, 1, \dots$ (see chapter 3). Table 14.2 gives permissible values for J .

Table 14.2. Values for J of the $so(6)$ limit depending on n_Δ and τ .

τ	n_Δ	J
0	0	0
1	0	2
2	0	2, 4
3	1	0
	0	3, 4, 6
4	1	2
	0	4, 5, 6, 8
5	1	2, 4
	0	5, 6, 7, 8, 10
6	2	0
	1	3, 4, 6
	0	6, 7, 8, 9, 10, 12

One treats the Casimir operator of the highest algebra, $u(6)$, in the same way as in the previously discussed special cases of the IBM1.

Now we turn to the Hamilton operator. For the present special case, the $so(6)$ limit, one deletes the following coefficients of the equation (9.39)

$$\varepsilon_d' = v_{nd} = v_d = v_q = 0. \quad (14.59)$$

Choosing $v_q = 0$ results in $v_2 = 0$ (9.38), which means (see 7.17) that the Hamilton operator is not able any longer to interchange dd - with sd -states i. e. that there is no interaction between these states. The remaining Hamilton operator, which defines the $so(6)$ - or γ -instable limit, reads

$$H^{(III)} = \varepsilon_n N + v_n N^2 + v_r R^2 + v_t T^2 + v_j J^2. \quad (14.60)$$

It represents a linear combination of the Casimir operators of the chain (14.21). Again the functions of the highest algebra are functions of the lower ones and therefore the energy eigenvalue is given according to (10.3), (10.13) and (14.55) by the following expression

$$E^{(III)} = \varepsilon_n N + v_n N^2 + v_r \rho(\rho + 4) + v_t \tau(\tau + 3) + v_j J(J + 1). \quad (14.61)$$

Figure 14.5 shows a typical energy spectrum of a nucleus of the $so(6)$ limit according to equation (14.61). The $2^+ - 4^+$ -doublet of the third and fourth level (with $\tau = 2$) is characteristic for this special case and appears also in the so-called γ -instable nuclei, for which reason this model is named γ -instable limit as well.

In figure 14.3 the regions with nuclei of the $so(6)$ type are marked with black dots. They join always regions with $su(5)$ nuclei (see figure 10.7) which corresponds to the noticeable similarity of both Hamiltonians.




Figure 14.5. Typical spectrum of the $so(6)$ limit. It is calculated by means of equation (14.61) and the values $N = 6$, $v_r = -30$ keV, $v_t = 40$ keV and $v_j = 5, 1$ keV. The parameters ρ and τ follow the rules (14.56) and (14.57). The values of n_Δ and J are given by (14.58). All levels have positive parity. Level groups with the same τ are joined together and labelled with the value of τ . The J -values can be taken from the columns on top of the figure.

Figures 14.6 and 14.7 compare measured with calculated levels of ^{196}Pt and ^{132}Ba .




Figure 14.6. Comparison between the measured and the calculated $so(6)$ -spectrum of ^{196}Pt (Iachello and Arima, 1987, p. 43). The parameters $N = 6$, $v_r = -42.75$ keV, $v_t = 50$ keV and $v_j = 10$ keV have been used with formula (14.61). The spins are given and the parity is positive. Broken lines represent calculated energies.




Figure 14.7. Comparison between the measured and the calculated $so(6)$ -spectrum of ^{132}Ba (Arima and Iachello, 1979, p. 486). The expression (14.61) and the parameters $\nu_t = 90$ keV and $\nu_j = 10$ keV have been used. Because only the states with $\rho = N = 6$ have been identified unambiguously, ν_r does not appear. Broken lines represent the calculated energies.

The $so(6)$ limit offers the chance to test the program PHINT (appendix 7). One chooses first the parameters in equation (14.60) and puts $\varepsilon_n = \nu_n = 0$, which is admissible when the spectrum of a single nucleus is looked at. By inserting in equation (9.38) and by transforming one obtains

$$\begin{aligned}
\varepsilon_d - \varepsilon_s &= 2 (3v_j + 2v_t), \\
c_0 &= 4 (- 3v_j - 2v_t - 2v_r), \\
c_2 &= 2 (-3v_j + v_t + v_r), \\
c_4 &= 2 (4v_j + v_t + v_r), \\
v_2 &= 0, \\
v_0 &= 2 \sqrt{5} v_r, \\
u_2 &= 2 \sqrt{5} v_r, \\
u_0 &= 0.
\end{aligned} \tag{14.62}$$

By means of (A7.1) one finds the input parameters for the program. Calculations using the parameters of figure 14.5, which are performed both according to (14.61) and with the help of PHINT and (14.62), agree perfectly.

According to Iachello and Arima (1987, p. 25) it can be shown that there exist only three chains of algebras, (14.11), (14.17) and (14.21), in the interacting boson model 1. Therefore there are only three types of dynamical symmetries.

15 The proton-neutron interacting boson model IBM2

In this interacting boson model, protons and neutrons are treated separately. In the sections 1 up to 3, we will show how the Hamilton operator of this model can be simplified and how the number of parameters is reduced. Sections 4 and 5 demonstrate the eigenenergies and transition probabilities and compare them with experimental data. There will be talk of group theoretical structures of the IBM2 in section 6.

15.1 The complete Hamiltonian of the IBM2

In the IBM1 the boson number of the protons is added to the boson number of neutrons (chapter 2) and later on, no distinction is made between these types of bosons. In chapter 7 the interaction terms don't contain Coulomb forces. Moreover, speaking in terms of the shell model, the fact is known that in intermediate and heavy nuclei protons and neutrons occupy different shells. This results in an interaction between proton- and neutron pairs, which differs from the interaction between identical pairs. This fact contrasts to the well-known charge independence of the short-range nuclear forces. Likewise, this effect is not taken into account in the IBM1.

Because of these simplifications, the parameters of the IBM1 fitted on experimental data change from nucleus to nucleus in a discontinuous and hardly interpretable way. On the other hand the interacting model 2, which distinguishes between proton- (π) and neutron (ν) bosons, yields results which can be interpreted by means of the SD-version of the shell model. The IBM2 has special practical significance because its parameters depend in a smoothed curve on π - and ν -boson numbers and it makes possible to calculate unknown nuclear spectra (Arima and Iachello, 1984, p. 177). The interest in the algebraic structure of the IBM2 has even grown since states with special magnetic dipole properties (Bohle, 1984) are known which can be explained with group theoretical methods.

The numbers of π -bosons N_π and ν -bosons N_ν are fixed equally as in chapter 2. There is no boson composed of a proton and a neutron. There exist 12 creation operators for bosons

$$\begin{aligned} b_{\pi, jm}^+ &= s_{\pi}^+, d_{\pi, m}^+ (m = -2, -1, \dots, 2) \text{ and} \\ b_{\nu, jm}^+ &= s_{\nu}^+, d_{\nu, m}^+ (m = -2, -1, \dots, 2). \end{aligned} \quad (15.1)$$

Analogously to (6.19), we have the same number of tensor operators for boson annihilation like this

$$b_{\pi, jm}^- = s_{\pi}^-, d_{\pi, m}^- , b_{\nu, jm}^- = s_{\nu}^-, d_{\nu, m}^- (m = -2, -1, \dots, 2). \quad (15.2)$$

In analogy with (5.33) and (5.35), every d - or s -operator for protons commutes with every d - or s -operator for neutrons. We demand that both N_π and N_ν are good quantum numbers i. e. the Hamilton operator must meet the condition

$$[H, N_\pi] = [H, N_\nu] = 0 \quad (15.3)$$

with the operators $N_\pi = \sqrt{5}[\mathbf{d}^+_\pi \times \mathbf{d}^\sim_\pi]^{(0)} + \mathbf{s}^+_\pi \mathbf{s}_\pi$ (6.37) and the corresponding N_ν . Generalising the equations (7.3), (7.5) and (7.6), we write the Hamilton operator as follows

$$H = H_\pi + H_\nu + V_{\pi\nu}. \quad (15.4)$$

Both H_π and H_ν have the form of the IBM1-Hamiltonian (7.3, 7.6, and 7.17) but there is nothing than π -operators in H_π and ν -operators in H_ν . We write the mixing potential operator $V_{\pi\nu}$ employing the parameter

$$\rho = \pi, \nu \quad (15.5)$$

analogously to (7.6) like this

$$V_{\pi\nu} = (1/2) \sum_{j_f j_g j_p j_q} \sum_{m_f m_g m_p m_q} \sum_{\rho_f \neq \rho_g} \sum_{\rho_p \neq \rho_q} \quad (15.6)$$

$$\langle \rho_f j_f m_f, \rho_g j_g m_g | \mathbf{W} | \rho_p j_p m_p, \rho_q j_q m_q \rangle \cdot \mathbf{b}^+_{\rho_f j_f m_f} \mathbf{b}^+_{\rho_g j_g m_g} \mathbf{b}_{\rho_p j_p m_p} \mathbf{b}_{\rho_q j_q m_q}$$

where $j = 0$ denotes a s -operator and $j = 2$ a d -operator. Both the creation and the annihilation operators in (15.6) comprise a π - and a ν -operator ($\rho_f \neq \rho_g, \rho_p \neq \rho_q$), otherwise the matrix element vanishes. Analogously to (7.6) $V_{\pi\nu}$ can be written by means of vector coupling as

$$V_{\pi\nu} = (1/2) \sum_{j_f j_g j_p j_q} \sum_{J=0}^4 \sum_{\rho_f \neq \rho_g} \sum_{\rho_p \neq \rho_q} \langle \rho_f j_f, \rho_g j_g, J | \mathbf{W} | \rho_p j_p, \rho_q j_q, J \rangle \cdot \sqrt{(2J+1)} [[\mathbf{b}^+_{\rho_f j_f} \times \mathbf{b}^+_{\rho_g j_g}]^{(J)} \times [\mathbf{b}_{\rho_p j_p} \times \mathbf{b}_{\rho_q j_q}]^{(J)}]^{(0)}. \quad (15.7)$$

Because the operators $\mathbf{b}^+_{\pi j m}$ and $\mathbf{b}^+_{\nu j m}$ create distinguishable bosons, the common state does not show symmetry according to (A2.5) and for $[\mathbf{d}^+_\pi \times \mathbf{d}^+_\nu]^{(J)}$ the statement (6.6) is not valid any more. On the contrary, for J all integers from 0 up to 4 are possible.

In the case of $j_p = j_q = 2$ for the two-boson state in the matrix element in (15.7) one obtains the following expression

$$| \pi 2, \nu 2, J M \rangle = \sum_{m_\pi m_\nu} (2 m_\pi 2 m_\nu | J M) | \pi 2 m_\pi, \nu 2 m_\nu \rangle = \quad (15.8)$$

$$(-1)^J \sum_{m_\pi m_\nu} (2 m_\nu 2 m_\pi | J M) | \nu 2 m_\nu, \pi 2 m_\pi \rangle = (-1)^J | \nu 2, \pi 2, J M \rangle.$$

Therefore, there are 12 independent matrix elements in (15.7) as follows

$$\begin{aligned}
W_5 &= \langle \pi 0, \nu 0, 0 | \mathbf{W} | \pi 0, \nu 0, 0 \rangle \\
W_6 &= \langle \pi 0, \nu 2, 2 | \mathbf{W} | \pi 0, \nu 2, 2 \rangle \\
W_7 &= \langle \pi 0, \nu 2, 2 | \mathbf{W} | \pi 2, \nu 0, 2 \rangle = \langle \pi 2, \nu 0, 2 | \mathbf{W} | \pi 0, \nu 2, 2 \rangle \\
W_8 &= \langle \pi 2, \nu 0, 2 | \mathbf{W} | \pi 2, \nu 0, 2 \rangle \\
W_9 &= \langle \pi 0, \nu 0, 0 | \mathbf{W} | \pi 2, \nu 2, 0 \rangle = \langle \pi 2, \nu 2, 0 | \mathbf{W} | \pi 0, \nu 0, 0 \rangle \\
W_{10} &= \langle \pi 0, \nu 2, 2 | \mathbf{W} | \pi 2, \nu 2, 2 \rangle = \langle \pi 2, \nu 2, 2 | \mathbf{W} | \pi 0, \nu 2, 2 \rangle \\
W_{11} &= \langle \pi 2, \nu 0, 2 | \mathbf{W} | \pi 2, \nu 2, 2 \rangle = \langle \pi 2, \nu 2, 2 | \mathbf{W} | \pi 2, \nu 0, 2 \rangle.
\end{aligned} \tag{15.9}$$

These matrix elements don't alter if one interchanges the π - and the ν -state on one side. The residual elements read

$$W_J = \langle \pi 2, \nu 2, J | \mathbf{W} | \pi 2, \nu 2, J \rangle \quad (J = 0, 1, 2, 3, 4). \tag{15.10}$$

Here interchanging the π - and the ν -state results in a factor $(-1)^J$ (see 15.8) i. e.

$$\begin{aligned}
\langle \pi 2, \nu 2, J | \mathbf{W} | \pi 2, \nu 2, J \rangle &= \langle \pi 2, \nu 2, J | \mathbf{W} | \nu 2, \pi 2, J \rangle (-1)^J = \\
\langle \nu 2, \pi 2, J | \mathbf{W} | \pi 2, \nu 2, J \rangle (-1)^J &= \langle \nu 2, \pi 2, J | \mathbf{W} | \nu 2, \pi 2, J \rangle.
\end{aligned} \tag{15.11}$$

The partial Hamilton operators H_π and H_ν contain, according to (7.17) and (7.33) respectively, 9 or 6 free parameters each. The considerations in chapter 7 that facilitated the reduction of this number can be applied to the 12 parameters in (15.9) and (15.11) and they can be reduced to 9 (Iachello and Arima, 1987, p. 138). Therefore the Hamilton operator (15.4) comprises 30 or 21 parameters respectively, which render this theory cumbersome and it is practically impossible to fit the parameters unambiguously on experimental data.

15.2 Basis states and the angular momentum operator of the IBM2

Before we deal with the simplification of the Hamilton operator, we turn to the eigenstates. For practical reasons, the basis states of the IBM2 are chosen as vectorial couplings of symmetrical, spherical IBM1-states (6.18) of π - and ν -bosons this way

$$| [(n_{s\pi}, n_{d\pi}, \tau_\pi, n_{\Delta\pi}, J_\pi) \times (n_{s\nu}, n_{d\nu}, \tau_\nu, n_{\Delta\nu}, J_\nu)]^{(J)}_M \rangle. \tag{15.12}$$

For example the complete set of states with $N_\pi = N_\nu = 1$ reads as follows

$$\begin{aligned}
&| [s_\pi \times s_\nu]^{(0)}_0 \rangle, | [s_\pi \times d_\nu]^{(2)}_\mu \rangle, | [d_\pi \times s_\nu]^{(2)}_{\mu'} \rangle, | [d_\pi \times d_\nu]^{(J)}_{\mu''} \rangle \text{ with} \\
&(J = 0, 1, \dots, 4).
\end{aligned} \tag{15.13}$$

The basis function of the IBM2 (15.12) demands a corresponding angular momentum operator according to (A1.21) and (A1.23). Analogously to (8.9) its components read

$$\mathbf{J}_\mu = \mathbf{J}_{\pi\mu} + \mathbf{J}_{\nu\mu} = \sqrt{10} ([\mathbf{d}^+_\pi \times \mathbf{d}^-_{\pi}]^{(1)}_\mu + [\mathbf{d}^+_\nu \times \mathbf{d}^-_{\nu}]^{(1)}_\mu) \quad (15.14)$$

and meet the commutation rules (8.7). According to (A1.22) and (A1.26) the states (15.12) are eigenfunctions of the operators \mathbf{J}_0 and $\mathbf{J}^2 = \mathbf{J}_0^2 - \mathbf{J}_0 - 2 \mathbf{J}_1 \mathbf{J}_1$ which are made up by the operators (15.14).

15.3 A reduced Hamiltonian of the IBM2

In section 15.1 we have seen that the IBM2-Hamiltonian comprises a large number of parameters. In practice it is reduced drastically in the course of which one is based on arguments of microscopic theories - mainly on the SD-version of the shell model - and on experiences made by fitting on measured data. A simplification of the Hamilton operator is also necessary because the sum \mathbf{H}_π and \mathbf{H}_ν generates a large number of energy levels. Because the IBM1 reproduces the spectrum of measured energies relatively well, the IBM2 may not generate essentially more levels.

A first simplification of the Hamiltonian consists in reducing \mathbf{H}_π and \mathbf{H}_ν to a special case (chapter 14). For that purpose, the vibrational limit (chapter 10) imposes itself because it is able to interpret most nuclei (compare figures 10.7 and 14.3). Thus in both IBM-Hamiltonians on drops \mathbf{R}^2 and \mathbf{Q}^2 (9.39). In practice in equation (7.17) not only v_0 and v_2 are deleted but also u_0 , u_2 and ε_s . Consequently the following expression is left over

$$\mathbf{H}_\rho = \varepsilon_\rho n_{d\rho} + \frac{1}{2} \sum_{J=0,2,4} c_{J\rho} \sqrt{(2J+1)} [[\mathbf{d}^+_\rho \times \mathbf{d}^+_\rho]^{(J)} \times [\mathbf{d}^-_\rho \times \mathbf{d}^-_\rho]^{(J)}]^{(0)} \\ \text{with } \rho = \pi, \nu. \quad (15.15)$$

The third term $\mathbf{V}_{\pi\nu}$ in the Hamilton operator (15.4) is simplified as well. Generally, one employs the following form (Iachello and Arima, 1987, p. 140)

$$\mathbf{V}_{\pi\nu} = \kappa \sqrt{5} [\mathbf{Q}_\pi(\chi_\pi) \times \mathbf{Q}_\nu(\chi_\nu)]^{(0)} + \lambda \mathbf{M}_{\pi\nu}. \quad (15.16)$$

The quadrupole operators $\mathbf{Q}_\rho(\chi_\rho)_\mu$ ($\rho = \pi, \nu$) correspond to the operators \mathbf{Q}_μ in (12.19) that is

$$\mathbf{Q}_\rho(\chi_\rho)_\mu = [\mathbf{d}^+_\rho \times \mathbf{s}_\rho]^{(2)}_\mu + [\mathbf{s}^+_\rho \times \mathbf{d}^-_\rho]^{(2)}_\mu + \chi_\rho [\mathbf{d}^+_\rho \times \mathbf{d}^-_\rho]^{(2)}_\mu. \quad (15.17)$$

For the quadrupole-quadrupole part of $\mathbf{V}_{\pi\nu}$ sometimes the symbol

$$\sqrt{5} [\mathbf{Q}_\pi(\chi_\pi) \times \mathbf{Q}_\nu(\chi_\nu)]^{(0)} \equiv \mathbf{Q}_\pi(\chi_\pi) \bullet \mathbf{Q}_\nu(\chi_\nu). \quad (15.18)$$

is employed. We write it explicitly

$$\begin{aligned}
& \sqrt{5} [\mathbf{Q}_\pi(\chi_\pi) \times \mathbf{Q}_\nu(\chi_\nu)]^{(0)} = \\
& \sqrt{5} \{ ([\mathbf{d}^+_\pi \times \mathbf{s}_\pi]^{(2)} \times [\mathbf{d}^+_\nu \times \mathbf{s}_\nu]^{(2)})^{(0)} + [[\mathbf{s}^+_\pi \times \mathbf{d}^+_\pi]^{(2)} \times [\mathbf{s}^+_\nu \times \mathbf{d}^+_\nu]^{(2)}]^{(0)} + \\
& ([[\mathbf{d}^+_\pi \times \mathbf{s}_\pi]^{(2)} \times [\mathbf{s}^+_\nu \times \mathbf{d}^+_\nu]^{(2)})^{(0)} + [[\mathbf{s}^+_\pi \times \mathbf{d}^+_\pi]^{(2)} \times [\mathbf{d}^+_\nu \times \mathbf{s}_\nu]^{(2)}]^{(0)} + \\
& \chi_\nu ([[\mathbf{d}^+_\pi \times \mathbf{s}_\pi]^{(2)} + [\mathbf{s}^+_\pi \times \mathbf{d}^+_\pi]^{(2)}) \times [\mathbf{d}^+_\nu \times \mathbf{d}^+_\nu]^{(2)} \}^{(0)} + \\
& \chi_\pi [[\mathbf{d}^+_\pi \times \mathbf{d}^+_\pi]^{(2)} \times ([\mathbf{d}^+_\nu \times \mathbf{s}_\nu]^{(2)} + [\mathbf{s}^+_\nu \times \mathbf{d}^+_\nu]^{(2)})]^{(0)} + \\
& \chi_\nu \chi_\pi [[\mathbf{d}^+_\pi \times \mathbf{d}^+_\pi]^{(2)} \times [\mathbf{d}^+_\nu \times \mathbf{d}^+_\nu]^{(2)}]^{(0)} \}. \tag{15.19}
\end{aligned}$$

The last term in equation (15.16) is named Majorana term and it reads

$$\begin{aligned}
\mathbf{M}_{\pi\nu} = & \sqrt{5} \{ ([[\mathbf{d}^+_\pi \times \mathbf{s}^+_\nu]^{(2)} \times [\mathbf{d}^+_\pi \times \mathbf{s}_\nu]^{(2)}]^{(0)} + [[\mathbf{s}^+_\pi \times \mathbf{d}^+_\nu]^{(2)} \times [\mathbf{s}_\pi \times \mathbf{d}^+_\nu]^{(2)}]^{(0)} - \\
& ([[\mathbf{d}^+_\pi \times \mathbf{s}^+_\nu]^{(2)} \times [\mathbf{s}_\pi \times \mathbf{d}^+_\nu]^{(2)}]^{(0)} + [[\mathbf{s}^+_\pi \times \mathbf{d}^+_\nu]^{(2)} \times [\mathbf{d}^+_\pi \times \mathbf{s}_\nu]^{(2)}]^{(0)}) \} + \\
& 2 \sum_{J=1,3} \sqrt{(2J+1)} [[\mathbf{d}^+_\pi \times \mathbf{d}^+_\nu]^{(J)} \times [\mathbf{d}^+_\pi \times \mathbf{d}^+_\nu]^{(J)}]^{(0)}. \tag{15.20}
\end{aligned}$$

Sometimes a further parameter is attached to the last expression. The terms in the expressions (15.19) and (15.20) appear individually in the original expression for $\mathbf{V}_{\pi\nu}$ (15.7), which can be pursued by means of table 15.1.

Table 15.1

Correspondence of terms in (15.7), (15.9) and (15.10) to the terms in (15.19) and (15.20).

Original terms of $\mathbf{V}_{\pi\nu}$ in (15.7), (15.9) and (15.10)	corresponding terms in (15.19) and (15.20), the numbers denote the lines.
Term with W_9 (see 15.9)	1. Term in $\sqrt{5} [\mathbf{Q}_\pi(\chi_\pi) \times \mathbf{Q}_\nu(\chi_\nu)]^{(0)}$, (15.19)
" " W_7	2. " " " ,
" " W_{10}	3. " " " ,
" " W_{11}	4. " " " ,
Terms with $W_0 \dots W_4$ (see 15.10)	5. " " " ,
" " W_6 and W_8	1. Term in $\mathbf{M}_{\pi\nu}$ (15.20),
Term with W_7	2. " " " ,
Terms with W_1 and W_3	3. " " " .

This comparison shows that in fact only the coefficients of the terms in $\mathbf{V}_{\pi\nu}$ have been altered. In doing so, individual terms had to be recoupled by means of (A4.1). The consistent correspondence reveals that expression (15.16) is not an ad hoc construction but the result of a reduction and remodelling of the operator (15.7).

The quadrupole-quadrupole operator has to replace to a certain degree the operators \mathbf{Q}_π^2 and \mathbf{Q}_ν^2 , which have been omitted in \mathbf{H}_π and \mathbf{H}_ν (15.15). Furthermore, one knows from the shell model that the strongly attractive interaction between proton and neutron breaks the so-called generalised seniority (Talmi, 1993, p. 906) and is responsible for the deformation of the nucleus (Eisenberg and Greiner, 1987, p. 443). This π - ν -interaction corresponds to the sd -terms in $\mathbf{Q}_\rho(\chi_\rho)_\mu$ (15.17), which influence the s - d -division of bosons.

In order to illustrate this behaviour we show that the term $\mathbf{Q}_\pi(\chi_\pi) \bullet \mathbf{Q}_\nu(\chi_\nu)$ (15.18, 15.19) is able to make up non vanishing matrix elements resting on the states $\mathbf{s}_\pi^+ \mathbf{s}_\nu^+ | \rangle$ and $[\mathbf{d}_\pi^+ \times \mathbf{d}_\nu^+]^{(0)} | \rangle$. We pursue this by means of the first term in (15.19), $[[\mathbf{d}_\pi^+ \times \mathbf{s}_\nu]^{(2)} \times [\mathbf{d}_\nu^+ \times \mathbf{s}_\pi]^{(2)}]^{(0)}$, as follows

$$\begin{aligned} & \langle | [\mathbf{d}_\pi^- \times \mathbf{d}_\nu^-]^{(0)} [[\mathbf{d}_\pi^+ \times \mathbf{s}_\nu]^{(2)} \times [\mathbf{d}_\nu^+ \times \mathbf{s}_\pi]^{(2)}]^{(0)} \mathbf{s}_\pi^+ \mathbf{s}_\nu^+ | \rangle = \\ & \langle | [\mathbf{d}_\pi^- \times \mathbf{d}_\nu^-]^{(0)} [\mathbf{d}_\pi^+ \times \mathbf{d}_\nu^+]^{(0)} \mathbf{s}_\pi \mathbf{s}_\nu \mathbf{s}_\pi^+ \mathbf{s}_\nu^+ | \rangle = \\ & \langle | [\mathbf{d}_\pi \times \mathbf{d}_\nu]^{(0)} [\mathbf{d}_\pi^+ \times \mathbf{d}_\nu^+]^{(0)} | \rangle = 1. \end{aligned} \quad (15.21)$$

Thus, the quadrupole-quadrupole operator creates non-diagonal matrix elements, which cause s - d -mixing.

The Majorana term $\mathbf{M}_{\pi\nu}$ (15.20) influences states with a special symmetry which is forbidden in the IBM1. We show that by means of states with $N_\pi = N_\nu = 1$ (15.13), which can be remodelled this way

$$[[\mathbf{s}_\pi \times \mathbf{s}_\nu]^{(0)}], ([[\mathbf{s}_\pi \times \mathbf{d}_\nu]^{(2)}_\mu] + [[\mathbf{d}_\pi \times \mathbf{s}_\nu]^{(2)}_\mu])/\sqrt{2}, [[\mathbf{d}_\pi \times \mathbf{d}_\nu]^{(J)}_\mu], J \text{ even}, \quad (15.22)$$

$$([[\mathbf{s}_\pi \times \mathbf{d}_\nu]^{(2)}_\mu] - [[\mathbf{d}_\pi \times \mathbf{s}_\nu]^{(2)}_\mu])/\sqrt{2}, [[\mathbf{d}_\pi \times \mathbf{d}_\nu]^{(J)}_\mu], J \text{ odd}. \quad (15.23)$$

The states (15.22) are symmetric with regard to the indices π and ν and the states (15.23) are antisymmetric. If we replace the labels π and ν by boson numbers 1 and 2, equation (15.22) represents the symmetric two-boson states of the IBM1 but the states (15.23) exist neither in this model nor in reality. By means of the Majorana term we can lift these intruder states in a higher energy than all decisive levels and make them insignificant.

For example, we consider first the influence of $\mathbf{M}_{\pi\nu}$ on one partial state in (15.23)

$$\begin{aligned} & (1/\sqrt{5}) \mathbf{M}_{\pi\nu} | [\mathbf{s}_\pi \times \mathbf{d}_\nu]^{(2)}_\mu \rangle = \\ & [[\mathbf{s}_\pi^+ \times \mathbf{d}_\nu^+]^{(2)} \times [\mathbf{s}_\pi \times \mathbf{d}_\nu^-]^{(2)}]^{(0)} \mathbf{s}_\pi^+ \mathbf{d}_{\nu\mu}^+ | \rangle - \\ & [[\mathbf{d}_\pi^+ \times \mathbf{s}_\nu^+]^{(2)} \times [\mathbf{s}_\pi \times \mathbf{d}_\nu^-]^{(2)}]^{(0)} \mathbf{s}_\pi^+ \mathbf{d}_{\nu\mu}^+ | \rangle. \end{aligned} \quad (15.24)$$

All other states vanish because each time an annihilation operator acts on an empty state $| \rangle$. We remodel the expression (15.24)

$$\begin{aligned}
(1/\sqrt{5}) \mathbf{M}_{\pi\nu} | [\mathbf{s}_\pi \times \mathbf{d}_\nu]^{(2)}_\mu \rangle = \\
\sum_\lambda \mathbf{d}^+_{\nu\lambda} \mathbf{d}^{\sim}_{\nu, -\lambda} (2 \lambda 2, -\lambda | 0 0) \mathbf{d}^+_{\nu\mu} \mathbf{s}^+_\pi \mathbf{s}^+_\pi | \rangle - \\
\sum_\lambda \mathbf{d}^+_{\pi\lambda} \mathbf{d}^{\sim}_{\nu, -\lambda} (2 \lambda 2, -\lambda | 0 0) \mathbf{d}^+_{\nu\mu} \mathbf{s}^+_\nu \mathbf{s}^+_\pi | \rangle.
\end{aligned} \tag{15.25}$$

Because of $\mathbf{d}^{\sim}_{-\lambda} = \mathbf{d}_\lambda (-1)^{-\lambda}$ and $(2 \lambda 2, -\lambda | 0 0) = (-1)^\lambda / \sqrt{5}$ the relation

$$(1/\sqrt{5}) \mathbf{M}_{\pi\nu} | [\mathbf{s}_\pi \times \mathbf{d}_\nu]^{(2)}_\mu \rangle = (1/\sqrt{5}) (\mathbf{d}^+_{\nu\mu} \mathbf{s}^+_\pi - \mathbf{d}^+_{\pi\mu} \mathbf{s}^+_\nu) | \rangle \text{ holds.} \tag{15.26}$$

Analogously one gets

$$\mathbf{M}_{\pi\nu} | [\mathbf{d}_\pi \times \mathbf{s}_\nu]^{(2)}_\mu \rangle = (\mathbf{d}^+_{\pi\mu} \mathbf{s}^+_\nu - \mathbf{d}^+_{\nu\mu} \mathbf{s}^+_\pi) | \rangle.$$

Therefore, for corresponding states of (15.22) and (15.23) we obtain

$$\mathbf{M}_{\pi\nu} (|[\mathbf{s}_\pi \times \mathbf{d}_\nu]^{(2)}_\mu \rangle + |[\mathbf{d}_\pi \times \mathbf{s}_\nu]^{(2)}_\mu \rangle) / \sqrt{2} = 0, \tag{15.27}$$

$$\begin{aligned}
\mathbf{M}_{\pi\nu} (|[\mathbf{s}_\pi \times \mathbf{d}_\nu]^{(2)}_\mu \rangle - |[\mathbf{d}_\pi \times \mathbf{s}_\nu]^{(2)}_\mu \rangle) / \sqrt{2} = \\
\sqrt{2} (|[\mathbf{s}_\pi \times \mathbf{d}_\nu]^{(2)}_\mu \rangle - |[\mathbf{d}_\pi \times \mathbf{s}_\nu]^{(2)}_\mu \rangle).
\end{aligned} \tag{15.28}$$

Now we investigate the effect of $\mathbf{M}_{\pi\nu}$ on other states of (15.22) and (15.23) like this

$$\begin{aligned}
\sqrt{(2J+1)} [[\mathbf{d}^+_\pi \times \mathbf{d}^+_\nu]^{(J)} \times [\mathbf{d}^{\sim}_\pi \times \mathbf{d}^{\sim}_\nu]^{(J)}]^{(0)} |[\mathbf{d}_\pi \times \mathbf{d}_\nu]^{(L)}_{M_L} \rangle = \\
\sum_M (-1)^{J-M} [\mathbf{d}^+_\pi \times \mathbf{d}^+_\nu]^{(J)}_M [\mathbf{d}^{\sim}_\pi \times \mathbf{d}^{\sim}_\nu]^{(J)}_{-M} (1/\sqrt{2}) |[\mathbf{d}_\pi \times \mathbf{d}_\nu]^{(L)}_{M_L} \rangle = \\
(1/\sqrt{2}) \sum_M (-1)^{J-M} [\mathbf{d}^+_\pi \times \mathbf{d}^+_\nu]^{(J)}_M \cdot \\
\sum_{\lambda_\pi \lambda_\nu \mu_\pi \mu_\nu} (2 \lambda_\pi 2 \lambda_\nu | J, -M) (2 \mu_\pi 2 \mu_\nu | L M_L) (-1)^{\lambda_\pi + \lambda_\nu} \mathbf{d}^+_{\pi, -\lambda_\pi} \mathbf{d}^+_{\nu, -\lambda_\nu} \mathbf{d}^+_{\pi \mu_\pi} \mathbf{d}^+_{\nu \mu_\nu} | \rangle = \\
(1/\sqrt{2}) \sum_M (-1)^J [\mathbf{d}^+_\pi \times \mathbf{d}^+_\nu]^{(J)}_M \cdot \sum_{\mu_\pi \mu_\nu} (2, -\mu_\pi 2, -\mu_\nu | J, -M) (2 \mu_\pi 2 \mu_\nu | L M_L) | \rangle = \\
(1/\sqrt{2}) \sum_M [\mathbf{d}^+_\pi \times \mathbf{d}^+_\nu]^{(J)}_M \cdot \sum_{\mu_\pi \mu_\nu} (2 \mu_\pi 2 \mu_\nu | J, -M) (2 \mu_\pi 2 \mu_\nu | L M_L) | \rangle = \\
(1/\sqrt{2}) [\mathbf{d}^+_\pi \times \mathbf{d}^+_\nu]^{(L)}_{M_L} | \rangle \delta_{JL}.
\end{aligned} \tag{15.29}$$

Therefore, we obtain

$$\mathbf{M}_{\pi\nu} |[\mathbf{d}_\pi \times \mathbf{d}_\nu]^{(L)}_{M_L} \rangle = + \sqrt{2} [\mathbf{d}^+_\pi \times \mathbf{d}^+_\nu]^{(L)}_{M_L} | \rangle (\delta_{1L} + \delta_{3L}). \tag{15.30}$$

For the first expression in (15.22) the relation

$$\mathbf{M}_{\pi\nu} | \mathbf{s}_\pi \mathbf{s}_\nu \rangle = 0 \tag{15.31}$$

holds. We see how the Majorana operator $\mathbf{M}_{\pi\nu}$ contributes to diagonal elements of the Hamilton matrix on the basis (15.22) and (15.23). According to (15.27), (15.28), (15.30) and (15.31) only the diagonal matrix elements of the antisymmetric states (15.23) are non-vanishing and positive if the coefficient λ in (15.16) is positive. By means of λ one lifts the levels of the states which are antisymmetric with respect to π and ν . Doing so, they lose all spectroscopic

significance. One can show that $\mathbf{M}_{\pi\nu}$ acting on more complex basis states has the same effect.

15.4 Eigenenergies and electromagnetic transitions within the IBM2

Having discussed the individual terms of the IBM2-Hamiltonian (see 15.15 up to 15.20) we put it together

$$\begin{aligned} \mathbf{H} = & \varepsilon_{\pi} \mathbf{n}_{d\pi} + \varepsilon_{\nu} \mathbf{n}_{d\nu} + \\ & \sum_{\rho=\pi,\nu} \frac{1}{2} \sum_{J=0,2,4} c_{J\rho} \sqrt{(2J+1)} [[\mathbf{d}_{\rho}^{+} \times \mathbf{d}_{\rho}^{+}]^{(J)} \times [\mathbf{d}_{\rho}^{-} \times \mathbf{d}_{\rho}^{-}]^{(J)}]^{(0)} + \\ & \kappa \sqrt{5} [\mathbf{Q}_{\pi}(\chi_{\pi}) \times \mathbf{Q}_{\nu}(\chi_{\nu})]^{(0)} + \lambda \mathbf{M}_{\pi\nu}. \end{aligned} \quad (15.32)$$

It contains 12 parameters. The eigenenergies of \mathbf{H} are calculated on a basis of functions analogous to (12.2) which are composed of the basis states (15.12) like this

$$|l, N_{\pi} N_{\nu} J M\rangle = \sum_i c^{(l)}_{N_{\pi} N_{\nu} J}(i) |i, N_{\pi} N_{\nu} J M\rangle. \quad (15.33)$$

The coefficients $c^{(l)}_{N_{\pi} N_{\nu} J}(i)$ of the l th state result from the diagonalisation procedure discussed in section 12.1. The sum comprises the configurations $i = (n_{s\pi} n_{d\pi} \tau_{\pi} n_{\Delta\pi} J_{\pi} n_{s\nu} n_{d\nu} \tau_{\nu} n_{\Delta\nu} J_{\nu})$ according to (15.12). The eigenvalue is labelled with $E^{(l)}_{N_{\pi} N_{\nu} J}$.

The probabilities for electromagnetic transitions between states of this kind are formulated analogously to the method described in chapter 11. The reduced transition probability $B(L, J_i \rightarrow J_f)$ is calculated by means of the initial and the final state of the type (15.33). For the operator $\mathbf{O}(E L M)$ for an electromagnetic transition with the angular momentum L , the form (11.4) is extended to a sum of the operator for π -bosons and the one for ν -bosons.

Specially for electric transitions with $L = 2$ (quadrupole radiation) we have analogously to (11.12) the following expression

$$\mathbf{O}(E 2 \mu) = \sum_{\rho=\pi,\nu} e_{\rho} (\mathbf{d}_{\rho\mu}^{+} \mathbf{s}_{\rho} + \mathbf{s}_{\rho}^{+} \mathbf{d}_{\rho\mu}^{-} + \chi'_{\rho} [\mathbf{d}_{\rho}^{+} \times \mathbf{d}_{\rho}^{-}]^{(2)}_{\mu}). \quad (15.34)$$

The operators in (15.34) correspond largely to the operator $\mathbf{Q}_{\rho}(\chi_{\rho})_{\mu}$ in (15.17), for which reason frequently one puts $\chi'_{\rho} = \chi_{\rho}$. This demands that this parameter is fitted simultaneously on energies and on transition probabilities. In this case, we have

$$\mathbf{O}(E 2 \mu) = e_{\pi} \mathbf{Q}_{\pi}(\chi_{\pi})_{\mu} + e_{\nu} \mathbf{Q}_{\nu}(\chi_{\nu})_{\mu}. \quad (15.35)$$

Frequently the quantity e_{π} is equated with e_{ν} . T. Otsuka and O. Scholten have written the computer program NPBOSS which performs IBM2-calculations essentially according to (15.32) (T. Otsuka, 1977, University Tokyo). The newer program BOSON2 (A. Novoselsky, 1985, Weizmann Institute, Israel) is able to handle with a great number of bosons.




Figure 15.1. Comparison between the measured and the calculated energies of the levels 2^+_{11} , 4^+_{11} , 6^+_{11} and 8^+_{11} of isotopes of ^{54}Xe with even numbers of neutrons 62 up to 78. The IBM2-Hamiltonian has been employed (Scholten, 1980, p. 55).

15.5 Comparisons with experimental data

Figures 15.1 up to 15.3 show experimental and calculated energies of levels of ^{54}Xe and ^{56}Ba isotopes. They are taken from the report of O. Scholten (1980). The Hamiltonian (15.32) has been employed with the simplification $e_\pi = e_\nu$. The Majorana term $M_{\pi\nu}$ has been formulated more generally.




Figure 15.2. Comparison between the measured and the calculated energies of the levels 5^+_1 , 4^+_2 , 3^+_1 and 2^+_2 of isotopes of ^{54}Xe with even numbers of neutrons 64 up to 78 (Scholten, 1980, p. 55).




Figure 15.3. Comparison between the measured and the calculated energies of 8 levels of ^{56}Ba isotopes with even numbers of neutrons 64 up to 78. The IBM2-Hamiltonian has been employed (Scholten, 1980, p. 55).

The corresponding parameters ε and κ of the Hamilton operator are represented in figure 15.4 as a function of the neutron number. The dependence of these quantities on the proton number (54 up to 58) could be neglected.




Figure 15.4. The parameters κ (with negative values) and ε employed for the IBM2-calculations shown in figures 15.1 up to 15.3 for nuclei with even neutron numbers from 54 up to 78 and proton numbers 54 up to 58 (Scholten, 1980, p. 57).




Figure 15.5. The parameters χ_v , c_{0v} and c_{2v} employed for the IBM2-calculations shown in figures 15.1 up to 15.3 for nuclei with even neutron numbers 54 up to 78 (Scholten, 1980, p. 57).

Figure 15.5 shows the separate dependence of the quantities χ_v , c_{0v} and c_{2v} on the neutron number. The coefficient c_{4v} was put equal to zero. Also the quantities $c_{0\pi}$, $c_{2\pi}$ and $c_{4\pi}$ are nearly zero, but χ_π descends from -0.81 to -1 for proton numbers 54 up to 58. Furthermore, O. Scholten (1980) showed that the terms of c_{0v} , c_{2v} and c_{4v} have little influence. Compared with the quadrupole term $\kappa\sqrt{5}[\mathbf{Q}_\pi(\chi_\pi) \times \mathbf{Q}_v(\chi_v)]^{(0)}$ they are only important if there are little neutrons beside many active protons or vice versa.

Reduced transition probabilities for quadrupole radiation are calculated according to (15.35) and shown in figure 15.6 together with experimental data.




Figure 15.6. Comparison between measured and calculated reduced transition probabilities for quadrupole radiation of even-even nuclei with neutron numbers 68 up to 78. The parameters of figures 15.4 and 15.5 have been employed (Scholten, 1980, p. 58).

Duval and Barrett (1981) have performed similar IBM2-calculations for tungsten isotopes. They employed the parameters $\varepsilon_\pi = \varepsilon_\nu$ and $c_{4\nu} = c_{0\pi} = c_{2\pi} = c_{4\pi} = 0$. Figures 15.7 and 15.8 show the comparison with experimental data.




Figure 15.7. Comparison between the measured and calculated energies of the levels 2^+_1 , 4^+_1 , 6^+_1 and 8^+_1 of isotopes from ^{74}W with even neutron numbers 96 up to 114 employing the IBM2 formalism (Duval and Barrett, 1981, p. 493).




Figure 15.8. Comparison between measured and calculated energies of 6 levels of ^{74}W with even numbers of neutrons 106 up to 116. The IBM2-Hamiltonian has been employed (Duval and Barrett, 1981, p. 494).

In figures 15.9 and 15.10, the behaviour of the corresponding IBM2-parameters depending on the neutron number is shown.




Figure 15.9. The parameters κ (with negative values) and ϵ employed for the IBM2-calculations shown in figures 15.7 and 15.8 for tungsten isotopes with even neutron numbers 96 up to 114 (Duval and Barrett, 1981, p. 495).




Figure 15.10. The parameters χ_v , c_{0v} and c_{2v} employed for the IBM2-calculations shown in figures 15.7 and 15.8 for tungsten isotopes with even neutron numbers 96 up to 114 (Duval and Barrett, 1981, p. 495).

Figure 15.11 shows some reduced transition probabilities $B(E2)$ for γ -radiation and compares calculated with measured values. Jolos et al. (1985) have performed similar calculations for isotopes of $_{44}\text{Ru}$ and $_{46}\text{Pd}$ with equal success.




Figure 15.11. Comparison between measured and calculated reduced transition probabilities for quadrupole radiation of tungsten isotopes with neutron numbers 106 up to 112 (Duval and Barrett, 1981, p. 495 up to p. 497). The employed parameters are shown in figures 15.9 and 15.10 and described in the text.

As we have mentioned above (Scholten, 1980) little changes if all coefficients $c_{J\rho}$ in (15.32) are equal to zero and $\varepsilon_\pi = \varepsilon_\nu$. In this case the Hamilton operator obtains the following form

$$H = \varepsilon (n_{d\pi} + n_{d\nu}) + \kappa \sqrt{5} [Q_\pi(\chi_\pi) \times Q_\nu(\chi_\nu)]^{(0)} + \lambda M_{\pi\nu}. \quad (15.36)$$

With this Hamiltonian, which contains essentially four parameters (frequently λ is fixed arbitrarily), for many nuclei good agreement could be achieved with measured data. Figure 0.1 puts together existing calculations up to 1988. The expression (15.36) is named Talmi's Hamiltonian (Iachello, Arima, 1987, p.140). In the same publication on page 140 a group of authors is given which have done calculations with this Hamilton operator.

15.6 Chains of Lie algebras in the IBM2

Owing to the relationship between the IBM2- and the IBM1-Hamiltonian we expect also chains of Lie algebras such as in (14.11), (14.17), (14.21) and table 14.1.

First, we consider the term $V_{\pi\nu}$ (15.7). It contains a sum comprising expressions of the following form

$$[[b^+_{\pi j_f} \times b^+_{\nu j_g}]^{(J)} \times [b^-_{\pi j_p} \times b^-_{\nu j_q}]^{(J)}]^{(0)}, (j_f, j_g, j_p, j_q = 0, 2) \quad (15.37)$$

where the labels π and ν are ordered. With the aid of the 9-j symbols, (A4.1) one obtains sums of the following expressions

$$[[b^+_{\pi j_f} \times b^-_{\pi j_p}]^{(J)} \times [b^+_{\nu j_g} \times b^-_{\nu j_q}]^{(J)}]^{(0)}. \quad (15.38)$$

On the other hand the terms H_ρ ($\rho = \pi, \nu$) in (15.4) consist according to (14.2), (14.3), (14.5) and (14.7) on sums of expressions such as

$$[b^+_{\rho j_f} \times b^-_{\rho j_p}]^{(0)} \text{ and } [[b^+_{\rho j_f} \times b^-_{\rho j_p}]^{(J)} \times [b^+_{\rho j_g} \times b^-_{\rho j_q}]^{(J)}]^{(0)}. \quad (15.39)$$

We single out special cases of Hamilton operators (15.4) which are able to constitute chains of Lie algebras.

First series of special cases of the IBM2

Here all corresponding coefficients in H_π and H_ν (14.2) are put equal i. e.

$$\varepsilon_{\pi n} = \varepsilon_{\nu n} \equiv \varepsilon_n, \dots, V_{\pi n} = V_{\nu n} \equiv V_n, \dots, V_{\pi q} = V_{\nu q} \equiv V_q. \quad (15.40)$$

If one replaces in H (14.2) (14.3), (14.5) and (14.7) every term of the form

$$[b^+_{j_a} \times b^-_{j_b}]^{(J)}_M, (j_a, j_b = 1, 2) \quad (15.41)$$

by expressions such as

$$G^{(J)}_M(j_a, j_b) = [b^+_{\pi j_a} \times b^-_{\pi j_b}]^{(J)}_M + [b^+_{\nu j_a} \times b^-_{\nu j_b}]^{(J)}_M, (j_a, j_b = 1, 2) \quad (15.42)$$

one obtains by means of (15.40) just the sum $H_\pi + H_\nu$ plus a sum of terms such as (15.38). As an additional restriction in this special case one demands that these residual terms cancel out with the terms of $V_{\pi\nu}$. This results in several conditions for the involved coefficients. Some authors leave a Majorana term (15.20) (Bonatsos, 1988, p. 78), (Frank and van Isacker, 1994, p. 369).

The remaining IBM2-Hamiltonian reads now analogously to (9.39) or (14.2)

$$H = \varepsilon_n N + v_n N^2 + \quad (15.43)$$

$$(\varepsilon_d' + v_{nd} N) n_{d, \pi+\nu} + v_d n_{d, \pi+\nu}^2 + v_r R_{\pi+\nu}^2 + v_t T_{\pi+\nu}^2 + v_j J_{\pi+\nu}^2 + v_q Q_{\pi+\nu}^2$$

where in each operator the basis operators (15.41) are replaced by (15.42). These make up the same chains of algebras as in the IBM1. The formulation of

these chains can be taken from (14.11), (14.17), (14.21) and table 14.1. They are put together in table 15.2 with usual markings.

Table 15.2. Chains of Lie algebras of the special cases I₁, II₁ and III₁ of the IBM2

Basis elements	Dimension	Algebra	Special case
$\mathbf{G}^{(2)}_{\mu}(2,0), \mathbf{G}^{(2)}_{\mu}(0,2), \mathbf{G}^{(0)}_0(0,0)$	$\left. \begin{array}{c} n \\ 36 \end{array} \right\}$	$u_{\pi+\nu}(6)$	I ₁
$\mathbf{G}^{(0)}_0(2,2), \mathbf{G}^{(2)}_{\mu}(2,2), \mathbf{G}^{(4)}_{\mu}(2,2)$	$\left. \begin{array}{c} 25 \end{array} \right\}$	$u_{\pi+\nu}(5)$	
$\mathbf{G}^{(3)}_{\mu}(2,2)$	$\left. \begin{array}{c} 10 \end{array} \right\}$	$so_{\pi+\nu}(5)$	
$\mathbf{G}^{(1)}_{\mu}(2,2)$	$\left. \begin{array}{c} 3 \end{array} \right\}$	$so_{\pi+\nu}(3)$	
$\mathbf{G}^{(2)}_{\mu}(2,0), \mathbf{G}^{(2)}_{\mu}(0,2), \mathbf{G}^{(0)}_0(0,0),$ $\mathbf{G}^{(0)}_0(2,2), \mathbf{G}^{(3)}_{\mu}(2,2), \mathbf{G}^{(4)}_{\mu}(2,2)$	$\left. \begin{array}{c} 36 \end{array} \right\}$	$u_{\pi+\nu}(6)$	II ₁
$\mathbf{G}^{(2)}_{\mu}(2,0) + \mathbf{G}^{(2)}_{\mu}(0,2) - (\sqrt{7}/2) \mathbf{G}^{(2)}_{\mu}(2,2)$	$\left. \begin{array}{c} 8 \end{array} \right\}$	$su_{\pi+\nu}(3)$	
$\mathbf{G}^{(1)}_{\mu}(2,2)$	$\left. \begin{array}{c} 3 \end{array} \right\}$	$so_{\pi+\nu}(3)$	
$\mathbf{G}^{(2)}_{\mu}(0,2), \mathbf{G}^{(0)}_0(0,0),$ $\mathbf{G}^{(0)}_0(2,2), \mathbf{G}^{(2)}_{\mu}(2,2), \mathbf{G}^{(4)}_{\mu}(2,2)$	$\left. \begin{array}{c} 36 \end{array} \right\}$	$u_{\pi+\nu}(6)$	III ₁
$\mathbf{G}^{(2)}_{\mu}(2,0) + \mathbf{G}^{(2)}_{\mu}(0,2)$	$\left. \begin{array}{c} 15 \end{array} \right\}$	$so_{\pi+\nu}(6)$	
$\mathbf{G}^{(3)}_{\mu}(2,2)$	$\left. \begin{array}{c} 10 \end{array} \right\}$	$so_{\pi+\nu}(5)$	
$\mathbf{G}^{(1)}_{\mu}(2,2)$	$\left. \begin{array}{c} 3 \end{array} \right\}$	$so_{\pi+\nu}(3)$	

As discussed in the preceding chapter every Lie algebra generates a set of basis functions. Their members are labelled with quantum numbers, which appear partly in the eigenvalues of the Casimir operators. In the IBM1 there is – except for $su(3)$ – only one quantum number per algebra, which is relevant for the Casimir operator. In the IBM2 there are mostly two such numbers as the investigation of van Isacker et al. (1984) shows. For the special case I₁ (table 15.2) result the following quantum numbers

$$\left| \begin{array}{l} u_{\pi+\nu}(6) \supset u_{\pi+\nu}(5) \supset so_{\pi+\nu}(5) \supset so_{\pi+\nu}(5) \\ [N_1, N_2] \quad [n_1, n_2] \quad [\tau_1, \tau_2], n_{\Delta 1}, n_{\Delta 2} \quad [J] \end{array} \right\rangle \quad l_1 \quad (15.44)$$

with $N_1 + N_2 = N$ ($N_1 \geq N_2$), $n_1 + n_2 = n_{d\pi} + n_{d\nu}$.

The quantum numbers in square brackets characterise the eigenvalues of the Casimir operators (see 15.47) and the remaining numbers serve to the classification of the states. Analogously to the IBM1, the relations

$$n_1 \leq N_1, n_2 \leq N_2, \tau_1 \leq n_1, \tau_2 \leq n_2 \quad (15.45)$$

hold together with further conditions for these quantum numbers and for the angular momentum. Iachello and Arima (1987, p. 152) have put them together.

By cancelling the coefficients v_r and v_q (15.43), one obtains analogously to the IBM1 (section 14.4) the combination of Casimir operators which belong to the first chain l_1 in table 15.2. In agreement with (14.37) the Hamilton operator can be written as

$$\begin{aligned} H^{(l_1)} = & \alpha_1 C_{1,u}(6), \pi+\nu + \alpha_2 C_{2,u}(6), \pi+\nu + \beta C_{1,u}(6), \pi+\nu C_{1,u}(5), \pi+\nu + \\ & \gamma_1 C_{1,u}(5), \pi+\nu + \gamma_2 C_{2,u}(5), \pi+\nu + \delta C_{2,so}(5), \pi+\nu + \varepsilon C_{2,so}(3), \pi+\nu. \end{aligned} \quad (15.46)$$

It contains Casimir operators which are formed by the operators $G^{(J)}_M(j_a, j_b)$ (15.42) in the same way as the operators in table 14.2. The formulas for eigenvalues of Casimir operators depending on quantum numbers are given by Iachello and Arima (1987, p. 37, p. 158). According to (15.46) the following eigenenergies for the special case l_1 result

$$\begin{aligned} E_{l_1} = & \alpha_1 (N_1 + N_2) + \alpha_2 (N_1(N_1 + 5) + N_2(N_2 + 3)) + \\ & \beta (N_1 + N_2)(n_1 + n_2) + \gamma_1 n_1(n_1 + 2) + \\ & \gamma_2 (n_1(n_1 + 4) + n_2(n_2 + 2)) + \delta (\tau_1(\tau_1 + 3) + \tau_2(\tau_2 + 1)) + \varepsilon J(J + 1). \end{aligned} \quad (15.47)$$

For large boson numbers N , one obtains a multiplicity of possible splittings in N_1 and N_2 but only the lowest levels are important. Figure 15.12 shows a spectrum according to (15.47) for two active π -bosons and a ν -boson. The part with $N_1 = 3$ and $N_2 = 0$ corresponds to the IBM1 spectrum for the vibrational limit.




Figure 15.12. Typical spectrum for the I_1 -special case of the IBM2. $N_\pi = 2$ and $N_\nu = 1$ (or vice versa) have been employed (Iachello and Arima, 1987, p.159).

For the chains of the algebras II_1 and III_1 (table 15.2) there exist representations of eigenvalues corresponding to (15.46) and (15.47) (Iachello and Arima, 1987, p. 159). In appropriate publications further chains of algebras are mentioned, in which the higher algebras are direct sums of π - and ν -algebras.

16 The interacting boson-fermion model IBFM

In the first section the extension of the IBM will be outlined which admits odd numbers of protons and/or neutrons. Creation and annihilation operators for nucleons are introduced and the corresponding Hamiltonian is brought in the usual form. The second section deals with a version of the IBMF which employs the boson-boson interaction of the $u(5)$ -limit. The simple case of one active d -boson coupled to a nucleon is discussed. In the third section we will explain how the number of parameters in the boson-nucleon interaction can be reduced realistically. Comparisons of IBFM calculations with measurements will follow.

16.1 The Hamiltonian of the IBFM

In the preceding chapters the nuclear model was restricted to nuclei with even proton- and neutron numbers. However, most of the nuclei are composed of odd-even (oe) or odd-odd (oo) nucleon numbers.

In order to include these nuclei, the interacting boson model is extended by attaching a nucleon to the boson configuration. For oo-nuclei, one has to add a proton and a neutron. The number of bosons is fixed in the same way as in chapter 2. If the active bosons are holes, the nucleon has to be taken in the hole state as well.

Nucleons are known to have total spins $j = 1/2, 3/2, 5/2, \dots$ with regard to the nucleus and they can show positive or negative parity. Coupling a spherical basis state for bosons (chapter 3, (6.18)) to a nucleon state creates the simplest basis state for oe-nuclei like this

$$|N_B n_d \tau n_\Delta J_B j J M\rangle \equiv [|(N_B n_d \tau n_\Delta)^{(J_B)} \times j|^{(j)}]_M. \quad (16.1)$$

For nucleons, there exist also creation and annihilation operators as follows

$$a_{jm}^+, a_{jm} \quad (16.2)$$

which are assigned to the nucleon state (j, m) . Analogously to (6.19) the annihilation operator can be put in tensor form like this

$$a_{jm}^- = (-1)^{j+m} a_{j, -m}. \quad (16.3)$$

Because the nucleon operators (16.2) create (or annihilate) antisymmetric states, the commutation rules (5.34) and (5.35) are not valid and the commutators have to be replaced by anticommutators. In analogy with the IBM2 (section 15.1) one demands that a_{jm}^+ and a_{jm}^- commute with the boson creation and annihilation operators. That is to say, bosons and nucleons are regarded as essentially different objects and one forgets that a boson represents a nucleon pair.

The Hamilton operator for bosons and nucleons contains, besides the interaction between the bosons, also nucleon-nucleon and nucleon-boson interactions. Analogously to (15.4) it reads

$$H = H_B + H_F + V_{BF}. \quad (16.4)$$

For the boson part H_B , mostly the IBM1-Hamilton operator (7.17) is employed. The pure nucleon Hamiltonian H_F is constructed analogously to (7.3) and (7.6) as follows

$$H_F = E_0 + \sum_j \varepsilon_j \sqrt{(2j+1)} [\mathbf{a}_j^+ \times \mathbf{a}_j^-]^{(0)} + \sum_{JJ'j_1j_2j_3j_4} c^{(J)}_{JJ'j_1j_2j_3j_4} [[\mathbf{a}_j^+ \times \mathbf{a}_j^-]^{(J)} \times [\mathbf{a}_{j_1}^- \times \mathbf{a}_{j_2}^-]^{(J)}]^{(0)}. \quad (16.5)$$

Henceforth we will restrict ourselves to oe-nuclei. Therefore, the last sum in (16.5) disappears because it does not contribute when it acts on a single nucleon. The quantity ε_j is the single-nucleon energy of the nucleon state (jm) . In analogy with (15.7) the boson-fermion interaction can be written as

$$V_{BF} = \sum_{l_1 l_2 j_1 j_2} v^{(j)}_{l_1 j_1 l_2 j_2} [[\mathbf{b}_{l_1}^+ \times \mathbf{a}_{j_1}^-]^{(j)} \times [\mathbf{b}_{l_2}^- \times \mathbf{a}_{j_2}^-]^{(j)}]^{(0)} \quad (16.6)$$

with $(l_1, l_2) = 0, 2$ and $(j_1, j_2) = 1/2, 3/2, 5/2, \dots$. The b -operators concern bosons. The coefficients $v^{(j)}_{l_1 j_1 l_2 j_2}$ are matrix elements of the boson-fermion interaction this way

$$v^{(j)}_{l_1 j_1 l_2 j_2} = \langle b_{l_1} a_{j_1} j | V_{BF} | b_{l_2} a_{j_2} j \rangle. \quad (16.7)$$

Analogously to (15.9) there is linear dependency between some of the quantities $v^{(j)}_{l_1 j_1 l_2 j_2}$. Frequently it is sufficient to employ only one nucleon spin $j_1 = j_2 = j_n$. The expression (16.6) can be recoupled by means of (A4.1), which yields

$$V_{BF} = \sum_{l_1 l_2 j_1 j_2} c^{(j)}_{l_1 j_1 l_2 j_2} [[\mathbf{b}_{l_1}^+ \times \mathbf{b}_{l_2}^-]^{(j)} \times [\mathbf{a}_{j_1}^+ \times \mathbf{a}_{j_2}^-]^{(j)}]^{(0)}. \quad (16.8)$$

The coefficients $c^{(j)}_{l_1 j_1 l_2 j_2}$ are linear combinations of $v^{(j)}_{l_1 j_1 l_2 j_2}$ and their factors contain 9-j symbols. We write (16.8) in detail employing d - and s -operators

$$\begin{aligned} V_{BF} = & \sum_j c^{(0)}_{00jj} [[\mathbf{s}^+ \times \mathbf{s}]^{(0)} \times [\mathbf{a}_j^+ \times \mathbf{a}_j^-]^{(0)}]^{(0)} + \\ & \sum_j c^{(0)}_{22jj} [[\mathbf{d}^+ \times \mathbf{d}]^{(0)} \times [\mathbf{a}_j^+ \times \mathbf{a}_j^-]^{(0)}]^{(0)} + \sum_{j_1 j_2} c^{(2)}_{02j_1 j_2} \cdot \\ & [([\mathbf{s}^+ \times \mathbf{d}]^{(2)} + [\mathbf{d}^+ \times \mathbf{s}]^{(2)} + (c^{(2)}_{22j_1 j_2} / c^{(2)}_{02j_1 j_2}) [\mathbf{d}^+ \times \mathbf{d}]^{(2)} \times [\mathbf{a}_{j_1}^+ \times \mathbf{a}_{j_2}^-]^{(2)}]^{(0)} + \\ & \sum_{j_1 j_2, J=1,3,4} c^{(J)}_{22j_1 j_2} [[\mathbf{d}^+ \times \mathbf{d}]^{(J)} \times [\mathbf{a}_{j_1}^+ \times \mathbf{a}_{j_2}^-]^{(J)}]^{(0)}. \end{aligned} \quad (16.9)$$

$$\text{The expressions } [[\mathbf{d}^+ \times \mathbf{d}]^{(J)} \times [\mathbf{a}_{j_1}^+ \times \mathbf{a}_{j_2}^-]^{(J)}]^{(0)} \quad (16.10)$$

in the last term of (16.9) are frequently written in another form. According to (A1.4) we have

$$[\mathbf{a}_{j_1}^+ \times \mathbf{a}_{j_2}^-]^{(J)}_M = (-1)^{j_1 + j_2 - J} \sum_{m_1 m_2} (j_2 \ m_2 \ j_1 \ m_1 | J \ M) \mathbf{a}_{j_1}^+ \mathbf{a}_{j_2}^-. \quad (16.11)$$

Now we are defining the operator of normal ordering of nucleon operators as follows

$$: \mathbf{a}_{j_2}^{\sim} \mathbf{a}_{j_1}^{\dagger} : \equiv \mathbf{a}_{j_1}^{\dagger} \mathbf{a}_{j_2}^{\sim}. \quad (16.12)$$

Therefore, we can write

$$[\mathbf{a}_{j_1}^{\dagger} \times \mathbf{a}_{j_2}^{\sim}]^{(J)}_M = (-1)^{j_1 + j_2 - J} : [\mathbf{a}_{j_2}^{\sim} \times \mathbf{a}_{j_1}^{\dagger}]^{(J)}_M :. \quad (16.13)$$

This means that the right hand side of (16.13) first is represented formally by means of (A1.1) and then the interchanging (16.12) is performed. Now the expression (16.10) is written with the aid of (A4.1) as follows

$$[[\mathbf{d}^{\dagger} \times \mathbf{d}^{\sim}]^{(J)} \times [\mathbf{a}_{j_1}^{\dagger} \times \mathbf{a}_{j_2}^{\sim}]^{(J)}]^{(0)} = \quad (16.14)$$

$$\sum_j (2J+1)(2j+1) \begin{pmatrix} 2 & 2 & J \\ j_1 & j_2 & J \\ j & j & 0 \end{pmatrix} (-1)^{j_2 - j} : [[\mathbf{d}^{\dagger} \times \mathbf{a}_{j_2}^{\sim}]^{(j)} \times [\mathbf{a}_{j_1}^{\dagger} \times \mathbf{d}^{\sim}]^{(j)}]^{(0)} :.$$

Therefore, the last sum in (16.9) obtains the following form

$$\sum_{j_1 j_2, J=1,3,4} c_{22j_1 j_2}^{(J)} [[\mathbf{d}^{\dagger} \times \mathbf{d}^{\sim}]^{(J)} \times [\mathbf{a}_{j_1}^{\dagger} \times \mathbf{a}_{j_2}^{\sim}]^{(J)}]^{(0)} = \quad (16.15)$$

$$\sum_{j j_1 j_2} \Lambda_{j_2 j_1}^{(j)} : [[\mathbf{d}^{\dagger} \times \mathbf{a}_{j_2}^{\sim}]^{(j)} \times [\mathbf{a}_{j_1}^{\dagger} \times \mathbf{d}^{\sim}]^{(j)}]^{(0)} :.$$

With the help of the number operators (6.35) up to (6.37) such as

$$[\mathbf{s}^{\dagger} \times \mathbf{s}]^{(0)} = \mathbf{N} - \mathbf{n}_d, \quad [\mathbf{d}^{\dagger} \times \mathbf{d}^{\sim}]^{(0)} = \mathbf{n}_d / \sqrt{5}, \quad [\mathbf{a}_j^{\dagger} \times \mathbf{a}_j^{\sim}]^{(0)} = n_j / \sqrt{(2j+1)}, \quad (16.16)$$

with the usual markings (Iachello and van Isacker, 1991, p. 12) like this

$$c_{00jj}^{(0)} \sqrt{5} - c_{22jj}^{(0)} = A_j, \quad c_{02j_1 j_2}^{(2)} = \Gamma_{j_1 j_2}, \quad (c_{22j_1 j_2}^{(2)} / c_{02j_1 j_2}^{(2)}) = \chi_{j_1 j_2} \quad (16.17)$$

and with the quadrupole operator (15.17)

$$\mathbf{Q}_B (\chi_{j_1 j_2})_{\mu} = [\mathbf{d}^{\dagger} \times \mathbf{s}]_{\mu}^{(2)} + [\mathbf{s}^{\dagger} \times \mathbf{d}]_{\mu}^{(2)} + \chi_{j_1 j_2} [\mathbf{d}^{\dagger} \times \mathbf{d}]_{\mu}^{(2)} \quad (16.18)$$

one obtains for \mathbf{V}_{BF} (16.9) the following expression

$$\mathbf{V}_{BF} = \sum_j c_{00jj}^{(0)} \mathbf{N} n_j - \sum_j A_j \mathbf{n}_d n_j / \sqrt{(5(2j+1))} + \quad (16.19)$$

$$\sum_{j_1 j_2} \Gamma_{j_1 j_2} [\mathbf{Q}_B (\chi_{j_1 j_2}) \times [\mathbf{a}_{j_1}^{\dagger} \times \mathbf{a}_{j_2}^{\sim}]^{(2)}]^{(0)} +$$

$$\sum_{j j_1 j_2} \Lambda_{j_2 j_1}^{(j)} : [[\mathbf{d}^{\dagger} \times \mathbf{a}_{j_2}^{\sim}]^{(j)} \times [\mathbf{a}_{j_1}^{\dagger} \times \mathbf{d}^{\sim}]^{(j)}]^{(0)} :.$$

The second term on the right hand side of (16.19) is named monopole term and the following are the quadrupole term and the exchange term.

16.2 The $u(5)$ -limit of the IBFM

If the part H_B of the Hamiltonian (16.4) has the special form (14.31) of the $u(5)$ -limit, the state (16.1) is an eigenfunction of H_B with the following eigenvalue (10.17)

$$E_B^{(1)} = \varepsilon_n N + v_n N^2 + (\varepsilon_d' + v_{nd} N) n_d + v_d n_d^2 + v_t \tau(\tau + 3) + v_j J_B(J_B + 1). \quad (16.20)$$

It can be written in the form (14.38) as well. The matrix of the operator H_B on the basis of functions (16.1) has therefore only diagonal elements. Naturally H_F (for one single nucleon) is diagonal and has the eigenvalues ε_j . In order to find the eigenenergies of the whole operator H (16.4) essentially only V_{BF} has to be diagonalised (see section 12.1). We investigate the influence of the individual terms of V_{BF} (16.19). Because the monopole term on the right hand side of (16.19) contains only number operators it gives an identical contribution to every diagonal matrix element with the same number of d -bosons and the same nucleon state $|j m\rangle$. The quadrupole term in V_{BF} contains the operators

$$\sum_{j_1 j_2} \Gamma_{j_1 j_2} [[\mathbf{d}^+ \times \mathbf{s} + \mathbf{s}^+ \times \mathbf{d}]^{(2)} \times [\mathbf{a}_{j_1}^+ \times \mathbf{a}_{j_2}^-]^{(2)}]^{(0)} \text{ and} \quad (16.21)$$

$$\sum_{j_1 j_2} \chi_{j_1 j_2} \Gamma_{j_1 j_2} [[\mathbf{d}^+ \times \mathbf{d}]^{(2)} \times [\mathbf{a}_{j_1}^+ \times \mathbf{a}_{j_2}^-]^{(2)}]^{(0)}. \quad (16.22)$$

The expression (16.21) generates non-diagonal elements in the energy matrix. Experience shows that its contributions can be neglected if ε_d' (16.20) is relatively large. The term (16.22) contributes to the diagonal matrix elements, which we now look into for the situation of a single d -boson coupled to a nucleon state $|j m\rangle$. Such a diagonal term reads

$$\begin{aligned} & \langle s^{N-1} [d \times j]^{(j)}_M | [[\mathbf{d}^+ \times \mathbf{d}]^{(2)} \times [\mathbf{a}_j^+ \times \mathbf{a}_j^-]^{(2)}]^{(0)} | s^{N-1} [d \times j]^{(j)}_M \rangle = \quad (16.23) \\ & \sum_{J'} 5(2J' + 1) \{^2_{j J'} \quad ^2_{j J'} \quad ^2_{20}\} \langle [d \times j]^{(j)}_M | [[\mathbf{d}^+ \times \mathbf{a}_j^+]^{(J')} \times [\mathbf{d}^- \times \mathbf{a}_j^-]^{(J')}]^{(0)} | [d \times j]^{(j)}_M \rangle = \\ & \sum_{J' M'} \sqrt{5} (-1)^{j+2J'-M'} \{^2_{j \quad j_2 \quad J'}\} \langle [d \times j]^{(j)}_M | [\mathbf{d}^+ \times \mathbf{a}_j^+]^{(J')}_{M'} [\mathbf{d}^- \times \mathbf{a}_j^-]^{(J')}_{-M'} | [d \times j]^{(j)}_M \rangle. \end{aligned}$$

The first transformation in (16.23) is a recoupling according to (A4.1) and the second employs the relations (4.2), (A1.1) and (A1.16). Furthermore, because of (16.3), (A1.1), (A1.5) and (A1.10) the following equation holds

$$\begin{aligned} & [\mathbf{d}^- \times \mathbf{a}_j^-]^{(J')}_{-M'} | [d \times j]^{(j)}_M \rangle = \\ & \sum_{\mu' m' \mu m} (-1)^{\mu' + j + m'} (2 \mu' j m' | J', -M') (2 \mu j m | J M) d_{\mu'} a_{j-m'} | d_{\mu} a_{j m} \rangle = \\ & \sum_{\mu' m' \mu m} (-1)^{j-M'} (2, -\mu j, -m | J', -M') (2 \mu j m | J M) \delta_{\mu, -\mu'} \delta_{m, -m'} = \\ & (-1)^{2j-J'-M'} \delta_{JJ'} \delta_{MM'}. \quad (16.24) \end{aligned}$$

Analogously we have

$$\langle [d \times j]^{(j)}_M | [\mathbf{d}^+ \times \mathbf{a}_j^+]^{(J')}_{-M'} = \delta_{JJ'} \delta_{MM'}. \quad (16.25)$$

Therefore, the term (16.22) yields the following contribution to the diagonal matrix element

$$\Delta E_{nd=1} = -\chi_{jj} \Gamma_{jj} \sqrt{5} (-1)^{J-j} \left\{ \begin{matrix} 2 & j & J \\ j & 2 & j \end{matrix} \right\}. \quad (16.26)$$

The energy of the state represented by a j -nucleon coupled up to a d -boson splits up into $2j + 1$ or 5 (if $j > 2$) levels starting from the energy $(\varepsilon_d' + v_{nd} N) + v_d + 4v_t + 6v_j$ (16.20). The splitting is proportional to the geometrical factors (16.26), which depend on J ($2 + j \geq J \geq |2 - j|$). The exchange term in \mathbf{V}_{BF} (16.19) yields a similar multiplicity of terms as (16.26).

Figure 16.1 shows a comparison between measured and calculated energies of the nucleus $^{133}_{57}\text{La}_{76}$ for d -boson configurations of the $u(5)$ -limit. The 2^+ -state of $^{152}_{56}\text{Ba}_{76}$ ($n_d = 1$) is split up into 5 levels by coupling a $j = 11/2$ -proton according to (16.26).




Figure 16.1. Comparison between measured and calculated energies of low-lying levels of ^{133}La . The energies of the 5 excited levels are calculated starting from the 2^+ -level of ^{132}Ba (consisting of one d -boson in the $u(5)$ -limit) with the aid of equation (16.26) and with the coefficient $\chi_{jj} \Gamma_{jj} = -0.94 \text{ MeV}$ (Scholten, 1980, p. 98).

Figure 16.2 compares experimental data with results of IBFM calculations in the $u(5)$ -limit of two Eu-isotopes.




Figure 16.2. Comparison between measured and calculated energy levels of ^{147}Eu and ^{151}Eu (Casten, 1980, p. 199). The calculations are performed according to the equation (16.19) and Arima and Iachello (1976b) respectively, by coupling a $11/2^-$ -proton up to $u(5)$ -states and by fitting the parameters. The ground states ($J = 11/2^-$) are drawn in, taking into account the binding energies.

Nuclei with boson configurations belonging to the other two special cases of dynamical symmetries are treated analogously.

16.3 Numerical treatment of the IBFM. Comparison with experimental data

The general operators H_B and H_F (16.4) and V_{BF} (16.19) comprise so many free parameters that the model becomes cumbersome. One has to reduce them suitably similarly as in the IBM2. In this section we deal specially with the term V_{BF} .

Owing to the microscopic theory, which connects the IBM with the shell model, and making use of the generalised seniority principle, Scholten (1981, p.286) gives the following instruction for the coefficients in (16.19), which represents an approximation

$$\begin{aligned} A_j &= -\sqrt{5(2j+1)} A_0, \\ \Gamma_{j_1 j_2} &= \sqrt{5} \gamma_{j_1 j_2} \Gamma_0, \quad \text{with } \gamma_{j_1 j_2} = (u_{j_1} u_{j_2} - v_{j_1} v_{j_2}) Q_{j_1 j_2}, \\ \Lambda_{j_1 j_2}^{(j)} &= -2 \sqrt{5/(2j+1)} \beta_{j_1 j} \beta_{j_2 j} \Lambda_0 \quad \text{with } \beta_{j' j} = (u_{j'} v_j + v_{j'} u_j) Q_{j' j}. \end{aligned} \quad (16.27)$$

The quantities $Q_{j' j}$ in (16.27) are reduced matrix elements of the quadrupole operator in the single-nucleon basis like this

$$Q_{j' j} = \langle j' || Y^{(2)} || j \rangle.$$

The quantities v_j^2 are occupation probabilities for the nucleon states (Ring and Schuck, 1980), (Otsuka and Arima, 1978). The relation $u_j = \sqrt{1 - v_j^2}$ holds. Apart from the constant term, the parameters of V_{BF} (16.19) are reduced to A_0 , Γ_0 and Λ_0 according to (16.27).

One finds the parameters of H_B in practice by fitting IBM1 calculations to the next lighter (or heavier) even-even nucleus which contains one nucleon (or nucleon hole) less than the even-odd nucleus.

Similarly as in chapter 12 the matrix elements of the Hamilton operator can be formed on the basis (16.1). By diagonalising, one obtains the eigenenergies and eigenstates.

Figure 16.3 shows results of such an IBFM calculation for $^{101}_{45}\text{Rh}_{56}$ compared with the measured level energies. States with positive parity are treated as a coupling of an IBM1 state to a $g_{9/2}$ -proton.




Figure 16.3. Comparison between measured and calculated low-lying levels with positive parity of ^{101}Rh (Scholten, 1981, p. 289). Not all experimental spin assignments are unambiguous. In the calculation a $g_{9/2^-}$ proton is coupled to a $^{100}_{44}\text{Ru}_{56}$ nucleus which has been treated with the IBM1 making use of the parameters $\varepsilon = 0.733$ MeV, $c_0' = -0.415$ MeV, $c_2' = -0.283$ MeV, $c_4' = -0.024$ MeV, $v_2 = 0.175$ MeV and $v_0 = -0.31$ MeV (7.31). For the fermion-boson term V_{BF} (16.19 and 16.27) the value $v_j^2 = 0.5$ has been chosen for which reason the quadrupole term vanishes.

The low-lying states of ^{191}Ir are calculated using a Hamiltonian containing an IBM1 part for bosons and parts with the proton state $(h_{11/2})^{-1}$. In figure 16.4, the results for negative parity are compared with experimental data.




Figure 16.4. Comparison between calculated and experimental spectra of the negative-parity states in ^{191}Ir (Bijker and Dieperink, 1982, p. 229). The broken lines denote calculated data. The levels are interpreted in the IBFM model as a $h_{11/2}^{-}$ proton hole state coupled to the $^{192}_{78}\text{Pt}_{114}$ core. The lines connecting the levels represent large E2 transitions by radiation.

Isotopes of gold of the odd-even type with 106 up to 116 neutrons have been treated as a coupling of an even-even Pt-core to a $h_{9/2}^-$ -proton. Figure 16.5 shows the lowest levels compared with experimental data.




Figure 16.5. Comparison between measured and calculated spectra with negative parity in ^{79}Au -isotopes (Bijker and Dieperink, 1982, p. 231). The neutron numbers comprise 104 up to 116. For the calculation, a $h_{9/2}^-$ -proton has been coupled to the corresponding ^{78}Pt -cores.

Figure 16.6 represents an IBFM-calculation for numerous levels with positive parity of ^{97}Tc . The even-even nucleus is coupled to a proton with two different shell states.

The IBFM comprises more dynamic symmetries than the IBM2. Because this book is only an introduction to the interacting boson model, we leave them out of consideration.




Figure 16.6. Comparison between the measured and the calculated spectrum with positive parity of ^{97}Tc (von Brentano, 1981, p. 307). In the calculation, a proton with the states $g_{9/2}$ and $d_{5/2}$ has been coupled to ^{96}Mo . This nucleus has been treated with the IBM1 by means of the parameters $\varepsilon = 0.795$ MeV, $c_0' = -0.25$ MeV, $c_2' = -0.085$ MeV, $c_4' = 0.047$ MeV, $v_2 = 0.113$ MeV and $v_0 = 0$ MeV (7.31). The fermion-boson term V_{BF} (16.18) has been calculated with expressions, which correspond largely to the equations (16.19) and (16.27).

Appendix

A1 Clebsch-Gordan coefficients and 3- j symbols

Some properties of the Clebsch-Gordan coefficients will be reviewed which are employed in the treatment of the IBM on hand. Descriptions of this topic in detail are given by Condon and Shortley (1953) and Talmi (1993).

Clebsch-Gordan coefficients arise when state functions containing a definite spin or angular momentum are coupled and constitute a joint state with good spin quantum numbers. Let the normalised and orthogonal function φ_{jm} describe the state of a particle or boson with spin or angular momentum j and projection m . In the language of quantum mechanics this means that φ_{jm} is an eigenfunction of the operators \mathbf{J}^2 and \mathbf{J}_z .

Starting from the functions $\varphi_{j_a m_a}$ and $\varphi_{j_b m_b}$ which belong to different objects a joint state $\Phi_{JM}(j_a, j_b)$ can be made up which is also an eigenfunction of the operators \mathbf{J}^2 and \mathbf{J}_z . Its total spin is characterized by J with the projection M . The new state reads

$$\Phi_{JM}(j_a, j_b) = \sum_{m_a m_b} (j_a m_a j_b m_b | J M) \varphi_{j_a m_a} \varphi_{j_b m_b} \quad (\text{A1.1})$$

$$\text{with } m_a + m_b = M \text{ and } |j_a - j_b| \leq J \leq j_a + j_b. \quad (\text{A1.2})$$

The factors $(j_a m_a j_b m_b | J M)$ are named Clebsch-Gordan coefficients or vector coupling coefficients. They are real and vanish if the conditions (A1.2) are violated. The coupling (A1.1) is frequently written as

$$[\varphi_{j_a m_a} \times \varphi_{j_b m_b}]^{(J)}_M \equiv \Phi_{JM}(j_a, j_b). \quad (\text{A1.3})$$

Clebsch-Gordan coefficients meet the following symmetry conditions

$$(j_a m_a j_b m_b | J M) = (-1)^{j_a + j_b - J} (j_b m_b j_a m_a | J M), \quad (\text{A1.4})$$

$$(j_a m_a j_b m_b | J M) = (-1)^{j_a + j_b - J} (j_a, -m_a, j_b, -m_b | J, -M), \quad (\text{A1.5})$$

$$(j_a m_a j_b m_b | J M) = (-1)^{j_a - m_a} ((2J + 1)/(2j_b + 1))^{1/2} (j_a m_a J, -M | j_b, -m_b). \quad (\text{A1.6})$$

For $j_a = j_b = j$ specially we can write

$$(j m j m' | J M) = (-1)^{2j - J} (j m' j m | J M). \quad (\text{A1.7})$$

If additionally $m = m'$ holds we have

$$(j m j m | J M) = (-1)^{2j - J} (j m j m | J M). \quad (\text{A1.8})$$

That is to say $(j m j m | J M) = 0$, if $2j - J$ is odd.

We demand that state functions such as $\Phi_{JM}(j_a, j_b)$ and $\Phi_{J'M'}(j_a, j_b)$ are normalised and orthogonal i. e.

$$\int \Phi_{JM}^*(j_a, j_b) \Phi_{J'M'}(j_a, j_b) d\sigma = \delta_{JJ'} \delta_{MM'}. \quad (\text{A1.9})$$

In (A1.9) the integration comprises all variables of the functions φ_{jm} . By inserting the relation (A1.1) and by means of the given orthonormality one obtains

$$\sum_{m_a m_b} (j_a m_a j_b m_b | J M) (j_a m_a j_b m_b | J' M') = \delta_{JJ'} \delta_{MM'}. \quad (\text{A1.10})$$

Making use of (A1.4) up to (A1.6) one obtains

$$\sum_{m_b M} (j_a m_a j_b m_b | J M) (j_a' m_a' j_b m_b | J M) = (2J + 1)(2j_a + 1)^{-1} \delta_{j_a j_a'} \delta_{m_a m_a'}. \quad (\text{A1.11})$$

Multiplying both sides of (A1.1) by $\varphi_{j_a m_a}^* \varphi_{j_b m_b}^*$ and integrating yield the relation

$$\int \varphi_{j_a m_a}^* \varphi_{j_b m_b}^* \Phi_{JM}(j_a, j_b) d\sigma = (j_a m_a' j_b m_b' | J M). \quad (\text{A1.12})$$

Because of the linearity of (A1.1) one expects an inverse transformation with real coefficients. We write it with conjugate complex functions like this

$$\varphi_{j_a m_a}^* \varphi_{j_b m_b}^* = \sum_{J' M'} C(j_a m_a j_b m_b J' M') \Phi_{J'M'}^*(j_a, j_b).$$

Multiplying by $\Phi_{JM}(j_a, j_b)$ and integrating one obtains with the aid of (A1.12)

$$C(j_a m_a j_b m_b J M) = \int \varphi_{j_a m_a}^* \varphi_{j_b m_b}^* \Phi_{JM}(j_a, j_b) d\sigma = (j_a m_a j_b m_b | J M). \quad (\text{A1.13})$$

Thus, we have

$$\varphi_{j_a m_a} \varphi_{j_b m_b} = \sum_{JM} (j_a m_a j_b m_b | J M) \Phi_{JM}(j_a, j_b) \quad (\text{A1.14})$$

as the inverse transformation to (A1.1). Multiplying (A1.14) by the expression $\varphi_{j_a m_a}^* \varphi_{j_b m_b}^* = \sum_{J' M'} (j_a m_a' j_b m_b' | J' M') \Phi_{J'M'}^*(j_a, j_b)$ and integrating one obtains the relation

$$\sum_{JM} (j_a m_a j_b m_b | J M) (j_a m_a' j_b m_b' | J M) = \delta_{m_a m_a'} \delta_{m_b m_b'}, \quad (\text{A1.15})$$

in analogy with (A1.10). We give the explicit expressions of frequently employed Clebsch-Gordan coefficients

$$\begin{aligned} (j_a m_a j_b m_b | 0 0) &= (-1)^{j_a - m_a} (2j_a + 1)^{-1/2} \delta_{j_a j_b} \delta_{m_a, -m_b}, \\ (2 m 2, -m | 1 0) &= (-1)^m m / \sqrt{10}, \end{aligned} \quad (\text{A1.16})$$

$$(2, m - 1, 2, -m | 1, -1) = (-1)^m \sqrt{(2 + m)(3 - m)} / \sqrt{20},$$

$$(2 0 2 m | 2 m) = (m^2 - 2) / \sqrt{14},$$

$$(j_a, m_a = j_a, j_b, m_b = j_b | J = j_a + j_b, M = j_a + j_b) = +1. \quad (\text{A1.17})$$

The last coefficient shows that the product of two states $|j, m = j\rangle$ is the stretched joint state lying in the z-direction. The choice of the positive sign on the right hand side of (A1.17) is arbitrary.

The symmetry properties of the Clebsch-Gordan coefficients are made visible especially by means of the 3-j symbol $\begin{pmatrix} j_a & j_b & J \\ m_a & m_b & -M \end{pmatrix}$, which is defined as follows

$$(j_a m_a j_b m_b | J M) = (-1)^{j_a - j_b + M} \sqrt{(2J+1)} (j_a m_a j_b m_b | J -M). \quad (A1.18)$$

The value of the 3- j symbol does not alter when its columns – here they are arranged at an angle – undergo an even number of interchanges. If this number is odd or if all projections m_a , m_b and $-M$ change their signs simultaneously, the new 3- j symbol differs from the original one in the phase factor $(-1)^{j_a + j_b - J}$. Edmonds (1964) and (1996) has given algebraic expressions for some 3- j symbols. Numerical tables for 3- j symbols were published by Rotenberg, Bivins et al. (1959) and Brussaard and Glaudemans (1977).

We come back to the statement (A1.1) and go into the angular momentum operators. If the spin operators $\mathbf{J}_{za} \equiv \mathbf{J}_{0a}$ and $\mathbf{J}_{zb} \equiv \mathbf{J}_{0b}$ act on the states $\varphi_{j_a m_a}$ and $\varphi_{j_b m_b}$ according to

$$\mathbf{J}_{0a} \varphi_{j_a m_a} = m_a \varphi_{j_a m_a}, \quad \mathbf{J}_{0b} \varphi_{j_b m_b} = m_b \varphi_{j_b m_b} \text{ and} \quad (A1.19)$$

$$\mathbf{J}_{0a} \varphi_{j_b m_b} = \varphi_{j_b m_b} \mathbf{J}_{0a}, \quad \mathbf{J}_{0b} \varphi_{j_a m_a} = \varphi_{j_a m_a} \mathbf{J}_{0b} \quad (A1.20)$$

the sum operator

$$\mathbf{J}_0 = \mathbf{J}_{0a} + \mathbf{J}_{0b} \quad (A1.21)$$

satisfies the following eigenvalue equation with the joint expression (A1.1) or (A1.3)

$$\mathbf{J}_0 \Phi_{JM}(j_a, j_b) \equiv \mathbf{J}_0 [\varphi_{j_a m_a} \times \varphi_{j_b m_b}]^{(J)}_M = M [\varphi_{j_a m_a} \times \varphi_{j_b m_b}]^{(J)}_M \quad (A1.22)$$

which can be seen with the help of (A1.1), (A1.19) and (A1.20). The other “ladder” components of \mathbf{J} read according to (8.6) as follows $\mathbf{J}_1 = -(\mathbf{J}_x + i\mathbf{J}_y)/\sqrt{2}$ and $\mathbf{J}_{-1} = (\mathbf{J}_x - i\mathbf{J}_y)/\sqrt{2}$. They meet the condition (13.26)

$$\mathbf{J}_{\pm 1, a} \varphi_{j_a m_a} = -(\pm)((j_a(j_a + 1) - m_a(m_a \pm 1))/2)^{1/2} \varphi_{j_a, m_a \pm 1},$$

which holds analogously for $\mathbf{J}_{\pm 1, b} \varphi_{j_b m_b}$. In analogy with (A1.21) we make up the operator

$$\mathbf{J}_{\pm 1} = \mathbf{J}_{\pm 1, a} + \mathbf{J}_{\pm 1, b} \quad (A1.23)$$

and expect that the similar relation

$$\begin{aligned} \mathbf{J}_{\pm 1} [\varphi_{j_a m_a} \times \varphi_{j_b m_b}]^{(J)}_M = \\ -(\pm)((J(J+1) - M(M \pm 1))/2)^{1/2} [\varphi_{j_a m_a} \times \varphi_{j_b m_b}]^{(J)}_{M \pm 1} \end{aligned} \quad (A1.24)$$

will hold. We write it in detail as follows

$$\begin{aligned} \sum_{m_a m_b} (j_a m_a j_b m_b | J M) \{ (j_a(j_a + 1) - m_a(m_a \pm 1))^{1/2} \varphi_{j_a, m_a \pm 1} \varphi_{j_b m_b} + \\ (j_b(j_b + 1) - m_b(m_b \pm 1))^{1/2} \varphi_{j_a m_a} \varphi_{j_b, m_b \pm 1} \} = \\ (J(J+1) - M(M \pm 1))^{1/2} [\varphi_{j_a m_a} \times \varphi_{j_b m_b}]^{(J)}_{M \pm 1}. \end{aligned} \quad (A1.25)$$

The relation (A1.25) cannot be explained solely by the properties of the Clebsch-Gordan coefficients put together in this appendix. In contrary, they constitute a special condition for these coefficients. Making use of the orthonormality of the state functions, from (A1.25) one obtains recursion formulas for the Clebsch-Gordan coefficients. They serve to calculate them numerically.

In case the operator of the total spin, \mathbf{J}^2 (8.8), acts on a coupled state, from (A1.22), (A1.24) and (8.8) follows the well-known relation

$$\begin{aligned} \mathbf{J}^2 [\varphi_{j_1 m_1} \times \varphi_{j_2 m_2}]^{(J)}_M &= (\mathbf{J}_0^2 - \mathbf{J}_0 - 2\mathbf{J}_1 \mathbf{J}_2) [\varphi_{j_1 m_1} \times \varphi_{j_2 m_2}]^{(J)}_M = \\ J(J+1) [\varphi_{j_1 m_1} \times \varphi_{j_2 m_2}]^{(J)}_M. \end{aligned} \quad (\text{A1.26})$$

A2 Symmetry properties of coupled spin states of identical objects

In this section the selection rules for the total spin of a coupled state of identical objects - fermions or bosons - will be looked into. According to (A1.1) such a coupling reads as follows

$$\Phi_{JM} j, 1, 2) = \sum_{m, m'} (j, m, j, m' | J, M) \varphi_{jm}(1) \varphi_{jm'}(2). \quad (\text{A2.1})$$

In (A2.1) provisionally we are considering these objects as distinguishable and label them with numbers. We now interchange both objects - that is to say all their variables - and obtain the following joint state

$$\Phi_{JM} j, 2, 1) = \sum_{m, m'} (j, m, j, m' | J, M) \varphi_{jm}(2) \varphi_{jm'}(1).$$

With (A1.7) we have

$$\begin{aligned} \Phi_{JM} j, 2, 1) &= (-1)^{2j-J} \sum_{m, m'} (j, m', j, m | J, M) \varphi_{jm'}(1) \varphi_{jm}(2) = \\ &(-1)^{2j-J} \Phi_{JM} j, 1, 2). \end{aligned} \quad (\text{A2.2})$$

First, we apply (A2.2) to fermions - for example to nucleons. They obey the Pauli principle i. e. the function of the whole state changes its sign if both fermions are interchanged as follows

$$\Phi_{JM} j, 2, 1) = -\Phi_{JM} j, 1, 2) \quad \text{i. e.} \quad (\text{A2.3})$$

$$(-1)^{2j-J} = -1. \quad (\text{A2.4})$$

Fermions have half-integer spin j , that is why $2j$ is odd. Consequently, the equations (A2.3) and (A2.4) are satisfied if J is even. That is to say, a pair of identical fermions can be represented by (A2.1) if J is even.

On the other hand, bosons behave symmetrically with respect to interchanging i. e. for a coupled pair of identical bosons the following relation holds

$$\Phi_{JM} j, 2, 1) = +\Phi_{JM} j, 1, 2) \quad \text{or} \quad (\text{A2.5})$$

$$(-1)^{2j-J} = +1. \quad (\text{A2.6})$$

Because here j is integer the total spin (or angular momentum) J must also be even. Specially for $j = 2$ the quantity J can amount to

$$J = 0, 2 \text{ or } 4. \quad (\text{A2.7})$$

A3 Racah-coefficients and 6- j symbols

Three state functions with spins can be coupled by combining the first and the second function and by adding the third one to the joint function. An other way to do it consists in coupling the second and the third function and in adding the result to the first one. By means of the Racah coefficients, it is possible to represent one configuration by linear combinations of the other type.

Three eigenfunctions of the spin - or angular momentum - operators \mathbf{J}^2 and \mathbf{J}_z are given and coupled in the following way (A1.1, A1.3)

$$[[\varphi_{ja} \times \varphi_{jb}]^{(J_{ab})} \times \varphi_{jc}]^{(J)}_M = \quad (\text{A3.1})$$

$$\sum_{m_a m_b M_{ab} m_c} (j_a m_a j_b m_b | J_{ab} M_{ab}) (J_{ab} M_{ab} j_c m_c | J M) \varphi_{ja m_a} \varphi_{jb m_b} \varphi_{jc m_c}.$$

With the help of (A1.14), we recouple the triple of functions on the right hand side in (A3.1) this way

$$[[\varphi_{ja} \times \varphi_{jb}]^{(J_{ab})} \times \varphi_{jc}]^{(J)}_M = \sum_{m_a m_b M_{ab} m_c} (j_a m_a j_b m_b | J_{ab} M_{ab}) (J_{ab} M_{ab} j_c m_c | J M) \cdot \quad (\text{A3.2})$$

$$\sum_{J_{bc} M_{bc}} (j_b m_b j_c m_c | J_{bc} M_{bc}) \cdot$$

$$\sum_{J' M'} (j_a m_a J_{bc} M_{bc} | J' M') [\varphi_{ja} \times [\varphi_{jb} \times \varphi_{jc}]^{(J_{bc})}]^{(J')}_{M'} \delta_{JJ'} \delta_{MM'}.$$

The symbol $\delta_{JJ'} \delta_{MM'}$ takes into account that both sides of (A3.2) are eigenfunctions of \mathbf{J}^2 and \mathbf{J}_z with the same corresponding eigenvalues. We can write (A3.2) like this

$$[[\varphi_{ja} \times \varphi_{jb}]^{(J_{ab})} \times \varphi_{jc}]^{(J)}_M = \sum_{J_{bc}} U(j_a j_b J j_c, J_{ab} J_{bc}) [\varphi_{ja} \times [\varphi_{jb} \times \varphi_{jc}]^{(J_{bc})}]^{(J)}_M \quad (\text{A3.3})$$

with $U(j_a j_b J j_c; J_{ab} J_{bc}) =$

$$\sum_{m_a m_b} (j_a m_a j_b m_b | J_{ab}, m_a + m_b) (J_{ab}, m_a + m_b, j_c, M - m_a - m_b | J M) \cdot$$

$$(j_b m_b j_c, M - m_a - m_b | J_{bc}, M - m_a) (j_a m_a J_{bc}, M - m_a | J M) \equiv \quad (\text{A3.4})$$

$$\sqrt{(2J_{ab} + 1)} \sqrt{(2J_{bc} + 1)} W(j_a j_b J j_c; J_{ab} J_{bc}).$$

The equation (A3.3) reveals that the coupling of three spin eigenfunctions according to (A3.1) can be represented as a linear combination of couplings where the second and the third functions are combined first. The factors U (A3.4) are named normalised Racah-coefficients. Apart from two factors they agree with the common Racah-coefficients W (A3.4). The following inverse relation can be shown

$$[\varphi_{j_a} \times [\varphi_{j_b} \times \varphi_{j_c}]^{(J_{bc})}]^{(J)}_M = \sum_{J_{ab}} U(j_a j_b J j_c, J_{ab} J_{bc}) [[\varphi_{j_a} \times \varphi_{j_b}]^{(J_{ab})} \times \varphi_{j_c}]^{(J)}_M. \quad (A3.5)$$

In order to visualise the symmetry properties of the Racah-coefficients, the 6- j symbol has been created as follows

$$\{j_a j_b j_c\} = (-1)^{j_a - j_b - l_a - l_b} (2j_c + 1)^{-1/2} (2l_c + 1)^{-1/2} U(j_a j_b l_b l_a; j_c l_c) \equiv (-1)^{j_a - j_b - l_a - l_b} W(j_a j_b l_b l_a; j_c l_c). \quad (A3.6)$$

The value of the 6- j symbol is unaltered when the columns - here they are arranged at an angle - are interchanged. It remains also the same when two columns are turned upside down. The following triples of quantities in the 6- j symbol have to form a triangle

$$(j_a j_b j_c), (j_a l_b l_c), (l_a l_b j_c) \text{ and } (l_a j_b l_c)$$

which is indicated by lines in the symbols as follows

$$\{j_a j_b j_c\}, \{j_a j_b l_c\}, \{j_a l_b j_c\}, \{j_a l_b l_c\}, \{l_a j_b j_c\}, \{l_a j_b l_c\}. \quad (A3.7)$$

From the orthonormality of the expression (A3.3), follows this orthogonality relation

$$\sum_{j_c} (2j_c + 1) \{j_a j_b j_c\} \{j_a j_b j_c\} = \delta_{j_c j_c} / (2l_c + 1). \quad (A3.8)$$

A similar relation reads like this

$$\sum_{l_c} (-1)^{l_c} (2l_c + 1) \{j_a j_b j_c\}^2 = \{j_a j_b j_c\} \text{ for } j_c. \quad (A3.9)$$

With the orthonormality of the Clebsch-Gordan coefficients the equations (A3.4) and (A3.6) can be brought in the following form

$$(-1)^{-l_a + l_b + n_c} \sqrt{(2l_a + 1)(2l_b + 1)} (l_a n_a l_b n_b | l_c n_c) \{j_a j_b j_c\} = \sum_{m_a m_b m_c} (-1)^{j_a + j_b + j_c + m_a + m_b + m_c} (j_a m_a j_b, -m_b | l_c, -n_c) (j_b m_b j_c, -m_c | l_a, -n_a) (j_c m_c j_a, -m_a | l_b, -n_b). \quad (A3.10)$$

Furthermore the relation

$$\sum_j (2j + 1) \{j_1 j_2 j_j\} = (-1)^{2(j_1 + j_2)}. \quad (A3.11)$$

holds. Some 6- j symbols are simple algebraic expressions. If l_c is zero, according to the triangle conditions (A3.7) we have $l_b = j_a$ and $l_a = j_b$. Then the 6- j symbol has the form

$$\{j_a j_b j_0\} = (-1)^{j_a + j_b + j_c} (2j_a + 1)^{-1} (2j_c + 1)^{-1}. \quad (A3.12)$$

The zero can be brought to any position by using the interchanging rules mentioned above. We refer to a further special 6- j symbol (Edmonds 1964, p. 154)

$$\{j_a j_b j_c\} = (-1)^{j_a+j_b+j_c+1} [j_a(j_a+1) + j_b(j_b+1) - j_c(j_c+1)] \cdot [4j_a(j_a+1)(2j_a+1)j_b(j_b+1)(2j_b+1)]^{-1/2}. \quad (\text{A3.13})$$

In the following table, numerical expressions of some 6- j symbols are put together (Rotenberg, Bivins et al. 1959), which appear in the IBM

J	J'	$\{ \begin{smallmatrix} 2 & 2 & J \\ 2 & 2 & J' \end{smallmatrix} \}$	
2	0	1/5	
2	1	-1/10	
2	2	-3/70	
2	3	4/35	
2	4	2/35	(A3.14)
4	0	1/5	
4	1	2/15	
4	2	2/35	
4	3	1/70	
4	4	1/630.	

A4 The 9- j symbol

If four spin eigenfunctions are coupled in pairs which are coupled additionally like this

$$[(\varphi_{j_a} \times \varphi_{j_b})]^{(J_{ab})} \times [\varphi_{j'} \times \varphi_j]^{(J')}]^{(J)}_M,$$

this configuration can be represented by means of equivalent expressions where φ_{j_b} and $\varphi_{j'}$ are interchanged. After several steps one obtains the following result

$$[(\varphi_{j_a} \times \varphi_{j_b})]^{(J_{ab})} \times [\varphi_{j'} \times \varphi_j]^{(J')}]^{(J)}_M = \sum_{J_a J_b} [(2J_{ab}+1)(2J'+1)(2J_a+1)(2J_b+1)]^{1/2}. \quad (\text{A4.1})$$

$$\{j_a j_{J_a} j_b j_{J_b} j_{J'} j_{J''}\} [(\varphi_{j_a} \times \varphi_{j_b})]^{(J_a)} \times [(\varphi_{j_b} \times \varphi_{j'})]^{(J_b)}]^{(J)}_M.$$

The matrix in braces is the 9- j symbol. It shows certain symmetries. An even permutation of its columns - here they are arranged at an angle - or of its rows leaves its value unaffected. An odd permutation of rows or columns introduces the phase factor $(-1)^s$, where s is equal to the sum of all arguments in the 9- j symbol. In every column and row, the triangle condition is satisfied.

If the 9- j symbol contains the argument $J = 0$, the relations $J_{ab} = J'$ and $J_a = J_b = J''$ hold. In this case the 6- j symbol reads

$$\{j_a j_{J''} j_b j_{J''} j_{J'} j_{J''}\} = (-1)^{j_b+j'+j''+J''} [(2J'+1)(2J''+1)]^{-1/2} \{j_a j_{J'} j_b j_{J''} j_{J'} j_{J''}\}. \quad (\text{A4.2})$$

Now we replace the functions φ in (A4.1) by operators \mathbf{b} and write the last brackets in detail with the help of (A1.1) and (A1.3). We leave the operators \mathbf{b} in the original sequence as follows

$$\begin{aligned} & [[\mathbf{b}_{j_a} \times \mathbf{b}_{j_b}]]^{(J_{ab})} \times [\mathbf{b}_j \times \mathbf{b}_{j'}]^{(J')} \bigg|^{(J)}_M = \\ & \sum_{J_a J_b} [(2J_{ab} + 1)(2J' + 1)(2J_a + 1)(2J_b + 1)]^{1/2} \cdot \{ \begin{matrix} j_a & j_b & J_{ab} \\ J_a & J_b & J \end{matrix} \}. \quad (\text{A4.3}) \\ & \sum_{m_a m_b m m' M_a M_b} (j_a m_a j m | J_a M_a) (j_b m_b j' m' | J_b M_b) (J_a M_a J_b M_b | J M) \cdot \\ & \mathbf{b}_{j_a m_a} \mathbf{b}_{j_b m_b} \mathbf{b}_{j m} \mathbf{b}_{j' m'}. \end{aligned}$$

If the operators \mathbf{b} commute, the equation (A4.3) can be transformed to the form (A4.1), otherwise this conversion is not possible.

A5 The Wigner-Eckart theorem

The IBM deals with matrix elements of coupled \mathbf{s} - and/or \mathbf{d} -operators (chapter 11 and 12). The Wigner-Eckart theorem makes possible to split up these matrix elements in a Clebsch-Gordan coefficient and an element which is independent on projection quantum numbers of the state functions and of the operators.

We start from the properties of the spherical harmonics $Y_{lm}(\varrho)$ depending on the direction vector $\varrho = \underline{r}/r$. They are eigenfunctions of the angular momentum operators \mathbf{J}^2 and \mathbf{J}_z and therefore they represent a part of the wave functions of particles in a central field. They are orthonormalised i. e.

$$\int Y_{lm}^*(\varrho) Y_{l'm'}(\varrho) d\varrho \equiv \int Y_{lm}^*(\vartheta, \varphi) Y_{l'm'}(\vartheta, \varphi) \sin \vartheta d\vartheta d\varphi = \delta_{ll'} \delta_{mm'}. \quad (\text{A5.1})$$

Spherical harmonics have the following properties

$$Y_{lm}(\varrho) = (-1)^m Y_{l,-m}^*(\varrho) \quad (\text{A5.2})$$

and $Y_{l_a m_a}(\varrho) Y_{l_b m_b}(\varrho) = \sum_{LM} [(2l_a + 1)(2l_b + 1)]^{1/2} [4\pi(2L + 1)]^{-1/2} \cdot$

$$(l_a m_a l_b m_b | L, -M) (l_a 0 l_b 0 | L 0) (-1)^{-M} Y_{LM}^*(\varrho). \quad (\text{A5.3})$$

Now we make up the following integral

$$\begin{aligned} & \int Y_{l_a m_a}^*(\varrho) Y_{LM}(\varrho) Y_{l_b m_b}(\varrho) d\varrho = (-1)^{m_a} \int Y_{LM}(\varrho) Y_{l_a, -m_a}(\varrho) Y_{l_b m_b}(\varrho) d\varrho = \\ & \sum_{L'M'} [(2l_a + 1)(2l_b + 1)]^{1/2} [4\pi(2L' + 1)]^{-1/2} \cdot \\ & (l_a, -m_a l_b m_b | L', -M') (l_a 0 l_b 0 | L' 0) (-1)^{-M'+m_a} \int Y_{L'M'}^*(\varrho) Y_{LM}(\varrho) d\varrho = \\ & (-1)^{-M'+m_a} (l_a, -m_a l_b m_b | L, -M) [(2l_a + 1)(2l_b + 1)]^{1/2} [4\pi(2L + 1)]^{-1/2} \cdot \\ & (l_a 0 l_b 0 | L 0). \end{aligned} \quad (\text{A5.4})$$

Doing this, (A5.1) and (A5.3) have been used. Making use of (A1.6) we obtain

$$\int Y_{l_a m_a}^*(\varrho) Y_{LM}(\varrho) Y_{l_b m_b}(\varrho) d\varrho = \quad (A5.5)$$

$$(-1)^{2L} (l_b m_b L M | l_a m_a) (2l_a + 1)^{-1/2} (-1)^{l_a + 2l_b + L} [(2l_a + 1)(2l_b + 1)/4\pi]^{1/2} (l_a 0 l_b 0 | L 0).$$

We write (A5.5) in the following form

$$\int Y_{l_a m_a}^*(\varrho) Y_{LM}(\varrho) Y_{l_b m_b}(\varrho) d\varrho \equiv \quad (A5.6)$$

$$\langle Y_{l_a m_a} | Y_{LM} | Y_{l_b m_b} \rangle = (-1)^{2L} (l_b m_b L M | l_a m_a) (2l_a + 1)^{-1/2} \langle Y_{l_a} || Y_L || Y_{l_b} \rangle.$$

The last quantity is named reduced matrix element. It is independent on the projection quantum numbers m_a , M and m_b and it reads

$$\langle Y_{l_a} || Y_L || Y_{l_b} \rangle = (-1)^{l_a + 2l_b + L} [(2l_a + 1)(2l_b + 1)/4\pi]^{1/2} (l_a 0 l_b 0 | L 0). \quad (A5.7)$$

The equations (A5.6) can be generalised. One can replace $Y_{l_a m_a}$ and $Y_{l_b m_b}$ by the eigenfunctions $| l_a m_a \rangle$ and $| l_b m_b \rangle$ of \mathbf{J}^2 and \mathbf{J}_z with a common potential. Instead of the function Y_{LM} , which is placed in the middle of the original matrix element (A5.5), one can put a tensor operator $\mathbf{T}^{(L)}_M$ of the rank L . These operators and spherical harmonics behave equally when the co-ordinate system rotates. After a rotation the new spherical harmonics are linear combinations of spherical harmonics belonging to the old system with the same rank L . For a given rotation, the corresponding coefficients for spherical harmonics and tensor operators are identical.

As an equivalent statement, one can say that tensor operators can be coupled to an operator with a defined angular momentum with the same procedure as the spherical harmonics i. e. by making use of Clebsch-Gordan coefficients. Examples for tensor operators are the angular momentum operator \mathbf{J}_i , the nabla operator ∇ , creation operators for bosons \mathbf{d}^+ , \mathbf{s}^+ (see the comment on (6.9)) and the corresponding modified annihilation operators \mathbf{d}^- and \mathbf{s} (see the comment on (6.19) and (6.20)).

One can show that (A5.6) remains valid if all three functions are replaced in the way just mentioned like this

$$\langle l_a m_a | \mathbf{T}^{(L)}_M | l_b m_b \rangle = (-1)^{2L} (l_b m_b L M | l_a m_a) (2l_a + 1)^{-1/2} \langle l_a || \mathbf{T}^{(L)} || l_b \rangle. \quad (A5.8)$$

This is the Wigner-Eckart theorem. It enables to split the matrix elements of a tensor operator in a reduced matrix element, which does not depend on the projection quantum numbers m and in a Clebsch-Gordan coefficient. It offers an essential simplification because as soon as a matrix element with a set of m -values is known it can be given for other sets of this kind. One obtains the reduced matrix element (on the right hand side of (A5.8)) by calculating the left side for one set of m -values and by remodelling.

Reduced matrix elements are also necessary for constituting the so-called coefficients of fractional parentage.

A6 A further multipole representation of the Hamiltonian

The multipole representation of the Hamilton operator according to (9.37) stands out due to an especially consequent choice of operators which permit to constitute the three special cases of the IBM in a direct manner. Frequently, in the literature the multipole representation appears, which was formulated originally by Iachello and Arima. In their review article, these authors use it again in 1987. This representation was built in the computer code PHINT (appendix 7) by O. Scholten in a simplified form.

This multipole representation contains the operators n_d and J^2 (9.1), which are already present in the Hamiltonian in (9.37). Instead of Q^2 (9.4) the more general form $Q^2(\chi)$ (12.19, 15.17) is employed. Its components read

$$Q_\mu(\chi) = d^+_\mu s + s^+ d^-_\mu + \chi [d^+ \times d^-]^{(2)}_\mu. \quad (A6.1)$$

Furthermore the operators $[[d^+ \times d^-]^{(J)} \times [d^+ \times d^-]^{(J)}]^{(0)}$ with $J = 3, 4$ occur, which come from the transformation of the term $[[d^+ \times d^-]^{(J)} \times (d^- \times d^-)^{(J)}]^{(0)}$ in (7.17).

In concrete terms, we make the claim that the Hamilton operator can be written as follows

$$H = \varepsilon' n_d + \frac{1}{2} \eta J^2 + \frac{1}{2} \kappa Q^2(\chi) - 5\sqrt{7} \omega [[d^+ \times d^-]^{(3)} \times (d^+ \times d^-)^{(3)}]^{(0)} + 5.3 \xi [[d^+ \times d^-]^{(4)} \times (d^+ \times d^-)^{(4)}]^{(0)}. \quad (A6.2)$$

Now we are working on the individual terms in (A6.2) in order to transform this equation in the basic form (7.17). With the help of (9.28) and (9.34) we can write $Q^2(\chi)$ in the following form

$$\begin{aligned} Q^2(\chi)/\sqrt{5} &= [d^+ \times d^-]^{(0)} s s + s^+ s^+ [d^- \times d^-]^{(0)} + \\ & s s^+ [d^+ \times d^-]^{(0)} + [d^- \times d^-]^{(0)} s^+ s + \chi^2 [[d^+ \times d^-]^{(2)} \times [d^+ \times d^-]^{(2)}]^{(0)} + \\ & 2 \chi ([[d^+ \times d^-]^{(2)} \times d^- s]^{(0)} + [s^+ d^+ \times [d^- \times d^-]^{(2)}]^{(0)}). \end{aligned} \quad (A6.3)$$

For the time being we are dealing with the second line in (A6.3). According to the commutation rules for d -operators we have

$$\begin{aligned} [d^- \times d^-]^{(0)} &= 5^{-1/2} \sum_{m=-1}^{+2} d^-_m d^+_{-m} (-1)^{-m} = 5^{-1/2} \sum_m d^-_m d^+_{-m} = \\ & 5/\sqrt{5} + 5^{-1/2} \sum_m d^+_{-m} d^-_m = \sqrt{5} + [d^+ \times d^-]^{(0)}. \end{aligned} \quad (A6.4)$$

Because of $s s^+ = 1 + s^+ s$ this line results in

$$\begin{aligned} & s s^+ [d^+ \times d^-]^{(0)} + [d^- \times d^-]^{(0)} s^+ s = \\ & [d^+ \times d^-]^{(0)} + s^+ s [d^+ \times d^-]^{(0)} + [d^+ \times d^-]^{(0)} s^+ s + \sqrt{5} s^+ s = \\ & n_d/\sqrt{5} + \sqrt{5}(N - n_d) + 2[d^+ s^+ \times d^- s]^{(0)}. \end{aligned} \quad (A6.5)$$

In order to handle the last expression in (A6.2) we need the relation (9.8)

$$\begin{aligned} & [[\mathbf{d}^+ \times \mathbf{d}^+]^{(J)} \times (\mathbf{d}^- \times \mathbf{d}^-)^{(J)}]^{(0)} = \\ & \sum_{J'} (-1)^{J'} \sqrt{(2J+1)(2J'+1)} \left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J' \end{matrix} \right\} [[\mathbf{d}^+ \times \mathbf{d}^+]^{(J')} \times [\mathbf{d}^- \times \mathbf{d}^-]^{(J')}]^{(0)} - \\ & (1/5) \sqrt{(2J+1)} \mathbf{n}_d. \end{aligned} \quad (\text{A6.6})$$

We multiply both sides of (A6.6) by $\sqrt{(2J+1)} \left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J'' \end{matrix} \right\}$, sum over J and obtain

$$\begin{aligned} & \sum_J \sqrt{(2J+1)} \left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J'' \end{matrix} \right\} [[\mathbf{d}^+ \times \mathbf{d}^+]^{(J)} \times (\mathbf{d}^- \times \mathbf{d}^-)^{(J)}]^{(0)} = \\ & \sum_{J'} (-1)^{J'} \sqrt{(2J'+1)} \langle \sum_J (2J+1) \left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J'' \end{matrix} \right\} \left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J'' \end{matrix} \right\} \rangle [[\mathbf{d}^+ \times \mathbf{d}^+]^{(J')} \times [\mathbf{d}^- \times \mathbf{d}^-]^{(J')}]^{(0)} \\ & - (1/5) \langle \sum_J (2J+1) \left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J'' \end{matrix} \right\} \rangle \mathbf{n}_d. \end{aligned} \quad (\text{A6.7})$$

The expressions in pointed brackets in (A6.7) read according to (A3.8) and (A3.11) like this

$$\sum_J (2J+1) \left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J'' \end{matrix} \right\} \left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J'' \end{matrix} \right\} = \delta_{J''J'} / (2J''+1)$$

$$\text{and } \sum_J (2J+1) \left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J'' \end{matrix} \right\} = 1.$$

Thus we obtain the inverse relation of (A6.6)

$$\begin{aligned} & [[\mathbf{d}^+ \times \mathbf{d}^+]^{(J')} \times [\mathbf{d}^- \times \mathbf{d}^-]^{(J')}]^{(0)} = \\ & (-1)^{J'} \sqrt{(2J'+1)} \sum_{J=0,2,4} \sqrt{(2J+1)} \left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J' \end{matrix} \right\} [[\mathbf{d}^+ \times \mathbf{d}^+]^{(J)} \times [\mathbf{d}^- \times \mathbf{d}^-]^{(J)}]^{(0)} + \\ & (-1)^{J'} (1/5) \sqrt{(2J'+1)} \mathbf{n}_d. \end{aligned} \quad (\text{A6.8})$$

According to (6.6) the summands with $J = 1, 3$ vanish. We now insert the expressions (9.1) and (A6.3) in (A6.2) and make use of (A6.5) and (A6.8). By means of the 6- j symbol $\left\{ \begin{matrix} 2 & 2 & J \\ 2 & 2 & J' \end{matrix} \right\}$ according to (A3.14) we obtain

$$\begin{aligned} \mathbf{H} = & (\varepsilon' + 3\eta + (1/2)(\chi^2 - 4)\kappa + 7\omega + 9\xi) \mathbf{n}_d + \\ & (-3\eta + (1/2)\chi^2\kappa - \omega + 9\xi) [[\mathbf{d}^+ \times \mathbf{d}^+]^{(0)} \times [\mathbf{d}^- \times \mathbf{d}^-]^{(0)}]^{(0)} + \\ & \sqrt{5} (-(3/2)\eta - (3/28)\chi^2\kappa + 4\omega + (18/7)\xi) [[\mathbf{d}^+ \times \mathbf{d}^+]^{(2)} \times [\mathbf{d}^- \times \mathbf{d}^-]^{(2)}]^{(0)} + \\ & 3(2\eta + (1/7)\chi^2\kappa + (1/2)\omega + (1/14)\xi) [[\mathbf{d}^+ \times \mathbf{d}^+]^{(4)} \times [\mathbf{d}^- \times \mathbf{d}^-]^{(4)}]^{(0)} + \\ & \sqrt{5}\chi\kappa ([[\mathbf{d}^+ \times \mathbf{d}^+]^{(2)} \times \mathbf{d}^- \mathbf{s}]^{(0)} + [\mathbf{s}^+ \mathbf{d}^+ \times [\mathbf{d}^- \times \mathbf{d}^-]^{(2)}]^{(0)}) + \\ & \sqrt{5}(1/2)\kappa ([[\mathbf{d}^+ \times \mathbf{d}^+]^{(0)} \mathbf{s} \mathbf{s} + \mathbf{s}^+ \mathbf{s}^+ [\mathbf{d}^- \times \mathbf{d}^-]^{(0)}) + \\ & \sqrt{5}\kappa [\mathbf{d}^+ \mathbf{s}^+ \times \mathbf{d}^- \mathbf{s}]^{(0)}. \end{aligned} \quad (\text{A6.9})$$

The comparison of (A6.9) with (7.17) yields

$$\begin{aligned}
\varepsilon_s &= 0, \\
\varepsilon_d - \varepsilon_s &= \varepsilon' + 3\eta + (1/2)(\chi^2 - 4)\kappa + 7\omega + 9\xi, \\
c_0 &= -6\eta + \chi^2\kappa - 14\omega + 18\xi, \\
c_2 &= -3\eta - (3/14)\chi^2\kappa + 8\omega + (36/7)\xi, \\
c_4 &= 4\eta + (2/7)\chi^2\kappa + \omega + (1/7)\xi, \\
v_2/\sqrt{2} &= \sqrt{5}\chi\kappa, \\
v_0/2 &= (1/2)\sqrt{5}\kappa, \\
u_2 &= \sqrt{5}\kappa, \\
u_0 &= 0.
\end{aligned} \tag{A6.10}$$

The ansatz (A6.2) comprises 6 parameters ε' , η , κ , ω , ξ and χ (contained in **Q**). The transformation (A6.10) reveals that the quantities ε_s and u_0 of the basic form (7.17) have been put equal to zero and that v_0 and u_2 are coupled. This reduction to six free parameters is permitted if one looks into the structure of individual nuclei disregarding the binding energies.

A7 The program package PHINT

The code PHINT calculates energies of nuclear states and reduced transition probabilities of the electromagnetic radiation in the interacting boson model. It was written by O. Scholten in FORTRAN, published in Computational Nuclear Physics 1 (O. Scholten, 1991, p.88) and made accessible in the diskette enclosed in that book.

The IBM calculation can be performed in the basic form of the Hamiltonian (7.17) with the program PCIBAXW. The input parameters (in capitals) correspond to the following coefficients in (7.17)

ε_s	0 (is put equal to zero)	
$\varepsilon_d - \varepsilon_s$	H BAR	
c_0, c_2, c_4	(C(1), C(2), C(3))	
$v_2/\sqrt{2}$	F	(A7.1)
$v_0/2$	G	
$u_2/\sqrt{5}$	CH2	
u_0	CH1.	

It is also possible to use the reduced form (7.31). In this case, the following assignments hold

ε	HBAR	
c_0', c_2', c_4'	C(1), C(2), C(3)	
$v_2/\sqrt{2}$	F	(A7.2)
$v_0/2$	G	
	CH2 = CH1 = 0.	

The calculation can be done in the multipole form of the Hamiltonian according to appendix 6, the coefficients of which correspond to the following input parameters

ε'	EPS	
η	ELL	
χ	CHQ/ $\sqrt{5}$	
κ	QQ	(A7.3)
ω	OCT	
ξ	HEX.	

To the technical notes of O. Scholten (1991, p. 103) the following remarks have to be added

- The subroutine RDPAR in the packet PCIBALIB has to be corrected in order to enable it to read the input parameters (C(1), C(2), C(3)) as a triplet (for example (3F10.4) has to be replaced by (3F7.4)).
- Energies should be given in MeV.
- In order to compile the source program no sensitive debug option can be used (for example for the FORTRAN compiler of Microsoft only the command fl and not fl/4Yb must be given).
- Increasing the boson number N to 14 or 16 calls for changes in COMMON(READMAT) and in the corresponding DATA in various places of the programs. Tips in the comments of the program texts have to be followed.

A8 Commutators of operators such as $[\mathbf{d}^+ \times \mathbf{d}^-]^{(J)}_M$

Commutators of the type $[[\mathbf{d}^+ \times \mathbf{d}^-]^{(J')}_{M'}, [\mathbf{d}^+ \times \mathbf{d}^-]^{(J'')}_{M''}]$ are employed in connection with the angular momentum operators (chapter 8) and with Lie algebras in the IBM (chapter 14).

By means of (6.19) and (A1.1) we can write

$$[[\mathbf{d}^+ \times \mathbf{d}^-]^{(J')}_{M'}, [\mathbf{d}^+ \times \mathbf{d}^-]^{(J'')}_{M''}] = \quad (A8.1)$$

$$\sum_{m' n' m'' n''} (2 m' 2 n' | J' M') (2 m'' 2 n'' | J'' M'') (-1)^{n'+m''} [\mathbf{d}^+_{m'} \mathbf{d}_{-n'}, \mathbf{d}^+_{m''} \mathbf{d}_{-n''}].$$

Using the commutation relations (5.36) we obtain

$$[\mathbf{d}^+_{m'} \mathbf{d}_{-n'}, \mathbf{d}^+_{m''} \mathbf{d}_{-n''}] = -\mathbf{d}^+_{m''} \mathbf{d}_{-n'} \delta_{m', n''} + \mathbf{d}^+_{m'} \mathbf{d}_{-n''} \delta_{m'', -n'}. \quad (A8.2)$$

Thus, from (A8.1) and (6.19) the following relation results

$$\begin{aligned}
& [[\mathbf{d}^+ \times \mathbf{d}^-]^{(J')}_{M'}, [\mathbf{d}^+ \times \mathbf{d}^-]^{(J'')}_{M''}] = \\
& \sum_{m' n' m'' n''} (2 m' 2 n' | J' M') (2 m'' 2 n'' | J'' M'') \cdot \\
& ((-1)^{n'} \mathbf{d}^+_{m'} \mathbf{d}^-_{-n'} \delta_{m', -n'} - (-1)^{n''} \mathbf{d}^+_{m''} \mathbf{d}^-_{-n''} \delta_{m'', -n''}) = \\
& \sum_{m' n' m'' n'' k \kappa} (2 m' 2 n' | J' M') (2 m'' 2 n'' | J'' M'') \cdot \\
& ((2 m' 2 n'' | k \kappa) [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}_{\kappa} (-1)^{n'} \delta_{m', -n'} - (2 m'' 2 n' | k \kappa) [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}_{\kappa} (-1)^{n''} \delta_{m'', -n''}).
\end{aligned} \tag{A8.3}$$

The last transformation was performed with the aid of (A1.14). By means of the relation (A3.10), one verifies the following result

$$\begin{aligned}
& [[\mathbf{d}^+ \times \mathbf{d}^-]^{(J')}_{M'}, [\mathbf{d}^+ \times \mathbf{d}^-]^{(J'')}_{M''}] = \sum_{k \kappa} [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}_{\kappa} \cdot \\
& (\sum_{n''} (2 m' 2, -m'' | J' M') (2 m'' 2 n'' | J'' M'') (2 m' 2 n'' | k \kappa) (-1)^{-m''} - \\
& (\sum_{n'} (2 m' 2, n' | J' M') (2 m'' 2, -m' | J'' M'') (2 m'' 2 n' | k \kappa) (-1)^{-m'}) = \\
& \sum_{k \kappa} [\mathbf{d}^+ \times \mathbf{d}^-]^{(k)}_{\kappa} \cdot \sqrt{(2J' + 1)} \sqrt{(2J'' + 1)} \cdot \\
& (J' M' J'' M'' | k \kappa) \left\{ \begin{matrix} J' & J'' & k \\ 2 & 2 & 2 \end{matrix} \right\} (-1)^k (1 - (-1)^{J'+J''+k}).
\end{aligned} \tag{A8.4}$$

A9 The commutator $[[\mathbf{d}^+ \times \mathbf{d}^-]^{(J')}_{M'}, [\mathbf{d}^+ \times \mathbf{d}^+]^{(J)}_M]$

This commutator is used in section 14.3. It is related to the commutator we dealt with in appendix A8. Analogously we transform as follows

$$\begin{aligned}
& [[\mathbf{d}^+ \times \mathbf{d}^-]^{(J')}_{M'}, [\mathbf{d}^+ \times \mathbf{d}^+]^{(J)}_M] = \\
& \sum_{m' n' m n} (2 m' 2 n' | J' M') (2 m 2 n | J M) (-1)^{n'} [\mathbf{d}^+_{m'} \mathbf{d}^-_{-n'}, \mathbf{d}^+_{-m} \mathbf{d}^+_n].
\end{aligned} \tag{A9.1}$$

$$\text{With } [\mathbf{d}^+_{m'} \mathbf{d}^-_{-n'}, \mathbf{d}^+_{-m} \mathbf{d}^+_n] = \mathbf{d}^+_{m'} \mathbf{d}^+_{-n} \delta_{-n', m} + \mathbf{d}^+_{m'} \mathbf{d}^+_{-m} \delta_{-n', n} \tag{A9.2}$$

one obtains by means of (A1.4)

$$\begin{aligned}
& [[\mathbf{d}^+ \times \mathbf{d}^-]^{(J')}_{M'}, [\mathbf{d}^+ \times \mathbf{d}^+]^{(J)}_M] = \\
& \sum_{m n} (2 m 2 n | J M) \sum_{m'} (2 m' 2, -m | J' M') (-1)^{-m} \mathbf{d}^+_{m'} \mathbf{d}^+_{-n} + \\
& (-1)^J \sum_{m n} (2 m 2 n | J M) \sum_{m'} (2 m' 2, -m | J' M') (-1)^{-n} \mathbf{d}^+_{m'} \mathbf{d}^+_{-m} = \\
& 2 \sum_m (2 m 2, M - m | J M) (2, M' + m, 2, -m | J' M') (-1)^m \mathbf{d}^+_{M'+m} \mathbf{d}^+_{M-m}.
\end{aligned} \tag{A9.3}$$

References

- Arima, A. and Iachello, F. (1975), Phys. Rev. Lett. 35, 1069.
- Arima, A. and Iachello, F. (1976a), Ann. Phys. 99, 253.
- Arima, A. and Iachello, F. (1976b), Phys. Rev. C14, 761.
- Arima, A. and Iachello, F. (1978), Ann. Phys. 111, 201.
- Arima, A. and Iachello, F. (1979), Ann. Phys. 123, 468.
- Arima, A. and Iachello, F. (1984), Adv. Nucl. Phys. 13, 139.
- Barrett, B., R. (1981), Revista mexicana de fisica 27, 533.
- Bayman, B. F. and Lande, A. (1966), Nucl. Phys. 77, 1.
- Bijker, R. and Dieperink, A. E. L. (1982), Nucl. Phys. A379, 221.
- Bohle, B., Richter, W., Steffen, W., Dieperink, A. E. L., Ludice, N. Lo., Palumbo, F. and Scholten, O. (1984), Phys. Lett. B137, 27.
- Bohr, A. and Mottelson, B. R. (1953), Dan. Vid. Selsk., Math.-Fys. Medd. No 16.
- Bohr, A. and Mottelson, B. R. (1975), *Nuclear Structure*, Vol. II, Benjamin, Reading, MA.
- Bonatsos, D. (1988), *Interacting Boson Models of Nuclear Structure*, Clarendon, Oxford.
- Brussaard, P. J. and Glaudemans, P. W. M. (1977), *Shell Model Applications in Nuclear Spectroscopy*, North Holland, Amsterdam.
- Castaños, O., Chacon, E., Frank, A. and Moshinsky, M. (1979), J. Math. Phys. 20, 35.
- Casten, R. F. (1980), Nucl. Phys. A347, 173.
- Casten, R. F. (1990), *Nuclear Structure from a Simple Perspective*, Oxford University Press.
- Casten, R. F. and Warner, D. D. (1988), Rev. Mod. Phys. 60, 389..
- Chacon, E., Moshinsky, M. and Sharp, R. T. (1976), J. Math. Phys. 17, 668.
- Chacon, E. and Moshinsky, M. (1977), J. Math. Phys. 18, 870.
- Condon, E. U. and Shortley, G. H. (1953), *The Theory of Atomic Spectra*, Cambridge University Press.
- Cornwell, J. F. (1990), *Group Theory in Physics*, Vol II, Academic Press.
- Duval, P. D. and Barrett, B. R. (1981), Phys. Rev. C 23, 492.
- Edmonds, A. R. (1964), *Drehimpulse in der Quantenmechanik*, Bibliographisches Institut, Mannheim.

- Edmonds, A. R. (1996), *Angular Momentum in Quantum Mechanics*, Princeton Landmark in Physics, Princeton.
- Eisenberg, J. M. and Greiner, W. (1987), *Nuclear Theory, Vol. 1, Nuclear Models*, 3. Ed, North-Holland.
- Elliott, J. P. (1958), Proc. Roy. Soc. A245,128; 562.
- Frank, A. and Van Isacker, P. (1994), *Algebraic Methods in Molecular and Nuclear Structure Physics*, John Wiley.
- Hamermesh, M. (1962), *Group Theory and its Applications to Physical Problems*, Addison-Wesley.
- Han, C. S., Chuu, D. S., Hsieh, S. T. (1990), Phys. Rev. C42, 280.
- Harper, R. P. Christy, A., Hall, I., Nagib, I. M. and Wakefield, B. (1971), Nucl. Phys. A162. 161.
- Hess, P. (1983), *Troisième cycle de la Physique*, Institut für Theoretische Physik der Universität Frankfurt/M, RFA.
- Iachello, F. and Arima, A. (1974), Phys. Lett. 53 B, 309.
- Iachello, F. and Arima, A. (1987), *The Interacting Boson Model*, Cambridge University Press.
- Iachello, F. (1988), in *International Conference on Contemporary Topics in Nuclear Structure Physics*, ed. by R. Casten, A. Frank, M. Moshinsky and S. Pittel, World Scientific, 109.
- Iachello, F. and Van Isacker, P. (1991), *The Interacting Boson-Fermion Model*, Cambridge University Press, Cambridge.
- Jolos, R. V., Lemberg, I. Kh. and Mikhailov, V. M. (1985), Sov. J. Part. Nucl. 16,2, Amer. Inst. of Phys.
- Kaup, U. (1983), *Anwendung des Interacting Boson Modells auf kollektive Zustände in mittelschweren Kernen*, Inaugural-Dissertation, Universität Köln.
- Krane, K. S. and Steffen, R. M. (1970), Phys. Rev. C2, 724.
- Landau, L. D. und Lifschitz, E. E. (1986), *Quantenmechanik*, Band 3, Akademie-Verlag, Berlin.
- Lawson, R. D. (1980), *Theory of the Nuclear Shell Model*, Clarendon Press, Oxford.
- Milner, W. T., Mc Gowan, F. K., Stelson, P. H., Robinson, R. L. and Sayer, R. O. (1969), Nucl. Phys. A129, 687.
- Otsuka, T. and Arima, A. (1978), Phys. Letters 77B, 1.
- Ring, P. and Schuck, P. (1980), *The Nuclear Many Body Problem*, Springer, New York.
- Rotenberg, M., Bivins, R., Metropolis, M. and Wooten Jr, J. K. (1959), *The 3-j and 6-j Symbols*, Technology Press MIT, Cambridge, MA.

- Scholten, O. (1980), *The Interacting Boson Approximation Model and Applications*. Proefschrift, Rijksuniversiteit, Groningen.
- Scholten, O. (1981), in *Interacting Bose Fermi Systems in Nuclei*, ed. by Iachello, F., 285, Plenum Press, New York.
- Scholten, O. (1991), in *Computational Nuclear Physics, Nuclear Structure*, Eds. K. Langanke, J. A. Maruhn, S. E. Koonin, Springer, Berlin.
- Talmi, I. (1993), *Simple Models of Complex Nuclei, The Shell Model and Interacting Boson Model*, Harwood Academic Publishers.
- Van Isacker, P., Frank, A. and Sun, H. Z. (1984), Ann. Phys. (NY) 157, 183.
- von Brentano, P., Gelberg, A. and Kaup, U. (1981), *Interacting Bose Fermi Systems in Nuclei*, Plenum Press (NY).

Index

9- j symbol	75	Hamilton operator of the IBM2	35
angular momentum operator	71	reduced Hamiltonian	38
of the IBM2	37	treatment of the complete	
boson-boson interaction		Hamiltonian	42
operator of the IBM2	36	IBFM basis states	58
boson-fermion interaction		IBFM calculation	
exchange term	60	Scholten instruction	64
monopole term	60	spectra of ^{79}Au isotopes	67
quadrupole term	60, 61	spectrum of ^{101}Rh	65
two-body operator	59	spectrum of ^{191}Ir	66
with d -, s - and a -operators	59	spectrum of ^{97}Tc	68
Casimir operator		IBFM calculation in the $u(5)$ limit	
of the subalgebra	11	low-lying levels of ^{133}La	62
$so(N)$ -eigenvalues	11	spectra of ^{147}Eu and ^{151}Eu	63
$su(3)$ -eigenvalues	11	IBM1 subalgebras	
Casimir operators	10	chains of algebras	18
of the $so(6)$ - or γ -instable limit ..	28	I. chain of Lie algebras	15
of the $su(3)$ - or rotational limit ..	23	II. chain of Lie algebras	16
of the $u(5)$ - or vibrational limit ..	22	III. chain of Lie algebras	17
Clebsch-Gordan coefficients	69	IBM2 basis states	37
3- j symbol	70	IBM2 calculation	
orthonormality	70	parameters ε and κ	46, 51
coefficients of fractional parentage		parameters $c_{0\nu}$, $c_{2\nu}$, χ_ν	47, 52
.....	77	spectra of ^{54}Xe isotopes	43, 44
electric quadrupole transition		spectra of ^{56}Ba isotopes	45
measured $B(E2)$ for ^{54}Xe , ^{56}Ba ,		spectra of ^{74}W isotopes	49, 50
^{58}Ce	48	typical spectrum for the I_1 -special	
measured $B(E2)$ for ^{74}W isotopes		case	57
.....	53	IBM2 subalgebras	54
operator of the IBM2	42	chains of Lie algebras	55
Hamilton operator in the $so(6)$ limit		eigenenergies of the chain I_1	56
.....	30	first series of special cases	54
eigenvalues	30	Lie algebra	
Hamilton operator in the $su(3)$ limit		basis functions	9, 10
.....	25	basis operators	10
eigenvalues	25	classical, dimensions	8
Hamilton operator in the $u(5)$ limit		definition	4
.....	20, 22	dimension n	6, 9
eigenvalues	22	isomorphic	5
Hamilton operator of the IBFM	58	n -dimensional vector space	5
$u(5)$ limit	61	$so(3)$ structure constants	8
Hamilton operator of the IBM1		$so(N)$	7
a further multipole representation		structure constants	5
.....	78	subalgebra	11

$u(N)$, $su(N)$	5	spectrum of ^{132}Ba	33
Majorana term	39, 40	spectrum of ^{196}Pt	32
matrices		typical nuclei	27
antihermitian	5	typical spectrum	30
antisymmetric	7	spherical basis	18
rank N	5	spherical harmonics	76
traces	6	$su(3)$ - or rotational limit	17
operator of normal ordering	60	spectrum of ^{170}Er	28
program BOSON2	42	typical nuclei	27
program NPBOS	42	typical spectrum	26
program package PHINT . 33, 78, 80		symmetry properties of coupled spin	
quadrupole-quadrupole operator		states	
of the IBM2	40	two bosons	72
Racah-coefficients	73	two fermions	72
6- j symbol	74	Talmi's Hamiltonian	53
normalised	73	tensor operator	77
reduced matrix element	77	Wigner-Eckart theorem	76
$so(6)$ - or γ -instable limit			