# HIGH- $P_T$ PION PRODUCTION IN HEAVY-ION COLLISIONS AT RHIC ENERGIES

G. G. BARNAFÖLDI, P. LÉVAI

KFKI Research Institute for Particle and Nuclear Physics, P.O. Box 49, Budapest, 1525, Hungary E-mail: bgergely@rmki.kfki.hu

G. PAPP

HAS Research Group for Theoretical Physics, Eötvös University Pázmány Péter sétány 1/A, Budapest, 1117 Hungary

G. FAI AND Y. ZHANG

Center for Nuclear Research, Department of Physics, Kent State University, Kent, OH 44242, USA

Perturbative QCD results on  $\pi$  production are presented in proton-proton, protonnucleus and nucleus-nucleus collisions from CERN SPS up to RHIC energy. A  $K_{jet}(s, p_T, Q)$  factor obtained from jet production is applied to perform nextto-leading order calculations. Using the intrinsic transverse momentum  $(k_T)$  we determined transverse momentum spectra for pions in wide energy region. We have investigated nuclear multiscattering and the Cronin effect at RHIC energies.

#### 1. Introduction

The calculation of high- $p_T$  particle production at SPS and RHIC energies requires solid theoretical background, based on perturbative QCD (pQCD). In these calculations, the choice of renormalization, factorization and fragmentation scales are typically chosen in the range  $p_T/3 \leq Q \leq 2p_T$ . Earlier we have found <sup>1</sup> that the theoretical reproduction of the measured nuclear effects in pA and AA collisions (specifically the Cronin effect <sup>2</sup>) strongly depends on the choice of the above scales. In the following we display new results analysing preliminary RHIC data <sup>3</sup> on pion production in  $AuAu \rightarrow \pi^0$ at  $\sqrt{s} = 130$  AGeV and 200 AGeV.

### 2. Perturbative QCD calculations with intrinsic $k_T$

In pQCD-improved parton model (including intrinsic  $k_T$ ) pion production was calculated in leading order (LO) <sup>4,5</sup> and next-to-leading order (NLO) <sup>6</sup>. An intermediate solution to speed up the time consuming full NLO calculations is to determine a  $K_{jet}(s, p_{T,jet}, Q)$  factor corresponds to higher order contributions at jet level in pp collisions <sup>7</sup> and apply it in the factorized pQCD equation. In this case the pion production in AA collision can be described as:

$$E_{\pi} \frac{\mathrm{d}\sigma_{\pi}^{AA}}{\mathrm{d}^{3}p} = \sum_{abcd} \int \mathrm{d}^{2}b \,\mathrm{d}^{2}r \,t_{A}(r) \,t_{B}(|\vec{b} - \vec{r}|) \int \mathrm{d}^{2}k_{T,a} \,\mathrm{d}^{2}k_{T,b} \,\mathrm{d}x_{a} \,\mathrm{d}x_{b} \,\mathrm{d}z_{c} \times \times g_{pA}(\vec{k}_{T,a}) \,f_{a/A}(x_{a},Q^{2}) \,g_{pA}(\vec{k}_{T,b}) \,f_{b/A}(x_{b},Q^{2}) \times \times \left[K_{jet}(s, p_{T,jet},Q) \,\frac{\mathrm{d}\sigma}{\mathrm{d}\hat{t}}^{ab \to cd}\right] \frac{D_{\pi/c}(z_{c},Q'^{2})}{\pi z_{c}^{2}} \,\hat{s} \,\delta(\hat{s} + \hat{t} + \hat{u}) \,.\,(1)$$

The partonic cross section,  $d\sigma/d\hat{t}$ , is the LO Born term, which is multiplied by  $K_{jet}(s, p_{T,jet}, Q)$  in eq. (1). ( $\hat{s}$ ,  $\hat{t}$  and  $\hat{u}$  are parton-level Mandelstam variables.) The nuclear thickness function,  $t_A(b) = \int dz \,\rho(b, z)$ , is normalized as  $\int d^2b t_A(b) = A$ . The nuclear parton distribution functions (PDF)  $f_{a/A}(x, Q^2)$  are based on the nucleon PDFs ( $f_{a/p}(x_a, Q^2)$ ) and modified in the nuclear environment ("shadowing")<sup>8</sup>. We are using NLO MRST <sup>9</sup> PDF set with  $Q = p_{T,jet}$  scale, where  $p_{T,jet} = p_T/z_c$ .

In a phenomenological approach, we introduced an extra 2-dimensional Gaussian transverse momentum distribution as the PDF's extension <sup>5</sup>:

$$g_{pA}(\vec{k}_T) = \frac{1}{\pi \langle k_T^2 \rangle_{pA}} e^{-k_T^2 / \langle k_T^2 \rangle_{pA}}, \text{ where } \langle k_T^2 \rangle_{pA} = \langle k_T^2 \rangle_{pp} + C \cdot h_{pA}(b).$$
(2)

We have found the Gaussian width  $(\langle k_T^2 \rangle_{pp})$  to be  $p_T$  independent in the window 2 GeV  $\leq p_T \leq 6$  GeV. Nuclear multiscatterings yield an extra broadening in pA collision, which can be related to the number of nucleonnucleon (NN) collisions in the medium and denoted by  $C \cdot h_{pA}(b)$ . Here C means the average increase in the transverse momentum width in one NN collision and  $h_{pA}(b)$  is a geometrical effectivity function. This latter quantity can be saturated <sup>4,5</sup>. The  $D_{\pi/c}(z_c, Q'^2)$  is the fragmentation function (FF), which gives the probability for parton c to fragment into  $\pi$  at momentum fraction  $z_c$  and fragmentation scale Q'. In this paper we fix  $Q' = p_T/2$  and use the KKP parametrization<sup>10</sup>.

#### 3. pQCD result on pp and AuAu at RHIC energies

In Figure 1 we display our results at  $\sqrt{s} = 130$  AGeV (*left panels*) and 200 AGeV (*right panels*). In the upper panels the dashed curves indicate the pp results which can be compared to the preliminary PHENIX data<sup>3</sup> (*full triangles*) at  $\sqrt{s} = 200$  GeV. Full dots indicate the  $\pi^0$  data from peripheral AuAu collisions, the calculated pion spectra (*solid lines*) approximately overlap with the data. We obtained  $N_{bin,p}^{130} = 19$  binary collisions for 60 – 80% peripheral case and  $N_{bin,p}^{200} = 10$  for 70 – 80% peripheral case. A satisfactory agreement between the data and the calculations demonstrate the validity of the improved pQCD calculation<sup>a</sup>.

Full squares indicate the data for central collisions. The number of binary collisions are  $N_{bin,c}^{130} = 857$  for 10% central AuAu case at  $\sqrt{s} = 130$ AGeV, and  $N_{bin,c}^{200} = 940$  at  $\sqrt{s} = 200$  AGeV. One can find a large difference between central data and theoretical spectra (full lines), a suppression factor of  $\sim 4-5$  appeared.

Lower panels display the nuclear modification factor, defined as

$$R_{AA}^{\pi} = \frac{1}{\langle N_{bin} \rangle} \frac{\sigma_{in}^{pp}}{\sigma_{in}^{AA}(b)} \frac{\mathrm{d}\sigma_{AA}/\mathrm{d}^{3}p_{\pi}}{\mathrm{d}\sigma_{pp}/\mathrm{d}^{3}p_{\pi}} \,. \tag{3}$$

This expression magnifies the enhancement of particle production in AuAu, relatively to the  $N_{bin}$  upscaled pp collision on a linear scale. Applying shadowing<sup>8</sup> and multiscattering, the Cronin peak can be seen clearly in both peripheral (dotted lines) and central (solid lines) collisions. The position of Cronin peak depends on the  $\langle k_T^2 \rangle_{pp}$  value, and the height depends on the extra broadening<sup>1</sup>,  $C \cdot h_{pA}(b)$ . One can see that the height of the peak in central case is larger than in peripheral one, which effect is caused by the number of nucleon-nucleon collisions inside a nucleus. We have fixed the PDF scale  $Q = p_{T,jet}$  and FF scale  $Q' = p_T/2$  to obtain the peak's maximum at  $p_T \approx 3$  GeV/c. The comparison of PHENIX data (full squares) with theoretical calculations display a difference of ~ 5. This suppression value is indicated on lower panels with a dashed lines. The origin of this suppression is jet quenching<sup>11</sup>.

<sup>a</sup>Parameters:  $\langle k_T^2 \rangle_{pp}^{130} = 1.8 \text{ GeV}^2$ ,  $\langle k_T^2 \rangle_{pp}^{200} = 1.5 \text{ GeV}^2$ ,  $C_{sat} = 0.4 \text{ and } \nu_{max} = 3$ .

0212111v3



Figure 1.  $K_{jet}$  based NLO pQCD  $\pi^0 p_T$ -spectra (upper panels) and the nuclear modification factor  $R_{AA}^{\pi}$  (lower panels) compared to preliminary PHENIX AuAu and pp data<sup>3</sup> at  $\sqrt{s} = 130$  AGeV (left) and 200 AGeV (right) RHIC energies. (See details in text.)

# Acknowledgments

This work was supported in part by U.S. DOE grant DE-FG02-86ER40251, NSF grant INT-0000211, FKFP220/2000, Hungarian grants OTKA T032796, T034842, and the HAS – Dubna agreement. Supercomputer time provided by BCPL in Norway and the EC – Access to Research Infrastructure action of the Improving Human Potential Programme is gratefully acknowledged, together with the ITOL at Eötvös University.

- References
  - 1. G.G. Barnaföldi et al., Heavy Ion Phys. (2003) nucl-th/0206006.
  - 2. J.W. Cronin et al., Phys. Rev. D 11, 3105 (1975).
  - 3. Adcox et al. (PHENIX) Phys. Rev. Lett. 88, 022301 (2001), D. d'Enterria for the PHENIX Coll. hep-ex/0209051 (2002), S. Mioduszewski for the PHENIX Coll. nucl-ex/0210021 (2002).

  - 4. G. Papp et al., Phys. Rev. C 61, 021902(R) (2000).
  - 5. Y. Zhang et al., Phys. Rev. C 65, 034903 (2002).
  - 6. G. Papp et al., hep-ph/0212249 (2002).
  - 7. G.G. Barnaföldi et al., J. Phys. G 27, 1767 (2001).
  - 8. X.N. Wang and M. Gyulassy, Phys. Rev. D 44, 3501 (1991).
  - 9. A.D. Martin et al., Eur. Phys. J. C 23, 73 (2002).
- 10. B.A. Kniehl, G. Kramer, and B. Pötter, Nucl. Phys. B 597, 337 (2001).
- 11. P. Levai et al., Nucl. Phys. A 698, 631c (2002).