Fragmentation or Recombination at High p_T ?

Rudolph C. Hwa¹ and C. B. Yang^{1,2}

¹Institute of Theoretical Science and Department of Physics University of Oregon, Eugene, OR 97403-5203, USA ²Institute of Particle Physics, Hua-Zhong Normal University, Wuhan 430079, P.

R. China

Abstract. All hadronization processes, including fragmentation, are shown to proceed through recombination. The shower partons in a jet turn out to play an important role in describing the p_T spectra of hadrons produced in heavy-ion collisions. Due to the recombination of the shower partons with the soft thermal partons, the structure of jets produced in AA collisions is not the same as that of jets produced in pp collisions.

The question is: "fragmentation or recombination?" The answer is not even fragmentation and recombination, but recombination only. What is usually called fragmentation is not a description of the hadronization process; it is a phenomenological parametrization of a black box represented by the function D(z)that gives the momentum fraction z of a hadron in a parton jet. We shall open up that black box and formulate the so-called fragmentation process in terms of recombination. To do so, we need to determine the distributions of the shower partons in a jet [1]. Those shower partons can recombine among themselves to form hadrons that are usually regarded as the products of the fragmentation process, but they can also recombine with the thermal partons not belonging to the jet. The latter process has hitherto been overlooked, but turns out to be very important in the moderate p_T range, $2 < p_T < 9$ GeV/c [2].

The fragmentation function (FF) can be expressed in the recombination model as a convolution [1]

$$zD_i^M(z) = \int \frac{dx_1}{x_1} \frac{dx_2}{x_2} F_{q\bar{q}'}^i(x_1, x_2) \ R^M(x_1, x_2, z) \ , \tag{1}$$

where M is the meson in a jet initiated by a hard parton i. $F_{q\bar{q}'}^i(x_1, x_2)$ is the joint distribution of the shower partons q and \bar{q}' in the jet, which are to recombine to form M. The recombination function $R^M(x_1, x_2, z)$ is known from previous studies [3, 4]. Since FFs are known from the parametrizations in [5], we can determine $F_{q\bar{q}'}^i$ from (1). Let S_i^j denote the shower parton distribution (SPD) of a parton j in the shower initiated by a hard parton i. On the assumption that $F_{q\bar{q}'}^i$ are factorizable into products of two SPDs, it is possible to reduce the 12 elements of the shower matrix S_i^j from hard parton i = u, d, s, g to shower parton j = u, d, s to just 5 basic distributions, $K_{\rm NS}, L, G, L_s$, and G_s , as follows:

$$S_i^j = \begin{pmatrix} K & L & L_s \\ L & K & L_s \\ L & L & K_s \\ G & G & G_s \end{pmatrix} , \qquad (2)$$

where $K = K_{\rm NS} + L$ and $K_s = K_{\rm NS} + L_s$. An example of K is $u \to u$ (valence + sea); L can be $u \to d$ or \bar{d} , and L_s can be $u \to s$. The five SPDs are shown in Fig. 1 as functions of their momentum fractions [1]. The anti-quarks \bar{u} , \bar{d} and \bar{s} have the same structure, and are related to u, d, s as sea, and vice-versa. The gluons are not included among the shower partons j because all gluons have been converted to quark pairs. Using those SPDs it is possible to calculate other FFs, both measured (e.g., for proton) and unmeasured (e.g., for two pions).



Figure 1. Shower parton distributions plotted as functions of their momentum fractions.

The above discussion on shower partons is independent of heavy-ion collisions. However, once the SPDs are known, their relevance to high- p_T hadrons produced at RHIC is conceptually unavoidable, since shower partons can recombine with thermal partons. To make this point transparent, consider the production of a u quark at $p_T = 8 \text{ GeV/c}$ by hard scattering. Among the many shower partons that it creates, we show in Fig. 2 by the solid line the distribution of u quarks in the shower. That distribution corresponds to K(z), which is the sum of the valence $K_{\text{NS}}(z)$ and the sea L(z) that are shown in Fig. 1. The thermal partons are shown by the dashed line in Fig. 2. It should be clear that the recombination of the u quark in the shower with a \overline{d} thermal quark to form a π^+ cannot be ignored, when the parton momenta are not too far apart. The thermal-shower recombination is important because the yield of that component is raised by the high density of soft partons, while the p_T of the produced hadrons is increased by the semi-hardness of the shower partons.

The invariant distribution of pions in a 1D description is

$$p\frac{dN^{\pi}}{dp} = \int \frac{dp_1}{p_1} \frac{dp_2}{p_2} F^i_{q\bar{q}'}(p_1, p_2) \ R^{\pi}(p_1, p_2, p) \ . \tag{3}$$

The $q\bar{q}'$ distribution has four components

$$F_{q\bar{q}'} = \mathcal{T}\mathcal{T} + \mathcal{T}\mathcal{S} + \mathcal{S}\mathcal{S} + (\mathcal{S}\mathcal{S})_2 , \qquad (4)$$

where \mathcal{T} denotes thermal partons and \mathcal{S} shower partons. Thus \mathcal{SS} represents two shower partons in one jet, and $(\mathcal{SS})_2$ signifies two shower partons from two jets. A sum over all hard partons *i* is implied. Mathematical details of what each component



Figure 2. Shower parton distribution of u quarks (in solid line) due to a hard u quark at 8 GeV/c. The dashed line shows the thermal parton distribution.

contains can be found in Ref. [2]. Letting p in (3) represent p_T and dividing that equation by p_T^2 we can calculate $dN^{\pi}/p_T dp_T$. The result for the pion p_T spectrum in central Au+Au collisions at $\sqrt{s} = 200$ GeV is shown in Fig. 3, exhibiting good agreement with the data [6]. Note that the thermal-shower component dominates over other components in the region $3 < p_T < 8$ GeV/c. The shower-shower (1 jet) component (i.e., SS) is the usual fragmentation and is not important until $p_T > 8$ GeV/c.



Figure 3. Transverse momentum distribution of π^0 in Au-Au collisions. Data are from [6]. The solid line is the sum of four contributions to the recombination of partons: \mathcal{TT} (dashed line); \mathcal{TS} (line with crosses); \mathcal{SS} , two shower partons in one jet (line with open circles); $(\mathcal{SS})_2$, two shower partons from two overlapping jets (line with squares).

Fragmentation or Recombination at High p_T ?

There is an essential parameter ξ that has been adjusted to fit the high- p_T data. It represents the fraction of hard partons that can get out of the dense medium created in central collisions to produce the showers. It is an average quantity that parametrizes the energy loss effect. Thus S and SS are proportional to ξ , but not \mathcal{T} . The value of $\xi = 0.07$ is determined by fitting the high- p_T spectrum. It is smaller than the value of the nuclear modification factor, $R_{AA}(p_T) \approx 0.2$, for $p_T > 4$ GeV/c. The reason for the difference is that the denominator of R_{AA} is the scaled pp spectrum (mainly fragmentation), which is smaller than our \mathcal{TS} recombination, as is evident in Fig. 3, the dominant component that determines ξ .

It is important to emphasize that because of the dominance of the \mathcal{TS} recombination, the jet structure in AA collisions cannot be the same as that in pp collisions, for which the SS recombination gives rise to the usual fragmentation. Thus there should be more particles associated with a trigger particle in AA collisions than in pp collisions for the simple reason that the abundance of soft thermal partons in the former case enhances the hadrons formed at moderate p_T through recombination. There exists experimental evidence for the excess of associated particles on the same side, given in terms of the quantity I_{AA} by STAR [7] for $3 < p_T^{\text{trig}} < 4 \text{ GeV/c}$. In that p_T range for the trigger particles it is found that $I_{AA} \approx 1.5$, whereas the value of 1 is expected if the jet structures in AA and pp collisions are identical.

The formalism described above for Au+Au collisions has very recently been extended to d+Au collisions. It is found that the recombination of soft and shower partons can account for the Cronin effect [8] that is usually attributed to the broadening of the parton k_T width in the initial state by multiple scattering [9]. In our study the enhancement of the pion spectrum at moderate p_T for more central d+Au collisions is due entirely to the final-state interaction, since there are more soft partons to recombine at higher centrality [10]. That is another instance where the adherence to the usual fragmentation picture leads one to a path that misses a crucial element of the hadronization process.

The recombination model has by now been shown to explain many features of hadron production in heavy-ion collisions [11]. What we add here is a new piece of the recombination products, arising from the interaction between the soft and shower partons. That new addition has far-reaching consequences. Theoretically, it means that the separation of partons into two noninteracting components, soft and hard, is an oversimplification. Experimentally, it means that jets in AA collisions are different from those in pp collisions. More precise quantification of that difference can provide further elucidation of the underlying hadronization process.

This work was supported, in part, by the U. S. Department of Energy under Grant No. DE-FG03-96ER40972 and by the Ministry of Education of China under Grant No. 03113.

- [1] R. C. Hwa and C. B. Yang, hep-ph/0312271.
- [2] R. C. Hwa and C. B. Yang, nucl-th/0401001.
- [3] R. C. Hwa, Phys. Rev. D 22, 1593 (1980).
- [4] R. C. Hwa and C. B. Yang, Phys. Rev. C 66, 025205 (2002).
- J. Binnewies, B. A. Kniehl and G. Kramer, Z. Phys. C 65, 471 (1995); B. A. Kniehl, G. Kramer and B. Pötter, Nucl. Phys. B582, 514 (2000).
- [6] S. S. Adler, PHENIX Collaboration, Phys. Rev. Lett. 91, 072301 (2003).
- [7] C. Adler et al., STAR Collaboration, Phys. Rev. Lett. 90, 082302 (2003).
- [8] J. W. Cronin et. al., Phys. Rev. D 11, 3105 (1975).
- [9] A. Accardi, hep-ph/0212148.
- [10] R. C. Hwa and C. B. Yang, nucl-th/0403001.
- [11] For a review, see R. J. Fries, Quark Matter 2004, in this volume.