# Calculations of the Exclusive Processes ${}^{2}H(e, e'p)n$ , ${}^{3}He(e, e'p){}^{2}H$ and ${}^{3}He(e, e'p)(pn)$ within a Generalized Eikonal Approximation

C.Ciofi degli Atti and L.P. Kaptari<sup>\*</sup>

Department of Physics, University of Perugia and Istituto Nazionale di Fisica Nucleare, Sezione di Perugia, Via A. Pascoli, I-06123, Italy

(Dated: December 7, 2018)

# Abstract

The exclusive processes  ${}^{2}H(e, e'p)n$ ,  ${}^{3}He(e, e'p){}^{2}H$  and  ${}^{3}He(e, e'p)(pn)$ , have been analyzed using realistic few-body wave functions and treating the final state interaction (FSI) within a Generalized Eikonal Approximation (GEA), based upon the direct calculation of the Feynman diagrams describing the rescattering of the struck nucleon with the nucleons of the A-1 system. The approach represents an improvement of the conventional Glauber approach (GA), in that it allows one to take into account the effects of the nuclear excitation of the A-1 system on the rescattering of the struck nucleon. Using realistic three-body wave functions corresponding to the AV18 interaction, the results of our parameter free calculations are compared with available experimental data. It is found that in some kinematical conditions FSI effects represent small corrections, whereas in other kinematics conditions they are very large and absolutely necessary to provide a satisfactory agreement between theoretical calculations and experimental data. It is shown that in the kinematics of the experimental data which have been considered, covering the region of missing momentum and energy with  $p_m \leq 0.6 \ GeV/c$  and  $E_m \leq 100 \ MeV$  in the perpendicular kinematics, the GA and GEA predictions differ only by less than  $\simeq 3 - 4\%$ .

<sup>\*</sup>On leave from Bogoliubov Lab. Theor. Phys.,141980, JINR, Dubna, Russia

#### I. INTRODUCTION

One of the main aims of nowadays hadronic physics is the investigation of the limits of validity of the so called *Standard Model* of nuclei, i.e. the description of nuclei in terms of the solution of the non relativistic Schrödinger equation containing realistic nucleon-nucleon interactions. To this end, exclusive lepton scattering could be very useful, for it might yield relevant information on the nuclear wave function, provided the initial and final states involved in the scattering process are described within a consistent, reliable approach. In the case of *few-body systems*, a consistent treatment of initial and final states is nowadays possible at low energies (see e.g. [1, 2] and References therein quoted), but at higher energies, when the number of partial waves sharply increases and nucleon-nucleon (NN) interaction becomes highly inelastic, the Schrödinger approach becomes impractical and other methods have to be employed. In the case of *complex nuclei*, additional difficulties arise due to the approximations which are still necessary to solve the many-body problem. As a matter of fact, whereas fundamental progress has been made in recent years in the calculation of various properties of light nuclei (see e.g. [3] and [4] and References therein quoted), much remains to be done for the treatment of the continuum, for which various approximate treatments of the final state cannot be avoided. In this context, it should be stressed that calculations involving few-body systems, where the ground state can be treated exactly, can also be very useful to investigate the limits of validity of various approximate schemes to treat the continuum and their possible extension to complex nuclei.

The aim of this paper is to present the results of a systematic theoretical investigation of the exclusive process A(e, e'p)B off <sup>2</sup>H (to be also denoted by D) and <sup>3</sup>He, based on a reliable description of:

- 1. initial state correlations (ISC), treated by the use of the status-of-the-art few-body wave functions [2] corresponding to the AV18 interaction [5];
- 2. final state interactions (FSI), treated within a relativistic framework based upon the calculation of the relevant Feynman diagrams which describe the rescattering of the struck nucleon by the other A 1 spectator nucleons of the target.

Whereas a correct treatment of ISC in few-body systems is automatically achieved by the use of realistic wave functions, the treatment of FSI at high energies is still matter of discussions. The approach we are going to use has several non trivial advantages, in that it allows one to work within a relativistic framework provided by the use of Feynman diagrams and, moreover, it can be applied, in principle, to the treatment of exclusive A(e, e'p)B processes off complex nuclei as well. It should be stressed, at this point, that the diagrammatic approach we are talking about is not a new one: it has been first formulated in Ref. [6] and [7] (see also Ref. [8]), within a spin-less treatment of particle-nucleus scattering, and applied subsequently to various types of high energy processes with nuclear targets. More recently, the diagrammatic approach has been generalized to the treatment of the FSI in exclusive A(e, e'p)B [9, 10, 11] and A(e, e'2p)B [12] processes, and a Feynman diagram approach has also been used in Ref. [13, 14] and [15] to take into account off-shell effects both in inclusive, A(e, e'p)X, and exclusive, A(e, e'p)B, processes.

The diagrammatic approach we are referring to, is a generalization of the standard Glauber Approach (GA) [16] based on the eikonal approximation, so that, following Refs. [12], we will call it Generalized Eikonal Approximation (GEA).

It is well known that the application of the GA to the treatment of A(e, e'p)B processes requires the following approximations: i) the NN scattering amplitude is obtained within the eikonal approximation; ii) the nucleons of the spectator system A - 1 are stationary during the multiple scattering with the struck nucleon (the *frozen approximation*), and iii) only perpendicular momentum transfer components in the NN scattering amplitude are considered. In the GEA the *frozen approximation* is partly removed by taking into account the excitation energy of the A - 1 system, which results in a correction term to the standard profile function of GA, leading to an additional contribution to the longitudinal component of the missing momentum.

In the present paper we apply both the GA and the GEA to the calculation of the processes  ${}^{2}H(e, e'p)n$ ,  ${}^{3}He(e, e'p){}^{2}H$ , and  ${}^{3}He(e, e'p)(np)$ , and compare our results with available experimental data [17, 18, 19, 20, 21, 22]. The  ${}^{3}He$  wave function of the Pisa group [2], corresponding to the AV18 interaction [5], will be used in the calculations. We will not consider, for the time being, Meson Exchange Currents (MEC),  $\Delta$ -Isobar Configurations, and similar effects, which have been the object of intensive theoretical studies in A(e, e'p)B processes off both few-body systems (see e.g. [23, 24]) and complex nuclei (see e.g. [25] and References therein quoted). As in Refs. [9, 10, 11, 12], we fully concentrate on the effects of the FSI but we will consider kinematical conditions for which the effects from meson

exchange currents (MEC) and  $\Delta$  excitation effects are expected to be small corrections, and, whenever possible, we will compare our results with the results by other authors which include these effects.

The structure of the paper is as follows: in Section II the basic formalism of leptonhadron scattering is briefly illustrated and the main formulae are obtained; in Section III the concept of Plane Wave Impulse Approximation (PWIA) and Spectral Function are recalled; in Section IV, the GEA is introduced, the relevant Feynman diagrams which one needs to take into account in the treatment of the full FSI are analyzed, and the problem of the factorization of the lepton-nucleus cross section within the GA and GEA is also discussed; the results of the calculations and their comparison with available experimental data are shown in Section V; eventually, the Summary and Conclusions are presented in Section VI. Some details concerning the formal aspects of our approach are given in Appendices A and B. Preliminary results of our calculations have been reported in Ref. [26] and [27].

### II. BASIC FORMULAE OF (e, e'p) SCATTERING OFF NUCLEI

The one-photon-exchange diagram for the process A(e, e'p)(A-1), where A-1 denotes a system of A-1 nucleons in a bound or continuum state, is presented in Fig. 1, where the relevant four-momenta in the scattering processes are shown, namely the electron momenta before and after interaction,  $k = (E, \mathbf{k})$  and  $k' = (E', \mathbf{k}')$ , the momentum of the target nucleus  $P_A = (E_A, \mathbf{P}_A)$  and, eventually, the momenta of the final proton and the final A-1system,  $p_1 = (\sqrt{\mathbf{p}_1^2 + M_N^2}, \mathbf{p}_1)$  and  $P_{A-1} = (\sqrt{\mathbf{P}_{A-1}^2 + (M_{A-1}^f)^2}, \mathbf{P}_{A-1})$ , where  $M_N$  is the nucleon mass,  $M_{A-1}^f = M_{A-1} + E_{A-1}^f$ , and  $E_{A-1}^f$  is the *intrinsic* excitation energy of the A-1 system.

Let us briefly recall some useful formulae regarding the process described by the diagram shown in Fig. 1. The differential cross section for the exclusive process has the following form (see e.g. [28])

$$\frac{d^6\sigma}{dE'd\Omega' \ d^3\mathbf{p}_1} = \sigma_{Mott} \tilde{l}^{\mu\nu} W^A_{\mu\nu},\tag{1}$$

where  $\sigma_{Mott} = \frac{4\alpha^2 E'^2 \cos^2 \frac{\theta}{2}}{Q^4}$  is the Mott cross section,  $\alpha$  the fine-structure constant,  $Q^2 = -q^2 = -(k-k')^2 = \mathbf{q}^2 - q_0^2 = 4EE' \sin^2 \theta/2$  the four-momentum transfer,  $\theta \equiv \theta_{\mathbf{k}\mathbf{k}'}$  the scattering angle. The quantities  $\tilde{l}_{\mu\nu}$  and  $W^A_{\mu\nu}$  are the reduced leptonic and hadronic tensors,

respectively; the former has the well known standard form ([28]), whereas the latter can be written as follows

$$W^{A}_{\mu\nu} = \frac{1}{4\pi M_{A}} \sum_{\alpha_{A}} \sum_{\alpha_{A-1},\alpha_{N}} (2\pi)^{4} \delta^{(4)} (P_{A} + q - P_{A-1} - p_{1}) \times \langle \alpha_{A} \mathbf{P}_{A} | \hat{J}^{A}_{\mu}(0) | \alpha_{N} \mathbf{p}_{1}, \alpha_{A-1} \mathbf{P}_{A-1} E^{f}_{A-1} \rangle \langle E^{f}_{A-1} \mathbf{P}_{A-1} \alpha_{A-1}, \mathbf{p}_{1} \alpha_{N} | \hat{J}^{A}_{\nu}(0) | \alpha_{A} \mathbf{P}_{A} \rangle ,$$
(2)

where  $\alpha_i$  denotes the set of discrete quantum numbers of systems A, A - 1 and N. In Eq. (2) the vector  $|\alpha_N \mathbf{p}_1, \alpha_{A-1} \mathbf{P}_{A-1} E_{A-1}^f\rangle$  consists asymptotically of a nucleus A - 1, with momentum  $\mathbf{P}_{A-1}$  and intrinsic excitation energy  $E_{A-1}^f$ , and a nucleon with momentum  $\mathbf{p}_1$ . Two relevant experimentally measurable quantities which characterize the process are the missing momentum  $\mathbf{p}_m$  (i.e. the momentum of the A - 1 system), and the missing energy  $E_m$  defined, respectively, by

$$\mathbf{p}_m = \mathbf{q} - \mathbf{p}_1 \quad E_m = \sqrt{P_{A-1}^2} + M_N - M_A = M_N + M_{A-1} - M_A + E_{A-1}^f = E_{min} + E_{A-1}^f, (3)$$

where  $E_{min} = E_A - E_{A-1} = M_N + M_{A-1} - M_A$ , and the (positive) ground-state energies of A and A - 1 are denoted by  $E_A$  and  $E_{A-1}$ , respectively. The exclusive cross section can then be written in the well-known form

$$\frac{d^6\sigma}{d\Omega' dE' \ d^3\mathbf{p}_m} = \sigma_{Mott} \sum_i V_i \ W_i^A(\nu, Q^2, \mathbf{p}_m, E_m), \tag{4}$$

where  $i \equiv \{L, T, LT, TT\}$ , and  $V_L, V_T, V_{LT}$ , and  $V_{TT}$  are well-known kinematical factors.

The evaluation of the nuclear response functions  $W_i^A$  requires the knowledge of the nuclear vectors  $|\alpha_A \mathbf{P}_A\rangle$  and  $|\alpha_N \mathbf{p}_1, \alpha_{A-1} \mathbf{P}_{A-1} E_{A-1}^f\rangle$  and the nuclear current operators  $\hat{J}_{\mu}^A(0)$ . Nowadays, there is no rigorous quantum field theory to describe, from first principles, a many body hadronic system, and one is forced to adhere to various approximations. Whereas at relatively low energies a consistent non relativistic treatment of the electro-disintegration of two- and three-body systems can be pursued, with increasing energy the treatment of the three- body final state requires proper approximations. In the present paper we describe the two- and three-body ground states in terms of realistic wave functions generated by modern two-body interactions [2], and treat the final state interaction by a diagrammatic approach of the elastic rescattering of the struck nucleon with the nucleons of the A - 1 system. The relevant diagrams which, within such an approximation, replace the One-Photon-Exchange diagram of Fig. 1, are shown in Fig. 2: the first one represents the *Plane Wave Impulse* Approximation (PWIA), whereas the other ones the final state rescattering (FSI). Although the PWIA appears to have a limited range of validity, it is useful to analyze its predictions since, within such an approximation, the cross section is directly related to a quantity, the *Spectral Function*, which, in the case of few-body systems, can be calculated with high degrees of accuracy (see [29, 30, 31, 32]). The relevant point here is that, provided the FSI of the struck nucleon with the A-1 system can be disregarded, the Spectral Function yields direct information on the nuclear wave function. For such a reason, we will present our results obtained within two distinct approaches:

1. the *PWIA* (Fig. 2a)), when the struck proton is described by a plane wave, whereas the A - 1 system in the final state, with momentum  $P_{A-1}$ , represents the bound or continuum state solutions of the Schrödinger equation with the same potential used to obtain the A-body wave function (note that some authors call PWIA the state in which *all* particles in the continuum are described by plane waves);

2. the full FSI approach (Fig. 2b,c)), when the A-1 system (in the ground or continuum states) is still described by the exact solution of the Schrödinger equation, and the interaction of the struck nucleon with the A-1 nucleons is treated by evaluating the Feynman diagrams of Fig. 2, either in the GA or the GEA approximations.

# III. THE PLANE WAVE IMPULSE APPROXIMATION AND THE NUCLEAR SPECTRAL FUNCTION

The main merit of the PWIA is that it allows one to express the *nuclear* response functions  $W_i^A$  in terms of the *nucleon* response functions which are very well known from e - N experiments. As a matter of fact, by expressing the hadronic tensor for the nucleus A (Eq. (2)) in terms of the hadronic tensor for the nucleon N

$$W_{\mu\nu}^{N} = \frac{1}{4\pi M_{N}} \sum_{\alpha_{N}} \sum_{\alpha_{N}'} (2\pi)^{4} \delta^{(4)}(p+q-p_{1}) \langle \alpha_{N} \mathbf{p}_{1} | \hat{J}_{\mu}^{N}(0) | \alpha_{N}' \mathbf{p}_{1}' \rangle \langle \alpha_{N}' \mathbf{p}_{1}' | \hat{J}_{\nu}^{N}(0) | \alpha_{N} \mathbf{p}_{1} \rangle, \quad (5)$$

the cross section assumes the following form (see e.g. Refs [33, 34, 35])

$$\frac{d^{\mathbf{o}}\sigma}{dE'd\Omega'd\mathbf{p}_m} = K(Q^2, x, \mathbf{p}_m)\,\sigma^{eN}(\bar{Q}^2, \mathbf{p}_m)P_A(|\mathbf{k}_1|, E),\tag{6}$$

where  $\bar{Q}^2 = \mathbf{q}^2 - \bar{q}_0^2$  ( $\bar{q}_0 = q_0 + M_A - \sqrt{(\mathbf{k}_1^2 + (M_{A-1}^f)^2 - \sqrt{\mathbf{k}_1^2 + M_N^2})}$ , and  $K(Q^2, x, \mathbf{p})$  a kinematical factor. In Eq. (6),  $\sigma^{eN}(\bar{Q}^2, \mathbf{p}_m)$  is the cross section describing electron scattering by an off-shell nucleon,  $x = Q^2/2M_Nq_0$  is the Bjorken scaling variable,  $\mathbf{k}_1 = -\mathbf{p}_m$  is the nucleon momentum before interaction,  $E \equiv E_{mis} = E_{min} + E_{A-1}^{f}$  is the removal energy and  $P(|\mathbf{k}_{1}|, E)$  is the nucleon Spectral Function, which can be written as follows:

$$P(|\mathbf{k}_{1}|, E) = \frac{1}{(2\pi)^{3}} \frac{1}{2J_{A}+1} \sum_{f} \sum_{\mathcal{M}_{A}, \mathcal{M}_{A-1}, \sigma_{N}} \left| \langle \alpha_{A} \mathbf{P}_{A} | \alpha_{N} \mathbf{k}_{1}, \alpha_{A-1} \mathbf{P}_{A-1} E_{A-1}^{f} \rangle \right|^{2} \times (7)$$
$$\times \delta \left( E - (E_{A-1}^{f} + E_{min}) \right),$$

where  $\mathcal{M}_A$ ,  $\mathcal{M}_{A-1}$ , and  $\sigma_n$ , are the spin projections, and the sum over f includes all possible discrete and continuum states of the A-1 system.

Whereas the Spectral Function for the Deuteron (D) has a particularly simple form, viz.

$$P_D(|\mathbf{k}_1|, E) = n_D(|\mathbf{k}_1|)\delta(E - \epsilon_D),$$
(8)

where  $\epsilon_D$  is the (positive) binding energy of the deuteron and  $n_D(|\mathbf{k_1}|) = (2\pi^2)^{-1} (u_S^2(|\mathbf{k_1}|) + u_D^2(|\mathbf{k_1}|))$  the nucleon momentum distribution, in the case of A=3, the proton Spectral Function consists of two parts,

$$P_{He}(|\mathbf{k}_1|, E) = P_{gr}(|\mathbf{k}_1|, E) + P_{ex}(|\mathbf{k}_1|, E),$$
(9)

The first one, or ground part  $P_{gr}$ , has the following form

$$P_{gr}(|\mathbf{k}_1|, E) = n_{gr}(|\mathbf{k}_1|)\delta(E - E_{min}), \tag{10}$$

where  $E_{min} = |E_3| - |E_2| \approx 5.49 MeV$ , and  $n_{gr}(|\mathbf{k}_1|)$ , which corresponds to the two-body break-up (2bbu) channel  ${}^{3}He \rightarrow D + p$ , is (hereafter, the projection of the spin of nucleon *i* will be denoted by  $s_i$ )

$$n_{gr}(|\mathbf{k}_1|) = \frac{1}{(2\pi)^3} \frac{1}{2} \sum_{\mathcal{M}_3, \mathcal{M}_2, s_1} \left| \int e^{-i\boldsymbol{\rho}\mathbf{k}_1} \chi_{\frac{1}{2}s_1}^{\dagger} \Psi_D^{\mathcal{M}_2\dagger}(\mathbf{r}) \Psi_{He}^{\mathcal{M}_3}(\boldsymbol{\rho}, \mathbf{r}) d\boldsymbol{\rho} d\mathbf{r} \right|^2.$$
(11)

In Eq. (11)  $\Psi_{He}^{\mathcal{M}_3}(\boldsymbol{\rho}, \mathbf{r})$  is the <sup>3</sup>*He* wave function,  $\mathcal{M}_3$  the projection of the spin of <sup>3</sup>*He*, and  $\mathbf{r}$  and  $\boldsymbol{\rho}$  the Jacobi coordinates describing, respectively, the motion of the spectator pair and the motion of the struck (active) nucleon with respect to the CM of the pair.

The second, or *excited* part  $P_{ex}$ , of  $P_{He}(|\mathbf{k}_1|, E)$ , corresponds to the three-body break-up (3bbu) channel  ${}^{3}He \rightarrow (np) + p$  and can be written as follows

$$P_{ex}(|\mathbf{k}_{1}|, E) = \frac{1}{(2\pi)^{3}} \frac{1}{2} \sum_{\mathcal{M}_{3}, S_{23}, s_{1}} \int \frac{d^{3}\mathbf{t}}{(2\pi)^{3}} \left| \int e^{-i\boldsymbol{\rho}\mathbf{k}_{1}} \chi^{\dagger}_{\frac{1}{2}s_{1}} \Psi^{\mathbf{t}\dagger}_{np}(\mathbf{r}) \Psi^{\mathcal{M}_{3}}_{He}(\boldsymbol{\rho}, \mathbf{r}) d\boldsymbol{\rho} d\mathbf{r} \right|^{2} \times \delta \left( E - \frac{\mathbf{t}^{2}}{M_{N}} - E_{3} \right),$$
(12)

where  $\Psi_{np}^{\mathbf{t}}(\mathbf{r})$  is the two-body continuum wave function characterized by spin projection  $S_{23}$ and by the relative momentum  $\mathbf{t} = \frac{\mathbf{k}_2 - \mathbf{k}_3}{2}$  of the np pair in the continuum. Obviously, for the neutron Spectral Function, only the excited part (12) contributes.

In Fig. 3 we show the Spectral Function of  ${}^{3}He$  obtained using the variational three-body wave function by the Pisa group [2] corresponding to the realistic AV18 potential [5] (see Appendix A). The two-body wave function entering Eq. (12) has been obtained by solving the Schrödinger equation for the continuum using the same AV18 two nucleon potential. Our results for the Spectral Function agree with the ones obtained in Ref. [31], where the same three-body wave function has been used. The normalization of the Proton Spectral Function has been fixed to 2 (two protons) and the normalization of the neutron Spectral Function to one. In Fig. 3 we also show the results predicted by the Plane Wave Approximation (PWA), which corresponds to the replacement of the continuum interacting (n-p) pair wave function with two plane waves. The left panel of Fig. 3 shows  $n_{gr}$  and the right panel  $P_{ex}$ . It can be seen that: i)  $P_{ex}$  exhibits maxima centered approximately at  $E_m \sim \frac{\mathbf{k}_1^2}{4M_N}$ , ii) around these values of  $E_m$  and  $\mathbf{k}_1$  the Spectral Functions, calculated disregarding the interaction in the NN-pair in the continuum (PWA) and taking it into account (PWIA), are almost identical, in agreement with the results obtained long ago [32] with the Spectral Function corresponding to the Reid Soft Core Interaction [36]. The region centered at  $E_m \sim \frac{{\bf k}_1^2}{4M_N}$  is the so-called two-nucleon correlation region [37], when one of the nucleons of the spectator NN-pair is fast, the other one being basically at rest (for an improved description which takes into account the motion of the third, uncorrelated nucleon, or the A-2 spectator system in case of heavier nuclei, see [38]). Then the fast nucleon becomes strongly correlated with the active nucleon (the proton, in the case of the proton Spectral Function, or the neutron, in the case of the neutron Spectral Function) forming a correlated pair which carries most of the nuclear momentum. In this case, it is intuitively expected that the slow nucleon acts as a passive spectator and, consequently, only the interaction in the correlated pair can be relevant for the Spectral Function. Hence, in this region the calculations including or omitting the interaction in the spectator pair, are expected to provide essentially the same results, as confirmed by present and previous calculations of the Spectral Function [26, 27, 32]. The situation which has been just described, is clearly illustrated in Fig. 4, where the three-dimensional neutron Spectral Function is presented.

The PWIA results suggest that experimental insight about the structure of the nuclear

wave function at short distances can be obtained from A(e, e'p)(A-1) processes, provided the PWIA entirely exhausts the reaction mechanism. Unfortunately, we know that in many cases the simplified PWIA mechanism fails to describe the experimental data (see, e.g. a recent discussion in Ref. [39]). However, a properly chosen kinematics can still leave room for studying NN correlations. It is clear, from Figs. 3 and 4, that such a kinematics should be located around the two-nucleon correlation region, in order to exclude the influence of the final state interaction between the spectator nucleons. This requires high values of the missing energy and momentum of the active nucleon (see Fig. 4). Another important condition is that the range of  $|\mathbf{q}|$  and  $q_0$  should not be too far from the quasi-elastic peak, where  $x \simeq 1$ . In this case the corrections from the off mass shell effects and meson production are minimized, and only the final state interaction of the hit nucleon with the spectators becomes relevant. However, it should always be kept in mind, that if a region exists where the interaction in the spectator pair (the A-1 system in case of complex nuclei) can be neglected, this is no guarantee that the interaction of the struck nucleon with the nucleons of the spectator pair (the A-1 system), can be neglected as well. It is clear therefore that one has to go beyond the PWIA, which is precisely the aim of the present paper. The effects of the full FSI on the process  ${}^{3}He(e, e'2p)n$  have been recently investigated, treating the FSI within the GA [40]. In the present paper we will investigate the same topic in A(e, e'p)Xprocess off  ${}^{2}H$  and  ${}^{3}He$  within both the GA and the GEA.

# IV. THE FULL FINAL STATE INTERACTION WITHIN THE GENERALIZED EIKONAL APPROXIMATION

Let us consider the interaction of the incoming virtual photon,  $\gamma^*$ , with a bound nucleon (the active nucleon) of low virtuality  $(p^2 \sim M_N^2)$  at a kinematics not very different from the quasi-elastic one, i.e. corresponding to  $x \sim 1$ . In the quasi-elastic kinematics, the virtuality of the struck nucleon after  $\gamma^*$ -absorption is also rather low and, provided  $\mathbf{p}_1$  is sufficiently high, nucleon rescattering with the "spectator" A - 1 can be described to a large extent in terms of multiple elastic scattering processes in the forward direction (in the system of reference where the target nucleon is at rest). These rescattering processes are diagrammatically depicted in Fig. 2 where, as in the rest of this paper, the internal and intermediate state momenta are denoted by  $k_i$ 's and the final state momenta by  $p_i$ 's. The diagrams essentially describe the process of multiple scattering in the most general case, within the assumption that all intermediate nucleons are on-shell. The low virtuality (before and after  $\gamma^*$  absorption) of the active nucleon, coupled with the forward propagation, allows one to simplify the description of the final state interaction, which can be treated within the eikonal approximation. Before illustrating in detail the approach we have used to treat FSI in (e, e'p) reactions, we would like to discuss an important related issue (see also [41]), namely the validity of the factorization approximation, frequently used in calculations at high  $Q^2$ , and consisting in factorizing the (e, e'p) cross section into an e.m. and a nuclear parts, in spite of the fact that factorization, holding exactly in PWIA, is violated when FSI is taken into account. In the next Section the factorization approximation will be discussed within the GA and the GEA.

## A. The FSI in A(e, e'p)B processes within a diagrammatic approach

Most of the problems one faces when trying to develop a fully covariant treatment of FSI, arise because of the hadrons' spins. Therefore, let us rewrite the hadronic tensor (Eq. (2)) in the following, fully equivalent form, which however exhibits explicitly the dependence upon the spin quantum numbers

$$W_{\mu\nu}^{A} = \frac{1}{4\pi M_{A}} \sum_{\alpha_{A}} \sum_{\alpha_{A-1}, s_{1}} T_{\mu}^{\dagger}(\mathcal{M}_{A}, \mathcal{M}_{A-1}, s_{1}) T_{\nu}(\mathcal{M}_{A}, \mathcal{M}_{A-1}, s_{1}) (2\pi)^{4} \delta^{(4)}(P_{A} + q - P_{A-1} - p_{1}),$$
(13)

where  $T_{\mu}$  is a short-hand notation for the transition matrix element

$$T_{\mu}(\mathcal{M}_{A}, \mathcal{M}_{A-1}, s_{1}) \equiv \langle \alpha_{A-1} \mathbf{P}_{A-1} E^{f}_{A-1}, s_{1} \mathbf{p}_{1} | \hat{J}^{A}_{\mu}(0) | \alpha_{A} \mathbf{P}_{A} \rangle.$$
(14)

The basic assumption underlying the eikonal diagrammatic treatment of FSI at high  $Q^2$  is that the transition matrix element  $T_{\mu}$  for a nucleus A can be written in the following form

$$T_{\mu}(\mathcal{M}_{A}, \mathcal{M}_{A-1}, s_{1}) = \sum_{n=0}^{A-1} T_{\mu}^{(n)}(\mathcal{M}_{A}, \mathcal{M}_{A-1}, s_{1}),$$
(15)

where the superscript (n) corresponds to the order of rescattering of the struck particle with the A - 1 nucleons (the "spectator" nucleons), namely  $T^{(0)}_{\mu}$  corresponds to the PWIA (no rescattering),  $T^{(1)}_{\mu}$  to the single rescattering of the struck nucleons with the spectator ones,  $T^{(2)}_{\mu}$  to double rescattering, and so on. Such an approach is expected to be valid either at high energies, when particles propagate mostly in the forward direction along the direction of the three-momentum transfer  $\mathbf{q}$ , or when the momentum of the struck nucleon  $\mathbf{p}_1$  relative to A - 1 is sufficiently high; in both cases the eikonal approximation could be applied. The calculation of the rescattering part of  $T_{\mu}$  in terms of Feynman diagrams appears in principle to be a prohibitive relativistic task due, as previously stressed, to the treatment of the spin. A relevant simplification occurs if the cross section factorizes into the e.m. and the nuclear parts and, as a matter of fact, many calculations performed within the eikonal approximation treatment of the FSI, simply assume factorization. Let us try to analyze the limits of validity of such an assumption, and to this end let us consider the deuteron. In this case the Feynman diagrams describing rescattering are given in Fig. 5, and the corresponding matrix element is

$$T_{\mu}(\mathcal{M}_2, s_1, s_2) = T_{\mu}^{(0)}(\mathcal{M}_2, s_1, s_2) + T_{\mu}^{(1)}(\mathcal{M}_2, s_1, s_2),$$
(16)

where  $\mathcal{M}_2$ ,  $s_1$  and  $s_2$  are the spin projection of the deuteron, and of nucleon "1" (the active nucleon) and nucleon "2" (the spectator nucleon) in the final state. Eq. (16) obviously states that in the deuteron the interaction between the struck and the spectator nucleon can occur only via single rescattering. The cross section of the process is given by

$$\frac{d^5\sigma}{dE'd\Omega'} = \sigma_{Mott} \tilde{l}^{\mu\nu} L^D_{\mu\nu} \frac{d^3p_1}{(2\pi)^3 2E_1} \frac{d^3p_2}{(2\pi)^3 2E_2},\tag{17}$$

where  $E_i = \sqrt{\mathbf{p}_i^2 + M^2}$ , and the hadronic tensor is as follows

$$L^{D}_{\mu\nu} = \frac{1}{2M_{D}} \frac{1}{3} \sum_{\mathcal{M}_{2}, s_{1}, s_{2}} T^{\dagger}_{\mu}(\mathcal{M}_{2}, s_{1}, s_{2}) T_{\nu}(\mathcal{M}_{2}, s_{1}, s_{2}) (2\pi)^{4} \delta^{(4)} \left(P_{D} + q - p_{1} - p_{2}\right).$$
(18)

Let us now obtain the factorization of the cross section, expressed in terms of the hadronic tensor (18), within a fully covariant approach.

#### 1. The PWIA and the factorization of the cross section.

The PWIA for the process  ${}^{2}H(e, e'p)n$  within a covariant Feynman diagram approach has been considered by various authors (see e.g. [42, 43, 44]). The matrix element  $T_{\mu} = T_{\mu}^{(0)}$ , in such a case, has the following form

$$T^{(0)}_{\mu}(\mathcal{M}_{2}, s_{1}, s_{2}) = \frac{1}{2M_{N}} \sum_{\tilde{s}_{1}} J^{eN}_{\mu}(Q^{2}, p_{1}, k_{1}, \tilde{s}_{1}, s_{1}) \left[ \bar{u}(\mathbf{k}_{1}, \tilde{s}_{1}) \Phi^{\mathcal{M}_{2}}_{D}(k_{1}, k_{2})(\hat{k}_{2} + M_{N})v(\mathbf{k}_{2}, s_{2}) \right], \quad (19)$$

where

$$J^{eN}_{\mu}(Q^2, p_1, k_1, \tilde{s_1}, s_1) = \langle \mathbf{p}_1, s_1 | \Gamma^{\gamma^*N}_{\mu}(Q^2, k_1^2) | \mathbf{k}_1, \tilde{s_1} \rangle,$$
(20)

 $\Gamma^{\gamma^*N}_{\mu}(Q^2, k_1^2)$  is the e.m. vertex, and  $\Phi^{\mathcal{M}_2}_D(k_1, k_2)$  the covariant deuteron amplitude corresponding to the  $D \to (pn)$  vertex. The explicit form of the amplitude  $\Phi^{\mathcal{M}_2}_D(k_1, k_2)$  depends upon the specific covariant model used to describe the deuteron and could be found elsewhere (see e.g., [42, 43, 45]). Here, without loss of generality, we will use the Bethe-Salpeter (BS) formalism according to Refs. [45] and [44].

When Eq. (19) is placed in Eq. (18), the e.m. and nuclear parts gets coupled by the summation over the intermediate spins  $\tilde{s}_1$  and  $\tilde{s}_1'$ . However, it can be shown (see Appendix B) that the square of the expression in brackets in Eq.(19) after summation over  $\mathcal{M}_2$  and  $s_2$  yields a  $\delta$  function  $\delta_{\tilde{s}_1\tilde{s}_1'}$ , i.e. becomes diagonal in  $\tilde{s}_1$ ; this leads to the decoupling between the e.m and the nuclear parts in Eq.(19), with the resulting hadronic tensor given by

$$L_{\mu\nu}^{D} = \frac{1}{2M_{D}} \frac{1}{3} \sum_{\mathcal{M}_{2}, s_{1}, s_{2}} T_{\mu}^{\dagger}(\mathcal{M}_{2}, s_{1}, s_{2}) T_{\nu}(\mathcal{M}_{2}, s_{1}, s_{2}) = = 2M_{D} \left( 2E_{\mathbf{p}_{1}} 2E_{\mathbf{k}_{1}} L_{\mu\nu}^{eN}(Q^{2}, p_{1}, k_{1}) \right) n_{D}(|\mathbf{k}_{1}|) (2\pi)^{4} \delta^{(4)} \left( P_{D} + q - p_{1} - p_{2} \right).$$
(21)

In Eq. (21),  $n_D$  is the deuteron momentum distribution given by

$$n_D(|\mathbf{k}_1|) = \frac{1}{3} \sum_{\mathcal{M}_2, \tilde{s}_1, s_2} \left| \left[ \bar{u}(\mathbf{k}_1, \tilde{s}_1) \Phi_D^{\mathcal{M}_2}(k_1, k_2) (\hat{k}_2 + M_N) v(\mathbf{k}_2, s_2) \right] \right|^2 = \frac{1}{3} \sum_{\mathcal{M}_2, \tilde{s}_1, s_2} \left| \left\langle \tilde{s}_1, s_2 | \Psi_D^{\mathcal{M}_2}(\mathbf{k}_1) \right\rangle \right|^2 = \frac{1}{2\pi^2} \left( u_S^2(|\mathbf{k}_1|) + u_D^2(|\mathbf{k}_1|) \right), \quad (22)$$

where the (covariant) deuteron wave function has been cast in a form similar to the non relativistic one with the scalar parts of the wave function,  $u_L(|\mathbf{k}|)$ 's, related to the corresponding vertex functions,  $G_L(k_1^2, k_2^2 = M_N^2)$ , by a well known definition (see Eqs. (B8) and (B9)) leading to

$$u_L(|\mathbf{k}|) \sim \sqrt{2M_N} \frac{G_L(|\mathbf{k}|, k_{10} = M_D - E_{\mathbf{k}})}{k^2 - M_N^2}.$$
(23)

Placing Eq. (21) in Eq. (17) the well known factorized form for the cross section is obtained, *viz.* 

$$\frac{d^6\sigma}{dE'd\Omega'd\mathbf{p}_m} = K(Q^2, x, \mathbf{p}_m) \,\sigma^{eN}(\bar{Q}^2, \mathbf{p}_m) n_D(|\mathbf{k}_1|)) \delta(q_0 + M_D - E_{\mathbf{k}_1 + \mathbf{q}} - E_{\mathbf{k}_1}).$$
(24)

We reiterate that factorization has been obtained because the sum over  $s_2$  and  $\mathcal{M}_2$  in (18) leads to the appearance of a delta function  $\delta_{\tilde{s}'_1,\tilde{s}_1}$ , which means, in turn, that the square of  $T^{(0)}_{\mu}$  becomes diagonal in  $\tilde{s}_1$ . This particular (exact) result is part of a more general assertion that within the PWIA the nuclear Spectral Function is always diagonal in spins [35]. Let us now consider FSI; in this case the tensor (18) is off-diagonal in spins and factorization does not occur. However, we will show that under certain kinematical conditions, satisfied to a large extent by the GA and GEA, factorization can be recovered.

#### 2. FSI: the single scattering contribution and factorization of the cross section.

Let us compute the second diagram of Fig. 5. To this end, we introduce a two-nucleon scattering operator  $\hat{T}$  in terms of which the elastic scattering amplitude  $f^{NN}$ , describing the elastic scattering of two on-shell nucleons, will be defined as follows

$$f_{\tilde{s}_1\tilde{s}_2;s_1,s_2}^{NN}(\mathbf{p}_1,\mathbf{p}_2;\mathbf{k}_1,\mathbf{k}_2) = \bar{u}(\mathbf{p}_1,s_1)\bar{u}(\mathbf{p}_2,s_2) \ \hat{T} \ u(\mathbf{k}_1,\tilde{s}_1)u(\mathbf{k}_2,\tilde{s}_2),$$
(25)

which is obviously the free *NN* scattering amplitude; for a bound nucleon one has in principle to consider off-shell effects but in the GEA no virtuality is considered; this could be done for example by the approach of Ref. [14], by introducing cut-off form factors in the corresponding nucleon lines, which formally leads to two Feynman diagrams with different "nucleonic" masses. In presence of FSI, the transition matrix element is

$$T_{\mu}(\mathcal{M}_2, s_1, s_2) = T_{\mu}^{(0)}(\mathcal{M}_2, s_1, s_2) + T_{\mu}^{(1)}(\mathcal{M}_2, s_1, s_2)$$
(26)

with  $T^{(0)}_{\mu}$  given again by Eq. (19), and  $T^{(1)}_{\mu}$  given by the following form (note that henceforth we always have  $\mathbf{k}_1 = -\mathbf{k}_2$ )

$$T_{\mu}^{(1)}(\mathcal{M}_{2}, s_{1}, s_{2}) = \frac{1}{2M_{N}} \sum_{\tilde{s}_{1}\tilde{s}_{1}'\tilde{s}_{2}} \int \frac{d^{4}k_{2}}{i(2\pi)^{4}} \frac{f_{\tilde{s}_{1}'\tilde{s}_{2};s_{1},s_{2}}^{NN}(\mathbf{p}_{1}, \mathbf{p}_{2}, \mathbf{k}_{1}', \mathbf{k}_{2})}{k_{1}'^{2} - M_{N}^{2} + i\varepsilon} \times \left[\bar{u}(\mathbf{k}_{1}', \tilde{s}_{1}')\Gamma_{\mu}^{\gamma^{*}N}(Q^{2}, k_{1}'^{2})u(\mathbf{k}_{1}, \tilde{s}_{1})\right] \left[\bar{u}(\mathbf{k}_{1}, \tilde{s}_{1})\Phi_{D}^{\mathcal{M}_{2}}(k_{1}, k_{2})v(\mathbf{k}_{2}, \tilde{s}_{2})\right].$$
(27)

The full matrix element will therefore be

$$T_{\mu}(\mathcal{M}_{2}, s_{1}, s_{2}) = \frac{1}{2M_{N}} \sum_{\tilde{s}_{1}} J_{\mu}^{eN}(Q^{2}, \mathbf{p}_{1}, \mathbf{p}_{m}, \tilde{s}_{1}, s_{1}) \left[ \bar{u}(\mathbf{k}_{1}, \tilde{s}_{1}) \Phi_{D}^{\mathcal{M}_{2}}(k_{1}, k_{2})(\hat{k}_{2} + M_{N})v(\mathbf{k}_{2}, s_{2}) \right] +$$

$$+ \frac{1}{2M_N} \sum_{\tilde{s}_1 \tilde{s}_1' \tilde{s}_2} \int \frac{d^4 k_2}{i(2\pi)^4} \frac{f_{\tilde{s}_1' \tilde{s}_2; s_1, s_2}^{NN}(\mathbf{p}_1, \mathbf{p}_2, \mathbf{k}_1', \mathbf{k}_2)}{k_1'^2 - M_N^2 + i\varepsilon} \times \left[ \bar{u}(\mathbf{k}_1', \tilde{s}_1') \Gamma_{\mu}^{\gamma^* N}(Q^2, k_1'^2) u(\mathbf{k}_1, \tilde{s}_1) \right] \left[ \bar{u}(\mathbf{k}_1, \tilde{s}_1) \Phi_D^{\mathcal{M}_2}(k_1, k_2) v(\mathbf{k}_2, \tilde{s}_2) \right].$$
(28)

When Eq. (28) is placed into Eq. (18), the resulting equation is not diagonal in the spin quantum numbers and factorization does not hold. Let us however consider the basic assumptions underlying the eikonal multiple scattering approach, viz.:

1. the momentum transfer  $\kappa$  in the elastic rescattering is small and mostly transverse i.e.

$$\boldsymbol{\kappa} = \mathbf{p}_1 - \mathbf{k}_1' = \mathbf{k}_2 - \mathbf{p}_2 \simeq \mathbf{k}_{2\perp} - \mathbf{p}_{2\perp} = \boldsymbol{\kappa}_\perp$$
(29)

2. the spin-flip part of the NN amplitude is very small, which means that, taking into account Point 1, one can write

$$f_{\tilde{s}_{1}'\tilde{s}_{2};s_{1},s_{2}}^{NN}(\mathbf{p}_{1},\mathbf{p}_{2},\mathbf{k}_{1}',\mathbf{k}_{2}) \approx \delta_{\tilde{s}_{1}',s_{1}}\delta_{\tilde{s}_{2},s_{2}}f^{NN}(\boldsymbol{\kappa}_{\perp})$$
(30)

which is realized either at high values of the three-momentum transfer  $\mathbf{q}$ , or at high values of the momentum  $\mathbf{p}_1$  of the struck nucleon relative to the A - 1 spectator nucleons.

If the above conditions are satisfied, Eq. (27) assumes the following form (cf. Appendix B)

$$T_{\mu}^{(1)}(\mathcal{M}_{2}, s_{1}, s_{2}) \simeq \sum_{\tilde{s}_{1}} J_{\nu}^{eN}(Q^{2}, \mathbf{p}_{m}, \mathbf{p}_{1}, \tilde{s}_{1}, s_{1}) \frac{1}{2M_{N}} \int \frac{d^{4}k_{2}}{i(2\pi)^{4}} \frac{f^{NN}(\boldsymbol{\kappa}_{\perp})}{k_{1}^{\prime 2} - M_{N}^{2} + i\varepsilon} \times \\ \times \left[ \bar{u}(\mathbf{k}_{1}, \tilde{s}_{1}) \Phi_{D}^{\mathcal{M}_{2}}(k_{1}, k_{2}) v(\mathbf{k}_{2}, s_{2}) \right]$$
(31)

and one can write

$$T_{\mu}(\mathcal{M}_{2}, s_{1}, s_{2}) \simeq \frac{1}{2M_{N}} \sum_{\tilde{s}_{1}} J_{\nu}^{eN}(Q^{2}, \mathbf{p}_{m}, \mathbf{p}_{1}, \tilde{s}_{1}, s_{1}) \times \\ \times \left\{ \left[ \bar{u}(\mathbf{k}_{1}, \tilde{s}_{1}) \Phi_{D}^{\mathcal{M}_{2}}(k_{1}, k_{2}) (\hat{k}_{2} + M_{N}) v(\mathbf{k}_{2}, s_{2}) \right] + \right. \\ \left. + \int \frac{d^{4}k_{2}}{i(2\pi)^{4}} \frac{f^{NN}(\boldsymbol{\kappa}_{\perp})}{k_{1}^{\prime 2} - M_{N}^{2} + i\varepsilon} \left[ \bar{u}(\mathbf{k}_{1}, \tilde{s}_{1}) \Phi_{D}^{\mathcal{M}_{2}}(k_{1}, k_{2}) v(\mathbf{k}_{2}, s_{2}) \right] \right\}.$$
(32)

It can be seen that  $T^{(0)}_{\mu}$  and  $T^{(1)}_{\mu}$  in Eq. (32) have very similar structures, except that in  $T^{(1)}_{\mu}$  the vector  $\mathbf{k}_2$  is now an integration variable, since  $\mathbf{k}_2 \neq \mathbf{p}_2$ . When Eq. (18) is evaluated, with  $T_{\mu}$  given by Eq. (32) and assuming soft NN rescattering ( low values of  $\boldsymbol{\kappa}_{\perp}$ ), the main

contribution in the integral over  $\mathbf{k}_2$  results from the region where  $\mathbf{k}_2 \sim \mathbf{p}_2$  and this, in turn, originates again a delta function  $\delta_{\tilde{s}_1 s_1}$  (See Appendix B) and the hadronic tensor becomes

$$L_{\mu\nu}^{D} = \frac{1}{2M_{D}} \frac{1}{3} \sum_{\mathcal{M}_{2}, s_{1}, s_{2}} T_{\mu}^{\dagger}(\mathcal{M}_{2}, s_{1}, s_{2}) \cdot T_{\nu}(\mathcal{M}_{2}, s_{1}, s_{2})(2\pi)^{4} \delta^{(4)}(P_{D} + q - p_{1} - p_{2}) \simeq$$

$$\simeq \frac{1}{2} \sum_{\tilde{s}_{2}, s_{1}} \left[ J_{\mu}^{eN\dagger}(Q^{2}, \mathbf{p}_{m}, \mathbf{p}_{1}, \tilde{s}_{2}, s_{1}) \cdot J_{\nu}^{eN}(Q^{2}, \mathbf{p}_{m}, \mathbf{p}_{1}, \tilde{s}_{2}, s_{1}) \right] \times$$

$$\times \frac{1}{(2M_{N})^{2}} \sum_{\mathcal{M}_{2}\tilde{s}_{1}, s_{2}} \left| \left[ \bar{u}(\mathbf{k}_{2}, \tilde{s}_{1}) \Phi_{D}^{\mathcal{M}_{2}}(k_{1}, k_{2})(\hat{k}_{2} + M_{N})v(\mathbf{k}_{2}, s_{2}) \right]_{k_{2} = p_{2}} +$$

$$+ \int \frac{d^{4}k_{2}}{i(2\pi)^{4}} \frac{f^{NN}(\boldsymbol{\kappa}_{\perp})}{k_{1}^{\prime 2} - M_{N}^{2} + i\varepsilon} \left[ \bar{u}(\mathbf{k}_{2}, \tilde{s}_{1}) \Phi_{D}^{\mathcal{M}_{2}}(k_{1}, k_{2})v(\mathbf{k}_{2}, s_{2}) \right] \right|^{2} \times$$

$$\times (2\pi)^{4} \delta^{(4)}(P_{D} + q - p_{1} - p_{2}) \qquad (33)$$

and the factorization of the e.m. and the nuclear parts is recovered. Eq. (33) could be written in a more familiar form if one integrates over  $k_{20}$  by taking into account the pole in the amplitude  $\Phi_D^{\mathcal{M}_2}(k_1, k_2)$  ( $k_{20} = E_{\mathbf{k}}$ ) and neglecting the pole from the active propagator, which is located at large values of  $k_{20}$  and does not contribute to the integral. Using (22)-(23) one obtains  $(\int d^4k/[i(2\pi)^4] \rightarrow \int d^3k/[(2E_{\mathbf{k}}(2\pi)^3])$ 

$$L_{\mu\nu}^{D} \simeq 2M_{D} \left[ 2E_{\mathbf{p}_{1}} 2E_{\mathbf{p}_{m}} L_{\mu\nu}^{N}(Q^{2}, \mathbf{p}_{m}, \mathbf{p}_{1}) \right] \times \\ \times \sum_{\mathcal{M}_{2}, s_{1}, s_{2}} \left| \left\langle s_{1}, s_{2} | \Psi_{D}^{\mathcal{M}_{2}}(\mathbf{k}_{2}) \right\rangle_{k_{2}=p_{2}} + \int \frac{d^{3}k_{2}}{2E_{\mathbf{k}_{2}}(2\pi)^{3}} \frac{f^{NN}(\boldsymbol{\kappa}_{\perp})}{k_{1}^{\prime 2} - M_{N}^{2} + i\varepsilon} \left\langle s_{1}, s_{2} | \Psi_{D}^{\mathcal{M}_{2}}(\mathbf{k}_{2}) \right\rangle \right|^{2} \times \\ \times \left( 2\pi \right)^{4} \delta^{(4)} \left( P_{D} + q - p_{1} - p_{2} \right).$$
(34)

By placing the above equation in Eq. (17), one obtains

$$\frac{d^6\sigma}{dE'd\Omega'd\mathbf{p}_m} = K(Q^2, x, \mathbf{p}_m)\sigma^{eN}(\bar{Q}^2, \mathbf{p}_m)n_D^{FSI}(\mathbf{p}_m))\delta(q_0 + M_D - E_{\mathbf{p}_1+\mathbf{q}} - E_{\mathbf{p}_2}), \quad (35)$$

where the Distorted Momentum Distribution  $n_D^{FSI}$  is

$$n_{D}^{FSI}(\mathbf{p}_{m}) = \frac{1}{3} \sum_{\mathcal{M}_{2},s_{1},s_{2}} \left| \langle s_{1}, s_{2} | \Psi_{D}^{\mathcal{M}_{2}}(\mathbf{k}_{2}) \rangle_{k_{2}=p_{2}} + \int \frac{d^{3}k_{2}}{2E_{\mathbf{k}_{2}}(2\pi)^{3}} \frac{f^{NN}(\boldsymbol{\kappa}_{\perp})}{k_{1}^{\prime 2} - M_{N}^{2} + i\varepsilon} \langle s_{1}, s_{2} | \Psi_{D}^{\mathcal{M}_{2}}(\mathbf{p}_{m}) \rangle \right|^{2} = \frac{1}{3} \sum_{\mathcal{M}_{2},s_{1},s_{2}} \left| \mathcal{T}_{D}^{(0)}(\mathcal{M}_{2}, s_{1}, s_{2}) + \mathcal{T}_{D}^{(1)}(\mathcal{M}_{2}, s_{1}, s_{2}) \right|^{2}$$
(36)

and the quantities

$$\mathcal{T}_D^{(0)}(\mathcal{M}_2, s_1, s_2) = \left\langle s_1, s_2 | \Psi_D^{\mathcal{M}_2}(\mathbf{k}_2) \right\rangle \tag{37}$$

and

$$\mathcal{T}_{D}^{(1)}(\mathcal{M}_{2}, s_{1}, s_{2}) = \int \frac{d^{3}k_{2}}{2E_{\mathbf{k}_{2}}(2\pi)^{3}} \frac{f^{NN}(\boldsymbol{\kappa}_{\perp})}{k_{1}^{\prime 2} - M_{N}^{2} + i\varepsilon} \left\langle s_{1}, s_{2} | \Psi_{D}^{\mathcal{M}_{2}}(\mathbf{p}_{2}) \right\rangle$$
(38)

can be called the *reduced (Lorentz index independent) amplitudes*; in the above equations  $\Psi_D^{\mathcal{M}_2}$  is the deuteron wave function and the spin wave function refers to the two particles in the continuum.

To sum up we have shown that:

- 1. the cross section which includes FSI factorizes provided: i) the spin flip part of the NN scattering amplitude can be disregarded, which is consistent with the high energies we are considering, and ii) the momentum transfer  $\kappa$  in the NN rescattering is small and transverse, so that in the integral (31) one has  $\mathbf{k}_2 \sim \mathbf{p}_2$  or, equivalently,  $\mathbf{k}_2 \simeq \mathbf{p}_m$ ; this is a reasonable approximation, thanks to the behaviour of the elastic NN scattering amplitude, which is sharply peaked in the forward direction;
- 2. in the eikonal approximation and neglecting the spin dependence (spin-flip part) of the NN-amplitude, the FSI is not affected by the spin structure of the wave functions of the deuteron and the two-body final state. This means that in computing the Feynman diagrams, the intermediate spin algebra can be disregarded, and only the scalar part of the corresponding vertex functions can be considered, using Eq. (23) to define the scalar parts of the wave functions. Then the resulting amplitude has to be merely sandwiched between the spin functions of initial and final particles.

These conclusions can be generalized to a nucleus A, for which the cross section of the process A(e, e'p)(A-1) is given by the following expression:

$$\frac{d^6\sigma}{dE'd\Omega'd\mathbf{p}_m} = K(Q^2, x, \mathbf{p}_m) \,\sigma^{eN}(\bar{Q}^2, \mathbf{p}_m) P_A^{FSI}(\mathbf{p}_m, E_m) \tag{39}$$

where  $P_A^{FSI}(\mathbf{p}_m, E_m)$  is the Distorted Spectral Function

$$P_{A}^{FSI}(\mathbf{p}_{m}, E_{m}) = \frac{1}{(2\pi)^{3}} \frac{1}{2J_{A}+1} \sum_{f} \sum_{M_{A}, \mathcal{M}_{A-1}, s_{1}} \left| \sum_{n=0}^{A-1} \mathcal{T}_{A}^{(n)}(\mathcal{M}_{A}, \mathcal{M}_{A-1}, s_{1}) \right|^{2} \times \delta\left( E_{m} - (E_{A-1}^{f} + E_{min}) \right)$$
(40)

and *n* denotes the order of rescattering. In what follows the distorted momentum distributions for the Deuteron (Eq. (36)) and the distorted Spectral Function for  ${}^{3}He$  (Eq. (40)) will be calculated within the GA and GEA.

# B. The process ${}^{2}H(e,e'p)n$ within the GA and GEA.

Let us now calculate the reduced amplitude  $\mathcal{T}_D^{(1)}$  in the process  ${}^2H(e, e'p)n$ , taking FSI into account by the GEA. This amounts to replace the energy denominator in Eq. (38) by its generalized eikonal approximation. To this end, we will consider both the "canonical" case, when the value of the 3-momenta transfers  $|\mathbf{q}|$  is so high that  $\mathbf{q} \simeq \mathbf{p}_1$ , with the z-axis naturally directed along  $\mathbf{q}$ , as well as the case of smaller values of  $\mathbf{q}$ , but high values of  $\mathbf{p}_1$ , when  $\mathbf{q}$  and  $\mathbf{p}_1$  may point to different directions, in which case the z-axis is oriented along  $\mathbf{p}_1$ .

Remembering that  $\kappa = p_1 - k'_1 = k_2 - p_2$ , the energy denominator can be written as follows

$$k_1'^2 - M_N^2 = (p_1 - \kappa)^2 - M_N^2 = -2p_1\kappa + \kappa^2 = 2|\mathbf{p}_1| \left(\kappa_z + \frac{\kappa_0(\kappa_0 - 2E_{\mathbf{p}_1})}{2|\mathbf{p}_1|} - \frac{\kappa^2}{2|\mathbf{p}_1|}\right) \approx \\ \approx 2|\mathbf{p}_1| \left(\kappa_z - \frac{E_{\mathbf{k}_1 + \mathbf{q}} + E_{\mathbf{p}_1}}{2|\mathbf{p}_1|}\kappa_0\right) \approx 2|\mathbf{p}_1| \left(\kappa_z + \Delta_z\right),$$

$$\tag{41}$$

where

$$\Delta_z = \frac{E_{\mathbf{k}_1+\mathbf{q}} + E_{\mathbf{p}_1}}{2|\mathbf{p}_1|} \left( E_m - |E_A| \right) \tag{42}$$

and the relation

$$\kappa_0 = E_{\mathbf{p}_1} - E_{\mathbf{k}_1 + \mathbf{q}} \approx -(E_m - |E_A|) \tag{43}$$

resulting from energy conservation  $q_0 + M_D = E_{\mathbf{p}_1} + E_{\mathbf{p}_2}$  has been used.

By changing the normalization of the NN amplitude from the covariant one to the non relativistic analogue  $(E_{\mathbf{p}} \simeq M_N)$ , one has

$$\frac{f^{NN}(\boldsymbol{\kappa}_{\perp})}{4E_{\mathbf{p}}|\mathbf{p}_{1}|} \approx \frac{f^{NN}(\boldsymbol{\kappa}_{\perp})}{4M_{N}|\mathbf{p}_{1}|} = a^{NR}(\boldsymbol{\kappa}_{\perp}) \equiv i \int d^{2}\mathbf{b}e^{i\boldsymbol{\kappa}_{\perp}\mathbf{b}} \Gamma(\mathbf{b})$$
(44)

and  $\mathcal{T}_D^{(1)}$  becomes

$$\mathcal{T}_{D}^{(1)}(\mathcal{M}_{2}, s_{1}, s_{2}) = \int \frac{d^{3}k}{(2\pi)^{3}} a^{NR}(\boldsymbol{\kappa}_{\perp}) \frac{1}{\kappa_{z} + \Delta_{z} + i\varepsilon} \left\langle s_{1}, s_{2} | \Psi_{D}^{\mathcal{M}_{2}}(\mathbf{p}) \right\rangle.$$
(45)

Using

$$\frac{1}{\kappa_z + \Delta_z + i\epsilon} = -i \int \theta(z) \, \mathrm{e}^{i(\kappa_z + \Delta_z) \cdot z} dz \tag{46}$$

we obtain, in coordinate space,

$$\mathcal{T}_{D}^{(0)}(\mathcal{M}_{2},s_{1},s_{2}) + \mathcal{T}_{D}^{(1)}(\mathcal{M}_{2},s_{1},s_{2}) = \left\langle s_{1},s_{2}\left(1-\theta(z)\mathrm{e}^{i\Delta_{z}z}\Gamma(\mathbf{b})\right)\mathrm{e}^{-i\mathbf{p}_{m}\mathbf{r}}|\Psi_{D}^{\mathcal{M}_{2}}(\mathbf{r})\right\rangle.$$
(47)

As a result, the cross section will read as follows

$$\frac{d^{6}\sigma}{dE'd\Omega'd\mathbf{p}_{m}} = K(Q^{2}, x, \mathbf{p}_{m}) \,\sigma^{eN}(\bar{Q}^{2}, \mathbf{p}_{m}) n_{D}^{FSI}(\mathbf{p}_{m}) \delta(M_{D} + \nu - E_{\mathbf{p}_{1}} - E_{\mathbf{p}_{m}})$$
(48)

with the distorted momentum distributions  $n_D^{FSI}$  defined by

$$n_D^{FSI}(\mathbf{p}_m) = \frac{1}{3} \frac{1}{(2\pi)^3} \sum_{\mathcal{M}_2, S_{23}} \left| \int d\mathbf{r} \chi_{S_{23}}^{\dagger} \Psi_D^{\mathcal{M}_2 \dagger}(\mathbf{r}) \mathcal{S}_{\Delta}^{FSI}(\mathbf{r}) \exp(-i\mathbf{p}_m \mathbf{r}) \right|^2, \tag{49}$$

where  $S_{\Delta}^{FSI}(\mathbf{r})$ , which describes the final state interaction between the hit nucleon and the spectator, is

$$S_{\Delta}^{FSI}(\mathbf{r}) = 1 - \theta(z) e^{i\Delta_z z} \Gamma(\mathbf{b})$$
(50)

with  $\mathbf{r} = (\mathbf{b}, z)$ . In the above formulae the z-axis is along  $\mathbf{p}_1$ ; it should be pointed out, however, that at large values of the momentum transfer, the hit nucleon propagates almost along  $\mathbf{q}$  so that by choosing the z-axis along the three-momentum transfer and neglecting the virtuality of the struck nucleon before and after interaction, one can write [10]

$$k_1'^2 - M_N^2 = (k_1 + q)^2 - M_N^2 \approx 2|\mathbf{q}| \left(\kappa_z + \Delta_z\right),$$
(51)

where

$$\Delta_z = \frac{q_0}{|\mathbf{q}|} E_m. \tag{52}$$

It can be seen that the FSI factor (50) in the GEA differs from the one of the standard GA [46, 47, 48], simply by the additional factor  $e^{i\Delta_z z}$ . It should be pointed out that whereas the well known factor  $\theta(z)$  [47, 48] originates from the non-relativistic reduction of the covariant Feynman diagrams and guarantees the correct time ordering of the rescattering processes, the quantity  $\Delta_z$  is of a pure nuclear structure origin and, as it can be seen from Eq. (47), represents a correction to the parallel component of the missing momentum. Therefore the corrections from  $\Delta_z$  are expected to be important in parallel kinematics at  $|\mathbf{p}_z| \simeq \Delta_z$ . As we shall see from the results of our calculations performed in perpendicular kinematics in the range  $|\mathbf{p}_m| \leq 600 \ MeV/c$  and  $E_m \leq 100 \ MeV$ , one always has  $|\Delta_z| \ll |\mathbf{p}|_{\perp}$  with  $q_0/|\mathbf{q}| \simeq 1$ , so that  $\Delta_z$  is always very small. We can therefore anticipate that effects of  $\Delta_z$  on the experimental data we have considered is also very small.

# C. The processes ${}^{3}H(e,e'p){}^{2}H$ and ${}^{3}H(e,e'p)(np)$ within the GA and GEA.

Let us now consider the three-body system. The distorted Spectral Function is given by Eq. (40)

$$P_{H_e}^{FSI}(\mathbf{p}_m, E_m) = \frac{1}{(2\pi)^3} \frac{1}{2} \sum_{f} \sum_{\mathcal{M}_3, \mathcal{M}_2, s_1} \left| \sum_{n=0}^2 \mathcal{T}_A^{(n)}(\mathcal{M}_3, \mathcal{M}_2, s_1) \right|^2 \times \delta\left( E_m - (E_2^f + E_{min}) \right),$$
(53)

where the magnetic quantum number  $\mathcal{M}_2$  refers either to the deuteron or to the two nucleon in the continuum, depending upon the break-up channel we are considering  $(E_{min} = E_3 - E_2 (E_{min} = E_3)$  for the two-body (three-body) break-up channel). The diagrams representing the rescattering processes are shown in Fig. 6. The evaluation of these diagrams follows the standard procedure adopted for the deuteron. Let us illustrate it in the case of the 3bbu considering, for ease of presentation, the single scattering diagram of Fig. 6 b). After integration over  $k_{20}$  and  $k_{30}$  in the corresponding poles of the propagators of the spectators  $(k_{20} = E_{\mathbf{k}_2} \text{ and } k_{30} = E_{\mathbf{k}_3})$ , we obtain

$$\mathcal{T}_{3}^{(1)}(\mathcal{M}_{3}, s_{1}, s_{2}, s_{3}) = \int \frac{d^{3}k_{2}}{2E_{\mathbf{k}_{2}}(2\pi)^{3}} \frac{d^{3}k_{3}}{2E_{\mathbf{k}_{3}}(2\pi)^{3}} \times \frac{G_{He \to 1(23)}(k_{1}, k_{2}, k_{3}, s_{1}, s_{2}, s_{3})}{(k_{1}^{2} - M_{N}^{2})} \frac{f_{NN}(p_{1} - k_{1}')}{k_{1}'^{2} - M_{N}^{2}} \frac{G_{(23) \to f}^{+}(k_{2}', k_{3}, s_{2}, s_{3})}{(k_{2}'^{2} - M_{N}^{2})},$$
(54)

where the overlaps of the vertex functions  ${\cal G}_i$  are

$$G_{He\to 1(23)}(k_1, k_2, k_3, s_1, s_2, s_3) = \langle \mathbf{k}_1, s_1, \mathbf{k}_2, s_2, \mathbf{k}_3, s_3 | G_{He\to 1(23)}(\mathcal{M}_3, \mathbf{P}_3) \rangle;$$
(55)

$$G_{(23)\to f}(k_2, k_3, s_2, s_3) = \langle \mathbf{k}_2, s_2, \mathbf{k}_3, s_3 | G_{(23)\to f}(\mathcal{M}_{23}, S_{23}, \mathbf{P}_2, E_2^f) \rangle;$$
(56)

The vertex functions  $G_i$  are replaced by the non-relativistic overlap functions according to the general convention (we omit for ease of presentation the proper normalization factors)

$$\langle s_1, s_2, s_3 | \Psi_{He}^{\mathcal{M}_3}(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3) \rangle \approx \frac{G_{He \to 1(23)}(k_1, k_2, k_3, s_1, s_2, s_3)}{(k_1^2 - M_N^2)}$$
(57)

and, using the completeness relation when summing over  $s_2$  and  $s_3$ , one gets

$$\mathcal{T}_{3}^{(1)}(Q^{2}, s_{1}, S_{23}) = \int \frac{d^{3}k_{2}}{2E_{\mathbf{k}_{2}}(2\pi)^{3}} \frac{d^{3}k_{3}}{2E_{\mathbf{k}_{3}}(2\pi)^{3}} \times \\ \times \Psi_{(23)}^{f}(\mathbf{k}_{3}, \mathbf{k}_{2}'; S_{23}) \frac{f_{NN}(\boldsymbol{\kappa})}{(k_{1}'^{2} - M_{N}^{2} + i\epsilon)} \langle s_{1} | \Psi_{He}^{\mathcal{M}_{3}}(\mathbf{k}_{1}, \mathbf{k}_{2}, \mathbf{k}_{3}) \rangle.$$
(58)

Following the procedure adopted for the deuteron, we obtain

$$\mathcal{T}_{3}^{(1)}(Q^{2}, s_{1}, S_{23}) = \int \frac{d^{3}\kappa}{(2\pi)^{3} 2E_{\mathbf{k}_{2}}} \Psi_{(23)}^{f}(\mathbf{k}_{3}, \mathbf{k}_{2}'; S_{23}) \frac{f_{NN}(\boldsymbol{\kappa})}{k_{1}^{\prime 2} - M_{N}^{2} + i\varepsilon} \langle s_{1} | \Psi_{He}^{\mathcal{M}_{3}}(\mathbf{k}_{1}, \mathbf{k}_{2}, \mathbf{k}_{3}) \rangle \approx \\ \approx \int \frac{d^{3}\kappa}{(2\pi)^{3}} \Psi_{(23)}^{f}(\mathbf{k}_{3}, \mathbf{k}_{2}'; S_{23}) \frac{f_{NN}(\boldsymbol{\kappa})/4M_{N} |\mathbf{p}_{1}|}{(\kappa_{z} + \Delta_{z} + i\epsilon)} \langle s_{1} | \Psi_{He}^{\mathcal{M}_{3}}(\mathbf{k}_{1}, \mathbf{k}_{2}, \mathbf{k}_{3}) \rangle,$$
(59)

where

$$\Delta_z = \frac{E_{\mathbf{k}_1 + \mathbf{q}} + E_{\mathbf{p}_1}}{2|\mathbf{p}_1|} (E_m - E_3).$$
(60)

Including also the 2buu channel, we can write, in coordinate space,

$$P_{He}^{FSI}(\mathbf{p}_m, E_m) = P_{gr}^{FSI}(\mathbf{p}_m, E_m) + P_{ex}^{FSI}(\mathbf{p}_m, E_m),$$
(61)

where

$$P_{gr}^{FSI}(\mathbf{p}_m, E_m) = n_{gr}^{FSI}(\mathbf{p}_m)\delta(E_m - (E_3 - E_2))$$
(62)

with

$$n_{gr}^{FSI}(\mathbf{p}_m) = \frac{1}{(2\pi)^3} \frac{1}{2} \sum_{\mathcal{M}_3, \mathcal{M}_2, s_1} \left| \int e^{i\boldsymbol{\rho}\mathbf{p}_m} \chi_{\frac{1}{2}s_1}^{\dagger} \Psi_D^{\mathcal{M}_2\dagger}(\mathbf{r}) \mathcal{S}_{\Delta}^{FSI}(\boldsymbol{\rho}, \mathbf{r}) \Psi_{He}^{\mathcal{M}_3}(\boldsymbol{\rho}, \mathbf{r}) d\boldsymbol{\rho} d\mathbf{r} \right|^2$$
(63)

and

$$P_{ex}^{FSI}(\mathbf{p}_m, E_m) = \frac{1}{(2\pi)^3} \frac{1}{2} \sum_{\mathcal{M}_3, S_{23}, s_1} \int \frac{d^3 \mathbf{t}}{(2\pi)^3} \left| \int e^{i\boldsymbol{\rho}\mathbf{p}_m} \chi_{\frac{1}{2}s_1}^{\dagger} \Psi_{np}^{\mathbf{t}\dagger}(\mathbf{r}) \mathcal{S}_{\Delta}^{FSI}(\boldsymbol{\rho}, \mathbf{r}) \Psi_{He}^{\mathcal{M}_3}(\boldsymbol{\rho}, \mathbf{r}) d\boldsymbol{\rho} d\mathbf{r} \right|^2 \times \delta \left( E_m - \frac{\mathbf{t}^2}{M_N} - E_3 \right).$$
(64)

The FSI factor  $S_{\Delta}^{FSI}$  describes the single and double rescattering of nucleon "1" with the spectators "2" and "3", and has the following form

$$\mathcal{S}_{\Delta}^{FSI}(\boldsymbol{\rho}, \mathbf{r}) = \mathcal{S}_{(1)}^{FSI}(\boldsymbol{\rho}, \mathbf{r}) + \mathcal{S}_{(2)}^{FSI}(\boldsymbol{\rho}, \mathbf{r})$$
(65)

with the single scattering contribution  $\mathcal{S}_{(1)}^{FSI}$  given by

$$\mathcal{S}_{(1)}^{FSI}(\boldsymbol{\rho}, \mathbf{r}) = 1 - \sum_{i=2}^{3} \theta(z_i - z_1) \mathrm{e}^{i\Delta_z(z_i - z_1)} \Gamma(\mathbf{b}_1 - \mathbf{b}_i)$$
(66)

and the double scattering contribution by (Ref. [10, 12]).

$$\mathcal{S}_{(2)}^{FSI}(\boldsymbol{\rho}, \mathbf{r}) = \left[\theta(z_2 - z_1)\theta(z_3 - z_2)e^{-i\Delta_3(z_2 - z_1)}e^{-i(\Delta_3 - \Delta_z)(z_3 - z_1)} + \theta(z_3 - z_1)\theta(z_2 - z_3)e^{-i\Delta_2(z_3 - z_1)}e^{-i(\Delta_2 - \Delta_z)(z_2 - z_1)}\right] \times \Gamma(\mathbf{b}_1 - \mathbf{b}_2)\Gamma(\mathbf{b}_1 - \mathbf{b}_3), \quad (67)$$

where  $\Delta_i = (q_0/|\mathbf{q}|)(E_{\mathbf{p}_i} - E_{\mathbf{k}'_i})$  and  $\Delta_z$  is given by Eq. (52).

When  $\Delta_z = 0$ , the familiar form for  $\mathcal{S}^{FSI}$  is obtained, namely

$$\mathcal{S}^{FSI}(\boldsymbol{\rho}, \mathbf{r}) = \prod_{i=2}^{3} \left[ 1 - \theta(z_i - z_1) \, \Gamma(\mathbf{b}_i - \mathbf{b}_1) \right], \tag{68}$$

and when  $\Gamma = 0$ , the distorted Spectral Function (61) transforms into the usual Spectral Function (9).

Using Eq. (61), the cross section of the process  ${}^{3}He(e,e'p)X$  (X = D or (np)) assumes the following form

$$\frac{d^6\sigma}{dE'd\Omega'd\mathbf{p}_m} = K(Q^2, x, \mathbf{p}_m) \,\sigma^{eN}(\bar{Q}^2, \mathbf{p}_m) P_{He}^{FSI}(\mathbf{p}_m, E_m).$$
(69)

## V. RESULTS OF THE CALCULATIONS

We have used Eqs. (48), (50), (69) and (67) to calculate the cross sections of the processes  ${}^{2}He(e, e'p)n$ ,  ${}^{3}He(e, e'p){}^{2}H$  and  ${}^{3}He(e, e'p)(np)$ . All calculations have been performed using the following well known parametrization of the profile function  $\Gamma(\mathbf{b})$ 

$$\Gamma(\mathbf{b}) = \frac{\sigma_{NN}^{tot} (1 - i\alpha_{NN})}{4\pi b_0^2} e^{-\mathbf{b}^2/2b_0^2},$$
(70)

where  $\sigma_{NN}^{tot}$  is the total NN cross section,  $\alpha_{NN}$  the ratio of the real to imaginary part of the forward NN amplitude, and  $b_0$  the slope of the differential elastic NN cross section. The values of the energy dependent quantities  $\sigma_{NN}^{tot}$  and  $\alpha_{NN}$  have been taken from Ref. [49]. For the electron-nucleon cross section  $\sigma^{eN}(\bar{Q}^2, \mathbf{p}_m)$  we used the De Forest  $\sigma_{cc1}^{eN}(\bar{Q}^2, \mathbf{p}_m)$  cross section [35]. All two- and three-body wave functions are direct solutions of the non relativistic Schrödinger equation, therefore our calculations are fully parameter free.

Calculations have been performed in PWIA and including the full rescattering within the GA and GEA by evaluating the Feynman diagrams shown in Figs. 5 and 6. It should be pointed out that, apart from minor differences (e.g. the structure of  $\Delta_z$  for complex nuclei) which do not affect the numerical results, our GEA is essentially the same as the one developed in [10, 12].

# A. The process ${}^{2}H(e,e'p)n$

Our results for the process  ${}^{2}H(e, e'p)n$  are compared in Figs. 7, 8 and 9 with three different sets of experimental data, covering different kinematical ranges, namely the experimental data from NIKHEF [17], SLAC [19], and Jlab [18]. The relevant kinematical variables in the three experiments are as follows: i)  $0.1 \leq Q^{2} \leq 0.3$ ,  $0.3 \leq x \leq 0.6$  [17]; ii)  $1.2 \leq Q^{2} \leq 6.8$ ,  $x \simeq 1$  [19]; iii)  $Q^{2} \simeq 0.665$   $(GeV/c)^{2}$ ,  $|\mathbf{q}| \simeq 0.7$  GeV/c,  $x \simeq 0.96$  [18]. In Figs. 7 and 9 the theoretical cross section corresponding to Eq. (48), namely

$$\frac{d^5\sigma}{dE'd\Omega'd\Omega_{\mathbf{p}_m}} = f_{rec} \,\mathcal{K}(Q^2, x, \mathbf{p}_m) \,\sigma_{cc1}^{eN}(\bar{Q}^2, \mathbf{p}_m) n_D^{FSI}(\mathbf{p}_m) \tag{71}$$

is compared with the corresponding data, whereas in Fig. 8 we compare, as in Ref. [18], the effective momentum distributions  $N_{eff}(p_m)$  (or reduced cross section) defined by [18]

$$N_{eff}(|\mathbf{p}_m|) = \frac{d^5 \sigma^{exp}}{d\Omega' dE' d\Omega_{\mathbf{p}_m}} \left[ f_{rec} \ \mathcal{K} \ \sigma^{eN}_{cc1} \right]^{-1}, \tag{72}$$

where in Eqs. (71) and (72)  $f_{rec}$  and  $\mathcal{K}$  are kinematical factors which arise from the integration over  $dT_{\mathbf{p}_1}$ .

The results presented in Figs. 7, 8 and 9 exhibit a general satisfactory agreement between theoretical calculations and experimental data, particularly in view of the wide range of kinematics covered by the data we have considered. Figs. 8 and 9 show however that quantitative disagreements with data exist in some regions. Particularly worth being noted is the disagreement in the region around  $|\mathbf{p}_m| \simeq 0.25 \ GeV/c$  appearing in Fig. 8. We did not try to remove such a disagreement by adjusting the quantities entering the profile function (48), but it turns out that  $n_D^{FSI}$  in the region around  $|\mathbf{p}_m| \simeq 0.25 \ GeV/c$  is rather sensitive to the value of  $\alpha_{NN}$ . The NIKHEF kinematics deserves a particular comment. As a matter of fact, the four momentum transfer in this experiment is rather low and one might rise doubts as to the validity of the eikonal approximation. In this respect, it should however be pointed out, that what really matters in GA and GEA is the relative three-momentum of the hit nucleon with respect to the A - 1 system; in the NIKHEF experiment, due to the large value of the energy transfer, the three momentum transfer is also large, and  $\gamma^*$ absorption occurs on a proton moving along **q**, with the recoiling neutron moving with low momentum against **q**; the resulting proton-neutron relative momentum is of the order of few hundreds MeV/c, which, though representing the lower limit for the validity of the eikonal approximation, still appears, according to our results, to be suitable for the application of the GEA: as a matter of fact our results appear to be in reasonable agreement with the ones obtained within approaches which are better justified at low energies, like, e.g., the ones presented, in Refs. [50, 51, 52, 53, 54].

Let us conclude this Section by stressing that, as far as the effects of MEC and  $\Delta$  isobar excitations are concerned, these have been found to be very small ( $\simeq 5 - 6\%$ ) in the SLAC kinematics (see [19]), with the results from [55] exhibiting the same trend also in the Jlab kinematics. Eventually, we should remark that the results of the GA and GEA differ by only few percent and cannot be distinguished in the Figures.

# **B.** The processes ${}^{3}He(e, e'p){}^{2}H$ and ${}^{3}He(e, e'p)(np)$ .

Calculations for the three-body systems are very involved, mainly because of the complex structure of the wave function of Ref. [2], which is given in a mixed  $(L_{\rho}, X, j_{23}, S_{23})$ representation, including angular momentum values up to  $L_{\rho} = 7$  and  $j_{23} = 8$  (a total of 58 configurations with different combinations of  $(L_{\rho}, X, j_{23}, S_{23})$  quantum numbers). Correspondingly, the wave function of the spectators (the deuteron or the continuum two-nucleon states) is given in a *JLS*-scheme (see Appendix A). We would like to stress, that no approximations have been made in the evaluation of the single and double scattering contributions to the FSI: proper intrinsic coordinates have been used and the energy dependence of the profile function has been taken into account in the properly chosen CM system of the interacting pair. The Feynman diagrams which have to be evaluated, both for the 2bbu and 3bbu channels are shown in Fig. 6.

# 1. The two-body break-up channel ${}^{3}He(e, e'p)^{2}H$ .

The 2bbu channel cross section

$$\frac{d^5\sigma}{dE'd\Omega'd\Omega_{\mathbf{p}_1}} = K_{2bbu}(Q^2, x, \mathbf{p}_m) \,\sigma_{cc1}^{eN}(\bar{Q}^2, \mathbf{p}_m) n_{gr}^{FSI}(\mathbf{p}_m)$$
(73)

obtained from Eq. (69), with  $n_{qr}^{FSI}(\mathbf{p}_m)$  given by Eq. (63), is compared in Fig. 10 with recent experimental data from Jlab Collaboration [21]. The relevant kinematical variables in the experiment are  $|\mathbf{q}| = 1.5 \ GeV/c, \ q_0 = 0.84 \ GeV, \ Q^2 = 1.55 \ (GeV/c)^2$ , and  $x \approx 1$ . The cross section is presented as a function of the missing momentum  $|\mathbf{p}_m|$  (which, for the  ${}^{3}He(e, e'p)D$ -process, exactly coincides with the final deuteron momentum). In PWIA the cross section is directly proportional to  $n_{gr}$  (Eq. (11)) shown in the left panel of Fig. 3. It can be seen that up to  $|\mathbf{p}_m| \sim 400 \ MeV/c$ , the PWIA and FSI results are almost the same and fairly well agree with the experimental data, which means, in turn, that the 2bbu  ${}^{3}He(e, e'p)^{2}H$  does provide information on  $n_{gr}$ ; on the contrary, at larger values of  $|\mathbf{p}_m| \ge 400 \ MeV/c$  the PWIA appreciably underestimates the experimental data. It is very gratifying to see that when FSI is taken into account, the disagreement is fully removed and an overall very good agreement between theoretical predictions and experimental data is obtained. It should be pointed out that the experimental data shown in Fig. 10 correspond to the perpendicular kinematics, when the deuteron momentum (the missing momentum) is always almost perpendicular to the momentum transfer  $\mathbf{q}$ ; in such a kinematics the effects from FSI are maximized, whereas in the so called parallel kinematics, they are minimized (see, e.g. [9], [48], [56]). The kinematics therefore reflects itself in the relevance of the calculated FSI; as a matter of fact, we have found that the effects of the FSI calculated either within the GA or GEA approximations, differ only by a few percent, which was expected in view of the observation that the factor  $\Delta_z$  (Eqs. (60) or (52)) affects only the longitudinal component of  $\mathbf{p}_m$  and therefore has minor effects on the data we have considered. The effects of MEC and  $\Delta$  isobar contributions have been estimated in [55] and found negligible up to about  $p_m \simeq 600 MeV/c$ .

# 2. The three-body break-up channel ${}^{3}He(e, e'p)(np)$ .

From Eq. (69), we obtain the cross section for the 3bbu in the following form

$$\frac{d^6\sigma}{dE'd\Omega'd\Omega_{\mathbf{p}_1}dE_m} = K_{3bbu}(Q^2, x, \mathbf{p}_m) \,\sigma_{cc1}^{eN}(\bar{Q}^2, \mathbf{p}_m) P_{ex}^{FSI}(\mathbf{p}_m, E_m),\tag{74}$$

where  $P_{ex}^{FSI}(\mathbf{p}_m, E_m)$  is given by Eq. (64). We have calculated Eq. (74) in correspondence of two different kinematical ranges: the one from Ref. [20] and the one corresponding to the experimental data from Jlab [22]. Contrary to the 2bbu channel, the 3bbu cross section depends upon an extra kinematical variable, the removal energy  $E_m$ , and corresponds to the process in which three particles interact in the continuum. We have considered three different theoretical approaches, namely:

- the Plane Wave Approximation (PWA), when FSI effects are completely ignored, i.
   e. the three particles in the continuum are described by plane waves;
- the Plane Wave Impulse Approximation (PWIA), in which the struck nucleon is described in the continuum by a plane wave and the spectator pair is described by the continuum solution of the Schrödinger equation (obviously, in the case of the deuteron the PWIA coincides with the PWA);
- 3. the full FSI, when the struck nucleon interacts in the continuum with the nucleons of the spectator pair via the standard GA or the more refined GEA.

In Fig. 11 the results of our calculations are compared with the experimental data from Ref. [20]. In the experiment, which corresponds to a relatively low beam energy  $(E = 0.560 \ GeV)$ , the scattering angle  $(\theta_e = 25^{\circ})$  and the energy transfer  $(q_0 = 0.32 GeV)$ were kept constant, and protons with different values of the missing momentum and energy were detected in correspondence of several values of the proton emission angle  $\theta_{\mathbf{p}_1}$ ,  $viz \ \theta_{\mathbf{p}_1} = 45^{\circ}, 60^{\circ}, 90.5^{\circ}, 112^{\circ}$  and  $142.5^{\circ}$ . The kinematics is far from the quasi elastic peak  $(x \simeq 0.1)$  and the values of the four- and three-momentum transfers are low  $(Q^2 \simeq 0.03 \ (GeV/c)^2$  and  $|\mathbf{q}| = 0.28 \ GeV/c)$ . At first glance this would invalidate the use of the eikonal approximation; however, a detailed analysis of the kinematics, shows that the value of both  $\mathbf{p}_1$  and  $\mathbf{p}_m$  are rather large  $(400 - 600 \ MeV/c)$ , and so is the value of the angle between them  $(\theta_{\widehat{\mathbf{p}_1\mathbf{p}_m}} \sim 150^{\circ})$ ; thus the momentum of the struck nucleon relative to the spectator pair is high enough to make the use of the eikonal approximation justified. Moreover, the values of the experimentally measured missing momenta and missing energy at each value of  $\theta_{\mathbf{p_1}}$ , always cover the kinematical range where the condition for two nucleon correlation mechanism  $E_m \sim \frac{\mathbf{p}_m^2}{4M_N}$  holds; as a matter of fact, as it can be seen from Fig. 11, the position of the bumps in the cross section are reasonably predicted by the PWA and PWIA.

The results presented in Fig. 11 clearly show that with increasing missing momentum, the experimental peak moves to higher values of missing energy, in qualitative agreement with the two-nucleon correlation mechanism. More important, it can be seen that at the highest value of  $|\mathbf{p}_m|$  ( $\theta_{\mathbf{p}_1} = 112^o$ ) the effects of FSI, both in the spectator pair and between the struck nucleon and the spectator pair, is very small. The reason for such a behaviour is as follows: the kinematics of the experiment is not purely perpendicular: the relation between  $|\mathbf{p}_{m\perp}|$  and  $|\mathbf{p}_m|$  is such that  $|\mathbf{p}_{m\perp}| \sim \frac{1}{2}|\mathbf{p}_m|$ , so that the dominant role played by FSI in the purely perpendicular kinematics is decreased with increased values of  $\theta_{\mathbf{p}_1}$ .

In Fig. 12 our results are compared with the recent data from Jlab [22], where the cross section was measured at fixed values  $|\mathbf{p}_m| vs$ . the missing energy  $E_m$ . As in the case of the Saclay data previously analyzed, even in this case the cross section exhibits bumps approximately located at values of  $E_m$  and  $|\mathbf{p}_m|$  satisfying the two-nucleon correlation mechanism relation ( $E_m \sim \frac{\mathbf{p}_m^2}{4M_N}$ ), and in agreement with the behaviour of the Spectral Function (see Figs. 3 and 4). However, contrary to the Saclay case, the PWIA dramatically underestimates the experimental data. This is clear evidence that the FSI between the struck nucleon and the nucleons of the spectator pair (Feynman diagrams b) and c) in Fig. 6) does play a relevant role, as the results of our calculations (the full line in Fig. 12) do indeed really show. Since, as already stressed, the Jlab experiment correspond to a perpendicular kinematics, this explains the larger effects of the FSI with respect to the Saclay experiment. The effects of the FSI calculated either within the GA or GEA approximations, differ only by a few percent, which was expected in view of the observation that the factor  $\Delta_z$  (Eqs. (60) or (52)) affects only the longitudinal component of  $\mathbf{p}_m$  and therefore has minor effects on the data we have considered.

There exist at present only two approaches to the calculations of the 2bbu and 3bbu channels at the Jlab kinematics: the one presented in this paper and the one by Laget reported in Refs. [21], [22] and [23]. A comparison of the results of the two approaches exhibits an encouraging agreement both in the 2bbu and 3bbu channels, with some minor differences which should most likely be ascribed to the different wave functions used in the two calculations. It is therefore gratifying to observe that different approximations to the treatment of FSI lead to very similar results.

The effects of MEC and  $\Delta$ , as previously pointed out, have not yet been considered in our approach; the calculation of Ref. [23], shows they reduce the cross section in the peak by about 10%, leaving the missing energy dependence and, consequently, our conclusions, practically unchanged.

#### VI. SUMMARY AND CONCLUSIONS

We have calculated the cross section of the processes  ${}^{2}H(e,e'p)n$ ,  ${}^{3}He(e,e'p)D$ , and  ${}^{3}He(e, e'p)(np)$ , using realistic wave functions for the ground state, which exhibits the very rich correlation structure generated by modern NN interactions; the FSI of the struck nucleon with the spectators has been treated within the standard Glauber eikonal approximation (GA) [16], as well as with its generalized version (GEA) [9, 10, 11]. The two approaches differ by a factor  $\Delta_z$  (Eqs. (42) and (52)) which modifies (see Eq. (65)) the FSI factor appearing in the standard GA (Eq. (68)). This factor takes into account in the NN scattering amplitude the removal energy of the struck nucleon, or, equivalently, the excitation energy of the system A - 1. By properly choosing the z-axis (along **q** or **p**<sub>1</sub>), we were able to calculate FSI effects either in the case of large values of the three-momentum transfer  $\mathbf{q}$ , or large values of the momentum of the struck nucleon  $\mathbf{p}_1$  relative to the A-1 system; by this way calculations could be extended successfully even at relatively low values of  $Q^2$ . As far as the three-body break-up channel in  ${}^{3}He$  is concerned, the FSI in the spectator pair was always calculated by the solution of the Schrödinger equation, whereas the interaction of the active, fast nucleon with the two nucleons of the spectator pair has been taken care of by the GA or GEA approximations. The method we have used is a very transparent one and fully parameter free: it is based upon Eqs. (49), (61), and (65, which only require the knowledge of the nuclear wave functions, since the FSI factor is fixed directly by NN scattering data. Of course with increasing A, the order of rescattering increases up to the (A-1)-th order; we have performed calculations in the three-body case exactly, and did not investigate the problem of the convergence of the multiple scattering series. This problem is under investigation in the case of  ${}^{4}He$ . Most of our calculations have been performed in kinematical conditions where the effects of MEC,  $\Delta$  isobar creation, etc. are minimized, as confirmed by calculations performed, e.g., in Refs. [23, 50, 51, 52, 53, 55, 57, 58]. As for the main results we have obtained, the following remarks are in order:

- 1. the agreement between the results of our calculations and the experimental data for both the deuteron and  ${}^{3}He$ , is a very satisfactory one, particularly in view of the lack of any adjustable parameter in our approach;
- 2. the effects of the FSI are such that they systematically bring theoretical calculations in better agreement with the experimental data. For some quantities, FSI simply improve the agreement between theory and experiment (cf. e.g. Figs. 7, 9 and 11), whereas for some other quantities, they play a dominant role (see e.g. Fig. 10 and 12);
- 3. a comparison of the PWA and the PWIA with the full FSI calculation, does show that proper kinematics conditions could be found corresponding to an overall very small effect of FSI, leaving thus room for the investigation of the details of the nuclear wave function; as a matter of fact, we always found that in the 3bbu channel in  ${}^{3}He$ ,  ${}^{3}He(e, e'p)(np)$ , the experimental values of  $p_m$  and  $E_m$  corresponding to the maximum values of the cross section, satisfy to a large extent the relation predicted by the twonucleon correlation mechanism [38], namely  $E_m \simeq p_m^2/4M_N + E_3$  (cf. Fig. 3, right panel), with the full FSI mainly affecting only the magnitude of the cross section; thus, quasi elastic one-nucleon emission A(e, e'p)B processes at  $x \simeq 1$ , together with processes at  $x \simeq 2$ , when the virtual photon is absorbed by a correlated two-nucleon "system", would represent valuable tool for the investigation of correlations in nuclei;
- 4. calculations of the 2bbu channel disintegration of  ${}^{4}He$ , i.e. the process  ${}^{4}He(e, e'p){}^{3}H$ , have already been performed [15] using realistic wave functions and taking exactly into account nucleon rescattering up to 3rd order, i.e by using the generalization of Eq. (65) to the four-particle case, *viz*

$$\mathcal{S}_{\Delta}^{FSI} = \mathcal{S}_{(1)}^{FSI}(\mathbf{R}, \mathbf{r}_{12}, \mathbf{r}_{34}) + \mathcal{S}_{(2)}^{FSI}(\mathbf{R}, \mathbf{r}_{12}, \mathbf{r}_{34}) + \mathcal{S}_{(3)}^{FSI}(\mathbf{R}, \mathbf{r}_{12}, \mathbf{r}_{34}),$$
(75)

where  $\mathbf{R}$ ,  $\mathbf{r}_{12}$ , and  $\mathbf{r}_{34}$  are four-body Jacobi coordinates. Calculations for the 3bbu and 4bbu channels are in progress and will be reported elsewhere [59]; they should in principle yield results appreciably differing from the predictions based upon shellmodel type four-body wave functions;

- 5. our results for  ${}^{3}He$  generally agree with the ones obtained in Ref. [23], so that it would appear that the problem of the treatment of FSI at high values of  $Q^{2}$  (or high  $\mathbf{p}_{1}$ ) is under control; nevertheless, a systematic comparison of the various approaches would be highly desirable;
- 6. we have given the criteria according to which at high energies the exclusive A(e, e'p)B cross section should factorize, and the similarity of our results with the ones based upon a non factorized cross section [23], confirm the validity of these criteria.
- 7. eventually, it appears that in the kinematical range we have considered only minor numerical differences were found between the conventional Glauber-eikonal approach and its generalized extension; this does not mean at all that the same will hold in other kinematical conditions (see e.g. [9, 10, 11]).

#### VII. ACKNOWLEDGMENTS

The authors are indebted to A. Kievsky for making available the variational three-body wave functions of the Pisa Group and to G. Salmè for useful discussions about their use. Thanks are due to M.A. Braun and D. Treleani for stimulating discussions on the Feynman diagram approach to nucleon rescattering. Many useful discussions with various members of the Jlab Experiment E-89-044 are gratefully acknowledged. We express our gratitude to M. Alvioli for a careful reading of the manuscript. L.P.K. is indebted to the University of Perugia and INFN, Sezione di Perugia, for a grant and for warm hospitality. This work was partially supported by the Italian Ministero dell'Istruzione, Università e Ricerca (MIUR), through the funds COFIN01, and by the Russian Fund for Basic Research 00-15-96737.

## APPENDIX A: THE NUCLEAR WAVE FUNCTIONS

In our calculations we have used two- and three-body wave functions corresponding to the AV18 potential [5].

# 1. The ground state wave function of ${}^{3}He$

For the <sup>3</sup>*He* wave function we have adopted the correlated variational wave function by the Pisa group [2] which is written in a mixed  $(L_{\rho}, X, j_{23}, S_{23})$ -representation, where  $j_{23}$  and  $S_{23}$  are the total angular momentum and the total spin of the pair "23", X is an intermediate angular momentum resulting from the coupling  $\mathbf{j}_{23} + \mathbf{s}_1$  and  $L_{\rho}$  is the radial angular momentum of the motion of the nucleon "1" relative to the pair "23". The explicit form of the wave function is

$$\Psi_{He}^{\mathcal{M}_{3}}(\boldsymbol{\rho},\mathbf{r}) = \sum_{\{\alpha\}} \sum_{\{m\}} \langle XM_{X} \ L_{\rho}m_{\rho} \ |\frac{1}{2}\mathcal{M}_{3} \rangle \langle j_{23}m_{23} \ \frac{1}{2}\sigma_{1} \ |XM_{X} \ \rangle \chi_{\frac{1}{2}\sigma_{1}} \ Y_{L_{\rho}M_{\rho}}(\hat{\boldsymbol{\rho}})$$

$$\langle l_{23}\mu_{23} \ S_{23}\nu_{23} | j_{23}m_{23} \rangle Y_{l_{23}\mu_{23}}(\hat{\mathbf{r}}) \chi_{S_{23}\nu_{23}} R_{\{\alpha\}}(r,\rho) \mathcal{I}_{\frac{1}{2}\frac{1}{2}}^{T_{23}},$$
(A1)

where  $\{\alpha\}$  labels all possible configurations in  ${}^{3}He$  with quantum numbers  $L_{\rho}, X, j_{23}, S_{23}$ , and  $T_{23}$  and  $\langle l_{1}m_{1}l_{2}m_{2}| l_{12}m_{12} \rangle$  is a Clebsch-Gordan coefficient. The total isospin function is  $\mathcal{I}_{\frac{1}{2}\frac{1}{2}}^{T_{23}} = \sum \langle T_{23}\tau_{23} \frac{1}{2}\tau_{1}| \frac{1}{2}\frac{1}{2} \rangle \mathcal{J}_{T_{23}\tau_{23}}\eta_{\frac{1}{2}\tau_{1}}$ , where  $\mathcal{J}_{T_{23}\tau_{23}}$  and  $\eta_{\frac{1}{2}\tau_{1}}$  are the isospin functions of the pair and the nucleon, respectively. Obviously, because of Pauli principle and parity constraints, the allowed configurations in eq. (A1) are those that satisfy the following conditions:

$$L_{\rho} + l_{23}$$
 is even and  $l_{23} + S_{23} + T_{23}$  is odd (A2)

The corresponding radial part of the wave function,  $R_{\{\alpha\}}(r,\rho)$ , has been obtained [2] by a variational method using the AV18 potential including into the calculations values of  $L_{\rho}, l_{12} = 0 \dots 9$  (a total of 58 different configurations  $L_{\rho}, X, j_{23}, l_{23}, S_{23}$  have been considered).

## 2. The two-body continuum wave function $\Psi_{23}^{t}(\mathbf{r})$ .

With the representation (A1) of the <sup>3</sup>He wave function, it was convenient to adopt for the two-nucleon scattering state  $\Psi_{23}^{\mathbf{t}}(\mathbf{r})$  the spin-channel representation  $\Psi_{S_{23}\nu_{23}}^{\mathbf{t}}(\mathbf{r})$ , characterized by the total (conserved in the scattering process) spin  $S_{23}$  and its projection  $\nu_{23}$ . For spin  $S_{23} = 1$  one has

$$\Psi_{S_{f}\nu_{f}}^{\mathbf{t}}(\mathbf{r}) = 4\pi \sum_{J_{f}M_{f}} \sum_{l_{0}l_{f}} \langle l_{0}\mu_{0} \ S_{f}\nu_{f} | J_{f}M_{f} \rangle Y_{l_{0}\mu_{0}}(\hat{\mathbf{t}}) R_{J_{f},l_{0}l_{f}}^{|\mathbf{t}|}(r) i^{l_{f}} \mathcal{Y}_{1l_{f}}^{J_{f}M_{f}}(\hat{\mathbf{r}}) \mathcal{J}_{T_{23}\tau_{23}}, \quad (A3)$$

where  $l_0, l_f = J_f \pm 1$ ,  $J_f$ . Note that the presence of tensor forces in the NN-potential leads to an admixture of partial waves with  $l = J_f - 1$  and  $l = J_f + 1$ . This hinders the use of real phase shifts for the asymptotic behaviour of the radial functions  $R_{J_f,l_0l_f}^{[t]}(r)$  and, consequently, the Schrödinger equation cannot be solved in terms of real solutions. However, a unitary transformation V allows one to define new radial functions  $\tilde{R} = VR$  which are eigenfunctions of the scattering problem, i.e., solutions of the Schrödinger equation with the proper asymptotic behaviour.

# 3. Wave function overlaps and the spectral function $P(|\mathbf{k}_1|, E)$ of ${}^{3}He$ .

The Spectral function for the three-body break-up channel can be expressed in terms of the overlap between the three-body and two-body radial functions by substituting (A1)-(A3) into Eq. (12). Using the orthogonality of the spherical harmonics  $Y_{lm}(\hat{\mathbf{t}})$  and the completeness of the Clebsch-Gordan coefficients, one obtains that only diagonal ({ $\alpha$ } = { $\alpha_N$ }) matrix elements contribute to the spectral function, viz.

$$P_{ex}(|\mathbf{k}_{1}|, E) = \frac{1}{2} \sum_{\mathcal{M}_{3}} \sum_{\sigma_{f}, S_{f}, \nu_{f}} \int \frac{d^{3}\mathbf{t}}{(2\pi)^{3}} \left| \int d\boldsymbol{\rho} d\mathbf{r} \Psi_{He}^{\mathcal{M}_{3}}(\boldsymbol{\rho}, \mathbf{r}) \Psi_{S_{f}\nu_{f}}^{\mathbf{t}}(\mathbf{r}) \mathrm{e}^{-i\boldsymbol{\rho}\mathbf{k}_{1}} \right|^{2} \delta \left( E_{m} - \frac{\mathbf{t}^{2}}{M_{N}} - E_{3} \right) = \frac{M_{N} \sqrt{M_{N} E_{rel}}}{2\pi^{3}} f_{iso} \sum_{\{\alpha\}} \left| \int \rho^{2} d\rho j_{L_{\rho}}(p\rho) \mathcal{O}_{\{\alpha\}}^{E_{rel}}(\rho) \right|^{2},$$
(A4)

where  $f_{iso} = 3(1)$  for the pair in the isosinglet (isotriplet) final state,  $j_{L_{\rho}}(p\rho)$  is the spherical Bessel functions, and the dimensionless overlap integrals  $\mathcal{O}_{\{\alpha\}}^{E_{rel}}(\rho)$  are defined as follows

$$\mathcal{O}_{\{\alpha\}}^{E_{rel}}(\rho) = \int R_{\{\alpha\}}(r,\rho) \ \tilde{R}_{\{\alpha\}}^{[\mathbf{t}]}(r)r^2 dr \tag{A5}$$

The normalization of the proton spectral function (A4)-(A5) is

$$\int d^3 \mathbf{k}_1 dEP(|\mathbf{k}_1|, E) \approx \begin{cases} 0.15 \text{ for } T_{23} = 0\\ 0.50 \text{ for } T_{23} = 1 \end{cases}$$
(A6)

so that the two-body break-up channel is normalized to  $\approx 1.35$ . Since the FSI factors  $S^{FSI}$  and  $S^{FSI}_{\Delta}$  (Eqs. (65) and (68)) are no spherically symmetric, the distorted Spectral Function  $P_{ex}^{FSI}(\mathbf{p}_m, E_m)$  (Eq.64) is not longer diagonal with respect to the  $(L_{\rho}, X, j_{23}, S_{23})$  configurations. Except for parity constraints (A2), any values of angular momenta of the pair in the final state contribute to  $P_{ex}^{FSI}(\mathbf{p}_m, E_m)$ .

#### APPENDIX B: FACTORIZATION OF THE COVARIANT CROSS SECTION

In this Appendix we will show, within a fully covariant approach, that under certain kinematical conditions the cross section for the process A(e, e'p)X process factorizes even in presence of FSI. We shall consider, to this end, the deuteron (D) treated within the Bethe-Salpeter (BS) formalism. As mentioned, the factorization depends upon the spin structure of the square of the matrix element  $\left[\bar{u}(\mathbf{k}_1, \tilde{s}_1)\Phi_D^{\mathcal{M}_2}(k_1, k_2)(\hat{k}_2 + M_N)v(\mathbf{k}_2, s_2)\right]$  appearing in Eq. (19) or, in case of FSI, upon the structure of  $\left[\bar{u}(\mathbf{k}_1, \tilde{s}_1)\Phi_D^{\mathcal{M}_2}(k_1, k_2)v(\mathbf{k}_2, s_2)\right]$  (cf Eq. (28)). The relevant spin parts can be evaluated directly by using the explicit form of the Dirac spinors, u and v, and the explicit expressions for the amplitudes  $\Phi_D^{\mathcal{M}_2}(k_1, k_2)$  (cf. Refs. [42, 43, 45]).

#### 1. The PWIA

In Ref. [44] the Feynman diagrams for the process D(e, e'p)X have been evaluated including all BS components. Here we re-calculate the diagrams for the  ${}^{3}S_{1}^{++}$  and  ${}^{3}D_{1}^{++}$ components in a slightly different manner which will be useful when FSI effects are considered.

In PWIA the cross section reads as follows

$$\frac{d^5\sigma}{dE'd\Omega'} = \sigma_{Mott} \tilde{l}^{\mu\nu} L^D_{\mu\nu} \frac{d^3p_1}{(2\pi)^3 2E_1} \frac{d^3p_2}{(2\pi)^3 2E_2},\tag{B1}$$

where  $\tilde{l}^{\mu\nu}$  and  $L^{D}_{\mu\nu}$  are the leptonic and hadronic tensors, respectively, the latter being

$$L^{D}_{\mu\nu} = \frac{1}{2M_{D}} \frac{1}{3} \sum_{\mathcal{M}_{2}, s_{1}, s_{2}} T_{\mu}(\mathcal{M}_{2}, s_{1}, s_{2}) T_{\nu}(\mathcal{M}_{2}, s_{1}, s_{2}) (2\pi)^{3} \delta^{(4)} \left(P_{D} + q - p_{1} - p_{2}\right) =$$
(B2)

$$= \frac{1}{2M_D} \frac{1}{3} \sum_{\mathcal{M}_2, s_1, s_2} \left\langle \mathcal{M}_2 | \hat{J}^N_\mu | p_2, s_2, p_1, s_1 \right\rangle \left\langle p_1, s_1, p_2, s_2 | \hat{J}^N_\nu | \mathcal{M}_2 \right\rangle (2\pi)^3 \delta^{(4)} \left( P_D + q - p_1 - p_2 \right),$$

where  $\hat{J}^N_{\mu}$  is the nucleon electromagnetic current operator. The amplitude  $T_{\mu}$  could be written in the following form

$$T_{\mu}(\mathcal{M}_2, s_1, s_2) = \bar{u}(\mathbf{p}_1, s_1) \Gamma_{\mu}^{\gamma^* N}(Q^2, k_1^2) \Phi_D^{\mathcal{M}_2}(k_1, k_2) \tilde{S}^{-1}(\hat{k}_2) v(\mathbf{p}_2, s_2),$$
(B3)

where  $\Phi_D^{\mathcal{M}_2}$  is a short-hand notation for the main BS amplitudes  $\Phi_{3S_1^{++}}$  and  $\Phi_{3D_1^{++}}$ , corresponding to L = 0 and L = 2, respectively (see Refs. [42, 45]),  $k_2 = p_2$ ,  $\hat{S}^{-1}(\hat{k}_2) = \hat{k}_2 + m$ ,

and  $\Gamma_{\mu}^{\gamma^*N}(Q^2, k_1^2)$  is the electromagnetic eN vertex which, for an off-mass-shell nucleon, depends not only upon  $Q^2$ , but upon  $k_1^2 \neq m^2$  as well.

By introducing between  $\Gamma_{\mu}(Q^2, p_1, k_1)$  and  $\Phi_D^{\mathcal{M}_D}(k_1, k_2)$  the complete set of the Dirac spinors

$$\frac{1}{2M_N} \sum_{\tilde{s}_1} \left[ u(\mathbf{k}_1, \tilde{s}_1) \bar{u}(\mathbf{k}_1, \tilde{s}_1) - v(\mathbf{k}_1, \tilde{s}_1) \bar{v}(\mathbf{k}_1, \tilde{s}_1) \right]$$
(B4)

and bearing in mind that for the  ${}^{3}S_{1}^{++}$  and  ${}^{3}D_{1}^{++}$  partial waves the second term in (B4) does not contribute, we obtain

$$T_{\mu}(\mathcal{M}_{2}, s_{1}, s_{2}) = \frac{1}{2M_{N}} \sum_{\tilde{s}_{1}} J_{\mu}^{eN}(Q^{2}, p_{1}, k_{1}, \tilde{s}_{1}, s_{1}) \left[ \bar{u}(\mathbf{k}_{1}, \tilde{s}_{1}) \Phi_{D}^{M}(k_{1}, k_{2}) \tilde{S}^{-1}(\hat{k}_{2}) v(\mathbf{p}_{2}, s_{2}) \right], (B5)$$

where  $J^{eN}_{\mu}(Q^2, p_1, k_1) = \langle \mathbf{p}_1, s_1 | \Gamma^{\gamma^*N}_{\mu}(Q^2, k_1^2) | \mathbf{k}_1, \tilde{s}_1 \rangle.$ 

Let us evaluate Eq. (B5) for the D wave. One has

$$\left[\bar{u}(\mathbf{k}_{1},\tilde{s}_{1})\Phi_{^{3}D_{1}^{++}}^{\mathcal{M}_{2}}(k_{1},k_{2})\tilde{S}^{-1}(\hat{k}_{2})v(\mathbf{p}_{2},s_{2})\right] = -\frac{NN_{1}^{2}}{\sqrt{2}}(k_{2}^{2}-M_{N}^{2})\phi_{D}(k_{0},|\mathbf{k}|) \ 2m\left\langle\chi_{\tilde{s}_{1}}^{\dagger}\left|\left\{-(\boldsymbol{\sigma}\boldsymbol{\xi}^{M})+3(\mathbf{n}\boldsymbol{\xi}^{M})(\mathbf{n}\boldsymbol{\sigma})\right\}\right|\tilde{\chi}_{s_{2}}\right\rangle, \quad (B6)$$

where **n** is a unit vector along **k**, i.e.,  $\mathbf{n} = \frac{\mathbf{k}}{|\mathbf{k}|} = \frac{\mathbf{k}_1}{|\mathbf{k}_1|}$ . When Eq. (B6) is inserted in the expression for the cross section, one obtains

$$\frac{1}{3} \sum_{\mathcal{M}_{2}, s_{2}} \left\langle \chi_{\tilde{s}_{1}} \left| \left\{ -(\boldsymbol{\sigma}\boldsymbol{\xi}^{\mathcal{M}_{2}}) + 3(\mathbf{n}\boldsymbol{\xi}^{\mathcal{M}_{2}})(\mathbf{n}\boldsymbol{\sigma}) \right\} \right| \tilde{\chi}_{s_{2}} \right\rangle \left\langle \tilde{\chi}_{s_{2}} \left| \left\{ -(\boldsymbol{\sigma}\boldsymbol{\xi}^{+\mathcal{M}_{2}}) + 3(\mathbf{n}\boldsymbol{\xi}^{+}M)(\mathbf{n}\boldsymbol{\sigma}) \right\} \right| \chi_{\tilde{s}_{1}} \right\rangle \\
= 2\delta_{\tilde{s}_{1}\tilde{s}_{1}'}.$$
(B7)

The last relation ensures factorization of the cross section; as a matter of fact, by performing the same procedure for the S-wave, it easy to show that, thanks to Eq. (B7), the cross section (Eq. (B1)) factorizes, assuming the form (24) with  $n_D$  given by Eq. (22). In obtaining the above equations we expressed the BS amplitudes  $\phi_L(k_0, |\mathbf{k}|)$  in terms of the BS vertices  $G_{3L_1^{++}}(k_0, |\mathbf{k}|)$  and the radial functions  $u_L$ , by the relations

$$\frac{NN_1^2}{\sqrt{2}}(k_2^2 - M_N^2)\phi_D(k_0, |\mathbf{k}|) = \frac{NN_1^2 2E_{\mathbf{k}}}{\sqrt{2}} \frac{G_{^3D_1^{++}}(k_0, |\mathbf{k}|)}{M_D - 2E_{\mathbf{k}}},$$
(B8)

where  $k_0 = \frac{M_D}{2} - E_{\mathbf{k}}$ , and

$$u_{S(D)} = \frac{G_{3S_1^{++}(^{3}D_1^{++})}(k_0, |\mathbf{k}|)/(4\pi)}{\sqrt{2M_D}(M_D - 2E_{\mathbf{k}})}.$$
(B9)

Note that in Eq. (B9) the normalization of the wave function is chosen so as to correspond to the non relativistic deuteron wave function

$$\frac{2}{\pi} \int |\mathbf{k}|^2 d|\mathbf{k}| \left( u_S^2(|\mathbf{k}|) + u_D^2(|\mathbf{k}|) \right) \approx 1.$$
(B10)

We reiterate that factorization in PWIA occurs because the sum over  $s_2$  and  $\mathcal{M}_2$  of the square of the matrix element in Eq. (B7) becomes diagonal with respect to  $\tilde{s}_1$ .

When FSI is taken into account, instead of Eq. (B7), one obtains for the *D*-wave (for the S-wave the spin structure is trivial)

$$\frac{1}{3} \sum_{\mathcal{M}_{2}, s_{2}} \left[ \bar{u}(\mathbf{k}_{1}, s_{1}) \Phi_{D}^{\mathcal{M}_{2}}(k_{1}, k_{2}) v(\mathbf{k}_{2}, s_{2}) \right]^{\dagger} \left[ \bar{u}(\mathbf{k}_{1}', \tilde{s}_{1}) \Phi_{D}^{\mathcal{M}_{2}}(k_{1}', k_{2}') v(\mathbf{k}_{2}', s_{2}) \right] \simeq 
\frac{1}{3} \sum_{\mathcal{M}, s_{2}} \left\langle \chi_{s_{1}} \left| \left\{ -(\boldsymbol{\sigma}\boldsymbol{\xi}^{\mathcal{M}_{2}}) + 3(\mathbf{n}\boldsymbol{\xi}^{\mathcal{M}_{2}})(\mathbf{n}\boldsymbol{\sigma}) \right\} \right| \tilde{\chi}_{s_{2}} \right\rangle \left\langle \tilde{\chi}_{s_{2}} \left| \left\{ -(\boldsymbol{\sigma}\boldsymbol{\xi}^{+\mathcal{M}_{2}}) + 3(\mathbf{n}'\boldsymbol{\xi}^{+\mathcal{M}_{2}})(\mathbf{n}'\boldsymbol{\sigma}) \right\} \right| \chi_{\tilde{s}_{1}} \right\rangle = 
= \frac{1}{3} \sum_{\mathcal{M}_{2}} \left\langle \chi_{s_{1}} \left| \left\{ -(\boldsymbol{\sigma}\boldsymbol{\xi}^{\mathcal{M}_{2}}) + 3(\mathbf{n}\boldsymbol{\xi}^{\mathcal{M}_{2}})(\mathbf{n}\boldsymbol{\sigma}) \right\} \left\{ -(\boldsymbol{\sigma}\boldsymbol{\xi}^{+\mathcal{M}_{2}}) + 3(\mathbf{n}'\boldsymbol{\xi}^{+\mathcal{M}_{2}})(\mathbf{n}'\boldsymbol{\sigma}) \right\} \right| \chi_{\tilde{s}_{1}} \right\rangle \tag{B11}$$

where  $\mathbf{n}$  ( $\mathbf{n}'$ ) is a unit vector along  $\mathbf{k}_1$  ( $\mathbf{k}'_2$ ).

By taking into account the completeness of the polarization vectors  $\boldsymbol{\xi}$ , the only spin dependence remaining in Eq. (B11) is contained in the term

$$\frac{1}{3}\sum_{M}(\mathbf{n}\boldsymbol{\xi}^{M})(\mathbf{n}\boldsymbol{\sigma})(\mathbf{n}'\boldsymbol{\xi}^{+M})(\mathbf{n}'\boldsymbol{\sigma}) = (\mathbf{n}\mathbf{n}')\left((\mathbf{n}\mathbf{n}') - i\boldsymbol{\sigma}\left[\mathbf{n}\times\mathbf{n}'\right]\right),\tag{B12}$$

so that in case of rescattering with low momentum transfer, when in the integral over  $\mathbf{k}_2$  the main contribution comes from  $\mathbf{k}_2 \sim \mathbf{p}_2$ ,  $\mathbf{k}'_2 \sim \mathbf{p}_2$ , one has  $\boldsymbol{\sigma} [\mathbf{n} \times \mathbf{n}'] = 0$  and factorization is approximately recovered, with the S and D waves adding incoherently.

Thus, to sum up, factorization is compatible with FSI if:

- 1. The spin-flip part of the NN amplitude should be very small, as it occurs when either the three-momentum transfer  $\mathbf{q}$ , or the momentum  $|\mathbf{p}_1|$  are large;
- 2. the momentum transfer  $\kappa$  in the NN rescattering has to be small small so that in the integral  $\mathbf{k}_2 \sim \mathbf{p}_2$ . This appears to be the case since the NN amplitude is sharply peaked forward.
- 3. The contribution from  $N\bar{N}$  pair currents can be neglected, which is to a large extent legitimate due to the smallness of the P wave in the deuteron.

- [1] W. Glöckle. H. Witala, D. Huber, H. Kamada, and J. Golak, Phys. Rep. 274 (1996) 107.
- [2] A. Kievsky, S. Rosati and M. Viviani, Nucl. Phys. A551 (1993) 241; A. Kievsky, Private communication.
- [3] S. C. Pieper, R. B. Wiringa, Ann. Rev. Nucl. Part. Sci. 51, (2001) 53.
- [4] S.C. Pieper, R.B. Wiringa and V.R. Pandharipande, Phys. Rev. C46, 1741 (2000).
- [5] R. B. Wiringa, V. G. J. Stoks and R. Schiavilla, Phys. Rev. C51 (1995) 38.
- [6] V.N.Gribov, Sov. Phys. JETP, **30** (1970) 709.
- [7] L. Bertocchi, Nuovo Cimento, **11A** (1972) 45.
- [8] J. H. Weis, Acta Phys. Pol. **B7**(1976)851.
- [9] L.L. Frankfurt, W.R. Greenberg, G.A. Miller, M.M. Sargsian, M.I. Strikman, Z. Phys. A352 (1995) 97.
- [10] L.L. Frankfurt, M.M. Sargsian, M.I. Strikman, Phys. Rev. C56 (1997) 1124.
- [11] M.M. Sargsian, Int. J. Mod. Phys. **E10**(2001)405.
- [12] M.M. Sargsian, T. V. Abrahamyan, M.I. Strikman, L.L. Frankfurt, arxiv: nucl-th/0406020.
- [13] M.A. Braun, C. Ciofi degli Atti, and D. Treleani, Phys. Rev. C62, 034606 (2000).
- [14] M. Braun, C. Ciofi degli Atti, L.P. Kaptari, Eur. J. Phys. A19 (2004) 143; arxiv: nucl-th/0311060.
- [15] H. Morita, M. Braun, C. Ciofi degli Atti and D. Treleani, Nucl. Phys. A699 328c (2002).
- [16] R. J. Glauber, in *Lectures in Theoretical Physics*, Vol. I p.315, edited by W. E. Brittin and L. G. Dunham. Wiley Interscience, New York (1959).
- [17] W. -J. Kasdorp *et al*, Few Body Syst. **25** (1998) 115.
- [18] P.E. Ulmer *et al.*, Phys. Rev. Lett., **89** (2002) 062301.
- [19] H.J. Bulten *et al*, Phys. Rev. Lett., **74** (1995) 4775.
- [20] C. Marchand *et al*, Phys. Rev. Lett. **60** (1988) 1703.
- [21] M. M. Rvachev et al,arxiv: nucl-ex/0409005v2.
- [22] F. Benmokhtar *et al*, arxiv: nucl-ex/0408015v2.
- [23] J.-M. Laget, arxiv: nucl-th/0410003.
- [24] J. J. van Leeuwe *et al*, Phys. Lett. **B523** (2001) 6.
- [25] S. Janssen, J. Ryckebusch, W. Van Nespen, D. Debruyne, Nucl. Phys. A 672 (2000) 285.

- [26] C.Ciofi degli Atti and L.P. Kaptari, in International Workshop on Probing Nucleons and Nuclei via the (e, e'p) reaction, Eds. D. Higinbotham, J. M. Laget and E. Voutier, The Print House, New York 2004, p. 121; arxiv: nucl-th/0312043.
- [27] M. Alvioli, C. Ciofi degli Atti, L.P. Kaptari, H. Morita, Proceedings of the 6th Workshop on Electromagnetically induced Two Hadron Emission, Pavia, Italy, 24-27 Sep 2003; arxiv: nucl-th/0312123.
- [28] S. Boffi, C. Giusti and F.D. Pacati, Phys. Rep. **226** (1993) 1.
- [29] R. -W. Schulze and P. E. Sauer, Phys. Rev. C226 (1993)38.
- [30] C. Ciofi degli Atti, E. Pace and G. Salmè, Phys. Rev. C21 (1980) 805.
- [31] A. Kievski, E. Pace, G. Salmè and M. Viviani, Phys. Rev. C56(1997)64.
- [32] C. Ciofi degli Atti, E. Pace and G. Salmè, in *Lecture Notes in Physics Vol.* 86 (Springer, Berlin, 1978) p. 315;
- [33] H. Meier-Hajduk, Ch. Hajduk, P. E. Sauer and W. Theis, Nucl. Phys. A395 (1983) 332;
   U. Oelfke, P. U. Sauer and F. Coester, Nucl. Phys. A518 (1990)593.
- [34] C. Ciofi degli Atti, L. Kaptari and S. Scopetta, Eur. Phys. Jou. A5 (1999) 191.
- [35] T. de Forest Jr., Nucl. Phys. A392 (1983) 232.
- [36] R.V. Reid Jr., Ann. Phys. **50** (1968) 411.
- [37] L. L. Frankfurt and M.I. Strikman, Phys. Rep. 160 (1988) 235.
- [38] C. Ciofi degli Atti, S. Simula, L.L. Frankfurt and M.I. Strikman, Phys. Rev. C 44 (1991) R7;
  C. Ciofi degli Atti and S. Simula, Phys. Rev. C53 (1996) 1689.
- [39] J. Golak, H. Witala, R. Skininski, W. Glöckle, A. Nogga, H. Kamada, Phys. Rev. C 70, 034005 (2004).
- [40] C. Ciofi degli Atti, L. P. Kaptari, Phys. Rev. C66 (2002)044004.
- [41] S. Jeschonnek, Phys. Rev. C63 (2001) 034609;
   J. Ryckebusch *et al*, Nucl. Phys. A728 (2003) 226.
- [42] B.D. Keister, J.A. Tjon, Phys. Rev. C26 (1982) 578;
  G. Rupp, J.A. Tjon, Phys. Rev. C41 (1990) 472;
  J.J. Kubis, Phys. Rev. D6 (1972) 547;
  M.J. Zuilhof, J.A. Tjon, Phys. Rev. C22 (1980) 2369.
- [43] F. Gross, Phys. Rev. 186 (1969) 1448;
  W.W. Buck, F. Gross, Phys. Rev. C20 (1979) 2361;

F. Gross, J.V. Van Orden, K. Holinde, *ibid* **45** (1992) 2094.

- [44] C. Ciofi degli Atti, D. Faralli, A.Yu. Umnikov and L.P. Kaptari, Phys. Rev. C60 (1999) 034003.
- [45] L.P. Kaptari, B. Kampfer, S.M. Dorkin, S.S. Semikh, Few Body Syst. 27 (1999) 189;
   L.P. Kaptari, B. Kampfer, S.M. Dorkin, S.S. Semikh, Phys.Rev. C 57 (1998) 1097.
- [46] C. Ciofi degli Atti, L.P. Kaptari and D. Treleani Phys. Rev. C63 (2001) 044601.
- [47] N. N. Nikolaev, J. Speth, B. G. Zakharov, JETP 82, (1996) 1046; arxiv: nucl-th/9501016.
- [48] A. Bianconi, S. Jeschonnek, N. N. Nikolaev and B. G. Zacharov, Phys. Lett. B343 (1995)13.
- [49] R.A. Arndt et al, "Partial-Wave Analysis Facility (SAID)", http://said.phys.vt.edu/.
- [50] J.-M. Laget, Phys. Lett. **B199** (1987) 1046.
- [51] F. Ritz, H. Göller, T. Wilbois and H. Arenhövel, Phys. Rev. C55(1997)2214.
- [52] G. Beck and H. Arenhövel, Few-Body Systems 13(1992)165.
- [53] G. Beck, T. Wilbois and H. Arenhövel, Few-Body Systems 17(1994)91.
- [54] E. Hummel, J. A. Tjon, Phys. Rev. C49(1994) 21.
- [55] J.-M. Laget, Few-Body Systems Suppl. 15 (2003) 171; arxiv:nucl-th/0303052.
- [56] H. Morita, C.Ciofi degli Atti and D. Treleani, Phys. Rev. C60(1999) 34603.
- [57] J.-M. Laget, Phys. Rev. **D579** (1994) 333.
- [58] J.-M. Laget, in New Vistas in Electronuclear Physics, edited by E. Tomusiak et al, Plenum N.Y.,1986, p.361.
- [59] M. Alvioli, C. Ciofi degli Atti, L. Kaptari and H. Morita, in preparation.



FIG. 1: The one-photon exchange approximation for the process A(e, e'p)(A-1).



c)

FIG. 2: Feynman diagrams for the process A(e, e'p)(A-1). (a) describes the Plane Wave Impulse Approximation (PWIA); (b) the single rescattering; (c) the full A - 1 rescattering. The fourmomenta of particle *i* before and after rescattering are denoted by  $k_i$ ,  $k'_i$ ,  $k''_i$ , etc., respectively. The black oval spots denote the elastic nucleon-nucleon scattering matrix..



FIG. 3: The proton Spectral Function of  ${}^{3}He$  (Eq.(9)). Left panel:  $n_{gr}$  (Eq. (11)) vs  $p \equiv |\mathbf{k}_{1}|$ . Right panel:  $P_{ex}$  (Eq. (12)) vs the excitation energy of the two-nucleon system in the continuum  $E_{rel} = \frac{\mathbf{t}^{2}}{M_{N}} = E_{2}^{f} = E - E_{min}$ , for various values of  $p \equiv |\mathbf{k}_{1}|$ . The dot-dashed curves represent the Plane Wave Approximation (PWA), when the three particles in the continuum are described by plane waves, whereas the full curves correspond to the PWIA, when the interaction in the spectator proton-neutron pair is taken into account. The arrows indicate the position of the peak ( $\sim p^{2}/4M_{N}$ ) predicted by the two-nucleon correlation model for the Spectral Function [38] (threebody wave function from Ref. [2], AV18 interaction [5]).



FIG. 4: The neutron Spectral Function of  ${}^{3}He$  (Eq. (12))  $vs \ p \equiv |\mathbf{k}_{1}|$  and the excitation energy of the two-nucleon system in the continuum  $E_{rel} = \frac{\mathbf{t}^{2}}{M_{N}} = E_{2}^{f} = E - E_{min}$ . The dotted curves represent the *PWA*, when the three particles in the continuum are described by plane waves, whereas the full curves correspond to the *PWIA*, when the interaction in the spectator protonproton pair is taken into account (three-body wave function from Ref. [2], *AV*18 interaction [5]).



FIG. 5: The Feynman diagrams for the process  ${}^{2}H(e, e'p)n$  representing the PWIA) (a)), and the single (b)) rescattering in the final state.  $f^{NN}$  denotes the elastic NN scattering amplitude.



a)



b)



c)

FIG. 6: The Feynman diagrams representing the PWIA (a)), the single (b)), and double (c)) rescattering in the processes  ${}^{3}He(e,e'p)D$  and  ${}^{3}He(e,e'p)(np)$ . In the former case the final twonucleon state is a deuteron with momentum  $\mathbf{P}_{D} = \mathbf{P}_{2}$ , whereas in the latter case the final state represents two free nucleons with momenta  $\mathbf{p}_{2}$  and  $\mathbf{p}_{3}$ , with  $\mathbf{P}_{2} = \mathbf{p}_{2} + \mathbf{p}_{3}$ . The trivial single and double rescattering diagrams with nucleons "2" and "3" interchanged are not drawn. The black oval spots denote the elastic nucleon-nucleon (NN) scattering matrix  $\hat{T}$  (See Eq. (25)).



FIG. 7: The process  ${}^{2}H(e, e'p)n$ : the NIKHEF experimental data [17] vs the missing momentum  $p_{m} \equiv |\mathbf{p}_{m}|$  are compared with our theoretical calculations; the dotted line represents the *PWIA*, whereas the full line include the final state rescattering. The curves labelled *L*01, *L*02, and *L*03, correspond to  $Q^{2} = 0.1$ , 0.2, and 0.3  $(GeV/c)^{2}$ , respectively, and  $x \simeq 0.3 - 0.6$  (in this Figure and in Figs. 8-12,  $p' \equiv |\mathbf{p}_{1}|$ ).



FIG. 8: The process  ${}^{2}H(e,e'p)n$ : the Jlab experimental data [18] ( $N_{eff}$  defined by Eq. (72)) vs the missing momentum  $p_m \equiv |\mathbf{p}_m|$ , compared with our theoretical calculations. The dotted line represents the PWIA whereas the full line includes the final state rescattering. The experimental data correspond to the perpendicular kinematics, with  $Q^2 \simeq 0.665$  (GeV/c)<sup>2</sup>,  $|\mathbf{q}| \simeq 0.7$  GeV/c, and  $x \simeq 0.96$ .



FIG. 9: The process  ${}^{2}H(e, e'p)n$ : the SLAC experimental data at  $x \simeq 1$  [19] vs the missing momentum  $p_m \equiv |\mathbf{p}_m|$ , compared with our theoretical calculations. The dotted lines represent the PWIA and the full lines include the final state rescattering. The positive and negative values of  $p_m$  correspond to values of the azimuthal angle  $\phi = \pi$  and 0, respectively.



FIG. 10: The process  ${}^{3}He(e, e'p)D$ : the experimental data from JLab (JLab Experiment E-89-044 [21]) vs  $p_m \equiv |\mathbf{p}_m|$  compared, at  $Q^2 = 1.55 \ (GeV/c)^2$  and x = 1, with our theoretical results. The dashed line corresponds to the PWIA and the full line includes the full FSI calculated using Eq. (63); the predictions by Eq.(65) (GEA) and Eq. (68) (GA), differ by at most 4% and cannot be distinguished in the Figure (three-body wave function from [2], AV18 interaction [5]).



FIG. 11: The process  ${}^{3}He(e, e'p)(np)$ : the experimental data from Saclay [20] vs  $E_m$  and for various values of the proton emission angle  $\theta_{\mathbf{p}_1}$  ( $p' \equiv |\mathbf{p}_1|$ ), are compared with our theoretical results. The dotted lines correspond to the PWA, when the three nucleons in the final state are described by plane waves, the dashed lines correspond to the PWIA, when the interaction in the spectator neutron-proton pair is taken into account, and the full lines include the full FSI calculated using Eq. (63); the predictions by Eq.(65) (GEA) and Eq. (68) (GA), differ by at most 4% and cannot be distinguished in the Figure. Note that the values of the experimental  $p_m$  and  $E_m$  corresponding to the maxima of the cross section, satisfy to a large extent the relation predicted by the two-nucleon correlation mechanism [38], namely  $E_m \simeq p_m^2/4M_N$  (cf. Fig. 3, right panel), with the full FSI mainly affecting only the magnitude of the cross section ( three-body wave function from [2], AV18interaction [5]).



FIG. 12: The process  ${}^{3}He(e, e'p)(np)$ : the same as in Fig. 11 but for the Jlab experimental data [22]) at  $Q^{2} = 1.55 \ (GeV/c)^{2}$  and x = 1. Note, that unlike what shown in Fig. 11, here the differential cross section is plotted vs the excitation energy of the two-nucleon system in the continuum i.e.  $E_{rel} = \frac{\mathbf{t}^{2}}{M_{N}} = E_{2}^{f} = E_{m} - E_{3}$ .