

# Mass and $K\Lambda$ coupling of $N^*(1535)$

B.C.Liu<sup>1</sup>, B. S. Zou<sup>2,1</sup>

<sup>1</sup> Institute of High Energy Physics, P.O.Box 918(4), Beijing 100049, China

<sup>2</sup> CCAST (World Lab.), P.O.Box 8730, Beijing 100080, China

(Dated: March 29, 2005)

## Abstract

Using resonance isobar model and effective Lagrangian approach, from recent BES results on  $J/\psi \rightarrow \bar{p}p\eta$  and  $\psi \rightarrow \bar{p}K^+\Lambda$ , we deduce the ratio between effective coupling constants of  $N^*(1535)$  to  $K\Lambda$  and  $p\eta$  to be  $R \equiv g_{N^*(1535)K\Lambda}/g_{N^*(1535)p\eta} = 1.3 \pm 0.3$ . With previous known value of  $g_{N^*(1535)p\eta}$ , the obtained new value of  $g_{N^*(1535)K\Lambda}$  is shown to reproduce recent  $pp \rightarrow pK^+\Lambda$  near-threshold cross section data as well. Taking into account this large  $N^*K\Lambda$  coupling in the coupled channel Breit-Wigner formula for the  $N^*(1535)$ , its Breit-Wigner mass is found to be around 1400 MeV, much smaller than previous value of about 1535 MeV obtained without including its coupling to  $K\Lambda$ . The implication on the nature of  $N^*(1535)$  is discussed.

The properties and the nature of the lowest spin-1/2 negative parity ( $J^P = 1/2^-$ ) nucleon resonance  $N^*(1535)$  are of great interests in many aspects of light hadron physics. In conventional constituent quark models, the lowest  $1/2^-$   $N^*$  resonance should be the first  $L = 1$  orbital excitation state. But it has been a long-standing problem for these conventional constituent quark models to explain why the mass of  $N^*(1535)$  has a mass higher than the lowest  $J^P = 1/2^+$  radial excitation state  $N^*(1440)$  [1]. This was used to argue in favor of the Goldstone-boson exchange quark models [2]. In the recent Jaffe-Wilczek diquark picture [3] for the  $\theta$  pentaquark, a  $J^P = 1/2^-$   $N^*$  pentaquark of mass around 1460 MeV is expected [4]. Another outstanding property of the  $N^*(1535)$  is its extraordinary strong coupling to  $\eta N$  [5], which lead to a suggestion that it is a quasi-bound ( $K\Sigma$ - $K\Lambda$ )-state [6]. This picture predicts also large effective couplings of  $N^*(1535)$  to  $K\Lambda$  and  $K\Sigma$  [7]. Experiment knowledge on these kaon-hyperon couplings is poor, partly because lack of data on experimental side and partly due to the complication of various interfering t-channel exchange contributions [8]. Better knowledge on these couplings is definitely useful for understanding the nature of  $N^*(1535)$ , the underneath quark dynamics, and also the strangeness production in relativistic heavy-ion collisions as a signature of the quark-gluon plasma [9, 10, 11].

Recently various  $N^*$  production processes from  $J/\psi$  decays have been investigated by BES collaboration [12, 13, 14, 15]. In the  $J/\psi \rightarrow \bar{p}p\eta$  [12, 13] and  $\psi \rightarrow \bar{p}K^+\Lambda + c.c.$  [14, 15] reactions, there are clear peak structures with  $J^P = 1/2^-$  in the  $p\eta$  and  $K\Lambda$  invariant mass spectra around  $p\eta$  and  $K\Lambda$  thresholds. A nature source for the peak structures is  $N^*(1535)$  coupling to  $N\eta$  and  $K\Lambda$ . In this letter, assuming the  $1/2^-$   $K\Lambda$  threshold peak to be dominantly from the tail of the  $N^*(1535)$ , we deduce the ratio between effective coupling constants of  $N^*(1535)$  to  $K\Lambda$  and  $p\eta$ ,  $R \equiv g_{N^*(1535)K\Lambda}/g_{N^*(1535)p\eta}$  from the new branching ratio results from BES experiment on  $J/\psi \rightarrow \bar{p}p\eta$  and  $\psi \rightarrow \bar{p}K^+\Lambda$ , then check the compatibility with recent  $pp \rightarrow pK^+\Lambda$  near-threshold data [16, 17]. Taking into account the large  $N^*K\Lambda$  coupling in the coupled channel Breit-Wigner formula for the  $N^*(1535)$ , we show it gives a very large influence to the Breit-Wigner mass of the  $N^*(1535)$ .

In the effective Lagrangian approach for the resonance isobar model, the Feynman diagram for  $\psi \rightarrow \bar{p}K^+\Lambda$  through  $N^*(1535)$  intermediate is shown in Fig.1. For  $\psi \rightarrow \bar{p}p\eta$ , besides a similar diagram through  $N^*(1535)$ , a diagram through  $\bar{N}^*(1535)$  should be added




FIG. 1: Feynman diagram for  $\psi \rightarrow \bar{p}K^+\Lambda$  through  $N^*$  resonance

simultaneously. The relevant interaction Lagrangians are [18, 19]

$$\mathcal{L}_{N^*\Lambda K} = -ig_{N^*\Lambda K}\bar{\Psi}_\Lambda\Phi_K\Psi_{N^*} + h.c., \quad (1)$$

$$\mathcal{L}_{N^*N\eta} = -ig_{N^*N\eta}\bar{\Psi}_N\Phi_\eta\Psi_{N^*} + h.c., \quad (2)$$

$$\mathcal{L}_{\psi NN^*}^{(1)} = \frac{ig_T}{M_{N^*} + M_p}\bar{\Psi}_{N^*}\gamma_5\sigma_{\mu\nu}p_\psi^\nu\Psi_N\varepsilon^\mu + h.c., \quad (3)$$

$$\mathcal{L}_{\psi NN^*}^{(2)} = -g_V\bar{\Psi}_{N^*}\gamma_5\gamma_\mu\Psi_N\varepsilon^\mu + h.c. \quad (4)$$

where  $\Psi_{N^*}$  represents the resonance  $N^*(1535)$  with mass  $M_{N^*}$ ,  $\Psi_N$  for proton with mass  $M_p$  and  $\varepsilon^\mu$  for  $J/\psi$  with four-momentum  $p_\psi$ . According to [13], the  $\mathcal{L}_{\psi NN^*}^{(2)}$  term given by Eq.(4) makes insignificant contribution for  $N^*(1535)$ , hence we drop this kind of coupling in our calculation. The amplitudes for  $J/\psi \rightarrow \bar{p}K^+\Lambda$  and  $\bar{p}p\eta$  via  $N^*(1535)$  resonance are then

$$M_{\psi \rightarrow \bar{p}K^+\Lambda} = \frac{ig_T g_{N^*K\Lambda}}{M_{N^*} + M_p} \bar{u}(p_\Lambda, s_\Lambda)(\not{p}_{N^*} + m_{N^*})BW(p_{N^*})\gamma_5\sigma_{\mu\nu}p_\psi^\nu\varepsilon^\mu v(p_{\bar{p}}, s_{\bar{p}}), \quad (5)$$

$$M_{\psi \rightarrow \bar{p}p\eta} = \frac{ig_T g_{N^*N\eta}}{M_{N^*} + M_p} \bar{u}(p_p, s_p)[(\not{p}_{N^*} + m_{N^*})BW(p_{N^*})\gamma_5\sigma_{\mu\nu}p_\psi^\nu\varepsilon^\mu + \gamma_5\sigma_{\mu\nu}p_\psi^\nu\varepsilon^\mu(-\not{p}_{\bar{N}^*} + m_{N^*})BW(p_{\bar{N}^*})]v(p_{\bar{p}}, s_{\bar{p}}), \quad (6)$$

respectively. Here  $BW(p_{N^*})$  is the Breit-Wigner formula for the  $N^*(1535)$  resonance

$$BW(p_{N^*}) = \frac{1}{M_{N^*}^2 - s - iM_{N^*}\Gamma_{N^*}(s)} \quad (7)$$

with  $s = p_{N^*}^2$ . According to PDG [5], the dominant decay channels for the  $N^*(1535)$  are  $N\pi$  and  $N\eta$ . For a resonance with mass close to some threshold of its dominant decay channel, the approximation of a constant width is not very good. Since the  $N^*(1535)$  is quite close to the  $\eta N$  threshold, we take the commonly used phase space dependent width for the resonance as the following

$$\Gamma_{N^*}(s) = \Gamma_{N^*}^0 \left( 0.5 \frac{\rho_{\pi N}(s)}{\rho_{\pi N}(M_{N^*}^2)} + 0.5 \frac{\rho_{\eta N}(s)}{\rho_{\eta N}(M_{N^*}^2)} \right) = \Gamma_{N^*}^0 [0.8\rho_{\pi N}(s) + 2.1\rho_{\eta N}(s)], \quad (8)$$

where  $\rho_{\pi N}(s)$  and  $\rho_{\eta N}(s)$  are the phase space factors for  $\pi N$  and  $\eta N$  final states, respectively, e.g.,

$$\rho_{\eta N}(s) = \frac{2q_{\eta N}(s)}{\sqrt{s}} = \frac{\sqrt{(s - (M_N + M_\eta)^2)(s - (M_N - M_\eta)^2)}}{s} \quad (9)$$

where  $q_{\eta N}$  is the momentum of  $\eta$  or  $N$  in the center-of-mass system of  $\eta N$ . According to PDG [5],  $M_{N^*} \approx 1535 \text{ MeV}$  and  $\Gamma_{N^*}^0 = \Gamma_{N^*}(M_{N^*}^2) \approx 150 \text{ MeV}$ .

From the amplitudes given above, we can calculate the decay widths of  $J/\psi \rightarrow \bar{p}K^+\Lambda$  and  $J/\psi \rightarrow \bar{p}p\eta$  via  $N^*(1535)$  resonance, and get their ratio as

$$\frac{\Gamma(\psi \rightarrow \bar{p}N^* \rightarrow \bar{p}K^+\Lambda)}{\Gamma(\psi \rightarrow \bar{p}N^* + p\bar{N}^* \rightarrow \bar{p}p\eta)} = \frac{1}{12.6} \left| \frac{g_{N^*K\Lambda}}{g_{N^*N\eta}} \right|^2. \quad (10)$$

On the other hand, from PDG and recent BES results, we have  $J/\psi$  decay branching ratio for the  $\bar{p}K^+\Lambda$  channel as  $(0.89 \pm 0.16) \times 10^{-3}$  [5] with  $(15 \sim 22)\%$  [15] via the near threshold  $N^*$  resonance and for the  $\bar{p}p\eta$  channel as  $(2.09 \pm 0.18) \times 10^{-3}$  [5] with  $(56 \pm 15)\%$  [13] via the  $N^*(1535)$  resonance. Therefore

$$\frac{\Gamma(\psi \rightarrow \bar{p}N^* \rightarrow \bar{p}K^+\Lambda)}{\Gamma(\psi \rightarrow \bar{p}N^* + p\bar{N}^* \rightarrow \bar{p}p\eta)} = \frac{(0.89 \pm 0.16) \times (15 \sim 22)}{(2.09 \pm 0.18) \times (56 \pm 15)}. \quad (11)$$

From Eq(10) and Eq(11), we get

$$R \equiv \left| \frac{g_{N^*(1535)K\Lambda}}{g_{N^*(1535)N\eta}} \right| \approx 1.3 \pm 0.3. \quad (12)$$

Previous knowledge on this ratio from  $\pi N \rightarrow K\Lambda$  and  $\gamma N \rightarrow K\Lambda$  reactions is poor. While Ref.[8] gave a range of  $0.8 \sim 2.6$ , others found the contribution from the  $N^*(1535)$  is not important for reproducing the data [11]. It seems that those data are not sensitive to the  $N^*(1535)$  contribution due to the complication of various interfering t-channel contributions which are absent in the  $J/\psi$  decays. Another relevant reaction is  $pp \rightarrow pK^+\Lambda$ . Some very precise near-threshold data are now available from COSY experiments [16, 17]. In the following we will check the compatibility of the large  $R$  value given by Eq.(12) with the recent  $pp \rightarrow pK^+\Lambda$  near-threshold data.

The relevant Feynman diagrams for the process  $pp \rightarrow pK^+\Lambda$  are shown in Fig.2. Since we are mainly interested in the near-threshold behavior where contribution from  $\pi$  and  $\eta$  meson exchange dominates [20], here for simplicity we ignore the small contribution from heavier mesons. We adopt the relevant effective Lagrangian and form factors used in Ref. [20].




FIG. 2: Feynman diagrams for reaction  $pp \rightarrow pK^+\Lambda$

First we have reproduced the results of Ref.[20] by including  $N^*(1650)1/2^-$ ,  $N^*(1710)1/2^+$  and  $N^*(1720)3/2^+$  resonances. Their prediction prior COSY data [16, 17] is shown by the dotted line in Fig.3, which is obviously underestimating the near-threshold data of COSY. In their work, all parameters have been fixed by previous study on other relevant reactions. A natural reason for the underestimation is their ignorance of the contribution from  $N^*(1535)$ . Here we calculate the contribution from  $N^*(1535)1/2^-$  for the process. The coupling constants for the vertices  $N^*(1535)N\pi$  and  $N^*(1535)N\eta$  are determined by the relevant partial decay width [5]. Then the coupling constant for the  $N^*K\Lambda$  is obtained by our new result  $|g_{N^*(1535)K\Lambda}/g_{N^*(1535)N\eta}| = 1.3$  from BES data. The result is shown by the dashed line in Fig.3 (left). Adding the contribution to the previous results of Ref.[20], the solid line in Fig.3 (left) reproduces the COSY near-threshold data very well. So the ratio given by Eq.(12) is also compatible with the data on  $pp \rightarrow pK^+\Lambda$ . Note we have not introduce any free parameters in this calculation.

The large  $|g_{N^*(1535)K\Lambda}/g_{N^*(1535)N\eta}|$  ratio has important implications on other properties of the  $N^*(1535)$ . First, in previous calculations, the coupling of  $N^*(1535)$  to  $K\Lambda$  channel is usually ignored in the Breit-Wigner formula for the  $N^*(1535)$ . Considering this coupling, the width in its Breit-Wigner formula should be

$$\Gamma_{N^*}(s) = \Gamma_{N^*}^0 [0.8\rho_{\pi N}(s) + 2.1\rho_{\eta N}(s) + 3.5\rho_{\Lambda K}(s)] \quad (13)$$

instead of Eq.(8). In order to give a similar Breit-Wigner amplitude squared  $|BW(p_{N^*})|^2$  as using Eq.(8) with  $M_{N^*} = 1535MeV$  and  $\Gamma_{N^*}^0 = 150MeV$ , we need  $M_{N^*} \approx 1400MeV$  and




FIG. 3: The cross section of the reaction  $pp \rightarrow pK^+\Lambda$  as a function of the excess energy with data from Refs.[16] (circle), [17] (triangle) and [21] (square). The dashed and dotted lines represent the contribution from  $N^*(1535)$  and other  $N^*$  resonances, respectively. The solid line is the sum. The left and right graphs are the results without and with including  $\Lambda K$  term in the  $\Gamma_{N^*}(s)$  for  $N^*(1535)$ .

$\Gamma_{N^*}^0 = 270 MeV$  when using Eq.(13). Note that the two-body phase space factors  $\rho_{\eta N}(s)$  and  $\rho_{\Lambda K}(s)$  are extended to below their corresponding thresholds to be pure imaginary as the Flatté formulation for  $f_0(980)$  meson [22].

In Fig.4, we show the Breit-wigner amplitude squared vs  $s^{1/2}$  for the two cases without (dashed line) and with (dotted line)  $\Lambda K$  channel contribution included in the energy-dependent width for the  $N^*(1535)$ . As a comparison, we also show the case assuming a constant width  $\Gamma_{N^*}(s) = 98 MeV$  with  $M_{N^*} = 1515 MeV$  (solid line). The three kinds of parametrization for the  $N^*(1535)$  amplitude give a similar amplitude squared, hence do not influence much on previous calculations on various processes involving the  $N^*(1535)$  resonance by using the Breit-Wigner formula without including the  $\Lambda K$  channel in the width. As an example, we show in Fig.3 (right) the results including the  $\Lambda K$  channel in  $\Gamma_{N^*}(s)$ . Comparing results in Fig.3 (left) without including the  $\Lambda K$  channel in  $\Gamma_{N^*}(s)$ , while the fit to the data for the energies between 10 MeV and 400 MeV improves a little bit, the overall shape looks very similar. However, the important point is that by including the large  $N^* K \Lambda$  coupling in the coupled channel Breit-Wigner formula for the  $N^*(1535)$ , its Breit-Wigner mass is reduced to be around 1400 MeV, much smaller than previous value of about 1535




FIG. 4: Breit-wigner amplitude squared vs  $s^{1/2}$  with a constant width (solid line), energy-dependent width without (dashed line) and with (dotted line)  $\Lambda K$  channel contribution included.

MeV obtained without including its coupling to  $K\Lambda$ . This will have important implication on various model calculations on its mass.

The second important implication of the large  $N^*K\Lambda$  coupling is that the  $N^*(1535)$  should have large  $s\bar{s}$  component in its wave function. It has been suggested to be a quasi-bound ( $K\Sigma$ - $K\Lambda$ )-state [6]. Based on this picture, the effective coupling of  $N^*(1535)$  to  $K\Lambda$  is predicted to be about 0.5  $\sim$  0.7 times of that for  $N^*(1535)$  to  $\eta N$  [7], which is about a factor 2 smaller than the value obtained here. Alternatively, the strangeness may mix into the  $N^*(1535)$  in the form of some pentaquark configuration [23]. According to Ref.[23], the  $[4]_X[31]_{FS}[211]_F[22]_S(qqq s\bar{s})$  pentaquark configuration has the largest negative flavor-spin dependent hyperfine interaction for  $1/2^-$   $N^*$  resonance. Hence the  $1/2^-$   $N^*(1535)$  resonance may have much larger  $(qqqs\bar{s})$  pentaquark configuration than  $1/2^+$   $N^*$  resonances, for which the penta-quark configurations with the largest negative flavor-spin dependent hyperfine interaction are non-strange ones, such as  $[31]_X[4]_{FS}[22]_F[22]_S(qqqq\bar{q})$  configuration [23]. This will result in a large  $N^*K\Lambda$  coupling. A concrete calculation in this picture should be very useful for understanding the nature of the  $N^*(1535)$ . A recent study of the strangeness in the proton [24] suggests that the strangeness in the nucleon and its excited states  $N^*$  are

most likely in the form of pentaquark instead of meson-cloud configurations.

Another implication of the large  $N^*(1535)K\Lambda$  coupling is that many previous calculations on various  $K\Lambda$  production processes without including this coupling properly should be re-examined. A proper treatment of the  $N^*(1535)$  contribution may help to extract properties of other  $N^*$  resonances more reliably.

In summary, from the recent BES data on  $J/\psi \rightarrow \bar{p}p\eta$  and  $\psi \rightarrow \bar{p}K^+\Lambda$ , the  $g_{N^*(1535)K\Lambda}/g_{N^*(1535)p\eta}$  ratio is deduced to be  $1.3 \pm 0.3$  which is also compatible with data from  $pp \rightarrow pK^+\Lambda$ ,  $\pi p \rightarrow K\Lambda$  and  $\gamma p \rightarrow K\Lambda$  processes. By including the large  $N^*(1535)K\Lambda$  coupling into the Breit-Wigner formula for the  $N^*(1535)$ , a much lower Breit-Wigner mass ( $\sim 1400 MeV$ ) is obtained for the  $N^*(1535)$ . These new properties have important implication on the nature of the lowest negative-parity  $N^*$  resonance. The  $N^*(1535)1/2^-$  could be the lowest  $L = 1$  orbital excited ( $3q$ ) state with a large admixture of  $[4]_X[31]_{FS}[211]_F[22]_S(qqq\bar{s})$  pentaquark component while the  $N^*(1440)$  could be the lowest radial excited ( $3q$ ) state with a large admixture of  $[31]_X[4]_{FS}[22]_F[22]_S(qqqq\bar{q})$  pentaquark component. While the lowest  $L = 1$  orbital excited ( $3q$ ) state should have a mass lower than the lowest radial excited ( $3q$ ) state, the  $(qqq\bar{s})$  pentaquark component has a higher mass than  $(qqqq\bar{q})$  pentaquark component. This makes the  $N^*(1535)$  having an almost degenerate mass with the  $N^*(1440)$ .

We thank S.Jin, D.Riska and A.Sibirtsev for useful discussion. This work is partly supported by the National Nature Science Foundation of China under grants Nos. 10225525, 10435080 and by the Chinese Academy of Sciences under project No. KJCX2-SW-N02.

- 
- [1] S.Capstick and W.Roberts, Prog. Part. Nucl. Phys. **45** S241 (2000), and references therein.
  - [2] L.Y.Glozman and D.O.Riska, Phys. Rept. **268**, 263 (1996); L.Y.Glozman, Nucl. Phys. **A663-664**, 103 (2000).
  - [3] R.Jaffe and F.Wilczek, Phys. Rev. Lett. **91**, 232003 (2003).
  - [4] A.Zhang et al., hep-ph/0403210.
  - [5] Particle Data Group, Phys. Lett. **B592**, 1 (2004), and relevant references therein.
  - [6] N.Kaiser, P.B.Siegel and W.Weise, Phys. Lett. **B362**, 23 (1995); N.Kaiser, T.Waas and W.Weise, Nucl. Phys. **A612**, 297 (1997).

- [7] T.Inoue, E.Oset and J.Vicente Vacas, Phys. Rev. **C65**, 035204 (2002).
- [8] G.Penner and U.Mosel, Phys. Rev. **C66**, 055211-055212 (2002).
- [9] S.Capstick and W.Roberts, Phys. Rev. **D58**, 074011 (1998).
- [10] J.Q. Wu and C.M. Ko, Nucl. Phys. **A499**, 810 (1989).
- [11] W.T.Chiang, B.Saghai, F.Tabakin, T.-S.H.Lee, Phys. Rev. **C69**, 065208 (2004); Phys. Lett. **B517**, 101 (2001).
- [12] H.B.Li et al., (BES Collaboration), Nucl. Phys. **A675**, 189c (2000); M.Ablikim et al., (BES Collaboration), hep-ex/0405030.
- [13] J.Z.Bai et al., (BES Collaboration), Phys. Lett. **B510**, 75 (2001).
- [14] B.S.Zou et al., (BES Collaboration), hep-ph/0110264, Proc. of MENU2001,  $\pi N$  Newsletter **16**, 174 (2002).
- [15] H.X.Yang et al., (BES Collaboration), Proc. of MENU2004, Int. J. Mod. Phys. **A20** (2005), in press.
- [16] J.Balewski et al., Phys. Lett. **B420**, 211 (1998); R.Bilger et al., Phys. Lett. **B420**, 217 (1998); S.Sewerin et al., Phys. Rev. Lett. **83**, 682 (1999).
- [17] P.Kowina et al., Eur. Phys. J. **A22**, 293 (2004).
- [18] M.Benmerrouche, N.C.Mukhopadhyay and J.F.Zhang, Phys. Rev. Lett. **77**, 4716 (1996); Phys. Rev. **D51**, 3237 (1995).
- [19] M.G.Olsson and E.T.Osypowski, Nucl. Phys. **B87**, 399 (1975); Phys. Rev. **D17**, 174 (1978).
- [20] K.Tsushima, A.Sibirtsev, A.W.Thomas, Phys. Rev. **C59**, 369 (1999).
- [21] Landolt-Börnstein, New Series, I/12, Ed. H. Schopper, Springer-Verlag (1988).
- [22] S.Flatté, Phys. Lett. **B63**, 224 (1976); B.S.Zou and D.V.Bugg, Phys. Rev. **D48**, 3948 (1993); N.A.Törnqvist, Z. Phys. **C68**, 647 (1995).
- [23] L.Hannelius and D.O.Riska, Phys. Rev. **C62**, 045204 (2000).
- [24] B.S.Zou and D.O.Riska, hep-ph/0502225.