Variational treatment of quenched QED using the worldline technique

A. W. Schreiber^a, C. Alexandrou^b and R. Rosenfelder^c

^aDepartment of Physics and Mathematical Physics and Research Centre for the Subatomic Structure of Matter, University of Adelaide, Adelaide, S. A. 5005, Australia

^bDepartment of Natural Sciences, University of Cyprus, CY-1678 Nicosia, Cyprus

^cPaul Scherrer Institute, CH-5232 Villigen PSI, Switzerland

1. INTRODUCTION

Perturbative Quantum Electrodynamics has been successfully tested to unparalleled precision for around half a century. In addition, and rather more recently, the theory's behaviour in its strong coupling limit has attracted a growing amount of attention. The reason for this is not only that it serves as a simple prototype gauge theory for testing nonperturbative calculational techniques which one then goes on to apply to QCD, but also because it exhibits interesting behaviour in its own right. In particular, it now appears to be quite well established that QED breaks chiral symmetry dynamically as long as the coupling is sufficiently large.

Traditionally, strongly coupled QED has been studied using Schwinger-Dyson, Lattice and renormalization group approaches. In this contribution we describe a rather different method, namely a variational approach using the worldline formulation of field theory. We have developed this technique in the context of a scalar field theory [1] and we applied it here, for the first time, to a gauge theory with spin- $\frac{1}{2}$ particles. Although it is too early for quantitative results, qualitatively the approach seems promising: The method is formulated covariantly in Minkowski space-time, the Ward-Takahashi identities are satisfied, the connection between the bare- and pole-masses of the electron is gauge independent, the known exact result for the infrared dependence of the on-shell renormalized propagator is reproduced and corrections to the variational approximation can be calculated systematically.

2. THE WORLDLINE TECHNIQUE

In recent years a rather large literature on the worldline technique has developed [2], so it suffices here to just provide a general outline. We shall concentrate on Green functions within quenched QED involving only one fermion line – i.e. the propagator, vertex function, Compton scattering etc. The theory is quenched by integrating out the fermions and setting the resulting determinant equal to unity. After differentiating the appropriate generating functional Z with respect to the external fermion's source one obtains the generating functional for the general (2 + n)-point function

$$Z'[j,y] = \int \mathcal{D}A < y \mid \frac{1}{i\partial - g\mathcal{A} - M_0} \mid 0 > \exp\left\{iS_0[A] + (j,A)\right\}.$$
(1)

Here j is the source for the photon field A, M_0 is the bare mass of the electron, g is the coupling constant and $S_0[A]$ is the gauge part of the standard QED action with Lagrangian

$$\mathcal{L}_0(A) = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} m_\gamma^2 A^2 - \frac{1}{2\xi} (\partial \cdot A)^2 \,. \tag{2}$$

In order to regulate infrared divergences a photon mass m_{γ} has been introduced and the gauge has been fixed covariantly by employing the gauge fixing parameter ξ .

We would like to integrate out the photon field. It is not possible to do this at this stage because it appears in the electron propagator in Eq. (1) in a non-gaussian way. In order to proceed, the fermion is re-introduced in terms of a path integral over its worldline x(t) as well as a corresponding path integral over a Grassmann valued function $\zeta(t)$ which contains the information about the electron's spin degree of freedom [3]. One obtains

$$< y \mid \frac{1}{i\partial - g\mathcal{A} - M_0} \mid 0 > = \int_0^\infty dT \, N(T) \, \int d\chi \int \mathcal{D}x \mathcal{D}\zeta \, e^{\,iS[x,\zeta]} \quad , \tag{3}$$

where N(T) provides the proper normalization, the integration parameter χ is a Grassmannian partner to the proper time T, the four-dimensional bosonic path integral has the boundary conditions x(0) = 0 and x(T) = y while the five dimensional fermionic path integral has the single boundary condition $\zeta^{\mu}(0) + \zeta^{\mu}(T) = i\gamma^{5}\gamma^{\mu}$, $\zeta^{5}(0) + \zeta^{5}(T) = i\gamma^{5}$. The worldline action $S[x, \zeta]$ is given by

$$S[x,\zeta] = S_0[x,\zeta] + g \int_0^T dt \left\{ -\dot{x}_\mu(t) A^\mu[x(t)] + i F_{\mu\nu}[x(t)] \zeta^\mu(t) \zeta^\nu(t) \right\} \quad .$$
(4)

As expected for a relativistic particle the free part of the action $S_0[x, \zeta]$ includes, apart from a kinetic term, further contributions which provide coupling between orbital and spin motion as well as a kinetic term for the spin

$$S_0[x,\zeta] = \int_0^T dt \left\{ -\frac{\dot{x}^2(t)}{2} - \frac{i}{T} \left[\dot{x}^\mu(t)\zeta_\mu(t) - M_0\zeta^5(t) \right] \chi - i\zeta_m(t)\dot{\zeta}^m(t) - \frac{i}{T}\zeta_m(T)\zeta^m(0) \right\}.$$
(5)

Here m = 0, 1, 2, 3, 5 and the metric $g^{mn} = \text{diag}(1, -1, -1, -1, -1)$ is used for short-hand notation. Note that the gauge field A appears linearly in Eq. (4) so that we may now integrate out the photons in the generating functional (1). Differentiating n times with respect to the photon source j (and then setting j = 0) as well as taking the Fourier transform yields the general Green function of an electron interacting with n external photons. With p defined as the ingoing electron momentum and p' ($\{q_i\}$) the outgoing electron (photon) momenta, and the external photon legs as well as the momentum conserving delta function removed, one obtains

$$G_{2,n}^{\{\mu_i\}}(p,p';\{q_i\}) = \text{const.} \int_0^\infty dT \int d\chi \exp\left[-\frac{i}{2}M_0^2T + \frac{i}{2}M_0\gamma_5\chi\right] \int \mathcal{D}\tilde{x}\mathcal{D}\zeta \ e^{i[S_{\text{eff}}+p'\cdot y]} \\ \cdot \prod_{i=1}^n \left[g\int_0^T d\tau_i \ e^{iq_i\cdot x(\tau_i)} \ J^{\mu_i}[-q_i, x(\tau_i), \zeta(\tau_i)]\right] \gamma_5 \quad , \qquad (6)$$

where the $\mathcal{D}\tilde{x}$ implies an integration over the endpoint y in addition to the path integral $\mathcal{D}x$ (similarly, we will henceforth define \tilde{S}_{eff} to be $S_{\text{eff}} + p' \cdot y$). Because the 5th component of the Grassmann function ζ is non-interacting, this integration has been carried out in this equation. The effective action contains the free term in Eq. (5) as well as an interaction dependent piece

$$S_{I}[x,\zeta] = -\frac{g^{2}}{2} \int_{0}^{T} dt_{1} dt_{2} \int \frac{d^{4}q}{(2\pi)^{4}} G^{\mu\nu}(q) J_{\mu}[q,x(t_{1}),\zeta(t_{1})] J_{\nu}[-q,x(t_{2}),\zeta(t_{2})] \\ \cdot \exp\left[-iq \cdot (x(t_{1})-x(t_{2}))\right]$$
(7)

Here $G^{\mu\nu}$ is the photon propagator and the fermion current is

$$J_{\mu}[q, x(t), \zeta(t)] = \dot{x}_{\mu}(t) + 2\zeta_{\mu}(t) q \cdot \zeta(t) \quad , \tag{8}$$

which corresponds to the usual Gordon decomposition for the Dirac current of a particle of unit mass into a convection and a spin current.

It is a nice feature of the worldline representation that it is trivial to verify the Ward-Takahashi identities for an arbitrary Green function. Because of the Grassmannian nature of $\zeta(t)$ a contraction of an external momentum q^{μ} into the appropriate current J_{μ} just yields a total derivative, i.e. $q_i \cdot J_i = q_i \cdot \dot{x}(\tau_i)$, which allows one to do the τ_i integration associated with q_i , i.e.

$$\int_{0}^{T} d\tau_{i} e^{iq_{i} \cdot x(\tau_{i})} q_{i} \cdot J[-q_{i}, x(\tau_{i}), \zeta(\tau_{i})] = \frac{1}{i} \left(e^{iq_{i} \cdot y} - 1 \right) \quad .$$
(9)

The remaining q_i dependence here may be absorbed in a shifted momentum p' in the momentum space action $S_{\text{eff}} + p' \cdot y$ and so $q_{\mu_j} \cdot G_{2,n}^{\{\mu_i\}}$ is related to the difference of two (2+(n-1))-point functions:

$$\frac{i}{g}q_{\mu_j} \cdot G_{2,n}^{\{\mu_i\}}(p,p';\{q_i\}) = G_{2,n-1}^{\{\mu_i\}}(p+q_i,p'+q_i;\{q_i\}) - G_{2,n-1}^{\{\mu_i\}}(p,p';\{q_i\}) \quad , \quad (10)$$

where the notation should be self explanatory. This is the correct result for the Ward-Takahashi identity for the (2 + n)-point function untruncated in the external fermion legs. Because of the simplicity of the above argument, based only on the structure of the fermionic current, it is quite easy to ensure that the variational approximation doesn't violate it.

3. THE VARIATIONAL APPROXIMATION

Eq. (6) provides an exact representation of all Green functions occurring in quenched QED involving only one fermion line. It is not possible to carry out the remaining path integrals over x(t) and $\zeta(t)$ because they do not occur quadratically in the action (see Eq. (7)). We shall calculate the Green functions of the theory in an approximate manner through the use of the Feynman-Jensen variational principle. This technique has a rather long history, having its origin in the polaron problem [4] and it was applied, more recently, to a scalar relativistic field theory by us [1]. In order to illustrate the idea, let us consider the electron propagator (i. e. n = 0 in Eq. (6)). We rewrite the path integral by adding and subtracting to \tilde{S}_{eff} a trial action \tilde{S}_t :

$$\int \mathcal{D}\tilde{x}\mathcal{D}\zeta e^{i\tilde{S}_{\text{eff}}} = \left(\int \mathcal{D}\tilde{x}\mathcal{D}\zeta \ e^{i\tilde{S}_{t}}\right) \frac{\int \mathcal{D}\tilde{x}\mathcal{D}\zeta \ e^{i(S_{\text{eff}}-S_{t})} \ e^{iS_{t}}}{\int \mathcal{D}\tilde{x}\mathcal{D}\zeta \ e^{i\tilde{S}_{t}}} \equiv \left(\int \mathcal{D}\tilde{x}\mathcal{D}\zeta \ e^{i\tilde{S}_{t}}\right) \langle e^{i\Delta\tilde{S}} \rangle_{\tilde{S}_{t}} \quad , (11)$$

i.e. we need to evaluate the average of the exponential of $i\Delta \tilde{S}$, weighted with the function $e^{i\tilde{S}_t}$. The approximation consists of replacing the average of the exponential by the exponential of the average, i.e.

$$\langle e^{i\Delta\tilde{S}}\rangle_{\tilde{S}_t} \approx e^{i\langle\Delta\tilde{S}\rangle_{\tilde{S}_t}}$$
 , (12)

where \approx indicates equality at the stationary point of the right hand side under unrestricted variations of the functional \tilde{S}_t . If one restricts \tilde{S}_t to depend only on (non-local) quadratic functions of x(t) and $\zeta(t)$ then the path integrals may be performed analytically. However, only the parameters in the trial action are then left for variation and so it is important to note that one can systematically calculate the corrections to the variational approximation by including higher terms in the cumulant expansion of which Eq. (12) is the first term. Of course, in order for the approximation to be a useful one, the essential physics should already be contained in the leading term.

4. THE VARIATIONAL PROPAGATOR

In order to assess the feasibility of the program outlined above we have calculated the electron propagator, using a very simple trial action

$$\tilde{S}_t = \lambda' p \cdot y + S_0[x, \zeta] + \int_0^T dt_1 dt_2 f(t_1 - t_2) \left[x(t_1) - x(t_2) \right]^2 \quad , \tag{13}$$

i.e. one which has the same dependence on the fermionic degrees of freedom $\zeta(t)$ as the free action. Only the dependence on the bosonic degrees of freedom x(t) has been changed from the free action through the use of the variational parameter λ' and the variational function f which describes the retardation effects on the electron due to multiple photon emission and re-absorption. This is certainly not an optimal trial action and it only serves to illustrate some of the qualitative features of what can be expected. In particular, the trial action (13) does not respect the supersymmetry between orbital and spin degrees of freedom which is present in the exact wordline action [5]. Work with a more general trial action is underway. Also, we shall only calculate the propagator near its mass-shell: This has the great advantage that in Fourier space the trial action effectively diagonalizes, i.e.

$$\int_{0}^{T} dt_{1} dt_{2} f(t_{1} - t_{2}) \left[x(t_{1}) - x(t_{2}) \right]^{2} = \sum_{k,k'=0}^{\infty} A_{kk'} b_{k} \cdot b_{k'} \stackrel{\text{onshell}}{\Longrightarrow} \sum_{k=0}^{\infty} A_{kk} b_{k}^{2} \quad , \ (14)$$

where b_k are the Fourier components of the path x(t) and A_{kk} become the variational parameters taking the place of the function $f(t_1 - t_2)$ (not to be confused with the photon field which has been integrated out).

With this trial action, and employing Pauli-Villars regularization as well as on-shell renormalization, the connection between the physical mass M_{phys} and the bare mass M_0 becomes

$$M_{\rm phys}^2 \left\{ 1 - (1-\lambda)^2 - 2 \, \frac{\Omega[A,\lambda] + V[A,\lambda]}{M_{\rm phys}^2} \right\} - M_0^2 = 0 \tag{15}$$

where $\lambda = \lambda'/A_{00}$ (first discussed by Mano [6]). The variational parameters should be fixed through the variational equation

$$\delta \left\{ M_{\text{phys}}^2 \left[1 - (1 - \lambda)^2 \right] - 2 \Omega[A, \lambda] + V[A, \lambda] \right\} = 0 \quad .$$

$$\tag{16}$$

Here $V[A, \lambda]$ is a functional which essentially arises from the required averaging of S_I (see Eq. (11)) while $\Omega[A, \lambda]$ gets contributions from the averaging of $\tilde{S}_0 - \tilde{S}_t$ and the quadratic fluctuations of the trial action (apart from the Minkowski formulation it is identical with the expression given in Ref. [1]). Both V and Ω are gauge independent, as they should be, and hence the variational parameters are also gauge independent. If, for $M_0 = 0$, Eqs. (15) and (16) have a common solution with $M_{\rm phys} \neq 0$ then the theory exhibits dynamical mass generation. At present it is not known whether the above trial action admits a solution such as this.

The contributions to the residue, on the other hand, may conveniently be broken up into a term F_g^{χ} which is proportional to χ and two terms, F_g^I and F_{qq}^I , which are not. (The subscripts g and qq on these functions indicate whether they arise from the $g_{\mu\nu}$ or $q_{\mu}q_{\nu}$ terms in the photon propagator.) We obtain

$$Z_2 = \frac{N_0(A)}{2} \left[1 + \frac{M_0}{\lambda M_{\text{phys}}} + \frac{1-\lambda}{\lambda A_{0,0}} + \frac{2i}{\lambda} F_g^{\chi}(A,\lambda) \right] e^{i \left[F_g^I(A,\lambda) + F_{qq}^I \right]} , \qquad (17)$$

where $N_0(A)$ is an overall factor of no interest here. The important thing to note is that this expression only contains infrared divergences in the functions occurring in the exponential factor, and these may in fact be extracted *analytically*. Indeed, the only gauge dependent function F_{qq}^I has the closed form

$$F_{qq}^{I} = -i\frac{\alpha}{4\pi} \left(1 - \xi\right) \log \frac{\Lambda^2}{m_{\gamma}^2} \quad , \tag{18}$$

resulting (together with the additional infrared divergence contained in F_g^I) in a residue which contains all its infrared and gauge dependent behaviour in the overall prefactor

$$Z_2 \propto \left(\frac{\Lambda}{m_{\gamma}}\right)^{\frac{\alpha}{2\pi}(3-\xi)}$$
 (19)

This is the well-known Bloch-Nordsieck result [7].

In conclusion, we have summarized here the preliminary results which have been obtained in the first application of the polaron variational technique to QED. We believe that the qualitative success shown so far warrants a more quantitative study using a more realistic trial action. Work on this is underway and will be published elsewhere.

REFERENCES

- 1. R. Rosenfelder and A. W. Schreiber, Phys. Rev. D 53 (1996) 3337; *ibid.* p. 3354.
- 2. For example, see C. Schubert, Acta Phys. Polon. **B** 27 (1996) 3965.
- 3. E. S. Fradkin and D. M. Gitman, Phys. Rev. D 44 (1991) 3230.
- 4. R. P. Feynman, Phys. Rev. **97** (1955) 660.
- L. Brink, P. Di Vecchia and P. Howe, Nucl. Phys. B 118 (1977) 76; F. Ravndal, Phys. Rev. D 21 (1980) 2823.
- 6. K. Mano, Progr. Theor. Phys. **14** (1955) 435.
- 7. See e.g. L. S. Brown: Quantum Field Theory, Cambridge University Press, 1992.